

GAUGE-INVARIANT IDEALS OF C^* -ALGEBRAS OF BOOLEAN DYNAMICAL SYSTEMS

TOKE MEIER CARLSEN AND EUN JI KANG[†]

ABSTRACT. We enlarge the class of C^* -algebras of Boolean dynamical systems in order to include all weakly left-resolving normal labelled space C^* -algebras in it. We prove a gauge-invariant uniqueness theorem and classify all gauge-invariant ideals of these C^* -algebras of generalized Boolean dynamical systems and describe the corresponding quotients as C^* -algebras of relative generalized Boolean dynamical systems.

1. INTRODUCTION

Inspired by the C^* -algebra of labelled graphs introduced in [2], the class of C^* -algebras of Boolean dynamical systems for which each action has compact range and closed domain was introduced in [4]. In the setting of [4], the class of C^* -algebras of Boolean dynamical systems contains many labelled graph C^* -algebras, but not all of them.

In this paper, we enlarge the class of C^* -algebras of Boolean dynamical systems so that it contains all weakly left-resolving normal labelled space C^* -algebras. As a result, this class of C^* -algebras contains all graph C^* -algebras, all ultragraph C^* -algebras, C^* -algebras of shift spaces and homeomorphism C^* -algebras over 0-dimensional compact spaces, among others.

This is done by considering pairs consisting of a Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta)$ and a family $(\mathcal{I}_\alpha)_{\alpha \in \mathcal{L}}$ of ideals in \mathcal{B} such that $\theta_\alpha(\mathcal{B}) \subseteq \mathcal{I}_\alpha$ for each α , and associate a universal C^* -algebra $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ to each such pair. We call such a pair $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ a *generalized Boolean dynamical system*.

If $(\mathcal{B}, \mathcal{L}, \theta)$ is a Boolean dynamical system with compact range \mathcal{R}_α and closed domain as in [4] and we let $\mathcal{I}_\alpha = \{B \in \mathcal{B} : B \subseteq \mathcal{R}_\alpha\}$ for each α , then $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ is canonically isomorphic to $C^*(\mathcal{B}, \mathcal{L}, \theta)$. If $(E, \mathcal{L}, \mathcal{B})$ is a weakly left-resolving normal labelled space as in [1] where (E, \mathcal{L}) is a labelled graph over \mathcal{A} , and $C^*(E, \mathcal{L}, \mathcal{B})$ is the C^* -algebra associated with $(E, \mathcal{L}, \mathcal{B})$ in [1, Definition 2.5], then $C^*(\mathcal{B}, \mathcal{A}, \theta, \mathcal{I}_{r(\alpha)})$ is canonically isomorphic to $C^*(E, \mathcal{L}, \mathcal{B})$ where θ is the action of \mathcal{A} on \mathcal{B} given by $\theta_\alpha(A) = r(A, \alpha)$ and $\mathcal{I}_{r(\alpha)} = \{B \in \mathcal{B} : B \subseteq r(\alpha)\}$.

The second goal of the paper is to give a description of the gauge-invariant ideals of our C^* -algebras and thus generalize the description of the gauge-invariant ideals of C^* -algebras of set-finite, receiver set-finite and weakly left-resolving labelled spaces

2000 *Mathematics Subject Classification.* 46L05, 46L55.

Key words and phrases. Relative Cuntz-Pimsner algebras, Cuntz-Pimsner algebras, C^* -algebras of Boolean dynamical systems, labelled graph C^* -algebras, gauge-invariant uniqueness theorem, gauge-invariant ideals.

Research partially supported by NRF-2017R1D1A1B03030540[†].

given in [7] and the description of the gauge-invariant ideals of C^* -algebras of locally finite Boolean dynamical systems given in [4].

Working with gauge-invariant ideals of generalized Boolean dynamical system, we found it convenient to use certain extensions of C^* -algebras of generalized Boolean dynamical systems. Such an extension is constructed from a generalized Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ together with an ideal \mathcal{J} of \mathcal{B}_{reg} . We call such a system a *relative generalized Boolean dynamical system*, and we associate a universal C^* -algebra $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ to it. These C^* -algebras of relative generalized Boolean dynamical systems are generalizations of relative graph C^* -algebras introduced by Muhly and Tomforde in [12]. If $\mathcal{J} = \mathcal{B}_{reg}$, then $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J}) = C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$.

By imitating a construction in [1], we show that if $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is a relative generalized Boolean dynamical system, then $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ can be constructed as relative Cuntz–Pimsner algebra, and then use Katsura’s gauge-invariant uniqueness theorem for relative Cuntz–Pimsner algebras [9, Corollary 11.8] to obtain a gauge-invariant uniqueness theorem for $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$. We then use this to show that there is a one-to-one correspondences between gauge-invariant ideals of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ and pairs $(\mathcal{H}, \mathcal{S})$ where \mathcal{H} is a hereditary \mathcal{J} -saturated ideal of \mathcal{B} and \mathcal{S} is an ideal of $\{A \in \mathcal{B} : \theta_\alpha(A) \in \mathcal{H} \text{ for all but finitely many } \alpha\}$ such that $\mathcal{H} \cup \mathcal{J} \subseteq \mathcal{S}$. We show in addition that the quotient by the ideal corresponding to $(\mathcal{H}, \mathcal{S})$ is isomorphic to the C^* -algebra of relative generalized Boolean dynamical system that can be constructed from $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ and $(\mathcal{H}, \mathcal{S})$.

We also use a construction from [9] to show that the C^* -algebra $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ of the relative generalized Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is canonically isomorphic to the C^* -algebra $C^*(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha)$ of a generalized Boolean dynamical system $(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha)$, and we show that if $(\mathcal{B}, \mathcal{L}, \theta)$ is a Boolean dynamical system, \mathcal{J} is an ideal of \mathcal{B}_{reg} , and $(\mathcal{I}_\alpha)_{\alpha \in \mathcal{L}}$ is a family of ideals of \mathcal{B} such that $\theta_\alpha(\mathcal{B}) \subseteq \mathcal{I}_\alpha$ for each α , then $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is a full hereditary C^* -subalgebra of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{B}; \mathcal{J})$. It follows that $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ and $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{R}_\alpha; \mathcal{J})$ where $\mathcal{R}_\alpha = \{A \in \mathcal{B} : A \subseteq \theta_\alpha(B) \text{ for some } B \in \mathcal{B}\}$, are Morita equivalent. We call the latter C^* -algebra the *C^* -algebra of the relative Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta; \mathcal{J}) := (\mathcal{B}, \mathcal{L}, \theta, \mathcal{R}_\alpha; \mathcal{J})$* or just the *$C^*$ -algebra of the Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta)$* if $\mathcal{J} = \mathcal{B}_{reg}$. We thus have that the C^* -algebra of any relative generalized Boolean dynamical system is Morita equivalent to the C^* -algebra of a Boolean dynamical system.

Précis. This rest of the paper is organized as follows.

In section 2, we recall the notions of Boolean algebras and Boolean dynamical systems. In section 3, we introduce the definition of a generalized Boolean dynamical system (Definition 3.2) and its C^* -algebra.

In section 4, we show that the C^* -algebras of Boolean dynamical systems with compact range and closed domain introduced in [4] and the C^* -algebras of weakly left-resolving normal labelled spaces considered in [1] are all C^* -algebras of generalized Boolean dynamical systems (Example 4.1 and Example 4.2).

In section 5, we construct from each relative generalized Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$, a relative Cuntz–Pimsner algebra that is isomorphic to $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ (Theorem 5.5). As a corollary, we get that $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ is isomorphic to a Cuntz–Pimsner algebra (Corollary 5.6).

In section 6, we use Katsura's gauge-invariant uniqueness theorem for relative Cuntz–Pimsner algebras [9, Corollary 11.8] to obtain a gauge-invariant uniqueness theorem for $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ (Theorem 6.1). As a corollary, a gauge-invariant uniqueness theorem for $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ will also be given (Corollary 6.2). We then show that $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is a full hereditary C^* -subalgebra of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{B}; \mathcal{J})$ (Proposition 6.3), and we construct a Boolean dynamical system $(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta})$ such that $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ and $C^*(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha)$ are isomorphic (Proposition 6.4).

In section 7, we classify the gauge-invariant ideals of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$. We show that a pair $(\mathcal{H}, \mathcal{S})$ where \mathcal{H} is a hereditary and \mathcal{J} -saturated ideal in \mathcal{B} and \mathcal{S} is an ideal of $\{A \in \mathcal{B} : \theta_\alpha(A) \in \mathcal{H} \text{ for all but finitely many } \alpha\}$ such that $\mathcal{H} \cup \mathcal{J} \subseteq \mathcal{S}$ give rises to a gauge-invariant ideal $I_{(\mathcal{H}, \mathcal{S})}$ of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ (Lemma 7.1), and prove that the quotient $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})/I_{(\mathcal{H}, \mathcal{S})}$ of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ by the gauge-invariant ideal $I_{(\mathcal{H}, \mathcal{S})}$ is canonically isomorphic to a relative Boolean C^* -algebra $C^*(\mathcal{B}/\mathcal{H}, \mathcal{L}, \theta, [\mathcal{I}_\alpha]; [\mathcal{S}])$ (Proposition 7.3). Using this result, we show that the gauge-invariant ideals of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ are in one-to-one correspondence with the pairs $(\mathcal{H}, \mathcal{S})$ (Theorem 7.4).

2. PRELIMINARIES

In this section, we review the notions of Boolean algebras and Boolean dynamical systems. For the most part we use the notational conventions of [4].

2.1. Boolean algebras. A *Boolean algebra* is a set \mathcal{B} with a distinguished element \emptyset and maps $\cap : \mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B}$, $\cup : \mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B}$ and $\setminus : \mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B}$ such that $(\mathcal{B}, \cap, \cup)$ is a distributive lattice, $A \cap \emptyset = \emptyset$ for all $A \in \mathcal{B}$, and $(A \cap B) \cup (A \setminus B) = A$ and $(A \cap B) \cap (A \setminus B) = \emptyset$ for all $A, B \in \mathcal{B}$. The Boolean algebra \mathcal{B} is called *unital* if there exists $1 \in \mathcal{B}$ such that $1 \cup A = 1$ and $1 \cap A = A$ for all $A \in \mathcal{B}$. (often, Boolean algebras are assumed to be unital and what we here call a Boolean algebra is often called a *generalized Boolean algebra*).

We call $A \cup B$ the *union* of A and B , $A \cap B$ the *intersection* of A and B , and $A \setminus B$ the *relative complement* of B with respect to A . A subset $\mathcal{B}' \subseteq \mathcal{B}$ is called a *Boolean subalgebra* if $\emptyset \in \mathcal{B}'$ and \mathcal{B}' is closed under taking union, intersection and the relative complement. A Boolean subalgebra of a Boolean algebra is itself a Boolean algebra.

We define a partial order on \mathcal{B} as follows: for $A, B \in \mathcal{B}$,

$$A \subseteq B \text{ if and only if } A \cap B = A.$$

Then (\mathcal{B}, \subseteq) is a partially ordered set, and $A \cup B$ and $A \cap B$ are the least upper-bound and the greatest lower-bound of A and B with respect to the partial order \subseteq . If a family $\{A_\lambda\}_{\lambda \in \Lambda}$ of elements from \mathcal{B} has a least upper-bound, then we denote it by $\cup_{\lambda \in \Lambda} A_\lambda$. If $A \subseteq B$, then we say that A is a *subset* of B .

A non-empty subset \mathcal{I} of \mathcal{B} is called an *ideal* [4, Definition 2.4] if

- (i) if $A, B \in \mathcal{I}$, then $A \cup B \in \mathcal{I}$,
- (ii) if $A \in \mathcal{I}$ and $B \in \mathcal{B}$, then $A \cap B \in \mathcal{I}$.

An ideal \mathcal{I} of a Boolean algebra \mathcal{B} is a Boolean subalgebra. For $A \in \mathcal{B}$, the ideal generated by A is defined by $\mathcal{I}_A := \{B \in \mathcal{B} : B \subseteq A\}$.

2.2. Boolean dynamical systems. A map $\phi : \mathcal{B} \rightarrow \mathcal{B}'$ between two Boolean algebras is called a *Boolean homomorphism* ([4, Definition 2.1]) if $\phi(A \cap B) = \phi(A) \cap \phi(B)$, $\phi(A \cup B) = \phi(A) \cup \phi(B)$, and $\phi(A \setminus B) = \phi(A) \setminus \phi(B)$ for all $A, B \in \mathcal{B}$.

A map $\theta : \mathcal{B} \rightarrow \mathcal{B}$ is called an *action* ([4, Definition 3.1]) on a Boolean algebra \mathcal{B} if it is a Boolean homomorphism with $\theta(\emptyset) = \emptyset$.

Given a set \mathcal{L} and any $n \in \mathbb{N}$, we define $\mathcal{L}^n := \{(\alpha_1, \dots, \alpha_n) : \alpha_i \in \mathcal{L}\}$ and $\mathcal{L}^* := \bigcup_{n \geq 0} \mathcal{L}^n$, where $\mathcal{L}^0 := \{\emptyset\}$. For $\alpha \in \mathcal{L}^n$, we write $|\alpha| := n$. For $\alpha = (\alpha_1, \dots, \alpha_n), \beta = (\beta_1, \dots, \beta_m) \in \mathcal{L}^*$, we will usually write $\alpha_1 \dots \alpha_n$ instead of $(\alpha_1, \dots, \alpha_n)$ and use $\alpha\beta$ to denote the word $\alpha_1 \dots \alpha_n \beta_1 \dots \beta_m$ (if $\alpha = \emptyset$, then $\alpha\beta := \beta$; and if $\beta = \emptyset$, then $\alpha\beta := \alpha$). For $k \in \mathbb{N}$, we let $\alpha^k := \alpha \alpha \dots \alpha$ where the concatenation on the right has k terms. Similarly we let $\alpha^0 := \emptyset$. For $1 \leq i \leq j \leq |\alpha|$, we denote by $\alpha_{[i,j]}$ the sub-word $\alpha_i \dots \alpha_j$ of $\alpha = \alpha_1 \alpha_2 \dots \alpha_{|\alpha|}$, where $\alpha_{[i,i]} = \alpha_i$.

Definition 2.1. A *Boolean dynamical system* is a triple $(\mathcal{B}, \mathcal{L}, \theta)$ where \mathcal{B} is a Boolean algebra, \mathcal{L} is a set, and $\{\theta_\alpha\}_{\alpha \in \mathcal{L}}$ is a set of actions on \mathcal{B} such that for $\alpha = \alpha_1 \dots \alpha_n \in \mathcal{L}^* \setminus \{\emptyset\}$, the action $\theta_\alpha : \mathcal{B} \rightarrow \mathcal{B}$ is defined as $\theta_\alpha := \theta_{\alpha_n} \circ \dots \circ \theta_{\alpha_1}$. We also define $\theta_\emptyset := \text{Id}$.

Remark 2.2. Given a Boolean algebra \mathcal{B} , we say that an action θ on \mathcal{B} has *compact range* [4, Definition 3.1] if $\{\theta(A)\}_{A \in \mathcal{B}}$ has a least upper-bound. We denote by \mathcal{R}_θ this least upper-bound if it exists. We say that an action θ has *closed domain* [4, Definition 3.1] if there exists $\mathcal{D}_\theta \in \mathcal{B}$ such that $\theta(\mathcal{D}_\theta) = \mathcal{R}_\theta$. In [4, Definition 3.3], a triple $(\mathcal{B}, \mathcal{L}, \theta)$ is called a Boolean dynamical system if θ_α has compact range $\mathcal{R}_{\theta_\alpha}$ and closed domain $\mathcal{D}_{\theta_\alpha}$ for each $\alpha \in \mathcal{L}$.

Notice that when we call $(\mathcal{B}, \mathcal{L}, \theta)$ a Boolean dynamical system in this paper, we do not assume that θ_α has compact range and closed domain.

For $B \in \mathcal{B}$, we define

$$\Delta_B^{(\mathcal{B}, \mathcal{L}, \theta)} := \{\alpha \in \mathcal{L} : \theta_\alpha(B) \neq \emptyset\} \text{ and } \lambda_B^{(\mathcal{B}, \mathcal{L}, \theta)} := |\Delta_B^{(\mathcal{B}, \mathcal{L}, \theta)}|.$$

We will often just write Δ_B and λ_B instead of $\Delta_B^{(\mathcal{B}, \mathcal{L}, \theta)}$ and $\lambda_B^{(\mathcal{B}, \mathcal{L}, \theta)}$.

We say that $A \in \mathcal{B}$ is *regular* ([4, Definition 3.5]) if for any $\emptyset \neq B \in \mathcal{I}_A$, we have $0 < \lambda_B < \infty$. If $A \in \mathcal{B}$ is not regular, then it is called a *singular* set. We write $\mathcal{B}_{reg}^{(\mathcal{B}, \mathcal{L}, \theta)}$ or just \mathcal{B}_{reg} for the set of all regular sets. Notice that $\emptyset \in \mathcal{B}_{reg}$.

2.3. Quotient Boolean dynamical systems. Let $(\mathcal{B}, \mathcal{L}, \theta)$ be a Boolean dynamical system and let \mathcal{J} be an ideal of \mathcal{B}_{reg} and \mathcal{H} an ideal of \mathcal{B} . As in [4], we say that \mathcal{H} is *hereditary* if $\theta_\alpha(A) \in \mathcal{H}$ for $A \in \mathcal{H}$ and $\alpha \in \mathcal{L}$, and we say that \mathcal{H} is \mathcal{J} -*saturated* if $A \in \mathcal{H}$ whenever $A \in \mathcal{J}$ and $\theta_\alpha(A) \in \mathcal{H}$ for all $\alpha \in \Delta_A$. When $\mathcal{J} = \mathcal{B}_{reg}$, then we often say *saturated* instead of \mathcal{J} -saturated.

If \mathcal{H} is an ideal of a Boolean algebra \mathcal{B} , then the relation

$$A \sim B \iff \mathbf{R} : A \cup A' = B \cup B' \text{ for some } A', B' \in \mathcal{H} \quad (1)$$

defines an equivalent relation on \mathcal{B} ([4, Definition 2.5]).

Remark 2.3. We have that $A \sim B$ if and only if either (and thus both) of the following two equivalent conditions hold.

R1 $A \cup A' = B \cup B'$ for some $A', B' \in \mathcal{H}$ with $A \cap A' = B \cap B' = \emptyset$.

R2 $A \cup C = B \cup C$ for some $C \in \mathcal{H}$.

Proof. Clearly, **R1** \implies **R** and **R2** \implies **R**. Suppose that $A \cup A' = B \cup B'$ for some $A', B' \in \mathcal{H}$. Then $A' \setminus A, B' \setminus B, A' \cup B' \in \mathcal{H}$. One sees that

$$A \cup (A' \cup B') = (A \cup A') \cup B' = (B \cup B') \cup B' = B \cup B',$$

$$B \cup (A' \cup B') = (A' \cup B') \cup B = A' \cup (A \cup A') = A \cup A',$$

and hence, $A \cup (A' \cup B') = B \cup (A' \cup B')$. Thus, **R** \implies **R2**.

Also, we have that $A \cup (A' \setminus A) = B \cup (B' \setminus B)$ with $A \cap (A' \setminus A) = B \cap (B' \setminus B) = \emptyset$. Thus, **R** \implies **R1**. \square

We denote the equivalent class of $A \in \mathcal{B}$ with respect to \sim by $[A]$ (or $[A]_{\mathcal{H}}$ if we need to specify the ideal \mathcal{H}) and the set of all equivalent classes of \mathcal{B} by \mathcal{B}/\mathcal{H} . It is easy to check that \mathcal{B}/\mathcal{H} is a Boolean algebra with operations defined by

$$[A] \cap [B] = [A \cap B], [A] \cup [B] = [A \cup B] \text{ and } [A] \setminus [B] = [A \setminus B].$$

The partial order \subseteq on \mathcal{B}/\mathcal{H} is characterized by

$$\begin{aligned} [A] \subseteq [B] &\iff A \subseteq B \cup W \text{ for some } W \in \mathcal{H} \\ &\iff [A] \cap [B] = [A]. \end{aligned}$$

If in addition \mathcal{H} is hereditary, and we define $\theta_{\alpha}([A]) = [\theta_{\alpha}(A)]$ for all $[A] \in \mathcal{B}/\mathcal{H}$ and $\alpha \in \mathcal{L}$, then $(\mathcal{B}/\mathcal{H}, \mathcal{L}, \theta)$ becomes a Boolean dynamical system. We call it a *quotient Boolean dynamical system* of $(\mathcal{B}, \mathcal{L}, \theta)$.

3. C^* -ALGEBRAS OF GENERALIZED BOOLEAN DYNAMICAL SYSTEMS

In this section, we introduce a definition of generalized Boolean dynamical systems and their C^* -algebras. Let $(\mathcal{B}, \mathcal{L}, \theta)$ be a Boolean dynamical system and let

$$\mathcal{R}_{\alpha}^{(\mathcal{B}, \mathcal{L}, \theta)} := \{A \in \mathcal{B} : A \subseteq \theta_{\alpha}(B) \text{ for some } B \in \mathcal{B}\}$$

for each $\alpha \in \mathcal{L}$. Note that each $\mathcal{R}_{\alpha}^{(\mathcal{B}, \mathcal{L}, \theta)}$ is an ideal of \mathcal{B} .

We will often, when it is clear which Boolean dynamical system we are working with, just write \mathcal{R}_{α} instead of $\mathcal{R}_{\alpha}^{(\mathcal{B}, \mathcal{L}, \theta)}$.

Remark 3.1. In [4], the notation \mathcal{R}_{α} is used to denote the least-upper bound of $\{\theta_{\alpha}(A)\}_{A \in \mathcal{B}}$ when θ_{α} has compact range.

Definition 3.2. A *generalized Boolean dynamical system* is a quadruple $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha})$ where $(\mathcal{B}, \mathcal{L}, \theta)$ is a Boolean dynamical system and $\{\mathcal{I}_{\alpha} : \alpha \in \mathcal{L}\}$ is a family of ideals in \mathcal{B} such that $\mathcal{R}_{\alpha} \subseteq \mathcal{I}_{\alpha}$ for each $\alpha \in \mathcal{L}$. A *relative generalized Boolean dynamical system* is a pentamerous $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J})$ where $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha})$ is a generalized Boolean dynamical system and \mathcal{J} is an ideal of \mathcal{B}_{reg} .

Definition 3.3. Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J})$ be a relative generalized Boolean dynamical system. A $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J})$ -*representation* is a family of projections $\{P_A : A \in \mathcal{B}\}$ and a family of partial isometries $\{S_{\alpha, B} : \alpha \in \mathcal{L}, B \in \mathcal{I}_{\alpha}\}$ such that for $A, A' \in \mathcal{B}$, $\alpha, \alpha' \in \mathcal{L}$, $B \in \mathcal{I}_{\alpha}$ and $B' \in \mathcal{I}_{\alpha'}$,

$$(i) \ P_{\emptyset} = 0, \ P_{A \cap A'} = P_A P_{A'}, \text{ and } P_{A \cup A'} = P_A + P_{A'} - P_{A \cap A'};$$

- (ii) $P_A S_{\alpha,B} = S_{\alpha,B} P_{\theta_\alpha(A)}$;
- (iii) $S_{\alpha,B}^* S_{\alpha',B'} = \delta_{\alpha,\alpha'} P_{B \cap B'}$;
- (iv) $P_A = \sum_{\alpha \in \Delta_A} S_{\alpha,\theta_\alpha(A)} S_{\alpha,\theta_\alpha(A)}^*$ for all $A \in \mathcal{J}$.

Given a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation $\{P_A, S_{\alpha,B}\}$ in a C^* -algebra \mathcal{A} , we denote by $C^*(P_A, S_{\alpha,B})$ the C^* -subalgebra of \mathcal{A} generated by $\{P_A, S_{\alpha,B}\}$.

We will show in Section 5 that there exists a universal $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation $\{p_A, s_{\alpha,B} : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_\alpha\}$ in the sense that if $\{P_A, S_{\alpha,B}\}$ is a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation in a C^* -algebra \mathcal{A} , then there exists a unique $*$ -homomorphism $\pi_{S,P} : C^*(p_A, s_{\alpha,B}) \rightarrow \mathcal{A}$ such that $\pi_{S,P}(p_A) = P_A$ and $\pi_{S,P}(s_{\alpha,B}) = S_{\alpha,B}$ for $A \in \mathcal{B}$, $\alpha \in \mathcal{L}$ and $B \in \mathcal{I}_\alpha$. We write $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ for $C^*(p_A, s_{\alpha,B})$ and call it the C^* -algebra of $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha, \mathcal{J})$.

By a *Cuntz–Krieger representation* of $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ we mean a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{B}_{reg})$ -representation, and by a *Toeplitz representation* of $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ we mean a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \emptyset)$ -representation. We write $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ for $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{B}_{reg})$ and call it the C^* -algebra of $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$, and we write $\mathcal{T}(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ for $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \emptyset)$ and call it the *Toeplitz C^* -algebra* of $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$.

When $(\mathcal{B}, \mathcal{L}, \theta)$ is a Boolean dynamical system, then we write $C^*(\mathcal{B}, \mathcal{L}, \theta)$ for $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{R}_\alpha)$ and call it the C^* -algebra of $(\mathcal{B}, \mathcal{L}, \theta)$. If moreover \mathcal{J} is an ideal of \mathcal{B}_{reg} , then we write $C^*(\mathcal{B}, \mathcal{L}, \theta; \mathcal{J})$ for $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{R}_\alpha; \mathcal{J})$ and call it the C^* -algebra of $(\mathcal{B}, \mathcal{L}, \theta; \mathcal{J})$.

We shall in Proposition 6.4 see that the C^* -algebra $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{R}_\alpha; \mathcal{J})$ of a relative generalized Boolean dynamical system is isomorphic to the C^* -algebra of a (different) generalized Boolean dynamical system $(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta}, \tilde{\mathcal{R}}_\alpha)$. Moreover, it follows from Proposition 6.3 that $C^*(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta}, \tilde{\mathcal{R}}_\alpha)$ and $C^*(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta})$ are Morita equivalent. We thus have that the C^* -algebra of any relative generalized Boolean dynamical system is Morita equivalent to the C^* -algebra of a Boolean dynamical system.

Remark 3.4. It follows from the universal property of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J}) = C^*(p_A, s_{\alpha,B})$ that there is a strongly continuous action $\gamma : \mathbb{T} \rightarrow \text{Aut}(C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J}))$, which we call the *gauge action*, such that

$$\gamma_z(p_A) = p_A \quad \text{and} \quad \gamma_z(s_{\alpha,B}) = z s_{\alpha,B}$$

for $A \in \mathcal{B}$, $\alpha \in \mathcal{L}$ and $B \in \mathcal{I}_\alpha$.

Remark 3.5. Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ be a relative generalized Boolean dynamical system. The fact that a family $\{P_A, S_{\alpha,B} : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_\alpha\}$ satisfies (i)–(iii) in Definition 3.3 is equivalent to that the family satisfies that

- (a) $P_\emptyset = 0$, $P_{A \cap A'} = P_A P_{A'}$, and $P_{A \cup A'} = P_A + P_{A'} - P_{A \cap A'}$;
- (b) $P_A S_{\alpha,B} = S_{\alpha,\theta_\alpha(A) \cap B}$;
- (c) $S_{\alpha,B} P_A = S_{\alpha,B \cap A}$;
- (d) $S_{\alpha,B}^* S_{\alpha',B'} = \delta_{\alpha,\alpha'} P_{B \cap B'}$

for $A, A' \in \mathcal{B}$, $\alpha, \alpha' \in \mathcal{L}$, $B \in \mathcal{I}_\alpha$ and $B' \in \mathcal{I}_{\alpha'}$.

Proof. By (b) and (c), we see that $P_A S_{\alpha,B} = S_{\alpha,\theta_\alpha(A) \cap B} = S_{\alpha,B} P_{\theta_\alpha(A)}$.

For the converse, note that $S_{\alpha,B}^* P_A = (P_A S_{\alpha,B})^* = (S_{\alpha,B} P_{\theta_\alpha(A)})^* = P_{\theta_\alpha(A)} S_{\alpha,B}^*$. It then follows that

$$\begin{aligned}
& \|P_A S_{\alpha,B} - S_{\alpha,\theta_\alpha(A) \cap B}\|^2 \\
&= \|(P_A S_{\alpha,B} - S_{\alpha,\theta_\alpha(A) \cap B})^* (P_A S_{\alpha,B} - S_{\alpha,\theta_\alpha(A) \cap B})\| \\
&= \|S_{\alpha,B}^* P_A S_{\alpha,B} - S_{\alpha,B}^* P_A S_{\alpha,\theta_\alpha(A) \cap B} - S_{\alpha,\theta_\alpha(A) \cap B}^* P_A S_{\alpha,B} + P_{\theta_\alpha(A) \cap B}\| \\
&= \|S_{\alpha,B}^* S_{\alpha,B} P_{\theta_\alpha(A)} - P_{\theta_\alpha(A)} S_{\alpha,B}^* S_{\alpha,\theta_\alpha(A) \cap B} - S_{\alpha,\theta_\alpha(A) \cap B}^* S_{\alpha,B} P_{\theta_\alpha(A)} + P_{\theta_\alpha(A) \cap B}\| \\
&= \|P_{\theta_\alpha(A) \cap B} - P_{\theta_\alpha(A) \cap B} - P_{\theta_\alpha(A) \cap B} + P_{\theta_\alpha(A) \cap B}\| = 0.
\end{aligned}$$

Thus, we have $P_A S_{\alpha,B} = S_{\alpha,\theta_\alpha(A) \cap B}$. One also sees that $\|S_{\alpha,B} P_A - S_{\alpha,B \cap A}\|^2 = 0$, and hence, we have $S_{\alpha,B} P_A = S_{\alpha,B \cap A}$. \square

For $\alpha = \alpha_1 \alpha_2 \cdots \alpha_n \in \mathcal{L}^* \setminus \{\emptyset\}$, we define

$$\mathcal{I}_\alpha := \{A \in \mathcal{B} : A \subseteq \theta_{\alpha_2 \cdots \alpha_n}(B) \text{ for some } B \in \mathcal{I}_{\alpha_1}\}.$$

For $\alpha = \emptyset$, we define $\mathcal{I}_\emptyset := \mathcal{B}$

Definition 3.6. Let $\{P_A, S_{\alpha,B} : A \in \mathcal{B}, \alpha \in \mathcal{L}, B \in \mathcal{I}_\alpha\}$ be a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation. For $\alpha = \alpha_1 \alpha_2 \cdots \alpha_n \in \mathcal{L}^* \setminus \{\emptyset\}$ and $A \in \mathcal{I}_\alpha$, we define

$$S_{\alpha,A} := S_{\alpha_1,B} S_{\alpha_2,\theta_{\alpha_2}(B)} S_{\alpha_3,\theta_{\alpha_2\alpha_3}(B)} \cdots S_{\alpha_n,A},$$

where $B \in \mathcal{I}_{\alpha_1}$ is such that $A \subseteq \theta_{\alpha_2 \cdots \alpha_n}(B)$. For $\alpha = \emptyset$, we also define $S_{\emptyset,A} := P_A$.

Remark 3.7. In the above definition, $S_{\alpha,A}$ is independent of the choice of $B \in \mathcal{I}_{\alpha_1}$. It is enough to show that for $|\alpha| = 2$. Put $\alpha = \alpha_1 \alpha_2 \in \mathcal{L}^*$, $A \subseteq \theta_{\alpha_2}(B_1)$ and $A \subseteq \theta_{\alpha_2}(B_2)$ for some $B_1, B_2 \in \mathcal{I}_{\alpha_1}$. Then we see that

$$\begin{aligned}
& \|S_{\alpha_1,B_1} S_{\alpha_2,A} - S_{\alpha_1,B_2} S_{\alpha_2,A}\|^2 \\
&= \|(S_{\alpha_2,A}^* S_{\alpha_1,B_1}^* - S_{\alpha_2,A}^* S_{\alpha_1,B_2}^*)(S_{\alpha_1,B_1} S_{\alpha_2,A} - S_{\alpha_1,B_2} S_{\alpha_2,A})\| \\
&= \|S_{\alpha_2,A}^* P_{B_1} S_{\alpha_2,A} - S_{\alpha_2,A}^* P_{B_1 \cap B_2} S_{\alpha_2,A} \\
&\quad - S_{\alpha_2,A}^* P_{B_1 \cap B_2} S_{\alpha_2,A} + S_{\alpha_2,A}^* P_{B_2} S_{\alpha_2,A}\| \\
&= \|P_{A \cap \theta_{\alpha_2}(B_1)} - P_{A \cap \theta_{\alpha_2}(B_1 \cap B_2)} - P_{A \cap \theta_{\alpha_2}(B_1 \cap B_2)} + P_{A \cap \theta_{\alpha_2}(B_2)}\| \\
&= \|P_A - P_A - P_A + P_A\| = 0.
\end{aligned}$$

Thus, it follows that $S_{\alpha_1,B_1} S_{\alpha_2,A} = S_{\alpha_1,B_2} S_{\alpha_2,A}$.

It is straightforward to check the following lemma.

Lemma 3.8. Let $\{P_A, S_{\alpha,B} : A \in \mathcal{B}, \alpha \in \mathcal{L}, B \in \mathcal{I}_\alpha\}$ be a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation. For $\alpha \in \mathcal{L}^*$, $A \in \mathcal{B}$ and $B \in \mathcal{I}_\alpha$, we have

- (1) $P_A S_{\alpha,B} = S_{\alpha,B} P_{\theta_\alpha(A)}$,
- (2) $P_A S_{\alpha,B} = S_{\alpha,\theta_\alpha(A) \cap B}$,
- (3) $S_{\alpha,B} P_A = S_{\alpha,B \cap A}$.

Lemma 3.9. *Let $\{P_A, S_{\alpha,B} : A \in \mathcal{B}, \alpha \in \mathcal{L}, B \in \mathcal{I}_\alpha\}$ be a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation. For $\alpha, \beta \in \mathcal{L}^*$, $A \in \mathcal{I}_\alpha$ and $B \in \mathcal{I}_\beta$, we have*

$$S_{\alpha,A}^* S_{\beta,B} = \begin{cases} P_{A \cap B} & \text{if } \alpha = \beta \\ S_{\alpha', A \cap \theta_{\alpha'}(B)}^* & \text{if } \alpha = \beta \alpha' \\ S_{\beta', B \cap \theta_{\beta'}(A)} & \text{if } \beta = \alpha \beta' \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Let $\alpha = \alpha_1 \alpha_2 \cdots \alpha_n$, $\beta = \beta_1 \beta_2 \cdots \beta_m \in \mathcal{L}^*$ and put

$$\begin{aligned} S_{\alpha,A} &= S_{\alpha_1,A'} S_{\alpha_2, \theta_{\alpha_2}(A')} S_{\alpha_3, \theta_{\alpha_2 \alpha_3}(A')} \cdots S_{\alpha_n,A}, \\ S_{\beta,B} &= S_{\beta_1,B'} S_{\beta_2, \theta_{\beta_2}(B')} S_{\beta_3, \theta_{\beta_2 \beta_3}(B')} \cdots S_{\beta_m,B}, \end{aligned}$$

where $A' \in \mathcal{I}_{\alpha_1}$ such that $A \subseteq \theta_{\alpha_2 \cdots \alpha_n}(A')$ and $B' \in \mathcal{I}_{\beta_1}$ such that $B \subseteq \theta_{\beta_2 \cdots \beta_m}(B')$. If $\alpha = \beta$, we have

$$\begin{aligned} S_{\alpha,A}^* S_{\alpha,B} &= (S_{\alpha_n,A}^* \cdots S_{\alpha_2, \theta_{\alpha_2}(A')}^* S_{\alpha_1,A'}^*) (S_{\alpha_1,B'} S_{\alpha_2, \theta_{\alpha_2}(B')} \cdots S_{\alpha_n,B}) \\ &= S_{\alpha_n,A}^* \cdots S_{\alpha_2, \theta_{\alpha_2}(A')}^* P_{A' \cap B'} S_{\alpha_2, \theta_{\alpha_2}(B')} \cdots S_{\alpha_n,B} \\ &= S_{\alpha_n,A}^* \cdots S_{\alpha_2, \theta_{\alpha_2}(A')}^* S_{\alpha_2, \theta_{\alpha_2}(B')} P_{\theta_{\alpha_2}(A' \cap B')} \cdots S_{\alpha_n,B} \\ &= S_{\alpha_n,A}^* \cdots P_{\theta_{\alpha_2}(A' \cap B')} \cdots S_{\alpha_n,B} \\ &\vdots \\ &= S_{\alpha_n,A}^* S_{\alpha_n,B} P_{\theta_{\alpha_2 \cdots \alpha_n}(A' \cap B')} \\ &= P_{A \cap B}. \end{aligned}$$

If $\alpha = \beta \alpha'$, we first note that

$$\begin{aligned} S_{\beta \alpha', A} &= (S_{\beta_1, A'} S_{\beta_2, \theta_{\beta_2}(A')} \cdots S_{\beta_{|\beta|}, \theta_{\beta_2 \cdots \beta_{|\beta|}}(A')}) S_{\alpha'_1, \theta_{\beta_2 \cdots \beta_{|\beta|}} \alpha'_1(A')} \cdots S_{\alpha'_{|\alpha'|}, A} \\ &= S_{\beta, \theta_{\beta_2 \cdots \beta_{|\beta|}}(A')} S_{\alpha', A}. \end{aligned}$$

Thus, we have

$$\begin{aligned} S_{\beta \alpha', A}^* S_{\beta, B} &= (S_{\beta, \theta_{\beta_2 \cdots \beta_{|\beta|}}(A')} S_{\alpha', A})^* S_{\beta, B} \\ &= S_{\alpha', A}^* S_{\beta, \theta_{\beta_2 \cdots \beta_{|\beta|}}(A')}^* S_{\beta, B} \\ &= S_{\alpha', A}^* P_{(\theta_{\beta_2 \cdots \beta_{|\beta|}}(A')) \cap B} \\ &= S_{\alpha', (\theta_{\beta_2 \cdots \beta_{|\beta|}} \alpha')(A')}^* \cap \theta_{\alpha'}(B) \cap A \\ &= S_{\alpha', A \cap \theta_{\alpha'}(B)}^*. \end{aligned}$$

Similarly, we also have $S_{\alpha,A}^* S_{\alpha \beta', B} = S_{\beta', B \cap \theta_{\beta'}(A)}$. Otherwise, $S_{\alpha,A}^* S_{\beta,B} = 0$ by Definition 3.3(iii). \square

As a corollary of Lemma 3.9, we have the following.

Lemma 3.10. *Let $\{P_A, S_{\alpha,B} : A \in \mathcal{B}, \alpha \in \mathcal{L}, B \in \mathcal{I}_\alpha\}$ be a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation. For $\alpha, \beta, \mu, \nu \in \mathcal{L}^*$, $A \in \mathcal{I}_\alpha$, $B \in \mathcal{I}_\beta$, $C \in \mathcal{I}_\mu$ and $D \in \mathcal{I}_\nu$, we*

have

$$(S_{\alpha,A}S_{\beta,B}^*)(S_{\mu,C}S_{\nu,D}^*) = \begin{cases} S_{\alpha,A \cap B \cap C}S_{\nu,D \cap B \cap C}^* & \text{if } \beta = \mu \\ S_{\alpha,A}S_{\nu\beta',\theta_{\beta'}(C \cap D) \cap B}^* & \text{if } \beta = \mu\beta' \\ S_{\alpha\mu',\theta_{\mu'}(A \cap B) \cap C}S_{\nu,D}^* & \text{if } \mu = \beta\mu' \\ 0 & \text{otherwise.} \end{cases}$$

Remark 3.11. Let $\{P_A, S_{\alpha,B} : A \in \mathcal{B}, \alpha \in \mathcal{L}, B \in \mathcal{I}_\alpha\}$ be a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation. We then have that

$$C^*(P_A, S_{\alpha,B}) = \overline{\text{span}}\{S_{\alpha,A}S_{\beta,B}^* : \alpha, \beta \in \mathcal{L}^* \text{ and } A \in \mathcal{I}_\alpha, B \in \mathcal{I}_\beta\} \quad (2)$$

$$= \overline{\text{span}}\{S_{\alpha,A}S_{\beta,A}^* : \alpha, \beta \in \mathcal{L}^* \text{ and } A \in \mathcal{I}_\alpha \cap \mathcal{I}_\beta\}. \quad (3)$$

Proof. It follows from Lemma 3.10 that the right-hand side of (2) is a C^* -subalgebra of $C^*(P_A, S_{\alpha,B})$, and since it contains all the generators of $C^*(P_A, S_{\alpha,B})$, they must be equal. To see the last equality, we only need to observe that $S_{\alpha,A}S_{\beta,B}^* = S_{\alpha,A}P_{A \cap B}S_{\beta,B}^* = S_{\alpha,A \cap B}S_{\beta,A \cap B}^*$ for $\alpha, \beta \in \mathcal{L}^*$ and $A \in \mathcal{I}_\alpha, B \in \mathcal{I}_\beta$. \square

4. EXAMPLES

The class of C^* -algebras of Boolean dynamical system described in Example 4.1 below was introduced in [4] to study labelled graph C^* -algebras from a more general point of view. It is shown in [4] that the class of C^* -algebras of Boolean dynamical system contains weakly left-resolving normal labelled graph C^* -algebras under some assumption. We show in the following two examples that our C^* -algebras of generalized Boolean dynamical systems contains all C^* -algebras of Boolean dynamical systems and all weakly left-resolving normal labelled graph C^* -algebras.

Example 4.1. Let $(\mathcal{B}, \mathcal{L}, \theta)$ be a Boolean dynamical system with assumptions that θ_α has compact range $\mathcal{R}_{\theta_\alpha}$ and closed domain $\mathcal{D}_{\theta_\alpha}$ for each $\alpha \in \mathcal{L}$ (see [4, Definition 3.3] or Remark 2.2). We denote the universal Cuntz–Krieger Boolean C^* -algebra constructed in [4] by $C^*(s_\alpha, p_A)$. We then show that

$$C^*(\mathcal{B}, \mathcal{L}, \theta) \cong C^*(s_\alpha, p_A).$$

We check that

$$\{p_A, s_\alpha p_B : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{R}_\alpha\}$$

is a Cuntz–Krieger representation of $(\mathcal{B}, \mathcal{L}, \theta)$ in $C^*(s_\alpha, p_A)$:

- (i) p_A 's clearly satisfy Definition 3.3(i),
- (ii) $p_A(s_\alpha p_B) = s_\alpha p_{\theta_\alpha(A)} p_B = (s_\alpha p_B) p_{\theta_\alpha(A)}$,
- (iii) $(s_\alpha p_B)^*(s_{\alpha'} p_{B'}) = p_B s_\alpha^* s_{\alpha'} p_{B'} = \delta_{\alpha, \alpha'} p_B p_{\mathcal{R}_{\theta_\alpha}} p_{B'} = \delta_{\alpha, \alpha'} p_{B \cap B'}$,
- (iv) $p_A = \sum_{\alpha \in \Delta_A} s_\alpha p_{\theta_\alpha(A)} s_\alpha^* = \sum_{\alpha \in \Delta_A} (s_\alpha p_{\theta_\alpha(A)}) (s_\alpha p_{\theta_\alpha(A)})^*$ for all $A \in \mathcal{B}_{reg}$.

Then the universal property of $C^*(\mathcal{B}, \mathcal{L}, \theta)$ gives a $*$ -homomorphism

$$\phi : C^*(\mathcal{B}, \mathcal{L}, \theta) \rightarrow C^*(s_\alpha, p_A)$$

defined by

$$\phi(p_A) = p_A \text{ and } \phi(s_{\alpha,B}) = s_\alpha p_B$$

for all $A \in \mathcal{B}, \alpha \in \mathcal{L}$ and $B \in \mathcal{R}_\alpha$. Since $\theta_\alpha(\mathcal{D}_{\theta_\alpha}) = \mathcal{R}_{\theta_\alpha}$ for each $\alpha \in \mathcal{L}$, we have $\mathcal{R}_{\theta_\alpha} \in \mathcal{R}_\alpha$ for each $\alpha \in \mathcal{L}$. Thus it follows from the equation $s_\alpha = s_\alpha p_{\mathcal{R}_{\theta_\alpha}}$ that the family $\{p_A, s_\alpha p_B : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{R}_\alpha\}$ contains all generators of $C^*(s_\alpha, p_A)$.

Thus ϕ is surjective. Since $p_A \neq 0$ for all $A \neq \emptyset$ and $C^*(s_\alpha, p_A)$ admits a gauge action $\beta : \mathbb{T} \rightarrow \text{Aut}(C^*(s_\alpha, p_A))$ such that $\beta_z(p_A) = p_A$ and $\beta_z(s_\alpha p_B) = z s_\alpha p_B$ for every $A \in \mathcal{B}$, $\alpha \in \mathcal{L}$ and $B \in \mathcal{R}_\alpha$, the gauge-invariant uniqueness theorem 6.2 implies that ϕ is an isomorphism.

Example 4.2 (Weakly left-resolving normal labelled spaces). We refer the reader to [1, 2, 3, 7] for the basic definitions and terminology of labelled graphs and their C^* -algebras. Let (E, \mathcal{L}) be a labelled graph over \mathcal{A} consisting of a directed graph E and a labelling map $\mathcal{L} : E^1 \rightarrow \mathcal{A}$ which is assumed to be onto. We then consider a weakly left-resolving normal labelled space $(E, \mathcal{L}, \mathcal{B})$ and put $C^*(E, \mathcal{L}, \mathcal{B}) := C^*(p_A, s_\alpha)$. Then \mathcal{B} is a Boolean algebra and for each $\alpha \in \mathcal{A}$, the map $\theta_\alpha : \mathcal{B} \rightarrow \mathcal{B}$ defined by

$$\theta_\alpha(A) := r(A, \alpha) \quad (4)$$

is an action on \mathcal{B} (see [4, Example 11.1]). Then the triple $(\mathcal{B}, \mathcal{A}, \theta)$ is a Boolean dynamical system. It is clear that $\mathcal{R}_\alpha \subseteq \mathcal{I}_{r(\alpha)} (= \{A \in \mathcal{B} : A \subseteq r(\alpha)\})$ for each $\alpha \in \mathcal{L}$. We claim that

$$C^*(E, \mathcal{L}, \mathcal{B}) \cong C^*(\mathcal{B}, \mathcal{A}, \theta, \mathcal{I}_{r(\alpha)}).$$

It is straightforward to check that

$$\{p_A, s_\alpha p_B : A \in \mathcal{B}, \alpha \in \mathcal{A} \text{ and } B \in \mathcal{I}_{r(\alpha)}\}$$

is a Cuntz–Krieger representation of $(\mathcal{B}, \mathcal{A}, \theta, \mathcal{I}_{r(\alpha)})$. Then the universal property of $C^*(\mathcal{B}, \mathcal{A}, \theta, \mathcal{I}_{r(\alpha)})$ gives a $*$ -homomorphism

$$\phi : C^*(\mathcal{B}, \mathcal{A}, \theta, \mathcal{I}_{r(\alpha)}) \rightarrow C^*(E, \mathcal{L}, \mathcal{B})$$

defined by

$$\phi(p_A) = p_A \text{ and } \phi(s_\alpha p_B) = s_\alpha p_B$$

for all $A \in \mathcal{B}, \alpha \in \mathcal{A}$ and $B \in \mathcal{I}_{r(\alpha)}$. Since $s_\alpha = s_\alpha p_{r(\alpha)}$, the family $\{p_A, s_\alpha p_B : A \in \mathcal{B}, \alpha \in \mathcal{A} \text{ and } B \in \mathcal{I}_{r(\alpha)}\}$ generates $C^*(E, \mathcal{L}, \mathcal{B})$, and hence, the map ϕ is onto. Applying the gauge-invariant uniqueness theorem 6.2, we conclude that ϕ is an isomorphism.

Remark 4.3. We continue Example 4.2. We claim that

- (1) If, in addition, E has no source and $(E, \mathcal{L}, \mathcal{B})$ is receiver set-finite, then we have $C^*(E, \mathcal{L}, \mathcal{B}) \cong C^*(\mathcal{B}, \mathcal{A}, \theta)$.
- (2) In general, $C^*(E, \mathcal{L}, \mathcal{B})$ is not isomorphic to $C^*(\mathcal{B}, \mathcal{A}, \theta)$.
- (3) $C^*(E, \mathcal{L}, \mathcal{B})$ is Morita equivalent to $C^*(\mathcal{B}, \mathcal{A}, \theta)$.

Proof. (1): For each $\alpha \in \mathcal{A}$, put $\mathcal{R}_\alpha = \{B \in \mathcal{B} : B \subseteq r(A, \alpha) \text{ for some } A \in \mathcal{B}\}$. Then

$$\{p_A, s_\alpha p_B : A \in \mathcal{B}, \alpha \in \mathcal{A} \text{ and } B \in \mathcal{R}_\alpha\}$$

is a Cuntz–Krieger representation of $(\mathcal{B}, \mathcal{A}, \theta)$. Clearly, p_A 's satisfy Definition 3.3

(i). We show that the family satisfies Definition 3.3(ii)–(iv):

- (ii) $p_A(s_\alpha p_B) = s_\alpha p_{r(A, \alpha)} p_B = (s_\alpha p_B) p_{r(A, \alpha)}$,
- (iii) $(s_\alpha p_B)^*(s_{\alpha'} p_{B'}) = p_B s_\alpha^* s_{\alpha'} p_{B'} = \delta_{\alpha, \alpha'} p_B p_{r(\alpha)} p_{B'} = \delta_{\alpha, \alpha'} p_{B \cap B'}$,
- (iv) $p_A = \sum_{\alpha \in \mathcal{L}(AE^1)} s_\alpha p_{r(A, \alpha)} s_\alpha^* = \sum_{\alpha \in \Delta_A} (s_\alpha p_{r(A, \alpha)})(s_\alpha p_{r(A, \alpha)})^*$ for all $A \in \mathcal{B}_{reg}$.

Then the universal property of $C^*(\mathcal{B}, \mathcal{A}, \theta)$ gives a $*$ -homomorphism

$$\phi : C^*(\mathcal{B}, \mathcal{A}, \theta) \rightarrow C^*(p_A, s_\alpha p_B)$$

defined by

$$\phi(p_A) = p_A \text{ and } \phi(s_{\alpha,B}) = s_\alpha p_B$$

for all $A \in \mathcal{B}, \alpha \in \mathcal{A}$ and $B \in \mathcal{R}_\alpha$. Since $p_A \neq 0$ for all $A \neq \emptyset$ and $C^*(p_A, s_\alpha p_B)$ admits a gauge action $\beta : \mathbb{T} \rightarrow \text{Aut}(C^*(E, \mathcal{L}, \mathcal{B}))$ such that $\beta_z(p_A) = p_A$ and $\beta_z(s_\alpha p_B) = z s_\alpha p_B$ for every $A \in \mathcal{B}, \alpha \in \mathcal{A}$ and $B \in \mathcal{R}_\alpha$, the gauge-invariant uniqueness theorems 6.2 implies that ϕ is an isomorphism.

If E has no sources and $(E, \mathcal{L}, \mathcal{B})$ is reciever set-finite, then we have


$$r(\alpha) = r(s(\alpha), \alpha) = r(\cup_{i=1}^m [w_i]_1, \alpha) = \cup_{i=1}^m r([w_i]_1, \alpha)$$

for some $w_i \in s(\alpha)$ ($i = 1, \dots, m$). The union is disjoint since $(E, \mathcal{L}, \mathcal{B})$ is weakly left-resolving. We then see that


$$s_\alpha = s_\alpha p_{r(\alpha)} = \sum_{i=1}^m s_\alpha p_{r([w_i]_1, \alpha)}$$

for some $r([w_i]_1, \alpha) \in \mathcal{R}_\alpha$. Thus, the family $\{p_A, s_\alpha p_B : A \in \mathcal{B}, \alpha \in \mathcal{A} \text{ and } B \in \mathcal{R}_\alpha\}$ generates $C^*(E, \mathcal{L}, \mathcal{B})$, and hence, the map ϕ becomes a $*$ -homomorphism from $C^*(\mathcal{B}, \mathcal{A}, \theta)$ onto $C^*(E, \mathcal{L}, \mathcal{B})$. The gauge-invariant uniqueness theorems 6.2 implies that ϕ is an isomorphism.

(2): Let $E^0 := \mathbb{N} \times \{1, 2\}$, $E^1 := \mathbb{N}$ and define $r, s : E^1 \rightarrow E^0$ by $s(e) = (e, 1)$, $r(e) = (e, 2)$. Then $E := (E^0, E^1, r, s)$ is the following directed graph.



Put $\mathcal{A} := \{a\}$ and define $\mathcal{L} : E^1 \rightarrow \mathcal{A}$ by $\mathcal{L}(e) = a$. Then we have the following labelled graph (E, \mathcal{L}) .



For $A, B \subseteq \mathbb{N}$, we write (A, B) for the subset $A \times \{1\} \cup B \times \{2\}$ of E^0 . Let

$\mathcal{B} := \{(A, B) : A \text{ is a finite subset of } \mathbb{N}, B \text{ is a subset of } \mathbb{N} \text{ such that } B \text{ or } \mathbb{N} \setminus B \text{ is finite}\}$.

Then $(E, \mathcal{L}, \mathcal{B})$ is a weakly left-resolving normal labelled space. Define $\theta_a : \mathcal{B} \rightarrow \mathcal{B}$ by $\theta_a((A, B)) := r((A, B), a) = (\emptyset, A)$. Then $(\mathcal{B}, \mathcal{A}, \theta)$ is a Boolean dynamical system such that $\mathcal{R}_a = \{(\emptyset, A) : A \text{ is a finite subset of } \mathbb{N}\}$.

Let $\widehat{\mathbb{N}}$ be the one-point compactification of \mathbb{N} , and let A be the C^* -algebra of 2×2 matrices over $C(\widehat{\mathbb{N}})$. For $(A, B) \in \mathcal{B}$, let

$$p_{(A,B)} = \begin{pmatrix} 1_A & 0 \\ 0 & 1_B \end{pmatrix}.$$

For a finite subset C of \mathbb{N} , let

$$s_{a,(\emptyset,C)} = \begin{pmatrix} 0 & 1_C \\ 0 & 0 \end{pmatrix},$$

and let

$$s_a = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Then it is straightforward to check that

$$\{p_{(A,B)} : (A, B) \in \mathcal{B}\} \cup \{s_a\}$$

is a representation of $(E, \mathcal{L}, \mathcal{B})$ and

$$\{p_{(A,B)} : (A, B) \in \mathcal{B}\} \cup \{s_{a,(\emptyset,C)} : C \text{ is a finite subset of } \mathbb{N}\}$$

is a Cuntz-Krieger representation of $(\mathcal{B}, \mathcal{A}, \theta)$. The C^* -algebra generated by

$$\{p_{(A,B)} : (A, B) \in \mathcal{B}\} \cup \{s_a\}$$

is A , while the C^* -algebra generated by

$$\{p_{(A,B)} : (A, B) \in \mathcal{B}\} \cup \{s_{a,(\emptyset,C)} : C \text{ is a finite subset of } \mathbb{N}\}$$

is

$$\left\{ \begin{pmatrix} f_{11} & f_{12} \\ f_{21} & f_{22} \end{pmatrix} \in A : f_{11}, f_{12}, f_{21} \in C_0(\mathbb{N}) \right\}$$

(where we consider $C_0(\mathbb{N})$ to be an ideal of $C(\widehat{\mathbb{N}})$). This shows that $C^*(\mathcal{B}, \mathcal{A}, \theta)$ is not isomorphic to $C^*(E, \mathcal{L}, \mathcal{B})$.

(3): By Example 4.2 and Proposition 6.3, we deduce that $C^*(E, \mathcal{L}, \mathcal{B})$ is Morita equivalent to $C^*(\mathcal{B}, \mathcal{A}, \theta)$. \square

5. C^* -ALGEBRAS OF RELATIVE GENERALIZED BOOLEAN DYNAMICAL SYSTEMS ARE RELATIVE CUNTZ-PIMSNER ALGEBRAS

In this section, we shall prove that for each relative generalized Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$, there exists a unique universal $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation by realizing $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ as a relative Cuntz-Pimsner algebra. We do that by closely imitating the approach used in [1, Section 3].

We first construct a C^* -correspondence of a generalized Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$. Most of the results can be obtained by arguments similar to arguments used in [1]. For completeness, we provide detailed proof for the results which we need to modify compared to [1].

Let $(\mathcal{B}, \mathcal{L}, \theta)$ be a Boolean dynamical system. For each $A \in \mathcal{B}$, we define

$$\chi_A := 1_{Z(A)} \in C_0(\widehat{\mathcal{B}}).$$

Let $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ denote the C^* -subalgebra of $C_0(\widehat{\mathcal{B}})$ generated by $\{\chi_A : A \in \mathcal{B}\}$. We then have that

$$\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta) = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}\}.$$

The following lemma will be frequently used throughout this section. The lemma can be proved in a way similar to have [1, Lemmas 3.2 and 3.3] is proved. We therefore omit the proof.

Lemma 5.1. *Let $(\mathcal{B}, \mathcal{L}, \theta)$ be a Boolean dynamical system.*

- (1) *Let $\{p_A : A \in \mathcal{B}\}$ be a family of projections in a C^* -algebra \mathcal{X} such that $p_{A \cap B} = p_A p_B$, $p_{A \cup B} = p_A + p_B - p_{A \cap B}$ for all $A, B \in \mathcal{B}$ and $p_\emptyset = 0$. Then there is a unique $*$ -homomorphism $\phi : \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta) \rightarrow \mathcal{X}$ such that $\phi(\chi_A) = p_A$ for all $A \in \mathcal{B}$.*
- (2) *If \mathcal{I} is an ideal of $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$, then we have $\mathcal{I} = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}, \chi_A \in \mathcal{I}\}$.*

Let $\{\mathcal{I}_\alpha : \alpha \in \mathcal{L}\}$ be a family of ideals of \mathcal{B} such that $\mathcal{R}_\alpha \subseteq \mathcal{I}_\alpha$ for all $\alpha \in \mathcal{L}$. For each $\alpha \in \mathcal{L}$, we let X_α denote the ideal of $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ generated by $\{\chi_A : A \in \mathcal{I}_\alpha\}$, that is,

$$X_\alpha = \overline{\text{span}}\{\chi_A : A \in \mathcal{I}_\alpha\}.$$

It is straightforward to see that X_α is a right Hilbert $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ -module with right action given by the usual multiplication in $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ and inner product defined by $\langle x, y \rangle = x^* y$ for $x, y \in X_\alpha$.

We define

$$\begin{aligned} X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha) &:= \bigoplus_{\alpha \in \mathcal{L}} X_\alpha \\ &= \left\{ (x_\alpha)_{\alpha \in \mathcal{L}} \in \prod_{\alpha \in \mathcal{L}} X_\alpha : \sum_{\alpha \in \mathcal{L}} x_\alpha^* x_\alpha \text{ converges in } \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta) \right\} \end{aligned}$$

to be the right Hilbert $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ -module with inner product defined by

$$\langle (x_\alpha)_{\alpha \in \mathcal{L}}, (y_\alpha)_{\alpha \in \mathcal{L}} \rangle := \sum_{\alpha \in \mathcal{L}} \langle x_\alpha, y_\alpha \rangle = \sum_{\alpha \in \mathcal{L}} x_\alpha^* y_\alpha$$

and the right action given by $(x_\alpha)_{\alpha \in \mathcal{L}} \cdot f := (x_\alpha f)_{\alpha \in \mathcal{L}}$ for $(x_\alpha)_{\alpha \in \mathcal{L}}, (y_\alpha)_{\alpha \in \mathcal{L}} \in X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ and $f \in \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$. It is easy to see that

$$\begin{aligned} X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha) &= \overline{\text{span}}\{e_{\alpha, A} : \alpha \in \mathcal{L} \text{ and } A \in \mathcal{I}_\alpha\} \\ &= \overline{\text{span}}\{e_{\alpha, A} : \alpha \in \mathcal{L} \text{ and } A \in F \text{ for some } F \in \mathcal{F}\}, \end{aligned}$$

where $e_{\alpha, A} := (\delta_{\alpha, \beta} \chi_A)_{\beta \in \mathcal{L}} \in \bigoplus_{\alpha \in \mathcal{L}} X_\alpha$ and \mathcal{F} is the collection of finite subsets F of mutually disjoint elements in \mathcal{I}_α .

We then have a unique $*$ -homomorphism $\phi_\alpha : \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta) \rightarrow X_\alpha$ defined by

$$\phi_\alpha(\chi_A) = \chi_{\theta_\alpha(A)}$$

for every $A \in \mathcal{B}$ and for each $\alpha \in \mathcal{L}$ (see [1, Lemma 3.4]). If we define a map

$$\phi : \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta) \rightarrow \mathcal{L}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))$$

by

$$\phi(f)((x_\alpha)_{\alpha \in \mathcal{L}}) = (\phi_\alpha(f)x_\alpha)_{\alpha \in \mathcal{L}}$$

for $f \in \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ and $(x_\alpha) \in X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$, where $\mathfrak{L}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))$ denotes the C^* -algebra of adjointable operators on $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$, then the map ϕ is a $*$ -homomorphism ([1, Lemma 3.6]). Thus, $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ is a C^* -correspondence over $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$.

5.1. Relative Cuntz-Pimsner algebras of generalized Boolean dynamical systems. Let $\mathfrak{K}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))$ denote the ideal of $\mathfrak{L}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))$ consisting of generalized compact operators, that is,

$$\mathfrak{K}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)) = \overline{\text{span}}\{\Theta_{x,y} \in \mathfrak{L}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)) : x, y \in X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)\}, \quad (5)$$

where $\Theta_{x,y}$ is the operator that maps z to $x\langle y, z \rangle$. As in [5, Definition 1.1], we define an ideal $J(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))$ of $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ by

$$J(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)) := \phi^{-1}(\mathfrak{K}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))).$$

We also define an ideal $J_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)}$ of $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ by

$$\begin{aligned} J_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)} &:= J(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)) \cap \ker(\phi)^\perp \\ &= \phi^{-1}(\mathfrak{K}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))) \cap \ker(\phi)^\perp \end{aligned}$$

([8, Definition 3.2]).

We first characterize the ideals $J(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))$ and $J_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)}$ as follows.

Lemma 5.2. *Let $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ be the C^* -correspondence over $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$.*

- (1) $J(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)) = \overline{\text{span}}\{\chi_A : A \in \mathcal{B} \text{ such that } \lambda_A < \infty\}$.
- (2) $J_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)} = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}_{\text{reg}}\}$.

Proof. (1): By Lemma 5.1(2), we only need to show that for $A \in \mathcal{B}$,

$$\phi(\chi_A) \in \mathfrak{K}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)) \iff \lambda_A < \infty.$$

For the "if" direction, choose $A \in \mathcal{B}$ such that $\lambda_A < \infty$. We then have

$$\phi(\chi_A) = \sum_{\alpha \in \Delta_A} \Theta_{e_{\alpha, \theta_\alpha(A)}, e_{\alpha, \theta_\alpha(A)}} \in \mathfrak{K}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)).$$

For the "only if" direction, we first claim that for each $\eta \in \mathfrak{K}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))$, the set $\{\alpha \in \mathcal{L} : \|\langle e_{\alpha, \theta_\alpha(A)}, \eta(e_{\alpha, \theta_\alpha(A)}) \rangle\| \geq 1\}$ is finite. By (5), it is enough to check that for $\eta = \theta_{x,y}$. For each $\alpha \in \mathcal{L}$ and $x = (x_\alpha)$, $y = (y_\alpha) \in X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$, it is straightforward to check that $\langle e_{\alpha, \theta_\alpha(A)}, \theta_{x,y}(e_{\alpha, \theta_\alpha(A)}) \rangle = x_\alpha y_\alpha^* \chi_{\theta_\alpha(A)}$. Then our claim follows since $\|x_\alpha y_\alpha^*\| \geq 1$ for only a finitely many α 's.

Now choose $A \in \mathcal{B}$ so that $\phi(\chi_A) \in \mathfrak{K}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))$. Then the set $\Delta_A = \{\alpha \in \mathcal{L} : \theta_\alpha(A) \neq \emptyset\}$ must be finite because $\|\langle e_{\alpha, \theta_\alpha(A)}, \phi(\chi_A)(e_{\alpha, \theta_\alpha(A)}) \rangle\| = \|\chi_{\theta_\alpha(A)}\| = 1$.

(2): It is enough to prove that for $A \in \mathcal{B}$, we have $\chi_A \in J_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)} \iff A \in \mathcal{B}_{\text{reg}}$. We first claim that

$$\ker(\phi)^\perp = \overline{\text{span}}\{\chi_A : A \in \mathcal{B} \text{ such that } 0 < \lambda_B \text{ for any } \emptyset \neq B \in \mathcal{I}_A\}. \quad (6)$$

Since $\ker(\phi) = \bigcap_{\alpha \in \mathcal{L}} \ker(\phi_\alpha)$, we see that

$$\begin{aligned} \ker(\phi) &= \overline{\text{span}}\{\chi_B : B \in \mathcal{B} \text{ with } \theta_\alpha(B) = \emptyset \text{ for all } \alpha \in \mathcal{L}\} \\ &= \overline{\text{span}}\{\chi_B : B \in \mathcal{B} \text{ with } \Delta_B = \emptyset\}. \end{aligned}$$

It then follows that

$$\ker(\phi)^\perp = \overline{\text{span}}\{\chi_A : A \cap B = \emptyset \text{ for all } B \in \mathcal{B} \text{ with } \Delta_B = \emptyset\}.$$

We prove (6) by showing that

$$\begin{aligned} \{A \in \mathcal{B} : A \cap B = \emptyset \text{ for all } B \in \mathcal{B} \text{ with } \Delta_B = \emptyset\} \\ = \{A \in \mathcal{B} : 0 < \lambda_B \text{ for any } \emptyset \neq B \in \mathcal{I}_A\}. \end{aligned}$$

For the inclusion (\subseteq), we choose $\emptyset \neq B \in \mathcal{I}_A$. Then $\emptyset \neq B = A \cap B$, so $\Delta_B \neq \emptyset$, which means that $\lambda_B > 0$. For the reverse inclusion (\supseteq), choose $B \in \mathcal{B}$ with $\Delta_B = \emptyset$. Since $\theta_\alpha(A \cap B) \subseteq \theta_\alpha(B) = \emptyset$ for all $\alpha \in \mathcal{L}$, we have $\Delta_{A \cap B} = \emptyset$, meaning $\lambda_{A \cap B} = 0$. Thus $A \cap B = \emptyset$.

Since for $A \in \mathcal{B}$, it is obvious that $\lambda_A < \infty \iff \lambda_B < \infty$ for all $\emptyset \neq B \in \mathcal{I}_A$. Then using the result (1) and the equation (6), we have our result. \square

Recall that a (*Toeplitz*) *representation* ([6] or [8, Definition 2.1]) of the C^* -correspondence $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ over $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ on a C^* -algebra \mathcal{X} is a pair (π, t) where $\pi : \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta) \rightarrow \mathcal{X}$ is a $*$ -homomorphism and $t : X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha) \rightarrow \mathcal{X}$ is a linear map satisfying

- (1) $t(x)^*t(y) = \pi(\langle x, y \rangle)$ for $x, y \in X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$,
- (2) $\pi(f)t(x) = t(\phi(f)x)$ for $f \in \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ and $x \in X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$.

Given a representation (π, t) , there is a $*$ -homomorphism $\psi_t : \mathfrak{K}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)) \rightarrow \mathcal{X}$ such that

$$\psi_t(\Theta_{x,y}) = t(x)t(y)^*$$

for $x, y \in X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$. Let K be an ideal in $J(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))$. A representation (π, t) is called *K-coisometric* if it satisfies that

$$\pi(f) = \psi_t(\phi(f))$$

for all $f \in K$ ([11, Theorem 2.19] or [5, Definition 1.1]). The *relative Cuntz-Pimsner algebra* $\mathcal{O}_{(K, X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))}$ determined by the ideal K is the C^* -algebra generated by a universal *K-coisometric* representation of $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$.

Remark 5.3. When K is the zero ideal of $J(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))$, the algebra $\mathcal{O}_{(K, X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))}$ is the Toeplitz algebra $\mathcal{T}_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)}$.

Remark 5.4. For an ideal \mathcal{J} of \mathcal{B} , we let $K_{\mathcal{J}}$ denote the ideal of $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ generated by $\{\chi_A : A \in \mathcal{J}\}$, that is,

$$K_{\mathcal{J}} := \overline{\text{span}}\{\chi_A : A \in \mathcal{J}\}.$$

Then the map $\mathcal{J} \mapsto K_{\mathcal{J}}$ is a bijective map between the set of all ideals of \mathcal{B} and the set of all ideals of $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$: For an ideal K of $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$, we define $\mathcal{J}_K := \{A \in \mathcal{B} : \chi_A \in K\}$. Then it is easy to check that \mathcal{J}_K is an ideal of \mathcal{B} and $K_{\mathcal{J}_K} = \overline{\text{span}}\{\chi_A : A \in \mathcal{J}_K\} = K$ since $K = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}, \chi_A \in K\}$ by Lemma 5.1(2). Thus the map is onto. The injectivity of the map is rather obvious since we have $\mathcal{J}_{K_{\mathcal{J}}} = \mathcal{J}$. Note that $K \mapsto \mathcal{J}_K$ is the inverse map of $\mathcal{J} \mapsto K_{\mathcal{J}}$.

The following theorem asserts that for a relative generalized Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$, there exists a universal $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation.

Theorem 5.5. *Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ be a relative generalized Boolean dynamical system. Then there is a one-to-one correspondence between $K_{\mathcal{J}}$ -coisometric representations of $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ and $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representations that takes a $K_{\mathcal{J}}$ -coisometric representation (π, t) of $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ to the $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation $\{\pi(\chi_A), t(e_{\alpha, B}) : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_\alpha\}$. Thus, we have $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J}) \cong \mathcal{O}_{(K_{\mathcal{J}}, X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha))}$.*

Proof. Let (π, t) be a $K_{\mathcal{J}}$ -coisometric representation of $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$. We show that

$$\{\pi(\chi_A), t(e_{\alpha, B}) : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_\alpha\}$$

is a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation: It is straightforward to check that $\pi(\chi_A)$'s satisfy Definition 3.3(i). For any $\alpha, \alpha' \in \mathcal{L}$, $B \in \mathcal{I}_\alpha$ and $B' \in \mathcal{I}_{\alpha'}$, we observe for Definition 3.3(ii) and (iii) that

$$\pi(\chi_A)t(e_{\alpha, B}) = t(\phi(\chi_A)e_{\alpha, B}) = t(e_{\alpha, B}\chi_{\theta_\alpha(A)}) = t(e_{\alpha, B})\pi(\chi_{\theta_\alpha(A)})$$

and that

$$t(e_{\alpha, B})^*t(e_{\alpha', B'}) = \pi(\langle e_{\alpha, B}, e_{\alpha', B'} \rangle) = \delta_{\alpha, \alpha'}\pi(\chi_{B \cap B'}).$$

For Definition 3.3(iv), choose $A \in \mathcal{J}$. Then $\chi_A \in K_{\mathcal{J}}$. Hence, it follows that

$$\begin{aligned} \pi(\chi_A) &= \psi_t(\phi(\chi_A)) \\ &= \psi_t\left(\sum_{\alpha \in \Delta_A} \Theta_{e_{\alpha, \theta_\alpha(A)}, e_{\alpha, \theta_\alpha(A)}}\right) \\ &= \sum_{\alpha \in \Delta_A} t(e_{\alpha, \theta_\alpha(A)})t(e_{\alpha, \theta_\alpha(A)})^*. \end{aligned}$$

Thus, $\{\pi(\chi_A), t(e_{\alpha, B})\}$ is a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation.

For the opposite direction, suppose that $\{P_A, S_{\alpha, B} : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_\alpha\}$ is a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation in a C^* -algebra \mathcal{X} . Since we have $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta) = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}\}$ and

$$X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha) = \overline{\text{span}}\{e_{\alpha, A} : \alpha \in \mathcal{L} \text{ and } A \in F \text{ for some } F \in \mathcal{F}\}, \quad (7)$$

where \mathcal{F} is the collection of finite subsets F of mutually disjoint elements in \mathcal{I}_α , there can be at most one representation (π, t) of $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ such that $\pi(\chi_A) = P_A$ and $t(e_{\alpha, B}) = S_{\alpha, B}$ for $A \in \mathcal{B}$, $\alpha \in \mathcal{L}$ and $B \in \mathcal{I}_\alpha$. We shall construct such a representation (π, t) .

As [1, Lemma 3.2], we can see that there is a unique $*$ -homomorphism

$$\pi : \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta) \rightarrow \mathcal{X} \text{ given by } \pi(\chi_A) = P_A$$

for $A \in \mathcal{B}$. We construct a linear map $t : X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha) \rightarrow \mathcal{X}$. Given a finite subset F in \mathcal{F} of mutually disjoint elements in \mathcal{I}_α and a finite subset F' of \mathcal{L} , we define

$$t\left(\sum_{\alpha \in F', B \in F} c_{\alpha, B} e_{\alpha, B}\right) := \sum_{\alpha \in F', B \in F} c_{\alpha, B} S_{\alpha, B},$$

where $c_{\alpha,B} \in \mathbb{C}$. Then we see that

$$\begin{aligned}
& t\left(\sum_{\alpha \in F', B \in F} c_{\alpha,B} e_{\alpha,B}\right)^* t\left(\sum_{\alpha' \in F', B' \in F} d_{\alpha',B'} e_{\alpha',B'}\right) \\
&= \left(\sum_{\alpha \in F', B \in F} c_{\alpha,B} S_{\alpha,B}\right)^* \left(\sum_{\alpha' \in F', B' \in F} d_{\alpha',B'} S_{\alpha',B'}\right) \\
&= \sum_{\alpha \in F' \text{ and } B, B' \in F} \overline{c_{\alpha,B}} d_{\alpha,B'} S_{\alpha,B}^* S_{\alpha,B'} \\
&= \sum_{\alpha \in F' \text{ and } B \in F} \overline{c_{\alpha,B}} d_{\alpha,B} P_B \\
&= \sum_{\alpha \in F' \text{ and } B, B' \in F} \overline{c_{\alpha,B}} d_{\alpha,B'} \pi(\chi_{B \cap B'}) \\
&= \sum_{\alpha \in F' \text{ and } B, B' \in F} \overline{c_{\alpha,B}} d_{\alpha,B'} \pi(\chi_B) \pi(\chi_{B'}) \\
&= \pi\left(\left\langle \sum_{\alpha \in F', B \in F} c_{\alpha,B} e_{\alpha,B}, \sum_{\alpha' \in F', B' \in F} d_{\alpha',B'} e_{\alpha',B'} \right\rangle\right).
\end{aligned}$$

Thus, t extends to a linear map from $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ to \mathcal{X} which satisfies

$$t(x)^* t(y) = \pi(\langle x, y \rangle)$$

for all $x, y \in X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$. We next show that we have

$$\pi(f) t(x) = t(\phi(f)x)$$

for $f \in \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ and $x \in X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$. For $A \in \mathcal{B}$, $\alpha \in \mathcal{L}$ and $B \in \mathcal{I}_\alpha$, we have

$$\pi(\chi_A) S_{\alpha,B} = P_A S_{\alpha,B} = S_{\alpha,B} P_{\theta_\alpha(A)} = S_{\alpha,B} \pi(\phi_\alpha(\chi_A)).$$

Since $\mathcal{A}(\mathcal{B}, \mathcal{L}, \theta) = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}\}$, we have $\pi(f) S_{\alpha,B} = S_{\alpha,B} \pi(\phi_\alpha(f))$ for any $f \in \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$ and $\alpha \in \mathcal{L}$. Thus we see that

$$\begin{aligned}
\pi(f) t(e_{\alpha,B}) &= \pi(f) S_{\alpha,B} = S_{\alpha,B} \pi(\phi_\alpha(f)) \\
&= t(e_{\alpha,B}) \pi(\phi_\alpha(f)) = t(\phi(f) e_{\alpha,B})
\end{aligned}$$

for all $f \in \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta)$, $\alpha \in \mathcal{L}$ and $B \in \mathcal{I}_\alpha$. Our claim then follows from (7).

If $A \in \mathcal{J}$, then

$$\begin{aligned}
\pi(\chi_A) &= P_A = \sum_{\alpha \in \Delta_A} S_{\alpha, \theta_\alpha(A)} S_{\alpha, \theta_\alpha(A)}^* \\
&= \sum_{\alpha \in \Delta_A} t(e_{\alpha, \theta_\alpha(A)}) t(e_{\alpha, \theta_\alpha(A)})^* \\
&= \sum_{\alpha \in \Delta_A} \psi_t(\Theta_{e_{\alpha, \theta_\alpha(A)}, e_{\alpha, \theta_\alpha(A)}}) \\
&= \psi_t\left(\sum_{\alpha \in \Delta_A} \Theta_{e_{\alpha, \theta_\alpha(A)}, e_{\alpha, \theta_\alpha(A)}}\right) \\
&= \psi_t(\phi(\chi_A)).
\end{aligned}$$

Hence, $\pi(f) = \psi_t(\phi(f))$ for all $f \in K_{\mathcal{J}}$. We have proved that (π, t) is a $K_{\mathcal{J}}$ -coisometric representation of $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha})$.

Therefore, we have $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J}) \cong \mathcal{O}_{(K_{\mathcal{J}}, X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}))}$ by the universal nature of each algebra. \square

Corollary 5.6. *Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha})$ be a generalized Boolean dynamical system. Then we have*

- (1) $\mathcal{T}(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}) \cong \mathcal{T}_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha})}$,
- (2) $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}) \cong \mathcal{O}_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha})}$.

Proof. (1): Taking $\mathcal{J} = \emptyset$ in Theorem 5.5, we have

$$C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \emptyset) \cong \mathcal{O}_{(K_{\emptyset}, X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}))} = \mathcal{O}_{(\{0\}, X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}))} = \mathcal{T}_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha})}.$$

(2): If $\mathcal{J} = \mathcal{B}_{reg}$, then $K_{\mathcal{B}_{reg}} = \overline{\text{span}}\{\chi_A : A \in \mathcal{B}_{reg}\} = J_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha})}$ by Lemma 5.2(2). Thus it follows that

$$C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{B}_{reg}) \cong \mathcal{O}_{(K_{\mathcal{B}_{reg}}, X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}))} = \mathcal{O}_{X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha})}.$$

\square

Corollary 5.7. *Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J})$ be a relative generalized Boolean dynamical system. Let $\{p_A, s_{\alpha, B} : A \in \mathcal{B}, \alpha \in \mathcal{L}, B \in \mathcal{I}_{\alpha}\}$ be the universal $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J})$ -representation. Then $p_A \neq 0$ for $A \in \mathcal{B} \setminus \{\emptyset\}$, and $p_A = \sum_{\alpha \in \Delta_A} s_{\alpha, \theta_{\alpha}(A)} s_{\alpha, \theta_{\alpha}(A)}^*$ if and only if $A \in \mathcal{J}$.*

Proof. This follows from Theorem 5.5 and [9, Corollary 11.8]. \square

6. THE GAUGE-INVARIANT UNIQUENESS THEOREM

In this section we shall prove a gauge-invariant uniqueness theorem of C^* -algebras of relative generalized Boolean dynamical systems. We also show that the class of C^* -algebras of relative generalized Boolean dynamical systems is the same as the class of C^* -algebras of generalized Boolean dynamical systems, and that the C^* -algebra of any relative generalized Boolean dynamical system is Morita equivalent to the C^* -algebra of a Boolean dynamical system.

Theorem 6.1 (Gauge-Invariant Uniqueness for $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J})$). *Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J})$ be a relative generalized Boolean dynamical system and let $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J}) = C^*(p_A, s_{\alpha, B})$. Suppose that $\{P_A, S_{\alpha, B} : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_{\alpha}\}$ is a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J})$ -representation in a C^* -algebra \mathcal{X} and let $\pi : C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J}) \rightarrow \mathcal{X}$ be the unique $*$ -homomorphism such that $\pi(p_A) = P_A$ and $\pi(s_{\alpha, B}) = S_{\alpha, B}$ for all $A \in \mathcal{B}$, $\alpha \in \mathcal{L}$ and $B \in \mathcal{I}_{\alpha}$. Then π is injective if and only if the following properties hold:*

- (1) $P_A \neq 0$ whenever $A \neq \emptyset$,
- (2) $P_A - \sum_{\alpha \in \Delta_A} S_{\alpha, \theta_{\alpha}(A)} S_{\alpha, \theta_{\alpha}(A)}^* \neq 0$ for all $A \in \mathcal{B}_{reg} \setminus \mathcal{J}$,
- (3) there exists for each $z \in \mathbb{T}$ a $*$ -homomorphism $\beta_z : C^*(P_A, S_{\alpha, B}) \rightarrow C^*(P_A, S_{\alpha, B})$ such that $\beta_z(P_A) = P_A$ and $\beta_z(S_{\alpha, B}) = z S_{\alpha, B}$ for $A \in \mathcal{B}$, $\alpha \in \mathcal{L}$ and $B \in \mathcal{I}_{\alpha}$.

Proof. If $\{P_A, S_{\alpha, B} : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_{\alpha}\}$ is a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha}; \mathcal{J})$ -representation in a C^* -algebra \mathcal{X} , then by Theorem 5.5, we have a $K_{\mathcal{J}}$ -coisometric representation (π', t') of $X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_{\alpha})$. The conditions (1)-(3) are equivalent to (1)'-(3)' stated below, respectively:

- (1)' $\ker \pi' = 0$,
- (2)' $\{a \in \mathcal{A}(\mathcal{B}, \mathcal{L}, \theta) : \phi(a) \in \mathcal{K}(X(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)) \text{ and } \pi'(a) = \psi_{t'}(\phi(a))\} = K_{\mathcal{J}}$,
- (3)' (π', t') admits a gauge action.

Thus, the result follows from [9, Corollary 11.8]. \square

As a corollary, we obtain a version of the gauge-invariant uniqueness theorem for C^* -algebras of generalized Boolean dynamical systems. It is a generalization of [1, Corollary 3.10], [4, Theorem 5.10], and [7, Theorem 4.2].

Corollary 6.2 (Gauge-Invariant Uniqueness for $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$). *Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ be a generalized Boolean dynamical system, and let $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha) := C^*(p_A, s_{\alpha, B})$. Suppose that $\{P_A, S_{\alpha, B} : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_\alpha\}$ is a Cuntz–Krieger representation of $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$ in a C^* -algebra \mathcal{X} and let $\pi : C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha) \rightarrow \mathcal{X}$ be the unique $*$ -homomorphism such that $\pi(p_A) = P_A$ and $\pi(s_{\alpha, B}) = S_{\alpha, B}$ for all $A \in \mathcal{B}, \alpha \in \mathcal{L}$ and $B \in \mathcal{I}_\alpha$. Then π is injective if and only if the following properties hold:*

- (1) $P_A \neq 0$ whenever $A \neq \emptyset$,
- (2) *there exists for each $z \in \mathbb{T}$ a $*$ -homomorphism $\beta_z : C^*(P_A, S_{\alpha, B}) \rightarrow C^*(P_A, S_{\alpha, B})$ such that $\beta_z(P_A) = P_A$ and $\beta_z(S_{\alpha, B}) = zS_{\alpha, B}$ for $A \in \mathcal{B}, \alpha \in \mathcal{L}$ and $B \in \mathcal{I}_\alpha$.*

Proof. Taking $\mathcal{J} = \mathcal{B}_{reg}$ in Theorem 6.1, we have the result. \square

The following proposition has the consequence that the C^* -algebra of any relative generalized Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is Morita equivalent to $C^*(\mathcal{B}, \mathcal{L}, \theta; \mathcal{J}) (= C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{R}_\alpha; \mathcal{J}))$.

Proposition 6.3. *Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ be a relative generalized Boolean dynamical system. Then $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is (isomorphic to) a full hereditary C^* -subalgebra of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{B}; \mathcal{J})$.*

Proof. Let $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J}) := C^*(q_A, t_{\alpha, B})$ and $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{B}; \mathcal{J}) := C^*(p_A, s_{\alpha, B})$. Since $\{p_A, s_{\alpha, B} : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_\alpha\}$ is a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation, by the universal property of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ there exists a $*$ -homomorphism $\iota : C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J}) \rightarrow C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{B}; \mathcal{J})$ such that

$$\iota(q_A) = p_A \text{ and } \iota(t_{\alpha, B}) = s_{\alpha, B}$$

for $A \in \mathcal{B}, \alpha \in \mathcal{L}$ and $B \in \mathcal{I}_\alpha$. Since $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{B}; \mathcal{J})$ admits a gauge action, $\iota(q_A) = p_A \neq 0$ for $A \neq \emptyset$, and $p_A - \sum_{\alpha \in \Delta_A} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^* \neq 0$ for all $A \in \mathcal{B}_{reg} \setminus \mathcal{J}$ by Corollary 5.7, the homomorphism ι is injective by Theorem 6.1.

We identify $\iota(q_A)$ with p_A and $\iota(t_{\alpha, B})$ with $s_{\alpha, B}$ so that $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is a subalgebra of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{B}; \mathcal{J})$. We then have that $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is hereditary because $p_A s_{\alpha, B} p_B = s_{\alpha, B \cap \theta_\alpha(A)} \in C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ for any $A, B \in \mathcal{B}$ and any $\alpha \in \mathcal{L}$, and it is full because any ideal of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{B})$ that contains $\{p_A : A \in \mathcal{B}\}$ must be equal to $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{B}; \mathcal{J})$ since $s_{\alpha, B} p_B = s_{\alpha, B}$ for any $B \in \mathcal{B}$ and $\alpha \in \mathcal{L}$. \square

Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ be a relative generalized Boolean dynamical system. By imitating a construction in [9, Section 6], we shall now construct a generalized Boolean dynamical system $(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha)$ such that $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is isomorphic to $C^*(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha)$.

Given a relative generalized Boolean dynamical system $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$, we let

$$\tilde{\mathcal{B}} = \{(A, [B]_{\mathcal{J}}) : A, B \in \mathcal{B} \text{ and } [A]_{\mathcal{B}_{reg}} = [B]_{\mathcal{B}_{reg}}\}.$$

Then $\tilde{\mathcal{B}}$ is a Boolean algebra with empty set $(\emptyset, \emptyset) := (\emptyset, [\emptyset]_{\mathcal{J}})$ and operations defined by

$$\begin{aligned} (A_1, [B_1]_{\mathcal{J}}) \cup (A_2, [B_2]_{\mathcal{J}}) &:= (A_1 \cup A_2, [B_1 \cup B_2]_{\mathcal{J}}), \\ (A_1, [B_1]_{\mathcal{J}}) \cap (A_2, [B_2]_{\mathcal{J}}) &:= (A_1 \cap A_2, [B_1 \cap B_2]_{\mathcal{J}}), \\ (A_1, [B_1]_{\mathcal{J}}) \setminus (A_2, [B_2]_{\mathcal{J}}) &:= (A_1 \setminus A_2, [B_1 \setminus B_2]_{\mathcal{J}}). \end{aligned}$$

For $\alpha \in \mathcal{L}$, we define $\tilde{\theta}_\alpha : \tilde{\mathcal{B}} \rightarrow \tilde{\mathcal{B}}$ by

$$\tilde{\theta}_\alpha(A, [B]_{\mathcal{J}}) = (\theta_\alpha(A), [\theta_\alpha(A)]_{\mathcal{J}}).$$

Then $(\tilde{\mathcal{B}}, \mathcal{L}, \tilde{\theta})$ is a Boolean dynamical system. Since $\Delta_{(\emptyset, [B]_{\mathcal{J}})}^{(\tilde{\mathcal{B}}, \mathcal{L}, \tilde{\theta})} = \emptyset$ for any $B \in \mathcal{B}$, we have that $[B]_{\mathcal{J}} = \emptyset$ if $(A, [B]_{\mathcal{J}}) \in \tilde{\mathcal{B}}_{reg}^{(\tilde{\mathcal{B}}, \mathcal{L}, \tilde{\theta})}$. It follows that

$$\tilde{\mathcal{B}}_{reg} := \tilde{\mathcal{B}}_{reg}^{(\tilde{\mathcal{B}}, \mathcal{L}, \tilde{\theta})} = \{(A, \emptyset) : A \in \mathcal{B}_{reg}\}.$$

Proposition 6.4. *Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ be a relative generalized Boolean dynamical system. Then we have*

$$C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J}) \cong C^*(\tilde{\mathcal{B}}, \mathcal{L}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha),$$

where $\tilde{\mathcal{I}}_\alpha = \{(A, [A]_{\mathcal{J}}) : A \in \mathcal{I}_\alpha\}$ for $\alpha \in \mathcal{L}$.

Proof. Let $\{p_A, s_{\alpha, B} : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_\alpha\}$ be the universal $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation, and let $\{p_{(A, [B]_{\mathcal{J}})}, s_{\alpha, (A, [A]_{\mathcal{J}})} : (A, [B]_{\mathcal{J}}) \in \tilde{\mathcal{B}}, \alpha \in \mathcal{L} \text{ and } (A, [A]_{\mathcal{J}}) \in \tilde{\mathcal{I}}_\alpha\}$ be the universal Cuntz-Krieger representation of $(\tilde{\mathcal{B}}, \mathcal{L}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha)$. It is easy to see that $\{p_{(A, [A]_{\mathcal{J}})}, s_{\alpha, (B, [B]_{\mathcal{J}})} : A \in \mathcal{B}, \alpha \in \mathcal{L} \text{ and } B \in \mathcal{I}_\alpha\}$ is a $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ -representation. There is therefore a $*$ -homomorphism

$$\phi : C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J}) \rightarrow C^*(\tilde{\mathcal{B}}, \mathcal{L}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha)$$

such that

$$\phi(p_A) = p_{(A, [A]_{\mathcal{J}})} \text{ and } \phi(s_{\alpha, B}) = s_{\alpha, (B, [B]_{\mathcal{J}})}$$

for $A \in \mathcal{B}$, $\alpha \in \mathcal{L}$ and $B \in \mathcal{I}_\alpha$.

We shall construct an inverse to ϕ . For each $(A, [B]_{\mathcal{J}}) \in \tilde{\mathcal{B}}$, $\alpha \in \mathcal{L}$ and $(A, [A]_{\mathcal{J}}) \in \tilde{\mathcal{I}}_\alpha$, we define

$$P_{(A, [B]_{\mathcal{J}})} := p_A + p_C - \sum_{\alpha \in \Delta_C} s_{\alpha, \theta_\alpha(C)} s_{\alpha, \theta_\alpha(C)}^* - p_D + \sum_{\alpha \in \Delta_D} s_{\alpha, \theta_\alpha(D)} s_{\alpha, \theta_\alpha(D)}^*,$$

$$S_{\alpha, (A, [A]_{\mathcal{J}})} := s_{\alpha, A},$$

where $C, D \in \mathcal{B}_{reg}$ are such that $A \cup C = B \cup D$ and $A \cap C = B \cap D = \emptyset$.

We first prove that the projection $P_{(A, [B]_{\mathcal{J}})}$ does not depend on the choice of B , C , and D . Supposet that $[B]_{\mathcal{J}} = [B']_{\mathcal{J}}$, $A \cup C = B \cup D$ with $A \cap C = B \cap D = \emptyset$, and $A \cup C' = B' \cup D'$ with $A \cap C' = B' \cap D' = \emptyset$. To ease notion we let

$$q_A := p_A - \sum_{\alpha \in \Delta_A} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^*$$

when $A \in \mathcal{B}_{reg}$. Then $q_A = 0$ if $A \in \mathcal{J}$. In particular, if $A \subseteq B \setminus B'$ or $A \subseteq B' \setminus B$, then $q_A = 0$. Thus,

$$q_{C' \setminus (C \cup D')} = q_{C \cap C' \cap D' \setminus D} = q_{D' \setminus (C' \cup D)} = 0$$

since $C' \setminus (C \cup D') \subseteq B' \setminus B$, $C \cap C' \cap D' \setminus D \subseteq B \setminus B'$ and $D' \setminus (C' \cup D) \subseteq B' \setminus B$. We also have that $q_{D \cap C' \cap D' \setminus C} = 0$ because $D \cap C' \cap D' \setminus C \subseteq A \cap C' = \emptyset$. It then follows that

$$\begin{aligned} q_{C' \setminus C} &= q_{C' \setminus (C \cup D')} + q_{C' \cap D' \setminus C} = q_{C' \cap D' \setminus C} = q_{D \cap C' \cap D' \setminus C} + q_{C' \cap D' \setminus (C \cup D)} \\ &= q_{C' \cap D' \setminus (C \cup D)} = q_{C' \cap D' \setminus (C \cup D)} + q_{C \cap C' \cap D' \setminus D} = q_{C' \cap D' \setminus D} \\ &= q_{C' \cap D' \setminus D} + q_{D' \setminus (C' \cup D)} = q_{D' \setminus D}. \end{aligned}$$

Thus, $q_C + q_{D \cup D'} = q_C + q_D + q_{D' \setminus D} = q_C + q_D + q_{C' \setminus C} = q_{C \cup C'} + q_D$ from which we have that

$$p_A + q_C - q_D = p_A + q_{C \cup C'} - q_{D \cup D'} = p_A + q_{C'} - q_{D'}.$$

So, $P_{(A, [B]_{\mathcal{J}})} = P_{(A, [B']_{\mathcal{J}})}$.

The following calculations show that the family $\{P_{(A, [B]_{\mathcal{J}})}, S_{\alpha, (A, [A]_{\mathcal{J}})}\}$ is a Cuntz–Krieger representation of $(\tilde{\mathcal{B}}, \mathcal{L}, \tilde{\theta}, \tilde{\mathcal{I}}_{\alpha})$:

(i): Clearly, $P_{(\emptyset, \emptyset)} = 0$. Given $(A_1, [B_1]_{\mathcal{J}})$ and $(A_2, [B_2]_{\mathcal{J}})$ in $\tilde{\mathcal{B}}$, we can choose $C_1, D_1, C_2, D_2 \in \mathcal{B}_{reg}$ such that

$$\begin{aligned} A_1 \cup C_1 &= B_1 \cup D_1 \text{ with } A_1 \cap C_1 = B_1 \cap D_1 = \emptyset, \\ A_2 \cup C_2 &= B_2 \cup D_2 \text{ with } A_2 \cap C_2 = B_2 \cap D_2 = \emptyset, \end{aligned}$$

and

$$\begin{aligned} (A_1 \cap A_2) \cup \{(A_1 \cap C_2) \cup (D_1 \cap D_2) \cup (C_1 \cap A_2) \cup (C_1 \cap C_2)\} \\ = (B_1 \cap B_2) \cup \{(C_1 \cap D_2) \cup (D_1 \cap A_2) \cup (D_1 \cap C_2) \cup (A_1 \cap D_2)\}, \end{aligned}$$

with

$$\begin{aligned} (A_1 \cap A_2) \cap \{(A_1 \cap C_2) \cup (D_1 \cap D_2) \cup (C_1 \cap A_2) \cup (C_1 \cap C_2)\} &= \emptyset, \\ (B_1 \cap B_2) \cap \{(C_1 \cap D_2) \cup (D_1 \cap A_2) \cup (D_1 \cap C_2) \cup (A_1 \cap D_2)\} &= \emptyset. \end{aligned}$$

Using the fact that $p_A q_B = q_{A \cap B}$ for all $A \in \mathcal{B}$ and $B \in \mathcal{B}_{reg}$, it easily follows that

$$\begin{aligned} &P_{(A_1, [B_1]_{\mathcal{J}})} P_{(A_2, [B_2]_{\mathcal{J}})} \\ &= (p_{A_1} + q_{C_1} - q_{D_1})(p_{A_2} + q_{C_2} - q_{D_2}) \\ &= p_{A_1 \cap A_2} + q_{A_1 \cap C_2} + q_{D_1 \cap D_2} + q_{C_1 \cap A_2} + q_{C_1 \cap C_2} \\ &\quad - q_{C_1 \cap D_2} - q_{D_1 \cap A_2} - q_{D_1 \cap C_2} - q_{A_1 \cap D_2} \\ &= P_{(A_1 \cap A_2, [B_1 \cap B_2]_{\mathcal{J}})}. \end{aligned}$$

The last equality holds true since one can choose a finite family $\{B_i\}_{i=1}^n$ of mutually disjoint elements in \mathcal{B}_{reg} so that

$$(A_1 \cap C_2) \cup (D_1 \cap D_2) \cup (C_1 \cap A_2) \cup (C_1 \cap C_2) = \cup_{i=1}^n B_i.$$

Let $X_1 = A_1 \cup C_1 = B_1 \cup D_1$ and $X_2 = A_2 \cup C_2 = B_2 \cup D_2$. We then have that $(C_1 \cap C_2) \cup (C_1 \setminus X_2) \cup (C_2 \setminus X_1), (D_1 \cap D_2) \cup (D_1 \setminus X_2) \cup (D_2 \setminus X_1) \in \mathcal{B}_{reg}$,

$$\begin{aligned} & (A_1 \cup A_2) \cap ((C_1 \cap C_2) \cup (C_1 \setminus X_2) \cup (C_2 \setminus X_1)) \\ &= (B_1 \cup B_2) \cap ((D_1 \cap D_2) \cup (D_1 \setminus X_2) \cup (D_2 \setminus X_1)) = \emptyset \end{aligned}$$

and that

$$\begin{aligned} X_1 \cup X_2 &= A_1 \cup A_2 \cup ((C_1 \cap C_2) \cup (C_1 \setminus X_2) \cup (C_2 \setminus X_1)) \\ &= B_1 \cup B_2 \cup ((D_1 \cap D_2) \cup (D_1 \setminus X_2) \cup (D_2 \setminus X_1)). \end{aligned}$$

It then follows that

$$\begin{aligned} & P_{(A_1 \cup A_2, [B_1 \cup B_2]_{\mathcal{J}})} \\ &= p_{A_1 \cup A_2} + q_{(C_1 \cap C_2) \cup (C_1 \setminus X_2) \cup (C_2 \setminus X_1)} - q_{(D_1 \cap D_2) \cup (D_1 \setminus X_2) \cup (D_2 \setminus X_1)}. \end{aligned}$$

Since $C_1 \setminus (A_2 \cup C_2) = C_1 \setminus ((C_1 \cap A_2) \cup (C_1 \cap C_2))$, it follows that

$$q_{C_1 \setminus X_2} = q_{C_1} - q_{C_1 \cap A_2} - q_{C_1 \cap C_2}.$$

Similarly, we have

$$\begin{aligned} q_{C_2 \setminus X_1} &= q_{C_2} - q_{C_2 \cap A_1} - q_{C_2 \cap C_1}, \\ q_{D_1 \setminus X_2} &= q_{D_1} - q_{D_1 \cap A_2} - q_{D_1 \cap C_2}, \\ q_{D_2 \setminus X_1} &= q_{D_2} - q_{D_2 \cap A_1} - q_{D_2 \cap C_1}. \end{aligned}$$

We thus have that

$$\begin{aligned} & P_{(A_1, [B_1]_{\mathcal{J}})} + P_{(A_2, [B_2]_{\mathcal{J}})} - P_{(A_1 \cap A_2, [B_1 \cap B_2]_{\mathcal{J}})} \\ &= (p_{A_1} + q_{C_1} - q_{D_1}) + (p_{A_2} + q_{C_2} - q_{D_2}) \\ &\quad - p_{A_1 \cap A_2} - q_{C_1 \cap A_2} - q_{C_2 \cap A_1} - q_{C_1 \cap C_2} - q_{D_1 \cap D_2} \\ &\quad + q_{D_1 \cap A_2} + q_{D_1 \cap C_2} + q_{D_2 \cap A_1} + q_{D_2 \cap C_1} \\ &= (p_{A_1} + p_{A_2} - p_{A_1 \cap A_2}) \\ &\quad + q_{C_1 \cap C_2} + (q_{C_1} - q_{C_1 \cap A_2} - q_{C_1 \cap C_2}) + (q_{C_2} - q_{C_2 \cap A_1} - q_{C_2 \cap C_1}) \\ &\quad - q_{D_1 \cap D_2} - (q_{D_1} - q_{D_1 \cap A_2} - q_{D_1 \cap C_2}) - (q_{D_2} - q_{D_2 \cap A_1} - q_{D_2 \cap C_1}) \\ &= p_{A_1 \cup A_2} + q_{(C_1 \cap C_2) \cup (C_1 \setminus X_2) \cup (C_2 \setminus X_1)} - q_{(D_1 \cap D_2) \cup (D_1 \setminus X_2) \cup (D_2 \setminus X_1)} \\ &= P_{(A_1 \cup A_2, [B_1 \cup B_2]_{\mathcal{J}})}. \end{aligned}$$

(ii): Using $q_A s_{\alpha, A'} = s_{\alpha, A' \cap \theta_{\alpha}(A)} - s_{\alpha, A' \cap \theta_{\alpha}(A)} = 0$ for all $A \in \mathcal{B}_{reg}$ and $A' \in \mathcal{I}_{\alpha}$, we see that

$$\begin{aligned} P_{(A, [B]_{\mathcal{J}})} S_{\alpha, (A', [A']_{\mathcal{J}})} &= (p_A + q_C - q_D) s_{\alpha, A'} \\ &= s_{\alpha, A'} p_{\theta_{\alpha}(A)} \\ &= S_{\alpha, (A', [A']_{\mathcal{J}})} P_{\theta_{\alpha}(A, [B]_{\mathcal{J}})}. \end{aligned}$$

$$(iii): S_{\alpha, (A, [A]_{\mathcal{J}})}^* S_{\alpha', (A', [A']_{\mathcal{J}})}^* = s_{\alpha, A}^* s_{\alpha', A'} = \delta_{\alpha, \alpha'} p_{A \cap A'} = \delta_{\alpha, \alpha'} P_{(A \cap A', [A \cap A']_{\mathcal{J}})}.$$

(iv): For $(A, \emptyset) \in \tilde{\mathcal{B}}_{reg}$, we have

$$\begin{aligned} P_{(A, \emptyset)} = p_A &= \sum_{\alpha \in \Delta_{(A, \emptyset)}} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^* \\ &= \sum_{\alpha \in \Delta_{(A, \emptyset)}} S_{\alpha, (\theta_\alpha(A), [\theta_\alpha(A)]_{\mathcal{J}})} S_{\alpha, (\theta_\alpha(A), [\theta_\alpha(A)]_{\mathcal{J}})}^*. \end{aligned}$$

Thus we have a $*$ -homomorphism

$$\rho : C^*(\tilde{\mathcal{B}}, \mathcal{L}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha) \rightarrow C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$$

such that

$$\rho(p_{(A, [B]_{\mathcal{J}})}) = p_A + q_C - q_D \text{ and } \rho(s_{\alpha, (A, [A]_{\mathcal{J}})}) = s_{\alpha, A}$$

for $(A, [B]_{\mathcal{J}}) \in \tilde{\mathcal{B}}$, $\alpha \in \mathcal{L}$ and $(A, [A]_{\mathcal{J}}) \in \tilde{\mathcal{I}}_\alpha$.

It is then easy to check that

$$\begin{aligned} \rho \circ \phi(p_A) &= \rho(p_{(A, [A]_{\mathcal{J}})}) = p_A, \\ \rho \circ \phi(s_{\alpha, B}) &= \rho(s_{\alpha, (B, [B]_{\mathcal{J}})}) = s_{\alpha, B} \end{aligned}$$

for all $A \in \mathcal{B}$, $\alpha \in \mathcal{L}$ and $B \in \mathcal{I}_\alpha$. Hence, $\rho \circ \phi = id$. Also, we see that

$$\begin{aligned} &\phi \circ \rho(p_{(A, [B]_{\mathcal{J}})}) \\ &= \phi(p_A + q_C - q_D) \\ &= p_{(A, [A]_{\mathcal{J}})} + p_{(C, [C]_{\mathcal{J}})} - \sum_{\alpha \in \Delta_C} s_{\alpha, (\theta_\alpha(C), [\theta_\alpha(C)]_{\mathcal{J}})} s_{\alpha, (\theta_\alpha(C), [\theta_\alpha(C)]_{\mathcal{J}})}^* \\ &\quad - p_{(D, [D]_{\mathcal{J}})} + \sum_{\alpha \in \Delta_D} s_{\alpha, (\theta_\alpha(D), [\theta_\alpha(D)]_{\mathcal{J}})} s_{\alpha, (\theta_\alpha(D), [\theta_\alpha(D)]_{\mathcal{J}})}^* \\ &= p_{(A, [A]_{\mathcal{J}})} \\ &\quad + p_{(\emptyset, [C]_{\mathcal{J}})} + p_{(C, \emptyset)} - \sum_{\alpha \in \Delta_{(C, \emptyset)}} s_{\alpha, (\theta_\alpha(C), [\theta_\alpha(C)]_{\mathcal{J}})} s_{\alpha, (\theta_\alpha(C), [\theta_\alpha(C)]_{\mathcal{J}})}^* \\ &\quad - p_{(\emptyset, [D]_{\mathcal{J}})} - p_{(D, \emptyset)} + \sum_{\alpha \in \Delta_{(D, \emptyset)}} s_{\alpha, (\theta_\alpha(D), [\theta_\alpha(D)]_{\mathcal{J}})} s_{\alpha, (\theta_\alpha(D), [\theta_\alpha(D)]_{\mathcal{J}})}^* \\ &= p_{(A, [A]_{\mathcal{J}})} + p_{(\emptyset, [C]_{\mathcal{J}})} - p_{(\emptyset, [D]_{\mathcal{J}})} \\ &= p_{(A, [A \cup C]_{\mathcal{J}})} - p_{(\emptyset, [D]_{\mathcal{J}})} \\ &= p_{(A, [B]_{\mathcal{J}})}. \end{aligned}$$

for $(A, [B]_{\mathcal{J}}) \in \tilde{\mathcal{B}}$. The last equality holds since

$$p_{(A, [A \cup C]_{\mathcal{J}})} = p_{(A, [B \cup D]_{\mathcal{J}})} = p_{(A, [B]_{\mathcal{J}})} + p_{(\emptyset, [D]_{\mathcal{J}})}.$$

For $\alpha \in \mathcal{L}$ and $(A, [A]_{\mathcal{J}}) \in \tilde{\mathcal{I}}_\alpha$, we have

$$\phi \circ \rho(s_{\alpha, (A, [A]_{\mathcal{J}})}) = \phi(s_{\alpha, A}) = s_{\alpha, (A, [A]_{\mathcal{J}})}.$$

Hence, $\phi \circ \rho = id$. □

Remark 6.5. Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ be a relative generalized Boolean dynamical system. By Proposition 6.4, $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is isomorphic to $C^*(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha)$, and it follows from Proposition 6.3 that $C^*(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta}, \tilde{\mathcal{I}}_\alpha)$ is Morita equivalent to $C^*(\tilde{\mathcal{B}}, \tilde{\mathcal{L}}, \tilde{\theta})$. We thus have that the C^* -algebra of any relative generalized Boolean dynamical system is Morita equivalent to the C^* -algebra of a Boolean dynamical system.

7. GAUGE-INVARIANT IDEALS IN $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$

In this section, we give a complete list of the gauge-invariant ideals of C^* -algebras of $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ and describe the quotients as C^* -algebras of relative generalized Boolean dynamical systems, thereby generalizing [4, Proposition 10.11 and Theorem 10.12] and [7, Theorem 5.2].

Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ be a relative generalized Boolean dynamical system. Given a hereditary \mathcal{J} -saturated ideal \mathcal{H} of \mathcal{B} , we define

$$\mathcal{B}_{\mathcal{H}} := \{A \in \mathcal{B} : [A] \in (\mathcal{B}/\mathcal{H})_{reg}\}.$$

It is easy to see that $\mathcal{B}_{\mathcal{H}}$ is an ideal of \mathcal{B} and $\mathcal{H} \cup \mathcal{B}_{reg} \subseteq \mathcal{B}_{\mathcal{H}}$.

Fix a hereditary \mathcal{J} -saturated ideal \mathcal{H} of \mathcal{B} and an ideal \mathcal{S} of $\mathcal{B}_{\mathcal{H}}$ such that $\mathcal{H} \cup \mathcal{J} \subseteq \mathcal{S}$. Note that \mathcal{S} is also an ideal of \mathcal{B} . We let $I_{(\mathcal{H}, \mathcal{S})}$ denote the ideal of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J}) := C^*(p_A, s_{\alpha, B})$ generated by the family of projections (where we just write $\Delta_{[A]}$ for $\Delta_{[A]}^{(\mathcal{B}/\mathcal{H}, \mathcal{L}, \theta)}$)

$$\left\{ p_A - \sum_{\alpha \in \Delta_{[A]}} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^* : A \in \mathcal{S} \right\}.$$

Note that the family contains the family of projections $\{p_A : A \in \mathcal{H}\}$ (because if $A \in \mathcal{H}$, then $\Delta_{[A]} = \emptyset$).

We put $p_{A, \mathcal{H}} := \sum_{\alpha \in \Delta_{[A]}} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^*$ throughout this section.

Lemma 7.1. *The ideal $I_{(\mathcal{H}, \mathcal{S})}$ is gauge-invariant and*

$$I_{(\mathcal{H}, \mathcal{S})} = \overline{\text{span}}\{s_{\alpha, B}(p_A - p_{A, \mathcal{H}})s_{\beta, C}^* : A \in \mathcal{S}, \alpha, \beta \in \mathcal{L}^*, B \in \mathcal{I}_\alpha \text{ and } C \in \mathcal{I}_\beta\}. \quad (8)$$

Proof. Since $I_{(\mathcal{H}, \mathcal{S})}$ is generated by a set that is gauge-invariant, $I_{(\mathcal{H}, \mathcal{S})}$ is gauge-invariant.

It follows from Lemma 3.9 that the right-hand side J of (8) is an ideal of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha)$. Since $s_{\alpha, B}(p_A - p_{A, \mathcal{H}})s_{\beta, C}^* = s_{\alpha, B}p_A s_{\beta, C}^*$ for all $A \in \mathcal{H}$, J contains the generators of $I_{(\mathcal{H}, \mathcal{S})}$. Thus $I_{(\mathcal{H}, \mathcal{S})} \subseteq J$. The opposite inclusion is clear. \square

We shall prove in Theorem 7.3 that every gauge-invariant ideals of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is of the form $I_{(\mathcal{H}, \mathcal{S})}$ for a hereditary \mathcal{J} -saturated ideal \mathcal{H} and an ideal \mathcal{S} of $\mathcal{B}_{\mathcal{H}}$ with $\mathcal{H} \subseteq \mathcal{S}$ and $\mathcal{J} \subseteq \mathcal{S}$, and show that the quotient of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ by the ideal $I_{(\mathcal{H}, \mathcal{S})}$ fall into the class of C^* -algebras of relative generalized Boolean dynamical systems. To do this, we first observe the following.

Lemma 7.2. *Let I be a nonzero ideal in $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$.*

(1) *The set $\mathcal{H}_I := \{A \in \mathcal{B} : p_A \in I\}$ is a hereditary and \mathcal{J} -saturated ideal of \mathcal{B} .*

(2) The set

$$\mathcal{S}_I := \left\{ A \in \mathcal{B}_{\mathcal{H}_I} : p_A - \sum_{\alpha \in \Delta_{[A]}} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^* \in I \right\}$$

is an ideal of $\mathcal{B}_{\mathcal{H}_I}$ (and hence an ideal of \mathcal{B}) with $\mathcal{H}_I \subseteq \mathcal{S}_I$ and $\mathcal{J} \subseteq \mathcal{S}_I$.

Proof. (1): Suppose $A, B \in \mathcal{H}_I$. Then $p_{A \cup B} = p_A + p_B - p_{A \cap B} \in I$, so $A \cup B \in \mathcal{H}_I$.

Suppose then that $B \in \mathcal{H}_I$ and $A \in \mathcal{B}$ with $A \subseteq B$. Then $p_A = p_{A \cap B} = p_A p_B \in I$, so $A \in \mathcal{H}_I$.

This shows that \mathcal{H}_I is an ideal of \mathcal{B} . To show that \mathcal{H}_I is hereditary, suppose $A \in \mathcal{H}_I$. Then $p_A s_{\alpha, \theta_\alpha(A)} = s_{\alpha, \theta_\alpha(A)} \in I$ for all $\alpha \in \mathcal{L}$. Thus, $s_{\alpha, \theta_\alpha(A)}^* s_{\alpha, \theta_\alpha(A)} = p_{\theta_\alpha(A)} \in I$ for all $\alpha \in \mathcal{L}$, that is, $\theta_\alpha(A) \in \mathcal{H}_I$ for all $\alpha \in \mathcal{L}$.

Suppose now that $A \in \mathcal{J}$ and $\theta_\alpha(A) \in \mathcal{H}_I$ for all $\alpha \in \Delta_A$. Then $s_{\alpha, \theta_\alpha(A)} = s_{\alpha, \theta_\alpha(A)} p_{\theta_\alpha(A)} \in I$ for all $\alpha \in \Delta_A$. Thus, $p_A = \sum_{\alpha \in \Delta_A} s_{\alpha, \theta_\alpha(A)}^* s_{\alpha, \theta_\alpha(A)} \in I$, which means that $A \in \mathcal{H}_I$. Hence, the hereditary set \mathcal{H}_I is \mathcal{J} -saturated.

(2): If $A \in \mathcal{S}_I$ and $B \in \mathcal{B}_{\mathcal{H}_I}$, then obviously $A \cap B \in \mathcal{B}_{\mathcal{H}_I}$, and

$$\begin{aligned} I &\ni p_B \left(p_A - \sum_{\alpha \in \Delta_{[A]}} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^* \right) \\ &= p_{A \cap B} - \sum_{\alpha \in \Delta_{[A]}} s_{\alpha, \theta_\alpha(A \cap B)} s_{\alpha, \theta_\alpha(A \cap B)}^* \\ &= p_{A \cap B} - \sum_{\alpha \in \Delta_{[A \cap B]}} s_{\alpha, \theta_\alpha(A \cap B)} s_{\alpha, \theta_\alpha(A \cap B)}^*. \end{aligned}$$

The last equality holds true since $p_B s_{\alpha, \theta_\alpha(A)} = s_{\alpha, \theta_\alpha(A \cap B)} = 0$ for $\alpha \in \Delta_{[A]}$ with $\theta_\alpha(A \cap B) = \emptyset$. Thus, $A \cap B \in \mathcal{S}_I$. To show that \mathcal{S}_I is closed under finite unions, suppose $A, B \in \mathcal{S}_I$. Then clearly, $A \cup B \in \mathcal{B}_{\mathcal{H}_I}$ and we see that

$$\begin{aligned} p_{A \setminus B} - \sum_{\alpha \in \Delta_{[A \setminus B]}} s_{\alpha, \theta_\alpha(A \setminus B)} s_{\alpha, \theta_\alpha(A \setminus B)}^* &\in I, \\ p_{A \cap B} - \sum_{\alpha \in \Delta_{[A \cap B]}} s_{\alpha, \theta_\alpha(A \cap B)} s_{\alpha, \theta_\alpha(A \cap B)}^* &\in I, \\ p_{B \setminus A} - \sum_{\alpha \in \Delta_{[B \setminus A]}} s_{\alpha, \theta_\alpha(B \setminus A)} s_{\alpha, \theta_\alpha(B \setminus A)}^* &\in I. \end{aligned}$$

Adding the 3 elements above and using the fact that

$$p_{A \cup B} = \sum_{\alpha \in \Delta_{[A \setminus B]}} s_{\alpha, \theta_\alpha(A \setminus B)} s_{\alpha, \theta_\alpha(A \setminus B)}^* + \sum_{\alpha \in \Delta_{[A \cap B]}} s_{\alpha, \theta_\alpha(A \cap B)} s_{\alpha, \theta_\alpha(A \cap B)}^* + \sum_{\alpha \in \Delta_{[B \setminus A]}} s_{\alpha, \theta_\alpha(B \setminus A)} s_{\alpha, \theta_\alpha(B \setminus A)}^*, \quad (9)$$

we see that $p_{A \cup B} - \sum_{\alpha \in \Delta_{[A \cup B]}} s_{\alpha, \theta_\alpha(A \cup B)} s_{\alpha, \theta_\alpha(A \cup B)}^* \in I$. Hence, $A \cup B \in \mathcal{S}_I$.

Since $\Delta_{[A]} = \emptyset$ for all $A \in \mathcal{H}_I$, it is rather obvious that $\mathcal{H}_I \subseteq \mathcal{S}_I$. For any $A \in \mathcal{J} \setminus \mathcal{H}_I$, we have $p_A + I = \sum_{\alpha \in \Delta_{[A]}} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^* + I$, and hence, $\mathcal{J} \subseteq \mathcal{S}_I$. \square

Proposition 7.3. *Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ be a relative generalized Boolean dynamical system. Suppose that I is an ideal of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$.*

There is then a surjective $*$ -homomorphism

$$\phi_I : C^*(\mathcal{B}/\mathcal{H}_I, \mathcal{L}, \theta, [\mathcal{I}_\alpha]; [\mathcal{S}_I]) \rightarrow C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})/I$$

such that

$$\phi_I(p_{[A]_{\mathcal{H}_I}}) = p_A + I \quad \text{and} \quad \phi_I(s_{\alpha, [B]_{\mathcal{H}_I}}) = s_{\alpha, B} + I,$$

where $[\mathcal{I}_\alpha] := \{[A] \in \mathcal{B}/\mathcal{H}_I : A \in \mathcal{I}_\alpha\}$ and $[\mathcal{S}_I] := \{[A] \in \mathcal{B}/\mathcal{H}_I : A \in \mathcal{S}_I\}$. Moreover, the following are equivalent.

- (1) I is gauge-invariant.
- (2) The map ϕ_I is an isomorphism.
- (3) $I = I_{(\mathcal{H}_I, \mathcal{S}_I)}$.

Proof. Since $p_A \in I$ and $s_{\alpha, A} = s_{\alpha, A}p_A \in I$ if $A \in \mathcal{H}_I$, it follows that $\{p_A + I : A \in \mathcal{B}\} \cup \{s_{\alpha, B} + I : \alpha \in \mathcal{L}, B \in \mathcal{I}_\alpha\}$ is a $(\mathcal{B}/\mathcal{H}_I, \mathcal{L}, \theta, [\mathcal{I}_\alpha]; [\mathcal{S}_I])$ -representation. So, the universal property of $C^*(\mathcal{B}/\mathcal{H}_I, \mathcal{L}, \theta, [\mathcal{I}_\alpha]; [\mathcal{S}_I])$ gives us a surjective $*$ -homomorphism

$$\phi_I : C^*(\mathcal{B}/\mathcal{H}_I, \mathcal{L}, \theta, [\mathcal{I}_\alpha]; [\mathcal{S}_I]) \rightarrow C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})/I$$

such that

$$\phi_I(p_{[A]_{\mathcal{H}_I}}) = p_A + I \quad \text{and} \quad \phi_I(s_{\alpha, [B]_{\mathcal{H}_I}}) = s_{\alpha, B} + I.$$

That (1) \implies (2) follows from Theorem 6.1, and that (3) \implies (1) follows from Lemma 7.1.

(2) \implies (3): Since $I_{(\mathcal{H}_I, \mathcal{S}_I)} \subseteq I$, there is a surjective $*$ -homomorphism

$$q : C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})/I_{(\mathcal{H}_I, \mathcal{S}_I)} \rightarrow C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})/I$$

such that $q(p_A + I_{(\mathcal{H}_I, \mathcal{S}_I)}) = p_A + I$ and $q(s_{\alpha, [B]_{\mathcal{H}_I}} + I_{(\mathcal{H}_I, \mathcal{S}_I)}) = s_{\alpha, B} + I$. An argument similar to the one used to construct ϕ_I gives us a surjective $*$ -homomorphism

$$\phi_{I_{(\mathcal{H}_I, \mathcal{S}_I)}} : C^*(\mathcal{B}/\mathcal{H}_I, \mathcal{L}, \theta, [\mathcal{I}_\alpha]; [\mathcal{S}_I]) \rightarrow C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})/I_{(\mathcal{H}_I, \mathcal{S}_I)}$$

such that $\phi_{I_{(\mathcal{H}_I, \mathcal{S}_I)}}(p_{[A]_{\mathcal{H}_I}}) = p_A + I_{(\mathcal{H}_I, \mathcal{S}_I)}$ and $\phi_{I_{(\mathcal{H}_I, \mathcal{S}_I)}}(s_{\alpha, [B]_{\mathcal{H}_I}}) = s_{\alpha, B} + I_{(\mathcal{H}_I, \mathcal{S}_I)}$. We have that $\phi_I = q \circ \phi_{I_{(\mathcal{H}_I, \mathcal{S}_I)}}$. It follows that if ϕ_I is an isomorphism, then q is an isomorphism and $I = I_{(\mathcal{H}_I, \mathcal{S}_I)}$. \square

The set of pairs $(\mathcal{H}, \mathcal{S})$, where \mathcal{H} is a hereditary \mathcal{J} -saturated ideal of \mathcal{B} and \mathcal{S} is an ideal of $\mathcal{B}_{\mathcal{H}}$ with $\mathcal{H} \cup \mathcal{J} \subseteq \mathcal{S}$ is a lattice with respect to the order relation defined by $(\mathcal{H}_1, \mathcal{S}_1) \leq (\mathcal{H}_2, \mathcal{S}_2) \iff (\mathcal{H}_1 \subseteq \mathcal{H}_2 \wedge \mathcal{S}_1 \subseteq \mathcal{S}_2)$. The set of gauge-invariant ideals of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ is a lattice with the order given by set inclusion (the meet of I_1 and I_2 is the ideal $I_1 \cap I_2$, and the join of I_1 and I_2 is the ideal generated by $I_1 \cup I_2$).

Theorem 7.4. *Let $(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ be a relative generalized Boolean dynamical system. Then the map $(\mathcal{H}, \mathcal{S}) \mapsto I_{(\mathcal{H}, \mathcal{S})}$ is a lattice isomorphism between the lattice of all pairs $(\mathcal{H}, \mathcal{S})$, where \mathcal{H} is a hereditary \mathcal{J} -saturated ideal of \mathcal{B} and \mathcal{S} is an ideal of $\mathcal{B}_{\mathcal{H}}$ with $\mathcal{H} \cup \mathcal{J} \subseteq \mathcal{S}$, and the lattice of all gauge-invariant ideals of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$.*

Proof. It follows from Theorem 7.3 the map $(\mathcal{H}, \mathcal{S}) \mapsto I_{(\mathcal{H}, \mathcal{S})}$ is onto.

To see the map $(\mathcal{H}, \mathcal{S}) \mapsto I_{(\mathcal{H}, \mathcal{S})}$ is injective, we show that

$$\mathcal{H}_{I_{(\mathcal{H}, \mathcal{S})}} = \mathcal{H} \quad \text{and} \quad \mathcal{S}_{I_{(\mathcal{H}, \mathcal{S})}} = \mathcal{S}.$$

It is easy to see that $\mathcal{H} \subseteq \mathcal{H}_{I(\mathcal{H}, \mathcal{S})}$ and $\mathcal{S} \subseteq \mathcal{S}_{I(\mathcal{H}, \mathcal{S})}$. It follows from the universal property of $\{p_A, s_{\alpha, B} : A \in \mathcal{B}, \alpha \in \mathcal{L}, B \in \mathcal{I}_\alpha\}$ that there is a $*$ -homomorphism $\phi : C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J}) \rightarrow C^*(\mathcal{B}/\mathcal{H}, \mathcal{L}, \theta, [\mathcal{I}_\alpha]; [\mathcal{S}])$ such that $\phi(p_A) = p_{[A]}$ and $\phi(s_{\alpha, B}) = s_{\alpha, [B]}$. Let $I' = \ker(\phi)$. Then $I_{(\mathcal{H}, \mathcal{S})} \subseteq I'$. It follows from Corollary 5.7 that $\mathcal{H}_{I'} = \mathcal{H}$ and $\mathcal{S}_{I'} = \mathcal{S}$. It follows that $\mathcal{H}_{I(\mathcal{H}, \mathcal{S})} \subseteq \mathcal{H}_{I'} = \mathcal{H}$ and $\mathcal{S}_{I(\mathcal{H}, \mathcal{S})} \subseteq \mathcal{S}_{I'} = \mathcal{S}$.

Next we show that $(\mathcal{H}, \mathcal{S}) \mapsto I_{(\mathcal{H}, \mathcal{S})}$ is order preserving. For this suppose that for $i = 1, 2$, \mathcal{H}_i is a hereditary \mathcal{J} -saturated ideal of \mathcal{B} and \mathcal{S}_i is an ideal of $\mathcal{B}_{\mathcal{H}_i}$ such that $\mathcal{H}_i \cup \mathcal{J} \subseteq \mathcal{S}_i$, and that $\mathcal{H}_1 \subseteq \mathcal{H}_2$ and $\mathcal{S}_1 \subseteq \mathcal{S}_2$. Suppose $A \in \mathcal{S}_1$. We then have that $\theta_\alpha(A) \in \mathcal{H}_2$ for $\alpha \in \Delta_{[A]}^{(\mathcal{B}/\mathcal{H}_1, \mathcal{L}, \theta)} \setminus \Delta_{[A]}^{(\mathcal{B}/\mathcal{H}_2, \mathcal{L}, \theta)}$, and thus that $p_A - \sum_{\alpha \in \Delta_{[A]}^{(\mathcal{B}/\mathcal{H}_1, \mathcal{L}, \theta)}} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^* \in I_{(\mathcal{H}_2, \mathcal{S}_2)}$. It follows that $I_{(\mathcal{H}_1, \mathcal{S}_1)} \subseteq I_{(\mathcal{H}_2, \mathcal{S}_2)}$.

For the converse suppose I_1 and I_2 are gauge-invariant ideals of $C^*(\mathcal{B}, \mathcal{L}, \theta, \mathcal{I}_\alpha; \mathcal{J})$ such that $I_1 \subseteq I_2$. Then clearly, $\mathcal{H}_{I_1} = \{A \in \mathcal{B} : p_A \in I_1\} \subseteq \{A \in \mathcal{B} : p_A \in I_2\} = \mathcal{H}_{I_2}$. Suppose $A \in \mathcal{S}_{I_1}$. Then $\theta_\alpha(A) \in \mathcal{H}_{I_2}$ for $\alpha \in \Delta_{[A]}^{(\mathcal{B}/\mathcal{H}_{I_1}, \mathcal{L}, \theta)} \setminus \Delta_{[A]}^{(\mathcal{B}/\mathcal{H}_{I_2}, \mathcal{L}, \theta)}$. It follows that

$$\begin{aligned} p_A - \sum_{\alpha \in \Delta_{[A]}^{(\mathcal{B}/\mathcal{H}_{I_2}, \mathcal{L}, \theta)}} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^* &= p_A - \sum_{\alpha \in \Delta_{[A]}^{(\mathcal{B}/\mathcal{H}_{I_1}, \mathcal{L}, \theta)}} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^* \\ &+ \sum_{\alpha \in \Delta_{[A]}^{(\mathcal{B}/\mathcal{H}_{I_1}, \mathcal{L}, \theta)} \setminus \Delta_{[A]}^{(\mathcal{B}/\mathcal{H}_{I_2}, \mathcal{L}, \theta)}} s_{\alpha, \theta_\alpha(A)} s_{\alpha, \theta_\alpha(A)}^* \in I_2. \end{aligned}$$

This shows that $\mathcal{S}_{I_1} \subseteq \mathcal{S}_{I_2}$. We thus have that $(\mathcal{H}, \mathcal{S}) \mapsto I_{(\mathcal{H}, \mathcal{S})}$ is a lattice isomorphism. \square

Acknowledgements. This research was partially supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education(NRF-2017R1D1A1B03030540).

The authors would like to thank the referee for valuable comments and suggestions.

REFERENCES

- [1] T. Bates, T. M. Carlsen, and D. Pask, *C^* -algebras of labelled graphs III— K -theory computations*, Ergod. Th. and Dynam. Sys. **37**(2017), Issue 2, 337–368.
- [2] T. Bates and D. Pask, *C^* -algebras of labelled graphs*, J. Operator Theory. **57**(2007), Issue 1, 207–226.
- [3] T. Bates and D. Pask, *C^* -algebras of labelled graphs II - simplicity results*, Math. Scand. **104**(2009), No. 2, 249–274.
- [4] T. M. Carlsen, E. Ortega, and E. Pardo, *C^* -algebras associated to Boolean dynamical systems*, J. Math. Anal. Appl. **450**(2017), Issue 1, 727–768.
- [5] N. J. Fowler, P.S. Muhly and I. Raeburn, *Representations of Cuntz-Pimsner algebras*, Indiana Univ. Math. J. **52**(2003), No. 3, 569–605.

- [6] N. J. Fowler and I. Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, Indiana Univ. Math. J. **48**(1999), No. 1, 155–181.
- [7] J. A. Jeong, S. H. Kim and G. H. Park, *The structure of gauge-invariant ideals of labelled graph C^* -algebras*, J. Funct. Anal. **262**(2012), Issue 4, 1759–1780.
- [8] T. Katsura, *On C^* -algebras associated with C^* -correspondences*, J. Funct. Anal. **217**(2004), Issue 2, 366–401.
- [9] T. Katsura, *Ideal structure of C^* -algebras associated with C^* -correspondences*, Pacific J. Math. **230**(2007), No. 2, 107–145.
- [10] A. Kumjian, D. Pask, I. Raeburn and J. Renault *Graphs, groupoids, and Cuntz–Krieger algebras*, J. Funct. Anal. **144**(1997), Issue 2, 505–541.
- [11] P.S. Muhly and B. Selel, *Tensor algebras over C^* -correspondences: representations, dilations, and C^* -envelopes*, J. Funct. Anal. **158**(1998), Issue 2, 389–457.
- [12] P.S. Muhly and M. Tomforde, *Adding Tails to C^* -correspondences*, Doc. Math. **9**(2004), 79–106.
- [13] I. Raeburn, *Graph Algebras*, CBMS Regional Conference Series in Mathematics, vol. 103, Amer. Math. Soc., Providence, RI 2005.

DEPARTMENT OF SCIENCES AND TECHNOLOGY, UNIVERSITY OF FAROE ISLANDS, NÓATÚN 3,
FO-100 TÓRSHAVN, FAROE ISLANDS
Email address: `toke.carlsen@gmail.com`

RESEARCH INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KOREA
Email address: `kkang33@snu.ac.kr`