

What is the B mode in CMB polarization?

Jack Y. L. Kwok¹, King Lau², Edward Young³ and Kenneth Young¹

¹ Department of Physics, The Chinese University of Hong Kong, Hong Kong, China

² Department of Physics, University of Minnesota, Minneapolis, MN 55455, USA

³ Department of Physics, Stanford University, Stanford, CA 94305, USA

E-mail: jackkwok@link.cuhk.edu.hk, kennylau@umn.edu, eyy@stanford.edu, kyong@cuhk.edu.hk

Keywords: CMB, polarization, B mode, inflation

2 December 2019

Abstract.

A linear polarization field on a surface is expressed in terms of scalar functions, providing an invariant separation into two components; one of these is the B mode, important as a signature of primordial gravitational waves, which would lend support to the inflation hypothesis. The case of a plane already exhibits the key ideas, including the formal analogy with a vector field decomposed into gradient plus curl, with the B mode like the latter. The formalism is generalized to a spherical surface using cartesian coordinates. Analysis of global data provides a path to vector and tensor spherical harmonics.

1. Introduction

1.1. Physical context

The isotropic black-body Cosmic Microwave Background (CMB) [1, 2, 3] indicates a dense and hot early universe and launched modern big-bang cosmology. With a dipole angular dependence [4, 5] due to observer motion ($v/c \sim 10^{-3}$) removed, the remaining anisotropies are at the $\sim 10^{-4}$ level [6] even across widely separated directions, thus prompting the inflation hypothesis [7, 8]. Better data [9] have enabled precision cosmology. Recent attention has turned to polarization [10], conventionally separated into the so-called E and B modes, at relative levels of $\sim 10^{-6}$ and $\sim 10^{-7}$. The CMB E modes, with sharper acoustic peaks, provide additional constraints on the standard Lambda cold dark matter (Λ CDM) model parameters and help to determine the optical depth to re-ionization.

The tiny B mode is important in cosmology: to first order it cannot be produced by scalar (e.g., density) perturbations, but only by tensor perturbations [11, 12], i.e., strong gravitational waves. Thus a cosmological B mode would support the

inflation hypothesis. However, detected signals so far [13] are not above foreground contamination [14, 15] (including lensing from E modes). Many further experiments are in progress [16] to pin down the cosmological B mode. Setting aside the enormous observational challenges (e.g., instrumental, low signal-to-noise, foreground contamination), there is a key conceptual question that deserves to be better understood by a wider community: How is the B mode defined and hence extracted from the polarization data?

The E and B modes are heuristically likened to a gradient and a curl, or TE/TM modes. Typical statements refer to “a twisting pattern, or spin, known as a curl or B -mode” [17]; or to “the curl component (B mode)” [18]; or to components having respectively “vanishing curl” and “vanishing divergence” [19]. If interpreted naively, these statements are misleading — the usual conception of gradient and curl does not apply to polarization. This paper explains the decomposition of the polarization field in an elementary way. By avoiding the apparatus of curved manifolds (celestial sphere) and in particular emphasizing the case of a plane, the account should be accessible to a wide audience.

1.2. Polarization

Consider a wave propagating along $+z$, with electric field components E_j , $j = 1, 2$, in the usual complex notation. Quadratic averages $S_{ij} = \text{Av}(E_i^* E_j)$ form a 2×2 Hermitian matrix with $S_{11} = I + Q$, $S_{22} = I - Q$, $S_{12} = S_{21}^* = U + iV$. Of the four Stokes parameters [20], the overall intensity I is considered separately. In CMB, V is usually ignored, because primordial circular polarization is not expected in standard models [21, 22]; with few exceptions that place upper limits [23, 24], V is assumed to be zero as a systematic check [22]. With I, V removed, polarization is described by the symmetric traceless tensor

$$\mathbb{P} = \begin{pmatrix} Q & U \\ U & -Q \end{pmatrix}. \quad (1)$$

All the information in \mathbb{P} is contained in its first column $\mathbf{P} = (P_1, P_2)^T = (Q, U)^T$, to be called a doublet. The linear polarization of magnitude $|\mathbf{P}|$ is along the direction χ , with $\tan 2\chi = U/Q$. Upon rotation by an angle α , $Q \pm iU$ gain a factor $e^{\pm 2i\alpha}$, i.e., they transform as spin ± 2 .

1.3. The central issue

Polarization \mathbb{P} , a rank-2 tensor, cannot be “similar to” a vector \mathbf{V} , for which one speaks of gradient and curl. A vector field (arrows) returns to itself upon rotation by 2π . A linear polarization field (*unsigned* line segments) returns to itself upon rotation by π . There is no polarization analogue to $\int \mathbf{V} \cdot d\mathbf{r}$.

If a scalar field varies along \mathbf{k} , then a vector field that is a gradient (curl) makes an angle $n\pi/2$ with \mathbf{k} , with n even (odd). However, E (B) mode polarization makes an

angle of $n\pi/4$ with \mathbf{k} , with n even (odd).

Thus a factor of 2 appears in the rank of the tensor, the phase under rotations and the angle relative to \mathbf{k} — all for the same reason. To avoid the strained analogy, henceforth the two components will be simply called the *A* and *B* modes and denoted with superscripts a and b .

The mathematical formalism in the literature [11, 19] that supports the heuristic analogy is typically bundled with additional elements. First, reference to inflation [18] is not needed, just as an electric field can be decomposed into a gradient plus a curl without reference to Maxwell’s equations. The complications of a sphere [19] can be avoided by (first) considering a plane, useful for narrow-field observations [25] or as a stepping stone to a sphere. A global harmonic analysis [19] is also not strictly necessary: the separation can in principle be implemented locally. This paper strips away these complications to provide a simple account of the separation of a linear polarization field into two components, for access by a wider audience.

1.4. Overview

The paper is structured so that readers can stop at various points depending on the level of technical detail desired. The formalism is first developed for the xy plane using the polar representation (Section 2.1): a vector field is expressed as $(\partial_1 + i\partial_2)$ and a polarization field as $(\partial_1 + i\partial_2)^2$ acting on scalar fields, with equations (2) and (6) exhibiting the analogy. As far as concepts are concerned, this is the end of the story.

The formalism, rewritten in cartesian coordinates (Sect. 2.2), is ported first to a plane with unit normal $\hat{\mathbf{n}}$ (Sect. 2.3) and then, upon $\hat{\mathbf{n}} \mapsto \mathbf{r}$, to a *small* part of a unit sphere (Sect. 3). The resultant formalism, involving the angular momentum \mathbf{L} rather than ∇ , applies “classically”, i.e., if commutators are ignored, valid for mode number $\ell \gg 1$. The proper ordering of operators yields lower-order corrections in \mathbf{L} . Global analysis (Sect. 4) leads to vector and tensor spherical harmonics, and provides a tool for disentangling different contributions to each polarization component.

2. Formalism for a plane

The objective is to express either a vector field \mathbf{V} or a polarization field \mathbb{P} on a surface as derivatives of two scalar fields.

2.1. Polar representation

2.1.1. Vector case. The decomposition of a vector field into a gradient and a curl (Helmholtz’s theorem) need not be repeated, but there are special features in two dimensions. Given a real vector field $\mathbf{V} = (V_1, V_2)^T$ on the xy plane (say in the domain $0 \leq x \leq \mathcal{L}_x$, $0 \leq y \leq \mathcal{L}_y$ with periodic boundary conditions), we seek a representation (the only form consistent with the angular momentum structure)

$$V_1 \pm iV_2 = (\partial_1 \pm i\partial_2) (\Phi^a \pm i\Phi^b) \quad (2)$$

in terms of real scalar functions $\Phi^{a,b}$. Henceforth only one of the conjugate equations will be shown. To prove such a representation, consider

$$\varphi^a + i\varphi^b \equiv (\partial_1 - i\partial_2)(V_1 + iV_2). \quad (3)$$

Since $(\partial_1 - i\partial_2)$ has -1 unit of angular momentum, $\varphi^{a,b}$ are scalars. The key identity

$$(\partial_1 - i\partial_2)(\partial_1 + i\partial_2) = \nabla_{\perp}^2 \quad (4)$$

leads to

$$\varphi^a + i\varphi^b = \nabla_{\perp}^2 (\Phi^a + i\Phi^b). \quad (5)$$

Thus, given \mathbf{V} , first finds (locally) $\varphi^{a,b}$ by (3); then invert the laplacian to give $\Phi^{a,b}$. With global data, the inversion is unique, proved by expanding in the eigenfunctions of ∇_{\perp}^2 — provided $\varphi^{a,b}$ have no projection on the null space of ∇_{\perp}^2 . The latter condition is obvious from (3) and the periodic boundary condition; another proof is given in (16) below. If data are confined to a subset, one can imagine any extension of $\varphi^{a,b}$ to the entire plane, and then invert; different extensions yield different $\Phi^{a,b}$.

2.1.2. Tensor case. Since $Q \pm iU$ have ± 2 units of angular momentum, the analogous representation must take the form

$$Q \pm iU = (\partial_1 \pm i\partial_2)^2 (\Phi^a \pm i\Phi^b), \quad (6)$$

involving a second-order operator. Define

$$\varphi^a + i\varphi^b \equiv (\partial_1 - i\partial_2)^2 (Q + iU). \quad (7)$$

Using the square of the identity (4), we find

$$\varphi^a + i\varphi^b = K (\Phi^a + i\Phi^b), \quad (8)$$

where K and its eigenvalue N_K (for plane waves with wave number \mathbf{k}) are

$$K = \nabla_{\perp}^4, \quad N_K = k^4. \quad (9)$$

The algorithm is the same: use (7) to obtain (locally) $\varphi^{a,b}$ and invert K to obtain $\Phi^{a,b}$.

Thus polarization on a plane is expressible in terms of two scalars, which importantly do not depend on the coordinate choice [11], in a manner analogous to the case of vector fields; as immediately shown in (11), the latter is the same as a gradient plus a curl.

2.1.3. Local separation. Whether for the vector or the tensor case, one can determine $\varphi^{a,b}$ locally and in principle assess the relative importance of the B mode on any subset Ω of the plane by the ratio R^b/R^a , where $R^{a,b} = \int_{\Omega} dS |\varphi^{a,b}|^2$.

2.2. Cartesian representation

2.2.1. *Vector case.* In cartesian coordinates, (2) is equivalent to

$$\mathbf{V} = \mathbf{d}\Phi^a - \tilde{\mathbf{d}}\Phi^b \quad (10)$$

$$= \nabla_{\perp}\Phi^a + \nabla_{\perp} \times (-\hat{\mathbf{n}}\Phi^b), \quad (11)$$

i.e., as a gradient plus a curl, where

$$\mathbf{d} = \nabla_{\perp} \quad , \quad \tilde{\mathbf{d}} = -\hat{\mathbf{n}} \times \mathbf{d}, \quad (12)$$

in which $\hat{\mathbf{n}} = \hat{\mathbf{e}}_z$ is the unit normal to the plane. Duals can equivalently be defined by

$$\tilde{\mathbf{d}} = \mathbb{E}\mathbf{d} \quad , \quad \mathbb{E} \equiv \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (13)$$

This latter definition for duals also applies to doublets and tensors.

The key identity (4) is equivalent to

$$\mathbf{d}^{\dagger} \cdot \mathbf{d} = \tilde{\mathbf{d}}^{\dagger} \cdot \tilde{\mathbf{d}} = -\nabla_{\perp}^2 \quad , \quad \tilde{\mathbf{d}}^{\dagger} \cdot \mathbf{d} = 0. \quad (14)$$

The orthogonality relation allows different components to be projected out. The definition of $\varphi^{a,b}$ in (3) becomes

$$\varphi^a = -\mathbf{d}^{\dagger} \cdot \mathbf{V} = \nabla_{\perp} \cdot \mathbf{V} \quad , \quad \varphi^b = \tilde{\mathbf{d}}^{\dagger} \cdot \mathbf{V} = (\nabla_{\perp} \times \mathbf{V}) \cdot \hat{\mathbf{n}}. \quad (15)$$

It can now be proved more formally that $\varphi^{a,b}$ have no projection on the null space \mathcal{S} of ∇_{\perp}^2 . Consider $F \in \mathcal{S}$ and define inner products in the usual way; then, e.g.,

$$(F, \varphi^a) = (F, \nabla_{\perp} \cdot \mathbf{V}) = -(\nabla_{\perp} F, \mathbf{V}). \quad (16)$$

But $(\nabla_{\perp} F, \nabla_{\perp} F) = (F, -\nabla_{\perp}^2 F) = 0$. This proof generalizes to tensors, and to the case of a sphere (for which $\varphi^{a,b}$ must have no projection on $\ell = 0, 1$).

2.2.2. *Tensor case.* Write the second-order differential operator as

$$(\partial_1 + i\partial_2)^2 = (\partial_1^2 - \partial_2^2) + i(2\partial_1\partial_2) \equiv D_1 + iD_2.$$

Define

$$\mathbb{D} = \begin{pmatrix} D_1 & D_2 \\ D_2 & -D_1 \end{pmatrix} \quad (17)$$

and its dual $\tilde{\mathbb{D}} = \mathbb{E}\mathbb{D}$. Denote the two columns of \mathbb{D} as $\mathbf{D} = (D_1, D_2)^T$ and $\tilde{\mathbf{D}} = \mathbb{E}\mathbf{D} = (D_2, -D_1)^T$.

The second-order analog to (14) is

$$\mathbb{D}^{\dagger}\mathbb{D} = \tilde{\mathbb{D}}^{\dagger}\tilde{\mathbb{D}} = K\mathbb{I} \quad , \quad \tilde{\mathbb{D}}^{\dagger}\mathbb{D} = K\tilde{\mathbb{I}}, \quad (18)$$

where \mathbb{I} is the identity, $\tilde{\mathbb{I}} = \mathbb{E}\mathbb{I} = \mathbb{E}$. Only a weaker version of (18) is needed:

$$\text{tr } \mathbb{D}^{\dagger}\mathbb{D} = \text{tr } \tilde{\mathbb{D}}^{\dagger}\tilde{\mathbb{D}} = 2K \quad , \quad \text{tr } \tilde{\mathbb{D}}^{\dagger}\mathbb{D} = 0. \quad (19)$$

The first- and second-order operators are related by

$$\begin{aligned} \mathbb{D} + i\tilde{\mathbb{D}} &= (\mathbf{d} + i\tilde{\mathbf{d}}) \otimes (\mathbf{d} + i\tilde{\mathbf{d}}), \\ \mathbb{D} &= \mathbf{d} \otimes \mathbf{d} - \tilde{\mathbf{d}} \otimes \tilde{\mathbf{d}} \quad , \quad \tilde{\mathbb{D}} = \mathbf{d} \otimes \tilde{\mathbf{d}} + \tilde{\mathbf{d}} \otimes \mathbf{d}, \end{aligned} \quad (20)$$

where $\mathbf{a} \otimes \mathbf{b}$ has matrix elements $a_i b_j$. Then (6) becomes

$$\mathbb{P} = \mathbb{D} \Phi^a - \tilde{\mathbb{D}} \Phi^b . \quad (21)$$

To prove this entirely using cartesian components, note that in analogy to (7) and (8),

$$\varphi^a + i\varphi^b \equiv (1/2) \text{tr} [(\mathbb{D}^\dagger - i\tilde{\mathbb{D}}^\dagger)\mathbb{P}] = K (\Phi^a + i\Phi^b) , \quad (22)$$

in which (19) has been used. All other steps are obvious. In analogy to (16), $\varphi^{a,b}$ have no projection on the null space of K .

2.3. Porting to another plane

To port the formalism to any plane with a unit normal $\hat{\mathbf{n}}$, remove explicit reference to ∇_\perp by adopting the equivalent definition (referring to ∇ but not to ∇_\perp)

$$\tilde{\mathbf{d}} = -\hat{\mathbf{n}} \times \nabla \quad , \quad \mathbf{d} = \hat{\mathbf{n}} \times \tilde{\mathbf{d}} . \quad (23)$$

All other formulas remain the same. However, \mathbf{V} , \mathbf{d} and $\tilde{\mathbf{d}}$ should now be regarded not as 2-vectors defined on a plane, but as 3-vectors satisfying a tangential condition: $n_i V_i = n_i d_i = n_i \tilde{d}_i = 0$. Likewise, \mathbb{P} , \mathbb{D} , $\tilde{\mathbb{D}}$ are 3×3 tensors satisfying $n_i P_{ij} = n_i D_{ij} = n_i \tilde{D}_{ij} = 0$.

3. Formalism for a sphere

3.1. Mapping the differential operator

A *small* part of the surface of a unit sphere can be regarded as a plane with normal $\hat{\mathbf{n}} = \mathbf{r}$. Then using (23), we have, in terms of the angular momentum \mathbf{L} ($\hbar = 1$):

$$\tilde{\mathbf{d}} = -i\mathbf{L} \quad , \quad \mathbf{d} = \hat{\mathbf{n}} \times (-i\mathbf{L}) \equiv i\tilde{\mathbf{L}} . \quad (24)$$

3.2. Vector field

The planar formula (10) applies, but with \mathbf{d} and $\tilde{\mathbf{d}}$ replaced as above. The orthogonality relations are

$$\mathbf{d}^\dagger \cdot \mathbf{d} = \tilde{\mathbf{d}}^\dagger \cdot \tilde{\mathbf{d}} = \mathbf{L}^2 \quad , \quad \tilde{\mathbf{d}}^\dagger \cdot \mathbf{d} = \mathbf{L} \cdot \tilde{\mathbf{L}} = 0 . \quad (25)$$

Only some key steps in transcribing the formalism will be sketched; the rest is obvious. For example, (10) becomes, upon the replacement (24),

$$\mathbf{V} = (i\tilde{\mathbf{L}}) \Phi^a + (i\mathbf{L}) \Phi^b . \quad (26)$$

Define

$$\varphi^a \equiv (i\tilde{\mathbf{L}})^\dagger \cdot \mathbf{V} = L^2 \Phi^a \quad , \quad \varphi^b \equiv (i\mathbf{L})^\dagger \cdot \mathbf{V} = L^2 \Phi^b . \quad (27)$$

Inversion of L^2 then gives $\Phi^{a,b}$. The inversion depends on $\varphi^{a,b}$ having zero projection on the null space of L^2 , proved in a manner similar to (16).

3.3. Polarization field

Equation (21) applies with the replacement (24), so that (absorbing some signs):

$$\mathbb{D} \approx (\mathbf{L} \otimes \mathbf{L} - \tilde{\mathbf{L}} \otimes \tilde{\mathbf{L}}) \quad , \quad \tilde{\mathbb{D}} \approx (\tilde{\mathbf{L}} \otimes \mathbf{L} + \mathbf{L} \otimes \tilde{\mathbf{L}}) . \quad (28)$$

The symbol \approx indicates that problems arising out of the ordering of operators are ignored (see below). The orthogonality relation (19) again holds, except that K and its eigenvalue are now given by $K \approx L^4$, $N_K \approx [\ell(\ell+1)]^2$. Everything in the planar case can be repeated with the replacement of K .

The results indicated with \approx are not exact: the order of the operators leads to ambiguities proportional to the commutators. Schematically $[L, L] = i\hbar L$, so corrections are ‘‘quantum’’ and lower-order in ℓ . In practice, many experiments targeting the cosmological B mode focus on $\ell \sim 80$ or larger [13], so classical (i.e., $\ell \gg 1$) formulas are already useful as a start.

To determine the subsidiary terms, from (28), one can write $D_{ij} = X_{ij} - Y_{ij}$ and $\tilde{D}_{ij} = Z_{ij}$, where to leading order $X_{ij} \approx L_i L_j$, $Y_{ij} \approx \tilde{L}_i \tilde{L}_j$ and $Z_{ij} \approx L_i \tilde{L}_j$, but these have to be made (a) symmetric and (b) tangential by adding terms that are lower-order in \mathbf{L} . The straightforward construction (Appendix A) gives

$$\begin{aligned} 2X_{ij} &= L_i L_j + L_j L_i + in_i \tilde{L}_j + in_j \tilde{L}_i , \\ 2Y_{ij} &= \tilde{L}_i \tilde{L}_j + \tilde{L}_j \tilde{L}_i - in_i \tilde{L}_j - in_j \tilde{L}_i , \\ 2Z_{ij} &= L_i \tilde{L}_j + L_j \tilde{L}_i - in_i L_j - in_j L_i . \end{aligned} \quad (29)$$

The operator K defined by (19) and its eigenvalue N_K now take on the exact values

$$K = L^2(L^2 - 2) \quad , \quad N_K = \ell(\ell+1)[\ell(\ell+1) - 2] , \quad (30)$$

consistent with the expression $\nabla^2(\nabla^2+2)$ in eqn. (9) in Ref. [19].

The formalism can be verified by constructing any tangential traceless symmetric tensor \mathbb{P} on a sphere, and then (a) evaluating $\varphi^{a,b}$ by (22); (b) inverting K to obtain $\Phi^{a,b}$; and then (c) recovering \mathbb{P} by (21).

4. Global analysis

With global data, the standard mode expansion for scalar functions leads naturally to corresponding mode expansions for vector and tensor functions. The modal characteristics allow different contributions to the polarization (e.g., cosmological versus foreground contributions) to be disentangled.

4.1. Planar case

In the case of a plane, the scalar functions $\Phi^{a,b}$ can be expanded in terms of plane waves $F_{\mathbf{k}} \propto \exp(i\mathbf{k} \cdot \mathbf{r})$:

$$\Phi^{a,b}(\mathbf{r}) = \sum_{\mathbf{k}} C_{\mathbf{k}}^{a,b} F_{\mathbf{k}}(\mathbf{r}) . \quad (31)$$

Then from (10) and (21), a vector field can be expressed in terms of $\mathbf{d}F_{\mathbf{k}}$ and $\tilde{\mathbf{d}}F_{\mathbf{k}}$ (vector planar harmonics), and a polarization field in terms of $\mathbb{D}F_{\mathbf{k}}$ and $\tilde{\mathbb{D}}F_{\mathbf{k}}$ (tensor planar harmonics). The straightforward derivation will not be shown, except to note that, acting on a plane wave, $D_1 \mapsto -(k_x^2 - k_y^2) = -k^2 \cos 2\phi$, $D_2 \mapsto -2k_x k_y = -k^2 \sin 2\phi$, where ϕ is the direction of \mathbf{k} , so

$$\mathbf{P} = \begin{pmatrix} Q \\ U \end{pmatrix} = - \sum_{\mathbf{k}} k^2 \left[C_{\mathbf{k}}^a \begin{pmatrix} \cos 2\phi \\ \sin 2\phi \end{pmatrix} + C_{\mathbf{k}}^b \begin{pmatrix} -\sin 2\phi \\ \cos 2\phi \end{pmatrix} \right] F_{\mathbf{k}}, \quad (32)$$

equivalent to eqn. (23) in Ref. [19]. Our approach is more accessible because the planar case is obtained on a stand-alone basis, and not as a limit of the spherical case. The polarization direction χ is given by $\tan 2\chi = U/Q$, and its relationship with ϕ follows trivially from (32).

4.2. Spherical case

On a sphere, the scalar functions can be expanded in spherical harmonics:

$$\Phi^{a,b} = \sum_{\ell m} C_{\ell m}^{a,b} Y_{\ell m}. \quad (33)$$

Then a vector field can be expressed in vector spherical harmonics $\mathbf{Y}_{\ell m}^{a,b}$, and a polarization field in tensor spherical harmonics $\mathbb{Y}_{\ell m}^{a,b}$:

$$\begin{pmatrix} \mathbf{Y}_{\ell m}^a \\ \mathbf{Y}_{\ell m}^b \end{pmatrix} = \frac{1}{\sqrt{\ell(\ell+1)}} \begin{pmatrix} i\tilde{\mathbf{L}} \\ -i\mathbf{L} \end{pmatrix} Y_{\ell m}, \quad (34)$$

$$\begin{pmatrix} \mathbb{Y}_{\ell m}^a \\ \mathbb{Y}_{\ell m}^b \end{pmatrix} = \frac{1}{\sqrt{2(\ell^2+\ell)(\ell^2+\ell-2)}} \begin{pmatrix} \mathbb{D} \\ \tilde{\mathbb{D}} \end{pmatrix} Y_{\ell m}, \quad (35)$$

where the prefactor in (35) is just $(2N_K)^{-1/2}$ with N_K given by (30). The vector spherical harmonics agree with well-known results [26]. The tensor harmonics start with $\ell = 2$ because, from (30), \mathbb{D} and $\tilde{\mathbb{D}}$ annihilate $\ell = 0, 1$.

5. Discussion

The representation of linear polarization fields in terms of two scalar functions involves *second-order* derivatives and cannot be likened to gradient and curl in the conventional sense. Nevertheless, a simple analogy exists formally, as shown in the simplest case by (2) and (6). The transcription of the formalism from a plane to the surface of a sphere is straightforward.

Vector and tensor fields on a sphere can be more neatly described using intrinsic coordinates (θ and ϕ). Nevertheless most physicists are more familiar with the cartesian representation (e.g., $L_x = yp_z - zp_y$). The present account has kept to the same spirit: embedding in 3-D space but enforcing tangential conditions. The vector and tensor spherical harmonics emerge naturally, without invoking the full apparatus of Newman–Penrose spin- s spherical harmonics [27, 28].

In closing, it is useful to recall the steps in the data analysis. (a) The measured polarization \mathbb{P} on a sphere is separated into two components, in some sense a generalization of Helmholtz’s theorem: from a spin-1 field to a spin-2 field, and from a plane to a sphere — the main focus of the present pedagogical exposition. (b) With global data, each polarization component can be decomposed into modes labelled by (ℓ, m) , as sketched in Section 4. (c) Finally model calculations (beyond the purpose of the present paper) would provide the different characteristics (in mode space, “map space” and frequency dependence) of the various dynamical contributions and thereby allow any *cosmological* contribution to the B mode to be isolated — which is the target of many experiments in progress.

Acknowledgments

We thank CS Chu, PT Leung, WT Ni and WM Suen for discussions. The work of EY is supported by the Kavli Foundation.

Appendix A. Lower-order corrections

The dual is defined as $\tilde{\mathbf{L}} = -\hat{\mathbf{n}} \times \mathbf{L}$, with $\hat{\mathbf{n}}$ on the left, to guarantee that $\tilde{\mathbf{L}}\Phi$ are tangential: $n_i \tilde{L}_i \Phi = 0$.

To leading order $X_{ij} \approx L_i L_j$, $Y_{ij} \approx \tilde{L}_i \tilde{L}_j$. These expressions when symmetrized

$$2X_{ij} \approx L_i L_j + L_j L_i \quad , \quad 2Y_{ij} \approx \tilde{L}_i \tilde{L}_j + \tilde{L}_j \tilde{L}_i \quad (\text{A.1})$$

are not tangential. To fix this problem, we add correction terms that are (a) symmetric and (b) lower-order in \mathbf{L} . The only possibility is a term proportional to

$$n_i \tilde{L}_j + n_j \tilde{L}_i \quad . \quad (\text{A.2})$$

All terms in (A.1) contain an even number of cross products (when expressed in terms of position and momenta), so in (A.2) we have to use \tilde{L} and not L .

The conditions $n_i X_{ij} = n_i Y_{ij} = 0$ determine the multiple of (A.2) to be added, giving the *exact* expressions in (29). A factor i appears in the subsidiary terms because \mathbf{L} and $\tilde{\mathbf{L}}$ are imaginary. Any zero-order terms $\propto n_i n_j$ would not be tangential.

Similarly,

$$2Z_{ij} \approx L_i \tilde{L}_j + L_j \tilde{L}_i \quad ,$$

in which equality is maintained to the accuracy implied by \approx . Enforcing the tangential condition gives the result in (29). In this case, all terms have an odd number of cross products, so the subsidiary terms involve L but not \tilde{L} .

The three operators \mathbb{X} , \mathbb{Y} and \mathbb{Z} are by construction symmetric and tangential, and these properties are inherited by \mathbb{D} and $\tilde{\mathbb{D}}$. Moreover, $\text{tr } \mathbb{X} = \text{tr } \mathbb{Y} = L^2$, $\text{tr } \mathbb{Z} = 0$. Thus \mathbb{D} and $\tilde{\mathbb{D}}$ are traceless. Further, $-\epsilon_{ijk} n_j (X_{km} - Y_{km}) = Z_{im}$; i.e., $\tilde{\mathbb{D}} = \mathbb{E}\mathbb{D}$ as expected.

References

- [1] A. A. Penzias and R. W. Wilson. A measurement of excess antenna temperature at 4080 Mc/s. *ApJ* **142**, 419–421 (1965). doi: 10.1086/148307
- [2] A. A. Penzias and R. W. Wilson. Measurement of the flux density of Cas A at 4080 Mc/s. *ApJ* **142**, 1149–1155 (1965). doi: 10.1086/148384
- [3] R. H. Dicke, P. J. E. Peebles, P. G. Roll, and D. T. Wilkinson. Cosmic black-body radiation. *ApJ* **142**, 414–419 (1965). doi: 10.1086/148306
- [4] D. J. Fixsen and J. C. Mather. The spectral results of the far-infrared absolute spectrophotometer instrument on COBE. *ApJ* **581**, 817–822 (2002). doi: 10.1086/344402
- [5] D. J. Fixsen. The temperature of the cosmic microwave background. *ApJ* **707**, 916 (2009). doi: 10.1088/0004-637X/707/2/916
- [6] G. F. Smoot et al. First results of the COBE satellite measurement of the cosmic microwave background radiation. *Adv. Space Res.* **11** (2), 193–205 (1991).
- [7] A. H. Guth. Inflationary universe: A possible solution to the horizon and flatness problems. *Phys. Rev. D* **23**, 347–356 (1981). doi: 10.1103/PhysRevD.23.347
- [8] A. D. Linde. A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems. *Physics Letters B*, **108**, 389–393 (1982). doi: 10.1016/0370-2693(82)91219-9
- [9] WMAP Highlights. <https://map.gsfc.nasa.gov>
- [10] A. Kogut. WMAP polarization results. *New Astronomy Reviews*, **47**, 977–986 (2003). doi: 10.1016/j.newar.2003.09.029
- [11] U. Seljak and M. Zaldarriaga. Signature of gravity waves in the polarization of microwave background. *Phys. Rev. Lett.* **78**, 2054–2057 (1997). doi: 10.1103/PhysRevLett.78.2054
- [12] M. Kamionkowski, A. Kosowsky and A. Stebbins. A probe of primordial gravity waves and vorticity. *Phys. Rev. Lett.* **78**, 2058 (1997). doi: 10.1103/PhysRevLett.78.2058
- [13] P. A. R. Ade et al. Detection of *B*-mode polarization at degree angular scales by BICEP2. *Phys. Rev. Lett.* **112**, 241101 (2014). doi: 10.1103/PhysRevLett.112.241101
- [14] BICEP2/Keck, Planck Collaboration: P. A. R. Ade et al. A Joint analysis of BICEP2/Keck array and Planck data. *Phys. Rev. Lett.* **114**, 101301 (2015). doi: 10.1103/PhysRevLett.114.101301
- [15] Planck Collaboration: Y. Akrami et al. Planck intermediate results. LIV. Polarized dust foregrounds. arXiv:1801.04945v1
- [16] For example, CMB-S4, Simons Observatory, BICEP Array, CHIME, Litebird, SPIDER2.
- [17] R. Cowen, Telescope captures view of gravitational waves: Images of the infant Universe reveal evidence for rapid inflation after the Big Bang. *Nature* **507**, Issue 7492 (2014).
<https://www.nature.com/news/telescope-captures-view-of-gravitational-waves-1.14876>
- [18] M. Kamionkowski and E. D. Kovetz. The quest for *B* modes from inflationary gravitational waves. *Annual Review of Astronomy and Astrophysics*, **54**, 227–269 (2016). doi: 10.1146/annurev-astro-081915-023433
- [19] E. F. Bunn, M. Zaldarriaga, M. Tegmark and A. de Oliveira-Costa. *E/B* decomposition of finite pixelized CMB maps. *Phys. Rev. D* **67**, 023501 (2003). doi: 10.1103/PhysRevD.67.023501
- [20] J. D. Jackson, *Classical Electrodynamics*, Second Edition. Wiley, New York. (1975). Section 7.2.
- [21] S. De and H. Tashiro. Circular polarization of the CMB: a probe of the first stars. *Phys. Rev. D* **92**, 123506 (2015). doi: 10.1103/PhysRevD.92.123506
- [22] S. King and P. Lubin. Circular polarization of the CMB: foregrounds and detection prospects. *Phys. Rev. D* **94**, 023501 (2016). doi: 10.1103/PhysRevD.94.023501
- [23] D. T. Chuss et al. Properties of a variable-delay polarization modulator. *Applied Optics* **51** (2), 197–208 (2012). doi: 10.1364/AO.51.000197
- [24] L. Padilla et al. Two-year cosmology large angular scale surveyor (CLASS) observations: A measurement of circular polarization at 40 GHz. arXiv:1911.00391v1.
- [25] H. C. Chiang et al. Measurement of cosmic microwave background polarization power spectra from

- two years of BICEP data. *ApJ* **711**, 1123–1140 (2010). doi: 10.1088/0004-637X/711/2/1123
- [26] J. D. Jackson, *Classical Electrodynamics*, Second Edition. Wiley, New York. (1975). Section 16.2.
- [27] E. T. Newman and R. Penrose. Ten exact gravitationally-conserved quantities. *Phys. Rev. Lett.* **15**, 231–233 (1965). doi: 10.1103/PhysRevLett.15.231
- [28] W. B. Campbell. Tensor and spinor spherical harmonics and the spin- s harmonics ${}_sY_{\ell m}(\theta, \phi)$. *J. Math. Phys.* **12**, 1763–1770 (1971). doi: 10.1063/1.1665802