

Preprint no. NJU-INP 011/19

## Two photon transition form factors of neutral pseudoscalar mesons

Minghui Ding\* and Daniele Binosi

*European Centre for Theoretical Studies in Nuclear Physics and Related Areas (ECT\*) and Fondazione Bruno Kessler*

*Villa Tambosi, Strada delle Tabarelle 286, I-38123 Villazzano (TN) Italy*

*\*E-mail: mding@ectstar.eu*

Khépani Raya and Lei Chang

*School of Physics, Nankai University, Tianjin 300071, China*

Craig D. Roberts

*School of Physics, Nanjing University, Nanjing, Jiangsu 210093, China*

*Institute for Nonperturbative Physics, Nanjing University, Nanjing, Jiangsu 210093, China*

Starting from the axial vector Ward-Green-Takahashi identity, and the necessity of preserving this identity when attempting to calculate the properties describing pseudoscalar mesons, we describe a Bethe-Salpeter kernel which takes into account the non-Abelian anomaly and therefore is suitable to describe the flavor mixing states  $\eta, \eta'$ . We then report results for the two photon transition form factors of  $\eta, \eta'$ , and compare them with existing experimental data.

*Keywords:* Pseudoscalar meson; Transition form factor; Non-Abelian anomaly.

### 1. Ward-Green-Takahashi identity

Ward's identity<sup>1</sup> and its generalization by Green<sup>2</sup> and Takahashi<sup>3</sup>, ought to be preserved by any approach that is willing to capture QCD's fundamental features. In its axial vector formulation this identity reads

$$P^\mu \Gamma_{5\mu}^a(k; P) = S^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a S^{-1}(k_-) - 2i\mathcal{M}^{ab} \Gamma_5^b(k; P) - \mathcal{A}^a(k; P), \quad (1)$$

where:  $P(k)$  is the total (relative) momentum between the amputated quark legs;  $\{\mathcal{F}^a | a = 0, \dots, N_f^2 - 1\}$  are the generators of  $U(N_f)$  in the fundamental representation, orthonormalized according to  $\text{tr} \mathcal{F}^a \mathcal{F}^b = \frac{1}{2} \delta^{ab}$ ;  $\mathcal{M}^{ab} = \text{tr}_F [\{\mathcal{F}^a, \mathcal{M}^{\text{bm}}\} \mathcal{F}^b]$ , with  $\mathcal{M}^{\text{bm}} = \text{diag}[m_u^{\text{bm}}, m_d^{\text{bm}}, m_s^{\text{bm}}, \dots]$  the current-quark bare masses matrix;  $\mathcal{S} = \text{diag}[S_u, S_d, S_s, \dots]$  the dressed-quark propagator matrix, with nonzero entries determined by the corresponding Dyson-Schwinger equation;  $\Gamma_{5\mu}^a$  and  $\Gamma_5^a$  the axial-vector and

pseudoscalar vertex respectively, both satisfying the corresponding inhomogeneous Bethe-Salpeter equations; and, finally,  $\mathcal{A}^a$  in an expression of the non-Abelian axial anomaly and can be defined through the topological charge density operator. Notice that only  $\mathcal{A}^{a=0}$  is nonzero: therefore, flavor-nonsinglet neutral pseudoscalar mesons satisfy the standard identity, while flavor-singlet state satisfies the anomalous one<sup>4</sup>.

One of the far-reaching consequences of the Ward-Green-Takahashi (WGT) identity (1) is a mass formula for pseudoscalar mesons which is valid irrespectively from the magnitude of the current-quark masses. It reads

$$m_M^2 f_M^a = 2 \mathcal{M}^{ab} \rho_M^b + \delta^{a0} n_M, \quad (2)$$

with:  $m_M$  the meson mass;  $f_M^a$  the decay constant;  $\rho_M^b$  the pseudoscalar projection of the meson's Bethe-Salpeter wave function onto the origin in configuration space; and  $n_M$  a quantity derived from topological charge density operator. This one single equation has a wide range of effects on Goldstone modes, excited states as well as system(s) described by the non-Abelian axial anomaly<sup>4-6</sup>; in particular, it shows that:

- The Nambu-Goldstone pion is a consequence of Dynamical Chiral Symmetry Breaking (DCSB).
- In the chiral limit, the decay constants of the pion's radial excitations are zero.
- $\eta'$  is split from the octet pseudoscalars by an amount that depends on QCD's topological susceptibility.

## 2. Kernels for the bound-state equations

Yet another consequence of Eq. (1) is that it leads to a symmetry-preserving truncation scheme, the simplest of which is the Rainbow-Ladder (RL) approximation. RL truncation is known to be accurate for ground-state light-quark vector- and isospin-nonzero-pseudoscalar-mesons<sup>7-10</sup>, and has been used in a recent study on pion parton distribution functions<sup>11</sup>. Implemented in the Bethe-Salpeter and gap equations, RL truncation is expressed via the following kernels, respectively:

$$[\mathcal{K}_N(k, q; P)]_{\ell_1 \ell_2}^{\ell_1' \ell_2'} = -\frac{4}{3} \mathcal{G}_{\mu\nu}(t) [i\gamma_\mu]_{\ell_1 \ell_1'} [i\gamma_\nu]_{\ell_2' \ell_2}, \quad (3a)$$

$$\mathcal{G}_{\mu\nu}(t = k - q) = \tilde{\mathcal{G}}(t^2) T_{\mu\nu}(t), \quad (3b)$$

where:  $t^2 T_{\mu\nu} = t^2 \delta_{\mu\nu} - t_\mu t_\nu$ ; and  $\tilde{\mathcal{G}}(t^2)$  is the interaction consistent with that determined in studies of QCD's gauge sector<sup>12</sup>.

However, RL truncation fails for the  $\eta$ - and  $\eta'$ -mesons as they do not produce vertices that satisfy the anomalous axial-vector WGT identity. Indeed, whilst the RL kernel is constrained by a large body of successful phenomenology, the form of the anomalous kernel is unknown. Considering the structure of the non-Abelian anomaly, it readily becomes apparent that no related contribution to the Bethe-Salpeter kernel can contain external quark or antiquark lines which are simply connected to the internal lines: purely gluonic configurations must mediate. Moreover, no finite sum of diagrams can be sufficient<sup>4</sup>. On general grounds, its contribution to the Bethe-Salpeter equation for pseudoscalar mesons must take the following form<sup>13</sup>:

$$[\mathcal{K}_A(k, q; P)]_{\ell_1 \ell_2}^{\ell'_1 \ell'_2} = - \sum_{i=1}^4 \sum_{f=l,s} a_i^f(k, q; P) \times [\mathcal{F}^f \mathcal{D}_i(q; P)]_{\ell'_2 \ell'_1} [\mathcal{F}^f \mathcal{D}_i(k; P)]_{\ell_1 \ell_2}, \quad (4)$$

where:  $\{\mathcal{D}_i(k; P) | i = 1, \dots, 4\}$  constitute a Dirac basis of the pseudoscalar meson Bethe-Salpeter wave function; and  $\{a_i^f(k, q; P) | i = 1, \dots, 4; f = l, s\}$  are scalar functions. An educated Ansatz for  $\mathcal{K}_A$  had been put forward in Ref. 13, which being essentially nonperturbative in nature, provided material support at infrared momenta. In this way the corresponding Bethe-Salpeter kernel becomes the sum of two terms:  $\mathcal{K} = \mathcal{K}_V + \mathcal{K}_A$ , and with the bound-state kernel thus defined, results for the  $\gamma^* \gamma \rightarrow \eta, \eta'$  transition form factors have been obtained.

### 3. Two-photon transition form factors

The two-photon transition form factor provides a window for exploring the internal structure of neutral pseudoscalar meson<sup>14–16</sup>. The parameters in the gap and Bethe-Salpeter equations can be fixed by requiring that the solutions of the coupled-channels bound-state problems reproduce the empirical  $\eta, \eta'$  masses and the four phenomenologically-determined values of the light- and strange-quark  $\eta, \eta'$  decay constants<sup>13</sup>. We find  $m_\eta = 0.56$  GeV and  $m_{\eta'} = 0.96$  GeV, to be compared with the experimental values<sup>17</sup> 0.55 GeV and 0.96 GeV, respectively; other observables like decay constants, mixing angle and two-photon decay widths are reported in Ref. 13.

The transition form factors  $F_{\eta, \eta'}(Q^2)$  obtained within the framework sketched above, are shown in Fig. 1 and compared with the asymptotic value of the  $\pi^0$  form factor,  $2f_\pi = 0.186$  GeV, drawn as the dotted (red) curve in both panels. There are marked similarities between  $F_\eta(Q^2)$  and the  $\pi^0$  tran-




Fig. 1.  $\gamma^*\gamma \rightarrow \eta, \eta'$  transition form factors: left panel,  $\eta$ ; right panel,  $\eta'$ . Curves: solid (green), our prediction with complete evolution, with the shaded (green) band indicating an uncertainty in our prediction owing to variations in the value of  $f_\eta^s$ ; dashed (blue), asymptotic limit, with the banded-shading (blue) region indicating the impact of uncertainty in  $f_\eta^s$  on the asymptotic behaviour. The broader, shaded (grey) band combines this with the uncertainty owing to omission of  $\varphi_M^0$ - $\varphi_M^g$  mixing ( $\varphi_M^0$  is the flavor singlet distribution amplitude and  $\varphi_M^g$  is the two-gluon distribution amplitude). In both panels, the dotted (red) curve is the  $\pi^0$  asymptotic limit,  $2f_\pi$ . Experimental data are<sup>18</sup>: diamonds (blue) “CLEO”; circles (red) “BaBar”.

sition form factor (see, for example, Ref. 14, Fig. 2); furthermore, our full  $F_{\eta, \eta'}(Q^2)$  result (solid, green curve) agree well with existing data. Finally, at asymptotically large momentum transfers, *i.e.* on  $\tau^2 := \Lambda_{\text{QCD}}^2/Q^2 \simeq 0$ , our full results meet the asymptotic trajectory (dashed blue curve).

The scope of the analysis sketched herein could also be extended to include the doubly off-shell transition form factors, for which the first data  $\gamma^*(k_1)\gamma^*(k_2) \rightarrow \eta'$  now exist<sup>19</sup>.

*Acknowledgments:* Work partially supported by Jiangsu Province *Hundred Talents Plan for Professionals*.

## References

1. J. C. Ward, An Identity in Quantum Electrodynamics, *Phys. Rev.* **78**, 182 (1950).
2. H. S. Green, A Pre-renormalized quantum electrodynamics, *Proc. Phys. Soc.* **A66**, 873-880 (1953).
3. Y. Takahashi, On the generalized Ward identity, *Nuovo Cim.* **6**, 371 (1957).
4. M. S. Bhagwat, L. Chang, Y. X. Liu, C. D. Roberts, and P. C. Tandy, Flavour symmetry breaking and meson masses, *Phys. Rev.*

- C76**, 045203 (2007).
5. P. Maris, C. D. Roberts, and P. C. Tandy, Pion mass and decay constant, *Phys. Lett.* **B420**, 267-273 (1998).
  6. A. Holl, A. Krassnigg, C. D. Roberts, and S. V. Wright, On the complexification of pseudoscalar mesons, *Int. J. Mod. Phys.* **A20**, 1778-1784 (2005).
  7. T. Horn and C. D. Roberts, The pion: an enigma within the Standard Model, *J. Phys.* **G43**, no.7, 073001 (2016).
  8. G. Eichmann, H. Sanchis-Alepuz, R. Williams, R. Alkofer, and C. S. Fischer, Baryons as relativistic three-quark bound states, *Prog. Part. Nucl. Phys.* **91**, 1-100 (2016).
  9. C. S. Fischer, QCD at finite temperature and chemical potential from DysonSchwinger equations, *Prog. Part. Nucl. Phys.* **105**, 1-60 (2019).
  10. V. D. Burkert, and C. D. Roberts, Colloquium : Roper resonance: Toward a solution to the fifty year puzzle, *Rev. Mod. Phys.* **91**, 011003 (2019).
  11. M. Ding, K. Raya, D. Binosi, L. Chang, C. D. Roberts and S. M. Schmidt, Symmetry, symmetry breaking, and pion parton distributions, (arXiv:1905.05208 [nucl-th]).
  12. D. Binosi, L. Chang, J. Papavassiliou and C. D. Roberts, Bridging a gap between continuum-QCD and ab initio predictions of hadron observables, *Phys. Lett.* **B742**, 183-188 (2015).
  13. M. Ding, K. Raya, A. Bashir, D. Binosi, L. Chang, M. Chen and C. D. Roberts,  $\gamma^*\gamma \rightarrow \eta, \eta'$  transition form factors, *Phys. Rev.* **D99**, 014014 (2019).
  14. K. Raya *et al.*, Structure of the neutral pion and its electromagnetic transition form factor, *Phys. Rev.* **D93**, 074017 (2016).
  15. K. Raya, M. Ding, A. Bashir, L. Chang, and C. D. Roberts, Partonic structure of neutral pseudoscalars via two photon transition form factors, *Phys. Rev.* **D95**, 074014 (2017).
  16. J. Chen, M. Ding, L. Chang and Y. X. Liu, Two Photon Transition Form Factor of  $\bar{c}c$  Quarkonia, *Phys. Rev.* **D95**, 016010 (2017).
  17. M. Tanabashi *et al.*, Review of Particle Physics, *Phys. Rev.* **D98**, 030001 (2018).
  18. P. del Amo Sanchez *et al.*, Measurement of the  $\gamma\gamma^* \rightarrow \eta$  and  $\gamma\gamma^* \rightarrow \eta'$  transition form factors, *Phys. Rev.* **D84**, 052001 (2011).
  19. J. P. Lees *et al.*, Measurement of the  $\gamma^*\gamma^* \rightarrow \eta'$  transition form factor, *Phys. Rev.* **D98**, 112002 (2018).