

INFERENCE UNDER RANDOM LIMIT BOOTSTRAP MEASURES*

GIUSEPPE CAVALIERE^{a,b} AND ILIYAN GEORGIEV^{a,c}

November 30, 2019

ABSTRACT

Asymptotic bootstrap validity is usually understood as consistency of the distribution of a bootstrap statistic, conditional on the data, for the unconditional limit distribution of a statistic of interest. From this perspective, randomness of the limit bootstrap measure is regarded as a failure of the bootstrap. We show that such limiting randomness does not necessarily invalidate bootstrap inference if validity is understood as control over the frequency of correct inferences in large samples. We first establish sufficient conditions for asymptotic bootstrap validity in cases where the unconditional limit distribution of a statistic can be obtained by averaging a (random) limiting bootstrap distribution. Further, we provide results ensuring the asymptotic validity of the bootstrap as a tool for conditional inference, the leading case being that where a bootstrap distribution estimates consistently a conditional (and thus, random) limit distribution of a statistic. We apply our framework to several inference problems in econometrics, including linear models with possibly non-stationary regressors, functional CUSUM statistics, conditional Kolmogorov-Smirnov specification tests, the ‘parameter on the boundary’ problem and tests for constancy of parameters in dynamic econometric models.

KEYWORDS: Bootstrap, random measures, weak convergence in distribution, asymptotic inference.

* We thank Matias Cattaneo, Graham Elliott, Michael Jansson, Sren Johansen, Ye Lu, Adam McCloskey, Marcelo Moreira, Ulrich Mller, Rasmus Sndergaard Pedersen, Anders Rahbek, Mervyn Silvapulle, Michael Wolf, for comments and suggestions. We also thank seminar participants at Boston U, Cambridge, ETH, Exeter, FGV Sao Paulo, Geneve, Hitotsubashi, Kyoto, LSE, Melbourne, Oxford, PUC Rio, Queen’s, QUT, SMU, Tilburg, UCL, UPF, Verona, Vienna, as well as participants to the 7th Italian Congress of Econometrics and Empirical Economics (ICEEE), the 14th SETA meeting and the NBER/NSF 2018 Time Series Conference. Financial support from the University of Bologna, ALMA IDEA 2017 grants, and from the Danish Council for Independent Research (DSF Grant 7015-00028), is gratefully acknowledged. Address correspondence to: Giuseppe Cavaliere, Department of Economics, University of Bologna, Piazza Scaravilli 2, 40126 Bologna, Italy; email: giuseppe.cavaliere@unibo.it.

^a Department of Economics, University of Bologna, Italy.

^b Department of Economics, Exeter Business School, UK.

^c Nova School of Business and Economics, Universidade Nova de Lisboa, Portugal.

1 INTRODUCTION

CONSIDER A DATA SAMPLE D_n of size n and a statistic $\tau_n := \tau_n(D_n)$, say a test statistic or a parameter estimator, possibly normalized. Interest is in a distributional approximation of τ_n . Let a bootstrap procedure generate a bootstrap analogue τ_n^* of τ_n ; i.e., computed on a bootstrap sample. Assume that τ_n converges in distribution to a non-degenerate random variable [rv], say τ . In classic bootstrap inference, asymptotic bootstrap validity is usually understood and established as convergence in probability (or almost surely) of the cumulative distribution function [cdf] of the bootstrap statistic τ_n^* conditional on the data D_n , say F_n^* , to the unconditional cdf of τ , say F . This convergence, along with continuity of F , implies by Polya's theorem that $\sup_{x \in \mathbb{R}} |F_n^*(x) - F(x)| \rightarrow 0$, in probability (or almost surely).

In many applications, however, the bootstrap statistic τ_n^* may possess, conditionally on the data, a *random* limit distribution. Cases of random bootstrap limit distributions appear in various areas of econometrics and statistics; for instance, they are documented for infinite-variance processes (Athreya, 1987; Knight, 1989; Aue, Berkes and Horváth, 2008; Cavaliere, Georgiev and Taylor, 2016), time series with unit roots (Basawa, Mallik, McCormick, Reeves and Taylor, 1991; Cavaliere, Nielsen and Rahbek, 2015), parameters on the boundary of the parameter space (Andrews, 2000), subsample inference based on fixed- b asymptotics (Shao and Politis, 2013), cube-root consistent estimators (Sen, Banerjee and Woodroffe, 2010; Cattaneo, Jansson and Nagasawa, 2017), Hodges-LeCam superefficient estimators (Beran, 1997). In most of these cases, the occurrence of a random limit distribution for the bootstrap statistic τ_n^* given the data – in contrast to a non-random limit of the unconditional distribution of the original statistic τ_n – is taken as evidence of failure of the bootstrap.

In this paper we show that randomness in the limiting distribution of a bootstrap statistic need not invalidate bootstrap inference. On the contrary, although the bootstrap no longer estimates the limiting unconditional distribution of the statistic of interest, it may still deliver hypothesis tests (or confidence intervals) with the desired null rejection probability (or coverage probability) when the sample size diverges. This happens because asymptotic control over the frequency of wrong inferences can be guaranteed by the asymptotic distributional uniformity of the bootstrap p -values, which in its turn can occur without the convergence in probability (or almost surely) of the bootstrap cdf F_n^* of τ_n^* to the asymptotic cdf F of τ .

Therefore, instead of assessing bootstrap validity in terms of the convergence of F_n^* to F , in cases where the limit of the bootstrap distribution is random we study bootstrap validity in terms of the property of asymptotic distributional uniformity of bootstrap p -values. Specifically, let p_n^* denote the bootstrap p -value, usually defined as $p_n^* := F_n^*(\tau_n)$. We define ‘bootstrap validity’ or ‘unconditional bootstrap validity’ the fact that

$$P(p_n^* \leq q) \rightarrow q \tag{1.1}$$

for $q \in (0, 1)$. The focus on this property is not new in the literature on bootstrap and simulation-based inference (see, e.g., Hansen, 1996, and Lockhart, 2012, among others).

Our first set of results provides sufficient conditions for bootstrap validity in the sense of (1.1) in situations where the bootstrap distribution is random in the limit.

Classic results for bootstrap validity when the limit bootstrap measure is not random can be obtained as special cases. The main requirement in our results is that the unconditional limit distribution of τ_n should be an average of the random limit distribution of τ_n^* given the data.

It is often the case that bootstrap validity can be addressed through the lens of a conditioning argument. In this regard, our second set of results concerns the possibility that, for a sequence of random elements X_n , it holds that the bootstrap p -value is uniformly distributed in large sample *conditionally* on X_n :

$$P(p_n^* \leq q | X_n) \xrightarrow{P} q \tag{1.2}$$

for $q \in (0, 1)$. This property, that we define ‘bootstrap validity conditional on X_n ’, implies unconditional validity in the sense of (1.1). Moreover, conditional bootstrap validity given X_n implies that the bootstrap replicates asymptotically the property of conditional tests and confidence intervals to have, conditionally on X_n , constant null rejection probability and coverage probability, respectively (for further roles of conditioning in inference, like the relevance of the drawn inferences and information recovery, see Reid, 1995, and the references therein). The leading case where we show (1.2) to hold – under regularity conditions that will be discussed in the paper – is that where the (random) limit of the conditional distribution of τ_n given X_n matches the (random) limit distribution of the bootstrap statistic. A property like (1.2) was initially established by LePage and Podgorski (1996) for permutation tests in location models. Their approach has not been developed further in the bootstrap literature, in particular because it requires probabilistic tools that are not widely popular in this field.

When dealing with random limit distributions, the usual convergence concept employed to establish bootstrap validity, i.e. weak convergence in probability, can only be employed in some very special cases. Instead, our formal discussion makes extensive use of the probabilistic concept of weak convergence of random measures; see e.g. Kallenberg (2017, Ch.4). To our knowledge, in the bootstrap context this concept has so far been mostly used to obtain negative results of lack of validity for specific bootstrap procedures (as e.g. in Knight, 1989, and Basawa et al., 1991), rather than positive validity results, as we do here. As an ingredient of our analysis, we also present some novel results on the weak convergence of conditional expectations.

To illustrate the practical relevance of our results, we initially present them by using a simple linear model with either stationary or non-stationary regressors, and later we analyze four well-known cases in the econometric literature where the bootstrap features a random limit distribution. The first is a standard CUSUM-type test of the i.i.d. property for a random sequence with infinite variance. This is a case where the limit distribution of the CUSUM statistic depends on unknown nuisance parameters (e.g., the tail index) and bootstrap or permutation tests fail to estimate this distribution consistently. We argue that a simple bootstrap based on permutations, albeit having a random limit distribution and hence being invalid in the usual sense, provides *exact* conditional inference and hence is also unconditionally valid in the sense of (1.1). The second application considers a Kolmogorov-Smirnov-type test for correct specification of the conditional distribution of a response variable given a vector of covariates. Andrews (1997) considers a parametric bootstrap implementation where the covariates are kept

fixed across bootstrap samples. While in the independent case the limit of the bootstrap distribution is non-random, this is not the case in general. Using our theory we discuss conditions for validity of the bootstrap within this framework. The third application is the implementation of the bootstrap in ‘parameter on the boundary’ problems (Andrews, 1999,2000). Taking hypothesis testing in a predictive regression framework as an illustration, we show that the kind of randomness in the limit distribution of the bootstrap statistics of interest depends, when the true parameter lies on the boundary of the parameter space, on how well the mutual position of the boundary, the set identified by the null hypothesis and the true parameter value, is approximated in the bootstrap world. Although the standard bootstrap may fail to be valid in the sense of (1.1), we provide conditions for the validity of alternative bootstrap schemes. The fourth application includes an analysis of the much applied bootstrap ‘sup F ’ tests of parameter constancy in regression models where the design matrix could be random but be conditioned upon; see Hall (1992, p.170). In the resampling process forming the bootstrap sample, it appears natural to take the design matrix as fixed across the bootstrap repetitions. Under a set of assumptions proposed by Hansen (2000), we argue that the fixed-regressor bootstrap ‘sup F ’ statistic has a random limit distribution, thus invalidating previous claims in the literature that the bootstrap is consistent for the unconditional limit distribution of the original ‘sup F ’ test statistic. We then provide conditions under which the fixed-regressor bootstrap is unconditionally valid and, additionally, valid conditionally on the chosen set of regressors.

STRUCTURE OF THE PAPER

The paper is organized as follows. In Section 2 we outline the main concepts and ideas using a simple linear regression model. Our main theoretical results are presented in Section 3. Section 4 contains the four applications of the theory, whereas Section 5 concludes. The paper has two Appendices. In Appendix A we collect some results on weak convergence in distribution which are useful to prove our main theorems and develop the applications. Appendix B contains the proofs of all theory results given in the paper. The proofs of the results from Appendix A and some additional material are collected in the accompanying supplement, Cavaliere and Georgiev (2019).

NOTATION AND DEFINITIONS

We use the following notation throughout. The spaces of càdlàg functions $[0, 1] \rightarrow \mathbb{R}^n$, $[0, 1] \rightarrow \mathbb{R}^{m \times n}$ and $\mathbb{R} \rightarrow \mathbb{R}$ (all equipped with the respective Skorokhod J_1 -topologies; see Kallenberg, 1997, Appendix A2), are denoted by \mathcal{D}_n , $\mathcal{D}_{m \times n}$ and $\mathcal{D}(\mathbb{R})$, respectively; for the first one, when $n = 1$ the subscript is suppressed. Integrals are over $[0, 1]$ unless otherwise stated, Φ is the standard Gaussian cdf, $U(0, 1)$ is the uniform distribution on $[0, 1]$ and $\mathbb{I}_{\{\cdot\}}$ is the indicator function. If F is a (random) cdf, F^{-1} stands for the right-continuous generalized inverse, i.e., $F^{-1}(u) := \sup\{v \in \mathbb{R} : F(v) \leq u\}$, $u \in \mathbb{R}$. Unless differently specified, limits are for $n \rightarrow \infty$.

Polish (i.e., complete and separable metric) spaces are always equipped with their Borel σ -algebras. Throughout, we assume that all random elements are Polish-space

valued and that well-defined conditional distributions exist. For random elements of a Polish space, the existence of regular conditional distributions is guaranteed and we assume without loss of generality that conditional probabilities are regular (Kallenberg, 1997, Theorem 5.3). Equality of conditional distributions is understood in the almost sure [a.s.] sense and, for random cdf's as random elements of $\mathcal{D}(\mathbb{R})$, equalities are up to indistinguishability.

Let $\mathcal{C}_b(\mathcal{S})$ be the set of all continuous and bounded real-valued functions on a metric space \mathcal{S} . For random elements Z, Z_n ($n \in \mathbb{N}$) of a metric space \mathcal{S}_Z , we employ the usual notation $Z_n \xrightarrow{w} Z$ for the property that the distribution of Z_n weakly converges to the distribution of Z , defined by the convergence $E\{g(Z_n)\} \rightarrow E\{g(Z)\}$ for all $g \in \mathcal{C}_b(\mathcal{S}_Z)$. For random elements $(Z, X), (Z_n, X_n)$ of the metric spaces $\mathcal{S}_Z \times \mathcal{S}$ and $\mathcal{S}_Z \times \mathcal{S}_n$ ($n \in \mathbb{N}$), and defined on a common probability space, we denote by $Z_n|X_n \xrightarrow{w_p} Z|X$ (resp. $Z_n|X_n \xrightarrow{w_{a.s.}} Z|X$) the fact that $E\{g(Z_n)|X_n\} \rightarrow E\{g(Z)|X\}$ in probability (resp. a.s.) for all $g \in \mathcal{C}_b(\mathcal{S}_Z)$. In the special case where $E\{g(Z_n)|X_n\} \xrightarrow{w} E\{g(Z)\}$ in probability (resp. a.s.) for all $g \in \mathcal{C}_b(\mathcal{S}_Z)$, we write $Z_n|X_n \xrightarrow{w_p} Z$ (resp. $Z_n|X_n \xrightarrow{w_{a.s.}} Z$). In such a case the weak limit (in probability or a.s.) of the random conditional distribution $Z_n|X_n$ is the non-random distribution of Z , thus reducing our definition to the one of weak convergence in probability (resp. a.s.) usually employed in the bootstrap literature.

In order to deal with random limit measures, we need a further convergence concept. For $(Z, X), (Z_n, X_n)$ ($n \in \mathbb{N}$) defined on possibly different probability spaces, we denote by $Z_n|X_n \xrightarrow{w_w} Z|X$ the fact that $E\{g(Z_n)|X_n\} \xrightarrow{w} E\{g(Z)|X\}$ for all $g \in \mathcal{C}_b(\mathcal{S}_Z)$ and label it 'weak convergence in distribution'. It coincides with the probabilistic concept of weak convergence of random measures (here, of the random conditional distributions $Z_n|X_n$; see Kallenberg, 2017, Ch.4). Whenever Z_n and Z are rv's and the conditional distribution of Z given X is *diffuse* (non-atomic), this is equivalent to the weak convergence $P(Z_n \leq \cdot | X_n) \xrightarrow{w} P(Z \leq \cdot | X)$ of the random cdf's as random elements of $\mathcal{D}(\mathbb{R})$ (see Kallenberg, 2017, Theorem 4.20). Finally, on probability spaces where both the data D_n and the auxiliary variates used in the construction of the bootstrap data are defined, we use $Z_n \xrightarrow{w_p^*} Z|X$ (resp. $\xrightarrow{w_{a.s.}^*}, \xrightarrow{w_w^*}$) interchangeably with $Z_n|D_n \xrightarrow{w_p} Z|X$ (resp. $\xrightarrow{w_{a.s.}}, \xrightarrow{w_w}$), and write $P^*(\cdot)$ for $P(\cdot|D_n)$.

2 A LINEAR REGRESSION EXAMPLE

In this section we provide an overview of the main results established in the sections below, and the concepts employed, by using a simple linear regression model. Further applications will be given in Section 4. We observe that even for this basic model bootstrap statistics may have a random limit distribution. Then, we show that convergence of the bootstrap statistic to a random limit may imply (asymptotic) bootstrap validity in the unconditional sense of eq. (1.1). Finally, we illustrate the possibility that bootstrap inference may have a conditional interpretation.

2.1 MODEL, BOOTSTRAP AND RANDOM LIMIT BOOTSTRAP MEASURES

Assume that the data are given by $D_n := \{y_t, x_t\}_{t=1}^n$ and consider the linear model

$$y_t = \beta x_t + \varepsilon_t \quad (t = 1, 2, \dots, n) \quad (2.1)$$

where x_t, y_t are scalar rv's and ε_t are unobservable zero-mean errors with $\omega_\varepsilon := \text{Var}(\varepsilon_t) \in (0, \infty)$, $t = 1, \dots, n$. Assume that $M_n := \sum_{t=1}^n x_t^2 > 0$ a.s. for all n ; further assumptions will be introduced gradually. Interest is in inference on β based on $T_n := \hat{\beta} - \beta$, with $\hat{\beta}$ the OLS estimator of β ; for instance, a confidence interval or a test of a null hypothesis of the form $H_0 : \beta = 0$.

The classic (parametric) fixed-design bootstrap, see e.g. Hall (1992), entails generating a bootstrap sample $\{y_t^*, x_t\}_{t=1}^n$ as

$$y_t^* = \hat{\beta} x_t + \hat{\omega}_\varepsilon^{1/2} \varepsilon_t^* \quad (t = 1, 2, \dots, n) \quad (2.2)$$

where $\{\varepsilon_t^*\}_{t=1}^n$ are i.i.d. $N(0, 1)$, independent of the original data, and $\hat{\omega}_\varepsilon$ is an estimator of ω_ε , e.g., the residual variance $n^{-1} \sum_{t=1}^n (y_t - \hat{\beta} x_t)^2$. The OLS estimator of β from the bootstrap sample is denoted by $\hat{\beta}^*$ and, conditionally on the original data, $T_n^* := \hat{\beta}^* - \hat{\beta} \sim N(0, \hat{\omega}_\varepsilon M_n^{-1})$. As is standard, the distribution of T_n is approximated by the distribution of T_n^* conditional on the data. With F_n^* denoting the cdf of T_n^* under P^* (the probability measure induced by the bootstrap; i.e., conditional on the original data), the bootstrap p -value is given by $p_n^* := F_n^*(T_n)$.

REMARK 2.1 A special case where the ensuing bootstrap inference is exact in finite samples, such that p_n^* is uniformly distributed for finite n , obtains when the original ε_t 's are $N(0, \omega_\varepsilon)$, independent of $X_n := \{x_t\}_{t=1}^n$, and ω_ε is known to the econometrician (hence $\hat{\omega}_\varepsilon = \omega_\varepsilon$). Then the conditional distribution of T_n^* given the data D_n equals the distribution of the original statistic T_n conditional on the regressor X_n (equivalently, on the ancillary statistic M_n): $T_n^* | D_n \stackrel{d}{=} T_n | X_n \sim N(0, \omega_\varepsilon M_n^{-1}) | M_n$. Put differently,

$$F_n^*(u) := P(T_n^* \leq u | D_n) = P(T_n \leq u | X_n) = \Phi(\omega_\varepsilon^{-1/2} M_n^{1/2} u), \quad u \in \mathbb{R}.$$

Then, as $\omega_\varepsilon^{-1/2} M_n^{1/2} T_n | M_n \sim N(0, 1)$, it is straightforward that in this special case bootstrap inference is exact: $p_n^* = F_n^*(T_n) = \Phi(\omega_\varepsilon^{-1/2} M_n^{1/2} T_n) \stackrel{d}{=} \Phi(N(0, 1)) \sim U(0, 1)$, and that this result also holds conditionally on M_n : $p_n^* | M_n \sim U(0, 1)$. \square

Although bootstrap inference is not exact in general, it may be still be asymptotically valid. To show this, we distinguish between the cases of a stationary and a non-stationary regressor x_t . It is the second case that anticipates the main results of the paper. We assume $\hat{\omega}_\varepsilon \xrightarrow{P} \omega_\varepsilon$ throughout.

2.1.1 CLASSIC BOOTSTRAP VALIDITY WHEN THE REGRESSOR IS STATIONARY

Suppose initially that $\{x_t\}_{t \in \mathbb{N}}$ is weakly stationary and $n^{-1} M_n \xrightarrow{P} M := E x_1^2 > 0$. Define $\tau_n := n^{1/2}(\hat{\beta} - \beta)$ and $\tau_n^* := n^{1/2}(\hat{\beta}^* - \hat{\beta})$; the bootstrap p -values based on (τ_n, τ_n^*)

and (T_n, T_n^*) are identical. The distribution of the bootstrap statistic τ_n^* conditional on the original data D_n satisfies

$$P^*(\tau_n^* \leq u) = \Phi(n^{-1/2}\hat{\omega}_\varepsilon^{-1/2}M_n^{1/2}u) \xrightarrow{p} \Phi(\omega_\varepsilon^{-1/2}M^{1/2}u), \quad u \in \mathbb{R}. \quad (2.3)$$

Hence, $\tau_n^* \xrightarrow{w^*} \tau \sim N(0, \omega_\varepsilon M^{-1})$ and the limit distribution is non-random.

If the initial assumptions are strengthened such that a central limit theorem [CLT] holds for $\{x_t \varepsilon_t\}_{t \in \mathbb{N}}$; that is, $n^{-1/2} \sum_{t=1}^n x_t \varepsilon_t \xrightarrow{w} N(0, \omega_\varepsilon M)$, then it also holds that $\tau_n \xrightarrow{w} \tau \sim N(0, \omega_\varepsilon M^{-1})$. Hence, the bootstrap distribution of τ_n^* consistently estimates the unconditional limit distribution of τ_n in the usual sense that $\sup_{u \in \mathbb{R}} |P^*(\tau_n^* \leq u) - P(\tau \leq u)| \xrightarrow{p} 0$, by Polya's theorem. As the limit cdf is continuous, the p -value p_n^* associated with (τ_n, τ_n^*) is asymptotically uniformly distributed; i.e., (1.1) holds.

2.1.2 RANDOM LIMIT BOOTSTRAP MEASURES WHEN THE REGRESSOR IS NON-STATIONARY

Suppose now that $\{x_t\}_{t \in \mathbb{N}}$ is such that, for some constant α , $n^{-\alpha}M_n \xrightarrow{w} M$, with $M > 0$ a.s. having a non-degenerate distribution. A well-known special case is that where x_t is a finite-variance random walk and $\alpha = 2$. Redefine $\tau_n := n^{\alpha/2}(\hat{\beta} - \beta)$ and $\tau_n^* := n^{\alpha/2}(\hat{\beta}^* - \hat{\beta})$; bootstrap p -values remain unchanged. Now the bootstrap distribution of τ_n^* , conditional on the data, remains random in the limit. Specifically, by the continuous mapping theorem [CMT],

$$P^*(\tau_n^* \leq u) = \Phi(n^{-\alpha/2}\hat{\omega}_\varepsilon^{-1/2}M_n^{1/2}u) \xrightarrow{w} \Phi(\omega_\varepsilon^{-1/2}M^{1/2}u), \quad u \in \mathbb{R}, \quad (2.4)$$

which is a random cdf. In terms of weak convergence in distribution, this amounts to

$$\tau_n^* \xrightarrow{w^*} N(0, \omega_\varepsilon M^{-1}) | M. \quad (2.5)$$

As a result, with τ_n^* and M generally defined on different probability spaces, weak convergence in probability of τ_n^* does not occur. Moreover, whatever the (unconditional) limit distribution of τ_n is, provided that it exists, $P(\tau_n \leq u)$, $u \in \mathbb{R}$, will tend to a deterministic cdf. Therefore, the bootstrap cannot estimate consistently the limit distribution of τ_n and it cannot hold that $\sup_{u \in \mathbb{R}} |P^*(\tau_n^* \leq u) - P(\tau \leq u)| \xrightarrow{p} 0$. Nevertheless, bootstrap inference need not become meaningless, as it may even be exact (see Remark 2.1). We proceed, therefore, to identify in what sense bootstrap inference could remain meaningful.

2.2 BOOTSTRAP VALIDITY

Within the framework of the linear regression model, we discuss two concepts of bootstrap validity in the case of a random limit bootstrap measure. These are employed to interpret the bootstrap as a tool for unconditional or conditional inference.

2.2.1 UNCONDITIONAL BOOTSTRAP VALIDITY

Under the assumption in Section 2.1.2, consider the random-walk special case, where $x_t := \sum_{s=1}^t \eta_s$ with $e_t := (\varepsilon_t, \eta_t)'$ forming a stationary, ergodic and conditionally

homoskedastic martingale difference sequence [mds] with p.d. variance matrix $\Omega := \text{diag}\{\omega_\varepsilon, \omega_\eta\}$.¹ Then, for $\beta \neq 0$ eq. (2.1) is an instance of a cointegration regression. It holds that $(n^{-1/2} \sum_{t=1}^{\lfloor n \rfloor} e_t', n^{-1} \sum_{t=1}^n x_{t-1} \varepsilon_t) \xrightarrow{w} (B_\varepsilon, B_\eta, \int B_\eta dB_\varepsilon)$ in $\mathcal{D}_2 \times \mathbb{R}$, where $(B_\varepsilon, B_\eta)'$ is a bivariate Brownian motion with covariance matrix Ω ; see Theorem 2.4 of Chan and Wei (1988). Moreover, $n^{-2} M_n \xrightarrow{w} M := \int B_\eta^2$ by the CMT, jointly with the convergence to a stochastic integral above, so that the assumption in Section 2.1.2 holds with $\alpha = 2$ and

$$\tau_n := n(\hat{\beta} - \beta) \xrightarrow{w} \left(\int B_\eta^2 \right)^{-1} \int B_\eta dB_\varepsilon \sim N(0, \omega_\varepsilon M^{-1}), \quad (2.6)$$

the limit being (by independence of B_η and B_ε) a variance mixture of normals, with mixing variable M^{-1} and cdf $\int_{\mathbb{R}} \Phi(\omega_\varepsilon^{-1/2} M^{1/2} u) dP(M)$.

A comparison between the limit distributions of τ_n^* and τ_n , resp. in (2.5) and (2.6), shows that the bootstrap mimics a component of the mixture limit distribution of τ_n , since the limit distribution of τ_n can be recovered by integrating over M the conditional limit distribution of τ_n^* given the data. This turns out to be sufficient for bootstrap unconditional validity in the sense of eq. (1.1). A direct argument is as follows: the bootstrap p -value $p_n^* := P^*(\tau_n^* \leq \tau_n)$ satisfies, by the CMT,

$$\begin{aligned} p_n^* &= \Phi(\hat{\omega}_\varepsilon^{-1/2} M_n^{1/2} (\hat{\beta} - \beta)) \xrightarrow{w} \Phi((\omega_\varepsilon \int B_\eta^2)^{-1/2} \int B_\eta dB_\varepsilon) \\ &\stackrel{d}{=} \Phi(N(0, 1)) \sim U(0, 1). \end{aligned} \quad (2.7)$$

Thus, when inference on β is based on the distribution of τ_n^* conditional on the data, the large-sample frequency of wrong inferences can be controlled.

2.2.2 CONDITIONAL BOOTSTRAP VALIDITY

In the case of unconditional bootstrap validity, it may be possible to find an interpretation of bootstrap inference as also valid in the sense of (1.2), *i.e.* conditionally on some X_n defined on the probability space of the original data D_n (for instance, but not necessarily, the regressor $X_n := \{x_t\}_{t=1}^n$).

In the linear regression case considered here, conditional bootstrap validity with respect to the regressor X_n can be obtained under a tightening of our previous assumptions such that the invariance principle $n^{-1/2} \sum_{t=1}^{\lfloor n \rfloor} e_t \xrightarrow{w} (B_\varepsilon, B_\eta)'$ holds *conditionally* (on X_n for finite n and on B_η in the limit, in the sense of weak convergence in distribution). A sufficient condition for the conditional invariance principle is that, additionally to the assumptions on e_t in Section 2.2.1, ε_t is an mds with respect to $\mathcal{G}_t = \sigma(\{\varepsilon_s\}_{s=-\infty}^t \cup \{\eta_s\}_{s \in \mathbb{Z}})$, and that $n^{-1} \sum_{t=1}^n E(\varepsilon_t^2 | \{\eta_s\}_{s \in \mathbb{Z}}) \rightarrow \omega_\varepsilon$ a.s. (see the proof of Theorem 2 in Rubshtein, 1996). Then, by using Theorem 3 of Georgiev, Harvey, Leybourne and Taylor (2018), it follows that

$$\tau_n | X_n \xrightarrow{w} N(0, \omega_\varepsilon M^{-1}) | M,$$

¹Non-diagonal Ω could be handled by augmenting the estimated regression with Δx_t (as we do in section 4.3), leading to no qualitative differences from the case of diagonal Ω .

which compared to (2.5) shows that the distribution of τ_n^* conditional on the data estimates consistently the random limit distribution of τ_n conditional on the regressor X_n . This fact is stated more precisely in Remark 3.12 where it is concluded that $p_n^*|X_n \xrightarrow{w}_p U(0, 1)$, *i.e.*, the bootstrap is valid conditionally on the regressor.

2.2.3 A NUMERICAL ILLUSTRATION

The result in Section 2.2.2 implies that unconditional bootstrap validity can sometimes be established by means of a conditioning argument; for example, by showing validity conditional on the regressor X_n . To illustrate, in Figure 1, panels (a) and (b), we summarize for two different data generating processes [DGPs] the cdf's of $p_n^*|X_n$ across $M = 1,000$ independent realizations of X_n for samples of size $n = 10$ (upper panels) and $n = 1,000$ (lower panels). Specifically, the DGP used for panel (i) is based on i.i.d. shocks, while the one for (ii) features ARCH-type shocks (details are reported in the accompanying Supplement, Section S.4). In both cases, the conditions of Section 2.2.2 are satisfied. For both DGPs, the conditional distributions of p_n^* given X_n are, as expected, close to the 45° line, which corresponds to the implied asymptotic $U(0, 1)$ distribution. Unconditional validity follows accordingly.




FIGURE 1: *Fan chart of the simulated cdfs (conditional on X_n) of the bootstrap p-values for the three DGPs (i)–(iii) and $n = 10$ (upper panels), 1000 (lower panels).*

Nevertheless, unconditional validity may also hold without validity conditional on an apparently ‘natural’ conditioning variable X_n , like the regressor in a fixed-regressor

bootstrap design. For instance, suppose that for the DGP in Sections 2.2.1 and 2.2.2 it holds that $\eta_t = \xi_t(1 + \mathbb{I}_{\{\varepsilon_t < 0\}})$, with $\{\varepsilon_t\}$ and $\{\xi_t\}$ two independent i.i.d. sequences of zero-mean, unit-variance rv's. Since η_t is informative about the sign of ε_t , the ε_t 's conditionally on their own past and the regressor X_n do not form an mds. It is shown in Appendix B.1, eq. (B.4), that this endogeneity fact, not replicated in the bootstrap world, induces the original statistic τ_n to satisfy

$$\tau_n | X_n \xrightarrow{w} M^{-1/2} (\omega_{\varepsilon|\eta}^{1/2} \xi_1 + (1 - \omega_{\varepsilon|\eta})^{1/2} \xi_2) \Big| (M, \xi_2), \quad (2.8)$$

where $\omega_{\varepsilon|\eta} := E\{\text{Var}(\varepsilon_s | \eta_s)\} \in (0, 1)$, and M, ξ_1, ξ_2 are jointly independent with $\xi_i \sim N(0, 1), i = 1, 2$. The limit in (2.8) contains more randomness (through ξ_2) than the bootstrap limit in eq. (2.5), thus resulting in a random limit for the distribution of the bootstrap p -value p_n^* conditional on the regressor X_n ; see panel (iii) of Figure 1, where for this DGP the cdf's of $p_n^* | X_n$ are reported for 1,000 realizations of X_n . These cdf's display substantial dispersion around the 45° line, and this feature does not vanish as n increases. However, and in agreement with the earlier discussion, their unconditional average (plotted in black) is very close to the 45° line, showing indeed unconditional validity of the bootstrap. This follows because $e_t := (\varepsilon_t, \eta_t)'$ is a zero-mean i.i.d. sequence with a diagonal covariance matrix and $p_n^* \xrightarrow{w} U(0, 1)$ as derived in Section 2.2.1.

REMARK 2.2 Although not valid conditionally on the regressor X_n , in the previous example the bootstrap may be valid conditionally on a non-trivial function of the regressor. See, in particular, Section 3.3 and Remark 3.15 therein. \square

3 MAIN RESULTS

We provide general conditions for bootstrap validity in cases where a bootstrap statistic conditionally on the data possesses a random limit distribution. Before all else, we formally distinguish between two concepts of bootstrap validity.

3.1 DEFINITIONS

The following definition employs the bootstrap p -value as a summary indicator of the accuracy of bootstrap inferences (see also Remark 3.3 below). The original and the bootstrap statistic are denoted by τ_n and τ_n^* , respectively.

DEFINITION 1 Let $\tau_n := \tau_n(D_n)$ and $\tau_n^* := \tau_n^*(D_n, W_n^*)$, $n \in \mathbb{N}$, where D_n denotes the data whereas W_n^* are auxiliary variates defined jointly with D_n on a possibly extended probability space. Let $p_n^* := P(\tau_n^* \leq \tau_n | D_n)$ be the bootstrap p -value.

We say that the bootstrap based on τ_n and τ_n^* is valid unconditionally if p_n^* is asymptotically $U(0, 1)$ distributed:

$$P(p_n^* \leq q) \rightarrow q, \quad q \in (0, 1), \quad (3.1)$$

where $P(\cdot)$ denotes probability w.r.t. the distribution of D_n .

Let further X_n be a random element defined on the probability space of D_n and W_n^* . We say that the bootstrap based on τ_n and τ_n^* is valid conditionally on X_n if p_n^* is asymptotically $U(0,1)$ distributed conditionally on X_n :

$$P(p_n^* \leq q | X_n) \xrightarrow{P} q, q \in (0,1), \quad (3.2)$$

where $P(\cdot | X_n)$ is determined up to a.s. equivalence by the distribution of (D_n, X_n) .

REMARK 3.1 Bootstrap validity conditionally on some X_n implies unconditional validity, by the dominated convergence theorem. In applications, therefore, the discussion of conditional validity may represent an intermediate step to assess unconditional validity.

REMARK 3.2 The validity properties in Definition 1 ensure correct asymptotic null rejection probability, unconditionally or conditionally on some X_n , for bootstrap hypothesis tests which reject the null when the bootstrap p -value p_n^* does not exceed a chosen nominal level, say $\alpha \in (0,1)$. If $P(\tau_n^* \leq \cdot | D_n)$ converges weakly in $\mathcal{D}(\mathbb{R})$ to a sample-path continuous random cdf, then correct asymptotic null rejection probability is ensured also for bootstrap tests rejecting the null hypothesis when $\tilde{p}_n^* := P(\tau_n^* \geq \tau_n | D_n) \leq \alpha$ (for applications, see Sections 4.3 and 4.4).

REMARK 3.3 Validity as in Definition 1 has also implications on the properties of bootstrap (percentile) confidence sets. Suppose, for instance, that T_n is an estimator of a population (scalar) parameter, whose true value is denoted by θ_0 , and assume for simplicity that τ_n is of the form $\tau_n = \rho(n)(T_n - \theta_0)$, where $\rho(n)$ is a normalizing factor such that τ_n has a non-degenerate limiting distribution (see Horowitz, 2001, p.3174). Its bootstrap analog is denoted by τ_n^* , and we assume that the bootstrap is valid in the unconditional sense of (3.1). Interest is in constructing a right-sided confidence interval for θ_0 , with (asymptotic) coverage $1 - \alpha \in (0,1)$, using a simple bootstrap percentile method. With $F_n^*(x) := P(\tau_n^* \leq x | D_n)$, let $q_n^*(1 - \alpha) := \inf\{x \in \mathbb{R} : F_n^*(x) \geq 1 - \alpha\}$ be the $(1 - \alpha)$ quantile of the bootstrap distribution F_n^* . Then, it is straightforward to show that, if F_n^* converges weakly to a sample-path continuous random cdf, then

$$P(\tau_n \leq q_n^*(1 - \alpha)) = P(p_n^* \leq 1 - \alpha) + o(1) \rightarrow 1 - \alpha$$

This implies that a confidence interval of the form $[T_n - \rho(n)^{-1}q_n^*(1 - \alpha), +\infty)$ has (unconditional) asymptotic coverage probability of $1 - \alpha$. If the bootstrap is valid conditionally on some X_n , as in (3.2), then the (asymptotic) coverage is $1 - \alpha$ also conditionally on this X_n . \square

Our main results make extensive use of *joint* weak convergence in distribution. Should the related notation not be self-explanatory, we refer the reader to Appendix A for the formal definitions.

3.2 UNCONDITIONAL BOOTSTRAP VALIDITY

The unconditional validity results in this section have in common the requirement, explicit or implicit, that the unconditional limit distribution of τ_n should be an average of the random limit distribution of τ_n^* given the data. Applications of Theorem 3.1 do not require a conditional analysis of τ_n , in contrast to applications of Theorem 3.2.

THEOREM 3.1 *Let there exist a rv τ and a random element X , both defined on the same probability space, such that $(\tau_n, F_n^*) \xrightarrow{w} (\tau, F)$ in $\mathbb{R} \times \mathcal{D}(\mathbb{R})$ for $F_n^*(u) := P(\tau_n^* \leq u | D_n)$ and $F(u) := P(\tau \leq u | X)$, $u \in \mathbb{R}$. If the (possibly) random cdf F is sample-path continuous, then the bootstrap based on τ_n and τ_n^* is valid unconditionally.*

Some remarks are in order.

REMARK 3.4 A trivial special case of Theorem 3.1 is obtained for independent τ and X . In this case the bootstrap distribution of τ_n^* estimates consistently the limiting unconditional distribution of τ_n and the bootstrap is valid in the usual sense.

REMARK 3.5 An important special case of Theorem 3.1 involves stable convergence of the original statistic τ_n (see Häusler and Luschgy, 2015, p.33, for a definition). With the notation of Theorem 3.1, let the data D_n and the random element X be defined on the same probability space, whereas the rv τ be defined on an extension of this probability space. Assume that $\tau_n \rightarrow \tau$ stably and $F_n^* \xrightarrow{p} F$. Then $(\tau_n, F_n^*) \xrightarrow{w} (\tau, F)$ by Theorem 3.7(b) of Häusler and Luschgy (2015). For instance, in the statistical literature on integrated volatility, a result of the form $\tau_n \rightarrow \tau$ stably is contained in Theorem 3.1 of Jacod, Mykland, Podolskij and Vetter (2009) for τ_n defined as a t -type statistic for integrated volatility, whereas the corresponding $F_n^* \xrightarrow{p} F$ result is established in Theorem 3.1 of Hounyo, Gonçalves and Meddahi (2017) for a combined wild and blocks-of-blocks bootstrap introduced in the latter paper.

REMARK 3.6 More generally, if $\tau_n^* \xrightarrow{w^*} \tau | X$ and $(\tau_n^*, \tau_n, X_n) \xrightarrow{w} (\tau^*, \tau, X)$ with D_n -measurable X_n ($n \in \mathbb{N}$), then the joint convergence $((\tau_n^* | D_n), \tau_n, X_n) \xrightarrow{w} ((\tau^* | X), \tau, X)$ follows (see Lemma A.1(b) in Appendix A). If $\tau^* | X \stackrel{d}{=} \tau | X$ and F is sample-path continuous, then $(\tau_n, F_n^*) \xrightarrow{w} (\tau, F)$ by Lemma A.2(b) in Appendix A.

REMARK 3.7 Alternatively, the convergence $(\tau_n, F_n^*) \xrightarrow{w} (\tau, F)$ could be obtained from (a) the convergence $(\tau_n, X_n) \xrightarrow{w} (\tau, X)$ for some D_n -measurable random elements X_n of the space of X , and (b) the implication (were it to hold) from the strong version $(\tau_n, X_n) \xrightarrow{a.s.} (\tau, X)$ to $\tau_n^* \xrightarrow{w^*} \tau | X$. The idea is to choose X_n such that τ_n^* depends on the data essentially through X_n . Applications of Theorem 3.1 along these lines could proceed in two steps: (i) prove that $(\tau_n, X_n) \xrightarrow{w} (\tau, X)$; (ii) consider, by extended Skorokhod coupling (Corollary 5.12 of Kallenberg, 1997), a representation of D_n and (τ, X) such that, with an abuse of notation, $(\tau_n, X_n) \xrightarrow{a.s.} (\tau, X)$ and, on a product extension of the Skorokhod-representation space, prove that $\tau_n^* \xrightarrow{w^*} \tau | X$. The latter conditional assertion, due to the product structure of the probability space, reduces to a collection of unconditional assertions by fixing the outcomes in the factor-space of the data. It then holds that $(\tau_n, F_n^*) \xrightarrow{p} (\tau, F)$ on the Skorokhod-representation space, whereas on a general probability space $(\tau_n, F_n^*) \xrightarrow{w} (\tau, F)$. We proceed like this in the applications of Section 4.2 (eq. (4.6)) and Section 4.4 (Theorem 4.3 under Assumption \mathcal{H}). \square

Unconditional bootstrap validity could also be established by means of an auxiliary conditional analysis of the original statistic τ_n . In the next theorem the conditioning

sequence X_n is chosen such that the bootstrap statistic τ_n^* depends on the data D_n approximately through X_n (condition (\dagger)). Then, the main requirement for bootstrap validity is that the limit bootstrap distribution should be a conditional average of the limit distribution of τ_n given X_n .

THEOREM 3.2 *With the notation of Definition 1, let X_n be D_n -measurable ($n \in \mathbb{N}$). Let it hold that*

$$(P(\tau_n \leq \cdot | X_n), P(\tau_n^* \leq \cdot | D_n)) \xrightarrow{w} (F, F^*) \quad (3.3)$$

in $\mathcal{D}(\mathbb{R}) \times \mathcal{D}(\mathbb{R})$, where F and F^* are sample-path continuous random cdf's, and let

(\dagger) there exist random elements X', X'_n such that F^* is X' -measurable, X'_n are X_n -measurable and $X'_n \xrightarrow{w} X'$ jointly with (3.3).

Then, if $E\{F(\cdot) | F^*\} = F^*(\cdot)$, the bootstrap based on τ_n and τ_n^* is valid unconditionally.

REMARK 3.8 Condition (\dagger) of Theorem 3.2 implies that τ_n^* depends on the data D_n approximately through X_n alone, for under this condition $P(\tau_n^* \leq \cdot | X_n)$ and $P(\tau_n^* \leq \cdot | D_n)$ are both close to $P(\tau_n^* \leq \cdot | X'_n)$. Condition (\dagger) is trivially satisfied in the case $F = F^*$ with the choice $X'_n = P(\tau_n \leq \cdot | X_n)$. It is also satisfied if $\tau_n^* = \tilde{\tau}_n^* + o_p(1)$ for some $\tilde{\tau}_n^*$ which is a measurable transformation of X_n and W_n^* , w.r.t. the probability measure on the space where D_n and W_n^* are jointly defined. In this case, $X'_n = P(\tilde{\tau}_n^* \leq \cdot | X_n)$ satisfies condition (\dagger) ; see Appendix B.2. An example of a pair $\tau_n^*, \tilde{\tau}_n^*$ is given in eq. (4.3) in Section 4.2. \square

Convergence (3.3) in Theorem 3.2 could be deduced from the weak convergence of the conditional distributions of τ_n and τ_n^* , as in the next corollary.

COROLLARY 3.1 *Let D_n and X_n ($n \in \mathbb{N}$) be as in Theorem 3.2. Let the rv τ and the random elements X, X' be defined on a single probability space and*

$$(\tau_n | X_n, \tau_n^* | D_n) \xrightarrow{w} (\tau | X, \tau | X') \quad (3.4)$$

in the sense of eq. (A.1). Let further $F(u) := P(\tau \leq u | X)$ and $F^*(u) := P(\tau \leq u | X')$, $u \in \mathbb{R}$, define sample-path continuous random cdf's. Then convergence (3.3) holds. Moreover, the bootstrap based on τ_n and τ_n^* is valid unconditionally provided that one of the following extra conditions holds:

- (a) $X' = X$;
- (b) $X = (X', X'')$ and $X'_n \xrightarrow{w} X'$ jointly with (3.4) for some X_n -measurable random elements X'_n .

REMARK 3.9 An instance of (3.3) with $F \neq F^*$ is implied by the setup of Section 2.2.3. There (3.4) holds with $\tau := M^{-1/2}(\omega_{\varepsilon|\eta}^{1/2}\xi_1 + (1 - \omega_{\varepsilon|\eta})^{1/2}\xi_2)$ and $X = (X', X'') = (M, (1 - \omega_{\varepsilon|\eta})^{1/2}\xi_2)$. Moreover, (3.4) is joint with the convergence $X'_n \xrightarrow{w} X'$ for $X'_n = n^{-2}M_n$ (see Appendix B.2). Hence, Corollary 3.1(b) implies that the bootstrap is unconditionally valid, as was already concluded in Section 2.2.1.

REMARK 3.10 Convergence (3.4) could be proved by replacing in Remark 3.7 the convergence $(\tau_n, X_n) \rightarrow (\tau, X)$ (weakly and a.s.) by $((\tau_n|X_n), X'_n) \rightarrow ((\tau|X), X')$ (weakly in distribution and weakly a.s.) Other ways of proving (3.4), that could be relevant if conditional bootstrap validity is of interest, are discussed in the next section. \square

3.3 CONDITIONAL BOOTSTRAP VALIDITY

Theorem 3.3 below states the asymptotic behavior of the bootstrap p -value conditional on an X_n chosen to satisfy condition (\dagger) of Theorem 3.2. It also characterizes the cases where the bootstrap is valid conditionally on such an X_n . Should validity conditional on such an X_n fail, in Corollary 3.2(b) we provide a result for validity conditional on a transformation of it.

THEOREM 3.3 *Under the conditions of Theorem 3.2, the bootstrap p -value p_n^* satisfies*

$$P(p_n^* \leq q | X_n) \xrightarrow{w} F(F^{*-1}(q)) \quad (3.5)$$

for almost all $q \in (0, 1)$, and the bootstrap based on τ_n and τ_n^* is valid conditionally on X_n if and only if $F = F^*$ such that

$$\sup_{u \in \mathbb{R}} |P(\tau_n \leq u | X_n) - P(\tau_n^* \leq u | D_n)| \xrightarrow{p} 0. \quad (3.6)$$

REMARK 3.11 Under (3.6), the bootstrap distribution of τ_n^* consistently estimates the limit of the conditional distribution of τ_n given X_n . Although under condition (\dagger) the proximity of $P(\tau_n \leq \cdot | X_n)$ and $P(\tau_n^* \leq \cdot | D_n)$ is necessary for bootstrap validity conditional on X_n , no such proximity is necessary for conditional validity in the general case. In fact, validity conditional on some X_n implies validity conditional on any measurable transformation $X'_n = \psi_n(X_n)$ and an analogue of (3.6) with X'_n in place of X_n cannot generally hold for all ψ_n , unless F^* is non-random. This is similar to what happens with unconditional bootstrap validity which, according to Theorem 3.1, may occur even if $P(\tau_n \leq \cdot)$ and $P(\tau_n^* \leq \cdot | D_n)$ are not close. \square

A corollary in the terms of weak convergence in distribution is given next.

COROLLARY 3.2 *Let D_n, X_n ($n \in \mathbb{N}$), τ, F, F^* be as in Corollary 3.1. Let (3.4) hold and F, F^* be sample-path continuous random cdf's. Then:*

(a) *If $X' = X$, the bootstrap based on τ_n and τ_n^* is valid conditionally on X_n and (3.6) holds.*

(b) *If $X = (X', X'')$, $(X'_n, X''_n) \xrightarrow{w} (X', X'')$ jointly with (3.4) for some X_n -measurable random elements (X'_n, X''_n) , and $X''_n | X'_n \xrightarrow{w} X'' | X'$, then the bootstrap is valid conditionally on X'_n and (3.6) holds with X_n replaced by X'_n .*

Corollary 3.2 requires checking the joint convergence in (3.4); see Remark 3.10. We provide further strategies to establish this convergence, useful if interest is in conditional bootstrap validity, in Remarks 3.13–3.14 below.

REMARK 3.12 Consider the linear regression example under the extra assumptions of Section 2.2.2 and set $\tau = (\int B_\eta^2)^{-1} \int B_\eta dB_\varepsilon$, $X = M$. It then follows (by using Theorem 3 of Georgiev *et al.*, 2018) that condition (3.4) holds in the form

$$(\tau_n|X_n, \tau_n^*|D_n) \xrightarrow{w} (\tau|B_\eta, \tau|B_\eta) \stackrel{d}{=} (1, 1)N(0, \omega_\varepsilon M^{-1})|M, \quad (3.7)$$

where $X_n := \{x_t\}_{t=1}^n$; equivalently, (3.3) holds with $F = F^* = \Phi(\omega_\varepsilon^{-1/2} M^{1/2}(\cdot))$. Hence, the bootstrap is consistent for the limit distribution of τ_n conditional on the regressor and, by Corollary 3.2(a), the bootstrap is valid conditionally on the regressor.

REMARK 3.13 The joint convergence in (3.4) would follow from the separate convergence facts $\tau_n|X_n \xrightarrow{w} \tau|X$, $\tau_n^* \xrightarrow{w^*} \tau^*|X'$ and $(\tau_n, \tau_n^*, \phi_n(X_n), \psi_n(D_n)) \xrightarrow{w} (\tau, \tau^*, X, X')$ for some measurable ϕ_n, ψ_n , provided that $\tau|X' \stackrel{d}{=} \tau^*|X'$; see Appendix B.2. We use this approach in Section 4.2, point (ii). The convergence $\tau_n|X_n \xrightarrow{w} \tau|X$ is the new ingredient compared to Remark 3.6.

REMARK 3.14 Convergence (3.4) would also follow from $(\tau_n, \phi_n(X_n), \psi_n(D_n)) \xrightarrow{w} (\tau, X, X')$ and $\tau_n|X_n \xrightarrow{w} \tau|X$ together with the implication (were it to hold) from $\psi_n(D_n) \xrightarrow{a.s.} X'$ to $\tau_n^* \xrightarrow{w^*} \tau^*|X'$, with $\tau|X' \stackrel{d}{=} \tau^*|X'$. A possible implementation strategy is: (i) prove that $\tau_n|X_n \xrightarrow{w} \tau|X$ and $(\tau_n, \phi_n(X_n), \psi_n(D_n)) \xrightarrow{w} (\tau, X, X')$; (ii) consider a Skorokhod representation of D_n and (τ, X, X') such that, maintaining the notation, $(\tau_n, \phi_n(X_n), \psi_n(D_n)) \xrightarrow{a.s.} (\tau, X, X')$ and, as a result, $\tau_n|X_n \xrightarrow{w} \tau|X$ strengthens to $\tau_n|X_n \xrightarrow{w} \tau|X$ (see Lemma A.1 in Appendix A); (iii) redefine the bootstrap variates W_n^* on a product extension of the Skorokhod-representation space and prove there that $\tau_n^* \xrightarrow{w^*} \tau^*|X'$. Then (3.4) holds on a general probability space. We proceed like this in the proof of Theorem 4.3 under Assumption \mathcal{C} . The convergence $\tau_n|X_n \xrightarrow{w} \tau|X$ is the extra ingredient compared to Remark 3.7. Notice also that if $\phi_n(X_n) = (X'_n, X''_n)$ and $\psi_n(D_n) = X'_n$, then the convergence $(X'_n, X''_n) \xrightarrow{w} (X', X'')$ in Corollary 3.2(b) would be joint with (3.4).

REMARK 3.15 In the setup of Section 2.2.3, (3.4) holds with τ and $X = (X', X'')$ given in Remark 3.9. Moreover, (3.4) is joint with the convergence $(X'_n, X''_n) \xrightarrow{w} (X', X'')$ for $X'_n = n^{-2}M_n$ and $X''_n = M_n^{-1/2} \sum_{t=1}^n x_t E(\varepsilon_t|\eta_t)$ (see Appendix B.2). By Corollary 3.2(b), the bootstrap would be valid conditionally on M_n if it additionally holds that $X''_n|M_n \xrightarrow{w} (1 - \omega_{\varepsilon|\eta})^{1/2} \xi_2|M \stackrel{d}{=} N(0, 1 - \omega_{\varepsilon|\eta})$. \square

3.4 LOCAL POWER OF BOOTSTRAP TESTS

When the limit bootstrap measure is random, the power function of the bootstrap test, conditionally on the data, is also random, even asymptotically. Its unconditional power function can be investigated using the following generalization of Theorem 3.1.

THEOREM 3.4 *Let there exist rv's τ, τ^* and a random element X , the three defined on the same probability space, such that $(\tau_n, F_n^*) \xrightarrow{w} (\tau, F^*)$ in $\mathbb{R} \times \mathcal{D}(\mathbb{R})$ for $F_n^*(u) := P(\tau_n^* \leq u|D_n)$ and $F^*(u) := P(\tau^* \leq u|X)$, $u \in \mathbb{R}$. If F^* is sample-path continuous, then*

the bootstrap p -value p_n^* satisfies $P(p_n^* \leq q) \rightarrow E\{F(F^{*-1}(q))\}$, $q \in (0, 1)$ with $F(\cdot) := P(\tau \leq \cdot | X)$.

To illustrate, with y_t , x_t and ε_t as in Section 2.2.2, let interest be in the large-sample behavior of the bootstrap test for the hypothesis $H_0 : \beta = 0$ against $H_1 : \beta < 0$, under the local alternative $\beta = \beta_n := b/n$ in (2.1). The original test statistic is $\tau_n = n\hat{\beta}$.

Without recourse to the explicit expression in (2.3) for the bootstrap p -value p_n^* in terms of the Gaussian cdf, which in many applications may have no analogue, we can instead use, for $\tau_n = b + n(\hat{\beta} - \beta)$ and τ_n^* , the joint convergence²

$$(\tau_n, (\tau_n^* | D_n)) \xrightarrow{w} (\tau, (\tau^* | M)),$$

with $\tau := b + \tau^*$ and $\tau^* := \omega_\varepsilon^{1/2} M^{-1/2} \xi$ for $\xi \sim N(0, 1)$ independent of M . By using Lemma A.2 in Appendix A, we can conclude that the conditions of Theorem 3.4 hold with $X = M$, $F(u) = P(\tau \leq u | M) = \Phi(\omega_\varepsilon^{-1/2} M^{1/2}(u - b))$ and $F^*(u) = \Phi(\omega_\varepsilon^{-1/2} M^{1/2}u)$. The unconditional asymptotic local power function of the one-sided, q -level, bootstrap test then follows as

$$P(p_n^* \leq q) \rightarrow E\{F(F^{*-1}(q))\} = E\{\Phi(\Phi^{-1}(q) - \omega_\varepsilon^{-1/2} M^{1/2}b)\}. \quad (3.8)$$

Notice that this power function is distinct from the asymptotic local power of the unconditional test based on critical values from the null asymptotic (unconditional) distribution of τ_n , which is that of $\omega_\varepsilon^{1/2} M^{-1/2} \xi$. Hence, when the limit bootstrap measure is random, the bootstrap test in general does not replicate, in terms of (unconditional) power, the standard asymptotic test (in this specific case, numerical evidence shows that for small b , where both local powers are relatively low, the bootstrap test is more powerful, whereas for large negative b , where the local power of both tests is high, the asymptotic test is preferable).

The unconditional power function in (3.8) can also be derived through a conditioning argument. This can be done using the results in Section 3.3 by considering the joint convergence

$$(\tau_n | X_n, \tau_n^* | D_n) \xrightarrow{w} (b, 0) + (1, 1)N(0, \omega_\varepsilon M^{-1}) | M,$$

see (3.7), which implies (3.3) with F and F^* as defined above. Hence, by Theorem 3.3, (3.5) holds and

$$P(p_n^* \leq q | X_n) \xrightarrow{w} \Phi(\omega_\varepsilon^{-1/2} M^{1/2}(F^{*-1}(q) - b)) = \Phi(\Phi^{-1}(q) - \omega_\varepsilon^{-1/2} M^{1/2}b) \quad (3.9)$$

for $q \in (0, 1)$. The latter expression is the (random) asymptotic local power, conditional on X_n , of the one-sided, q -level test, bootstrap test. By averaging the rhs of (3.9) over M , the unconditional power function in (3.8) follows.

²The conditional analysis of τ_n^* , needed to show that under local alternatives it behaves asymptotically as under H_0 , is straightforward and is omitted.

4 APPLICATIONS

4.1 A PERMUTATION CUSUM TEST UNDER INFINITE VARIANCE

Consider a standard CUSUM test for the null hypothesis (say, H_0) that $\{\varepsilon_t\}_{t=1}^n$ is a sequence of i.i.d. random variables. The test statistic is of the form

$$\tau_n := \nu_n^{-1} \max_{t=1, \dots, n} \left| \sum_{i=1}^t (\varepsilon_i - \bar{\varepsilon}_n) \right|, \quad \bar{\varepsilon}_n := n^{-1} \sum_{t=1}^n \varepsilon_t,$$

where ν_n is a permutation-invariant normalization sequence. Standard choices are $\nu_n^2 = \sum_{t=1}^n (\varepsilon_t - \bar{\varepsilon}_n)^2$ in the case where $E\varepsilon_t^2 < \infty$, and $\nu_n = \max_{t=1, \dots, n} |\varepsilon_t|$ when $E\varepsilon_t^2 = \infty$. If ε_t is in the domain of attraction of a strictly α -stable law with $\alpha \in (0, 2)$, such that $E\varepsilon_t^2 = \infty$, the asymptotic distribution of τ_n depends on unknown parameters (e.g., the characteristic exponent α), which makes the test difficult to apply (see also Politis, Romano and Wolf, 1999, and the references therein). To overcome this problem, Aue *et al.* (2008) consider a permutation analogue of τ_n , defined as

$$\tau_n^* := \nu_n^{-1} \max_{t=1, \dots, n} \left| \sum_{i=1}^t (\varepsilon_{\pi(i)} - \bar{\varepsilon}_n) \right|$$

where π is a (uniformly distributed) random permutation of $\{1, 2, \dots, n\}$, independent of the data.³ In terms of Definition 1, the data is $D_n := \{\varepsilon_t\}_{t=1}^n$ and the auxiliary ‘bootstrap’ variate is $W_n^* := \pi$. With $X_n := \{\varepsilon_{(t)}\}_{t=1}^n$ denoting the vector of order statistics of $\{\varepsilon_t\}_{t=1}^n$, there exists a random permutation ϖ of $\{1, \dots, n\}$ (under H_0 , uniformly distributed conditionally on X_n) for which it holds that $\varepsilon_t = \varepsilon_{(\varpi(t))}$ ($t = 1, \dots, n$), whereas the ‘bootstrap’ sample is $\{\varepsilon_{\pi(t)}\}_{t=1}^n$. The results in Aue *et al.* (2008, Corollary 2.1, Theorem 2.4) imply that, if H_0 holds and ε_t is in the domain of attraction of a strictly α -stable law with $\alpha \in (0, 2)$, then $\tau_n \xrightarrow{w} \rho_\alpha(S)$ and $\tau_n^* \xrightarrow{w^*} \rho_\alpha(S)|S$ for a certain random function ρ_α and $S = (S_1, S_2)'$, with $S_i = \{S_{ij}\}_{j=1}^\infty$ ($i = 1, 2$) being partial sums of sequences of i.i.d. standard exponential rv’s, and with ρ_α independent of S .⁴

Aue *et al.* (2008) do not report the fact that statistical inferences are not invalidated by the failure of the permutation procedure to estimate consistently the distribution of $\rho_\alpha(S)$. In fact, the situation is similar to that of Remark 2.1, as $\tau_n|X_n \stackrel{d}{=} \tau_n^*|D_n$ under H_0 . As a consequence, under H_0 the permutation test implements *exact*⁵ finite-sample inference conditional on X_n and, additionally, the distribution of τ_n^* given the data estimates consistently the limit of the conditional distribution $\tau_n|X_n$, in the sense of joint weak convergence in distribution (see eq. (A.1)):

$$(\tau_n|X_n, \tau_n^*|D_n)' \xrightarrow{w^*} (\rho_\alpha(S)|S, \rho_\alpha(S)|S). \quad (4.1)$$

³The normalization of ν_n is only of theoretical importance for obtaining non-degenerate limit distributions. In practice, any bootstrap procedure comparing τ_n to the quantiles of τ_n^* is invariant to the choice of ν_n and can be implemented by setting $\nu_n = 1$.

⁴To avoid centering terms, Aue *et al.* (2008) assume additionally that the location parameter of the limit stable law is zero when $\alpha \in [1, 2)$. Moreover, although they provide conditional convergence results only for the finite-dimensional distributions of the CUSUM process, these could be strengthened to conditional functional convergence as in Proposition 1 of LePage *et al.* (1997) in order to obtain the conditional convergence of τ_n^* .

⁵Here by ‘exact’ we mean that bootstrap inference replicates the finite-sample (conditional) distribution of the test statistic for any sample size with no error.

CUSUM tests can also be applied to residuals from an estimated model in order to test for correct model specification or stability of the parameters (see e.g., Ploberger and Krämer, 1992). Consider thus the case where $\{\varepsilon_t\}_{t=1}^n$ are the disturbances in a statistical model (e.g., the regression model of Section 2), and we observe residuals $\hat{\varepsilon}_t$ obtained upon estimation of the model using a sample D_n not containing the unobservable $\{\varepsilon_t\}_{t=1}^n$. The residual-based CUSUM statistic is $\hat{\tau}_n := \hat{\nu}_n^{-1} \max_{t=1, \dots, n} |\sum_{i=1}^t (\hat{\varepsilon}_i - \bar{\hat{\varepsilon}}_n)|$, where $\hat{\nu}_n$ and $\bar{\hat{\varepsilon}}_n$ are the analogues of ν_n and $\bar{\varepsilon}_n$ computed from $\hat{\varepsilon}_t$ instead of ε_t . The bootstrap statistic could be defined as $\hat{\tau}_n^* := \hat{\nu}_n^{-1} \max_{t=1, \dots, n} |\sum_{i=1}^t (\hat{\varepsilon}_{\pi(i)} - \bar{\hat{\varepsilon}}_n)|$. If $\hat{\tau}_n - \tau_n \xrightarrow{p} 0$ and $(\hat{\tau}_n^* - \tau_n^*)|D_n \xrightarrow{w} 0$ under H_0 (e.g., due to consistent parameter estimation), then also $(\hat{\tau}_n - \tau_n)|X_n \xrightarrow{w} 0$, such that the (Lévy) distances between the pairs of conditional distributions $\hat{\tau}_n|X_n$ and $\tau_n|X_n$ on the one hand, and $\hat{\tau}_n^*|D_n$ and $\tau_n^*|D_n$ on the other hand, converge in probability to zero. Hence, in view of (4.1), and under the conjecture that $P(\rho_\alpha(S) \leq \cdot|S)$ defines a sample-path continuous cdf, the residual-based permutation procedure is consistent in the sense that

$$(\hat{\tau}_n|X_n, \hat{\tau}_n^*|D_n) \xrightarrow{w} (\rho_\alpha(S)|S, \rho_\alpha(S)|S) \quad (4.2)$$

for $X_n := \{\varepsilon_{(t)}\}_{t=1}^n$ again. It follows that:

- (i) The permutation residual-based test is valid conditionally on X_n , by Corollary 3.2(a) with condition (3.4) taking the form (4.2).
- (ii) This test is valid unconditionally, as a results of either the validity conditional on X_n , or by Corollary 3.1.

4.2 A PARAMETRIC BOOTSTRAP GOODNESS-OF-FIT TEST

The parametric bootstrap is a standard technique for the approximation of a conditional distribution of goodness-of-fit test statistics (Andrews, 1997; Lockhart, 2012). When these are discussed in the i.i.d. finite-variance setting, the limit of the bootstrap distribution is non-random. However, if we return to the relation (2.1), there exist relevant settings where a random limit of the normalized M_n implies that parametrically bootstrapped goodness-of-fit test statistics have random limit distributions.

4.2.1 SET UP AND A RANDOM LIMIT BOOTSTRAP MEASURE

Let the null hypothesis of interest, say H_0 , be that the standardized errors $\omega_\varepsilon^{-1/2}\varepsilon_t$ in (2.1) have a certain known density f with mean 0 and variance 1. For expositional ease we assume that $\omega_\varepsilon = 1$ and is known to the econometrician. Then, the Kolmogorov-Smirnov statistic based on OLS residuals $\hat{\varepsilon}_t$ is

$$\tau_n := n^{1/2} \sup_{u \in \mathbb{R}} \left| n^{-1} \sum_{t=1}^n \mathbb{I}_{\{\hat{\varepsilon}_t \leq u\}} - \int_{-\infty}^u f \right|.$$

A (parametric) bootstrap counterpart, τ_n^* , of τ_n could be constructed under H_0 by (i) drawing $\{\varepsilon_t^*\}_{t=1}^n$ as i.i.d. from f , independent of the data; (ii), regressing them on x_t , thus obtaining an estimator $\hat{\beta}^*$ and associated residuals $\hat{\varepsilon}_t^*$; and (iii) calculating τ_n^* as $\tau_n^* := n^{1/2} \sup_{u \in \mathbb{R}} |n^{-1} \sum_{t=1}^n \mathbb{I}_{\{\hat{\varepsilon}_t^* \leq u\}} - \int_{-\infty}^u f|$.

To see that the distribution of the bootstrap statistic τ_n^* conditional on the data $D_n := \{x_t, y_t\}_{t=1}^n$ may have a random limit, consider the Gaussian case, $f = \Phi'$. Under the assumptions of Johansen and Nielsen (2016, Sec. 4.1-4.2), it holds (*ibidem*) that $\tau_n^* = \tilde{\tau}_n^* + o_p(1)$ under the product probability on the product probability space where the data and $\{\varepsilon_t^*\}$ are jointly defined, with

$$\tilde{\tau}_n^* := \sup_{u \in [0,1]} \left| n^{-1/2} \sum_{t=1}^n (\mathbb{I}_{\{\varepsilon_t^* \leq q(u)\}} - u) + \Phi'(q(u)) \hat{\beta}^* n^{-1/2} \sum_{t=1}^n x_t \right|, \quad (4.3)$$

where $q(u) = \Phi^{-1}(u)$ is the u -th quantile of Φ . The expansion of τ_n^* holds also conditionally on the data, i.e., $\tau_n^* - \tilde{\tau}_n^* \xrightarrow{w^*} 0$, since convergence in probability to a constant is preserved upon such conditioning. Hence, if $\tilde{\tau}_n^* | D_n$ converges to a random limit, so does $\tau_n^* | D_n$ for the same limit. Assume that $X_n := n^{-\alpha/2} x_{\lfloor n \cdot \rfloor} \xrightarrow{w} X$ in \mathcal{D} for some $\alpha > 0$ and that $M := \int X^2 > 0$ a.s. (e.g., $X = B_\eta$ if $x_t = \sum_{s=1}^{t-1} \eta_s$ with $\{\eta_t\}$ introduced in Section 2.2). Then $(M_n, \xi_n) := (\sum_{t=1}^n x_t^2, \sum_{t=1}^n x_t)$ satisfies $(n^{-\alpha-1} M_n, n^{-\alpha/2-1} \xi_n) \xrightarrow{w} (M, \xi)$, $\xi := \int X$. Furthermore, if $W_n^*(u) := n^{-1} \sum_{t=1}^n (\mathbb{I}_{\{\varepsilon_t^* \leq q(u)\}} - u)$, $u \in [0, 1]$, is the bootstrap empirical process in probability scale, then W_n^* and $M_n^{1/2} \hat{\beta}^*$ are independent of the data individually (the second one being conditionally standard Gaussian), but not jointly independent of the data, because

$$\text{Cov}^*(n^{1/2} W_n^*(u), M_n^{1/2} \hat{\beta}^*) = (n^{-\alpha-1} M_n)^{-1/2} n^{-\alpha/2-1} \xi_n \psi(u) \xrightarrow{w} M^{-1/2} \xi \psi(u),$$

$u \in [0, 1]$, where $\psi(\cdot) := E^*[\varepsilon_1^* \mathbb{I}_{\{\varepsilon_1^* \leq q(\cdot)\}}] = -\Phi'(q(\cdot))$ is a trimmed mean function, with $\text{Cov}^*(\cdot)$ and $E^*(\cdot)$ calculated under P^* . It is shown in Appendix B.3 that, more strongly,

$$(n^{1/2} W_n^*, n^{(\alpha+1)/2} \hat{\beta}^*, n^{-\alpha/2-1} \xi_n) \xrightarrow{w^*} (W, M^{-1/2} b, \xi) \Big| (M, \xi) \quad (4.4)$$

on $\mathcal{D} \times \mathbb{R}^2$, where (W, b) is a pair of a standard Brownian bridge and a standard Gaussian rv individually independent of X (and thus, of M, ξ), but with Gaussian joint conditional (on X) distributions having covariance $\text{Cov}(W(u), b | X) = M^{-1/2} \xi \psi(u)$, $u \in [0, 1]$. Combining the expansion of τ_n^* , (4.3) and (4.4) with the extended CMT (Theorem A.1 in Appendix A) yields

$$\tau_n^* \xrightarrow{w^*} \left\{ \sup_{u \in [0,1]} |W(u) + \Phi'(q(u)) M^{-1/2} b \xi| \right\} (M, \xi) \stackrel{d}{=} \tau | (M, \xi), \quad (4.5)$$

where $\tau := \sup_{u \in [0,1]} |\tilde{W}(u)|$ for a process \tilde{W} which conditionally on X (and thus, on M, ξ), is a zero-mean Gaussian process with $\tilde{W}(0) = \tilde{W}(1) = 0$ a.s. and conditional covariance function $K(u, v) = u(1-v) - M^{-1} \xi^2 \psi(u) \psi(v)$ for $0 \leq u \leq v \leq 1$. In summary, the limit bootstrap distribution is random because the latter conditional covariance is random whenever M or ξ are such.

4.2.2 BOOTSTRAP VALIDITY

We now discuss in what sense τ_n^* can provide a distributional approximation of τ_n and whether the bootstrap can be valid in the sense of Definition 1.

(i) Under H_0 that $\varepsilon_t \sim \text{i.i.d. } N(0, 1)$, the bootstrap could be shown to be unconditionally valid using Theorem 3.1. Specifically, under H_0 , the assumptions and results of Johansen and Nielsen (2016, Sec. 4.1-4.2) guarantee that τ_n has the expansion $\tau_n = \tilde{\tau}_n + o_p(1)$, with $\tilde{\tau}_n := \sup_{u \in [0,1]} |n^{-1/2} \sum_{t=1}^n (\mathbb{I}_{\{\varepsilon_t \leq q(u)\}} - u) + \Phi'(q(u))(\hat{\beta} - \beta)n^{-1/2} \sum_{t=1}^n x_t|$ defined similarly to $\tilde{\tau}_n^*$. Assume that $\hat{\beta}$ is asymptotically mixed Gaussian, such that jointly with $n^{-\alpha/2} x_{\lfloor n \cdot \rfloor} \xrightarrow{w} X$ it holds that

$$(n^{-1/2} \sum_{t=1}^n (\mathbb{I}_{\{\varepsilon_t \leq q(u)\}} - u), n^{(\alpha+1)/2}(\hat{\beta} - \beta), n^{-\alpha/2-1}\xi_n) \xrightarrow{w} (W, M^{-1/2}b, \xi) ;$$

then $\tau_n = \tilde{\tau}_n + o_p(1) \xrightarrow{w} \tau = \sup_{u \in [0,1]} |\tilde{W}(u)|$. Thus, the unconditional limit of τ_n obtains by averaging (over M, ξ) the conditional limit of τ_n^* . This is the main prerequisite for establishing unconditional bootstrap validity via Theorem 3.1. More precisely, it is proved in Appendix B.3 that

$$(\tau_n, F_n^*) \xrightarrow{w} (\tau, F), F_n^*(\cdot) := P^*(\tau_n^* \leq \cdot), F(\cdot) := P(\tau \leq \cdot | M, \xi). \quad (4.6)$$

As F is sample-path continuous (e.g., by Proposition 3.2 of Linde, 1989, applied conditionally on M, ξ), Theorem 3.1 guarantees the unconditional validity of the bootstrap.

(ii) As $\tau_n = \tilde{\tau}_n + o_p(1)$ under H_0 , with $\tilde{\tau}_n$ related to (M_n, ξ_n) through the same functional form as $\tilde{\tau}_n^*$, it is possible for $\tau_n | X_n$ to have the same random limit distribution under H_0 as τ_n^* given the data, i.e., $\tau_n | X_n \xrightarrow{w} \tau | (M, \xi)$. For instance, this occurs if $\{\varepsilon_t\}$ is an i.i.d. sequence independent of X_n , by the same argument as for $\tilde{\tau}_n^*$. According to Remark 3.13, the convergence $\tau_n | X_n \xrightarrow{w} \tau | (M, \xi)$ and the convergence $(\tau_n, \tau_n^*, n^{-\alpha-1}M_n, n^{-\alpha/2-1}\xi_n) \xrightarrow{w} (\tau, \tau^*, M, \xi)$ with $\tau^* | (M, \xi) \stackrel{d}{=} \tau | (M, \xi)$ (shown in the proof of (4.6), see Appendix B.3) are sufficient for eq. (3.4) to hold in the form

$$(\tau_n | X_n, \tau_n^* | D_n) \xrightarrow{w} (\tau | (M, \xi), \tau | (M, \xi)).$$

As F is sample-path continuous, the bootstrap is valid conditionally on X_n by Corollary 3.2(a).

4.3 PARAMETERS ON THE BOUNDARY IN PREDICTIVE REGRESSION

Here we consider an instance of the ‘parameter on the boundary’ problem in the framework of predictive regressions for financial returns; see e.g. Phillips (2014) and the references therein. While in this context the bootstrap is potentially useful (e.g., when there is uncertainty about the degree of persistence of the posited predicting variable), its application is not straightforward if some of the parameters may lie on the boundary of the parameter space; see Andrews (2000).

We show that in the presence of parameters on the boundary, the distribution of the bootstrap statistic may be random in the limit. Moreover, the type of randomness induced by parameters on the boundary depends on how well the bootstrap scheme approximates the mutual position of three objects, namely (i) the boundary, (ii) the set identified by the null hypothesis, and (iii) the true parameter value. Standard bootstrap approximations may not be sufficiently precise, giving rise to complex conditioning in

the limit bootstrap distribution, with ensuing unconditional bootstrap validity only for special statistics. Conversely, non-standard, or ad hoc, bootstrap schemes, designed to provide a better match with the original geometry, may feature limit bootstrap distributions where no randomness attributable to the possibly boundary value of a parameter is present.

4.3.1 GENERAL SETUP

Consider the predictive regression

$$y_t = \theta_1 + \theta_2 x_{n,t-1} + \varepsilon_t \quad (t = 1, \dots, n; n = 1, 2, \dots) \quad (4.7)$$

under Assumption 1 of Georgiev et al. (2018), specialized for simplicity to unconditionally homoskedastic errors. The posited predicting variable $x_{n,t}$ is such that, in \mathcal{D} , $x_{n,[n \cdot]} \xrightarrow{w} X$, e.g. a Brownian motion or an Ornstein-Uhlenbeck process, and hence features low frequency variability in the sense of Müller and Watson (2008). We assume that the parameter space, say Θ , is defined by an inequality constraint and that the true value of the parameter $\theta := (\theta_1, \theta_2)'$, say $\theta_0 := (\theta_{1,0}, \theta_{2,0})'$, may lie on the boundary of Θ . An important example is when θ is assumed to belong to the set $\mathbb{R} \times [0, \infty)$, with the boundary corresponding to the case $\theta_2 = 0$ of no predictability of y_t by $x_{n,t-1}$ and the interior corresponding to (sign-restricted) predictability.

More specifically, assume that $\Theta := \{\theta \in \mathbb{R}^2 : g(\theta) \geq 0\}$, where $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a real function, continuously differentiable on some neighborhood of θ_0 and with gradient $\frac{\partial}{\partial \theta} g(\theta) \neq 0$ on that neighborhood, with $\dot{g} := \frac{\partial}{\partial \theta} g(\theta_0)$. The boundary of Θ is denoted by $\partial\Theta := \{\theta \in \mathbb{R}^2 : g(\theta) = 0\}$. The aforementioned example $\theta_2 \geq 0$ is obtained by setting $g(\theta) = (0, 1)\theta = \theta_2$.

Interest is in bootstrap inference on a null hypothesis H_0 identifying a set of parameter values that has a non-empty intersection with the boundary of the parameter space. In particular, we consider the following mutual positions of the boundary, the parameter set identified by H_0 and the true value θ_0 :

\mathcal{G}_1 . H_0 is the hypothesis that θ_0 belongs to the boundary: $H_0 : g(\theta_0) = 0$;

\mathcal{G}_2 . H_0 is a simple null hypothesis on the boundary: $H_0 : \theta_0 = \bar{\theta}, g(\bar{\theta}) = 0$;

\mathcal{G}_3 . $H_0 : h(\theta_0) = 0$, where $\{\theta \in \mathbb{R}^2 : h(\theta) = 0\}$ is not a subset of the boundary $\partial\Theta$, but meets $\partial\Theta$ at a singleton set.

For example, let again $g(\theta) = \theta_2$. Then the hypothesis of no predictability $H_0 : \theta_{2,0} = 0$ falls under \mathcal{G}_1 ; the hypothesis $H_0 : \theta_0 = (0, 0)'$ that y_t is unpredictable with zero mean falls under \mathcal{G}_2 ; the hypothesis $H_0 : (1, 1)'\theta_0 = \theta_{1,0} + \theta_{2,0} = 0$ falls under \mathcal{G}_3 . In the latter case, the intersection point of the boundary and H_0 is $(0, 0)'$ which might, but need not, be the true value under H_0 .

Let $\hat{\theta}$ be the OLS estimator of the first two coefficients in the equation

$$y_t = \theta_1 + \theta_2 x_{n,t-1} + \delta \Delta x_{n,t} + e_t \quad (4.8)$$

subject to the constraint $\hat{\theta} \in \Theta$, i.e. $g(\hat{\theta}) \geq 0$ (here $\Delta x_{n,t}$ is included in order to obtain residuals asymptotically uncorrelated with the innovations driving $x_{n,t}$). It holds that $n^{1/2}(\hat{\theta} - \theta_0) \xrightarrow{w} \ell(\theta_0)$, with $\ell(\theta_0)$ depending on the position of θ_0 relative to the boundary $\partial\Theta$. Thus, $\ell(\theta_0) = \tilde{\ell} := M^{-1/2}\xi$ if $\theta_0 \in \text{int}(\Theta) := \Theta \setminus \partial\Theta$, where $M := \int \tilde{X}\tilde{X}'$, $\tilde{X} := (1, X)'$, $\xi \sim N(0, \sigma_e^2 I_2)$ is independent of X , and $\sigma_e > 0$, whereas (see Section 12 in the working paper version of Andrews, 1999),

$$n^{1/2}(\hat{\theta} - \theta_0) \xrightarrow{w} \ell(\theta_0) = \ell := \arg \min_{\lambda \in \Lambda := \{\lambda \in \mathbb{R}^2: \dot{g}'\lambda \geq 0\}} \|\lambda - M^{-1/2}\xi\|_M \quad (4.9)$$

if $g(\theta_0) = 0$, where we use the notation $\|x\|_M := (x'Mx)^{1/2}$ for $x \in \mathbb{R}^2$.

Consider now a bootstrap sample generated as

$$y_t^* = \hat{\theta}_1 + \hat{\theta}_2 x_{n,t-1} + \varepsilon_t^*, \quad (4.10)$$

where $\varepsilon_t^* = \hat{\varepsilon}_t w_t^*$, $t = 1, \dots, n$, with $\hat{\varepsilon}_t$ the residuals of (4.8) and w_t i.i.d. $N(0, 1)$, independent of the original data.⁶ Then the distribution of $n^{1/2}(\hat{\theta} - \theta_0)$ could be tentatively approximated by the distribution of $n^{1/2}(\hat{\theta}^* - \hat{\theta})$ conditional on the original data, where $\hat{\theta}^*$ is obtained by regressing y_t^* on $(1, x_{n,t-1})'$ (the term $\Delta x_{n,t}$ is no longer necessary) under the constraint $\hat{\theta}^* \in \Theta^* = \Theta$ (as for the original estimator), i.e., $g(\hat{\theta}^*) \geq 0$; see Andrews (2000).

For $\theta_0 \in \text{int}(\Theta)$, it turns out that the bootstrap statistic converges to a conditional version of the limit of $n^{1/2}(\hat{\theta} - \theta_0)$ found earlier:

$$n^{1/2}(\hat{\theta}^* - \hat{\theta}) = n^{1/2}(\tilde{\theta}^* - \hat{\theta}) + o_p(1) \xrightarrow{w^*} \tilde{\ell}|M, \quad (4.11)$$

where $\tilde{\theta}^*$ denotes the unconstrained OLS estimator from the bootstrap sample.

On the other hand, if $\theta_0 \in \partial\Theta$ the bootstrap statistic converges as follows, jointly with $n^{1/2}(\hat{\theta} - \theta_0)$:

$$n^{1/2}(\hat{\theta}^* - \hat{\theta}) \xrightarrow{w^*} \ell^*|(M, \ell), \quad \ell^* := \arg \min_{\lambda \in \Lambda_\ell^* := \{\lambda \in \mathbb{R}^2: \dot{g}'\lambda \geq -\dot{g}'\ell\}} \|\lambda - M^{-1/2}\xi^*\|_M, \quad (4.12)$$

where $\xi^* \sim N(0, 1)$ is independent of (M, ℓ) ; see Theorem 4.1 below. In contrast with the case $\theta_0 \in \text{int}(\Theta)$, the limit in (4.12) is not a conditional version of the limit of $n^{1/2}(\hat{\theta} - \theta_0)$, inasmuch as Λ_ℓ^* in (4.12) is a *random* half-plane, rather than the original set Λ of (4.9). The reason is that the standard bootstrap does not approximate well the original mutual position of the true value and the boundary, unless $g(\hat{\theta}) = 0$. Other, non-standard bootstraps may be designed in order to provide better approximations, at least under the null hypothesis. This is analyzed next.

4.3.2 UNCONDITIONALLY VALID BOOTSTRAP SCHEMES

In order to unify the discussion of several bootstrap schemes for inference on H_0 under the three cases \mathcal{G}_1 , \mathcal{G}_2 and \mathcal{G}_3 , consider a bootstrap sample generated as in (4.10) and,

⁶The conclusions do not change if, instead of this wild (fixed regressor) bootstrap, a standard residual-based i.i.d. bootstrap or a parametric bootstrap is used.

more generally than before, a bootstrap OLS estimator $\hat{\theta}^*$ constrained to belong to the (random) set

$$\Theta^* := \{\theta \in \mathbb{R}^2 : g(\theta) \geq g^*(\hat{\theta})\}$$

where the function $g^* : \mathbb{R}^2 \rightarrow \mathbb{R}$ is continuously differentiable on some neighborhood of θ_0 and satisfies $g^*(\theta) \leq g(\theta)$ for $\theta \in \Theta$. The standard bootstrap considered in Section 4.3.1 obtains by setting $g^* = 0$ (such that $\Theta^* = \Theta$, the original parameter space). Alternatively, setting $g^* = g$ restricts the bootstrap true value $\hat{\theta}$ to lie on the boundary of the bootstrap parameter space Θ^* (as, in this case, $\Theta^* = \{\theta \in \mathbb{R}^2 : g(\theta) \geq g(\hat{\theta})\}$); see Cavaliere, Nielsen and Rahbek (2017) for an application of this ‘restricted’ bootstrap to the location model. Finally, setting $g^* = g - |g|^{1+\kappa}$ for some $\kappa > 0$ introduces a correction, in the spirit of an alternative to the standard bootstrap mentioned in Andrews (2000,p.403, Method two) and Fang and Santos (2019, Example 2.1), where the bootstrap true value either shrinks to the boundary of the bootstrap parameter space at a proper rate or remains bounded away from this boundary, according to whether θ_0 belongs to the original boundary $\partial\Theta$ or not.⁷

In general, the limit distribution of the resulting bootstrap estimator is random, with randomness depending on both the stochastic regressor and the position of θ_0 relative to the boundary. This distribution is given in the following theorem, where $\dot{g}^* := \frac{\partial}{\partial\theta'} g^*(\theta_0)$.

THEOREM 4.1 *Under the assumptions and the notation introduced above, let a null hypothesis H_0 as in \mathcal{G}_1 – \mathcal{G}_3 hold. Let also $\xi^*|(M, \ell(\theta_0)) \sim N(0, 1)$. Then*

$$(n^{1/2}(\hat{\theta} - \theta_0), (n^{1/2}(\hat{\theta}^* - \hat{\theta})|D_n)) \xrightarrow{w} (\ell(\theta_0), (\ell^*(\theta_0)|(M, \ell(\theta_0)))) , \quad (4.13)$$

where in the case $g^*(\theta_0) < g(\theta_0)$,

$$\ell^*(\theta_0) = \tilde{\ell}^* := M^{-1/2}\xi^* \text{ with } \tilde{\ell}^*|(M, \ell(\theta_0)) \stackrel{d}{=} \tilde{\ell}|M, \quad (4.14)$$

whereas in the case $g^*(\theta_0) = g(\theta_0)$,

$$\ell^*(\theta_0) = \ell^* := \arg \min_{\lambda \in \Lambda_\ell^*} \|\lambda - M^{-1/2}\xi^*\|_M, \quad \Lambda_\ell^* := \{\lambda \in \mathbb{R}^2 : \dot{g}'\lambda \geq (\dot{g}^* - \dot{g})'\ell(\theta_0)\}. \quad (4.15)$$

The following conclusions could be drawn.

(i) Consider first configurations \mathcal{G}_1 and \mathcal{G}_2 under H_0 , such that $g(\theta_0) = 0$. Consider the magnitude order, in probability, of the distance between the bootstrap ‘true’ value $\hat{\theta}$ and the bootstrap boundary $\partial\Theta^*$ as a precision measure for a bootstrap approximation to the geometry of \mathcal{G}_1 and \mathcal{G}_2 . As seen above, the standard bootstrap (corresponding to $g^* = 0$) approximates the geometry up to an exact magnitude order of $n^{-1/2}$, resulting in a situation where the belonging of θ_0 to the boundary contributes to the randomness of limit bootstrap distribution given by (4.12) and (4.15) via conditioning on the rv $\ell(\theta_0) = \ell$. Conversely, bootstrap schemes employing $g^*(\theta_0) = g(\theta_0)$ and $\dot{g}^* = \dot{g}$, such that the bootstrap boundary is tangent to the original boundary at θ_0 , give rise to

⁷Instead of setting $g^* = g - |g|^{1+\kappa}$, one could alternatively set $g^* := g - n^{-\kappa}|g|$ for $\kappa \in (0, \frac{1}{2})$. The results would be unchanged.

approximations of order $o_p(n^{-1/2})$ and all the randomness in the bootstrap limit is due to the properties of the stochastic regressor (via the rv M , as now $\ell^*|(M, \ell) \stackrel{d}{=} \ell|M$; see (4.9) and (4.15)). Moreover, for such schemes the bootstrap mimics a conditional version of the asymptotic distribution of the original estimator: $n^{1/2}(\hat{\theta}^* - \hat{\theta}) \xrightarrow{w^*} \ell|M$. Examples are the ‘restricted’ bootstrap based on $g^* = g$, which replicates the geometry of the original data under \mathbf{H}_0 by putting $\hat{\theta}$ on the bootstrap boundary, and the choices $g^* = g - |g|^{1+\kappa}$ for some $\kappa > 0$.

(ii) Consider now the case in \mathcal{G}_3 , such that $g(\theta_0) = 0$ need not, but may hold under \mathbf{H}_0 . Among the bootstraps considered in (i), the standard one would fail to mimic a conditional version of the original distribution if $g(\theta_0) = 0$, while the ‘restricted’ one would fail if $g(\theta_0) > 0$. As an alternative, consider the bootstrap based on $g^* = g - |g|^{1+\kappa}$ for some $\kappa > 0$, see above. If $\theta_0 \in \partial\Theta$, then this choice puts the bootstrap true value $\hat{\theta}$ at an (asymptotically negligible) distance of $o_p(n^{-1/2})$ from the bootstrap boundary, whereas if $\theta_0 \in \text{int}(\Theta)$, then θ is bounded away from the bootstrap boundary, in probability. This guarantees bootstrap unconditional validity, see (iii) below.

(iii) In general, bootstrap unconditional validity can be evaluated through the following corollary of Theorem 4.1.

COROLLARY 4.1 *Under the assumptions of Theorem 4.1, a necessary and sufficient condition for the convergence*

$$\left(n^{1/2}(\hat{\theta} - \theta_0), (n^{1/2}(\hat{\theta}^* - \hat{\theta})|D_n) \right) \xrightarrow{w} (\ell(\theta_0), (\ell(\theta_0)|M)) \quad (4.16)$$

is that: (i) under \mathcal{G}_1 and \mathcal{G}_2 , $g(\theta_0) = g^*(\theta_0)$ and $\dot{g} = \dot{g}^*$; (ii) under \mathcal{G}_3 , either $g(\theta_0) = g^*(\theta_0)$ and $\dot{g} = \dot{g}^*$, or $g(\theta_0) > \max\{0, g^*(\theta_0)\}$.

Moreover, under (4.16) the bootstrap is unconditionally valid for any pair of statistics $\tau = \phi(n^{1/2}(\hat{\theta} - \theta_0)) + o_p(1)$ and $\tau^* = \phi(n^{1/2}(\hat{\theta}^* - \hat{\theta})) + o_p(1)$, where ϕ is a continuous real function such that the cdf of $\phi(\ell(\theta_0))|M$ is continuous.

The class of functions $g^* = g - |g|^{1+\kappa}$ for $\kappa > 0$ satisfies both conditions (i) and (ii) of the previous corollary; hence, the ensuing bootstrap inference is unconditionally valid under all of \mathcal{G}_1 - \mathcal{G}_3 . In contrast, the standard bootstrap violates condition (i) and, in general, is asymptotically invalid if $g(\theta_0) = 0$. An exception is when the discrepancy between the original and the bootstrap geometry is offset by the use of a test statistic that takes into account the geometric position of the null hypothesis in the original parameter space. The next section focuses on this setup.

4.3.3 UNCONDITIONAL VALIDITY OF ONE-SIDED STANDARD BOOTSTRAP TESTS

Under case \mathcal{G}_1 , consider testing $\mathbf{H}_0 : g(\theta_0) = 0$ against the alternative $\mathbf{H}_1 : g(\theta_0) > 0$ using the standard bootstrap (i.e., with $g^* = 0$). For a test statistic of the form $\tau_n := n^{1/2}g(\hat{\theta})$,⁸ its bootstrap counterpart is given by $\tau_n^* := n^{1/2}(g(\hat{\theta}^*) - g(\hat{\theta}))$ and the (one-sided) bootstrap test rejects for *large* values of the bootstrap p -value $p_n^* := P^*(\tau_n^* \leq \tau_n)$;

⁸What follows easily generalizes to statistics of the form $\tau_n := \phi(n^{1/2}g(\hat{\theta}))$ with $\phi(\cdot)$ strictly increasing and normalized by $\phi(0) = 0$.

equivalently, for small values of $\tilde{p}_n^* := 1 - p_n^*$ (see Remark 3.2). As for $\hat{\theta}^*$, also τ_n^* is affected in the limit by extra randomness due to θ_0 being on the boundary. From (4.13), which reduces to (4.9) and (4.12), it follows by the Delta method that

$$(\tau_n, (\tau_n^* | D_n)) \xrightarrow{w} (\dot{g}'\ell, (\dot{g}'\ell^* | (M, \ell))) = (\dot{g}'\ell, (\max\{-\dot{g}'\ell, \dot{g}'\tilde{\ell}^*\} | (M, \ell))).$$

For τ_n^* , however, the randomness induced by conditioning on ℓ affects the sample paths of the associated random cdf on the negative half-line alone (because $\dot{g}'\ell \geq 0$), and is thus irrelevant for bootstrap tests with nominal levels in $(0, \frac{1}{2})$. Put differently, the bootstrap p -values \tilde{p}_n^* are (asymptotically) uniformly distributed below $\frac{1}{2}$. This follows rigorously from the next generalization of Theorem 3.1 (the proof being analogous), where conditions for unconditional bootstrap validity restricted to a subset of nominal testing levels are formulated.

THEOREM 3.1*. *Let there exist a rv τ and a random element X , both defined on the same probability space, such that the support of τ_n is contained in a closed interval \mathbb{T} (finite or infinite), and $(\tau_n, F_n^*) \xrightarrow{w} (\tau, F)$ in $\mathbb{R} \times D(\mathbb{T})$ for $F_n^*(u) := P(\tau_n^* \leq u | D_n)$ and $F(u) := P(\tau \leq u | X)$, $u \in \mathbb{T}$. If the (possibly random) cdf F is sample-path continuous on \mathbb{T} , then the bootstrap p -value $p_n^* := F_n^*(\tau_n)$ satisfies*

$$P(p_n^* \leq q) \rightarrow q$$

for q such that $q \in F(\mathbb{T})$ a.s.

By Theorem 3.1* with $\mathbb{T} = [0, \infty)$ (which corresponds to the support of τ_n and $\tau := \dot{g}'\ell$), it follows that the standard bootstrap applied to the one-sided statistic τ_n is unconditionally valid for nominal levels in $(0, \frac{1}{2})$.

4.4 BOOTSTRAP TESTS OF PARAMETER CONSTANCY

4.4.1 GENERAL SET UP

Here we apply the results of Section 3 to the classic problem of parameter constancy testing in regression models (Chow, 1960; Quandt, 1960; Nyblom, 1989; Andrews, 1993; Andrews and Ploberger, 1994). Specifically, we deal with bootstrap implementations when the moments of the regressors may be unstable over time; see Hansen (2000) and Zhang and Wu (2012), *inter alia*.

Consider a linear regression model for $y_{nt} \in \mathbb{R}$ given $x_{nt} \in \mathbb{R}^m$, in triangular array notation:

$$y_{nt} = \beta_t' x_{nt} + \varepsilon_{nt} \quad (t = 1, 2, \dots, n). \quad (4.17)$$

The null hypothesis of parameter constancy is $H_0 : \beta_t = \beta_1$ ($t = 2, \dots, n$), which is tested here against the alternative $H_1 : \beta_t = \beta_1 + \theta \mathbb{I}_{\{t \geq n^*\}}$ ($t = 2, \dots, n$), where $n^* := \lfloor r^* n \rfloor$ and $\theta \neq 0$ respectively denote the timing and the magnitude of the possible break,⁹ both assumed unknown to the econometrician. The so-called break fraction r^* belongs to a known closed interval $[\underline{r}, \bar{r}]$ in $(0, 1)$. In order to test H_0 against H_1 , it is customary

⁹We suppress the possible dependence of $\beta_t = \beta_{nt}$ on n with no risk of ambiguities.

to consider the ‘sup F ’ (or ‘sup Wald’) test (Quandt, 1960; Andrews, 1993), based on the statistic $\mathcal{F}_n := \max_{r \in [\underline{r}, \bar{r}]} F_{\lfloor nr \rfloor}$, where $F_{\lfloor nr \rfloor}$ is the usual F statistic for testing the auxiliary null hypothesis that $\theta = 0$ in the regression

$$y_{nt} = \beta' x_{nt} + \theta' x_{nt} \mathbb{I}_{\{t \geq \lfloor rn \rfloor\}} + \varepsilon_{nt}.$$

We make the following assumption, allowing for non-stationarity in the regressors (see also Hansen, 2000, Assumptions 1 and 2).

ASSUMPTION \mathcal{H} . The following conditions on $\{x_{nt}, \varepsilon_{nt}\}$ hold:

- (i) (*mda*) ε_{nt} is a martingale difference array with respect to the current value of x_{nt} and the lagged values of $(x_{nt}, \varepsilon_{nt})$;
- (ii) (*wlln*) ε_{nt}^2 satisfies the law of large numbers $n^{-1} \sum_{t=1}^{\lfloor nr \rfloor} \varepsilon_{nt}^2 \xrightarrow{p} r(E\varepsilon_{nt}^2) = r\sigma^2 > 0$, for all $r \in (0, 1]$;
- (iii) (*non-stationarity*) in $\mathcal{D}_{m \times m} \times \mathcal{D}_{m \times m} \times \mathcal{D}_m$:

$$\left(\frac{1}{n} \sum_{t=1}^{\lfloor n \cdot \rfloor} x_{nt} x'_{nt}, \frac{1}{n\sigma^2} \sum_{t=1}^{\lfloor n \cdot \rfloor} x_{nt} x'_{nt} \varepsilon_{nt}^2, \frac{1}{n^{1/2}\sigma} \sum_{t=1}^{\lfloor n \cdot \rfloor} x_{nt} \varepsilon_{nt} \right) \xrightarrow{w} (M, V, N),$$

where M and V are a.s. continuous and (except at 0) strictly positive-definite valued processes, whereas N , conditionally on $\{V, M\}$, is a zero-mean Gaussian process with covariance kernel $E\{N(r_1)N(r_2)'\} = V(r_1)$ ($0 \leq r_1 \leq r_2 \leq 1$).

REMARK 4.1 A special case of Assumption \mathcal{H} is obtained when the regressors satisfy the weak convergence $x_{n\lfloor n \cdot \rfloor} \xrightarrow{w} U(\cdot)$ in \mathcal{D}_m , such that $M(\cdot) = \int_0^\cdot UU'$. Under extra conditions (e.g., if $\sup_n \sup_{t=1, \dots, n} E|E(\varepsilon_{nt}^2 - \sigma^2 | \mathcal{F}_{n, t-i})| \rightarrow 0$ as $i \rightarrow \infty$ for some filtrations $\mathcal{F}_{n, t}$, $n \in \mathbb{N}$, to which $\{\varepsilon_{nt}^2\}$ is adapted), also $V(\cdot) = \int_0^\cdot UU'$ (see Theorem A.1 of Cavaliere and Taylor, 2009). \square

The null asymptotic distribution of \mathcal{F}_n under Assumption \mathcal{H} is provided in Hansen (2000, Theorem 2):

$$\mathcal{F}_n \xrightarrow{w} \sup_{r \in [\underline{r}, \bar{r}]} \{\tilde{N}(r)' \tilde{M}(r)^{-1} \tilde{N}(r)\} \quad (4.18)$$

with $\tilde{N}(u) := N(u) - M(u)M(1)^{-1}N(1)$ and $\tilde{M}(r) := M(r) - M(r)M(1)^{-1}M(r)$. In the case of (asymptotically) stationary regressors, \mathcal{F}_n converges to the supremum of a squared tied-down Bessel process; see Andrews (1993). In the general case, however, since the asymptotic distribution in (4.18) depends on the joint distribution of the limiting processes M, N, V , which is unspecified under Assumption \mathcal{H} , asymptotic inference based on (4.18) is unfeasible. Simulation methods as the bootstrap can therefore be appealing devices for computing p -values associated with \mathcal{F}_n .

4.4.2 BOOTSTRAP TEST AND RANDOM LIMIT BOOTSTRAP DISTRIBUTION

Following Hansen (2000), we consider here a fixed-regressor wild bootstrap introduced to accommodate possible conditional heteroskedasticity of ε_{nt} . It is based on the residuals \tilde{e}_{nt} from the OLS regression of y_{nt} on x_{nt} and $x_{nt}\mathbb{I}_{\{t \geq \lfloor \tilde{r}n \rfloor\}}$, where $\tilde{r} := \arg \max_{r \in [\underline{r}, \bar{r}]} F_{\lfloor nr \rfloor}$ is the estimated break fraction for the original sample. The bootstrap statistic is

$$\mathcal{F}_n^* := \max_{r \in [\underline{r}, \bar{r}]} F_{\lfloor nr \rfloor}^*,$$

where $F_{\lfloor nr \rfloor}^*$ is the F statistic for the auxiliary null hypothesis that $\theta^* = 0$ in the regression

$$y_t^* = \beta^{*l} x_{nt} + \theta^{*l} x_{nt} \mathbb{I}_{\{t \geq \lfloor rn \rfloor\}} + \text{error}_{nt}^*, \quad (4.19)$$

with bootstrap data $y_t^* := \tilde{e}_{nt} w_t^*$ for an i.i.d. $N(0,1)$ sequence of bootstrap multipliers w_t^* independent of the data.

The weak limit of the bootstrap statistic \mathcal{F}_n^* is stated in the next theorem.

THEOREM 4.2 *Under Assumption \mathcal{H} and under H_0 , it holds that*

$$\mathcal{F}_n^* \xrightarrow{w^*} \sup_{r \in [\underline{r}, \bar{r}]} \left\{ \tilde{N}(r)' \tilde{M}(r)^{-1} \tilde{N}(r) \right\} \Big| (M, V), \quad (4.20)$$

where $\tilde{M}(r)$, $\tilde{N}(r)$ are as in (4.18).

REMARK 4.2 Theorem 4.2 establishes that, in general, the weak limit of the fixed-regressor bootstrap statistic is *random*. In particular, it is distinct from the limit in eq. (4.18) and, as a result, the bootstrap does not estimate consistently the unconditional limit distribution of the statistic \mathcal{F}_n under H_0 (contrary to the claim in Theorem 6 of Hansen, 2000). To illustrate the limiting randomness, consider the case $M = V$ with a scalar regressor $x_{nt} \in \mathbb{R}$. By a change of variable (as in Theorem 3 of Hansen, 2000), convergence (4.20) reduces to

$$\mathcal{F}_n^* \xrightarrow{w^*} \sup_{u \in I(M, \underline{r}, \bar{r})} \left\{ \frac{W(u)^2}{u(1-u)} \right\} \Big| M \quad \text{for} \quad I(M, \underline{r}, \bar{r}) := \left[\frac{M(\underline{r})}{M(1)}, \frac{M(\bar{r})}{M(1)} \right],$$

where W is a standard Brownian bridge on $[0, 1]$, independent of M . As the maximization interval $I(M, \underline{r}, \bar{r})$ depends on M , so does the supremum itself. \square

4.4.3 BOOTSTRAP VALIDITY

Although under Assumption \mathcal{H} the bootstrap does not replicate the asymptotic (unconditional) distribution in (4.18), unconditional bootstrap validity can be established under no further assumptions than Assumption \mathcal{H} , by using the results in Section 3.2. In contrast, despite fixing the regressors across bootstrap samples, if interest is in achieving bootstrap validity conditional on the regressors $X_n := \{x_{nt}\}_{t=1}^n$, further conditions are required; e.g., the following Assumption \mathcal{C} .

ASSUMPTION C. Assumption \mathcal{H} holds and, as random measures on $\mathcal{D}_{m \times m} \times \mathcal{D}_{m \times m} \times \mathcal{D}_m$

$$\left(\frac{1}{n} \sum_{t=1}^{\lfloor n \rfloor} x_{nt} x'_{nt}, \frac{1}{n\sigma^2} \sum_{t=1}^{\lfloor n \rfloor} x_{nt} x'_{nt} \varepsilon_{nt}^2, \frac{1}{n^{1/2}\sigma} \sum_{t=1}^{\lfloor n \rfloor} x_{nt} \varepsilon_{nt} \right) \Bigg| X_n \xrightarrow{w} (M, V, N) | (M, V)$$

jointly with the convergence in Assumption \mathcal{H} (iii).

The results on the validity of the bootstrap parameter constancy tests are summarized in the following theorem.

THEOREM 4.3 *Let the parameter constancy hypothesis H_0 hold for model (4.17). Then, under Assumption \mathcal{H} , the bootstrap based on $\tau_n = \mathcal{F}_n$ and $\tau_n^* = \mathcal{F}_n^*$ is unconditionally valid. If Assumption C holds, then the bootstrap based on \mathcal{F}_n and \mathcal{F}_n^* is valid also conditionally on X_n .*

REMARK 4.3 In the proof of Theorem 4.3, we refer to Theorem 3.1 and Corollary 3.2(a) for establishing respectively unconditional and conditional bootstrap validity. Notice that Assumption C is stronger than Assumption \mathcal{H} due to the fact that – differently from the bootstrap variates w_t^* – the errors $\{\varepsilon_{nt}\}$ need not be independent of $\{x_{nt}\}$. The third DGP of Section 2.2.3 could be used to construct an example, with $x_{nt} := n^{-1/2}x_t$ and $\varepsilon_{nt} := \varepsilon_t$, where Assumption \mathcal{H} (iii) holds but Assumption C does not.

REMARK 4.4 The meaning of ‘jointly’ in Assumption C is given in eq. (A.2). By Lemma A.1(b), the convergence will be automatically joint if in $\mathcal{D}_{m \times m}$, $n^{-1}\sigma^{-2} \sum_{t=1}^{\lfloor n \rfloor} x_{nt} x'_{nt} (\varepsilon_{nt}^2 - E(\varepsilon_{nt}^2 | X_n)) = o_p(1)$, such that $n^{-1}\sigma^{-2} \sum_{t=1}^{\lfloor n \rfloor} x_{nt} x'_{nt} \varepsilon_{nt}^2$ is asymptotically equivalent to an X_n -measurable process. \square

5 CONCLUSIONS

When the distribution of a bootstrap statistic conditional on the data is random in the limit, the bootstrap fails to estimate consistently the asymptotic distribution of the original statistic. In this case, the bootstrap is usually regarded as invalid. Renormalization of the statistic of interest cannot always be used to eliminate the limiting bootstrap randomness (e.g., it cannot be used in any of the four applications discussed in Section 4). We have shown, however, that if (asymptotic) bootstrap validity is defined as (large sample) control over the frequency of correct inferences, then randomness of the limit bootstrap distribution does not imply invalidity of the bootstrap, even without renormalizing the original statistic.

For the asymptotic validity of bootstrap inference, in an unconditional or a conditional sense, we have established sufficient conditions and strategies to verify these conditions in specific applications. The conditions differ mainly in their demands on the dependence structure of the data, and are more restrictive for conditional validity to hold.

We have provided four applications to well-known econometric inference problems which feature randomness of the limit bootstrap distribution. Among the further applications where randomness of the limit bootstrap distribution is likely to appear, and

that could be analyzed using our approach, are bootstrap inference in weakly or partially identified models, inference in time series models with time-varying (stochastic) volatility, inference after model selection, and the bootstrap in high-dimensional models. In addition, the methods we provide for establishing conditional bootstrap validity could be useful in problems involving nuisance parameters that are not consistently estimable under the null hypothesis but where sufficient statistics are available (with the bootstrap being potentially valid conditionally on such statistics).

An important issue not analyzed in the paper is whether the bootstrap can deliver refinements over standard asymptotics in cases where the limit bootstrap measure is random. We have seen in Sections 2 and 4.1 that bootstrap inference in such cases could be exact or close to exact. This seems to suggest that a potential for refinements exists. Moreover, there is also a potential for the bootstrap to inherit the finite-sample refinements offered by conditional asymptotic expansions (in line with Barndorff-Nielsen's p^* -formula, see Barndorff-Nielsen and Cox, 1994, Sec. 6.2), as has been established for some bootstrap procedures (DiCiccio and Young, 2008) in the special case of correctly specified parametric models. The study of such questions requires mathematical tools different from those employed here, and is therefore left for further research.

REFERENCES

- ANDREWS, D.W.K. (1993): Tests for parameter instability and structural change with unknown change point, *Econometrica* 61, 821–856.
- (1997): A conditional Kolmogorov test, *Econometrica* 65, 1097–1128.
- (1999): Estimation when a parameter is on a boundary, *Econometrica*, 67, 1341–1383.
- (2000): Inconsistency of the bootstrap when a parameter is on the boundary of the parameter space, *Econometrica*, 68, 399–405.
- ANDREWS, D.W.K. AND W. PLOBERGER (1994): Optimal tests when a nuisance parameter is present only under the alternative, *Econometrica* 62, 1383–1414.
- ATHREYA, K.B.(1987): Bootstrap of the mean in the infinite variance case, *The Annals of Statistics* 15, 724–731.
- AUE, A., I. BERKES AND L. HORVÀTH (2008): Selection from a stable box, *Bernoulli* 14, 125–139.
- O.E.BARNDORFF-NIELSEN AND D.R. COX (1994): *Inference and Asymptotics*, Chapman & Hall.
- BASAWA, I.V., A.K. MALLIK, W.P. MCCORMICK, J.H. REEVES, AND R.L. TAYLOR (1991): Bootstrapping unstable first-order autoregressive processes, *The Annals of Statistics* 19, 1098–1101.

- BENTKUS, V. (2005): A Lyapunov-type bound in R^d , *Theory of Probability and Its Applications* 49(2), 311–323.
- BERAN, R. (1997): Diagnosing Bootstrap success, *Annals of the Institute of Statistical Mathematics* 49, 1–24.
- BILLINGSLEY, P. (1968): *Convergence of Probability Measures*, John Wiley & Sons, NY.
- CATTANEO, M., M. JANSSON AND K. NAGASAWA (2017): Bootstrap-based inference for cube root consistent estimators, *arXiv:1704.08066*.
- CAVALIERE, G. AND I. GEORGIEV (2019): Inference under random limit bootstrap measures: supplemental material.
- CAVALIERE, G., I. GEORGIEV AND A.M.R. TAYLOR (2016): Sieve-based inference for infinite-variance linear processes, *Annals of Statistics* 44, 1467–1494.
- CAVALIERE, G., H.B. NIELSEN AND A. RAHBK (2015): Bootstrap testing of hypotheses on co-integration relations in vector autoregressive models, *Econometrica*, 83, 813–831.
- (2017): On the consistency of bootstrap testing for a parameter on the boundary of the parameter space, *Journal of Time Series Analysis* 38, 513–534
- CAVALIERE, G. AND A.M.R. TAYLOR (2009): Heteroskedastic time series with a unit root, *Econometric Theory* 25, 1228–1276.
- CHAN, N.H. AND C.Z. WEI (1988): Limiting distributions of Least Squares estimates of unstable autoregressive processes, *Annals of Statistics* 16, 367–401.
- CHOW G. (1960): Tests of equality between sets of coefficients in two linear regressions, *Econometrica* 28, 591–605.
- CRIMALDI I. AND L. PRATELLI (2005): Convergence results for conditional expectations, *Bernoulli* 11, 737–745.
- DASGUPTA A. (2008): *Asymptotic Theory of Statistics and Probability*, Springer-Verlag: Berlin.
- DI CICCIO T. AND G.A. YOUNG (2008): Conditional properties of unconditional parametric bootstrap procedures for inference in exponential families, *Biometrika* 95, 747–758.
- FANG Z. AND A. SANTOS (2019): Inference on directionally differentiable functions, *Review of Economic Studies* 86, 377–412.
- GEORGIEV, I., D. HARVEY, S. LEYBOURNE AND A.M.R. TAYLOR (2019): A bootstrap stationarity test for predictive regression invalidity, *Journal of Business & Economic Statistics* 37, 528–541.

- HALL, P. (1992): *The Bootstrap and Edgeworth Expansion*, Springer-Verlag: Berlin.
- HANSEN B.E. (2000): Testing for structural change in conditional models. *Journal of Econometrics* 97, 93–115.
- HÄUSLER, E. AND H. LUSCHGY (2015): *Stable Convergence and Stable Limit Theorems*, Springer-Verlag: Berlin.
- HOROWITZ, J.L. (2001): The bootstrap. In Heckman, J.J. and E. Leamer (eds.) *Handbook of Econometrics* 5, chapter 52, Elsevier: Amsterdam.
- HOUNYO, U., S. GONÇALVES AND N. MEDDAHI (2017): Bootstrapping pre-averaged realized volatility under market microstructure noise, *Econometric Theory* 33, 791–838.
- JACOD, J., Y. LI, P. MYKLAND, M. PODOLSKIJ AND M. VETTER (2009): Microstructure noise in the continuous case: The pre-averaging approach. *Stochastic Processes and Their Applications* 119, 2249–2276.
- JOHANSEN S. AND B. NIELSEN (2016): Analysis of the Forward Search using some new results for martingales and empirical processes, *Bernoulli* 22, 1131–1183.
- KNIGHT, K. (1989): On the bootstrap of the sample mean in the infinite variance case, *The Annals of Statistics* 17, 1168–1175.
- KALLENBERG O. (1997): *Foundations of Modern Probability*, Springer-Verlag: Berlin.
- KALLENBERG O. (2017): *Random measures: theory and applications*, Springer-Verlag: Berlin.
- LEPAGE, R. AND K. PODGÓRSKY: (1996): Resampling permutations in regression without second moments, *Journal of Multivariate Analysis* 57, 119–141.
- LEPAGE, R., K. PODGÓRSKY AND M. RYZNAR (1997): Strong and conditional invariance principles for samples attracted to stable laws, *Probability Theory and Related Fields* 108, 281–298.
- LINDE, W. (1989): Gaussian measure of translated balls in a Banach space, *Theory of Probability and its Applications* 34, 349–359.
- LOCKHART R. (2012): Conditional limit laws for goodness-of-fit tests, *Bernoulli* 18, 857–882.
- MÜLLER, U.K. AND M. WATSON (2008): Testing models of low frequency variability, *Econometrica* 76, 979–1016.
- NYBLOM J., (1989): Testing for the constancy of parameters over time, *Journal of the American Statistical Association* 84, 223–230.
- PHILLIPS, P.C.B. (2014): On confidence intervals for autoregressive roots and predictive regression, *Econometrica* 82, 1177–1195.

- POLITIS, D., J. ROMANO AND M. WOLF (1999): *Subsampling*, Springer, Berlin.
- PLOBERGER, W. AND W. KRÄMER (1992): The CUSUM test with OLS residuals, *Econometrica* 60, 271–285.
- QUANDT R. (1960): Tests of the hypothesis that a linear regression system obeys two separate regimes, *Journal of the American Statistical Association* 55, 324–330.
- REID N. (1995): The roles of conditioning in inference, *Statistical Science* 10, 138–199.
- RUBSHTEIN B. (1996): A central limit theorem for conditional distributions, In Bergelson V., P. March, J. Rosenblatt (eds.), *Convergence in Ergodic Theory and Probability*, De Gruyter: Berlin.
- SEN, B., M. BANERJEE, AND M. WOODROOFE (2010): Inconsistency of Bootstrap: The Grenander Estimator, *Annals of Statistics* 38, 1953–1977.
- SHAO X. AND D.N. POLITIS (2013): Fixed b subsampling and the block bootstrap: improved confidence sets based on p -value calibration, *Journal of the Royal Statistical Society B* 75, 161–184.
- SWEETING T.J. (1989): On conditional weak convergence, *Journal of Theoretical Probability* 2, 461–474.
- ZHANG, T. AND W. WU (2012): Inference of time-varying regression models, *Annals of Statistics* 40, 1376–1402.

APPENDICES

A WEAK CONVERGENCE IN DISTRIBUTION

In this section we establish some properties of weak convergence in distribution for random elements of Polish spaces. They are useful in applications, in order to verify the high-level conditions of our main theorems, as well as to prove these very theorems. Recall the convention that, throughout, Polish spaces are equipped with their Borel sets. Finite n -tuples of random elements defined on the same probability space are considered as random elements of a product space with the product topology and σ -algebra.

Let (Z_n, X_n) and (Z, X) be random elements such that $Z_n = (Z'_n, Z''_n)$ and $Z = (Z', Z'')$ are $\mathcal{S}'_Z \times \mathcal{S}''_Z$ -valued, whereas $X_n = (X'_n, X''_n)$ and $X = (X', X'')$ are resp. $\mathcal{S}' \times \mathcal{S}''$ -valued and $\mathcal{S}'_X \times \mathcal{S}''_X$ -valued ($n \in \mathbb{N}$). We say that $Z'_n|X'_n \xrightarrow{w} Z'|X'$ and $Z''_n|X''_n \xrightarrow{w} Z''|X''$ jointly (denoted by $(Z'_n|X'_n, Z''_n|X''_n) \xrightarrow{w} (Z'|X', Z''|X'')$) if

$$(E\{h'(Z'_n)|X'_n\}, E\{h''(Z''_n)|X''_n\}) \xrightarrow{w} (E\{h'(Z')|X'\}, E\{h''(Z'')|X''\}) \quad (\text{A.1})$$

for all $h' \in \mathcal{C}_b(\mathcal{S}'_Z)$ and $h'' \in \mathcal{C}_b(\mathcal{S}''_Z)$. Even for $X'_n = X''_n$, this property is weaker than the convergence $(Z'_n, Z''_n)|X'_n \xrightarrow{w} (Z', Z'')|X$ defined by $E\{g(Z'_n, Z''_n)|X'_n\} \xrightarrow{w} E\{g(Z', Z'')|X\}$ for all $g \in \mathcal{C}_b(\mathcal{S}'_Z \times \mathcal{S}''_Z)$. We notice that for $Z'_n = X'_n$, (A.1) reduces to

$$(Z'_n, E\{h''(Z''_n)|X''_n\}) \xrightarrow{w} (Z', E\{h''(Z'')|X''\}) \quad (\text{A.2})$$

for all $h'' \in \mathcal{C}_b(\mathcal{S}_Z'')$ and in this case we write $(Z'_n, (Z''_n|X''_n)) \xrightarrow{w}_w (Z'', (Z'|X'))$ (see Corollary S.2.1 of the accompanying Supplement).

The first lemma given here is divided in two parts. In the first part, we provide conditions for strengthening weak convergence in distribution to weak convergence in probability. The second part, in its simplest form, provides conditions such that the two convergence facts $(Z_n, X_n) \xrightarrow{w} (Z, X)$ and $Z_n|X_n \xrightarrow{w}_w Z|X$ imply the joint convergence $((Z_n|X_n), Z_n, X_n) \xrightarrow{w}_w ((Z|X), Z, X)$.

LEMMA A.1 *Let $\mathcal{S}_Z, \mathcal{S}'_Z, \mathcal{S}_X$ and \mathcal{S}'_X be Polish spaces. Consider the random elements Z_n, Z (\mathcal{S}_Z -valued), Z'_n, Z' (\mathcal{S}'_Z -valued), X_n (\mathcal{S}_X -valued) and X'_n, X (\mathcal{S}'_X -valued) for $n \in \mathbb{N}$. Assume that X'_n are X_n -measurable and $Z_n|X_n \xrightarrow{w}_w Z|X$.*

(a) *If all the considered random elements are defined on the same probability space, $(Z_n, X'_n) \xrightarrow{w} (Z, X)$ and $X'_n \xrightarrow{p} X$, then $Z_n|X_n \xrightarrow{w}_p Z|X$.*

(b) *If $(Z_n, X'_n, Z'_n) \xrightarrow{w} (Z, X, Z')$, then the joint convergence $((Z_n|X_n), Z_n, X'_n, Z'_n) \xrightarrow{w}_w ((Z|X), Z, X, Z')$ holds in the sense that, for all $h \in \mathcal{C}_b(\mathcal{S}_Z)$,*

$$(E\{h(Z_n)|X_n\}, Z_n, X'_n, Z'_n) \xrightarrow{w} (E\{h(Z)|X\}, Z, X, Z'). \quad (\text{A.3})$$

Notice that, by choosing $Z'_n = Z' = 1$, a corollary of Lemma A.1(b) not involving Z'_n and Z' is obtained. It states that $Z_n|X_n \xrightarrow{w}_w Z|X$ and $(Z_n, X'_n) \xrightarrow{w} (Z, X)$ together imply the joint convergence $((Z_n|X_n), Z_n, X'_n) \xrightarrow{w}_w ((Z|X), Z, X)$, provided that X'_n are X_n -measurable.

By means of eq. (A.1) we defined *joint* weak convergence in distribution and denoted it by $(Z'_n|X'_n, Z''_n|X''_n) \xrightarrow{w}_w (Z'|X', Z''|X'')$. We now extend it to

$$((Z'_n|X'_n), (Z''_n|X''_n), Z'''_n) \xrightarrow{w}_w ((Z'|X'), (Z''|X''), Z'''_n), \quad (\text{A.4})$$

defined to mean that

$$(E\{h'(Z'_n)|X'_n\}, E\{h''(Z''_n)|X''_n\}, Z'''_n) \xrightarrow{w} (E\{h'(Z')|X'\}, E\{h''(Z'')|X''\}, Z'''_n) \quad (\text{A.5})$$

for all continuous and bounded real h', h'' with matching domain. The natural equivalence of $((Z'_n|X'_n), (Z''_n|X''_n), Z'''_n) \xrightarrow{w}_w ((Z'|X'), (Z''|X''), Z'''_n)$ and $((Z'_n|X'_n), Z'''_n) \xrightarrow{w}_w ((Z'|X'), Z'''_n)$ holds under separability of the space \mathcal{S}''' where Z'''_n, Z'''_n take values (see Remark S.2.1).

In Lemma A.2(b) below we relate (A.4) to the joint weak convergence of the respective conditional cdf's in the case of rv's Z'_n, Z''_n, Z' and Z'' . Before that, in Lemma A.2(a) we show how joint weak convergence can be strengthened to a.s. weak convergence on a special probability space. For a single convergence $Z'_n|X'_n \xrightarrow{w}_w Z'|X'$, part (a) implies that there exists a Skorokhod representation $(\tilde{Z}'_n, \tilde{X}'_n) \stackrel{d}{=} (Z'_n, X'_n)$, $(\tilde{Z}', \tilde{X}') \stackrel{d}{=} (Z', X')$ such that $\tilde{Z}'_n|\tilde{X}'_n \xrightarrow{w}_{a.s.} \tilde{Z}'|\tilde{X}'$.

LEMMA A.2 *Let $(Z'_n, Z''_n, Z'''_n, X'_n, X''_n)$ and $(Z', Z'', Z'''_n, X', X'')$ be random elements of the same Polish product space, defined on possibly different probability spaces ($n \in \mathbb{N}$).*

(a) *If (A.4)-(A.5) hold, then there exist a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ and random elements $(\tilde{X}'_n, \tilde{X}''_n, \tilde{Z}'_n, \tilde{Z}''_n, \tilde{Z}'''_n) \stackrel{d}{=} (X'_n, X''_n, Z'_n, Z''_n, Z'''_n)$, $(\tilde{X}', \tilde{X}'', \tilde{Z}', \tilde{Z}'', \tilde{Z}'''_n) \stackrel{d}{=} (X', X'', Z', Z'', Z'''_n)$*

(X', X'', Z', Z'', Z''') defined on this space such that $\tilde{Z}'_n | \tilde{X}'_n \xrightarrow{w}_{a.s.} \tilde{Z}' | \tilde{X}'$, $\tilde{Z}''_n | \tilde{X}''_n \xrightarrow{w}_{a.s.} \tilde{Z}'' | \tilde{X}''$ and $\tilde{Z}'''_n \xrightarrow{a.s.} \tilde{Z}'''$.

(b) Let Z', Z'' be rv's and Z''' be \mathcal{S}''' -valued. If the conditional distributions $Z' | X'$ and $Z'' | X''$ are diffuse, then (A.4)-(A.5) is equivalent to the weak convergence of the associated random cdf's:

$$(P(Z'_n \leq \cdot | X'_n), P(Z''_n \leq \cdot | X''_n), Z'''_n) \xrightarrow{w} (P(Z' \leq \cdot | X'), P(Z'' \leq \cdot | X''), Z''') \quad (\text{A.6})$$

as random elements of $\mathcal{D}(\mathbb{R}) \times \mathcal{D}(\mathbb{R}) \times \mathcal{S}'''$.

The definition of the convergence $Z_n | X_n \xrightarrow{w}_w Z | X$ implies that $h(Z_n) | X_n \xrightarrow{w}_w h(Z) | X$ for any continuous $h : \mathcal{S}_Z \rightarrow \mathcal{S}'_Z$ between Polish spaces. A generalization for functions h with a negligible set of discontinuities is provided in the following CMT (for weak convergence a.s. and weak convergence in probability, see Theorem 10 of Sweeting, 1989).

THEOREM A.1 *Let $\mathcal{S}_Z, \mathcal{S}'_Z, \mathcal{S}_X$ and \mathcal{S}'_X be Polish spaces and the random elements Z_n, Z be \mathcal{S}_Z -valued, X_n be \mathcal{S}_X -valued and X be \mathcal{S}'_X -valued. If $Z_n | X_n \xrightarrow{w}_w Z | X$ and $h : \mathcal{S}_Z \rightarrow \mathcal{S}'_Z$ has its set of discontinuity points D_h with $P(Z \in D_h | X) = 0$ a.s., then $h(Z_n) | X_n \xrightarrow{w}_w h(Z) | X$.*

Next, we prove in Theorem A.2 a weak convergence result for iterated conditional expectations. The theorem provides conditions under which the convergence $E(z_n | X_n) \xrightarrow{w} E(z | X', X'')$ implies, upon iteration of the expectations, that $E\{E(z_n | X_n) | X'_n\} \xrightarrow{w} E\{E(z | X', X'') | X'\}$ for rv's z_n, z and for X_n -measurable X'_n . In terms of weak convergence in distribution, the result allows to pass from $Z_n | X_n \xrightarrow{w}_w Z | (X', X'')$ to $Z_n | X'_n \xrightarrow{w}_w Z | X'$. The most transparent conclusions from Theorem A.2 are that, (i), $(E(z_n | X_n), X'_n) \xrightarrow{w} (E(z | X'), X')$ implies $E(z_n | X'_n) \xrightarrow{w} E(z | X')$, and (ii), that $((Z_n | X_n), X'_n) \xrightarrow{w}_w ((Z | X'), X')$ implies $Z_n | X'_n \xrightarrow{w}_w Z | X'$. We need, however, a more elaborate version for joint weak convergence.

THEOREM A.2 *For $n \in \mathbb{N}$, let z_n be integrable rv's, X_n, Y_n and (X'_n, X''_n) be random elements of Polish spaces (say, $\mathcal{S}_X, \mathcal{S}_Y$ and \mathcal{S}'_X), defined on the probability spaces $(\Omega_n, \mathcal{F}_n, P_n)$ and such that (X'_n, X''_n) are X_n -measurable ($n \in \mathbb{N}$). Let also z be an integrable rv and $Y, (X', X'')$ be random elements of the Polish spaces $\mathcal{S}_Y, \mathcal{S}'_X$ defined on a probability space (Ω, \mathcal{F}, P) . If*

$$(E(z_n | X_n), X'_n, X''_n, Y_n) \xrightarrow{w} (E(z | X', X''), X', X'', Y) \quad (\text{A.7})$$

and $X''_n | X'_n \xrightarrow{w}_w X'' | X'$, then $E(z_n | X'_n) \xrightarrow{w} E(z | X')$ jointly with (A.7).

Moreover, let Z_n, Z be random elements of a Polish space \mathcal{S}_Z defined resp. on $(\Omega_n, \mathcal{F}_n, P_n)$ and (Ω, \mathcal{F}, P) . If

$$((Z_n | X_n), X'_n, X''_n, Y_n) \xrightarrow{w}_w ((Z | (X', X'')), X', X'', Y) \quad (\text{A.8})$$

and $X''_n | X'_n \xrightarrow{w}_w X'' | X'$, then

$$((Z_n | X'_n), (Z_n | X_n), X'_n, X''_n, Y_n) \xrightarrow{w}_w ((Z | X'), (Z | (X', X'')), X', X'', Y). \quad (\text{A.9})$$

REMARK A.1 A special case with $X_n'' = X'' = 1$ and $Y_n = Y = 1$ is that where $(E(z_n|X_n), X_n) \xrightarrow{w} (E(z|X'), X')$ such that $X_n''|X_n' \xrightarrow{w} X''|X'$ is trivial, and thus, as a result, $E(z_n|X_n) \xrightarrow{w} E(z|X')$. In terms of conditional distributions, the joint convergence $((Z_n|X_n), X_n) \xrightarrow{w} ((Z|X'), X')$ implies that $((Z_n|X_n'), (Z_n|X_n), X_n') \xrightarrow{w} ((Z|X'), (Z|X'), X')$.

REMARK A.2 Theorem A.2 can be applied to the bootstrap p -value. Let (A.7) hold for $z_n = p_n^*$ and $Y_n = Y = 1$, and let G^* be the conditional cdf of $p^*|(X', X'')$. If $E(G^*|X')$ equals pointwise the cdf of the $U(0, 1)$ distribution, then the convergence $p_n^*|X_n' \xrightarrow{w} p^*|X'$ implied by Theorem A.2 under the condition $X_n''|X_n' \xrightarrow{w} X''|X'$ becomes $p_n^*|X_n' \xrightarrow{w} U(0, 1)$. \square

We conclude the section with a result that is used for establishing the joint convergence of original and bootstrap quantities as an implication of a marginal and a conditional convergence.

LEMMA A.3 Let $(\Omega \times \Omega^*, \mathcal{F} \times \mathcal{F}^*, P \times P^b)$ be a product probability space. Let $D_n : \Omega \rightarrow \mathcal{S}_D$, $W_n^* : \Omega^* \rightarrow \mathcal{S}_W$, $X : \Omega \rightarrow \mathcal{S}_X$ and $Z^* : \Omega \times \Omega^* \rightarrow \mathcal{S}_Z$ ($n \in \mathbb{N}$) be random elements of the Polish spaces \mathcal{S}_D , \mathcal{S}_W , $\mathcal{S}_X = \mathcal{S}'_X \times \mathcal{S}''_X$ and \mathcal{S}_Z . Assume further that X_n are D_n -measurable random elements of \mathcal{S}_X and Z_n are (D_n, W_n^*) -measurable random elements of \mathcal{S}_Z ($n \in \mathbb{N}$). If $X_n \xrightarrow{p} X = (X', X'')$ and $Z_n^*|D_n \xrightarrow{w} Z^*|X'$, then $(Z_n^*, X_n) \xrightarrow{w} (Z, X)$ and $(Z_n^*, X_n)|D_n \xrightarrow{w} (Z, X)|X$ on $\mathcal{S}_Z \times \mathcal{S}_X$, where Z is a random element of \mathcal{S}_Z such that $Z^*|X' \stackrel{d}{=} Z|X' \stackrel{d}{=} Z|X$.

The existence of a random element Z with the specified properties, possibly on an extension of the original probability space, is ensured by Lemma 5.9 of Kallenberg (1997). A well-known special case of Lemma A.3 is that where $X_n = (1, X_n'')$, $X = (1, X'')$, $X_n'' \xrightarrow{p} X''$ and $Z_n^*|X_n'' \xrightarrow{w} X''$ (such that $D_n = X_n''$). Then $(Z_n^*, X_n'') \xrightarrow{w} (Z, X'')$ with $Z|X'' \stackrel{d}{=} Z \stackrel{d}{=} X''$ reducing to the condition that X'' and Z are independent and distributed like X'' (DasGupta, 2008, p.475).

B PROOFS

B.1 PROOFS OF THE RESULTS IN SECTION 2

PROOF OF EQ. (2.8). Let $\hat{\varepsilon}_t := \varepsilon_t - E(\varepsilon_t|\eta_t)$, $t \in \mathbb{N}$. Then $(\hat{\varepsilon}_t, E(\varepsilon_t|\eta_t), \eta_t)'$ is an i.i.d. sequence with diagonal covariance matrix $\text{diag}(\omega_{\varepsilon|\eta}, 1 - \omega_{\varepsilon|\eta}, 1)$, $\omega_{\varepsilon|\eta} := E\{\text{Var}(\varepsilon_t|\eta_t)\} \in (0, 1)$, and it is a standard fact that

$$n^{-1/2}(\sum_{t=1}^{\lfloor n \cdot \rfloor} \hat{\varepsilon}_t, \sum_{t=1}^{\lfloor n \cdot \rfloor} E(\varepsilon_t|\eta_t), \sum_{t=1}^{\lfloor n \cdot \rfloor} \eta_t) \xrightarrow{w} (\omega_{\varepsilon|\eta}^{1/2} B_{y1}, (1 - \omega_{\varepsilon|\eta})^{1/2} B_{y2}, B_\eta) \quad (\text{B.1})$$

in \mathcal{D}_3 , where (B_{y1}, B_{y2}, B_η) is a standard Brownian motion in \mathbb{R}^3 . Further, by the conditional invariance principle of Rubshtein (1996),

$$n^{-1/2} \sum_{t=1}^{\lfloor n \cdot \rfloor} \hat{\varepsilon}_t \Big| \sum_{t=1}^{\lfloor n \cdot \rfloor} \eta_t \xrightarrow{w} \omega_{\varepsilon|\eta}^{1/2} B_{y1} \stackrel{d}{=} \omega_{\varepsilon|\eta}^{1/2} B_{y1} \Big| (B_{y2}, B_\eta) \quad (\text{B.2})$$

as a convergence of random measures on \mathcal{D} . Since $\sigma(\sum_{t=1}^{\lfloor n \rfloor} \eta_t) = \sigma(\sum_{t=1}^{\lfloor n \rfloor} E(\varepsilon_t | \eta_t))$, $\sum_{t=1}^{\lfloor n \rfloor} \eta_t = \sigma(X_n)$, the convergence

$$n^{-1/2} \left(\sum_{t=1}^{\lfloor n \rfloor} \hat{\varepsilon}_t, \sum_{t=1}^{\lfloor n \rfloor} E(\varepsilon_t | \eta_t), \sum_{t=1}^{\lfloor n \rfloor} \eta_t \right) \Big| X_n \xrightarrow{w} (\omega_{\varepsilon|\eta}^{1/2} B_{y1}, (1 - \omega_{\varepsilon|\eta})^{1/2} B_{y2}, B_\eta) \Big| (B_{y2}, B_\eta)$$

follows from (B.1) and (B.2) by Theorem 2.1 of Crimaldi and Pratelli (2005), for random measures on \mathcal{D}_3 . Notice that $(n^{-1} \sum_{t=1}^n \eta_t \hat{\varepsilon}_t, n^{-1} \sum_{t=1}^n \eta_t E(\varepsilon_t | \eta_t)) \xrightarrow{p} 0$ and the convergence is preserved upon conditioning on X_n . Then, by using conditional convergence to stochastic integrals (Theorem 3 of Georgiev et al., 2018), it further follows that

$$\begin{aligned} & \left(n^{-2} M_n, n^{-1} \sum_{t=1}^{\lfloor n \rfloor} x_t \hat{\varepsilon}_t, n^{-1} \sum_{t=1}^{\lfloor n \rfloor} x_t E(\varepsilon_t | \eta_t) \right) \Big| X_n = \\ & \left(n^{-2} M_n, n^{-1} \sum_{t=1}^{\lfloor n \rfloor} x_{t-1} \hat{\varepsilon}_t + o_p(1), n^{-1} \sum_{t=1}^{\lfloor n \rfloor} x_{t-1} E(\varepsilon_t | \eta_t) + o_p(1) \right) \Big| X_n \\ & \xrightarrow{w} (M, \omega_{\varepsilon|\eta}^{1/2} M^{1/2} \xi_1, (1 - \omega_{\varepsilon|\eta})^{1/2} M^{1/2} \xi_2) \Big| (B_{y2}, B_\eta) \end{aligned}$$

with $M := \int B_\eta^2$, $\xi_1 := (\int B_\eta^2)^{-1/2} \int B_\eta dB_{y1}$, $\xi_2 := (\int B_\eta^2)^{-1/2} \int B_\eta dB_{y2}$ jointly independent and $\xi_i \sim N(0, 1)$, $i = 1, 2$. Then, by Theorem A.1, τ_n of (2.6) satisfies

$$(\tau_n, n^{-2} M_n, n^{-1} \sum_{t=1}^n x_t E(\varepsilon_t | \eta_t)) \Big| X_n \xrightarrow{w} (\tau, M, (1 - \omega_{\varepsilon|\eta})^{1/2} M^{1/2} \xi_2) \Big| (B_{y2}, B_\eta) \quad (\text{B.3})$$

with $\tau := M^{-1/2} (\omega_{\varepsilon|\eta}^{1/2} \xi_1 + (1 - \omega_{\varepsilon|\eta})^{1/2} \xi_2)$. This yields (2.8). The bootstrap, instead of estimating consistently the limiting conditional distribution of τ_n given X_n , estimates the random distribution obtained by averaging this limit over ξ_2 . As a result, conditionally on X_n the bootstrap p -value is not asymptotically uniformly distributed:

$$p_n^* | X_n = \Phi(\hat{\omega}_\varepsilon^{-1/2} M_n^{1/2} (\hat{\beta} - \beta)) \Big| X_n \xrightarrow{w} \Phi(\omega_{\varepsilon|\eta}^{1/2} \xi_1 + (1 - \omega_{\varepsilon|\eta})^{1/2} \xi_2) \Big| \xi_2, \quad (\text{B.4})$$

which is not the cdf of a $U(0, 1)$ rv. \square

B.2 PROOFS OF THE RESULTS IN SECTION 3

PROOF OF THEOREM 3.1. The result is a special case of Theorem 3.4 (proved later in an independent manner) with $\tau^* = \tau$ and $F^* = F$. \square

PROOF OF THEOREM 3.2. The result (as well as Corollary 3.1) follows from Theorem 3.3, which is proved below in an independent manner. Specifically, as the conditions of Theorem 3.3 are satisfied, it holds that $P(p_n^* \leq q | X_n) \xrightarrow{w} F(F^{*-1}(q))$. Let $g(\cdot) = \min\{\cdot, 1\} \mathbb{1}_{\{\cdot \geq 0\}}$. By the definition of weak convergence,

$$\begin{aligned} P(p_n^* \leq q) &= E\{g(P(p_n^* \leq q | X_n))\} \xrightarrow{w} E\{g(F(F^{*-1}(q)))\} \\ &= E\{F(F^{*-1}(q))\} = E\{E[F(F^{*-1}(q)) | F^*]\} = E\{F^*(F^{*-1}(q))\} = q \end{aligned}$$

using for the penultimate equality the F^* -measurability of $F^{*-1}(q)$ and the relation $E(F(\gamma) | F^*) = F^*(\gamma)$ for F^* -measurable rv's γ . Thus, $P(p_n^* \leq q) \rightarrow q$ for almost all $q \in (0, 1)$, which proves that $p_n^* \xrightarrow{w} U(0, 1)$. \square

DETAILS OF REMARK 3.8. We justify an assertion of Remark 3.8 regarding condition (†). Let $\tilde{\tau}_n^*$ be a measurable transformation of X_n and W_n^* such that the expansion $\tau_n^* = \tilde{\tau}_n^* + o_p(1)$ holds w.r.t. the probability measure on the space where D_n and W_n^* are jointly defined. Then condition (†) is satisfied with $X'_n := P(\tilde{\tau}_n \leq \cdot | X_n) \in \mathcal{D}(\mathbb{R})$ and $X = F^*$, provided that F^* is sample-path continuous. In fact, under the assumed expansion, it holds that $(\tau_n^* - \tilde{\tau}_n^*)|D_n \xrightarrow{w} 0$ because convergence in probability to zero is preserved upon conditioning. As $\tilde{\tau}_n^*|D_n \stackrel{d}{=} \tilde{\tau}_n^*|X_n$, it follows that the Lévy distance between $F_n^*(\cdot) := P(\tau_n^* \leq \cdot | D_n)$ and $X'_n := P(\tilde{\tau}_n \leq \cdot | X_n)$ is $o_p(1)$, and since the weak limit F^* of F_n^* is sample-path continuous, $F_n^* = X'_n + o_p(1)$ in the uniform distance. Thus, also $X'_n \xrightarrow{w} F^*$. \square

PROOF OF COROLLARY 3.1. Convergence (3.4) implies condition (3.3) with the specified F, F^* by Lemma A.2, since the limit random measures are diffuse. Part (a) follows from Theorem 3.2 with $F = F^*$ (see Remark 3.8), and part (b) from Theorem 3.2 with $F^*(u) = E(F(u)|X')$, $u \in \mathbb{R}$. \square

PROOF OF THEOREM 3.3. Introduce $F_n(\cdot) := P(\tau_n \leq \cdot | X_n)$, $F_n^*(\cdot) := P(\tau_n^* \leq \cdot | D_n)$ and $\tilde{F}_n^*(\cdot) := P(\tau_n^* \leq \cdot | X_n)$ as random elements of $\mathcal{D}(\mathbb{R})$. On the probability space of X' , possibly upon extending it, define $\tau^* := F^{*-1}(\zeta)$ for a rv $\zeta \sim U(0, 1)$ which is independent of X' . Then the convergence $(F_n^*, X'_n) \xrightarrow{w} (F^*, X')$, where F^* is X' -measurable and sample-path continuous, implies that $((\tau_n^*|D_n), X'_n) \xrightarrow{w} ((\tau^*|X'), X')$ by Lemma A.2(b). Since X'_n is D_n -measurable, by Theorem A.2 (see also Remark A.1) it follows that $(\tau_n^*|D_n, \tau_n^*|X'_n) \xrightarrow{w} (\tau^*|X', \tau^*|X')$. Since the conditional cdf F^* of $\tau^*|X'$ is sample-path continuous, for $r_n(\cdot) := F_n^*(\cdot) - P(\tau_n^* \leq \cdot | X'_n)$ it follows that $\sup_{x \in \mathbb{R}} |r_n(x)| \xrightarrow{p} 0$, by using Lemma A.2(b). Then the D_n -measurability of X_n , the X_n -measurability of X'_n and Jensen's inequality yield

$$\left| \tilde{F}_n^*(u) - P(\tau_n^* \leq u | X'_n) \right| = |E\{r_n(u) | X_n\}| \leq E\{|r_n(u)| | X_n\} \leq E\{\sup_{x \in \mathbb{R}} |r_n(x)| | X_n\}$$

for every $u \in \mathbb{R}$, and further,

$$\sup_{\mathbb{R}} \left| \tilde{F}_n^* - P(\tau_n^* \leq \cdot | X'_n) \right| \leq E\{\sup_{x \in \mathbb{R}} |r_n(x)| | X_n\} \xrightarrow{p} 0$$

because the $o_p(1)$ property of $\sup_{x \in \mathbb{R}} |r_n(x)|$ is preserved upon conditioning and because $\sup_{x \in \mathbb{R}} |r_n(x)|$ is bounded. Therefore, $F_n^* = P(\tau_n^* \leq \cdot | X'_n) + r_n = \tilde{F}_n^* + o_p(1)$ uniformly. Then the convergence $(F_n, F_n^*) \xrightarrow{w} (F, F^*)$ in $\mathcal{D}(\mathbb{R})^{\times 2}$ extends to $(F_n, F_n^*, \tilde{F}_n^*) \xrightarrow{w} (F, F^*, F^*)$ in $\mathcal{D}(\mathbb{R})^{\times 3}$.

Fix a $q \in (0, 1)$ at which F^{*-1} is a.s. continuous; such q are all but countably many because F^{*-1} is càdlàg. Here F^{*-1} stands for the right-continuous generalized inverse of F^* , and similarly for other cdf's. It follows from the CMT that $(F_n, F_n^{*-1}(q), \tilde{F}_n^{*-1}(q)) \xrightarrow{w} (F, F^{*-1}(q), F^{*-1}(q))$ in $\mathcal{D}(\mathbb{R}) \times \mathbb{R}^2$. Hence, $F_n^{*-1}(q) = \tilde{F}_n^{*-1}(q) + o_p(1)$ such that $P(|F_n^{*-1}(q) - \tilde{F}_n^{*-1}(q)| < \epsilon) \rightarrow 1$ for all $\epsilon > 0$. With $\mathbb{I}_{n,\epsilon} := \mathbb{I}_{\{|F_n^{*-1}(q) - \tilde{F}_n^{*-1}(q)| < \epsilon\}} = 1 + o_p(1)$, it holds that

$$\begin{aligned} & |P(\tau_n \leq F_n^{*-1}(q) | X_n) - P(\tau_n \leq \tilde{F}_n^{*-1}(q) + \epsilon | X_n)| \\ & \leq \mathbb{I}_{n,\epsilon} |P(\tau_n \leq \tilde{F}_n^{*-1}(q) + \epsilon | X_n) - P(\tau_n \leq \tilde{F}_n^{*-1}(q) - \epsilon | X_n)| + (1 - \mathbb{I}_{n,\epsilon}) \end{aligned}$$

$$= \mathbb{I}_{n,\epsilon} |F_n(\tilde{F}_n^{*-1}(q) + \epsilon) - F_n(\tilde{F}_n^{*-1}(q) - \epsilon)| + (1 - \mathbb{I}_{n,\epsilon}),$$

the equality because $\tilde{F}_n^{*-1}(q) \pm \epsilon$ are X_n -measurable. Using the continuity of F and the CMT, we conclude that the upper bound in the previous display converges weakly to $|F(F^{*-1}(q) + \epsilon) - F(F^{*-1}(q) - \epsilon)|$, which in its turn converges in probability to zero as $\epsilon \rightarrow 0^+$ again by the continuity of F . Therefore,

$$\lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} P\left(|P(\tau_n \leq F_n^{*-1}(q)|X_n) - P(\tau_n \leq \tilde{F}_n^{*-1}(q) + \epsilon|X_n)| > \eta\right) = 0$$

for every $\eta > 0$. On the other hand, as it was already used, by the X_n measurability of $\tilde{F}_n^{*-1}(q) + \epsilon$ and the CMT,

$$P(\tau_n \leq \tilde{F}_n^{*-1}(q) + \epsilon|X_n) = F_n(\tilde{F}_n^{*-1}(q) + \epsilon) \xrightarrow[n \rightarrow \infty]{w} F(F^{*-1}(q) + \epsilon) \xrightarrow[\epsilon \rightarrow 0^+]{w} F(F^{*-1}(q)).$$

Theorem 4.2 of Billingsley (1968) thus yields $P(\tau_n \leq F_n^{*-1}(q)|X_n) \xrightarrow{w} F(F^{*-1}(q))$. The proof of (3.5) is concluded by noting that $P(p_n^* \leq q|X_n)$ differs from $P(\tau_n \leq F_n^{*-1}(q)|X_n)$ by no more than the largest jump of F_n^* , which tends in probability to zero because the weak limit of F_n^* is continuous.

Asymptotic validity of the bootstrap conditional on X_n requires that $F(F^{*-1}(q)) = q$ for almost all $q \in (0, 1)$, which by the continuity of F and F^* reduces to $F = F^*$. \square

PROOF OF COROLLARY 3.2. Part (a) follows from Theorem 3.3 with $F = F^*$ and Polya's theorem. Regarding part (b), $((\tau_n|X_n), (\tau_n^*|D_n), X'_n, X''_n) \xrightarrow{w_w} (\tau|X', X'')$, $(\tau|X'), X', X''$ and $X''_n|X'_n \xrightarrow{w_w} X''|X'$ imply, by Theorem A.2 with $Y_n = E\{g(\tau_n^*)|D_n\}$, $Y = E\{g(\tau)|X'\}$ and an arbitrary $g \in \mathcal{C}_b(\mathbb{R})$, that $(\tau_n|X'_n, \tau_n^*|D_n) \xrightarrow{w_w} (\tau|X', \tau|X')$. As the conditional distribution $\tau|X'$ is diffuse, the proof is completed as in part (a). \square

DETAILS OF REMARK 3.13. By extended Skorokhod coupling (Corollary 5.12 of Kallenberg, 1997), consider a Skorokhod representation of D_n and (τ, τ^*, X, X') such that $(\tau_n, \tau_n^*, \phi_n(X_n), \psi_n(D_n)) \xrightarrow{a.s.} (\tau, \tau^*, X, X')$. Then, by Lemma A.1(a), on the Skorokhod-representation space it holds that $\tau_n|X_n \xrightarrow{w_p} \tau|X$ and $\tau_n^*|D_n \xrightarrow{w_p} \tau^*|X'$, such that on a general probability space $(\tau_n|X_n, \tau_n^*|D_n) \xrightarrow{w_w} (\tau|X, \tau^*|X')$. \square

DETAILS OF REMARK 3.15. With (X'_n, X''_n) and (X', X'') as in Remark 3.15, and with the notation of Section B.1, we argue next that the weak convergence of $\tau_n|X_n, \tau_n^*|D_n$ and (X'_n, X''_n) is joint. Consider a Skorokhod representation of D_n and (M, ξ_1, ξ_2) on a probability space where convergence (B.3) is strengthened to $(\tau_n, X'_n, (X'_n)^{1/2} X''_n)|X_n \xrightarrow{w_{a.s.}} (\tau, M, (1 - \omega_{\varepsilon|\eta})M^{1/2}\xi_2)|(M, \xi_2)$ (by Lemma A.2(a)), and $\hat{\omega}_\varepsilon \xrightarrow{a.s.} \omega_\varepsilon$. Thus, on this space, $\tau_n|X_n \xrightarrow{w_{a.s.}} \tau|(M, \xi_2)$, $(X'_n, X''_n) \xrightarrow{a.s.} (M, (1 - \omega_{\varepsilon|\eta})\xi_2)$, and by (2.4), also $P^*(\tau_n^* \leq u) \xrightarrow{a.s.} \Phi(\omega_\varepsilon^{-1/2} M^{1/2} u)$, $u \in \mathbb{R}$, such that $\tau_n^*|D_n \xrightarrow{w_{a.s.}} \tau|M$. It follows that on a general probability space $((\tau_n|X_n), (\tau_n^*|D_n), X'_n, X''_n) \xrightarrow{w_w} (\tau|(M, \xi_2), (\tau|M), M, (1 - \omega_{\varepsilon|\eta})\xi_2)$. \square

PROOF OF THEOREM 3.4. The random element (τ, F^*) of $\mathbb{R} \times \mathcal{D}(\mathbb{R})$ is a measurable function of (τ, X) determined up to indistinguishability by the joint distribution of (τ, τ^*, X) . By extended Skorokhod coupling (Corollary 5.12 of Kallenberg, 1997), we can regard the data and (τ, τ^*, X) as defined on a special probability space where $(\tau_n, F_n^*) \rightarrow$

(τ, F^*) a.s. in $\mathbb{R} \times \mathcal{D}(\mathbb{R})$ and $F^*(\cdot) = P(\tau^* \leq \cdot | X)$ still holds. We can also replace the redefined F^* by a sample-path continuous random cdf that it is indistinguishable from it (and maintain the notation F^*).

Since F^* is sample-path continuous and F_n^*, F^* are (random) cdf's, $F_n^* \xrightarrow{a.s.} F^*$ in $\mathcal{D}(\mathbb{R})$ implies that $\sup_{u \in \mathbb{R}} |F_n^*(u) - F^*(u)| \xrightarrow{a.s.} 0$. Therefore, $F_n^*(\tau_n) - F^*(\tau_n) \xrightarrow{a.s.} 0$. Since $\tau_n \xrightarrow{a.s.} \tau$ and F^* is sample-path continuous, it holds further that $F^*(\tau_n) \xrightarrow{a.s.} F^*(\tau)$, so also $F_n^*(\tau_n) \xrightarrow{a.s.} F^*(\tau)$ on the special probability space. Hence, in general, $F_n^*(\tau_n) \xrightarrow{w} F^*(\tau)$.

We now find the cdf of $F^*(\tau)$. Again by continuity of F^* and by the choice of F^{*-1} as the right-continuous inverse, the equality of events $\{F^*(u) \leq q\} = \{u \leq F^{*-1}(q)\}$, $q \in (0, 1)$, holds and implies that

$$P(F^*(u)|_{u=\tau} \leq q | X) = P(\tau \leq F^{*-1}(q) | X) = F(F^{*-1}(q)),$$

the latter equality because $F^{*-1}(q)$ is X -measurable. We conclude that $F^*(\tau)$ has cdf $E\{F(F^{*-1}(q))\}$, $q \in (0, 1)$, as asserted. \square

B.3 PROOFS OF THE RESULTS IN SECTION 4

PROOF OF EQ. (4.4). By extended Skorokhod coupling (Corollary 5.12 of Kallenberg, 1997), we can regard the data and X as defined on a single probability space such that $n^{-\alpha/2}x_{[n \cdot]} \xrightarrow{a.s.} X$ in \mathcal{D} . Then, by a product-space construction, we can extend this space to define also an i.i.d. standard Gaussian sequence $\{\varepsilon_t^*\}$ independent of the data and (by Lemma 5.9 of Kallenberg, 1997), a random element (W, b) of $\mathcal{D} \times \mathbb{R}$ such that $(W, b) | X$ has the conditional distribution specified in the text. Consider outcomes (say ω) in the factor-space of $n^{-\alpha/2}x_{[n \cdot]}$ such that $(n^{-\alpha-1}M_n(\omega), n^{-\alpha/2-1}\xi_n(\omega)) \rightarrow (M(\omega), \xi(\omega))$, $n^{-(\alpha+1)/2} \sup |x_{[n \cdot]}(\omega)| \rightarrow 0$ and $M(\omega) > 0$; such outcomes have probability one. For every such outcome, $(n^{1/2}W_n^*, M_n^{1/2}\hat{\beta}^*)$ is tight in $\mathcal{D} \times \mathbb{R}$ because $n^{1/2}W_n^*$ and $M_n^{1/2}\hat{\beta}^*$ are tight in \mathcal{D} and \mathbb{R} resp., and its finite-dimensional distributions converge, by the multivariate Lyapunov CLT (Bentkus, 2005), to those of (W^ω, b^ω) , where W^ω and b^ω are resp. a standard Brownian bridge and a standard Gaussian rv with $\text{Cov}(W^\omega(u), b^\omega) = M(\omega)^{-1/2}\xi(\omega)\psi(u)$, $u \in [0, 1]$. It follows by disintegration (Theorem 5.4 of Kallenberg, 1997) that $(n^{1/2}W_n^*, M_n^{1/2}\hat{\beta}^*)|x_{[n \cdot]} \xrightarrow{w} a.s. (W, b) | X \stackrel{d}{=} (W, b) | (M, \xi)$, and further, that

$$(n^{1/2}W_n^*, M_n^{1/2}\hat{\beta}^*, n^{-\alpha-1}M_n, n^{-\alpha/2-1}\xi_n)|x_{[n \cdot]} \xrightarrow{w} a.s. (W, b, M, \xi) | (M, \xi)$$

by Lemma A.3, since $(n^{-\alpha-1}M_n, n^{-\alpha/2-1}\xi_n)$ are $x_{[n \cdot]}$ -measurable. Still further, by a CMT for a.s. weak convergence (Theorem 10 of Sweeting, 1989),

$$(n^{1/2}W_n^*, n^{(\alpha+1)/2}\hat{\beta}^*, n^{-\alpha/2-1}\xi_n)|x_{[n \cdot]} \xrightarrow{w} a.s. (W, M^{-1/2}b, \xi) | (M, \xi)$$

on the special probability space. This implies (4.4) on a general probability space. \square

PROOF OF EQ. (4.6). Under H_0 , by extended Skorokhod coupling (Corollary 5.12 of Kallenberg, 1997), we regard the data and (X, τ) as defined on a single probability space such that $(n^{-\alpha/2}x_{[n \cdot]}, \tau_n) \xrightarrow{a.s.} (X, \tau)$ in $\mathcal{D} \times \mathbb{R}$, and which is extended to support the independent bootstrap sequence $\{\varepsilon_t^*\}$ and (W, b) such that $(W, b) | X$ has the conditional distribution specified in the text. We have by the same argument as for eq. (4.5)

that, on this space, $\tau_n^* \xrightarrow{w^*}_p \tau|(M, \xi)$ so that $F_n^*(\cdot) := P(\tau_n^* \leq \cdot | D_n) \xrightarrow{p} F(\cdot) := P(\tau \leq \cdot | M, \xi)$ in $\mathcal{D}(\mathbb{R})$, because F is sample-path continuous (*e.g.*, by Proposition 3.2 of Linde, 1989, applied conditionally on M, ξ). As further $\tau_n \xrightarrow{a.s.} \tau$ on this space, we can collect the previous convergence facts into $(\tau_n, F_n^*) \xrightarrow{p} (\tau, F)$, which proves that on a general probability space eq. (4.6) holds.

Notice that (i) $(\tau_n, n^{-\alpha-1}M_n, n^{-\alpha/2-1}\xi_n) \xrightarrow{a.s.} (\tau, M, \xi)$, as implied by the a.s. convergence of τ_n and $n^{-\alpha/2}x_{[n \cdot]}$, and (ii) $\tau_n^* \xrightarrow{w^*}_p \tau|(M, \xi)$, jointly imply, by Lemma A.3, that $(\tau_n, \tau_n^*, n^{-\alpha-1}M_n, n^{-\alpha/2-1}\xi_n) \xrightarrow{w} (\tau, \tau^*, M, \xi)$ with $\tau|(M, \xi) \stackrel{d}{=} \tau^*|(M, \xi)$. The latter convergence remains valid on general probability spaces and is used in the discussion of conditional bootstrap validity. \square

PROOF OF THEOREM 4.1. Introduce $\tilde{x}_t := (1, x_{n,t-1})'$. Let $\mu_n := n^{1/2}(\hat{\theta} - \theta_0)$ and $N_n^* := n^{-1/2} \sum_{t=1}^n \varepsilon_t^* \tilde{x}_t$. Moreover, let the normalized bootstrap estimator be denoted by $\tilde{\mu}_n^* := n^{1/2}(\hat{\theta}^* - \hat{\theta})$; similarly, $\tilde{\mu}_n^* := n^{1/2}(\hat{\theta}^* - \hat{\theta})$, where $\hat{\theta}^*$ is the unrestricted (OLS) bootstrap estimator.

As in Theorem 3 of Georgiev *et al.* (2018), it follows, *mutatis mutandis*, that $(\mu_n, M_n, N_n^*) \xrightarrow{w^*}_w (\ell(\theta_0), M, M^{1/2}\xi^*)|(M, \ell(\theta_0))$ in $\mathbb{R}^{2 \times 4}$, where M is of full rank with probability one, $\xi^*|(M, \ell(\theta_0)) \sim N(0, \sigma_e^2 I_2)$ and σ_e^2 denotes the variance of ε_t corrected for $\Delta x_{n,t}$. To derive the main result, we analyze the properties of μ_n^* on a special probability space where (μ_n, M_n, N_n^*) given the data converge weakly a.s. rather than weakly in distribution. Specifically, by Lemma A.2(a) we can consider a probability space (where $\ell(\theta_0), M$ and, for every $n \in \mathbb{N}$, also the original data and the bootstrap sample can be redefined, maintaining their distribution), such that

$$\mu_n \xrightarrow{a.s.} \ell(\theta_0), M_n \xrightarrow{a.s.} M, N_n^* \xrightarrow{w^*}_{a.s.} M^{1/2}\xi^*|(M, \ell(\theta_0)) \stackrel{d}{=} M^{1/2}\xi^*|M. \quad (\text{B.5})$$

Let $q_n^*(\theta) := n^{-1} \sum_{t=1}^n (y_t^* - \theta' \tilde{x}_t)^2$ and notice that $\tilde{\theta}^* := \arg \min_{\theta \in \mathbb{R}^2} q_n^*(\theta)$ is such that $\tilde{\mu}_n^* := n^{1/2}(\tilde{\theta}^* - \hat{\theta}) = M_n^{-1} N_n^*$. On the special probability space, the asymptotic distribution of $\tilde{\mu}_n^*$ follows from (B.5) and a CMT (Theorem 10 of Sweeting, 1989) as

$$\tilde{\mu}_n^* \xrightarrow{w^*}_{a.s.} \tilde{\ell}^*|(M, \ell(\theta_0)) \stackrel{d}{=} \tilde{\ell}^*|M, \tilde{\ell}^* := \sigma_e^2 M^{-1/2} \xi^*. \quad (\text{B.6})$$

Let us turn to the bootstrap estimator $\hat{\theta}^*$. If $g(\theta_0) > g^*(\theta_0)$, then the consistency facts $\hat{\theta} \xrightarrow{a.s.} \theta_0$ (from (B.5)) and $\tilde{\theta}^* \xrightarrow{w^*}_{a.s.} \theta_0$ (from (B.5)-(B.6)), jointly with the continuity of g, g^* at θ_0 , imply that $P^*(g(\tilde{\theta}^*) \geq g^*(\hat{\theta})) \xrightarrow{a.s.} 1$. Hence, $P^*(\hat{\theta}^* = \tilde{\theta}^*) \xrightarrow{a.s.} 1$ and $P^*(\mu_n^* = \tilde{\mu}_n^*) \xrightarrow{a.s.} 1$. Using also (B.6), it follows that $\mu_n^* \xrightarrow{w^*}_{a.s.} \tilde{\ell}^*|M$ on the special probability space, and since $\mu_n \xrightarrow{a.s.} \ell(\theta_0)$ on this space, it follows further that $(\mu_n, (\mu_n^* | D_n)) \xrightarrow{w}_w (\ell(\theta_0), (\tilde{\ell}^* | M))$ on a general probability space, as asserted in (4.14).

In the case $g^*(\theta_0) = g(\theta_0)$, it still holds that $\hat{\theta}^* = \tilde{\theta}^*$ whenever $g(\tilde{\theta}^*) \geq g^*(\hat{\theta})$. However, the probability of the latter event no longer tends to one. Whenever $g(\tilde{\theta}^*) < g^*(\hat{\theta})$, the estimator $\hat{\theta}^*$ exists if and only if the bootstrap estimator restricted to the bootstrap boundary, say $\check{\theta}^*$, exists. Let $\mathbb{I}_n^* := \mathbb{I}_{\{h(\tilde{\theta}_n^*) \geq 0\}}$ with $h(\theta) := g(\theta) - g^*(\hat{\theta})$. In order to justify the equality

$$\hat{\theta}^* = \tilde{\theta}^* \mathbb{I}_n^* + \check{\theta}^* (1 - \mathbb{I}_n^*) \quad (\text{B.7})$$

in an appropriate sense, we show in Section S.3 of the accompanying Supplement that $\check{\theta}^*(1 - \mathbb{I}_n^*)$ is well-defined with P^* -probability approaching one a.s. We also show there that $\|\check{\theta}^* - \hat{\theta}\|(1 - \mathbb{I}_n^*) = O_{p^*}(n^{-1/2})$ a.s., and as a result, $\|\hat{\theta}^* - \hat{\theta}\| = O_{p^*}(n^{-1/2})$ a.s., using also (B.6). We do not discuss the uniqueness of $\check{\theta}^*$ but we show instead that the measurable minimizers of $q_n^*(\theta)$ over the bootstrap boundary are asymptotically equivalent, as they give rise to the same asymptotic distribution for $\hat{\theta}^*$.

To see this, we further establish in the Supplement that $\check{\theta}^*$ satisfies a first-order condition [foc] with P^* -probability approaching one a.s. Let dots over function names denote differentiation w.r.t. θ (e.g., $\dot{q}_n^*(\theta) := (\partial \dot{q}_n^*/\partial \theta')(\theta)$, a column vector). Then the foc takes the form

$$\{\dot{q}_n^*(\check{\theta}^*) + \check{\delta}_n \dot{h}(\check{\theta}^*)\}(1 - \mathbb{I}_n^*) = \{\dot{q}_n^*(\check{\theta}^*) + \check{\delta}_n \dot{g}(\check{\theta}^*)\}(1 - \mathbb{I}_n^*) = 0, \quad h(\check{\theta}^*) = 0,$$

where $\check{\delta}_n \in \mathbb{R}$. The foc implies, by means of a standard argument, that we can choose a measurable $\bar{\theta}^*$ between $\check{\theta}^*$ and $\hat{\theta}$ such that

$$\{n^{1/2}(\check{\theta}^* - \hat{\theta}) - (I_2 - A_n^* \dot{g}(\bar{\theta}^*))' \tilde{\mu}_n^* - A_n^* n^{1/2} h(\hat{\theta})\}(1 - \mathbb{I}_n^*) = 0,$$

where $A_n^* := M_n^{-1} \dot{g}(\check{\theta}^*) [\dot{g}(\bar{\theta}^*)' M_n^{-1} \dot{g}(\check{\theta}^*)]^{-1}$. As further $\|\check{\theta}^* - \hat{\theta}\|(1 - \mathbb{I}_n^*) = O_{p^*}(n^{-1/2})$ a.s., $\|\bar{\theta}^* - \hat{\theta}\|(1 - \mathbb{I}_n^*) = O_{p^*}(n^{-1/2})$ a.s. and $\hat{\theta} - \theta_0 = O(n^{-1/2})$ a.s., using the continuity of $\dot{g}(\theta)$ at θ_0 it follows that

$$\{n^{1/2}(\check{\theta}^* - \hat{\theta}) - [(I_2 - A^* \dot{g}') \tilde{\mu}_n^* - A^* (\dot{g} - \dot{g}')' n^{1/2} (\hat{\theta} - \theta_0)]\}(1 - \mathbb{I}_n^*) = o_{p^*}(1) \quad \text{a.s.},$$

where $A^* := M^{-1} \dot{g} [\dot{g}' M^{-1} \dot{g}]^{-1}$ and $P^*(|o_{p^*}(1)| > \eta) \xrightarrow{a.s.} 1$ for all $\eta > 0$.

Returning to (B.7), we conclude that

$$n^{1/2}(\hat{\theta}^* - \hat{\theta}) = \tilde{\mu}_n^* \mathbb{I}_n^* + \{(I_2 - A^* \dot{g}') \tilde{\mu}_n^* - A^* (\dot{g} - \dot{g}')' n^{1/2} (\hat{\theta} - \theta_0)\}(1 - \mathbb{I}_n^*) + o_{p^*}(1) \quad \text{a.s.} \quad (\text{B.8})$$

Consider the event indicated by \mathbb{I}_n^* . As $\|\hat{\theta}^* - \hat{\theta}\| = O_{p^*}(n^{-1/2})$ a.s. and $\hat{\theta} - \theta_0 = O(n^{-1/2})$ a.s., by the mean value theorem and the continuous differentiability of g, g^* it holds that

$$n^{1/2} h(\check{\theta}^*) = \dot{g}' \tilde{\mu}_n^* + (\dot{g} - \dot{g}')' \mu_n + c_n,$$

where $c_n = o_{p^*}(1)$ a.s. Then $\mathbb{I}_n \xrightarrow{w^* a.s.} \mathbb{I}_\infty | (M, \ell(\theta_0))$ with $\mathbb{I}_\infty := \mathbb{I}_{\{\dot{g}' \tilde{\ell}^* \geq (\dot{g}^* - \dot{g}')' \ell(\theta_0)\}}$, by (B.5)-(B.6) and the CMT for weak a.s. convergence (Theorem 10 of Sweeting, 1989), as the probability of the limiting discontinuities is 0: $P(\dot{g}' \tilde{\ell}^* = (\dot{g}^* - \dot{g}')' \ell(\theta_0) | (M, \ell(\theta_0))) = 0$ a.s. By exactly the same facts, passage to the limit directly in (B.8) yields

$$n^{1/2}(\hat{\theta}^* - \hat{\theta}) \xrightarrow{w^* a.s.} \{\tilde{\ell}^* \mathbb{I}_\infty + \check{\ell}^* (1 - \mathbb{I}_\infty)\} | (M, \ell(\theta_0)), \quad \check{\ell}^* := (I_2 - A^* \dot{g}') \tilde{\ell}^* - A^* (\dot{g} - \dot{g}')' \ell$$

on the special probability space, where also $\mu_n \xrightarrow{a.s.} \ell(\theta_0)$ by (B.5). Therefore, on a general probability space it holds that

$$(\mu_n, (n^{1/2}(\hat{\theta}^* - \hat{\theta}) | D_n)) \xrightarrow{w^*} (\ell(\theta_0), [\{\tilde{\ell}^* \mathbb{I}_\infty + \check{\ell}^* (1 - \mathbb{I}_\infty)\} | (M, \ell(\theta_0))]).$$

The proof is completed by checking directly that, as asserted in (4.15), $\tilde{\ell}^* \mathbb{I}_\infty + \check{\ell}^* (1 - \mathbb{I}_\infty)$ equals $\arg \min_{\{\dot{g}' \lambda \geq (\dot{g}^* - \dot{g}')' \ell\}} \|\lambda - \tilde{\ell}^*\|_M$ a.s. \square

PROOF OF THEOREM 4.2. Let $(M_n, \tilde{V}_n) := (n^{-1} \sum_{t=1}^{\lfloor n \rfloor} x_{nt} x'_{nt}, n^{-1} \sigma^{-2} \sum_{t=1}^{\lfloor n \rfloor} x_{nt} x'_{nt} \tilde{\varepsilon}_{nt}^2)$. As $\tilde{V}_n = n^{-1} \sigma^{-2} \sum_{t=1}^{\lfloor n \rfloor} x_{nt} x'_{nt} \varepsilon_{nt}^2 + o_p(1)$ under H_0 and Assumption \mathcal{H} , it further holds that $(M_n, \tilde{V}_n) \xrightarrow{w} (M, V)$ in $\mathcal{D}_{m \times m} \times \mathcal{D}_{m \times m}$. The data $D_n := \{x_{nt}, y_{nt}\}_{t=1}^n$ and the bootstrap multipliers $\{w_t^*\}_{t \in \mathbb{N}}$ can be regarded (upon padding with zeroes) as defined on the Polish space $(\mathbb{R}^\infty)^{k+2}$. Therefore, by Corollary 5.12 of Kallenberg (1997), there exists a special probability space where (M, V) , and for every $n \in \mathbb{N}$, also the original and the bootstrap data can be redefined, maintaining their distribution (we also maintain the notation), such that $(M_n, \tilde{V}_n) \xrightarrow{a.s.} (M, V)$.

Consider $N_n^* := n^{-1/2} \sigma^{-1} \sum_{t=1}^{\lfloor n \rfloor} x_{nt} y_t^*$. As N_n^* conditionally on the data is a zero-mean Gaussian process with independent increments and variance function \tilde{V}_n , the argument for Theorem 5 of Hansen (2000) yields the conditional convergence

$$N_n^* | D_n \stackrel{d}{=} N_n^* | (M_n, \tilde{V}_n) \xrightarrow{w}_{a.s.} N | (M, V) \quad (\text{B.9})$$

on the special probability space. The marginal convergence $(M_n, \tilde{V}_n) \xrightarrow{a.s.} (M, V)$ in $(\mathcal{D}_{m \times m})^2$ and (B.9) jointly imply, by Lemma A.3, that

$$(M_n, \tilde{V}_n, N_n^*) \Big| D_n \stackrel{d}{=} (M_n, \tilde{V}_n, N_n^*) \Big| (M_n, \tilde{V}_n) \xrightarrow{w}_p (M, V, N) | (M, V) \quad (\text{B.10})$$

as a convergence of random measures on $(\mathcal{D}_{m \times m})^2 \times \mathcal{D}_m$.

The proof is completed as in Theorems 5 and 6 of Hansen (2000), by using the following uniform expansion in $r \in [\underline{r}, \bar{r}]$: $F_{[nr]}^* = \tilde{F}_n(r) + o_p(1)$ with

$$\tilde{F}_n(r) = \left\| (M_n(r) - M_n(r) M_n(1)^{-1} M_n(r))^{-1/2} (N^*(r) - M_n(r) M_n(1)^{-1} N_n^*(1)) \right\|^2$$

and where convergence is w.r.t. the joint measure over the original and the bootstrap data. As $\tilde{F}_n(r)$ depends on the data only through M_n, \tilde{V}_n , it follows that

$$P^*(\max_{r \in [\underline{r}, \bar{r}]} \tilde{F}_n(r) \leq \cdot) = P(\max_{r \in [\underline{r}, \bar{r}]} \tilde{F}_n(r) \leq \cdot | M_n, \tilde{V}_n),$$

and since $\{\max_{r \in [\underline{r}, \bar{r}]} \tilde{F}_n(r)\} | (M_n, \tilde{V}_n) \xrightarrow{w}_p \mathcal{F} | (M, V)$ by (B.10) and a CMT for weak convergence in probability (Theorem 10 of Sweeting, 1989), with

$$\mathcal{F} := \sup_{r \in [\underline{r}, \bar{r}]} \{\tilde{N}(r)' \tilde{M}(1)^{-1} \tilde{N}(r)\}$$

as in of eq. (4.18), also $\max_{r \in [\underline{r}, \bar{r}]} \tilde{F}_n(r) \xrightarrow{w}_p \mathcal{F} | (M, V)$. Finally, as $\mathcal{F}_n^* := \max_{r \in [\underline{r}, \bar{r}]} F_{[nr]}^* = \max_{r \in [\underline{r}, \bar{r}]} \tilde{F}_n(r) + o_p(1)$ and $\langle \cdot \rangle \xrightarrow{p} 0$ becomes $\langle \cdot \rangle \xrightarrow{w}_p 0$ upon conditioning on the data, we conclude that $\mathcal{F}_n^* \xrightarrow{w}_p \mathcal{F} | (M, V)$ on the special probability space. Then $\mathcal{F}_n^* \xrightarrow{w} \mathcal{F} | (M, V)$ in general. \square

PROOF OF THEOREM 4.3. Additionally to the notation introduced in the proof of Theorem 4.2, let $V_n := n^{-1} \sigma^{-2} \sum_{t=1}^{\lfloor n \rfloor} x_{nt} x'_{nt} \varepsilon_{nt}^2$ and $X_n := \{x_{nt}\}_{t=1}^n$. Under Assumption \mathcal{H} , by Corollary 5.12 of Kallenberg (1997), consider a single probability space where, for

every $n \in \mathbb{N}$, the original and the bootstrap data are redefined together with (M, V, N) , maintaining their distribution (we also maintain the notation), such that

$$\left(M_n, V_n, \tilde{V}_n, \frac{1}{n^{1/2}\sigma} \sum_{t=1}^{\lfloor n \cdot \rfloor} x_{nt} \varepsilon_{nt}, \mathcal{F}_n\right) \xrightarrow{a.s.} (M, V, V, N, \mathcal{F}) \quad (\text{B.11})$$

in $(\mathcal{D}_{m \times m})^3 \times \mathcal{D}_m \times \mathbb{R}$, with $\mathcal{F} := \sup_{r \in [\underline{r}, \bar{r}]} \{\tilde{N}(r)' \tilde{M}(r)^{-1} \tilde{N}(r)\}$ of eq. (4.18). On this space also $\mathcal{F}_n^* \xrightarrow{w^*}_p \mathcal{F} | (M, V)$ holds, by the proof of Theorem 4.2, or equivalently, $P^*(\mathcal{F}_n^* \leq \cdot) \xrightarrow{p} P(\mathcal{F} \leq \cdot | M, V)$ in $\mathcal{D}(\mathbb{R})$, given that sample-path continuity of the conditional cdf $P(\mathcal{F} \leq \cdot | M, V)$ is guaranteed by Proposition 3.2 of Linde (1989) applied conditionally on M, V . We see that $(\mathcal{F}_n, P(\mathcal{F}_n^* \leq \cdot | D_n)) \xrightarrow{p} (\mathcal{F}, P(\mathcal{F} \leq \cdot | M, V))$ on the special probability space. This implies that $(\mathcal{F}_n, P^*(\mathcal{F}_n^* \leq \cdot)) \xrightarrow{w} (\mathcal{F}, P(\mathcal{F} \leq \cdot | M, V))$ in $\mathbb{R} \times \mathcal{D}(\mathbb{R})$ on general probability spaces. Theorem 3.1 becomes applicable and the conclusion of Theorem 4.2 about unconditional validity of the bootstrap follows.

Let now Assumption \mathcal{C} hold. Let the original and the bootstrap data be redefined on another probability space where, by Lemma A.2(a), (B.11) holds (and thus, $\mathcal{F}_n^* \xrightarrow{w^*}_p \mathcal{F} | (M, V)$ by the proof of Theorem 4.2), and additionally, the convergence in Assumption \mathcal{C} holds as an a.s. convergence of random probability measures:

$$\left(M_n, V_n, \frac{1}{n^{1/2}\sigma} \sum_{t=1}^{\lfloor n \cdot \rfloor} x_{nt} \varepsilon_{nt}\right) \Big| X_n \xrightarrow{w}_{a.s.} (M, V, N) | (M, V).$$

By expanding $F_{\lfloor nr \rfloor}$ similarly to $F_{\lfloor nr \rfloor}^*$ in the proof of Theorem 4.2 and applying the CMT of Sweeting (1989, Theorem 10), we can conclude that $\mathcal{F}_n | X_n \xrightarrow{w}_p \mathcal{F} | (M, V)$. Recalling that also $\mathcal{F}_n^* \xrightarrow{w^*}_p \mathcal{F} | (M, V)$, it follows that on a general probability space $(\mathcal{F}_n | X_n, \mathcal{F}_n^* | D_n) \xrightarrow{w}_w (\mathcal{F} | (M, V), \mathcal{F} | (M, V))$.

As previously, the continuity requirement of Corollary 3.2(a) (with $\tau := \mathcal{F}$ and $X = X' := (M, V)$) is satisfied by Proposition 3.2 of Linde (1989) applied conditionally. The bootstrap based on \mathcal{F}_n and \mathcal{F}_n^* is then concluded to be valid conditionally on X_n . \square

INFERENCE UNDER RANDOM LIMIT BOOTSTRAP MEASURES: SUPPLEMENTAL MATERIAL

GIUSEPPE CAVALIERE AND ILIYAN GEORGIEV

November 30, 2019

S.1 INTRODUCTION

This supplement contains in Section S.2 the proofs of the results formulated in Appendix A of Cavaliere and Georgiev (2019), CG hereafter. In Section S.3, it makes available some technical details referred to in the proof of Theorem 4.1 in CG. In Section S.4, the supplement provides a description of the Monte Carlo simulation design used in Section 2 of CG. For notation, see CG. Unless differently specified, all references are to sections, equations and results in CG. Additional references are reported at the end of this document.

S.2 WEAK CONVERGENCE IN DISTRIBUTION: PROOFS

Throughout this supplement, references to extended Skorokhod coupling are based on Corollary 5.12 of Kallenberg (1997). The exposition could sometimes be shortened by explicitly considering the random measures of interest as random elements of a Polish space of measures. To avoid an extra level of abstraction, we do not adopt this perspective.

PROOF OF LEMMA A.1. The proof of part (a) is a straightforward modification of step 1 in the proof of Theorem 2.1 in Crimaldi and Pratelli (2005), where $X'_n = X_n$ is considered. For part (b), let $X'_n = \phi_n(X_n)$ ($n \in \mathbb{N}$) for some measurable functions ϕ_n . Without loss of generality, we can consider that $\mathcal{S}_X = \mathcal{S}'_X$, for otherwise we could identify X_n and X with some random elements (X_n, Y_n) and (Y, X) of $\mathcal{S}_X \times \mathcal{S}'_X$ for arbitrary constant random elements Y_n (\mathcal{S}'_X -valued) and Y (\mathcal{S}_X -valued) defined on the probability spaces of resp. X_n and X . Then, by extended Skorokhod coupling, consider a single probability space supporting $(\tilde{Z}_n, \tilde{X}_n, \tilde{Z}'_n) \stackrel{d}{=} (Z_n, X_n, Z'_n)$ and $(\tilde{Z}, \tilde{X}, \tilde{Z}') \stackrel{d}{=} (Z, X, Z')$ with the respective $\tilde{X}'_n := \phi_n(\tilde{X}_n)$ such that $(\tilde{Z}_n, \tilde{X}'_n, \tilde{Z}'_n) \xrightarrow{a.s.} (\tilde{Z}, \tilde{X}, \tilde{Z}')$. Then also $\tilde{Z}_n | \tilde{X}_n \xrightarrow{w} \tilde{Z} | \tilde{X}$ because weak convergence in distribution is property of the distributions of $(\tilde{Z}_n, \tilde{X}_n)$ and (\tilde{Z}, \tilde{X}) . From part (a) it follows that $\tilde{Z}_n | \tilde{X}_n \xrightarrow{w} \tilde{Z} | \tilde{X}$ such that $E\{h(\tilde{Z}_n) | \tilde{X}_n\} \xrightarrow{p} E\{h(\tilde{Z}) | \tilde{X}\}$ for every $h \in \mathcal{C}_b(\mathcal{S}_Z)$. The convergence $(E\{h(\tilde{Z}_n) | \tilde{X}_n\}, \tilde{Z}_n, \tilde{X}'_n, \tilde{Z}'_n) \xrightarrow{p} (E\{h(\tilde{Z}) | \tilde{X}\}, \tilde{Z}, \tilde{X}, \tilde{Z}')$ for such h implies (A.3) on a general probability space. \square

REMARK S.2.1 We establish here the ‘natural’ fact that the convergence

$$((Z'_n|X'_n), (Z'_n|X'_n), Z''') \xrightarrow{w} ((Z'|X'), (Z'|X'), Z''')$$

is equivalent to $((Z'_n|X'_n), Z''') \xrightarrow{w} ((Z'|X'), Z''')$ under separability of the space \mathcal{S}''' where Z''', Z''' take values. In fact, in this case, (A.5) with $Z'_n = Z''_n$ and $X'_n = X''_n$ is equivalent to

$$(E\{h(Z'_n)|X'_n\}, Z''') \xrightarrow{w} (E\{h(Z')|X'\}, Z''') \quad (\text{S.1})$$

for all continuous and bounded real h with matching domain, since both are equivalent to $uE\{h(Z'_n)|X'_n\} + v\mathbb{I}_{\{Z''' \in A\}} \xrightarrow{w} uE\{h(Z')|X'\} + v\mathbb{I}_{\{Z''' \in A\}}$ for all such h , all continuity sets A of the distribution of Z''' and all $(u, v) \in \mathbb{R}^2$, by Theorem 3.1 of Billingsley (1968) and the Cramér-Wold theorem.

PROOF OF LEMMA A.2(a). Let (A.4)-(A.5) hold. Then $Z'_n \xrightarrow{w} Z'$ such that the sequence of probability measures $\{P'_n\}$ induced by Z'_n is tight. The sequence of conditional measures $Z'_n|X'_n$ has the tight sequence $\{P'_n\}$ as its sequence of average measures. As a result, there exists a countable set of continuous and bounded real functions, say $\{h'_i\}_{i \in \mathbb{N}}$, such that the convergence $E\{h'_i(Z'_n)|X'_n\} \xrightarrow{a.s.} E\{h'_i(Z')|X'\}$, were it to hold for all $i \in \mathbb{N}$, would imply $E\{h'(Z'_n)|X'_n\} \xrightarrow{a.s.} E\{h'(Z')|X'\}$ for all continuous and bounded real h' with the domain of h'_i (by Theorem 2.2 of Berti, Pratelli and Rigo, 2006). Similarly, there exists a sequence of continuous and bounded real functions $\{h''_i\}_{i \in \mathbb{N}}$, such that the convergence $E\{h''_i(Z''_n)|X''_n\} \xrightarrow{a.s.} E\{h''_i(Z'')|X''\}$ for all $i \in \mathbb{N}$ would imply $E\{h''(Z''_n)|X''_n\} \xrightarrow{a.s.} E\{h''(Z'')|X''\}$ for all continuous and bounded real h'' with the domain of h''_i .

Consider the measurable functions H_n with values in $\mathcal{S}''' \times \mathbb{R}^\infty$ defined by

$$H_n(X'_n, X''_n, Z''') = (Z''', \phi'_{n1}(X'_n), \phi''_{n1}(X''_n), \phi'_{n2}(X'_n), \phi''_{n2}(X''_n), \dots)$$

such that a version $\phi'_{ni}(X'_n)$ of $E\{h'_i(Z'_n)|X'_n\}$ and a version $\phi''_{ni}(X''_n)$ of $E\{h''_i(Z''_n)|X''_n\}$ appear resp. at positions $2i$ and $2i + 1$, and the analogous

$$H(X', X'', Z''') = (Z''', \phi'_1(X'), \phi''_1(X''), \phi'_2(X'), \phi''_2(X''), \dots),$$

where $\phi'_i(X')$ and $\phi''_i(X'')$ are versions of resp. $E\{h'_i(Z')|X'\}$ and $E\{h''_i(Z'')|X''\}$ ($i \in \mathbb{N}$). By separability and Theorem 3.1 of Billingsley (1968), $H_n(X'_n, X''_n, Z''') \xrightarrow{w} H(X', X'', Z''')$ in $\mathcal{S}''' \times \mathbb{R}^\infty$ would follow if

$$\begin{aligned} &(\mathbb{I}_{\{Z''' \in A\}}, \phi'_{n1}(X'_n), \phi''_{n1}(X''_n), \phi'_{n2}(X'_n), \phi''_{n2}(X''_n), \dots) \\ &\xrightarrow{w} (\mathbb{I}_{\{Z''' \in A\}}, \phi'_1(X'), \phi''_1(X''), \phi'_2(X'), \phi''_2(X''), \dots) \end{aligned}$$

in \mathbb{R}^∞ for every continuity set A of the distribution of Z''' . The previous is equivalent to weak convergence of the finite-dimensional distributions of the considered sequences for every such A (Billingsley, 1968, p.19). Any linear combination of finitely many functions among $\{h'_i\}_{i \in \mathbb{N}}$ is a bounded and continuous real function, and so for $\{h''_i\}_{i \in \mathbb{N}}$, and for any such two linear combinations (say $h' = \sum_{s=1}^m u_s h'_{i_s}$ and $h'' = \sum_{s=1}^l v_s h''_{j_s}$), it holds that $\sum_{s=1}^m u_s \phi'_{ni_s}(X'_n)$ and $\sum_{s=1}^l v_s \phi''_{nj_s}(X''_n)$ are versions of resp.

$E\{h'(Z'_n)|X'_n\}$ and $E\{h''(Z''_n)|X''_n\}$, whereas $\sum_{s=1}^m u_s \phi'_{i_s}(X')$ and $\sum_{s=1}^l v_s \phi''_{j_s}(X'')$ are versions of resp. $E\{h'(Z')|X'\}$ and $E\{h''(Z'')|X''\}$; therefore,

$$\begin{aligned} & \left(\mathbb{I}_{\{Z'''_n \in A\}}, \sum_{s=1}^m u_s \phi'_{i_s}(X'_n), \sum_{s=1}^l v_s \phi''_{j_s}(X''_n) \right) \\ & \xrightarrow{w} \left(\mathbb{I}_{\{Z''' \in A\}}, \sum_{s=1}^m u_s \phi'_{i_s}(X'), \sum_{s=1}^l v_s \phi''_{j_s}(X'') \right) \end{aligned}$$

by (A.5) and Theorem 3.1 of Billingsley (1968). By the Cramér-Wold theorem, this implies that the finite-dimensional distributions of $H_n(X'_n, X''_n, Z'''_n)$ weakly converge to those of $H(Z', X', X'')$. As a result, $H_n(X'_n, X''_n, Z'''_n) \xrightarrow{w} H(X', X'', Z''')$ in $\mathcal{S}''' \times \mathbb{R}^\infty$.

By extended Skorokhod coupling, $(X'_n, X''_n, Z'_n, Z''_n, Z'''_n)$ and (X', X'', Z', Z'', Z''') can be redefined, maintaining their distribution, on a new probability space where $H_n(X'_n, X''_n, Z'''_n) \xrightarrow{a.s.} H(X', X'', Z''')$ in $\mathcal{S}''' \times \mathbb{R}^\infty$ (we subsume the \sim -notation for the redefined variables). On the new probability space, the relevant components of H_n , H are still versions of the conditional expectations for the redefined variables, for conditional expectations are determined up to equivalence by the underlying joint distributions. As a result, $Z'''_n \xrightarrow{a.s.} Z'''$, $E\{h'_i(Z'_n)|X'_n\} \xrightarrow{a.s.} E\{h'_i(Z')|X'\}$ for all $i \in \mathbb{N}$, and similarly for h''_i . By the choice of $\{h'_i\}_{i \in \mathbb{N}}$ and $\{h''_i\}_{i \in \mathbb{N}}$, on this space $Z'_n|X'_n \xrightarrow{w}_{a.s.} Z'|X'$ and $Z''_n|X''_n \xrightarrow{w}_{a.s.} Z''|X''$.

PROOF OF LEMMA A.2(b). Let Z'_n, Z''_n, Z', Z'' ($n \in \mathbb{N}$) be rv's. By the proof of Kallenberg (2017, Theorem 4.20), on the Skorokhod-coupling space considered in the proof of part (a) it holds that $P(Z'_n \leq \cdot | X'_n) \xrightarrow{a.s.} P(Z' \leq \cdot | X')$ in $\mathcal{D}(\mathbb{R})$ and $P(Z''_n \leq \cdot | X''_n) \xrightarrow{a.s.} P(Z'' \leq \cdot | X'')$ in $\mathcal{D}(\mathbb{R})$. Since on this space also $Z'''_n \xrightarrow{a.s.} Z'''$, (A.6) follows on a general probability space.

Conversely, let (A.6) hold. Notice that $P(Z'_n \leq \cdot | X'_n)$, $P(Z''_n \leq \cdot | X''_n)$, $P(Z' \leq \cdot | X')$ and $P(Z'' \leq \cdot | X'')$ as random elements of $\mathcal{D}(\mathbb{R})$ are measurable transformations of resp. X'_n, X''_n, X' and X'' that are determined up to indistinguishability by the joint distributions of resp. (Z'_n, X'_n) , (Z''_n, X''_n) , (Z', X') and (Z'', X'') . By extended Skorokhod coupling, $(X'_n, X''_n, Z'_n, Z''_n, Z'''_n)$ and (X', X'', Z', Z'', Z''') can be redefined, maintaining their distribution, on a new probability space where $Z'''_n \xrightarrow{a.s.} Z'''$, $P(Z'_n \leq \cdot | X'_n) \xrightarrow{a.s.} P(Z' \leq \cdot | X')$ in $\mathcal{D}(\mathbb{R})$ and $P(Z''_n \leq \cdot | X''_n) \xrightarrow{a.s.} P(Z'' \leq \cdot | X'')$ in $\mathcal{D}(\mathbb{R})$. By the proof of Kallenberg (2017, Theorem 4.20), on this space

$$(E\{h'(Z'_n)|X'_n\}, E\{h''(Z''_n)|X''_n\}, Z'''_n) \xrightarrow{a.s.} (E\{h'(Z')|X'\}, E\{h''(Z'')|X''\}, Z''')$$

for all $h', h'' \in \mathcal{C}_b(\mathbb{R})$, and therefore, (A.5) holds on a general probability space. \square

The following corollary, in its simplest version, established the ‘natural’ equivalence of $Z_n \xrightarrow{w} Z$ and $Z_n|Z_n \xrightarrow{w} Z|Z$ for random elements Z_n, Z of a Polish space.

COROLLARY S.2.1 *Let (Z_n, X_n) and (Z, X) be random elements such that $Z_n = (Z'_n, Z''_n)$ and $Z = (Z', Z'')$ are $\mathcal{S}'_Z \times \mathcal{S}''_Z$ -valued, whereas X_n and X are resp. \mathcal{S} -valued and \mathcal{S}_X -valued ($n \in \mathbb{N}$), with all the mentioned spaces being Polish metric spaces. Then the convergence $(Z'_n|Z'_n, Z''_n|X_n) \xrightarrow{w} (Z'|Z', Z''|X)$ in the sense of (A.1) is equivalent to the*

convergence $(Z'_n, (Z''_n|X_n)) \xrightarrow{w} (Z', (Z''|X))$ in the sense of (A.2). If additionally $\mathcal{S}'_Z = \mathbb{R}$ and the conditional distribution $Z''|X$ is diffuse, then both convergence facts are equivalent to $(Z'_n, P(Z''_n \leq \cdot | X''_n)) \xrightarrow{w} (Z', P(Z'' \leq \cdot | X''))$ as random elements of $\mathcal{S}'_Z \times \mathcal{D}(\mathbb{R})$.

PROOF OF COROLLARY S.2.1. As in the proof of Lemma A.1, there is no loss of generality in assuming the equality $\mathcal{S}_X = \mathcal{S}$. First, let $(Z'_n|Z'_n, Z''_n|X_n) \xrightarrow{w} (Z'|Z', Z''|X)$. By Lemma A.2(a), consider a Skorokhod representation such that $\tilde{Z}'_n|\tilde{Z}'_n \xrightarrow{w}_{a.s.} \tilde{Z}'|\tilde{Z}'$ and $\tilde{Z}''_n|\tilde{X}_n \xrightarrow{w}_{a.s.} \tilde{Z}''|\tilde{X}$. Then $h'(\tilde{Z}'_n) \xrightarrow{a.s.} h'(\tilde{Z}')$ for every $h' \in \mathcal{C}_b(\mathcal{S}'_Z)$. This implies that $\tilde{Z}'_n \xrightarrow{p} \tilde{Z}'$ by the proof of Proposition 4.3(i) of Crimaldi and Pratelli (2005). As further $E\{h''(\tilde{Z}''_n)|\tilde{X}_n\} \xrightarrow{a.s.} E\{h''(\tilde{Z}'')|\tilde{X}\}$ for every $h'' \in \mathcal{C}_b(\mathcal{S}''_Z)$, the convergence $(\tilde{Z}'_n, E\{h''(\tilde{Z}''_n)|\tilde{X}_n\}) \xrightarrow{p} (\tilde{Z}', E\{h''(\tilde{Z}'')|\tilde{X}\})$ implies that $(Z'_n, E\{h''(Z''_n)|X_n\}) \xrightarrow{w} (Z', E\{h''(Z'')|X\})$ for every such h'' , which is (A.2). Second, let $(Z'_n, E\{h''(Z''_n)|X_n\}) \xrightarrow{w} (Z', E\{h''(Z'')|X\})$ hold for every $h'' \in \mathcal{C}_b(\mathcal{S}''_Z)$. Then $(h'(Z'_n), E\{h''(Z''_n)|X_n\}) \xrightarrow{w} (h'(Z'), E\{h''(Z'')|X\})$ for every $h' \in \mathcal{C}_b(\mathcal{S}'_Z)$, by the CMT. The latter statement is equivalent to (A.1) with $Z'_n = X'_n$. Finally, equivalence to the convergence involving the random cdf $P(Z'' \leq \cdot | X'')$ follows from Lemma A.2(b); see also S.2.1. \square

PROOF OF THEOREM A.1. As in the proof of Lemma A.1, without loss of generality, we can assume that $\mathcal{S}_X = \mathcal{S}'_X$. By Lemma A.2(a), consider a Skorokhod representation such that $\tilde{Z}_n|\tilde{X}_n \xrightarrow{w}_{a.s.} \tilde{Z}|\tilde{X}$. Then $h(\tilde{Z}_n)|\tilde{X}_n \xrightarrow{w}_{a.s.} h(\tilde{Z})|\tilde{X}$ on the Skorokhod-representation space by Theorems 8(i) and 10 of Sweeting (1989). Therefore, $h(Z_n)|X_n \xrightarrow{w} h(Z)|X$ on a general probability space. \square

PROOF OF THEOREM A.2. In terms of conditional expectations, the first part of the theorem asserts that if (A.7) holds and $E\{h(X''_n)|X'_n\} \xrightarrow{w} E\{h(X'')|X'\}$ for all continuous and bounded real functions h with matching domain, where (X'_n, X''_n) are X_n -measurable, then the iterated expectations

$$E(z_n|X'_n) = E\{E(z_n|X_n)|X'_n\} \text{ and } E(z|X') = E\{E(z|X', X'')|X'\}$$

satisfy the convergence

$$(E(z_n|X'_n), E(z_n|X_n), X'_n, X''_n, Y_n) \xrightarrow{w} (E(z|X'), E(z|X', X''), X', X'', Y). \quad (\text{S.2})$$

We set up the proof in these terms.

By Theorem 2.1 of Crimaldi and Pratelli (2005), $(X'_n, X''_n) \xrightarrow{w} (X', X'')$ and $X''_n|X'_n \xrightarrow{w} X''|X'$ imply $(X'_n, X''_n)|X'_n \xrightarrow{w} (X', X'')|X'$; i.e., for all $h \in \mathcal{C}_b(\mathcal{S}'_X)$, it holds that $E\{h(X'_n, X''_n)|X'_n\} \xrightarrow{w} E\{h(X', X'')|X'\}$.

Let ϕ_n and ϕ be measurable real functions such that $\phi_n(X_n)$ and $\phi(X', X'')$ are versions respectively of the conditional expectations $E(z_n|X_n)$ and $E(z|X', X'')$. We proceed in two steps. First, we argue that we can redefine (X_n, Y_n) and (X', X'', Y) , maintaining their distribution, on a new probability space where $(\phi_n(X_n), X'_n, X''_n, Y_n) \xrightarrow{a.s.} (\phi(X', X''), X', X'', Y)$ and $E\{h(X'_n, X''_n)|X'_n\} \xrightarrow{p} E\{h(X', X'')|X'\}$ for all $h \in \mathcal{C}_b(\mathcal{S}'_X)$. Second, we show that on this space $E\{\phi_n(X_n)|X'_n\} \xrightarrow{p} E\{\phi(X', X'')|X'\}$, which implies convergence (S.2) on a general probability space.

STEP 1. Let the measurable function ψ_n be such that $(X'_n, X''_n) = \psi_n(X_n)$, thus $(\phi_n(X_n), \psi_n(X_n), Y_n) \xrightarrow{w} (\phi(X', X''), X', X'', Y)$. By extended Skorokhod coupling,

there exist a probability space and random elements $(\tilde{X}_n, \tilde{Y}_n) \stackrel{d}{=} (X_n, Y_n)$, $(\tilde{X}', \tilde{X}'', \tilde{Y}) \stackrel{d}{=} (X', X'', Y)$ defined on this space such that $(\phi_n(\tilde{X}_n), \psi_n(\tilde{X}_n), \tilde{Y}_n) \xrightarrow{a.s.} (\phi(\tilde{X}', \tilde{X}''), \tilde{X}', \tilde{X}'', \tilde{Y})$. On this space it also holds that $E\{h(\tilde{X}'_n, \tilde{X}''_n)|\tilde{X}'_n\} \xrightarrow{w} E\{h(\tilde{X}', \tilde{X}'')|\tilde{X}'\}$ for all $h \in \mathcal{C}_b(\mathcal{S}'_X)$ and $(\tilde{X}'_n, \tilde{X}''_n) := \psi_n(\tilde{X}_n)$, as a consequence of the distributional equalities $(\tilde{X}'_n, \tilde{X}''_n) \stackrel{d}{=} (X'_n, X''_n)$ and $(\tilde{X}', \tilde{X}'') \stackrel{d}{=} (X', X'')$. Moreover, this convergence can be strengthened to $E\{h(\tilde{X}'_n, \tilde{X}''_n)|\tilde{X}'_n\} \xrightarrow{p} E\{h(\tilde{X}', \tilde{X}'')|\tilde{X}'\}$ for all $h \in \mathcal{C}_b(\mathcal{S}'_X)$ by Lemma A.1(a). The next step of the proof takes place in this special probability space (we subsume the $\tilde{\cdot}$ -notation).

STEP 2. As $\mathcal{C}_b(\mathcal{S}'_X)$ is dense in the real functions on \mathcal{S}'_X that are integrable w.r.t. the probability measure induced by (X', X'') , it follows that for every $\varepsilon \in (0, 1)$ there exists a $\phi_\varepsilon \in \mathcal{C}_b(\mathcal{S}'_X)$ such that $E|\phi_\varepsilon(X', X'') - \phi(X', X'')| < (\varepsilon/5)^2$. We decompose

$$\begin{aligned} |E\{\phi_n(X_n)|X'_n\} - E\{\phi(X', X'')|X'\}| &\leq E\{|\phi_n(X_n) - \phi(X', X'')||X'_n\} \\ &\quad + |E\{\phi(X', X'') - \phi_\varepsilon(X', X'')|X'_n\}| \\ &\quad + |E\{\phi_\varepsilon(X', X'') - \phi_\varepsilon(X'_n, X''_n)|X'_n\}| \\ &\quad + |E\{\phi_\varepsilon(X'_n, X''_n)|X'_n\} - E\{\phi_\varepsilon(X', X'')|X'\}| \\ &\quad + |E\{\phi_\varepsilon(X', X'') - \phi(X', X'')|X'\}| \end{aligned}$$

and label the addends on the right-hand side ρ_i , $i = 1, \dots, 5$, in order of appearance. The term ρ_1 is $o_p(1)$ because $|\phi_n(X_n) - \phi(X', X'')| \xrightarrow{p} 0$ and the $o_p(1)$ property is preserved upon conditioning. By Markov's inequality and the choice of ϕ_ε , it follows that $P(\rho_2 \geq \varepsilon/5) \leq \varepsilon/5$. Since $(X'_n, X''_n) \xrightarrow{a.s.} (X', X'')$ and ϕ_ε is continuous, it holds that $|\phi_\varepsilon(X', X'') - \phi_\varepsilon(X'_n, X''_n)| \xrightarrow{a.s.} 0$ and ρ_3 is $o_p(1)$ similarly to ρ_1 . For $h = \phi_\varepsilon$ it holds that $|E\{h(X'_n, X''_n)|X'_n\} - E\{h(X', X'')|X'\}| \xrightarrow{p} 0$ such that $\rho_4 = o_p(1)$. Finally, $P(\rho_5 \geq \varepsilon/5) \leq \varepsilon/5$ by Markov's inequality, similarly to ρ_2 . By combining these results, it follows that

$$P(|E\{\phi_n(X_n)|X'_n\} - E\{\phi(X', X'')|X'\}| \geq \varepsilon) < \varepsilon$$

for large enough n . This proves $E(z_n|X'_n) = E\{\phi_n(X_n)|X'_n\} \xrightarrow{p} E\{\phi(X', X'')|X'\} = E(z|X')$ on the special probability space, and since also $(E(z_n|X_n), X'_n, X''_n, Y_n) \xrightarrow{a.s.} (E(z|X'), X', X'', Y)$ on that space, (S.2) follows on the original probability spaces.

To prove the second part of the theorem, let $h', h'' \in \mathcal{C}_b(\mathcal{S}_Z)$ be arbitrary. By Remark S.2.1, (A.8) implies (A.7) with $z_n = h'(Z_n)$, $Y_n = E\{h''(Z_n)|X_n\}$, $z = h'(Z)$, $Y = E\{h''(Z)|X', X''\}$. By the first part of the theorem and the arbitrariness of h', h'' , (A.9) follows. \square

PROOF OF LEMMA A.3. Let $\{f_j\}_{j \in \mathbb{N}}$ be a convergence-determining countable set of bounded Lipschitz functions $\mathcal{S}_X \times \mathcal{S}_Z \rightarrow \mathbb{R}$ such that $(X_n, Z_n^*) \xrightarrow{w} (X, Z)$ is implied by the convergence

$$E f_j(X_n, Z_n^*) \rightarrow E f_j(X, Z) \text{ as } n \rightarrow \infty \text{ for all } j \in \mathbb{N}. \quad (\text{S.3})$$

The existence of such $\{f_j\}_{j \in \mathbb{N}}$ follows from the proof of Proposition 3.4.4 of Ethier and Kurtz (2005); see also Proposition 2.2 of Worm and Hille (2011). If $\{n_m\}$ is an arbitrary subsequence of the naturals, there exists a further subsequence $\{n_{m_k}\}$ such that

$X_{n_{m_k}} \xrightarrow{\text{a.s.}} X$ and the (random) conditional distribution of $Z_{n_{m_k}}^*$ given $D_{n_{m_k}}$ a.s. converges to the (random) conditional distribution of Z^* given X' (the latter by Corollary 2.4 of Berti, Pratelli and Rigo, 2006). In particular, $E\{h(Z_{n_{m_k}}^*)|D_{n_{m_k}}\} \xrightarrow{\text{a.s.}} E\{h(Z^*)|X'\}$ as $k \rightarrow \infty$ for every $h \in \mathcal{C}_b(\mathcal{S}_Z)$. If we show that $Ef_j(X_{n_{m_k}}, Z_{n_{m_k}}^*) \rightarrow Ef_j(X, Z)$ as $k \rightarrow \infty$ for all $j \in \mathbb{N}$, (S.3) will follow. Hence, without loss of generality we can take $X_n \xrightarrow{\text{a.s.}} X$ and $E\{h(Z_n^*)|D_n\} \xrightarrow{\text{a.s.}} E\{h(Z^*)|X'\}$ for every $h \in \mathcal{C}_b(\mathcal{S}_Z)$, and prove that, as a result, (S.3) holds.

Write $Z_n^* = Z_n^*(D_n, W_n^*)$ and define the measurable functions $\phi_{nj} : \mathcal{S}_X \times \mathcal{S}_D \rightarrow \mathbb{R}$ and $\phi_j : \mathcal{S}_X \times \Omega \rightarrow \mathbb{R}$ by

$$\phi_{nj}(x, d) := E_{P_b}\{f_j(x, Z_n^*(d, W_n^*))\} \text{ and } \phi_j(x, \omega) := \int_{\mathcal{S}_Z} f_j(x, z)\nu(dz, X'(\omega)),$$

where ν is a regular conditional distribution of Z^* given X' . First, we show that there exists an event $A \in \mathcal{F}$ with $P(A) = 1$ such that

$$\phi_{nj}(x, D_n(\omega)) \rightarrow \phi_j(x, \omega) \text{ for all } j \in \mathbb{N}, x \in \mathcal{S}_X, \omega \in A. \quad (\text{S.4})$$

Second, we conclude that $Ef_j(X, Z_n^*) \rightarrow Ef_j(X, Z)$ as $n \rightarrow \infty$ for all $j \in \mathbb{N}$, and then we obtain (S.3).

Let $\{x_i\}_{i \in \mathbb{N}}$ be a countable dense subset of \mathcal{S}_X . As $f_j(x_i, \cdot) \in \mathcal{C}_b(\mathcal{S}_Z)$, it holds that $E\{f_j(x_i, Z_n^*)|D_n\} \xrightarrow{\text{a.s.}} E\{f_j(x_i, Z^*)|X'\}$ (take $h = f_j(x_i, \cdot)$). Since $\phi_{nj}(x_i, D_n)$ and $\phi_j(x_i, \omega)$ are versions of $E\{f_j(x_i, Z_n^*)|D_n\}$ and $E\{f_j(x_i, Z^*)|X'\}$ respectively (see Ex. 10.1.9 of Dudley, 2004, p.341, for the former and Theorem 5.4 of Kallenberg, 1997, for both or the latter), there exist sets $A_{ij} \in \mathcal{F}$ with $P(A_{ij}) = 1$ such that $\phi_{nj}(x_i, D_n(\omega)) \rightarrow \phi_j(x_i, \omega)$ for all $\omega \in A_{ij}$ and every $i, j \in \mathbb{N}$. Define $A := \cap_{i, j \in \mathbb{N}} A_{ij}$ with $P(A) = 1$. It then holds that

$$\phi_{nj}(x_i, D_n(\omega)) \rightarrow \phi_j(x_i, \omega) \text{ for all } i, j \in \mathbb{N}, \omega \in A.$$

Since, for every $x \in \mathcal{S}_X$ and $j \in \mathbb{N}$, $|f_j(x_i, \cdot) - f_j(x, \cdot)| \leq L_j\{\rho_X(x_i, x) \wedge 1\}$ can be made arbitrarily small by an appropriate choice of x_i , where ρ_X is the metric on \mathcal{S}_X and $L_j \in \mathbb{R}$ are Lipschitz constants for f_j , from the definitions of ϕ_{nj} and ϕ_j it follows that $|\phi_{nj}(x_i, D_n(\omega)) - \phi_{nj}(x, D_n(\omega))|$ and $|\phi_j(x_i, \omega) - \phi_j(x, \omega)|$ for every fixed x, j can be made arbitrarily small uniformly over n, ω . From this fact and from the previous display, (S.4) follows for $A = \cap_{i, j \in \mathbb{N}} A_{ij}$.

For arbitrary $j, n \in \mathbb{N}$, (S.4) ensures that $\phi_{nj}(X(\omega), D_n(\omega)) \rightarrow \phi_j(X(\omega), \omega)$ for all $\omega \in A$, and thus, a.s. Since $\phi_{nj}(X, D_n)$ is a version of $E\{f_j(X, Z_n^*)|D_n, X\}$ (by the product structure of the probability space; see Ex. 10.1.9 of Dudley, 2004, p.341), it follows that

$$\begin{aligned} E\{f_j(X, Z_n^*)|D_n, X\} &\xrightarrow{\text{a.s.}} \phi_j(X, \omega) = E\{f_j(x, Z^*)|X'\}|_{x=X} \stackrel{(1)}{=} E\{f_j(x, Z)|X'\}|_{x=X} \\ &\stackrel{(2)}{=} E\{f_j(x, Z)|X\}|_{x=X} \stackrel{(3)}{=} E\{f_j(X, Z)|X\} \text{ a.s.,} \end{aligned}$$

equalities (1) and (2) from $Z^*|X' \stackrel{d}{=} Z|X'$ and equality (3) because for X -measurable ξ 's, $E\{f_j(x, Z)|X\}|_{x=\xi} = E\{f_j(\xi, Z)|X\}$. By the bounded convergence theorem, $Ef_j(X, Z_n^*) \rightarrow Ef_j(X, Z)$.

Next, $|Ef_j(X_n, Z_n^*) - Ef_j(X, Z_n^*)| \leq L_j E\{\rho_X(X_n, X) \wedge 1\} \rightarrow 0$ for every $j \in \mathbb{N}$, again by the bounded convergence theorem, as $X_n \xrightarrow{a.s.} X$. Thus, $Ef_j(X_n, Z_n^*) = Ef_j(X, Z_n^*) + o(1) \rightarrow Ef_j(X, Z)$ and (S.3) is proved. This establishes the convergence $(X_n, Z_n^*) \xrightarrow{w} (X, Z)$.

Finally, $(X_n, Z_n^*) \xrightarrow{w} (X, Z)$ and $Z_n^*|D_n \xrightarrow{w_p} Z|X$, where X_n are D_n -measurable, imply that $(\{(X_n, Z_n^*)|D_n\}, X_n) \xrightarrow{w_w} (\{(X, Z)|X\}, X)$, by a straightforward modification of the proof of Theorem 2.1 in Crimaldi and Pratelli (2005). By Theorem A.2 (see also Remark A.1), the latter convergence implies that

$$((X_n, Z_n^*)|D_n, (X_n, Z_n^*)|X_n) \xrightarrow{w_w} ((X, Z)|X, (X, Z)|X). \quad (\text{S.5})$$

In their turn, $X_n \xrightarrow{p} X$ and $(X_n, Z_n^*)|X_n \xrightarrow{w_w} (X, Z)|X$ imply, by Corollary 4.4 of Crimaldi and Pratelli (2005), that $(X_n, Z_n^*)|X_n \xrightarrow{w_p} (X, Z)|X$, which jointly with (S.5) yields the convergence $(X_n, Z_n^*)|D_n \xrightarrow{w_p} (X, Z)|X$. \square

S.3 DETAILS OF THE PROOF OF THEOREM 4.1

Here we establish the well-definition of $\check{\theta}^*$ in (B.7) and its consistency at the $n^{-1/2}$ rate in the sense that $\|\check{\theta}^* - \hat{\theta}\|(1 - \mathbb{I}_n^*) = O_{p^*}(n^{-1/2})$ a.s.

STEP 1. *Existence of a minimizer of q_n^* over a portion of the bootstrap boundary close to θ_0 .* The choice $(\theta, c) = (\theta_0, g(\theta_0))$ trivially satisfies the equation $g(\theta) = c$. Since g is continuously differentiable in a neighborhood of θ_0 and $\dot{g} = (\dot{g}_1(\theta_0), \dot{g}_2(\theta_0))' \neq 0$ (say that $\dot{g}_1(\theta_0) \neq 0$, with the subscript denoting partial differentiation), by the implicit function theorem there exist an $r > 0$ and a unique function $\gamma : [\theta_{2,0} - r, \theta_{2,0} + r] \times [g(\theta_0) - r, g(\theta_0) + r] \rightarrow [\theta_{1,0} - r, \theta_{1,0} + r]$ such that $\gamma(\theta_{2,0}, 0) = \theta_{1,0}$, $g(\gamma(\theta_2, c), \theta_2) = c$; moreover, γ is continuously differentiable. For outcomes such that $|g^*(\hat{\theta}) - g(\theta_0)| \leq r$, the (non-empty) portion of the bootstrap boundary $\partial\Theta^* = \{\theta \in \mathbb{R}^2 : g(\theta) = g^*(\hat{\theta})\}$ contained in the rectangle $\Pi := [\theta_{1,0} - r, \theta_{1,0} + r] \times [\theta_{2,0} - r, \theta_{2,0} + r]$ can be parameterized as $\theta_1 = \gamma(\theta_2, g^*(\hat{\theta}))$, $\theta_2 \in [\theta_{2,0} - r, \theta_{2,0} + r]$. Define $\check{\theta}^* := (\gamma(\check{\theta}_2^*, g^*(\hat{\theta}^r)), \check{\theta}_2^*)'$, where $\check{\theta}_2^*$ is a measurable minimizer of the continuous function $q_n^*(\gamma(\theta_2, g^*(\hat{\theta}^r)), \theta_2)$ over $\theta_2 \in [\theta_{2,0} - r, \theta_{2,0} + r]$, with $\hat{\theta}^r := \hat{\theta} \mathbb{I}_{\{|g^*(\hat{\theta}) - g(\theta_0)| \leq r\}} + \theta_0 \mathbb{I}_{\{|g^*(\hat{\theta}) - g(\theta_0)| > r\}}$. Since $\mathbb{I}_{\{|g^*(\hat{\theta}) - g(\theta_0)| \leq r\}} \xrightarrow{a.s.} 1$ under $g^*(\theta_0) = g(\theta_0)$, it follows that $\check{\theta}^*$ minimizes q_n^* over $\partial\Theta^* \cap \Pi$ with P^* -probability approaching one a.s.

STEP 2. *Minimization of q_n^* over the entire bootstrap boundary.* For outcomes in

$$A_n := \{|g^*(\hat{\theta}) - g(\theta_0)| \leq r\} \cap \{g(\check{\theta}^*) < g^*(\hat{\theta})\} \cap \{\|\hat{\theta} - \theta_0\| + \|\check{\theta}^* - \hat{\theta}\| \leq \frac{r}{2}\},$$

the minimum of q_n^* over the entire bootstrap boundary $\partial\Theta^*$ exists and is attained only in Π (e.g., at the bootstrap estimator $\check{\theta}^*$ defined in Step 1), provided that

$$\alpha_n := \lambda_{\min}(M_n) \frac{r^2}{4} - \lambda_{\max}(M_n) \|\check{\theta}^* - \hat{\theta}\|^2 > 0.$$

To see this, consider $\theta^c := c\hat{\theta} + (1-c)\check{\theta}^*$ where $c := \inf\{a \in [0, 1] : h(a\hat{\theta} + (1-a)\check{\theta}^*) = 0\}$; θ^c is well-defined whenever $g(\check{\theta}^*) < g^*(\hat{\theta})$ because $g(\hat{\theta}) \geq g^*(\hat{\theta})$ and h is continuous. Moreover, $\theta^c \in \Pi$ for outcomes in A_n because $\|\theta^c - \theta_0\| \leq \|\hat{\theta} - \theta_0\| + \|\check{\theta}^* - \hat{\theta}\| \leq \frac{r}{2}$

and, hence, $q_n^*(\theta^c) \geq q_n^*(\check{\theta}^*)$ for outcomes in A_n , by the minimizing property of $\check{\theta}^*$ on $\partial\Theta^* \cap \Pi$. For any $\theta \notin \Pi$ and outcomes in A_n , we therefore find that

$$\begin{aligned} q_n^*(\theta) - q_n^*(\check{\theta}^*) &\geq q_n^*(\theta) - q_n^*(\theta^c) = q_n^*(\theta) - q_n^*(\tilde{\theta}^*) + q_n^*(\tilde{\theta}^*) - q_n^*(\theta^c) \\ &\geq \lambda_{\min}(M_n)\|\theta - \tilde{\theta}^*\|^2 - \lambda_{\max}(M_n)\|\tilde{\theta}^* - \theta^c\|^2 \\ &\geq \lambda_{\min}(M_n)\{\|\theta - \theta_0\| - \|\tilde{\theta}^* - \theta_0\|\}^2 - \lambda_{\max}(M_n)\|\tilde{\theta}^* - \hat{\theta}\|^2 \\ &\geq \lambda_{\min}(M_n)\frac{r^2}{4} - \lambda_{\max}(M_n)\|\tilde{\theta}^* - \hat{\theta}\|^2 = \alpha_n. \end{aligned}$$

Thus, for outcomes in $A_n \cap \{\alpha_n > 0\}$, q_n^* out of Π is larger than $\min_{\theta \in \partial\Theta^* \cap \Pi} q_n^*(\theta)$, such that $\check{\theta}^*$ minimizes q_n^* over the entire bootstrap boundary.

We find the associated probability

$$\begin{aligned} P^* \left((1 - \mathbb{I}_n^*)q_n^*(\check{\theta}^*) \leq (1 - \mathbb{I}_n^*)q_n(\theta) \quad \forall \theta : h(\theta) = 0 \right) \\ \geq P^* \left(|g^*(\hat{\theta}) - g(\theta_0)| \leq r, \|\hat{\theta} - \theta_0\| \leq \frac{r}{4}, \|\tilde{\theta}^* - \hat{\theta}\| \leq \frac{r}{4}, \alpha_n \geq 0 \right) \\ = \mathbb{I}_{\{|g^*(\hat{\theta}) - g(\theta_0)| \leq r\} \cap \{\|\hat{\theta} - \theta_0\| \leq r/4\}} P^* \left(\|\tilde{\theta}^* - \hat{\theta}\| \leq \frac{r}{4}, \alpha_n \geq 0 \right) \xrightarrow{a.s.} 1 \end{aligned}$$

because $g(\hat{\theta}) \xrightarrow{a.s.} g(\theta_0)$, $\lambda_{\min}(M_n) \rightarrow \lambda_{\min}(M) > 0$ a.s., $\lambda_{\max}(M_n) \rightarrow \lambda_{\max}(M) < \infty$ a.s. and $\|\tilde{\theta}^* - \hat{\theta}\| \xrightarrow{w^*}_{a.s.} 0$. This establishes the fact that $\hat{\theta}^*$ of (B.7), with $\tilde{\theta}^*$ as defined in Step 1, minimizes q_n^* over the bootstrap parameter space Θ^* with P^* -probability approaching one a.s.

STEP 3. *Consistency rate of $\check{\theta}^*$.* Similarly to Step 2, for outcomes in A_n ,

$$0 \geq q_n^*(\check{\theta}^*) - q_n^*(\theta^c) \geq \lambda_{\min}(M_n)\|\check{\theta}^* - \tilde{\theta}^*\|^2 - \lambda_{\max}(M_n)\|\tilde{\theta}^* - \hat{\theta}\|^2,$$

the first inequality by the minimizing property of $\check{\theta}^*$ over $\partial\Theta^* \cap \Pi$. Therefore,

$$\begin{aligned} P^* \left((1 - \mathbb{I}_n^*)\|\check{\theta}^* - \tilde{\theta}^*\|^2 \leq (1 - \mathbb{I}_n^*)\frac{\lambda_{\max}(M_n)}{\lambda_{\min}(M_n)}\|\tilde{\theta}^* - \hat{\theta}\|^2 \right) \\ \geq P^* \left(|g^*(\hat{\theta}) - g(\theta_0)| \leq r, \|\hat{\theta} - \theta_0\| \leq \frac{r}{4}, \|\tilde{\theta}^* - \hat{\theta}\| \leq \frac{r}{4} \right) \\ = \mathbb{I}_{\{|g^*(\hat{\theta}) - g(\theta_0)| \leq r\} \cap \{\|\hat{\theta} - \theta_0\| \leq r/4\}} P^* \left(\|\tilde{\theta}^* - \hat{\theta}\| \leq \frac{r}{4} \right) \xrightarrow{a.s.} 1. \end{aligned}$$

As $\lambda_{\max}(M_n)/\lambda_{\min}(M_n) \xrightarrow{a.s.} \lambda_{\max}(M)/\lambda_{\min}(M)$ and $\|\tilde{\theta}^* - \hat{\theta}\| = O_{P^*}(n^{-1/2})$ P -a.s. (the latter, by (B.6)), it follows that $(1 - \mathbb{I}_n^*)\|\check{\theta}^* - \tilde{\theta}^*\| = O_{P^*}(n^{-1/2})$ P -a.s. and $\|\hat{\theta}^* - \tilde{\theta}^*\| = O_{P^*}(n^{-1/2})$ P -a.s. for $\hat{\theta}^*$ of (B.7). Thus, $\hat{\theta}^*$ has the same consistency rate as $\tilde{\theta}^*$. This argument applies to any $\check{\theta}^*$ which is measurable and minimizes q_n^* over $\partial\Theta^* \cap \Pi$ for outcomes in A_n . This completes Step 3.

Finally, consider the first-order condition [foc] for minimization of q_n^* on $\partial\Theta^*$. As $\|\check{\theta}^* - \theta_0\|(1 - \mathbb{I}_n^*) \leq \{\|\check{\theta}^* - \tilde{\theta}^*\| + \|\tilde{\theta}^* - \theta_0\|\}(1 - \mathbb{I}_n^*) \xrightarrow{w^*}_{a.s.} 0$, it follows that $\mathbb{I}_{\{\check{\theta}^* \in \text{int}(\Pi)\}}(1 - \mathbb{I}_n^*) + \mathbb{I}_n^* \xrightarrow{w^*}_{a.s.} 1$. As additionally $\dot{g}(\theta_0) \neq 0$, by continuity of $\dot{g}(\theta) := (\partial g / \partial \theta')(\theta)$, the foc takes the form

$$P^* \left(\{\dot{q}_n(\check{\theta}^*) + \check{\delta}_n \dot{g}(\check{\theta}^*)\}(1 - \mathbb{I}_n^*) = 0 \right) \xrightarrow{a.s.} 1,$$

where $\check{\delta}_n \in \mathbb{R}$ are measurable and can be determined from the constraint $h(\check{\theta}^*) = 0$.

S.4 SIMULATION DESIGN

We provide here a description of the Monte Carlo [MC] simulation design used for the linear regression model of Section 2. Data $D_n := \{y_t, x_t\}_{t=1}^n$ are generated according to eq. (2.1) and the object of interest is inference on β based on $\tau_n := n^{1/2}(\hat{\beta} - \beta)$, with $\hat{\beta}$ denoting the OLS estimator of β , see Section 2.1. We consider the case where $x_t = \sum_{s=1}^t \eta_s$ is a non-stationary ($I(1)$) process under the following three different distributional structures for (ε_t, η_t) :

- (i) (ε_t, η_t) is i.i.d. $N(0, I_2)$ such that if the true variance $\hat{\omega}_\varepsilon = 1$ was used in eq. (2.2), the bootstrap would perform exact conditional inference (see Remark 2.1);
- (ii) $\varepsilon_t = \zeta_t(1 + 0.3\varepsilon_{t-1}^2 + 0.3\eta_{t-1}^2)^{1/2}$ and $\eta_t = \xi_t(1 + 0.6\eta_{t-1}^2)^{1/2}$, where (ζ_t, ξ_t) is i.i.d. $N(0, I_2)$; this corresponds to a stationary and ergodic conditionally heteroskedastic process with non Gaussian unconditional marginals;
- (iii) $\eta_t = \xi_t(1 + \delta \mathbb{I}_{\{\varepsilon_t \leq 0\}})$, where (ε_t, ξ_t) is i.i.d. $N(0, I_2)$ and $\delta = 9$.

For model (ii) we initialize the process by setting the conditional variance equal to its unconditional expectation (results do not change if a burn-in method is employed instead). The bootstrap is implemented as in Section 2.1, with $\hat{\omega}_\varepsilon$ chosen as the OLS residual variance. For DGP (i) bootstrap inference is close to exact (see Remark 2.1) and it holds that the bootstrap p -value p_n^* satisfies $p_n^* \stackrel{d}{=} U(0, 1) + O_p(n^{-1/2})$. DGP (ii) satisfies the conditions discussed in Section 2.2.2 and bootstrap inference is valid conditionally on $X_n := \{x_t\}_{t=1}^n$; i.e., $p_n^* | X_n \xrightarrow{w} U(0, 1)$. For DGP (iii), on the other hand, bootstrap inference is not valid conditionally on this X_n , but it is valid unconditionally; see Section 2.2.3. Hence, $p_n^* \xrightarrow{w} U(0, 1)$ while $p_n^* | X_n$ has a random limit distribution. Notice that since the bootstrap statistic is conditionally Gaussian, p -values can be obtained without resorting to simulation.

Standard MC experiments generating $D_n = (y_1, \dots, y_n, X_n)$ at each MC iteration allow estimation of the unconditional distribution of the p -value p_n^* , rather than its distribution conditional on X_n (see e.g. Hansen, 2000, footnote 11). To simulate the distribution of p_n^* conditional on X_n , we implement a double MC design where, for each $m = 1, \dots, M$, we generate the regressors $X_n^{(m)} \sim X_n$ and then, for each $v = 1, \dots, N$, we generate data $(y_1^{(m,v)}, \dots, y_n^{(m,v)})$ from their distribution conditional on $X_n = X_n^{(m)}$. The respective statistics $\tau_n^{(m,v)}$ and the associated bootstrap p -values, $p_n^{*(m,v)}$, are used to estimate the conditional distribution of $p_n^* | \{X_n = X_n^{(m)}\}$, for each m , by the empirical cdf $N^{-1} \sum_{v=1}^N \mathbb{I}_{\{p_n^{*(m,v)} \leq \cdot\}}$. We set $M = 1,000$ and $N = 100,000$ throughout. Notice that for model (iii), once the regressor x_t (hence, η_t) is generated, simulation conditional on $\{x_t\}$ requires drawing from the conditional distribution of ε_t given η_t . A simple application of the Bayes rule yields that

$$P(\varepsilon_t \leq 0 | \eta_t) = \frac{1}{1 + (1 + \delta)e^{-\frac{\eta_t^2}{2} \frac{\delta(2+\delta)}{(1+\delta)^2}}} =: p_t$$

and hence that $\varepsilon_t | \{x_t\} \stackrel{d}{=} \varepsilon_t | \eta_t \sim |\xi_t| s_t$, where s_t is a random sign equal to -1 with probability p_t .

All computations have been performed using MATLAB R2019b. Code is available from the Authors upon request.

ADDITIONAL REFERENCES

- BERTI, P., L. PRATELLI AND P. RIGO (2006): Almost sure weak convergence of random probability measures, *Stochastics* 78, 91–97.
- CAVALIERE, G. AND I. GEORGIEV (2019): Inference under random limit bootstrap measures. Under revision at *Econometrica*.
- DUDLEY, R.M (2004): *Real Analysis and Probability*, Cambridge University Press, Cambridge.
- ETHIER, S AND T. KURTZ (2005): *Markov Processes. Characterization and Convergence*, John Wiley&Sons, Hoboken.
- WORM, D AND S. HILLE (2011): Ergodic decompositions associated with regular Markov operators on Polish spaces, *Ergodic Theory and Dynamical Systems* 31, 571-597.