

Equivalence Relations for Computing Permutation Polynomials

Sergey Bereg* Brian Malouf* Linda Morales* Thomas Stanley*
 I. Hal Sudborough* Alexander Wong*

November 25, 2021

Abstract

We present a new technique for computing permutation polynomials based on equivalence relations. The equivalence relations are defined by expanded normalization operations and new functions that map permutation polynomials (PPs) to other PPs. Our expanded normalization applies to almost all PPs, including when the characteristic of the finite field divides the degree of the polynomial. The equivalence relations make it possible to reduce the size of the space, when doing an exhaustive search. As a result, we have been able to compute almost all permutation polynomials of degree d at most 10 over $GF(q)$, where q is at most 97. We have also been able to compute nPPs of degrees 11 and 12 in a few cases. The techniques apply to arbitrary q and d . In addition, the equivalence relations allow the set of all PPs for a given degree and a given field $GF(q)$ to be succinctly described by their representative nPPs. We give several tables at the end of the paper listing the representative nPPs (*i.e.*, the equivalence classes) for several values of q and d . We also give several new lower bounds for $M(n, D)$, the maximum number of permutations on n symbols with pairwise Hamming distance D , mostly derived from our results on PPs.

1 Introduction

Let $GF(q)$ denote the finite field over $q = p^m$ elements, where p is prime and $m \geq 1$. The prime p is called the *characteristic* of the field. A polynomial $P(x)$ over $GF(q)$ is a *permutation polynomial* (PP) if it permutes the elements of $GF(q)$. Let $N_d(q)$ be the number of PPs of degree d over $GF(q)$. Lidl and Mullen [19, 20] posed the problem of computing $N_d(q)$ and they found the following boundary conditions for $N_d(q)$:

$$\begin{aligned} N_1(q) &= q(q-1) \\ N_d(q) &= 0 \text{ if } d \nmid (q-1) \text{ and } d > 1 \\ \sum_{d=1}^{q-2} N_d(q) &= q! \end{aligned}$$

Dickson [10] characterized all PPs of degree up to 6. Hou [15] gave a survey of recent results about PPs. Chu, Colbourn, and Dukes [8], using a table of all PPs of degree at most five given by Lidl and Mullen [19], counted the number of different PPs of degree at most 5. Shallue and

*Department of Computer Science, University of Texas at Dallas, Box 830688, Richardson, TX 75083 USA. Research of the first author is supported in part by NSF award CCF-1718994.

Wanless [24] described those of degree 6. Li, Chandler, and Xiang [18] described PPs of degree 6 and 7 over a field of characteristic 2.

An exceptional PP is a PP $P(x)$ for $GF(q)$, which, for infinitely many m , is also a PP for $GF(q^m)$. Fan [11] obtained a classification of all permutation polynomials of degree 7 over $GF(q)$ for any odd prime power $q > 7$ up to linear transformations and proved that there are no non-exceptional PPs of degree 7 for finite fields of order q when $q > 49$. Fan also [13] obtained a classification of all permutation polynomials of degree 8 over $GF(q)$ for any odd prime power $q > 8$ up to linear transformations and proved that there are no PPs of degree 8 for finite fields of order q when $q > 31$. In addition, Fan [12] described all PPs of degree 8 over finite fields of characteristic 2 and proved that non-exceptional PP's of degree 8 do not exist over $GF(2^r)$ if $r > 6$.

It is an interesting problem to enumerate all permutation polynomials of somewhat large degree. Let $P(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0$ be a degree d permutation polynomial over $GF(q)$. A brute force search for degree d permutation polynomials over $GF(q)$ would require $O(dq^{d+2})$ time as there are $d + 1$ coefficients, with q choices for each one, and for each of these possibilities, one needs to examine the list of q values formed by the d terms of the polynomial on each element of $GF(q)$ to see if the result is a permutation. One way to make the search more efficient is to look for all *normalized* PPs (denoted by *nPPs*), which fixes certain coefficients. In normalization, $a_d = 1$, $a_0 = 0$, and when the field characteristic p does not divide d , we have $a_{d-1} = 0$ [19]. It is known that any PP can be transformed into an nPP by certain algebraic operations, which we will describe shortly. In this paper, we use the names *c-normalization* to denote the case where $p \nmid d$, *m-normalization* when $p > 2 \mid d$, and *b-normalization* when $p = 2 \mid d$. Properties of the latter two types of normalization will be investigated in the next section. In all three cases of normalization, searching for nPPs takes $O(dq^{d-1})$ time as three coefficients are fixed.

We define transformations that map PPs to PPs, and derive new equivalence relations based these functions, which yield a more succinct classification of PPs. Using these functions, we are able to fix a fourth coefficient, making the search for nPPs take $O(dq^{d-2})$ time. Our new equivalence relations are defined for arbitrary q and d . We provide results for PPs of somewhat large q and d . Specifically, we characterize and count PPs of degree at most 9 for finite fields for primes and prime powers up to $q = 97$, PPs of degree 10 for primes and prime powers up to $q = 73$, and PPs of degree 11 for primes and prime powers up to 32. This is a near complete accounting of PPs up to the stated limits for q and d .

Let n be a positive integer and let σ and π be permutations over n symbols. A *permutation array (PA)* is a set of permutations on n symbols. σ and π have *Hamming distance at least D* , denoted by $hd(\sigma, \pi) \geq D$, if $\sigma(x) \neq \pi(x)$ in at least D different positions x . A PA A has Hamming distance D , denoted by $hd(A) \geq D$, if every pair of permutations in A has Hamming distance at least D . The maximum number of permutations in a PA A on n symbols with $hd(A) \geq D$ is denoted by $M(n, D)$.

We are interested in an application of permutation polynomials to permutation arrays with large Hamming distance. Much recent work has focused on computing large permutation arrays with a given lower bound for their pairwise Hamming distance [1, 2, 3, 4, 5, 6, 8, 9, 14, 16, 17, 23, 25, 27, 22]. Chu *et al.* [8] showed that PPs can be used for lower bounds on $M(n, D)$. Let $P(x)$ and $Q(x)$ be two degree d PPs over $GF(q)$. Basically, $P(x)$ and $Q(x)$ can agree in at most d points, because for every set of d points there is one and only one polynomial of degree d that passes through them. So, the corresponding permutations must disagree in at least $q - d$ positions. That is, the permutations have Hamming distance at least $q - d$. So, it follows that, if $\sum_{k=1}^d N_k(q) \geq T$, then

$M(q, q-d) \geq T$, for some positive integer T . In addition, Chu, *et al.* [8] showed that when q is a prime power:

- $M(q, q-d) \geq \sum_{k=1}^d N_k(q)$.
- If $q = 2^k \not\equiv 1 \pmod{3}$, then $M(q, q-3) \geq (q+2)q(q-1)$ and $M(q, q-4) \geq \frac{1}{3}q(q-1)(q^2 + 3q + 8)$.
- If $q \not\equiv 2 \pmod{3}$, then $M(q, q-2) \geq q^2$.

R. Sobhani, *et al.* [26] computed some values of $N_d(q)$ and used these to give lower bounds for some values of $M(n, D)$. Berge, *et al.* [5] give a table with several new lower bounds for $M(n, D)$ for $n \leq 550$ and a table for $M(n, n-1)$ for prime powers $n \leq 600$.

The paper is organized as follows. In Section 2, we discuss normalization of permutation polynomials. In Sections 2.1 and 2.2, we extend the concept of normalization by defining m-normalization and b-normalization, to include cases when the degree of the polynomial is a multiple of the field characteristic p . These new forms of normalization apply when $p \mid d$, to which classic c-normalization does not apply. In Section 2.1, we show that for all d such that $p \mid d$ and $p \neq 2$, one can limit a search for m-normalized PPs, where $a_d = 1$, $a_0 = 0$, and either $a_{d-1} = 0$ or $a_{d-2} = 0$. So, again such a search takes $O(dq^{d-1})$ time, as three coefficients are fixed. In Section 2.2, we show that when $p \mid d$, $p = 2$ and d is in an interval $2^i \leq d \leq 2^{i+1}$, for some i , then one can limit a search for b-normalized PPs, where $a_d = 1$, $a_0 = 0$, and either $a_{e-2} = 0$ or $a_{e-1} = 0$, for a specified position e . Again, a such search takes $O(dq^{d-1})$ time.

In Section 3, we consider other transformations that map nPPs to PPs, allowing an order of magnitude optimization of the search, as mentioned earlier. We call these transformations the *F-map* and the *G-map*. In Section 4, we use the *F-map* and the *G-map* to define more inclusive equivalence relations on PPs with larger equivalence classes, which permit a more succinct description of all PPs of degree d for a field $GF(q)$. As the *F-map* allows us to fix another coefficient in a search for nPPs, the time complexity is reduced to $O(dq^{d-2})$. In Section 5, we describe algorithms that implement the search for nPPs using equivalence relations.

Finally, in Section 6, we present our numeric results. We provide a table listing specific values for $N_{11}(q)$. We present several tables that list the new categorizations of PPs (using equivalence relations), as well as the number of PPs in each category, for q up to 97 and degrees up to 11. We summarize all of these results in a table that lists the number of nPPs, equivalence classes, and total number of PPs for $q \leq 97$ and degree d , $6 \leq d \leq 10$. We also include a table giving new lower bounds for $M(n, D)$.

Notation. In this paper, we use p to denote a prime and $q = p^m$ to denote a power of the prime p for some $m \geq 1$. We use d ($2 \leq d < q$) to denote the degree of a PP $P(x) = a_dx^d + a_{d-1}x^{d-1} + \dots + a_1x + a_0$ over the finite field $GF(p^m) = GF(q)$ with field characteristic p . Throughout the paper and in the tables at the end we use the following notation for the elements of $GF(q)$. $GF(q)$ has $q-1$ non-zero elements, all of which can be listed as t^0, t^1, \dots, t^{q-2} , where $t \neq 0$ represents a generator of the multiplicative group of nonzero elements of $GF(q)$. We use the notation $t^0 = 1, t^1 = 2, \dots, t^{q-2} = q-1$. Lidl and Niederreiter [21] give this as the second choice of notations to describe the elements of a finite field, with the first choice for $GF(p^m)$ being by degree m polynomials with coefficients from $GF(p)$. PPs can easily be converted from one notation to the other. As a primitive polynomial is needed to do the appropriate arithmetic, we give explicit primitive polynomials for our results.

2 Normalized permutation polynomials and equivalence relations

As described in the introduction, a c-normalized nPP is a PP where $a_d = 1$, $a_{d-1} = 0$, and $a_0 = 0$. This form of normalization can be achieved when $p \nmid d$. We introduce two new definitions for normalization for the case $p \mid d$, which we call *m-normalization* and *b-normalization*. Table 1 summarizes the three types of normalization.

normalization type	degree restriction	nPP properties
<i>c-normalization</i>	$p \nmid d$	monic, $P(0) = 0$, $a_{d-1} = 0$
<i>m-normalization</i>	$p \mid d$ and $p > 2$	monic, $P(0) = 0$, either $a_{d-1} = 0$ or $a_{d-2} = 0$
<i>b-normalization</i>	$p \mid d$ and $p = 2$	monic, $P(0) = 0$, if $2^i \leq d \leq 2^{i+1} - 3$ for some i then either $a_r = 0$ or $a_{r-1} = 0$, where $r = 2^i - 1$

Table 1: Types of normalization for PPs, $P(x) = a_dx^d + a_{d-1}x^{d-1} + \cdots + a_1x + a_0$, of degree d with field characteristic p .

Here are some examples of normalized PPs:

- The degree-9 PP $x^9 + 2x^7 + 3x^5$ over $GF(5^2)$ is c-normalized, as $a_9 = 1$, $a_8 = 0$ and $a_0 = 0$.
- The degree-6 PP is $x^6 + x^5 + x^3 + 5x^2 + 5x$ over $GF(3^2)$ is m-normalized, as $a_6 = 1$, $a_4 = 0$ and $a_0 = 0$.
- The degree-10 PP $x^{10} + x^9 + x^7 + 26x^5 + 30x^4 + 21x^2 + 31x$ over $GF(2^5)$ is b-normalized, as $a_{10} = 1$, $a_6 = 0$ and $a_0 = 0$ (with $i = 3$ and $r = 2^i - 1 = 7$).

The normalization definitions of nPPs allow nearly all PPs of degree d to be converted to nPPs of degree d by applying certain algebraic operations, called *normalization operations*, in some order. The normalization operations [19, 24] on a PP $P(x)$ are

- multiplication by a constant, *i.e.*, $aP(x)$, for some nonzero constant a in $GF(q)$,
- addition to the variable, *i.e.*, $P(x + b)$, for some constant b in $GF(q)$.
- addition of a constant, *i.e.*, $P(x) + c$, for some constant c in $GF(q)$, and

We include an extended normalization operation, namely

- multiplication of the variable, *i.e.*, $P(sx)$, for some nonzero constant s .

Extended normalization will come into play in Section 4. It is customary in the literature to refer to the normalization operations applied to a PP $P(x)$ by $aP(x + b) + c$, where a , b , and c are elements of the finite field, with a being nonzero. References to extended normalization applied to a PP $P(x)$, denoted by $aP(sx + b) + c$, where a, b, c , and s are elements of the finite field, with both a and s are nonzero, appear, for example, in [11, 12, 13] and have been called linear transformations.

Notice that each operation has an inverse. That is, if e is the multiplicative inverse of a in $GF(q)$, then $e(aP(x)) = P(x)$. If h is the additive inverse of b in $GF(q)$, then $P((x + b) + h) = P(x)$, etc. Let $aP(x + b) + c$ denote the result of performing normalization operations in any order. Note that the second operation, addition of a constant b to the variable x , when $P(x)$ is a PP, has the effect of permuting the order of its values. Note also that, if one adds c and then multiplies by a , the

result is $aP(x) + ac$. Whereas, if we multiply by a and then add c , the result is $aP(x) + c$. Since we consider adding all such constants, and a times c is a constant in $GF(q)$, the set of such PPs is the same regardless of the order of operations.

We define two equivalence relations, \mathcal{R} and \mathcal{R}_E , on PPs of degree d , for some $d > 1$, as follows.

Definition 1. Let $P(x)$ and $Q(x)$ be PPs of degree d , for some d . If $P(x)$ can be converted into $Q(x)$ by some sequence of normalization operations, then $P(x)$ and $Q(x)$ are related by \mathcal{R} . If $P(x)$ can be converted into $Q(x)$ by some sequence of normalization operations including the extended normalization operation, then $P(x)$ and $Q(x)$ are related by \mathcal{R}_E .

It is easy to see that \mathcal{R} and \mathcal{R}_E are equivalence relations. Observe that any PP $P(x)$ can be transformed to itself by the empty sequence of normalization operations. If $P(x)$ can be transformed into $Q(x)$ by some sequence, then $Q(x)$ can be transformed into $P(x)$ by the inverse of each step in the reversed sequence. Finally, if $P(x)$ can be transformed into $Q(x)$, and $Q(x)$ can be transformed into $T(x)$, then $P(x)$ can be transformed into $T(x)$ by the composition of the two sequences.

An nPP is a representative of an equivalence class defined by \mathcal{R} . We can make a more efficient search algorithm by searching for nPPs and the equivalence class they represent, rather than searching directly for PPs. In order to use equivalence classes to count PPs, we need to explore specific properties of normalization.

2.1 m-normalization

If the degree d of a PP $P(x)$ is a multiple of the field characteristic p , then there is no constant b which when added to the variable x will make the coefficient a_{d-1} of $P(x+b)$ equal to 0. This is due to the fact that in this case, the expansion of $(x+b)^d$, does not have a nonzero term involving x^{d-1} . So, when $p \mid d$, we cannot necessarily achieve c-normalization. However, if $a_{d-1} \neq 0$, then there is a constant b such that $P(x+b)$ has $a_{d-2} = 0$. That is, we can achieve m-normalization. This is similar to Fan's idea in Observation 8 [11]. Recall from Table 1 that m-normalization applies when $p \neq 2$ and d is a multiple of p . In this section, we show that m-normalization can be achieved for any PP satisfying these conditions.

As stated in Section 2, $P(x)$ is *m-normalized* when $a_d = 1$, $a_0 = 0$, and either $a_{d-1} = 0$ or $a_{d-2} = 0$. We show that any PP of degree d over $GF(p^m)$ where $p \mid d$ and $p \neq 2$ can be m-normalized. By the definition of characteristic of a finite field, for all $k \in GF(q)$, $\underbrace{k + k + \dots + k}_{d \text{ times}} = 0$.

Furthermore, for all $c \in GF(q)$, there exists a $k \in GF(q)$ such that $\underbrace{k + k + \dots + k}_{d \text{ times}} = c$. To see this let $\underbrace{1 + 1 + \dots + 1}_{d-1 \text{ times}} = s$. Note that $s \neq 0$, since $\underbrace{1 + 1 + \dots + 1}_{d \text{ times}} = 0$. Since $\underbrace{k + k + \dots + k}_{d-1 \text{ times}} = ks$, one wants $k(s) = c$ or, equivalently, $k = c/s$ where the arithmetic is in $GF(q)$.

Theorem 2. Any PP $P(x)$ where the degree d is a multiple of the field characteristic p , can be transformed to an m-normalized PP $Q(x)$ by the normalization operations.

Proof. Let $P(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_1 x + a_0$, and for some $a, b, c \in GF(q)$ with $a \neq 0$, let

$$\begin{aligned} Q(x) &= aP(x+b) + c \\ &= aa_d(x+b)^d + aa_{d-1}(x+b)^{d-1} + aa_{d-2}(x+b)^{d-2} + \dots + aa_1(x+b) + aa_0 + c \\ &= b_d x^d + b_{d-1} x^{d-1} + b_{d-2} x^{d-2} + \dots + b_1 x + b_0, \end{aligned}$$

Observe that the degree d term of $Q(x)$ has the coefficient $b_d = aa_d$. If we choose a to be the multiplicative inverse of a_d , then the degree d coefficient of $Q(x)$ will be 1.

If $a_{d-1} = 0$, then $b_{d-1} = 0$ and the desired property is true. So suppose that $a_{d-1} \neq 0$, and consider b_{d-2} in $Q(x)$. Since d is a multiple of p , the expansion of $a_d(x+b)^d$ will derive nonzero coefficients only for terms whose degrees are multiples of p . Since $p > 2$, this means that $(d-2) \nmid p$, so $a_d(x+b)^d$ will have a coefficient of 0 for the degree $d-2$ term. Hence b_{d-2} is calculated solely by the expansion of $aa_{d-1}(x+b)^{d-1} + aa_{d-2}(x+b)^{d-2}$.

The expansion of $aa_{d-1}(x+b)^{d-1}$ will produce a term of degree $d-2$ with coefficient $aa_{d-1}b'$ where $b' = \sum_{i=1}^{d-1} b$. The expansion of $aa_{d-2}(x+b)^{d-2}$ will produce a term of degree $d-2$ with coefficient $b_{d-2} = aa_{d-2}$. Therefore the coefficient of x^{d-2} in $Q(x)$ is $b_{d-2} = aa_{d-1}b' + aa_{d-2} = a(a_{d-1}b' + a_{d-2})$. And by algebra, b_{d-2} is zero if $a_{d-1}b' + a_{d-2} = 0$. Since $a_{d-1} \neq 0$ and $d-1$ is not a multiple of p , we can choose b such that b' is the additive inverse of a_{d-2}/a_{d-1} , making $b_{d-2} = 0$ in $Q(x)$. So in $Q(x)$, $b_d = 1, b_0 = 0$, and either $b_{d-1} = 0$ or $b_{d-2} = 0$.

If we choose c to be the additive inverse of the constant term of $aP(x+b)$, then the constant term becomes zero. So, we achieve m -normalization. □

2.2 b-normalization

In Section 2.1, we considered PPs whose degree d is a multiple of the field characteristic p and $p > 2$. We showed that under those conditions, m -normalization can be achieved. In this section, we consider the remaining case, namely, $p \mid d$ and $p = 2$, and show that, b -normalization can be achieved except when $d = 2^i - 2$, for some $i \geq 2$.

We say that the integer interval $[r, s]$ has a $[t, u]$ gap, if for all $d \in [r, s]$, the expansion of $(x+b)^d$, does not include any nonzero x^e monomials, where $e \in [t, u]$. For example, the integer interval $[8, 13]$ has a $[6, 7]$ gap as seen by:

$$\begin{aligned} (x+b)^8 &= x^8 + b^8 \\ (x+b)^9 &= x^9 + bx^8 + b^8x + b^9 \\ (x+b)^{10} &= x^{10} + b^2x^8 + b^8x^2 + b^{10} \\ (x+b)^{11} &= x^{11} + bx^{10} + b^2x^9 + b^3x^8 + b^8x^3 + b^9x^2 + b^{10}x + b^{11} \\ (x+b)^{12} &= x^{12} + b^4x^8 + b^8x^4 + b^{12} \\ (x+b)^{13} &= x^{13} + bx^{12} + b^4x^9 + b^5x^8 + b^8x^5 + b^9x^4 + b^{12}x + b^{13}. \end{aligned}$$

That is, in each of the exhibited expansions there are no x^6 or x^7 terms.

We use this observation in the proofs below regarding b -normalization. In Lemma 4, we show that in the expansion of $(x+b)^d$ when $d \in [2^i, 2^{i+1} - 3]$, the coefficients of the terms x^{2^i-2} and x^{2^i-1} term are always zero. That is, the interval $[2^i, 2^{i+1} - 3]$ has a $[2^i - 2, 2^i - 1]$ gap. In Lemma 5 we show that any PP $P(x)$ of characteristic 2 such that d is a multiple of 2, has a related PP $P(x+b)$ for which certain terms in its expansion always have a coefficient equal to zero. We use Lucas' Theorem (stated below for the reader's convenience) in the proof of Lemma 4.

Theorem 3. [Lucas' Theorem[7]] Let p be prime, and let $m = m_0 + m_1p + \dots + m_dp^d$ and $n = n_0 + n_1p + \dots + n_dp^d$, where $0 \leq m_i, n_i < p$ for $i = 0, 1, \dots, d$. Then

$$\binom{m}{n} \equiv \prod_{i=0}^d \binom{m_i}{n_i} \pmod{p}$$

Lemma 4. [Gap Lemma] For all $i > 1$, the expansion of $(x + b)^d$, for $d \in [2^i, 2^{i+1} - 3]$, has a $[2^i - 2, 2^i - 1]$ gap.

Proof. Consider the expansion $(x + b)^d = \sum_{k=0}^d \binom{d}{k} x^{d-k} b^k$. Let $k \in \{d - (2^i - 2), d - (2^i - 1)\}$. Clearly, b^k is not zero, so our job is to show that the expression $\binom{d}{k}$ is zero. Let $k' = d - k$, that is $k' \in \{2^i - 2, 2^i - 1\}$. By a well-known identity, we have $\binom{d}{k} = \binom{d}{d-k} = \binom{d}{k'}$. Represent d and k' by their base-2 $(i + 1)$ -tuples $\delta = (\delta_i, \delta_{i-1}, \dots, \delta_2, \delta_1, \delta_0)$ and $\kappa = (\kappa_i, \kappa_{i-1}, \dots, \kappa_2, \kappa_1, \kappa_0)$, respectively, where for all i , $\delta_i, \kappa_i \in \{0, 1\}$. Observe that at least one δ_j ($0 \leq j \leq i - 2$) must be 0 because $2^i \leq d \leq 2^{i+1} - 3$. Observe also that $\kappa_i = 1$ for all $i > 0$. Hence there is a j such that $\delta_j = 0$ and $\kappa_j = 1$, so by Lucas' Theorem, $\binom{d}{k'} = 0 = \binom{d}{k}$. It follows that $(x + b)^d$, for $d \in [2^i, 2^{i+1} - 3]$, has a $[2^i - 2, 2^i - 1]$ gap. \square

Lemma 5. Let $i > 1$. Let $d \in [2^i, 2^{i+1} - 3]$ be even. For any PP $P(x)$ over $GF(2^m)$, where $m > 2$, there is a constant b in $GF(2^m)$ such that in the PP $P(x + b)$, either the x^{2^i-1} term or the x^{2^i-2} term is zero.

Proof. By Lemma 4, the interval $[2^i, 2^{i+1} - 3]$ has a $[2^i - 2, 2^i - 1]$ gap. Let $P(x) = a_d x^d + a_{d-1} x^{d-1} + a_{d-2} x^{d-2} + \dots + a_1 x + a_0$, where $d \in [2^i, 2^{i+1} - 3]$ is even. Adding b to the argument gives: $P(x + b) = a_d (x + b)^d + a_{d-1} (x + b)^{d-1} + a_{d-2} (x + b)^{d-2} + \dots + a_1 (x + b) + a_0$.

If a_{2^i-1} is zero there is nothing to prove, so suppose a_{2^i-1} is not zero. Since $[2^i, 2^{i+1} - 3]$ has a $[2^i - 2, 2^i - 1]$ gap, each term $(x + b)^s$, for $s \in [2^i, 2^{i+1} - 3]$ has no x^e term for $e \in [2^i - 2, 2^i - 1]$. This means that $a_{2^i-1} (x + b)^{2^i-1}$ and $a_{2^i-2} (x + b)^{2^i-2}$ are the only possible terms whose expansion has a nonzero x^{2^i-2} term. By the binomial theorem, $a_{2^i-1} (x + b)^{2^i-1} = a_{2^i-1} x^{2^i-1} + a_{2^i-1} b x^{2^i-2} + \dots$, and $a_{2^i-2} (x + b)^{2^i-2} = a_{2^i-2} x^{2^i-2} + \dots$, where low order terms are not shown. Summing these two expansions and isolating the x^{2^i-2} term, we solve for the value of b such that $a_{2^i-1} b x^{2^i-2} + a_{2^i-2} x^{2^i-2} = 0$. We see that when $b = -a_{2^i-1}/a_{2^i-2}$, the coefficient of the x^{2^i-2} term of $P(x + b)$ is zero. \square

For example, let $d = 2^3$, and let $P(x) = a_8 x^8 + a_7 x^7 + a_6 x^6 + \dots + a_1 x + a_0$. Adding b to the argument gives:

$$\begin{aligned} P(x + b) &= a_8 (x + b)^8 + a_7 (x + b)^7 + a_6 (x + b)^6 + \dots + a_1 x + a_0 \\ &= a_8 (x^8 + b^8) + a_7 (x^7 + b x^6 + b^2 x^5 + \dots) + a_6 (x^6 + b^2 x^4 + \dots) + \dots \\ &= a_8 x^8 + a_8 b^8 + (a_7 x^7 + a_7 b x^6 + \dots) + (a_6 x^6 + a_6 b^2 x^4 + \dots) + \dots \end{aligned}$$

We want to solve for the value of b that makes the coefficient of the x^6 term of $P(x + b)$ zero. So $a_7 b x^6 + a_6 x^6 = 0$ is satisfied by $b = -a_7/a_6$.

Theorem 6. Any PP $P(x)$ over $GF(2^m)$ for some $m > 2$, and $2 \mid d$ can be transformed to an b -normalized PP $Q(x)$ by the normalization operations, except when $d = 2^i - 2$, for some $i \geq 2$.

Proof. Let $P(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_1 x + a_0$, and for some $a, b, c \in GF(q)$ with $a \neq 0$, let

$$\begin{aligned} Q(x) &= aP(x + b) + c \\ &= aa_d (x + b)^d + aa_{d-1} (x + b)^{d-1} + aa_{d-2} (x + b)^{d-2} + \dots + aa_1 (x + b) + aa_0 + c \\ &= b_d x^d + b_{d-1} x^{d-1} + b_{d-2} x^{d-2} + \dots + b_1 x + b_0, \end{aligned}$$

Observe that the degree d term of $Q(x)$ has the coefficient $b_d = aa_d$. If we choose a to be the multiplicative inverse of a_d , then the degree d coefficient of $Q(x)$ will be 1. If we choose c to be the additive inverse of the constant term of aa_0 , then the constant term of $Q(x)$ will be $b_0 = aa_0 + c = 0$. By Lemma 5, there is a b such that in $Q(x)$, the coefficient of either the degree $2^i - 1$ term or degree $2^i - 2$ term equal to 0, except when $d = 2^i - 2$, for some $i \geq 2$. Hence $Q(x)$ is b-normalized. \square

2.3 Counting PPs using equivalence classes based on normalization

When the degree d of a PP is not a multiple of the field characteristic p , the PP can be c-normalized. As shown below, for any such PP $P(x)$, there is a unique triple (a, b, c) such that $aP(x+b) + c$ is c-normalized. Moreover, each equivalence class contains exactly one nPP, and each PP belongs to exactly one equivalence class. These properties allow us to count, for a given q and d , the number of PPs in each equivalence class.

Observation 7. *Let $P(x)$ be PP where $p \nmid d$. Then there is a unique triple (a, b, c) such that $aP(x+b) + c$ is c-normalized.*

Proof. Let $Q(x) = aP(x+b) + c$. Since $Q(x)$ is c-normalized, the coefficient of its degree- d term is $aa_d = 1$. Hence $a = a_d^{-1}$. The degree- $(d-1)$ term of $Q(x)$ is $(aa_{d-1} + db)x^{d-1} = 0$, so $b = -aa_{d-1}/d$. (Note that $d \neq 0$). Finally, since the constant term of $Q(x)$, namely $Q(0)$, is 0, c is uniquely determined by the constant term of $aP(x+b)$. \square

Let $Q(x)$ be an nPP of degree d . The equivalence class under the relation \mathcal{R} containing $Q(x)$, denoted by $[Q]$, is the set

$$[Q] = \{aQ(x+b) + c \mid a, b, c \in GF(q) \text{ and } a \neq 0\}.$$

Lemma 8. *Let $Q(x)$ be an nPP of degree $d < q$ where $p \nmid d$. Then all $q^2(q-1)$ polynomials in $[Q]$ are different.*

Proof. Let $Q(x) = a_dx^d + a_{d-1}x^{d-1} + \dots + a_1x + a_0$ and assume that polynomials $P(x) = aQ(x+b) + c$ and $P'(x) = a'Q(x+b') + c'$ in $[Q]$ are equal. Since the degree- d terms of $P(x)$ and $P'(x)$ are equal, $a = a'$. Since the degree- $(d-1)$ terms are equal, $a(a_{d-1} + a_d db) = a'(a_{d-1} + a_d db')$. Then $a_{d-1} + a_d db = a_{d-1} + a_d db'$ and $db = db'$ and $b = b'$ (since d is not a multiple of p). Let e be the lowest degree term of $aQ(x+b)$. Then the lowest degree terms of $P(x)$ and $P'(x)$ are $e + c$ and $e + c'$, respectively. Thus, $c = c'$ and the claim follows. \square

Note that Lemma 8 implies that each equivalence class of \mathcal{R} contains one and only one nPP. Thus when d is not a multiple of p , each equivalence class contains exactly $q^2(q-1)$ members (including the representative nPP). Note that the equivalence classes by definition are disjoint. If the number of nPPs is k , there are $kq^2(q-1)$ PPs.

3 Mapping nPPs to nPPs

We now describe the F -map and the G -map, two new functions that map nPPs to nPPs. We will use these functions in Section 4 to define new equivalence relations on nPPs whose equivalence classes are unions of equivalence classes of \mathcal{R} .

3.1 The F -map

The F -map is the function that multiplies the degree $(d - k)$ term of $P(x)$ by t^k , for all k . The F -map allows one additional coefficient to be fixed, resulting in an order of magnitude speedup in the search for PPs.

Definition 9. Define the F -map by

$$F(P(x)) = t^0 a_d x^d + t^1 a_{d-1} x^{d-1} + \cdots + t^k a_{d-k} x^{d-k} + \cdots + t^{d-1} a_1 x + t^d a_0 = \sum_{k=0}^d t^k a_{d-k} x^{d-k}.$$

First, we show that the set of PPs is closed under the F -map.

Lemma 10. If $P(x)$ is a PP, then $F(P(x))$ is a PP.

Proof. We show that $F(P(x))$ is one-to-one. Observe that, for any nonzero t^i ,

$$\begin{aligned} F(P(t^i)) &= t^0 a_d (t^i)^d + t^1 a_{d-1} (t^i)^{d-1} + \cdots + t^{d-1} a_1 (t^i)^1 + t^d a_0 \\ &= t^d (a_d (t^i)^{d-i} + a_{d-1} (t^i)^{d-i-1} + \cdots + a_1 (t^i)^{d-i-1} + a_0) \\ &= t^d (a_d (t^{i-1})^d + a_{d-1} (t^{i-1})^{d-1} + \cdots + a_1 (t^{i-1})^1 + a_0) \\ &= t^d P(t^{i-1}) \\ &= t^d P(t^i/t). \end{aligned}$$

It follows that $F(P(x))$ is a permutation polynomial, since it is obtained from $P(x)$ by multiplying by the constant t^d and replacing the argument x by $t^{-1}x$. \square

Lemma 10 provides an alternative formulation for the F -map, namely, $F(P(x)) = t^d P(t^{-1}x) = t^d P(x/t)$. Let $F^i(P(x))$ denote the composition of the F -map with itself i times, for some i . Then

$$F^i(P(x)) = t^{di} P(t^{-i}x) = t^{di} P(x/t^i). \quad (1)$$

We will use this formulation in later proofs in this paper.

By definition, the F -map multiplies the coefficient a_d by $t^0 = 1$, so when $P(x)$ is an nPP, $F(P(x))$ is also an nPP. This is formalized in Corollary 11 below, which follows easily from Lemma 10.

Corollary 11. If $P(x)$ is an nPP over $GF(q)$, then $F(P(x))$ is an nPP.

Observe that the $q - 1$ non-zero elements of $GF(q)$ form a cyclic group, \mathcal{G}_{q-1} , under multiplication [21]. Moreover, for each k , there exists an r , ($0 < r \leq q - 1$), such that the iterates, $t^k, t^{2k}, \dots, (t^{rk \bmod (q-1)} = 1)$, form a cyclic subgroup, \mathcal{H}_{t^k} , of \mathcal{G}_{q-1} . By Lagrange's theorem, the number of elements in \mathcal{H}_{t^k} , i.e., $\text{ord}(\mathcal{H}_{t^k})$, is a divisor of $q - 1$.

Consider iterations of the F -map, namely, the sequence

$$P(x), F(P(x)), F^2(P(x)), \dots, F^i(P(x)), \dots, \quad (2)$$

where $F^i(P(x)) = \sum_{k=0}^d t^{ik} a_{d-k} x^{d-k}$. For the degree $(d - k)$ term, iterative use of the F -map yields the sequence of coefficients

$$a_{d-k}, t^k a_{d-k}, t^{2k} a_{d-k}, \dots, (t^{rk \bmod (q-1)} a_{d-k} = a_{d-k})$$

where the terms are simply the elements of \mathcal{H}_{t^k} multiplied by the common factor a_{d-k} . This forms a cycle of length $r \leq q - 1$ where r is smallest integer such that $t^{rk \bmod (q-1)} a_{d-k} = a_{d-k}$.

Definition 12. Let k be an integer such that $1 \leq k \leq d$. Define the **F_k -map** by $F_k(x) = xt^k$.

The F_k function gives us a third way to look at the F -map, namely, for each k , the F -map computes $F_k(a_{d-k}) = t^k a_{d-k}$. In other words, $F(P(x)) = a_d x^d + t^1 a_{d-1} x^{d-1} + \cdots + t^{d-1} a_1 x + t^d a_0 = a_d x^d + F_1(a_{d-1})x^{d-1} + \cdots + F_{d-1}(a_1)x + F_d(a_0)$. Note that iterations of the F_k -map on the element 1 yields the cyclic subgroup \mathcal{H}_{t^k} . We call this sequence of iterations the **F_k -cycle**. Define the length of the F_k -cycle to be $\text{ord}(\mathcal{H}_{t^k})$. In this paper, we are interested in those values of k for which the $(d-k)^{\text{th}}$ coefficient of an nPP $P(x)$ is not 0.

Observe that for any PP $P(x)$, there is an integer $s \geq 1$, such that the sequence shown in Equation (2) forms a cycle.

Definition 13. The sequence of iterates of the F -map on the PP $P(x)$, namely

$$P(x), F(P(x)), F^2(P(x)), \dots, F^s(P(x)) = P(x)$$

is called the **F -cycle on $P(x)$** .

Consider the values of k for which the coefficient a_{d-k} in $P(x)$ is nonzero. For all such k , ($1 \leq k \leq d$), let $g_k = \gcd(k, q-1)$, and let $j = \min_k \{g_k\}$. The length of the F -cycle on $P(x)$ is $s = (q-1)/j$. That is, the length of the F -cycle on $P(x)$ is the least common multiple of the orders of the subgroups \mathcal{H}_{t^k} for all k such that $a_{d-k} \neq 0$.

For example, consider $GF(5^2)$. The cyclic subgroups of \mathcal{G}_{5^2-1} are $\mathcal{H}_{t^0}, \mathcal{H}_{t^1}, \mathcal{H}_{t^2}, \mathcal{H}_{t^3}, \mathcal{H}_{t^4}, \mathcal{H}_{t^6}$,

\mathcal{H}_{t^8} , $\mathcal{H}_{t^{12}}$, and their orders are 1, 24, 12, 8, 6, 4, 3 and 2, respectively. To see this, observe that

$$\begin{aligned}
\mathcal{H}_1 = \mathcal{H}_{t^0} &= \{t^{i*0}\} = \{1\} \text{ for all } i & \mathcal{H}_7 = \mathcal{H}_{t^6} &= \{t^{i*6}\} \text{ for all } i \\
& & &= \{t^{0*6}, t^{1*6}, t^{2*6}, t^{3*6}, t^{4*6}\} \\
\mathcal{H}_2 = \mathcal{H}_{t^1} &= \{t^{i*1}\} \text{ for all } i & &= \{t^0, t^6, t^{12}, t^{18}, t^{24} = t^0\} \\
&= \{t^{0*1}, t^{1*1}, t^{2*1}, t^{3*1}, \dots\} & &= \{1, 7, 13, 19\} \\
&= \{t^0, t^1, t^2, t^3, \dots, t^{23}, t^{24} = t^0\} & &= \mathcal{H}_{19} \\
&= \{1, 2, 3, 4, \dots, 24\} & & \\
&= \mathcal{H}_6 = \mathcal{H}_8 = \mathcal{H}_{12} = \mathcal{H}_{14} = \mathcal{H}_{18} = \mathcal{H}_{20} = \mathcal{H}_{24} & \mathcal{H}_9 = \mathcal{H}_{t^8} &= \{t^{i*8}\} \text{ for all } i \\
\mathcal{H}_3 = \mathcal{H}_{t^2} &= \{t^{i*2}\} \text{ for all } i & &= \{t^{0*8}, t^{1*8}, t^{2*8}, t^{3*8}\} \\
&= \{t^{0*2}, t^{1*2}, t^{2*2}, t^{3*2}, \dots\} & &= \{t^0, t^8, t^{16}, t^{24} = t^0\} \\
&= \{t^0, t^2, t^4, t^6, \dots, t^{22}, t^{24} = t^0\} & &= \{1, 9, 17\} \\
&= \{1, 3, 5, 7, \dots, 23\} & &= \mathcal{H}_{17} \\
&= \mathcal{H}_{11} = \mathcal{H}_{15} = \mathcal{H}_{23} & \mathcal{H}_{13} = \mathcal{H}_{t^{12}} &= \{t^{i*12}\} \text{ for all } i \\
\mathcal{H}_4 = \mathcal{H}_{t^3} &= \{t^{i*3}\} \text{ for all } i & &= \{t^{0*12}, t^{1*12}, t^{2*12}\} \\
&= \{t^{0*3}, t^{1*3}, t^{2*3}, t^{3*3}, \dots\} & &= \{t^0, t^{12}, t^{24} = t^0\} \\
&= \{t^0, t^3, t^6, t^9, \dots, t^{21}, t^{24} = t^0\} & &= \{1, 13\} \\
&= \{1, 4, 7, 10, \dots, 22\} & & \\
&= \mathcal{H}_{10} = \mathcal{H}_{16} = \mathcal{H}_{22} & & \\
\mathcal{H}_5 = \mathcal{H}_{t^4} &= \{t^{i*4}\} \text{ for all } i & & \\
&= \{t^{0*4}, t^{1*4}, t^{2*4}, t^{3*4}, t^{4*4}, t^{5*4}, t^{6*4}\} & & \\
&= \{t^0, t^4, t^8, t^{12}, t^{16}, t^{20}, t^{24} = t^0\} & & \\
&= \{1, 5, 9, 13, 17, 21\} & & \\
&= \mathcal{H}_{21} & &
\end{aligned}$$

where *mod* 24 arithmetic is used in the exponents.

To illustrate the computation of F -cycles, consider the nPP $P(x) = x^9 + 2x^7 + 12x^5 + 4x^3 + 17x$ over $GF(25)$. For all k , ($1 \leq k \leq d$), the non-zero coefficients are a_7, a_5, a_3 and a_1 , which correspond to $k = 2, 4, 6$, and 8 and $g_k = \gcd(k, 24) = 2, 4, 6$, and 8, respectively. We next compute $j = \min_k \{g_k\} = \min\{2, 4, 6, 8\} = 2$. So, the length of the F -cycle on $P(x)$ is $(q-1)/j = 24/2 = 12$.

That $F^{12}(P(x)) = P(x)$ is easily verified. Note also that the length of each respective F_k -cycle is the order of the subgroup \mathcal{H}_{t^k} , which, referring to the list above, is $\text{ord}(\mathcal{H}_{t^2}) = 12$, $\text{ord}(\mathcal{H}_{t^4}) = 6$, $\text{ord}(\mathcal{H}_{t^6}) = 4$, and $\text{ord}(\mathcal{H}_{t^8}) = 3$, and their least common multiple is 12, which is the length of the F -cycle on $P(x)$.

In general, for larger degree nPPs over $GF(q)$, there is a k such that $\gcd(k, q-1) = 1$, so the length of the F -cycle is $q-1$. This means that if there is a nPP $a_d x^d + a_{d-1} x^{d-1} + \dots + a_1 x + a_0$ such that $a_{d-k} \neq 0$, then there is also one in which $a_{d-k} = 1$. This allows us to fix an additional coefficient a_{d-k} to the values in $\{0, 1\}$, thus reducing the search time for nPPs to $O(dq^{d-2})$.

3.2 The G -map

The G -map is the function that raises each coefficient in $P(x)$ to the power p . That is,

Definition 14. Define the G -map by

$$G(P(x)) = a_d^p x^d + a_{d-1}^p x^{d-1} + \cdots + a_1^p x + a_0^p = \sum_{k=0}^d a_{d-k}^p x^{d-k}.$$

We also consider the function G as a function on the elements of $GF(q)$, which follows from the case the polynomial is of degree 0.

We show that the set of PPs is closed under the G -map.

Lemma 15. If $P(x)$ is a PP, then $G(P(x))$ is a PP.

Proof. Observe that for all c, d in $GF(q)$, $(c + d)^p = c^p + d^p$. This is the so-called “Freshman Dream” idea, which follows from the binomial theorem and the fact that $\binom{p}{i} \equiv 0 \pmod{p}$ when p is prime [21]. So,

$$\begin{aligned} (P(x))^p &= (a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0)^p \\ &= a_d^p (x^p)^d + a_{d-1}^p (x^p)^{d-1} + \cdots + a_1^p x^p + a_0^p \\ &= G(P(x^p)). \end{aligned}$$

Also observe that if $(z_0, z_1, \dots, z_{q-1})$ is a permutation of $GF(q)$, then so is $(z_0^p, z_1^p, \dots, z_{q-1}^p)$. Suppose not. That is, suppose that $z_i^p = z_j^p$. Then $(z_i^p - z_j^p) = 0$, so, $(z_i - z_j)^p = 0$. Hence, $z_i - z_j = 0$, and so, $z_i = z_j$, a contradiction.

By assumption, $P(x)$ is a PP, so $\theta = (P(0), P(1), \dots, P(q-1))$ is a permutation. It follows that $\pi = (P(0)^p, P(1)^p, \dots, P(q-1)^p)$ is also a permutation. Since π is the permutation generated by the polynomial $(P(x))^p$, it follows that $(P(x))^p$ is a PP. Hence, $G(P(x^p))$ is a PP, since, as shown above, $(P(x))^p = G(P(x^p))$. This means the permutation generated by $G(P(x^p))$, namely, $(G(P(0^p)), G(P(1^p)), \dots, G(P((q-1)^p)))$ is identical to the permutation π . That is, $\pi = (P(0)^p, P(1)^p, \dots, P(q-1)^p) = (G(P(0^p)), G(P(1^p)), \dots, G(P((q-1)^p)))$.

Note also that $\sigma = (0^p, 1^p, \dots, (q-1)^p)$ is a permutation. Applying $P(x)$ to σ yields the permutation $\rho = (P(0^p), P(1^p), \dots, P((q-1)^p))$. Since ρ is a permutation, it is simply a reordering of the permutation $\theta = (P(0), P(1), \dots, P(q-1))$. Hence the sequence $\tau = (G(P(0)), G(P(1)), \dots, G(P(q-1)))$ is simply a reordering of the permutation $\pi = (G(P(0^p)), G(P(1^p)), \dots, G(P((q-1)^p)))$. That is, τ is a permutation. Finally, observe that τ is the permutation generated by applying the G -map to the PP $P(x)$. So it follows that $G(P(x)) = (a_d)^p x^d + (a_{d-1})^p x^{d-1} + \cdots + (a_1)^p x + (a_0)^p$ is a PP over $GF(q)$. \square

We now show that the set of nPPs is closed under the G -map.

Corollary 16. If $P(x)$ is a nPP then $G(P(x))$ is also an nPP.

Proof. By Lemma 15, we only need to show that $G(P(x))$ is normalized when $P(x)$ is normalized. Note that the coefficient of x^d in $P(x)$ is $a_d = 1$. Hence, by definition of the G -map, the coefficient of x^d in $G(P(x))$ is $a_d^p = 1^p = 1$. Also, since the coefficient of x^0 in $P(x)$ is $a_0 = 0$, then the coefficient of x^0 in $G(P(x))$ is $a_0^p = 0^p = 0$. For c-normalization, since the coefficient of x^{d-1} in $P(x)$ is $a_{d-1} = 0$, then the coefficient of x^{d-1} in $G(P(x))$ is $a_{d-1}^p = 0^p = 0$. For m-normalization, if the coefficient of x^{d-2} in $P(x)$ is $a_{d-2} = 0$, then the coefficient of x^{d-2} in $G(P(x))$ is $a_{d-2}^p = 0^p = 0$. For b-normalization, if for some specified j , $a_{d-j} = 0$, then the corresponding coefficient $a_{d-j}^p = 0^p = 0$ as well. \square

By definition, the G -map raises each coefficient in the PP $P(x)$ to the power p . Consider iterations of the G -map, namely, the sequence

$$P(x), G(P(x)), G^2(P(x)), \dots, G^i(P(x)), \dots, \quad (3)$$

where $G^i(P(x)) = \sum_{k=0}^d a_{d-k}^{p^i} x^{d-k}$. Iterative use of the G -map on the $(d-k)^{th}$ coefficient yields the sequence

$$a_{d-k} = a_{d-k}^{p^0}, a_{d-k}^{p^1}, a_{d-k}^{p^2}, \dots, a_{d-k}^{p^m \pmod{p^m-1}} = a_{d-k}$$

where $a_{d-k}^{p^m \pmod{p^m-1}} = a_{d-k}$, because $p^m = 1 \pmod{p^m-1}$. This forms a cycle of length m . Let $r_{a_{d-k}} \pmod{p^m-1}$ be the smallest integer such that $a_{d-k}^{p^{r_{a_{d-k}}}} = a_{d-k}$. The sequence of coefficients $a_{d-k}, a_{d-k}^{p^1}, \dots, a_{d-k}^{p^{r_{a_{d-k}}}} = a_{d-k}$ is called the **G -cycle on the coefficient a_{d-k}** . Define the length of the G -cycle on the coefficient a_{d-k} to be this integer r_k .

Observe that for any PP $P(x)$, there is an integer $r \geq 1$, such that the sequence of iterates of the G -map shown in (3) forms a cycle.

Definition 17. *The sequence of iterates of the G -map on the PP $P(x)$, namely*

$$P(x), G(P(x)), G^2(P(x)), \dots, G^r(P(x)) = P(x),$$

*is called the **G -cycle on $P(x)$** .*

Consider the values of k for which the coefficient a_{d-k} in $P(x)$ is nonzero. For all such k , ($1 \leq k \leq d$), let $r_{a_{d-k}} = \min_{1 \leq j \leq m} \{j \mid ip^j \pmod{p^m-1} = i\}$, where $a_{d-k} = t^i$ for some i . The length of the G -cycle on $P(x)$ is $r = \max_k \{r_{a_{d-k}}\}$. It should be noted that, for any element a in $GF(q)$, the length of the G -cycle containing a is a divisor of m . The length of the G -cycle on $P(x)$ is simply the least common multiple of the lengths of the G -cycles on all non-zero coefficients of $P(x)$.

To illustrate G -cycles on coefficients, consider $GF(2^4)$. The elements of $GF(2^4)$ are partitioned into 6 disjoint equivalence classes (*i.e.*, G -cycles on coefficients) by the G -map, namely the equivalence classes $[0]$, $[1]$, $[2]$, $[4]$, $[6]$, and $[8]$. To see this, observe that

$$\begin{aligned} [0] &= \{0^{2^i}\} = \{0\} \text{ for all } i & [6] &= [t^5] = \{(t^5)^{2^i}\} \text{ for all } i \\ & & &= \{(t^5)^{2^0}, (t^5)^{2^1}, (t^5)^{2^2}, \dots\} \\ [1] &= [t^0] = \{(t^0)^{2^i}\} = \{1\} \text{ for all } i & &= \{t^5, t^{10}, t^5, \dots\} \\ [2] &= [t^1] = \{(t^1)^{2^i}\} \text{ for all } i & &= \{6, 11\} \\ &= \{t^{2^0}, t^{2^1}, t^{2^2}, t^{2^3}, t^{2^4}, \dots\} & & \\ &= \{t^1, t^2, t^4, t^8, t^1, \dots\} & [8] &= [t^7] = \{(t^7)^{2^i}\} \text{ for all } i \\ &= \{2, 3, 5, 9\} & &= \{(t^7)^{2^0}, (t^7)^{2^1}, (t^7)^{2^2}, (t^7)^{2^3}, (t^7)^{2^4}, \dots\} \\ [4] &= [t^3] = \{(t^3)^{2^i}\} \text{ for all } i & &= \{t^7, t^{14}, t^{13}, t^{11}, t^7, \dots\} \\ &= \{(t^3)^{2^0}, (t^3)^{2^1}, (t^3)^{2^2}, (t^3)^{2^3}, (t^3)^{2^4}, \dots\} & &= \{8, 12, 14, 15\} \\ &= \{t^3, t^6, t^{12}, t^9, t^3, \dots\} & & \\ &= \{4, 7, 10, 13\} \end{aligned}$$

where $\text{mod } 15$ arithmetic is used in the exponents. The size of each equivalence class is the length of the corresponding G -cycle on the coefficient a_{d-k} . In this case, there are cycles of lengths 1, 2, and 4.

To illustrate the computation of the G -cycle on $P(x)$, consider the degree 7 nPP $P(x) = x^7 + x^5 + 8x^4 + 6x^2 + 4x$ over $GF(2^4)$. For all $k, (1 \leq k \leq d)$, the non-zero coefficients a_{d-k} are $a_5 = 1, a_4 = 8, a_2 = 6$ and $a_1 = 4$, which correspond to $k = 2, 3, 5$, and 6, respectively. For example, the G -cycle for the nonzero coefficient $a_4 = 8 = t^7$, is $[8] = [t^7] = \{8, 12, 14, 15\}$, which gives the 4 successive coefficients of x^4 in the iterates of the G -map on $P(x)$ shown below. We compute the length of the G -cycle for each nonzero coefficient a_{d-k} , namely, $r_{a_{d-k}} = \min_{1 \leq j \leq m} \{j \mid ip^j \pmod{p^m - 1} = i\}$ resulting in $r_{a_5} = r_1 = 1, r_{a_4} = r_8 = 4, r_{a_2} = r_6 = 2, r_{a_1} = r_4 = 4$. By Definition 17, the length of the G -cycle on $P(x)$ is $\max_k \{r_{a_{d-k}}\} = 4$, as verified below.

$$\begin{aligned}
P(x) &= x^7 + x^5 + 8x^4 + 6x^2 + 4x \\
G(P(x)) &= 1^2x^7 + 1^2x^5 + 8^2x^4 + 6^2x^2 + 4^2x \\
&= (t^0)^2x^7 + (t^0)^2x^5 + (t^7)^2x^4 + (t^5)^2x^2 + (t^3)^2x \\
&= (t^0)x^7 + (t^0)x^5 + (t^{14})x^4 + (t^{10})x^2 + (t^6)x \\
&= x^7 + x^5 + 15x^4 + 11x^2 + 7x \\
G^2(P(x)) &= (t^0)^2x^7 + (t^0)^2x^5 + (t^{14})^2x^4 + (t^{10})^2x^2 + (t^6)^2x \\
&= (t^0)x^7 + (t^0)x^5 + (t^{28})x^4 + (t^{20})x^2 + (t^{12})x \\
&= (t^0)x^7 + (t^0)x^5 + (t^{13})x^4 + (t^5)x^2 + (t^{12})x \\
&= x^7 + x^5 + 14x^4 + 6x^2 + 13x \\
G^3(P(x)) &= (t^0)^2x^7 + (t^0)^2x^5 + (t^{13})^2x^4 + (t^5)^2x^2 + (t^{12})^2x \\
&= (t^0)x^7 + (t^0)x^5 + (t^{26})x^4 + (t^{10})x^2 + (t^{24})x \\
&= (t^0)x^7 + (t^0)x^5 + (t^{11})x^4 + (t^{10})x^2 + (t^9)x \\
&= x^7 + x^5 + 12x^4 + 11x^2 + 10x \\
G^4(P(x)) &= (t^0)^2x^7 + (t^0)^2x^5 + (t^{11})^2x^4 + (t^{10})^2x^2 + (t^9)^2x \\
&= (t^0)x^7 + (t^0)x^5 + (t^{22})x^4 + (t^{20})x^2 + (t^{18})x \\
&= (t^0)x^7 + (t^0)x^5 + (t^7)x^4 + (t^5)x^2 + (t^3)x \\
&= x^7 + x^5 + 8x^4 + 6x^2 + 4x \\
&= P(x)
\end{aligned}$$

Note that the lengths of G -cycles on each of the nonzero coefficients $a_d - k, (1 \leq k \leq 7)$ are 1, 4, 2 and 4, respectively, and their least common multiple, namely 4, is the length of the G -cycle on $P(x)$.

3.3 Iterating the F -map and the G -map

We have introduced two functions, the F -map and G -map, that transform PPs into other PPs and nPPs into other nPPs. These functions can be applied sequentially. For example, we can represent the application of the F and G maps alternately two times on the PP $P(x)$ by the sequence

$(G \circ F \circ G \circ F)(P(x))$, meaning one first applies the F -map, then the G -map, the F -map, and finally the G -map again. The F -map and the G -map are used together to create larger equivalence classes, allowing for a faster search for PPs.

It is interesting to note that two different sequences of compositions can represent the same transformation. In the following we show that we can replace any sequence of compositions by an equivalent sequence in which all of the G -maps are applied first, followed by some number of F -maps. The number of F -maps is related to the field characteristic p . Our result is illustrated by the following diagram, which indicates that $(G \circ F)(P(x))$ is the same as $(F^p \circ G)(P(x))$, for all PPs $P(x)$.

$$\begin{array}{ccc} P(x) & \xrightarrow{G} & G(P(x)) \\ F \downarrow & & \downarrow F^p \\ F(P(x)) & \xrightarrow{G} & (F^p \circ G)(P(x)) \end{array}$$

In Lemma 18, we show that any sequence of F -maps and G -maps is equivalent to a sequence F^i, G^j , for some i ($0 \leq i \leq r$) and some j ($0 \leq j \leq s$), where r is the length of the F -cycle and s is the length of the G -cycle.

Lemma 18. *For any PP $P(x)$, $(G \circ F)(P(x)) = (F^p \circ G)(P(x))$.*

Proof. Let $P(x) = \sum_{k=0}^d a_{d-k} x^{d-k}$. If $a_{d-k} = 0$ then the coefficients of x^{d-k} in $(G \circ F)(P(x))$ and $(F^p \circ G)(P(x))$ are both 0. Suppose that $a_{d-k} \neq 0$. Then $a_{d-k} = t^j$, for some j , $0 \leq j \leq q-2$. Then the x^{d-k} -th term in $F(P(x))$, $G(P(x))$, $(G \circ F)(P(x))$, and $(F^p \circ G)(P(x))$, respectively, are:

$$\begin{aligned} F(P(x)) &= \dots + t^{j+k} x^{d-k} + \dots \\ G(P(x)) &= \dots + t^{pj} x^{d-k} + \dots \\ (G \circ F)(P(x)) &= \dots + t^{(j+k)p} x^{d-k} + \dots = \dots + t^{p(j+k)} x^{d-k} + \dots \\ (F^p \circ G)(P(x)) &= \dots + t^{pj+pk} x^{d-k} + \dots = \dots + t^{p(j+k)} x^{d-k} + \dots \end{aligned}$$

Hence, the coefficients of x^{d-k} , $0 \leq k \leq d$ in $(G \circ F)(P(x))$ and $(F^p \circ G)(P(x))$ are equal and the lemma follows. \square

For example, let $P(x)$ be an nPP over $GF(2^5)$. Consider $(G \circ F \circ G \circ F)(P(x))$. Since $p = 2$, by Lemma 18 $(G \circ F)(P(x)) = (F^2 \circ G)(P(x))$. So, $(G \circ F \circ G \circ F)(P(x)) = (G \circ F \circ F^2 \circ G)(P(x)) = (G \circ F^3 \circ G)(P(x))$. Then by an iterative use of Lemma 18, we get $(G \circ F^3 \circ G)(P(x)) = (F^6 \circ G \circ G)(P(x)) = (F^6 \circ G^2)(P(x))$.

4 Equivalence classes based on the F -map and the G -map

The equivalence relations \mathcal{R} and \mathcal{R}_E which we described in Section 2 allow a more efficient search for PPs by limiting the search to nPPs. More inclusive equivalence relations would optimize the search even further, by allowing the search to be restricted to representatives of equivalence classes. In this section, we introduce new equivalence relations, based on the F -map and the G -map and iterations of the maps, that merge equivalence classes, thereby compressing the search space considerably. We begin with an equivalence relation induced by the F -map.

Definition 19. Let $P(x)$ and $Q(x)$ be PPs. If $P(x)$ can be converted into $Q(x)$ by some sequence consisting of normalization operations and F -map operations, then $P(x)$ and $Q(x)$ are related by \mathcal{R}_F .

We have seen that the iterates of the F -map form a cycle. Moreover, if $P(x)$ is an nPP, then the F -cycle on $P(x)$ is a cycle of nPPs. We will show shortly that \mathcal{R}_F is an equivalence relation that is identical to the \mathcal{R}_E relation defined on nPPs (or PPs). We can choose any nPP in an \mathcal{R}_F equivalence class to be the representative for the class, but for convenience we usually designate the nPP with the smallest coefficient of a specific degree to be the representative. To illustrate, let $P(x)$ be an nPP in some equivalence class, let the specific degree be $d-k$ for some k , and consider the coefficient of x^{d-k} , namely a_{d-k} . Let the multiplicative inverse of a_{d-k} be t^{ik} for some i . Suppose the length of the F -cycle on a_{d-k} is $q-1$. Then, sequence $a_{d-k}, t^k a_{d-k}, t^{2k} a_{d-k}, \dots, t^{rk \bmod (q-1)} a_{d-k}$ includes every nonzero element in $GF(q)$. Specifically, one element in the F -cycle is $a_{d-k} t^{ik} = 1$. In other words, in the nPP $F^i(P(x))$, the coefficient of x^{d-k} takes the value 1. So, in this case we choose $F^i(P(x))$ to be the representative of the equivalence class, making the search for nPPs more efficient since the coefficient of x^{d-k} can be fixed to 1. That is, if there is a nPP whose $(d-k)^{th}$ coefficient is nonzero, there is also one whose $(d-k)^{th}$ coefficient is equal to 1. Notice that if the length of the F -cycle on $P(x)$ is less than $q-1$, then there may be a nPP with a nonzero $(d-k)^{th}$ coefficient, but not one with the $(d-k)^{th}$ coefficient equal to 1. In that case, if the cycle length is $(q-1)/j$, for some $j > 1$, then one needs to search for nPPs with a $(d-k)^{th}$ coefficient equal to each of the values $1, 2, 3, \dots, j-1$. Note that if there is an nPP whose $(d-k)^{th}$ coefficient is zero, it would be chosen as the representative for the equivalence class.

So again, in our search for all permutation polynomials over $GF(q)$ for a given q , we can reduce the search to the space of normalized PPs which are representatives of an F -map equivalence class. In other words, when the degree of the polynomial is not a multiple of the field characteristic, we restrict the search to polynomials $a_d x^d + a_{d-1} x^{d-1} + \dots + a_1 x + a_0$ where $a_d = 1, a_0 = a_{d-1} = 0$, and a_i ranges over all cycle representatives, where the i^{th} coefficient has the maximum length cycle. When the degree is a multiple of the field characteristic, and the characteristic is not 2, we restrict the search to polynomials $a_d x^d + a_{d-1} x^{d-1} + \dots + a_1 x + a_0$ where $a_d = 1, a_0 = 0$, and one of the following cases (1) $a_{d-1} = 0$, and (2) $a_{d-1} \neq 0$ and $a_{d-2} = 0$. There is a similar statement when the field characteristic is 2 and the degree of the polynomial is even.

Lemma 20. The equivalence relations \mathcal{R}_F and \mathcal{R}_E are the same.

Proof. Suppose that $P(x)$ and $Q(x)$ are \mathcal{R}_E -related. That is, suppose that $Q(x) = aP(sx+b)+c$ for $a, b, c, s \in GF(q)$ such that $a \neq 0$ and $s \neq 0$. We show that $P(x)$ and $Q(x)$ are \mathcal{R}_F -related. Since $s \neq 0$, we have $s = t^i$, for some i ($0 \leq i \leq q-2$). Consider $P'(x) = F^{q-i-1}(x) = t^{d(q-i-1)} P(x/t^{q-i-1})$, the last equality due to Equation (1). Then $P'(x) = t^{d(q-i-1)} P(sx)$ since t^i is the inverse of t^{q-i-1} . Take $a' = at^{di}$ and $b' = b/s$. Note that $a' \neq 0$. Clearly, P' and $P''(x) = a'P'(x+b') + c$ are \mathcal{R}_F -related. Then

$$\begin{aligned} P''(x) &= a'(t^{d(q-i-1)} P(s(x+b')) + c \\ &= at^{di}(t^{d(q-i-1)} P(sx + sb')) + c \\ &= aP(sx + b) + c \\ &= Q(x). \end{aligned}$$

Therefore $P(x)$ and $Q(x)$ are \mathcal{R}_F -related.

Suppose that $P(x)$ and $Q(x)$ are \mathcal{R}_F -related. We show that $P(x)$ and $Q(x)$ are \mathcal{R}_E -related. Let $P_1(x) = P(x), P_2(x), \dots, P_k(x) = Q(x)$ be a sequence of permutation polynomials such that, for any i , $P_i(x)$ and $P_{i+1}(x)$ are \mathcal{R}_F -related. Suppose $P_{i+1}(x) = F^j(P_i(x))$, for some j ($0 \leq j \leq q-2$). Then, by Equation (1) in Section 3.1, $P_{i+1}(x) = t^{dj} P_i(x/t^j) = aP_i(sx+b) + c$ for $a = t^{dj} \neq 0$, $s = t^{-j} = t^{q-j-1} \neq 0$, and $b = c = 0$. That is, $P_i(x)$ and $P_{i+1}(x)$ are \mathcal{R}_E -related. \square

Recall that one of our goals is to make the search for nPPs more efficient. Lemmas 21 and 22 below, show how iterations of the F -map can be used to restrict the value of one additional coefficient and hence decrease the size of the search space.

Lemma 21 shows that under certain conditions, a fourth coefficient can be restricted to the values 0 and 1.

Lemma 21. *Let s denote the position fixed for normalization, i.e. $s = d-1$ for c -normalization, $s = d-1$ or $s = d-2$ for m -normalization, and $s = d-r-1$ or $s = d-r$, where $r = 2^i - 1$ for some i , for b -normalization. If there is a $k \neq s$ relatively prime to $q-1$, then each equivalence class of \mathcal{R}_F contains an nPP $Q(x) = b_d x^d + b_{d-1} x^{d-1} + \dots + b_{d-k} x^{d-k} + \dots + b_1 x + b_0$ satisfying $b_d = 1$, $b_s = b_0 = 0$ and $b_{d-k} \in \{0, 1\}$, for some $k \in 1, \dots, d-1, k \neq s$.*

Proof. Let $P(x) = \sum_{k=0}^d a_{d-k} x^{d-k}$ be an arbitrary nPP from any equivalence class of \mathcal{R}_F . Then $a_d = 1, a_s = 0$, and $a_0 = 0$. On one hand, if a_{d-k} is 0, then $Q(x) = P(x)$, and the lemma is proved.

On the other hand, suppose that $a_{d-k} = t^i, i \in \{1, 2, \dots, q-2\}$. Since k is relatively prime to $q-1$, there is a positive integer j such that $i + jk \equiv 0 \pmod{q-1}$. Let Q be the polynomial obtained by applying the F map j times to P . That is, let $Q(x) = \sum_{k=0}^d a_{d-k} t^{jk} x^{d-k}$. Then the coefficient of the x^{d-k} term of Q is $b_{d-k} = a_{d-k} t^{kj} = t^{i+jk} = t^{i+kj} = 1$. Furthermore, $b_d = a_d = 1$, $b_0 = a_0 = 0$, and $b_s = a_s t^{(d-s)j} = 0 t^{(d-s)j} = 0$. Hence Q is an nPP with the desired coefficients. \square

It could be that there is no k in the range ($2 \leq k \leq d-2$) that is relatively prime to $q-1$ and Lemma 21 cannot be applied. For example, when $q = 31$ and $d = 8$, then none of the integers in $\{2, 3, 4, 5, 6\}$ are relatively prime to $q-1$. We now show that when this is the case, we can restrict the range of the coefficient of either x^{d-1} or x^{d-2} to at most 3 values.

Lemma 22. *Let s denote the position fixed for normalization, i.e. $s = d-1$ for c -normalization, $s = d-1$ or $s = d-2$ for m -normalization, and $s = d-r-1$ or $s = d-r$, where $r = 2^i - 1$ for some i , for b -normalization. Then each equivalence class of \mathcal{R}_F contains an nPP $Q(x) = b_d x^d + b_{d-1} x^{d-1} + \dots + b_1 x + b_0$ satisfying $b_d = 1$, $b_s = b_0 = 0$ and if $s = d-1$, then $b_{d-2} \in \{0, 1, 2\}$, otherwise, if $s \geq d-2$, then $b_{d-1} \in \{0, 1\}$.*

Proof. Let $P(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_1 x + a_0 = \sum_{k=0}^d a_{d-k} x^{d-k}$ be an arbitrary nPP from any equivalence class of \mathcal{R}_F . Then $a_d = 1, a_s = 0$, and $a_0 = 0$. We will obtain a new nPP $Q(x)$ from $P(x)$ by using the F -map iteratively. The coefficients of $Q(x)$ are changed from the given a_i to a new value b_i , for all i . Since the F -map does not change the first coefficient, $b_d = a_d = 1$. Since the F -map does not change a zero coefficient, $b_s = a_s = 0$, and $b_0 = a_0 = 0$. It remains to prove that if $s = d-1$ then $b_{d-2} \in \{0, 1, 2\}$, otherwise, if $s \geq d-2$, then $b_{d-1} \in \{0, 1\}$.

Case 1: $s = d-1$.

If $a_{d-2} = 0$, then the lemma is proved, as we can take $Q(x) = P(x)$.

So, suppose $a_{d-2} \neq 0$. If 2 is relatively prime to $q-1$, then Lemma 21 fixes b_{d-2} to a value in $\{0, 1\}$. So suppose 2 is not relatively prime to $q-1$. Then $q-1$ must be even, that is

$\gcd(2, q-1) = 2$. Hence, the length of F -cycle on a_{d-2} is $(q-1)/2$. Substituting $k=2$ into the definition of the F -cycle on a_{d-2} yields the sequence

$$a_{d-2}, t^2 a_{d-2}, t^4 a_{d-2}, \dots, (t^{2r} \bmod (q-1)) a_{d-2} = a_{d-2}$$

where $r = (q-1)/2$. So there are two possibilities for the values in F -cycle on a_{d-2} :

- the F -cycle contains $(q-1)/2$ odd values including 1, in which case b_{d-2} can be fixed at 1,
- the F -cycle contains $(q-1)/2$ even values including 2, in which case b_{d-2} can be fixed at 2.

Case 2: $s \geq d-2$.

If $a_{d-1} = 0$, then the lemma is proved, as we can take $Q(x) = P(x)$.

So, suppose $a_{d-1} \neq 0$. Since 1 and $q-1$ are relatively prime, Lemma 21 fixes b_{d-1} to a value in $\{0, 1\}$.

Hence $Q(x)$ is an nPP with the desired coefficients. \square

If Lemma 21 does not apply, then the F -cycle is not of length $q-1$. Since the length of the F -cycle must divide $q-1$, it follows from Lemma 22 that, when the coefficient a_{d-2} is not zero, the F -cycle is of length $(q-1)/2$. That is, as the coefficient a_{d-2} is multiplied by t^2 with each iteration of the F -map, to return in the cycle to the same coefficient a_{d-2} , the length of the cycle must be r , where r is the smallest integer such that $(t^2)^r = 1$. It follows that $r = (q-1)/2$, as this is the smallest r such that $2r = 0 \bmod (q-1)$.

By Lemmas 21 and 22 we have shown that every equivalence class of \mathcal{R}_F contains an nPP in which four of its coefficients are either fixed or are limited to at most three choices. We have implemented this in our search algorithm, **iBlast** (described in Section 5.1). **iBlast** searches for equivalence class representatives in $O(dq^{d-2})$ time rather than $O(dq^{d-1})$ time, as was previously described in Section 1.

In Section 3.3 we discussed iterations of the F -map and the G -map. We now introduce a new equivalence relation, $\mathcal{R}_{F,G}$, based on such iterations.

Definition 23. Let $P(x)$ and $Q(x)$ be nPPs of degree d , for some d . If $P(x)$ can be converted into $Q(x)$ by some sequence consisting of F -map and G -map operations, then $P(x)$ and $Q(x)$ are related by $\mathcal{R}_{F,G}$.

We present a theorem that characterizes the nPPs in the equivalence classes of $\mathcal{R}_{F,G}$, and gives the number of nPPs in each equivalence class.

Theorem 24. Let $P(x)$ be a nPP. Then the equivalence class of $\mathcal{R}_{F,G}$ containing $P(x)$ is

$$[P(x)] = \{F^i(G^j(P)) \mid 0 \leq i \leq r-1, 0 \leq j \leq s-1\},$$

where r and s are the lengths of the F - and G -cycles, respectively, and the number of nPPs in $[P(x)]$ is rs .

Proof. This follows from Lemma 18. Clearly, $[P(x)]$ is a subset of the equivalence class of $\mathcal{R}_{F,G}$ containing $P(x)$. Let $Q(x)$ be a polynomial of $[P(x)]$, that is, let $Q(x) = (F^i \circ G^j)(P)$, for some i, j . Observe that $F(Q(x)) = (F \circ F^i \circ G^j)(P(x)) = (F^{i+1} \circ G^j)(P(x))$, so $F(Q(x)) \in [P(x)]$. It

remains to show that $G(Q(x)) \in [P(x)]$. If $i = 0$ then $G(Q(x)) = G^{j+1}(P(x))$, so $G(Q(x)) \in [P(x)]$. Suppose $i \geq 1$. Then $G(Q(x)) = (G \circ F^i \circ G^j)(P(x))$. As the iterates of the F -map and the G -map form cycles of lengths r and s , respectively, the size of the exponents of F and G are computed by arithmetic $(\text{mod } r)$ and $(\text{mod } s)$, respectively. Applying Lemma 18 i times, we have

$$\begin{aligned} G(Q(x)) &= (G \circ F^i \circ G^j)(P(x)) \\ &= (F^p \circ G \circ F^{i-1} \circ G^j)(P(x)) \\ &= (F^{p+1} \circ G \circ F^{i-2} \circ G^j)(P(x)) \\ &= \dots \\ &= (F^{pi} \circ G^{j+1})(P(x)) \\ &\in [P(x)] \end{aligned}$$

By algebra, the number of nPPs in $[P(x)]$ is rs . The theorem follows. \square

For example, consider the nPP $P(x) = x^8 + 4x^4 + 16x^2 + 3x$ for $GF(2^5)$, with the primitive polynomial $x^5 + x^3 + x^2 + x + 1$. As there are 31 non-zero elements in $GF(31)$, and 31 is prime, the length of the F -cycle on $P(x)$ is 31. The iterates of G -map on the coefficient 16 give the G -cycle $\{16, 31, 30, 27, 24\}$, which has length 5. So, $[P(x)]$ denotes an equivalence class with $31 * 5 = 155$ nPPs.

Theorem 24 shows that combined iterations of the F -map and the G -map on an nPP $P(x)$ can condense smaller equivalence classes induced by the individual maps into fewer (and larger) equivalence classes. So although the G -map by itself (unlike the F -map) does not allow an additional coefficient to be fixed, when used with the F -map, it does allow the search to include fewer equivalence classes. This increases the efficiency of the search for PPs, although not by an order of magnitude.

5 Algorithms for computing equivalence classes of nPPs

5.1 Algorithm iBlast

We have implemented an algorithm called **iBlast** (whose name is derived from our permuted first name initials) that uses Lemmas 21 and 22, and Theorem 24 to make the enumeration of nPPs more efficient. As previously remarked, **iBlast** searches for equivalence class representatives in $O(dq^{d-2})$ time rather than $O(dq^{d-1})$ time (the latter being the time bound of previous algorithms), so we are able to compute new results for various q and d . We exhibit many of these in Table 3 and give specific details in Tables 5 through 28.

iBlast computes the set \mathcal{S} of all nPPs in $GF(q)$ of degree d . Define a *mask* as an array with values in $\{0, 1\}$ associated with a polynomial's coefficients. 0 designates that the corresponding coefficient should be fixed at a particular value, and 1 designates the coefficient is both unfixed and nonzero. The purpose of using masks is to maximize our usage of the F -map and G -map functions. Since both the F -map and G -map take the symbol 0 to itself, masks allow us to consider only nonzero elements when selecting an additional coefficient to fix.

For example, consider the search for nPPs of degree 5 over $GF(25)$. Since $p \mid d$ and $p > 2$, the masks must correspond to m-normalization, that is, to PPs in the form $a_5x^5 + a_4x^4 + a_3x^3 + a_2x^2 + a_1x + a_0$ where $a_5 = 1$, $a_0 = 0$, and either $a_4 = 0$ or $a_3 = 0$. Since a_5 is always fixed at 1,

Algorithm 1: iBlast

```
1 if  $p \nmid d$  then Create the set  $\mathcal{M}$  of all possible masks corresponding to c-normalization
2 else if  $p = 2$  then Create the set  $\mathcal{M}$  of all possible masks corresponding to b-normalization
3 else Create the set  $\mathcal{M}$  of all possible masks corresponding to m-normalization
4  $\mathcal{S} = \emptyset$ 
5 foreach mask  $m \in \mathcal{M}$  do
6   Use the  $F$ -map and  $G$ -map functions to select the optimal coefficient in  $m$  to
   additionally fix
7    $currentPolynomial$  = its default value where each fixed coefficient is assigned its
   designated value, and each unfixed coefficient is assigned 1
8   do
9     if  $currentPolynomial$  is a PP and  $currentPolynomial \notin \mathcal{S}$  then
10       $\mathcal{T}$  = the set of all PPs that are  $\mathcal{R}_{F,G}$ -related to  $currentPolynomial$ 
11      if  $p \mid d$  then
12        foreach PP  $P \in \mathcal{T}$  and  $b \in GF(q)$  do
13           $\mathcal{T} = \mathcal{T} \cup P(x + b)$ 
14        end
15      end
16       $\mathcal{S} = \mathcal{S} \cup \mathcal{T}$ 
17    end
18    increment  $currentPolynomial$ 
19  while  $currentPolynomial$  is not maximized
20 end
21 return  $\mathcal{S}$ 
```

and a_0 is always fixed at 0, we can first consider all cases where a_4 is fixed at 0. **iBlast** generates $2^3 = 8$ masks in the form $[a_5, a_4, a_3, a_2, a_1, a_0]$ to account for all combinations where a_3, a_2 , and a_1 are either fixed at 0 or are unfixed and nonzero. The eight masks are $[0, 0, 0, 0, 0, 0]$, $[0, 0, 0, 0, 1, 0]$, $[0, 0, 0, 1, 0, 0]$, $[0, 0, 0, 1, 1, 0]$, $[0, 0, 1, 0, 0, 0]$, $[0, 0, 1, 0, 1, 0]$, $[0, 0, 1, 1, 0, 0]$, $[0, 0, 1, 1, 1, 0]$. **iBlast** also creates an additional 4 masks for the cases where a_3 is fixed at 0, and a_2 is unfixed and nonzero, namely $[0, 1, 0, 0, 0, 0]$, $[0, 1, 0, 0, 1, 0]$, $[0, 1, 0, 1, 0, 0]$, $[0, 1, 0, 1, 1, 0]$.

iBlast then iterates through the masks to search for nPPs. Each mask has at least 3 coefficients fixed due to normalization, and Lemmas 21 and 22 fix an additional coefficient, giving a total of 4 fixed coefficients.

For example, consider the G -cycles and F -cycles for the mask $[0, 1, 0, 1, 0, 0]$ in Table 2. Observe that for the F -cycles of a_2 we can use the G -map to map every element in the F -cycle that contains 2 (*i.e.*, the F -cycle $\{2, 5, 8, 11, 14, 17, 20, 23\}$) to some element in the F -cycle that contains 6 (*i.e.*, the F -cycle $\{3, 6, 9, 12, 15, 18, 21, 24\}$). If we were to therefore consider fixing the coefficient a_2 , we would only need to check polynomials where a_2 had a value of 1 or 3. However, a_4 is the better choice as it can be fixed to the single value 1 because for a_4 , the F -cycle that contains 1 also contains all non-zero elements of $GF(25)$.

iBlast then begins the search by creating a polynomial variable, $currentPolynomial$, and setting its coefficients to their assigned value if they are fixed, or 1 if they are unfixed. For our example mask $[0, 1, 0, 1, 0, 0]$, this would correspond to the polynomial $x^5 + x^4 + x^2$. **iBlast**

<i>G</i> -cycles		<i>F</i> -cycles for a_4	<i>F</i> -cycles for a_2
{0}	{9, 17}	{0}	{0}
{1}	{10, 22}	{1, 2, 3, ..., 24}	{1, 4, 7, 10, 13, 16, 19, 22}
{2, 6}	{13}		{2, 5, 8, 11, 14, 17, 20, 23}
{3, 11}	{14, 18}		{3, 6, 9, 12, 15, 18, 21, 24}
{4, 16}	{15, 23}		
{5, 21}	{19}		
{7}	{20, 24}		
{8, 12}			

Table 2: *G*-cycles and *F*-cycles for the mask $[0, 1, 0, 1, 0, 0]$ in $GF(25)$ degree 5.

increments the polynomial as a $d + 1$ -tuple and adds 1 to the lowest degree, unfixed coefficient. If a coefficient exceeds its maximum value, **iBlast** resets it to 1, since unfixed coefficients are nonzero, and the next highest unfixed coefficient is incremented. Note that if the coefficient fixed by the *F*-map and *G*-map does not reduce to a single value, we can include it in our increment function, but it only increments through the minimum necessary values as determined by the *F*-cycles and *G*-cycles.

Each time *currentPolynomial* is incremented, it is checked to determine if it is a PP. If *currentPolynomial* is a new PP, *i.e.*, it is not already in the set \mathcal{S} , **iBlast** uses the *F*-map and *G*-map functions to create the set \mathcal{T} of all PPs that are $\mathcal{R}_{F,G}$ -related to *currentPolynomial*.

If $p \mid d$ and $p > 2$, m-normalization applies. In this case, for each PP in \mathcal{T} , **iBlast** computes $P(x + b)$, for all b in $GF(q)$, and puts the resulting PPs in the set \mathcal{T} . Note that this must be done for any search where $p \mid d$ since both m-normalization and b-normalization make use of this normalization operation. Finally **iBlast** adds all of the newly generated PPs to \mathcal{S} by setting $\mathcal{S} = \mathcal{S} \cup \mathcal{T}$ and then continues to increment *currentPolynomial* until it reaches its maximum configuration.

Reaching the maximum configuration signifies that the process can be repeated for the next mask. **iBlast** terminates when all of the masks have been considered. At this point, the set \mathcal{S} contains all nPPs in $GF(q)$ of degree d .

5.2 Algorithm 2

Given the set \mathcal{S} of all nPPs in $GF(q)$, the following algorithm will create a set of equivalence class representatives.

While \mathcal{S} is not empty, select an arbitrary $P(x)$ in \mathcal{S} . Compute the set \mathcal{T} of all nPPs that are $\mathcal{R}_{F,G}$ -related to $P(x)$. If $p \mid d$, additionally compute all possible combinations of $P(x + b)$ for each nPP in \mathcal{T} , adding each result to \mathcal{T} . Designate $P(x)$ as an equivalence class representative, and $\mathcal{S} = \mathcal{S} - \mathcal{T}$.

6 Results

As stated in Section 1, a brute force search for degree d permutation polynomials over $GF(q)$ would require $O(dq^{d+2})$ time. Normalization operations defined in [19], for PPs in which $p \nmid d$, fixes three

of the coefficients and therefore requires $O(dq^{d-1})$ time. We refer to this type of normalization as c-normalization. In this paper, we have succeeded in improving the time bound by an order of magnitude, that is, lowering it to $O(dq^{d-2})$, by fixing an additional coefficient. This improvement applies to a larger class of PPs, namely all c-normalized PPs, all PPs for which $p \nmid d$ and $p > 2$ (i.e., m-normalized PPs), and all PPs for which $p \nmid d$ and $p > 2$, except when $d = 2^i - 2$ for some $i \geq 2$ (i.e., b-normalized PPs). We have done this by expanding the definition of normalization to include m-normalization and b-normalization, and by introducing four new equivalence relations on PPs and nPPs, namely, \mathcal{R}_E , \mathcal{R}_F , \mathcal{R}_G , and $\mathcal{R}_{F,G}$. We have been able to reduce the search space for PPs by limiting the search to equivalence class representatives. In addition, equivalence classes allow a more succinct categorization of PPs, since the equivalence classes can include quite a number of PPs. Furthermore, our new techniques apply to arbitrary q and d .

We implemented our search for equivalence classes in the algorithm **iBlast** and computed many new results which are shown in Table 5. For almost all $q \leq 100$ and $d \leq 10$, **iBlast** found all nPPs and all equivalence classes. Table 5 lists the number of nPPs, the number of equivalence classes, and total number of PPs for $q \leq 97$ and degree d , where $6 \leq d \leq 10$. Note that Table 5 has columns for $d \in \{6, 7, 8, 9, 10\}$. For degrees $d \leq 5$, all PPs have been described, for example in [8]. More recent work [11, 12, 13, 18, 24] gives all PPs of degree $d \leq 8$; however, we list our results in Table 5 for completeness. Table 5 does not have columns for $d \geq 11$, because the computations become too time consuming (at least for large q). However, we have been able to compute all degree 11 PPs over $GF(q)$, for powers of primes q ($16 \leq q \leq 32$). The results are listed in Table 3. The sum of $N_d(q)$, for $1 \leq d \leq 11$, for prime powers q , also gives improved lower bounds for $M(q, q - 11)$ shown in Table 6. For example, we show that $M(16, 5) \geq 5,112,053,760$, which improves on the lower bound given in [27]. We also computed $N_{12}(17) = 68,126,982,656$, $N_{12}(19) = 46,631,675,376$, $N_{12}(23) = 13,755,394,444$. This yields $M(17, 5) \geq 72,377,516,320$, $M(19, 7) \geq$, and $M(23, 11) \geq 14,341,972,920$.

q	$N_{11}(q)$	number of nPPs	number of equivalence classes
16	4,751,093,760	1,237,264	20,663
17	4,001,494,000	865,375	54,225
19	2,431,915,488	374,256	20,874
23	0	0	0
25	6,509,295,000	433,953	9,266
27	2,826,989,100	149,150	2,060
29	1,014,518,484	43,083	1,639
31	385,053,480	13,356	507
32	190,940,160	6,015	51

Table 3: Number of PPs and nPPs for degree 11 polynomials over $GF(q)$.

We also provide a website that explicitly lists nPPs and PPs of degree $1 \leq d \leq 10$ and the values of $n \leq 100$ at <https://personal.utdallas.edu/~bdm170430/npps/>. The nPPs are computed using normalizations from Table 1. The equivalence classes are defined using the relations \mathcal{R} , \mathcal{R}_F , and $\mathcal{R}_{F,G}$. The number of equivalence classes is given in parentheses. The total number of PPs is computed from the number of nPPs. In looking at Tables 5, and Tables 7 through 28, one should keep in mind that the specific nPPs listed in the tables are for the stated primitive polynomial and

for our naming convention for elements of $GF(q)$. Our results for degrees 7 and 8 agree with those listed in [12], [13], and [11], except for differences caused by naming conventions.

In some cases, there is a small number of equivalence classes representing a large number of PPs. For example, $N_8(27) = 6,899,256$, and there are 364 nPPs, but only 6 equivalence classes as shown in Table 12. Compare our list of classes of degree 8 PPs over $GF(27)$ with the list given in [13], which has 26 nPPs. The difference is that we often combined three of his nPPs into one class by the use of our $R_{F,G}$ relation. This gives a method to make rather concise representations of large sets of PPs. Again, for example, $N_9(32) = 9,872,384$, and there are 311 nPPs, but only 7 equivalence classes as shown in Table 15. Observe that, for the second through sixth nPPs in Table 15, there is a term, either x^3 , x^6 , or x^7 , with coefficient 1. Since the number of non-zero elements is 31, which is prime, the length of the F -map has length 31. Since $G(1) = 1$, the G -map produces no other nPPs in the equivalence classes. So, each equivalence class has 31 elements, as defined by the F -map alone. For the last PP in Table 15, again the length of the F -cycle on that PP is 31, but now the length of the G -cycle is applied, for example, to the coefficient 16, has length 5, namely $(16,31,30,27,24)$. Thus, the equivalence class of nPPs produced by the F -map and G -map combined is 31 times 5 or 155.

We note that it was stated in [26] that 32 hours were required to compute the RS code for $q = 32$ and $d = 5$; for $q = 32$ and $d = 7$, the authors were able to compute only the size of the set of PPs, not the set itself. Our algorithm `iBlast` allows for the computation of all nPPs, for $q = 32$ and $d = 7$, in a few seconds and, for $q = 32$ and $d = 8$, in about 10 minutes.

It should be noted that in many of our tables (Tables 7 through 28) we use variables to further reduce the table's size. For example, for $q = 25$ and $d = 9$ (see Table 10) there are 38 equivalence classes, which is rather large. Instead, in Table 10, we use a variable a , with specified values, which allows the table to have 21 classes rather than 38. For example, the first class shown is $x^9 + ax$, where $a \in \{1, 7\}$. This is the union of two equivalence classes, namely those with the representatives $x^9 + x$ and $x^9 + 7x$. The variable a is also used in other tables. In Table 11, $\alpha(x)$ represents three sets of values for low order coefficients in the polynomial shown in the first column.

Our results on the number of PPs also give several new lower bounds for $M(n, D)$. These are given in Table 6. Several additional improved lower bounds can be obtained from those shown in Table 6 using the well known inequality $M(n, D - 1) \geq M(n, D)/n$ [8] or the operation of contraction [1].

Lidl and Mullen [19, 20] give a list of several interesting open questions about PPs. For example, they ask about *complete* PPs, where a PP $P(x)$ is called *complete* if $P(x) + x$ is also a PP. There are several complete PPs in our tables. For example, x^9 and $x^9 + x$ are both PPs for $q = 3^4$.

For the reader's convenience, we provide a master list of tables in Table 4.

Acknowledgments

We thank Professor Carlos Arreche of the Mathematics Department at UTD for bringing our attention to Galois field orbits. We also want to thank Professor Xiang Fan of Mathematics Department Sun Yat Sen University in Guangzhou, China for his valuable comments and suggestions, including the use of Lucas's theorem.

Table 3	Number of PPs and nPPs for degree 11 polynomials over $GF(q)$
Table 5	Number of nPPs, Equivalence Classes, and Total number of PPs for $q \leq 97$ and degree d , where $6 \leq d \leq 10$
Table 6	New lower bounds for $M(n, D)$
Table 7	nPP Equivalence Classes for $q = 11$, degree 7, and primitive polynomial $x + 4$
Table 8	nPP Equivalence Classes for $q = 23$, degree 7, and primitive polynomial $x + 12$
Table 9	nPP Equivalence Classes for $q = 25$, degree 7, and primitive polynomial $x^2 + 3x + 3$
Table 10	nPP Equivalence Classes for $q = 25$, degree 9, and primitive polynomial $x^2 + 3x + 3$
Table 11	nPP Equivalence Classes for $q = 25$, degree 10, and primitive polynomial $x^2 + 3x + 3$
Table 12	nPP Equivalence Classes for $q = 27$, degree 8, and primitive polynomial $x^3 + 2x + 1$
Table 13	nPP Equivalence Classes for $q = 27$, degree 9, and primitive polynomial $x^3 + 2x + 1$
Table 14	nPP Equivalence Classes for $q = 32$, degree 8, and primitive polynomial $x^5 + x^3 + x^2 + x + 1$
Table 15	nPP Equivalence Classes for $q = 32$, degree 9, and primitive polynomial $x^5 + x^3 + x^2 + x + 1$
Table 16	nPP Equivalence Classes for $q = 37$, degree 7, and primitive polynomial $x + 13$
Table 17	nPP Equivalence Classes for $q = 41$, degree 9, and primitive polynomial $x + 17$
Table 18	nPP Equivalence Classes for $q = 43$, degree 9, and primitive polynomial $x + 13$
Table 19	nPP Equivalence Classes for $q = 43$, degree 10, and primitive polynomial $x + 13$
Table 20	nPP Equivalence Classes for $q = 47$, degree 9, and primitive polynomial $x + 12$
Table 21	nPP Equivalence Classes for $q = 49$, degree 9, and primitive polynomial $x^2 + 6x + 3$
Table 22	nPP Equivalence Classes for $q = 49$, degree 10, and primitive polynomial $x^2 + 6x + 3$
Table 23	nPP Equivalence Classes for $q = 71$, degree 9, and primitive polynomial $x + 38$
Table 24	nPP Equivalence Classes for $q = 73$, degree 7, and primitive polynomial $x + 20$
Table 25	nPP Equivalence Classes for $q = 79$, degree 7, and primitive polynomial $x + 25$
Table 26	nPP Equivalence Classes for $q = 81$, degree 9, and primitive polynomial $x^4 + 2x^3 + 2$
Table 27	nPP Equivalence Classes for $q = 89$, degree 7, and primitive polynomial $x + 76$
Table 28	Number of nPPs in Equivalence Classes for prime powers q and degree d , for given primitive polynomial

Table 4: List of tables of results.

References

- [1] S. Bereng, A. Levy, and I. H. Sudborough. Constructing permutation arrays from groups. *Designs, Codes and Cryptography*, 86(5):1095–1111, 2018.
- [2] S. Bereng, Z. Miller, L. G. Mojica, L. Morales, and I. H. Sudborough. New lower bounds for permutation arrays using contraction. *Designs, Codes and Cryptography*, 87:2105–2128, 2019.
- [3] S. Bereng, L. G. Mojica, L. Morales, and H. Sudborough. Kronecker product and tiling of permutation arrays for hamming distances. In *the 2017 IEEE International Symposium on Information Theory (ISIT)*, pages 2198–2202, 2017.
- [4] S. Bereng, L. G. Mojica, L. Morales, and I. H. Sudborough. Parallel partition and extension. In *51st Annual Conference on Information Sciences and Systems, CISS 2017*, pages 1–6, 2017.

- [5] S. Berreg, L. G. Mojica, L. Morales, and I. H. Sudborough. Constructing permutation arrays using partition and extension. *Designs, Codes and Cryptography*, 2019.
- [6] S. Berreg, L. Morales, and I. H. Sudborough. Extending permutation arrays: improving MOLS bounds. *Designs, Codes and Cryptography*, 83(3):661–683, 2017.
- [7] P. J. Cameron. *Combinatorics: Topics, Techniques and Algorithms*. Cambridge University Press, 1995.
- [8] W. Chu, C. J. Colbourn, and P. Dukes. Constructions for permutation codes in powerline communications. *Designs, Codes and Cryptography*, 2004.
- [9] C. Colbourn, T. Kløve, and A. C. Ling. Permutation arrays for powerline communication and mutually orthogonal latin squares. *IEEE Transactions on Information Theory*, 50(6):1289–1291, 2004.
- [10] L. E. Dickson. The analytic representation of substitutions on a power of a prime number of letters with a discussion of the linear group, part II. *Ann. of Math*, 11:65–120, 1896-1897.
- [11] X. Fan. A classification of permutation polynomials of degree 7 over finite fields. *Finite Fields and Their Applications*, 59:1–21, 2019.
- [12] X. Fan. Permutation polynomials of degree 8 over finite fields of characteristic 2. *arXiv e-prints*, page arXiv:1812.06886, 2019.
- [13] X. Fan. Permutation polynomials of degree 8 over finite fields of odd characteristic. *Bulletin of the Australian Mathematical Society*, pages 1–16, 2019.
- [14] F. Gao, Y. Yang, and G. Ge. An improvement on the Gilbert-Varshamov bound for permutation codes. *IEEE Trans. Information Theory*, 59(5):3059–3063, 2013.
- [15] X. Hou. Permutation polynomials over finite fields - a survey of recent advances. *Finite Fields and their Applications*, 32:82–119, 2015.
- [16] I. Janiszczak, W. Lempken, P. R. J. Östergård, and R. Staszewski. Permutation codes invariant under isometries. *Des. Codes Cryptogr.*, 75(3):497–507, 2015.
- [17] I. Janiszczak and R. Staszewski. Isometry invariant permutation codes and mutually orthogonal Latin squares. *arXiv e-prints*, page arXiv:1812.06886, Dec 2018.
- [18] J. Li, D. B. Chandler, and Q. Xiang. Permutation polynomials of degree 6 or 7 over finite fields of characteristic 2. *Finite Fields Appl.*, 16:406–419, 2010.
- [19] R. Lidl and G. L. Mullen. When does a polynomial over a finite field permute the elements of the fields? *Am. Math. Monthly*, 95(3):243–246, 1988.
- [20] R. Lidl and G. L. Mullen. When does a polynomial over a finite field permute the elements of the fields?, II. *Am. Math. Monthly*, 100(1):71–74, 1993.
- [21] R. Lidl and H. Niederreiter. *Introduction to finite fields and their applications*. Cambridge University Press, revised edition, 1994.

- [22] G. Micheli and A. Neri. New lower bounds for permutation codes using linear block codes. *IEEE Transactions on Information Theory*, to appear, 2019.
- [23] Q. T. Nguyen. *Transitivity and hamming distance of permutation arrays*. Phd thesis, University of Texas at Dallas, Richardson, TX 75080, USA, 2013.
- [24] C. J. Shallue and I. M. Wanless. Permutation polynomials and orthomorphism polynomials of degree six. *Finite Fields and Their Applications*, 20:84–92, 2013.
- [25] D. H. Smith and R. Montemanni. A new table of permutation codes. *Designs, Codes and Cryptography*, 63(2):241–253, 2012.
- [26] R. Sobhani, A. Abdollahi, J. Bagherian, and M. Khatami. A note on good permutation codes from Reed–Solomon codes. *Designs, Codes and Cryptography*, Feb 2019.
- [27] X. Wang, Y. Zhang, Y. Yang, and G. Ge. New bounds of permutation codes under hamming metric and kendall’s τ -metric. *Des. Codes Cryptography*, 85(3):533–545, 2017.

Table 5: Number of nPPs, Equivalence Classes, and Total number of PPs for $q \leq 97$ and degree d , where $6 \leq d \leq 10$

q	6	7	8	9	10
11					
nPPs (ECs)	24 (4)	225 (28)	2,754 (277)	29,985 (3,036)	
Total PPs	29,040	272,250	3,332,340	36,281,850	
13					
nPPs (ECs)	0	115 (15)	1,380 (117)	16,740 (1,422)	218,020 (18,193)
Total PPs	0	233,220	2,798,640	33,948,720	442,144,560
16					
nPPs (ECs)	840 (3)	216 (7)	14,816 (57)	4,200 (74)	1,417,600 (1,786)
Total PPs	201,600	829,440	3,555,840	16,128,000	340,224,000
17					
nPPs (ECs)	0	209 (20)	0	3,023 (201)	50,608 (3,163)
Total PPs	0	966,416	0	13,978,352	234,011,392
19					
nPPs (ECs)	0	112 (12)	864 (48)	0	19,544 (1,094)
Total PPs	0	727,776	5,614,272	0	126,996,912
23					
nPPs (ECs)	0	89 (6)	154 (7)	3,092 (174)	50,402 (2,291)
Total PPs	0	1,035,782	1,792,252	35,984,696	586,578,476
25					
nPPs (ECs)	0	45 (5)	0	1,038 (38)	401,280 (341)
Total PPs	0	675,000	0	15,570,000	240,768,000
27					
nPPs (ECs)	702 (1)	14 (2)	364 (6)	29,550 (41)	7,098 (95)
Total PPs	492,804	265,356	6,899,256	20,744,100	134,535,492
29					
nPPs (ECs)	0	0	32 (2)	1,751 (67)	1,568 (56)
Total PPs	0	0	753,536	41,232,548	36,923,264
31					
nPPs (ECs)	0	106 (6)	30 (1)	630 (22)	0
Total PPs	0	3,055,980	864,900	18,162,900	0
32					
nPPs (ECs)	1,024 (2)	32 (2)	19,624 (68)	311 (7)	410,720 (91)
Total PPs	1,015,808	1,015,808	19,467,008	9,872,384	407,434,240
37					
nPPs (ECs)	0	37 (3)	0	0	216 (10)
Total PPs	0	1,823,508	0	0	10,645,344
Continued on next page					

Table 5 – continued from previous page

q	6	7	8	9	10
41					
nPPs (ECs)	0	1 (1)	0	331 (16)	0
Total PPs	0	67,240	0	22,256,440	0
43					
nPPs (ECs)	0	0	0	42 (2)	98 (3)
Total PPs	0	0	0	3,261,636	7,610,484
47					
nPPs (ECs)	0	47 (3)	0	116 (4)	0
Total PPs	0	4,775,858	0	11,787,224	0
49					
nPPs (ECs)	0	3,961 (10)	0	96 (3)	16 (1)
Total PPs	0	9,316,272	0	11,063,808	1,843,968
53					
nPPs (ECs)	0	53 (3)	0	53 (2)	0
Total PPs	0	7,741,604	0	7,741,604	0
59					
nPPs (ECs)	0	59 (3)	0	117 (4)	0
Total PPs	0	11,911,982	0	23,622,066	0
61					
nPPs (ECs)	0	61 (3)	0	0	0
Total PPs	0	13,618,860	0	0	0
64					
nPPs (ECs)	0	0	80,968 (214)	0	21,120 (8)
Total PPs	0	0	326,462,976	0	85,155,840
67					
nPPs (ECs)	0	67 (3)	0	0	0
Total PPs	0	19,850,358	0	0	0
71					
nPPs (ECs)	0	0	0	71 (2)	0
Total PPs	0	0	0	25,053,770	0
73					
nPPs (ECs)	0	73 (3)	0	0	0
Total PPs	0	28,009,224	0	0	0
79					
nPPs (ECs)	0	79 (3)	0	0	
Total PPs	0	38,457,042	0	0	
81					
nPPs (ECs)	0	81 (3)	0	471,891 (55)	0
Total PPs	0	42,515,280	0	3,057,853,680	0
Continued on next page					

Table 5 – continued from previous page

q	6	7	8	9	10
83					
nPPs (ECs)	0	1 (1)	0	83 (2)	
Total PPs	0	564,898	0	46,886,534	
89					
nPPs (ECs)	0	89 (3)	0	89 (2)	
Total PPs	0	62,037,272	0	62,037,272	
97					
nPPs (ECs)	0	1 (1)	0	0	
Total PPs	0	903,264	0	0	

Table 5: Number of nPPs, Equivalence Classes, and Total number of PPs for $q \leq 97$ and degree d , where $6 \leq d \leq 10$

n	D	$M(n, D) \geq$	Previous	n	D	$M(n, D) \geq$	Previous
16	5	5,112,053,760	143,866,479	41	32	22,392,560	1,565,096
17	6	4,250,533,664	143,866,479	43	38	3,341,100	397,074
18	9	72,480,384*	73,195,200	47	38	21,442,716	103,776
18	10	12,240,000*	1,269,376	47	40	9,655,492	103,776
19	8	2,565,261,288	143,866,479	47	42	4,879,634	103,776
24	20	24,288*	23,782	49	39	23,341,648	207,552
25	14	6,766,500,000	143,866,479	49	40	20,497,680	207,552
25	15	257,205,000	244,823,040	49	42	9,433,872	207,552
25	18	867,000	279,818	53	44	23,373,636	470,400
25	20	192,060	19,404	53	46	15,632,032	148,824
27	16	2,990,448,396	143,866,479	53	48	7,890,428	148,824
27	17	163,459,296	9,313,200	59	50	35,941,266	4,762,368
27	18	289,233,804	9,313,200	59	52	12,319,200	1,339,416
27	19	8,179,204	1,326,000	59	54	407,218	205,320
27	21	1,015,092	249,600	61	54	13,622,520	410,640
27	22	522,288	31,200	64	55	332,236,800	262,080
29	18	1,093,475,740	143,866,479	64	59	5,773,824	262,080
29	19	78,957,256	9,533,160	64	60	5,515,776	262,080
29	20	42,033,992	9,533,160	67	60	39,705,138	524,160
31	20	407,138,190	143,866,479	67	62	19,854,780	524,160
31	22	22,084,719	1,291,080	71	62	25,411,610	601,392
31	23	3,921,819	1,291,080	73	66	56,023,704	357,840
31	24	3,056,919	372,992	73	68	28,014,480	357,840
32	21	460,134,208	143,866,479	79	72	38,950,002	492,960
32	22	440,194,048	1,291,080	81	72	3,100,641,120	571,704
32	23	32,759,808	1,291,080	81	74	42,787,440	571,704
32	24	22,887,424	372,992	83	74	94,909,670	888,729
32	25	3,420,416	372,992	83	76	48,023,136	571,704
32	26	2,404,608	208,320	83	78	47,458,238	571,704
32	27	1,388,800	372,992	89	80	125,476,464	1,062,720
32	29	33,728	32,736	89	82	63,439,192	704,880
37	27	14,293,692	1,473,120	89	84	1,401,920	704,880
37	30	3,648,348	155,122	97	90	88,529,184	912,576
37	32	1,824,840	50,616	97	92	87,625,920	912,576

Table 6: New lower bounds for $M(n, D)$. Note: * denotes values that were obtained by coset search [5], not by permutation polynomials. We include these for the sake of completeness.

class	a	number of nPPs
x^7	-	1
$x^7 + ax^2$	{3,5}	4
$x^7 + x^4 + 7x$	-	10
$x^7 + x^4 + ax^2 + 3x$	{9,10}	20
$x^7 + x^5 + 3x^3 + ax$	{2,4,9}	15
$x^7 + x^5 + 3x^3 + 5x^2 + 8x$	-	10
$x^7 + x^5 + 2ax^4 + 3x^3 + 4a^{-3}x^2 + 3x$	{1,2}	20
$x^7 + x^5 + 2ax^4 + 3x^3 + 5x^2$	{1,4}	20
$x^7 + x^5 + 4x^4 + 3x^3 + 9x^2 + 10x$	-	10
$x^7 + 2x^5 + 5x^3 + ax$	{2,3,4}	15
$x^7 + 2x^5 + 5x^3 + 2x^2 + x$	-	10
$x^7 + 2x^5 + ax^4 + 5x^3 + 8a^{-6}x^2 + 4x$	{1,2}	20
$x^7 + 2x^5 + 2x^4 + 5x^3 + 3x^2 + 8x$	-	10
$x^7 + 2x^5 + 2ax^4 + 5x^3 + 10a^{-7}x^2 + 5a^4x$	{1,2}	20
$x^7 + 2x^5 + 4ax^4 + 5x^3 + 5a^{-2}x^2 + 9a^{-1}x$	{1,2}	20
$x^7 + 2x^5 + 5x^4 + 5x^3 + 8a^2x^2 + 5a^4x$	{1,2}	20
TOTAL		225

Table 7: nPP Equivalence Classes for $q = 11$, degree 7, and primitive polynomial $x + 4$.

class	number of nPPs
x^7	1
$x^7 + x^5 + 4x^3 + 9x$	11
$x^7 + x^5 + x^4 + 21x^2 + 7x$	22
$x^7 + x^5 + 6x^4 + 2x^3 + 6x^2 + 7x$	22
$x^7 + x^5 + 11x^4 + 3x^3 + 10x^2 + 22x$	22
$x^7 + 2x^5 + 6x^3 + 12x$	11
TOTAL	89

Table 8: nPP Equivalence Classes for $q = 23$, degree 7, and primitive polynomial $x + 12$.

class	number of nPPs
x^7	1
$x^7 + 2x$	8
$x^7 + x^5 + x^3 + 13x$	12
$x^7 + 2x^5 + 3x^3$	12
$x^7 + 2x^5 + 3x^3 + 16x$	12
TOTAL	45

Table 9: nPP Equivalence Classes for $q = 25$, degree 7, and primitive polynomial $x^2 + 3x + 3$.

class	a	number of nPPs
$x^9 + ax$	$\{1,7\}$	6
$x^9 + 2ax^5 + 9a^2x$	$\{1,3\}$	12
$x^9 + x^7 + 2ax^5 + 22x^3 + 8a^{-1}x$	$\{1,2\}$	48
$x^9 + x^7 + 13x^5 + 4x^3 + ax$	$\{5,6,22\}$	72
$x^9 + 2x^7 + 10ax^3 + 23a^{-1}x$	$\{1,7\}$	24
$x^9 + 2x^7 + 4x^5 + 4a^5x^3 + 21a^{-7}x$	$\{1,2\}$	48
$x^9 + 2x^7 + 12a^9x^5 + 4a^{18}x^3 + 17a^{-12}x$	$\{1,2\}$	36
$x^9 + 2x^7 + 3x^5$	-	12
$x^9 + 2x^7 + 9x^5 + 5x$	-	12
$x^9 + x^7 + 9x^5 + 16a^3x^3 + 10a^2x$	$\{1,3\}$	48
$x^9 + x^7 + 3a^5x^5 + 16a^2x^3 + 10a^2x$	$\{1,2\}$	48
$x^9 + x^7 + 14ax^5 + 3a^4x^4 + 12x^3 + 11a^2x^2$	$\{1,2\}$	96
$x^9 + 2x^7 + 5a^{11}x^5 + x^3 + 8a^{-7}x^4 + 18a^6x^3 +$ $+17a^6x^3 + 17a^{-1}x^2 + 9a^{-7}x$	$\{1,2\}$	96
$x^9 + x^7 + 3a^6x^5 + 9a^{-2}x^4 + 20a^{-8}x^3 + 9a^{-8}x^2 + 6a^7x$	$\{1,2\}$	96
$x^9 + 2x^7 + 9x^5 + 4a^6x^4 + 18a^{-14}x^3 + 18a^{-17}x^2 + 8a^{-4}x$	$\{1,2\}$	96
$x^9 + 2x^7 + 2a^5x^5 + 16a^{-6}x^3 + a^{14}x$	$\{1,2\}$	96
$x^9 + 2x^7 + 11x^5 + 3a^2x^4 + 8a^9x^3 + 23a^{-1}x^2 + 9x$	$\{1,2\}$	96
$x^9 + x^7 + 7x^5 + 4x^2 + x$	-	24
$x^9 + 2x^7 + 9x^5 + 16a^4x^3 + 5a^{12}x$	$\{1,2\}$	24
$x^9 + 2x^7 + 2x^5 + x^4 + 19x^3 + 14x^2 + 23x$	-	48
$x^9 + 2x^7 + 4x^5 + 11x^4 + 8x^3 + 9x$	-	48
TOTAL		1038

Table 10: nPP Equivalence Classes for $q = 25$, degree 9, and primitive polynomial $x^2 + 3x + 3$.

class	$\alpha(x)$	number of nPPs
$x^{10} + 2x^8 + \alpha(x)$	$\{3x^6 + 4x^5 + 4x^4 + x^3 + 22x^2 + 21x,$ $10x^6 + 4x^5 + 20x^4 + 6x^2 + 23x,$ $12x^6 + 10x^5 + 24x^4 + 6x^3 + 7x^2 + 16x\}$	3600
TOTAL		3600

Table 11: nPP Equivalence Classes for $q = 25$, degree 10, and primitive polynomial $x^2 + 3x + 3$.

class	number of nPPs
$x^8 + x^6 + 14x^4 + x^3 + 14x^2$	26
$x^8 + x^6 + x^5 + 14x^4 + x^3 + 14x^2$	26
$x^8 + x^6 + 2x^5 + 4x^4 + 15x^3 + 14x^2 + 7x$	78
$x^8 + x^6 + 2x^5 + 24x^4 + 9x^3 + 14x^2 + 11x$	78
$x^8 + x^6 + 8x^5 + 11x^4 + 16x^3 + 14x^2 + 6x$	78
$x^8 + 2x^6 + 6x^5 + 26x^4 + 6x^3 + 17x^2 + 2x$	78
TOTAL	364

Table 12: nPP Equivalence Classes for $q = 27$, degree 8, and primitive polynomial $x^3 + 2x + 1$.

class	a	number of nPPs
x^9	-	1
$x^9 + x$	-	13
$x^9 + x^3 + ax$	$\{0,2,3,5,8,9,14\}$	221
$x^9 + 2x^3 + ax$	$\{1,5,6,11,12,24\}$	182
$x^9 + x^5 + x^3 + x$	-	351
$x^9 + x^5 + 14x^3 + ax$	$\{0,1\}$	702
$x^9 + 2x^5 + 3x$	-	351
$x^9 + 2x^5 + 6a^2x^3 + 24ax$	$\{1,19\}$	1404
$x^9 + x^6 + ax^5 + x^3 + 14ax^2 + a^2x$	$\{1,8,9,15\}$	7020
$x^9 + x^7 + x^3 + ax$	$\{1,14\}$	702
$x^9 + x^7 + 3ax^3 + 12a^{-1}x$	$\{1,12\}$	1404
$x^9 + x^7 + a^2x^4 + x^3 + 14a^2x^2 + 14ax$	$\{1,3\}$	2808
$x^9 + x^7 + 3ax^4 + x^3 + 16ax^2 + 19a^{10}x$	$\{1,6\}$	4212
$x^9 + x^7 + 2x^4 + a^{16}x^3 + 14ax^2 + a^5x$	$\{1,10\}$	4212
$x^9 + 2x^7 + x$	-	351
$x^9 + 2x^7 + ax^3$	$\{4,17\}$	702
$x^9 + 2x^7 + 2x^4 + 11x^3 + x^2 + 14x$	-	2106
$x^9 + 2x^7 + 10a^5x^3 + 17ax^2 + 11ax$	$\{1,5\}$	2808
TOTAL		29550

Table 13: nPP Equivalence Classes for $q = 27$, degree 9, and primitive polynomial $x^3 + 2x + 1$.

class	a	number of nPPs
x^8	-	1
$x^8 + x^2 + ax$	$\{1,12,16\}$	341
$x^8 + x^4 + ax$	$\{1,2,8\}$	341
$x^8 + x^4 + x^2 + ax$	$\{0,2,6\}$	341
$x^8 + x^4 + 2x^2 + ax$	$\{0,3,6,7,8,13,14,16,18,19,24,25,31\}$	2015
$x^8 + x^4 + 4x^2 + ax$	$\{0,13,15,20,28,29,30\}$	1085
$x^8 + x^4 + 6x^2 + ax$	$\{4,7,11,12,15,20,23,24,28,29,30\}$	2015
$x^8 + x^4 + 8x^2 + ax$	$\{1,2,5,9,12,13,19,25,28\}$	1395
$x^8 + x^4 + 12x^2 + ax$	$\{3,4,6,11,20,29,30\}$	1085
$x^8 + x^4 + 16x^2 + ax$	$\{1,5,10,12,15,17,18,30\}$	1240
$x^8 + x^6 + 4x^5 + x^4 + 23x^3 + 2x^2 + 23x$	-	4960
$x^8 + x^6 + 6x^5 + 7x^3 + 2x^2$	-	4960
TOTAL		19624

Table 14: nPP Equivalence Classes for $q = 32$, degree 8, and primitive polynomial $x^5+x^3+x^2+x+1$.

class	number of nPPs
x^9	1
$x^9 + x^3 + x$	31
$x^9 + x^6 + x^2$	31
$x^9 + x^6 + x^3$	31
$x^9 + x^7 + x$	31
$x^9 + x^7 + x^6 + x^4 + x^3$	31
$x^9 + x^7 + 16x^6 + 22x^5 + 16x^4 + 30x^3 + 17x^2 + 5x$	155
TOTAL	311

Table 15: nPP Equivalence Classes for $q = 32$, degree 9, and primitive polynomial $x^5+x^3+x^2+x+1$.

class	number of nPPs
x^7	1
$x^7 + x^5 + 26x^3 + 5x$	18
$x^7 + 2x^5 + 28x^3 + 8x$	18
TOTAL	37

Table 16: nPP Equivalence Classes for $q = 37$, degree 7, and primitive polynomial $x + 13$.

class	a	number of nPPs
x^9	-	1
$x^9 + 2ax^5 + 3a^2x$	$\{1,2,3\}$	30
$x^9 + x^6 + 6x^3$	-	40
$x^9 + x^7 + 4a^9x^5 + 36a^{19}x^3 + 33a^{-1}x$	$\{1,2\}$	40
$x^9 + x^7 + 14a^{25}x^5 + 28a^{-2}x^3 + 5a^4x$	$\{1,2\}$	40
$x^9 + ax^7 + 19a^{-8}x^6 + 5a^{11}x^5 + 3a^{-7}x^4 +$ $+16a^{14}x^3 + 15a^7x^2 + 39a^{-10}x$	$\{1,2\}$	80
$x^9 + 2x^7 + 15a^{16}x^5 + 2a^{-3}x^3 + 31a^{22}x$	$\{1,2\}$	40
$x^9 + 2x^7 + 33ax^5 + 18a^{22}x^3 + 15a^{22}x$	$\{1,2\}$	40
$x^9 + 2x^7 + 39x^5 + 25x^3 + 35x$	-	20
TOTAL		331

Table 17: nPP Equivalence Classes for $q = 41$, degree 9, and primitive polynomial $x + 17$.

class	number of nPPs
$x^9 + 2x^7 + 28x^5 + 25x^3 + 2x$	21
$x^9 + 2x^7 + 39x^5 + 33x^3 + 30x$	21
TOTAL	42

Table 18: nPP Equivalence Classes for $q = 43$, degree 9, and primitive polynomial $x + 13$.

class	number of nPPs
$x^{10} + x^7 + 28x^4 + 42x$	14
$x^{10} + x^8 + 19x^7 + x^6 + 33x^5 + 29x^4 + 13x^3 + x^2 + 27x$	42
$x^{10} + 2x^8 + 12x^7 + 32x^6 + 32x^5 + 10x^4 + 39x^3 + 15x^2 + 10x$	42
TOTAL	98

Table 19: nPP Equivalence Classes for $q = 43$, degree 10, and primitive polynomial $x + 13$.

class	number of nPPs
x^9	1
$x^9 + x^6 + 45x^3$	46
$x^9 + x^6 + 13x^5 + 46x^4 + 44x^3 + 39x^2 + 19x$	46
$x^9 + 2x^7 + 13x^5 + 32x^3 + 22x$	23
TOTAL	116

Table 20: nPP Equivalence Classes for $q = 47$, degree 9, and primitive polynomial $x + 12$.

class	number of nPPs
$x^9 + 3x^5 + 10x$	24
$x^9 + 2x^7 + 3x^5 + 4x^3$	24
$x^9 + 2x^7 + 21x^5 + 26x^3 + 36x$	48
TOTAL	96

Table 21: nPP Equivalence Classes for $q = 49$, degree 9, and primitive polynomial $x^2 + 6x + 3$.

class	number of nPPs
$x^{10} + x^7 + 9x$	16
TOTAL	16

Table 22: nPP Equivalence Classes for $q = 49$, degree 10, and primitive polynomial $x^2 + 6x + 3$.

class	number of nPPs
x^9	1
$x^9 + x^6 + 3x^3$	70
TOTAL	71

Table 23: nPP Equivalence Classes for $q = 71$, degree 9, and primitive polynomial $x + 38$.

class	number of nPPs
x^7	1
$x^7 + x^5 + 44x^3 + 31x$	36
$x^7 + 2x^5 + 46x^3 + 34x$	36
TOTAL	73

Table 24: nPP Equivalence Classes for $q = 73$, degree 7, and primitive polynomial $x + 20$.

class	number of nPPs
x^7	1
$x^7 + x^5 + 72x^3 + 75x$	39
$x^7 + 2x^5 + 74x^3 + 78x$	39
TOTAL	79

Table 25: nPP Equivalence Classes for $q = 79$, degree 7, and primitive polynomial $x + 25$.

class	a	number of nPPs
$x^9 + ax$	$\{0, 2, 3, 5, 6\}$	71
$x^9 + ax^3 + ax$	$\{2, 3, 5, 8, 12, 17, 23, 24, 26, 27, 41, 45, 51\}$	1880
$x^9 + 2x^3 + ax$	$\{0, 3, 5, 6, 7, 9, 10, 12, 14, 19, 29, 30, 33, 39\}$	1760
$x^9 + x^2ax^5 + 3a^2x$	$\{1, 2\}$	4860
$x^9 + x^6 + 8ax^5 + x^3 + 48ax^2 + 15a^2x$	$\{1, 2\}$	38880
$x^9 + x^6 + 15ax^5 + x^3 + 55ax^2 + 29a^2x$	$\{1, 4\}$	51840
$x^9 + 2x^7 + 7a^2x^4 + 4x^3 + 48a^2x^2 + 67a^{-3}x$	$\{1, 2\}$	51840
$x^9 + x^6 + 24ax^5 + x^3 + 64ax^2 + 47a^2x$	$\{1, 3\}$	51840
$x^9 + x^6 + 42ax^5 + x^3 + 2ax^2 + 3a^2x$	$\{1, 4\}$	51840
$x^9 + 2x^7 + 4x^3 + 5x$		3240
$x^9 + 2x^7 + ax^4 + 4x^3 + 42ax^2 + 11a^{-5}x$	$\{1, 2\}$	51840
$x^9 + 2x^7 + 4a^2x^4 + 4x^3 + 45a^2x^2 + 33a^{-17}x$	$\{1, 2\}$	51840
$x^9 + 2x^7 + 10a^2x^4 + 4x^3 + 51a^2x^2 + 57a^{-3}x$	$\{1, 2\}$	51840
$x^9 + 2x^7 + 15a^3x^4 + 4x^3 + 56a^3x^2 + 21a^7x$	$\{1, 3\}$	51840
$x^9 + 2x^7 + 3x^5 + 4x^3 + 5x$		3240
$x^9 + 2x^7 + 43x^5 + 5x$		3240
TOTAL		471891

Table 26: nPP Equivalence Classes for $q = 81$, degree 9, and primitive polynomial $x^4 + 2x^3 + 2$.

class	number of nPPs
x^7	1
$x^7 + x^5 + 2x^3 + 59x$	44
$x^7 + 2x^5 + 4x^3 + 62x$	44
TOTAL	89

Table 27: nPP Equivalence Classes for $q = 89$, degree 7, and primitive polynomial $x + 76$.

q	d	class	primitive polynomial	number of nPPs
32	7	x^7	$x^5 + x^3 + x^2 + x$	1
		$x^7 + x^5 + x$		31
41	7	x^7	$x + 17$	1
53	7	x^7	$x + 51$	1
		$x^7 + x^5 + 40x^3 + 25x$		26
		$x^7 + 2x^5 + 42x^3 + 28x$		26
59	7	x^7	$x + 5$	1
		$x^7 + x^5 + 32x^3 + 53x$		29
		$x^7 + 2x^5 + 34x^3 + 56x$		29
	9	x^9	$x + 5$	1
		$x^9 + x^6 + 41x^3$		59
		$x^9 + x^7 + 45x^5 + 22x^3 + 21x$		29
		$x^9 + x^7 + 29x^5 + 4x^3 + 49x$		29
61	7	x^7	$x + 10$	1
		$x^7 + x^5 + 25x^3 + 15x$		30
		$x^7 + 2x^5 + 27x^3 + 18x$		30
67	7	x^7	$x + 17$	1
		$x^7 + x^5 + 45x^3 + 57x$		33
		$x^7 + 2x^5 + 47x^3 + 60x$		33
83	9	x^9	$x + 61$	1
		$x^9 + x^6 + 67x^3$		82
89	9	x^9	$x + 76$	1
		$x^9 + x^6 + 66x^3$		88
97	7	x^7	$x + 90$	1

Table 28: Number of nPPs in Equivalence Classes for prime powers q and degree d , for given primitive polynomial.