

R-TRIVIALITY OF GROUPS OF TYPE F_4 ARISING FROM THE FIRST TITS CONSTRUCTION

SEIDON ALSAODY, VLADIMIR CHERNOUSOV, AND ARTURO PIANZOLA

ABSTRACT. Any group of type F_4 is obtained as the automorphism group of an Albert algebra. We prove that such a group is R -trivial whenever the Albert algebra is obtained from the first Tits construction. Our proof uses cohomological techniques and the corresponding result on the structure group of such Albert algebras.

1. INTRODUCTION

The notion of path connectedness for topological spaces is natural and well-known. In algebraic geometry, this notion corresponds to that of R -triviality. Recall that the notion of R -equivalence, introduced by Manin in [9], is an important birational invariant of an algebraic variety defined over an arbitrary field and is defined as follows: if $\mathbf{X}$ is a variety over a field K with $\mathbf{X}(K) \neq \emptyset$, two K -points $x, y \in \mathbf{X}(K)$ are called *elementarily R -equivalent* if there is a path from x to y , i.e. if there exists a rational map $f : \mathbb{A}^1 \dashrightarrow \mathbf{X}$ defined at $0, 1$ and mapping 0 to x and 1 to y . This generates an equivalence relation R on $\mathbf{X}(K)$ and one can consider the set $\mathbf{X}(K)/R$.

If $\mathbf{X}$ has in addition a group structure, i.e. $\mathbf{X} = \mathbf{G}$ is an algebraic group over a field K , then the set $\mathbf{G}(K)/R$ has a natural structure of an (abstract) group and this gives rise to a functor

$$\mathbf{G}/R : Fields/K \longrightarrow Groups,$$

where $Fields/K$ is the category of field extensions of K , and $\mathbf{G}/R$ is given by $L/K \rightarrow \mathbf{G}(L)/R$. One says that $\mathbf{G}$ is R -trivial if $\mathbf{G}(L)/R = 1$ for all field extensions L/K .

The computation of the group $\mathbf{G}(K)/R$ is known only in a few cases. Here are two examples.

Example 1: Let A be a central simple algebra over a field K and $\mathbf{G} = \mathbf{GL}(1, A)$ the group of “invertible elements of A ”. Denote by $\mathbf{H}$ the kernel $\mathbf{SL}(1, A)$ of the reduced norm homomorphism $\text{Nrd} : \mathbf{G} \rightarrow \mathbf{G}_m$. Then by [18] the group $\mathbf{G}(K)/R$ equals $A^\times/[A^\times, A^\times]$, which is naturally isomorphic to the Whitehead group $K_1(A)$ in algebraic K-theory (cf., [11, § 3]). The group $\mathbf{H}(K)/R$ is known

S. Alsaody wishes to thank PIMS for their financial support.

V. Chernousov was partially supported by Alexander von Humboldt Foundation and an NSERC research grant.

A. Pianzola wishes to thank NSERC and CONICET for their continuous support.

as the reduced Whitehead group $SK_1(A)$. The groups $K_1(A)$ and $SK_1(A)$ have been extensively studied in the framework of algebraic K -Theory.

Example 2 (cf. [4]): let f be a quadratic form over a field F of characteristic not 2 and let $\mathbf{G} = \mathbf{Spin}(f)$. Consider the complex

$$\cdots \longrightarrow \coprod_{x \in \mathbf{X}_{(1)}} K_2 F(x) \xrightarrow{\partial} \coprod_{x \in \mathbf{X}_{(0)}} K_1 F(x) \xrightarrow{N} K_1 F = F^\times$$

where $\mathbf{X}$ is the projective quadric corresponding to f . Let $A_0(\mathbf{X}, K_1)$ be the last homology group of this complex. Then $\mathbf{G}(F)/R \simeq A_0(\mathbf{X}, K_1)$. This result allows one to use the machinery of algebraic K -theory when dealing with groups of R -equivalence classes and is useful in both directions. For instance, if f is a Pfister form, then $\mathbf{G}$ is a rational variety, hence $\mathbf{G}(F)/R = 1$ and therefore $A_0(\mathbf{X}, K_1) = 0$. This triviality result is originally due to M. Rost and was used in the proof of the Milnor conjecture. Conversely, if f is a generic quadratic form over the field $F = k(t_1, \dots, t_m)$, where $t_1, \dots, t_m$ are independent variables over the field k , then using techniques of Chow groups one shows that $A_0(\mathbf{X}, K_1) = 0$, implying $\mathbf{G}(F)/R = 1$.

These two examples show that the group $\mathbf{G}(K)/R$ encodes deep arithmetic properties of $\mathbf{G}$, and that its computation is of great importance in the theory of algebraic groups. It is in general a difficult question to determine whether $\mathbf{G}$ is R -trivial. For algebraic groups, this is the case if $\mathbf{G}$ is *rational*, i.e. birationally equivalent to affine space (cf. [5]). Also, it is known that semisimple groups of rank at most 2 are rational, hence R -trivial (cf. [18]).

The group $\mathbf{G}(K)/R$ for algebraic tori was computed in [5] and for adjoint classical semisimple groups in [10]. The latter computation enables one to construct first examples of non-rational adjoint groups. For simply connected semisimple groups the computation of the group of R -equivalence classes is known only for type A_n (cf. [3]), for spinor groups $\mathbf{Spin}(f)$ (see the above Example 2) and in some easy cases when this $\mathbf{G}/R$ is trivial because the corresponding algebraic group is rational (types C_n and G_2).

The case of exceptional groups is widely open. Beyond G_2 , which is rational by the above, not much is known. In particular it is not known whether groups of type F_4 are R -trivial. These groups arise as automorphism groups of Albert algebras (i.e. simple exceptional Jordan algebras, defined below). Recall that any Albert algebra can be obtained through the first or the second Tits construction. The following theorem is the main result of our paper.

Theorem 1.1. *Any algebraic group of type F_4 over a field K of characteristic not 2 or 3 arising from the first Tits construction is R -trivial.*

We conclude this introduction by mentioning three standard conjectures about the functor $\mathbf{G}/R$: (i) it takes values in the category of abelian groups, (ii) $\mathbf{G}(K)/R$ is a finite group if K is finitely generated over its prime subfield, and (iii) the functor $\mathbf{G}/R$ has transfers.

If $\mathbf{G}$ is an algebraic torus, then the first and third conjectures trivially hold and the second was proved by J.-L. Colliot-Thélène and J.-J. Sansuc [5]. For semisimple groups not much is known. Our main result provides an indication that all three conjectures are true for reductive algebraic groups.

Acknowledgements: We wish to thank Holger Petersson for his answers to our questions.

2. PRELIMINARIES ON ALGEBRAIC GROUPS


In this section we collect some facts about algebraic groups for later use. We start with an observation on subgroups of type D_4 inside the split group of type F_4 .

Proposition 2.1. *Let $\mathbf{H}$ be a split group of type F_4 defined over an arbitrary field K . Then any K -subgroup $\mathbf{M} \subset \mathbf{H}$ of type D_4 is quasi-split.*


Proof. Consider a Borel K -subgroup $\mathbf{B}$ of $\mathbf{H}$. Since the dimension of $\mathbf{B}$ is 28, it intersects the 28-dimensional subgroup $\mathbf{M}$ non-trivially. Thus $\mathbf{M}$ contains a split unipotent subgroup or a split torus $\mathbf{G}_m$. In both cases, $\mathbf{M}$ is K -isotropic.

Let $\mathbf{S} \subset \mathbf{M}$ be a maximal K -split torus. If $\dim(\mathbf{S}) \geq 3$, then $\mathbf{M}$ is necessarily quasi-split. It remains to check the cases where the dimension of $\mathbf{S}$ is 1 or 2. Note that if $\mathbf{M}$ is a triality group of type ${}^{3,6}D_4$ and if $\dim(\mathbf{S}) = 2$ then $\mathbf{M}$ is quasi-split. The remaining cases correspond to the following Tits K -indices (cf. [17]).


2.1. Type 1D_4 .



2.2. Type 2D_4 .



2.3. Type ${}^{3,6}D_4$.

(7) 

We will show that neither case can occur.

Case (1): Here $\dim(\mathbf{S}) = 1$. The semisimple anisotropic kernel $\mathbf{M}_{ss}$ of $\mathbf{M}$ is a group of type 1A_3 , and is thus of dimension 15. From the theory of algebraic groups, the centralizer $C_{\mathbf{H}}(\mathbf{S})$ in $\mathbf{H}$ is a K -split¹ reductive group of rank 4 and by construction contains $\mathbf{M}_{ss}$. Hence the semisimple part

$$C_{\mathbf{H}}(\mathbf{S})_{ss} = [C_{\mathbf{H}}(\mathbf{S}), C_{\mathbf{H}}(\mathbf{S})]$$

of $C_{\mathbf{H}}(\mathbf{S})$ is a semisimple K -split group of rank 3 containing $\mathbf{M}_{ss}$. Since $C_{\mathbf{H}}(\mathbf{S})_{ss}$ is split, $C_{\mathbf{H}}(\mathbf{S})_{ss} \neq \mathbf{M}_{ss}$. Therefore the type of $C_{\mathbf{H}}(\mathbf{S})_{ss}$ is B_3 or C_3 , and so its dimension is 21. In both cases, any Borel K -subgroup of $C_{\mathbf{H}}(\mathbf{S})_{ss}$ has dimension 12, and thus for dimension reasons intersects $\mathbf{M}_{ss}$ nontrivially, contradicting the assumption that $\mathbf{M}_{ss}$ is K -anisotropic.

Case (2): Here $\dim(\mathbf{S}) = 2$, and the semisimple anisotropic kernel $\mathbf{M}_{ss}$ of $\mathbf{M}$ is a group of type $A_1 \oplus A_1$, hence of dimension 6. Again, we consider the centraliser $C_{\mathbf{H}}(\mathbf{S})$; its semisimple part $C_{\mathbf{H}}(\mathbf{S})_{ss} = [C_{\mathbf{H}}(\mathbf{S}), C_{\mathbf{H}}(\mathbf{S})]$ is a K -split semisimple group of rank 2 containing $\mathbf{M}_{ss}$. Therefore it is either of type A_2 or $B_2 = C_2$. A dimension count shows that the intersection of a Borel K -subgroup of $C_{\mathbf{H}}(\mathbf{S})_{ss}$ and $\mathbf{M}_{ss}$ is nontrivial, contradicting anisotropy.

Case (3): Here, up to isogeny, $\mathbf{M}$ is of the form $\mathbf{O}^+(D, f)$ where D is a quaternion algebra and f is a skew-hermitian form over D of dimension 4 with Witt index 1. It is known that after passing to the function field of the Severi–Brauer variety of D , the anisotropic part of f remains anisotropic, hence the Tits index of $\mathbf{M}$ is of the form (2). But this case is impossible by the above consideration.

Cases (4), (5) and (6) are treated in the same way as cases (1), (2) and (3) respectively, since the dimensions of the corresponding subgroups are the same. In case (7), the uncircled vertices correspond to the Weil restriction $R_{L/K}(\mathrm{SL}(1, D))$ for a cubic field extension L of K and a quaternion L -algebra D . Extending scalars to L reduces this case to one of the previously treated cases (3) and (6). This completes the proof. $\square$

2.4. The norm principle. Let $\mathbf{G}$ be a semisimple group defined over a field K and $\mathbf{Z} \subset \mathbf{G}$ a central subgroup. Let moreover $c = [\xi]$ be an element in

$$(8) \quad \mathrm{Ker} [H^1(K, \mathbf{Z}) \longrightarrow H^1(K, \mathbf{G})].$$

Definition. We say that $[\xi]$ is *R-trivial* if there exists

$$[\xi(t)] \in \mathrm{Ker} [H^1(K(t), \mathbf{Z}) \longrightarrow H^1(K(t), \mathbf{G})],$$

¹It is K -split because every split subtorus in $\mathbf{H}$, in particular $\mathbf{S}$, is contained in a maximal split torus of $\mathbf{H}$.

where $K(t)$ is a purely transcendental extension of K , such that $\xi(t)$ is defined at $t = 0$ and $t = 1$, and satisfies $[\xi(0)] = 1$ and $[\xi(1)] = [\xi]$.

Remark 2.2. Here, if $\mathbf{J}$ is an algebraic group over K , we say that a cocycle in $Z^1(K(t), \mathbf{J})$ is *defined at 0 and 1* if it lies in the image of $Z^1(\mathcal{O}, \mathbf{J})$, where $\mathcal{O}$ is the intersection of the localizations $K[t]_{(t)}$ and $K[t]_{(t-1)}$ in $K(t)$. Using the evaluation maps $\varepsilon_0, \varepsilon_1 : \mathcal{O} \rightarrow K$, such a cocycle can be evaluated at 0 and 1.

Remark 2.3. The short exact sequence

$$1 \longrightarrow \mathbf{Z} \longrightarrow \mathbf{G} \longrightarrow \mathbf{G}/\mathbf{Z} \longrightarrow 1$$

induces an exact sequence

$$1 \longrightarrow \mathbf{Z}(K) \longrightarrow \mathbf{G}(K) \longrightarrow (\mathbf{G}/\mathbf{Z})(K) \xrightarrow{f} H^1(K, \mathbf{Z}) \longrightarrow H^1(K, \mathbf{G})$$

and the map f induces a bijection between the orbits of $\mathbf{G}(K)$ in $(\mathbf{G}/\mathbf{Z})(K)$ and the kernel in (8).

Example 2.4. Let $\mathbf{G}$ be a quasi-split (absolutely) simple simply connected group. Then

$$H^1(K, \mathbf{Z}) = \text{Ker} [H^1(K, \mathbf{Z}) \rightarrow H^1(K, \mathbf{G})]$$

and every $[\xi] \in H^1(K, \mathbf{Z})$ is R -trivial. Indeed, $\mathbf{Z}$ is contained in a maximal K -quasi-split torus T and $H^1(F, T) = 1$ for any field extension F/K . It remains to note that the variety $\mathbf{G}/\mathbf{Z}$ is K -rational, which implies that all elements in $H^1(K, \mathbf{Z})$ are R -trivial.

Example 2.5. Let $\mathbf{G} = \mathbf{SL}(1, D)$ where D is a central simple algebra of arbitrary degree n over K , and let $\mathbf{Z}$ be the centre of $\mathbf{G}$. Since $\mathbf{Z} \simeq \mu_n$, we have $H^1(K, \mathbf{Z}) \simeq K^\times / K^{\times n}$. Also, $H^1(K, \mathbf{G}) \simeq K^\times / \text{Nrd}(D^\times)$. Therefore,

$$\text{Ker} [H^1(K, \mathbf{Z}) \longrightarrow H^1(K, \mathbf{G})] = \text{Nrd}(D^\times) / K^{\times n}.$$

Since D is an affine space, any $[\xi]$ in the above kernel is R -trivial.

For any finite extension L/K we have the restriction map

$$\text{res}_K^L : H^1(K, \mathbf{Z}) \rightarrow H^1(L, \mathbf{Z})$$

and the corestriction map

$$\text{cor}_K^L : H^1(L, \mathbf{Z}) \rightarrow H^1(K, \mathbf{Z}).$$

Definition. Let L/K be a finite field extension. We say that *the norm principle holds for a cohomology class*

$$[\eta] \in \text{Ker} [H^1(L, \mathbf{Z}) \longrightarrow H^1(L, \mathbf{G})]$$

if

$$\text{cor}_K^L([\eta]) \in \text{Ker} [H^1(K, \mathbf{Z}) \longrightarrow H^1(K, \mathbf{G})].$$

We also say that the norm principle holds for the pair $(\mathbf{Z}, \mathbf{G})$ if it holds for all classes

$$[\eta] \in \text{Ker} [H^1(L, \mathbf{Z}) \longrightarrow H^1(L, \mathbf{G})]$$

for each finite field extension L/K .

Theorem 2.6. (P. Gille [6]) *Let L/K be a finite field extension and assume that $[\eta] \in \text{Ker}[H^1(L, \mathbf{Z}) \rightarrow H^1(L, \mathbf{G})]$ is R -trivial. Then the norm principle holds for $[\eta]$.*

3. PRELIMINARIES ON ALBERT ALGEBRAS

3.1. Albert Algebras. Let K be a field of characteristic different from 2 and 3. A *Jordan algebra* over K is a unital, commutative K -algebra² A in which the Jordan identity

$$(xy)(xx) = x(y(xx))$$

holds for all $x, y \in A$ (thus A is power associative).

If B is an associative algebra with multiplication denoted by $\cdot$, then the anticommutator $\frac{1}{2}(x \cdot y + y \cdot x)$ endows B with a Jordan algebra structure, which we denote by B^+ . A Jordan algebra A is called *special* if it is isomorphic to a Jordan subalgebra of B^+ for some associative algebra B , and *exceptional* otherwise. An *Albert algebra* is then defined as a simple, exceptional Jordan algebra. It is known that the dimension of any Albert algebra is 27, and that over separably closed fields, all Albert algebras are isomorphic [8, 37.11]. From this it follows that all Albert algebras over K are twisted forms of each other.

Over a general field K , any Albert algebra arises from one of the two *Tits constructions*. In this note, we will mainly be concerned with those arising from the first Tits construction. Recall that in this case, $A = D \oplus D \oplus D$ as a vector space, where D is a central simple algebra over K of degree 3. The multiplication of A is determined by a scalar $\mu \in K^*$. More precisely, define the cross product on D by

$$u \times v = uv + vu - T_D(u)v - T_D(v)u - T_D(uv) + T_D(u)T_D(v),$$

where juxtaposition denotes the associative multiplication of D , and T_D denotes the reduced trace, and write

$$\tilde{u} = T_D(u) - u$$

for $u \in D$. For any $\mu \in K^\times$, the product $(x, y, z)(x', y', z')$ in $A = D \oplus D \oplus D$ is then given by

$$\frac{1}{2}(xx' + x'x + \tilde{y}z' + \tilde{y}'z, \tilde{x}y' + \tilde{x}'y + \mu^{-1}z \times z', z\tilde{x}' + z'\tilde{x} + \mu y \times y').$$

One then writes $A = J(D, \mu)$ and says that A arises from D and μ via the first Tits construction.

In some arguments, we will also need to consider reduced Albert algebras, and we briefly recall their construction. Given an octonion algebra C over K

²We do not assume the algebra to be associative.

and $\Gamma = (\gamma_1, \gamma_2, \gamma_3) \in (K^\times)^3$, the set $H_3(C, \Gamma)$ of Γ -hermitian 3×3 -matrices over C , i.e. those of the form

$$\begin{pmatrix} \xi_1 & c_3 & \gamma_1^{-1} \gamma_3 \overline{c_2} \\ \gamma_2^{-1} \gamma_1 \overline{c_3} & \xi_2 & c_1 \\ c_2 & \gamma_3^{-1} \gamma_2 \overline{c_1} & \xi_3 \end{pmatrix},$$

with $\xi_i \in K$ and $c_i \in C$ for $i = 1, 2, 3$, and where $\overline{}$ denotes the involution on C , is an Albert algebra under the anticommutator of the matrix product. From [13] we know that an Albert algebra over K is a division algebra if and only if it is not of this form.

3.2. Automorphisms and Similarities. If A is an Albert algebra over K , then $\mathbf{H} = \mathbf{Aut}(A)$ is a simple algebraic group over K of type F_4 . In fact, the assignment $A \mapsto \mathbf{Aut}(A)$ establishes an equivalence between the category of Albert algebras over K and the category of simple K -groups of type F_4 . Moreover, A is endowed with a cubic form $N : A \rightarrow K$ known as the *norm* of A . The *structure group* $\mathbf{Str}(A)$ of A is the affine group scheme whose R -points are defined by

$$\mathbf{Str}(A)(R) = \{x \in \mathbf{GL}(A)(R) \mid x \text{ is an isotopy of } A \otimes R, \}$$

for every K -ring R . By an *isotopy* of an Albert algebra B over a ring R one understands an isomorphism of Jordan algebras from B to the isotope $B^{(p)}$ for some $p \in B^\times$ with $N(p) \in R^\times$, the algebra $B^{(p)}$ having the same underlying module as B , with multiplication

$$x \cdot_p y = x(y p) + y(x p) - (x y) p.$$

It is clear that every isotopy is a norm similarity of A . This holds for Albert algebras over commutative rings (see [12] for a discussion of such algebras) and implies that the structure group is a subgroup of the group of norm similarities. By [7, VI Theorem 7], if F is a field with more than 3 elements, then every norm similarity is an isotopy. Since these groups are smooth, it follows that they coincide, i.e. for any K -ring R ,

$$\mathbf{Str}(A)(R) = \{x \in \mathbf{GL}(A)(R) \mid N_R(x(a)) = \nu(x) N_R(a) \ \forall a \in A \otimes R \},$$

where N_R is the scalar extension of N to $A \otimes R$, and $\nu(x)$ is a scalar in $R^\times$ (called a multiplier) depending on x only. Note that $\mathbf{Str}(A)$ contains a (central) split torus $\mathbf{G}_m$ consisting of all homotheties of A .

It is well known that the derived subgroup

$$\mathbf{G} = [\mathbf{Str}(A), \mathbf{Str}(A)]$$

of $\mathbf{Str}(A)$ is a strongly inner form of a split simple simply connected algebraic group of type E_6 and that $\mathbf{Str}(A)$ is an almost direct product of $\mathbf{G}_m$ and $\mathbf{G}$ (their intersection is the centre of $\mathbf{G}$). Thus,

$$\overline{\mathbf{G}} = \mathbf{Str}(A)/\mathbf{G}_m$$

is an adjoint group of type E_6 . Since our main result is known for split or isotropic $\mathbf{G}$ we may assume without loss of generality that $\mathbf{G}$ is K -anisotropic

which is equivalent to saying that A is a division algebra or equivalently the norm map N is anisotropic, i.e. the equation $N(a) = 0$ has no non-zero solutions over K .

3.3. Action of the structure group $\mathbf{Str}(A)$ on A . We denote the group of K -points of $\mathbf{Str}(A)$ (resp. $\mathbf{G}, \overline{\mathbf{G}}, \mathbf{H}$) by $\mathrm{Str}(A)$ (resp. $G, \overline{G}, H$). Under the natural action of $\mathbf{Str}(A)$ on A , the group $\mathbf{H}$ coincides with the stabilizer of $1 \in A$, as follows from [14, 5.9.4]. This action of $\mathbf{Str}(A)$ on A induces an action on the open subset U defined by

$$U(R) = \{a \in A \otimes_K R \mid N_R(a) \in R^\times\}.$$

for any K -ring R .

Lemma 3.1. *Let A be split over K . Then the action $\mathrm{Str}(A) \times U \rightarrow U$ is transitive.*

It is known that the action is transitive over a separable closure of K . Therefore, to prove the lemma, it suffices to show that the natural map $H^1(K, \mathbf{H}) \rightarrow H^1(K, \mathbf{Str}(A))$ has trivial kernel or equivalently

$$\mathrm{Ker} [H^1(K, \mathbf{H}) \longrightarrow H^1(K, \mathbf{G})] = 1.$$

We will give two proofs of this statement, one using the structure theory of algebraic groups, and the other more Jordan theoretic in nature, to illuminate both sides of the statement.

First proof. Let $[\xi]$ be in the kernel. Consider the twisted group ${}^\xi \mathbf{H}$. It is a K -subgroup of the split group ${}^\xi \mathbf{G} \simeq \mathbf{G}$. Let $\mathbf{B} \subset \mathbf{G}$ be a Borel K -subgroup. A dimension count shows that ${}^\xi \mathbf{H} \cap \mathbf{B}$ is nontrivial. Hence ${}^\xi \mathbf{H}$ is isotropic, which implies that the invariants g_3 and f_5 of ${}^\xi \mathbf{H}$ are both trivial. Thus, up to equivalence in $H^1(K, \mathbf{H})$, we may assume that ξ takes values in a split subgroup of $\mathbf{H}$ of type G_2 which is in turn a subgroup of the split group $\mathbf{M}$ of type D_4 generated by the roots $\alpha_2, \dots, \alpha_5$. Since $[\xi]$ is trivial in $H^1(K, \mathbf{G})$, it follows from the Tits classification of isotropic groups of type ${}^1 E_6$ that it is trivial in $H^1(K, \mathbf{M})$, hence in $H^1(K, \mathbf{H})$. $\square$

Second proof. By [1, Theorem 2.5], this kernel parametrises the isomorphism classes of isotopes of A . But from [12, Corollary 60], it follows that the invariants mod 2 and 3 of any isotope of the split Albert algebra A are trivial. Thus all such isotopes are isomorphic, and the above kernel is trivial, as desired. $\square$

3.4. Subgroups and a cohomology class attached to a point in A . Let $a \in A$. We associate to it a subgroup of type D_4 in $\mathbf{G}$ and a 2-dimensional torus as follows. Let $L \subset A$ be a subalgebra generated by a . It is known that L is a subfield since A is a division algebra. Define the group $\mathbf{G}^L$ by

$$\mathbf{G}^L(R) = \{x \in \mathbf{G}(R) \mid x(l) = l \ \forall l \in L \otimes_K R\}$$

for any K -ring R . Since $\mathbf{G}^L$ stabilizes $1 \in L \subset A$ one gets $\mathbf{G}^L \subset \mathbf{H} \subset \mathbf{G}$. It is known that over a separable closure of K the group $\mathbf{G}^L$ is conjugate to the standard subgroup in $\mathbf{G}$ of type D_4 generated by roots $\alpha_2, \alpha_3, \alpha_4, \alpha_5$. Therefore

$\mathbf{S}'^L = C_{\mathbf{G}}(\mathbf{G}^L) \subset \mathbf{G}$ is a 2-dimensional torus over K . Using an explicit model of A over a separable closure of K one can easily verify that

$$\mathbf{Z}^L := \mathbf{S}'^L \cap \mathbf{H} = \mathbf{S}'^L \cap \mathbf{G}^L$$

is the centre of $\mathbf{G}^L$.

Let $\mathbf{S}^L \subset \mathbf{Str}(A)$ be a 3-dimensional torus generated by $\mathbf{S}'^L$ and $\mathbf{G}_m$. Note that $\mathbf{S}^L \cdot \mathbf{G}^L$ is the connected component of the stabilizer $\mathbf{N}^L = \mathbf{Str}(A, L)$ defined by

$$\mathbf{N}^L(R) = \mathbf{Str}(A, L)(R) = \{x \in \mathbf{Str}(A)(R) \mid x(L \otimes_K R) = L \otimes_K R\}$$

for each K -ring R , and that over the separable closure of K one has $\mathbf{N}^L/(\mathbf{S}^L \cdot \mathbf{G}^L) \simeq S_3$. Since the representatives of this quotient group can be chosen in $\mathbf{H}$, we have a natural identification $\mathbf{N}^L/(\mathbf{S}^L \cdot \mathbf{G}^L) \simeq \mathbf{Aut}(L/K)$.

Our next goal is to describe a structure of $\mathbf{S}^L$. Before doing so, we recall a standard fact about actions of algebraic groups on varieties and the corresponding transporters.

Let $L_1, L_2 \subset A$ be two cubic étale (commutative) subalgebras. Consider the transporter defined by

$$\mathbf{Transp}(L_1, L_2)(R) = \{x \in \mathbf{Str}(A)(R) \mid x(L_1 \otimes_K R) = L_2 \otimes_K R\}.$$

for each K -ring R . It is an $\mathbf{N}^{L_1}$ -torsor, hence it is represented by a cohomology class

$$[\xi] \in \text{Ker}[H^1(K, \mathbf{N}^{L_1}) \longrightarrow H^1(K, \mathbf{Str}(A))].$$

By construction, this class is trivial if and only if L_1 and L_2 are conjugate over K . Furthermore, let $[\bar{\xi}]$ be the image of $[\xi]$ in

$$H^1(K, \mathbf{N}^{L_1}/(\mathbf{S}^{L_1} \cdot \mathbf{G}^{L_1})) = H^1(K, \mathbf{Aut}(L_1/K)).$$

Then we have ${}^{\bar{\xi}}L_1 \simeq L_2$. In particular, if $L_1 \simeq L_2$, then $[\bar{\xi}]$ is trivial, hence up to equivalence in $H^1(K, \mathbf{N}^{L_1})$ we may assume that ξ takes values in $\mathbf{S}^{L_1} \cdot \mathbf{G}^{L_1}$. This observation leads us to the following.

Lemma 3.2. *Let $A = H_3(C, \Gamma)$ be a reduced Albert algebra, where C is an octonion algebra over K and $\Gamma \in (K^\times)^3$. Let $L_1 \subset A$ be the diagonal subalgebra and let $L_2 \subset A$ be an arbitrary split cubic subalgebra. Then L_1 and L_2 are conjugate in $\mathbf{Str}(A)$.*

Proof. By the discussion preceding the lemma, the obstacle for conjugacy is a cohomology class $[\xi] \in H^1(K, \mathbf{S}^{L_1} \cdot \mathbf{G}^{L_1})$ which maps to the trivial class in $H^1(K, \mathbf{Str}(A))$. Since L_1 consists of diagonal matrices in $H_3(C, \Gamma)$, it readily follows that $\mathbf{S}^{L_1}$ is a split K -torus, hence $\mathbf{S}^{L_1} \cdot \mathbf{G}^{L_1}$ is a Levi subgroup of a parabolic subgroup $\mathbf{P} \subset \mathbf{Str}(A)$. It is well known that both arrows

$$H^1(K, \mathbf{S}^{L_1} \cdot \mathbf{G}^{L_1}) \longrightarrow H^1(K, \mathbf{P}) \longrightarrow H^1(K, \mathbf{Str}(A))$$

have trivial kernel. It follows that $[\xi] = 1$, as desired. $\square$

We come back to an arbitrary Albert algebra A over K , and a cubic subfield $K \subset L \subset A$.

Proposition 3.3. *One has $\mathbf{S}^L \simeq R_{L/K}(\mathbf{G}_m)$ and $\mathbf{S}'^L \simeq R_{L/K}^{(1)}(\mathbf{G}_m)$.*

Proof. Let F/K (resp. F'/K) be the minimal splitting field of $R_{L/K}^{(1)}(\mathbf{G}_m)$ (resp. $\mathbf{S}'^L$). Let $\Gamma = \text{Gal}(F/K)$ and $\Gamma' = \text{Gal}(F'/K)$. These two Galois groups act naturally on the corresponding character lattices of the tori $R_{L/K}^{(1)}(\mathbf{G}_m)$ and $\mathbf{S}'^L$, respectively, and hence both embed naturally into $\text{GL}_2(\mathbb{Z})$. Recall that an arbitrary n -dimensional torus is determined uniquely by its minimal splitting field and the conjugacy class of the image of the corresponding Galois group in $\text{GL}_n(\mathbb{Z})$.

Step 1: $F' \subset F$. The base extension F/K completely splits L . Hence the g_3 -invariant of A becomes trivial over F , implying that $A_F \simeq H_3(C, \Gamma)$ where C is the octonion algebra over F corresponding to $f_3(A_F)$, and $\Gamma \in (F^\times)^3$. By the above lemma, up to conjugacy we may assume that $L \otimes_K F \subset A_F$ coincides with the diagonal subalgebra. Explicit computations show that $(\mathbf{S}^L)_F$ is a split 3-dimensional torus, and that so is $(\mathbf{S}'^L)_F$.

Step 2: $F \subset F'$. Since the field extension F'/K splits $\mathbf{S}^L$ it follows from Tits' classification of isotropic simple algebraic groups of type ${}^1\text{E}_6$ that the semisimple part $\mathbf{G}^L$ of $C_{\text{Str}(A)}(\mathbf{S}^L)$ is isomorphic to $\mathbf{Spin}(f)$ where f is the 3-fold Pfister form as above, which is either split over F' or splits over a quadratic field extension E/F' . Therefore, the same holds for $\text{Str}(A)$ and for A . In both cases the g_3 -invariant of $A_{F'}$, being an invariant mod 3, is trivial, implying that $A_{F'} \simeq H_3(C, \Gamma)$ for an octonion F' -algebra C and $\Gamma \in (F'^\times)^3$. Since all maximal split tori in $\text{Str}(A_{F'})$ are conjugate over F' , we may assume that, up to conjugacy, the split torus $(\mathbf{S}^L)_{F'}$ preserves the diagonal subalgebra of $H_3(C, \Gamma)$; then so does its centralizer in $\text{Str}(A_{F'})$. Furthermore, explicit computations show that

$$(\mathbf{G}^L)_{F'} = [C_{\text{Str}(A_{F'})}(\mathbf{S}^L)_{F'}, C_{\text{Str}(A_{F'})}(\mathbf{S}'^L)_{F'}]$$

acts trivially on the diagonal. Since L consists of precisely those points of A fixed pointwise under the action of $\mathbf{G}^L$, it follows that $L \otimes_K F' \subset A_{F'}$ coincides with the diagonal. Thus L is split over F' .

Step 3: the images of Γ and Γ' in $\text{GL}_2(\mathbb{Z})$ are conjugate. Recall that up to conjugacy the group $\text{GL}_2(\mathbb{Z})$ has two maximal finite subgroups, isomorphic to the dihedral groups D_4 and $D_6 \simeq D_3 \times \mathbb{Z}/2$. Since the orders of Γ and Γ' are divisible by 3 we may, up to conjugacy, assume that $\Gamma, \Gamma' \subset D_6 \subset \text{GL}_2(\mathbb{Z})$. If $|\Gamma| = |\Gamma'| = 3$, then these groups coincide with the unique 3-Sylow subgroup of D_6 and we are done. Otherwise Γ and Γ' have order 6, hence $\Gamma \simeq \Gamma' \simeq S_3$. The group $D_6 \simeq D_3 \times \mathbb{Z}/2 \simeq S_3 \times \mathbb{Z}/2$ has two subgroups isomorphic to S_3 . They are not conjugate in D_6 , but explicit computations shows that they are conjugate in $\text{GL}_2(\mathbb{Z})$. This completes the proof. $\square$

In the proof we used the fact that the inclusion

$$L \subset \{a \in A \mid g(a) = a \ \forall g \in \mathbf{G}^L(K)\}$$

is an equality. Hence the restriction of the action of $\mathbf{Str}(A)$ (on A) at $\mathbf{S}^L$ defines an action $\mathbf{S}^L \times L^\times \rightarrow L^\times$. Since $\mathbf{S}^L \cap \mathbf{H} = \mathbf{S}'^L \cap \mathbf{H} = \mathbf{Z}^L$, the stabilizer of $1 \in L^\times$ in $\mathbf{S}^L$ is equal to $\mathbf{Z}^L$. It follows that over a separable closure of K the action $\mathbf{S}^L \times L^\times \rightarrow L^\times$ is transitive. Thus we may identify the varieties $\mathbf{S}^L/\mathbf{Z}^L \simeq L^\times$. The exact sequence

$$1 \longrightarrow \mathbf{Z}^L \longrightarrow \mathbf{S}^L \xrightarrow{\phi} L^\times \longrightarrow 1$$

and the fact that $H^1(K, \mathbf{S}^L) = 1$ imply that we have a canonical surjective map $L^\times \rightarrow H^1(K, \mathbf{Z}^L)$. Thus to each point $a \in L^\times$ we can attach its image $[\xi_a] \in H^1(K, \mathbf{Z}^L)$.

Remark 3.4. By construction, the map ϕ is the restriction of the map $\mathbf{Str}(A) \rightarrow U$ sending g to $g(1)$.

4. R-TRIVIALITY

We keep the notation introduced in Section 3; in particular $U \subset A$ is the open subset consisting of the elements a in A whose norm $N(a)$ is invertible.

Proposition 4.1. *Let A be an Albert algebra arising from the first Tits construction. Then the group $\mathbf{Str}(A)$ acts transitively on U .*

Proof. Recall that the stabilizer of $1 \in A$ is $\mathbf{H}$. Thus we may identify the variety $\mathbf{G}/\mathbf{H}$ with U . It follows from the exact sequence

$$1 \longrightarrow \mathbf{H} \longrightarrow \mathbf{Str}(A) \longrightarrow U \longrightarrow 1$$

that there is a canonical one-to-one correspondence between the set of $\mathbf{Str}(A)$ -orbits in U and the set

$$\mathrm{Ker} [H^1(K, \mathbf{H}) \longrightarrow H^1(K, \mathbf{Str}(A))].$$

Therefore our assertion amounts to proving triviality of the above kernel.

Let $a \in U$ and let

$$[\eta] \in \mathrm{Ker} [H^1(K, \mathbf{H}) \longrightarrow H^1(K, \mathbf{Str}(A))]$$

be the corresponding cohomology class. We take a cubic subfield L containing a , construct the class $[\xi_a] \in H^1(K, \mathbf{Z}^L)$ as in the end of the previous section, and write $[\xi]_{\mathbf{H}}$ for its image under the composition

$$H^1(K, \mathbf{Z}^L) \rightarrow H^1(K, \mathbf{G}^L) \rightarrow H^1(K, \mathbf{H}).$$

By Remark 3.4, $[\eta] = [\xi]_{\mathbf{H}}$. The following claim completes the proof of the proposition.

Claim: the map $H^1(K, \mathbf{Z}^L) \rightarrow H^1(K, \mathbf{H})$ is trivial.

Indeed, let $[\xi] \in H^1(K, \mathbf{Z}^L)$. Take its restriction $[\xi]_L \in H^1(L, ((\mathbf{Z}^L)_L))$. Upon extending scalars to L , the g_3 -invariant of $\mathbf{H}$ becomes trivial. Since A is obtained from the first Tits construction, its invariants mod 2 are trivial, and thus the group $\mathbf{H}_L$ is split. Then by Lemma 2.1, the group $\mathbf{G}^L$ is quasi-split over L . Choose a maximal quasi-split torus T_L in $\mathbf{G}^L \otimes_K L$. It contains the

centre $(\mathbf{Z}^L)_L$ of $(\mathbf{G}^L)_L$ and has trivial cohomology in dimension 1. This implies that

$$[\xi]_L \in \text{Ker} [H^1(L, (\mathbf{Z}^L)_L) \longrightarrow H^1(L, (\mathbf{G}^L)_L)].$$

By Example 2.4, every element in $H^1(L, (\mathbf{Z}^L)_L)$ is R -trivial and therefore the norm principle holds for each such element. Thus

$$\text{cor}_K^L(\text{res}_L([\xi])) \in \text{Ker} [H^1(K, \mathbf{Z}^L) \rightarrow H^1(K, \mathbf{G}^L)].$$

The claim follows upon noting that

$$\text{cor}_K^L(\text{res}_L([\xi])) = [\xi]^3 = [\xi],$$

because $H^1(K, \mathbf{Z}^L)$ is a group of exponent 2. $\square$

In the course of the proof of the proposition we established the following.

Corollary 4.2. *Let F/K be an arbitrary field extension.*

- (a) *We have $\text{Ker} [H^1(F, \mathbf{H}) \rightarrow H^1(F, \mathbf{Str}(A))] = 1$.*
- (b) *The variety $\mathbf{H} \times U$ is birationally isomorphic to $\mathbf{Str}(A)$; in particular $\mathbf{H}$ is R -trivial if and only if so is $\mathbf{Str}(A)$.*

Proof. (a) was already established and (b) follows immediately from (a). $\square$

Corollary 4.3. *Let K be a field of characteristic $\neq 2, 3$. Let $\mathbf{H}$ be an arbitrary algebraic group of type F_4 over K arising from the first Tits construction. Then $\mathbf{H}$ is R -trivial.*

Proof. Let A be the corresponding Albert algebra. By [15] and [16], the structure group $\mathbf{Str}(A)$ is R -trivial. Hence the assertion follows from the above corollary. $\square$

Remark 4.4. In a forthcoming paper [2], we will deal with automorphism and structure groups of arbitrary Albert algebras.

REFERENCES

- [1] S. Alsaody, *Albert algebras over rings and related torsors*, arXiv:1901.04459 (2019).
- [2] S. Alsaody, V. Chernousov and A. Pianzola, *On the Tits–Weiss conjecture and the Kneser–Tits conjecture for $E_{7,1}^{78}$ and $E_{8,2}^{78}$* , preprint (2019).
- [3] V. Chernousov and A. Merkurjev, *R-equivalence and special unitary groups*, J. Algebra 209, 175–198 (1998).
- [4] V. Chernousov and A. Merkurjev, *R-equivalence in spinor groups*, J. Amer. Math. Soc., bf 14 (2001), 3, 509–534.
- [5] J.-L. Colliot-Thélène and J.-J. Sansuc, *La R-équivalence sur les tores*, Ann. Sci. École Norm. Sup. (4), 10, 175–229 (1977).
- [6] P. Gille, *La R-équivalence sur les groupes algébriques réductifs définis sur un corps global*, Publications mathématiques de l’I.H.E.S., **86** (1997), 199–235.
- [7] N. Jacobson, *Structure and Representations of Jordan Algebras*, AMS Colloquium Publications, Vol. 39, 1968.
- [8] M.-A. Knus, A. Merkurjev, M. Rost, J.-P. Tignol, *The book of involutions*, AMS Colloquium Publications, Vol. 44, 1998.
- [9] Yu. I. Manin, *Cubic forms*, North-Holland, Amsterdam, 1974.

- [10] A. Merkurjev, *R-equivalence and rationality problem for semisimple adjoint classical algebraic groups*, Publications mathématiques de l'I.H.É.S., **84** (1996), 189–213.
- [11] J. Milnor, *Introduction to algebraic K-theory*, Ann. Math. Stud., N 72, Princeton, 1971.
- [12] H. P Petersson, *A survey on Albert algebras*, Transformation Groups, **24** (2019), 219–278.
- [13] R. D. Schafer, *The exceptional simple Jordan algebras*, Amer. J. Math. **70** (1948), 82–94.
- [14] T. A. Springer and F. D. Veldkamp, *Octonion algebras, Jordan algebras and exceptional groups*, Springer Monographs in Mathematics, Springer-Verlag (2000).
- [15] M. Thakur, *Automorphisms of Albert algebras and a conjecture of Tits and Weiss*, Trans. AMS **365** (2013), 3041–3068.
- [16] M. Thakur, *Automorphisms of Albert algebras and a conjecture of Tits and Weiss II*, Trans. AMS **372** (2019), 4701–4728.
- [17] J. Tits, *Classification of algebraic semisimple groups*, In Algebraic Groups and discontinuous Subgroups, (eds. A. Borel and G.D. Mostow), Proc. Symp. Pure Math. **9** (1966), 33–62.
- [18] V. E. Voskresenskii, *Algebraic tori*, Nauka, Moscow, 1977.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON,
ALBERTA, CANADA T6G 2G1

E-mail address: `alsaody@ualberta.ca`

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON,
ALBERTA, CANADA T6G 2G1

E-mail address: `vladimir@ualberta.ca`

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON,
ALBERTA, CANADA T6G 2G1

E-mail address: `a.pianzola@ualberta.ca`