

# Global bounded solution of the higher-dimensional forager-exploiter model with/without growth sources<sup>1</sup>

Jianping Wang, Mingxin Wang<sup>2</sup>

School of Mathematics, Harbin Institute of Technology, Harbin 150001, PR China

**Abstract.** We investigate the higher-dimensional forager-exploiter model with homogeneous Neumann boundary condition

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla w) + \eta_1(u - u^m), & x \in \Omega, \quad t > 0, \\ v_t = \Delta v - \xi \nabla \cdot (v \nabla u) + \eta_2(v - v^l), & x \in \Omega, \quad t > 0, \\ w_t = \Delta w - \lambda(u + v)w - \mu w + r(x, t), & x \in \Omega, \quad t > 0, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), \quad w(x, 0) = w_0(x), & x \in \Omega, \end{cases}$$

where  $\Omega \subset \mathbb{R}^n$  is a bounded domain, the constants  $\chi, \xi, \lambda, \mu$  are positive,  $m, l > 1$  and  $\eta_1, \eta_2 \geq 0$ ,  $r \in C^1(\bar{\Omega} \times [0, \infty)) \cap L^\infty(\Omega \times (0, \infty))$ . The nonnegative initial functions  $u_0, w_0 \in W^{2,\infty}(\Omega) \setminus \{0\}$  and  $v_0 \in W^{1,\infty}(\Omega) \setminus \{0\}$ . It will be shown that, for  $\eta_1 = \eta_2 = 0$  and  $n \geq 2$ , on the one hand, there is  $\varepsilon > 0$  such that, if

$$\|u_0\|_{W_{2p}^1(\Omega)} + \|w_0\|_{W_{2(p+1)}^1(\Omega)} + \|r\|_{L^\infty(\Omega \times (0, \infty))} \leq \varepsilon \quad \text{with } p = \min \{k \in \mathbb{N} : k > n/2\},$$

then this system is globally solvable in the classical sense; On the other hand, the weak taxis effects rule out any blow-up. Secondly, for  $\eta_1 > 0, m \geq 2, \eta_2 = 0$  and  $n = 2$ , we establish the global solvability of this problem provided only  $\xi$  is sufficiently small. Finally, when  $n = 2, \eta_1, \eta_2 > 0$ , we find a condition for the logistic degradation rates  $m, l$  that ensures the global existence of the classical solutions.

**Keywords:** Forager-exploiter model; super-logistic source; classical solution; Global existence; Boundedness.

**AMS subject classifications (2010):** 35B40, 35K57, 92C17.

## 1 Introduction

This paper concerns the higher-dimensional forager-exploiter model with different taxis strategies for two groups in search of food ([1, 12, 14]):

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla w) + \eta_1(u - u^m), & x \in \Omega, \quad t > 0, \\ v_t = \Delta v - \xi \nabla \cdot (v \nabla u) + \eta_2(v - v^l), & x \in \Omega, \quad t > 0, \\ w_t = \Delta w - \lambda(u + v)w - \mu w + r(x, t), & x \in \Omega, \quad t > 0, \\ \partial_\nu u = \partial_\nu v = \partial_\nu w = 0, & x \in \partial\Omega, \quad t > 0, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), \quad w(x, 0) = w_0(x), & x \in \Omega, \end{cases} \quad (1.1)$$

where  $\Omega \subset \mathbb{R}^n$  is a bounded domain with smooth boundary  $\partial\Omega$ ,  $\partial_\nu$  denotes differentiation with respect to the outward normal vector  $\nu$  on  $\partial\Omega$ . The unknown functions  $u, v$  and  $w$  are the densities of the forager population, exploiter population and nutrient, respectively. The constants  $\chi, \xi, \lambda, \mu$

<sup>1</sup>This work was supported by NSFC Grants 11771110, 11971128

<sup>2</sup>Corresponding author. E-mail: mxwang@hit.edu.cn

are supposed to be positive,  $m, l > 1$  and  $\eta_1, \eta_2$  are nonnegative. The given nonnegative function  $r(x, t)$  is the production rate of nutrient and satisfies

$$r \in C^1(\bar{\Omega} \times [0, \infty)) \cap L^\infty(\Omega \times (0, \infty)). \quad (1.2)$$

It is assumed that, besides the random diffusions, the first species (foragers) move toward the increasing nutrient gradient direction, while the second species (exploiters) follow the foragers to find the food indirectly.

The model (1.1) was initially proposed by Tania et al. [12]. When  $n = 1$ ,  $\eta_1 = \eta_2 = 0$  and  $r$  is a nonnegative constant, Tao and Winkler in [14] proved the global existence of the classical solution to (1.1), and showed that the solution stabilizes to a positive constant equilibrium exponentially provided that  $\min\{\int_\Omega u_0, \int_\Omega v_0\}$  is small enough. When  $n \geq 1$  and  $\eta_1 = \eta_2 = 0$ , by taking into account the volume-filling effect (i.e.,  $u_t = \Delta u - \nabla \cdot (u(1-u)\nabla w)$ ,  $v_t = \Delta v - \nabla \cdot (v(1-v)\nabla u)$  and  $0 \leq u_0, v_0 \leq 1$ ), the global existence of the classical solution was built in [8]. For  $\eta_1 = \eta_2 = 0$  and  $n \geq 1$ , under an explicit condition linking  $r$  and  $w_0$ , the global existence of generalized solutions of system (1.1) has been obtained and the long time behavior of the solution was established when  $r$  decays suitably in time ([19]). When  $n = 2$  and  $\eta_1, \eta_2 > 0$ , Black [1] showed that (1.1) has at least one global generalized solution if  $m > \sqrt{2} + 1$  and  $\min\{m, l\} > (m+1)/(m-1)$ , and proved that the generalized solution actually becomes a classical one after some waiting time  $T_*$  (possibly large) under conditions:  $m, l > \sqrt{2} + 1$ ,  $r \geq 0$  satisfies (1.2) and  $r \in L^1((0, \infty); L^\infty(\Omega))$ .

It is noticed that, none of the above mentioned works involve the global existence of classical solutions to (1.1) in high dimensions. The motivation of this paper is to establish the global solvability of (1.1) in high dimensions in the classical sense.

In order to better describe our contribution, we recall some related works on the original minimal Keller-Segel system in bounded domain

$$\begin{cases} u_t = \Delta u - \nabla \cdot (u \nabla v), & x \in \Omega, \quad t > 0, \\ v_t = \Delta v - v + u, & x \in \Omega, \quad t > 0, \\ \partial_\nu u = \partial_\nu v = 0, & x \in \partial\Omega, \quad t > 0, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), & x \in \Omega, \end{cases} \quad (1.3)$$

where  $\Omega \subset \mathbb{R}^n$  ( $n \geq 2$ ),  $u_0 \in C(\bar{\Omega})$  and  $v_0 \in W_\sigma^1(\Omega)$  with  $\sigma > n$ . It was shown in [2] that, there exist  $\varepsilon, \lambda > 0$  such that, if

$$\|u_0\|_{L^{n/2}(\Omega)} \leq \varepsilon, \quad \|v_0\|_{W_n^1(\Omega)} \leq \varepsilon,$$

then (1.3) admits a unique nonnegative classical solution  $(u, v)$  which exists globally in time and satisfies

$$\|u(\cdot, t) - \bar{u}_0\|_{L^\infty(\Omega)} + \|v(\cdot, t) - \bar{v}_0\|_{L^\infty(\Omega)} \leq C e^{-\lambda t}, \quad t > 0$$

with  $\bar{u}_0 = \frac{1}{|\Omega|} \int_\Omega u_0$ ,  $\bar{v}_0 = \frac{1}{|\Omega|} \int_\Omega v_0$  and some  $C > 0$ . For the minimal Keller-Segel model in  $\mathbb{R}^n$ , the boundedness result holds provided  $\|u_0\|_{L^r(\mathbb{R}^n)}$  and  $\|v_0\|_{W_q^1(\mathbb{R}^n)}$  small enough with  $r > n/2$  and  $q \geq n$  ([3]). For the prey-taxis system

$$\begin{cases} u_t = \Delta u - \nabla \cdot (u \nabla v) + uv, & x \in \Omega, \quad t > 0, \\ v_t = \Delta v + v(k - v) - uv, & x \in \Omega, \quad t > 0, \\ \partial_\nu u = \partial_\nu v = 0, & x \in \partial\Omega, \quad t > 0, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), & x \in \Omega, \end{cases} \quad (1.4)$$

it was shown in [15, 20] that, under a small condition on  $\max\{\|v_0\|_{L^\infty(\Omega)}, k\}$ , the classical solution to (1.4) is global in time and bounded. Inspired by above mentioned works (the classical Keller-Segel model [2, 3] and prey-taxis system [6, 15, 20]), it is reasonable to conjecture the global existence of the classical solutions to (1.1) under some smallness conditions. However, unlike the simple structure in the minimal Keller-Segel system or the prey-taxis model, the different taxis strategies in the system (1.1) produce more mathematical challenges and make the analysis work more complicate and difficult.

In this paper we investigate the global bounded solution of (1.1) for the higher-dimensional case:  $n \geq 2$ . Throughout this paper, we use  $C$ ,  $C'$  and  $C_i$  to represent generic positive constants, which may be different in different places. And, for simplicity, we use  $\int_\Omega f$  and  $\|f\|_p$  to denote  $\int_\Omega f dx$  and  $\|f\|_{L^p(\Omega)}$ , respectively. We suppose from now on that

$$u_0, w_0 \in W_\infty^2(\Omega), \quad v_0 \in W_\infty^1(\Omega), \quad \text{and} \quad u_0, v_0, w_0 \geq 0, \neq 0 \quad \text{in } \bar{\Omega}.$$

Our first statement is that, for the fixed  $\chi, \xi > 0$ , the small initial data can prevent blow-up.

**Theorem 1.1.** *Let  $n \geq 2$ ,  $\eta_1 = \eta_2 = 0$  and  $p = \min\{k \in \mathbb{N} : k > n/2\}$ . For the fixed  $\chi, \xi > 0$ , there exists  $\varepsilon > 0$  such that, if*

$$\|u_0\|_{W_{2p}^1(\Omega)} + \|w_0\|_{W_{2(p+1)}^1(\Omega)} + \|r(x, t)\|_{L^\infty(\Omega \times (0, \infty))} \leq \varepsilon,$$

*then (1.1) admits a unique nonnegative global bounded classical solution  $(u, v, w) \in [C(\bar{\Omega} \times [0, \infty)) \cap C^{2,1}(\bar{\Omega} \times (0, \infty))]^3$ . Moreover, there exists  $C > 0$  such that*

$$\|u(\cdot, t)\|_{W_{2p}^1(\Omega)} + \|v(\cdot, t)\|_{W_{2p}^1(\Omega)} + \|w(\cdot, t)\|_{W_{2p}^1(\Omega)} \leq C, \quad \forall t \in (0, \infty). \quad (1.5)$$

The second conclusion is that, for the given initial data  $(u_0, v_0, w_0)$ , if both the taxis effects  $\chi, \xi$  are weak enough, then the classical solution of (1.1) exists globally and remain bounded.

**Theorem 1.2.** *Let  $n \geq 2$  and  $\eta_1 = \eta_2 = 0$ . For the given initial data  $(u_0, v_0, w_0)$ , there exists  $\chi_0, \xi_0 > 0$  such that, if*

$$\chi \leq \chi_0, \quad \xi \leq \xi_0,$$

*then (1.1) admits a unique nonnegative global bounded classical solution  $(u, v, w) \in [C(\bar{\Omega} \times [0, \infty)) \cap C^{2,1}(\bar{\Omega} \times (0, \infty))]^3$  which satisfies (1.5).*

It is noted that Theorems 1.1 and 1.2 will be proved by the same arguments and both of them can be easily deduced from Lemma 3.6 below.

Next, we observe that, when  $\eta_1 > 0, m \geq 2$  and  $\eta_2 = 0$ , the solution exists globally and remain bounded in 2 dimension provided only  $\xi$  is small (without any restriction on  $\chi$ ).

**Theorem 1.3.** *Let  $n = 2$ ,  $\eta_1 > 0, m \geq 2$  and  $\eta_2 = 0$ . Then there exists  $\xi_0 > 0$  such that, for  $\xi \leq \xi_0$ , the problem (1.1) admits a unique nonnegative global bounded classical solution  $(u, v, w) \in [C(\bar{\Omega} \times [0, \infty)) \cap C^{2,1}(\bar{\Omega} \times (0, \infty))]^3$  which satisfies, for some  $C > 0$ ,*

$$\|u(\cdot, t)\|_{W_4^1(\Omega)} + \|v(\cdot, t)\|_{W_4^1(\Omega)} + \|w(\cdot, t)\|_{W_4^1(\Omega)} \leq C, \quad \forall t \in (0, \infty). \quad (1.6)$$

The last aim of this article is to show that the suitable large damping rates can prevent blow-up in two dimensional case.

**Theorem 1.4.** *Let  $n = 2$  and  $\eta_1, \eta_2 > 0$ . If*

$$m, l \geq 2 \quad \text{and} \quad l \geq \max \{3, 3\tilde{m}/(2\tilde{m} - 3)\} \quad \text{with} \quad \tilde{m} = \min\{m, l\}, \quad (1.7)$$

*then the problem (1.1) admits a unique nonnegative global bounded classical solution  $(u, v, w) \in [C(\bar{\Omega} \times [0, \infty)) \cap C^{2,1}(\bar{\Omega} \times (0, \infty))]^3$ . Moreover, (1.6) holds as well.*

**Remark 1.1.** *The condition (1.7) is equivalent to*

$$2 \leq m < 3, \quad l \geq 3m/(2m - 3) \quad \text{or} \quad m, l \geq 3. \quad (1.8)$$

*We recall from the pioneer work [10] and recent researches [21, 22] that the logistic growth or some sub-logistic source is enough to rule out any blow-up in the Keller-Segel systems in two dimensional case. It would be rather meaningful to investigate whether or not the condition (1.8) or (1.7) is the optimal one for ensuring the classical global solvability of (1.1) in two dimension. We leave this challenging problem as the future work.*

**Remark 1.2.** *In the present paper, we use the special form of the generalized logistic source  $\eta_1(u - u^m)$  and  $\eta_2(v - v^l)$  for the simplicity. It seems worthwhile to mention that, for the generic constant parameters, i.e.,  $\eta_1(a_1u - b_1u^m)$  and  $\eta_2(a_2v - b_2v^l)$  with  $a_i, b_i > 0$  for  $i = 1, 2$ , the conclusions in Theorem 1.3 and Theorem 1.4 hold as well.*

The article is organized as follows. Section 2 provides some basic preliminaries, including the local solvability of (1.1) and a fundamental ODE result. Section 3 is devoted to prove that the small values of initial data or taxis coefficients can prevent blow-up in any dimensions when  $\eta_1 = \eta_2 = 0$  (Theorem 1.1 and Theorem 1.2). In section 4, we show that, in the case  $\eta_1 > 0, m \geq 2$  and  $\eta_2 = 0$ , the small value of  $\xi$  is enough to ensure the global existence of the solution to (1.1) (Theorem 1.3). We finally prove in Section 5 that, when  $\eta_1, \eta_2 > 0$  and logistic degradation rates  $m, l$  are sufficiently large, then (1.1) is globally solvable (Theorem 1.4).

## 2 Local existence and preliminaries

The local solvability of (1.1) are well established by the Amann theory (cf. [14, Lemma 2.1] or [17]). The positivity of solution and the uniform-in-time boundedness of  $w$  are due to the parabolic maximum principle.

**Lemma 2.1.** *Suppose that  $n \geq 1, \eta_i \geq 0$  with  $i = 1, 2$  and  $m, l > 1$ . Then there exist a  $T_m \in (0, \infty]$  and a unique solution  $(u, v, w)$  which solves (1.1) in  $[0, T_m)$ ,  $u, v, w \in C(\bar{\Omega} \times [0, T_m)) \cap C^{2,1}(\bar{\Omega} \times (0, T_m))$  and satisfies*

$$u, v > 0, \quad 0 < w \leq Q \quad \text{in} \quad \bar{\Omega} \times (0, T_m). \quad (2.1)$$

*Moreover, the “existence time  $T_m$ ” can be chosen maximal: either  $T_m = \infty$ , or  $T_m < \infty$  and*

$$\limsup_{t \rightarrow T_m} (\|u(\cdot, t)\|_{W_q^1(\Omega)} + \|v(\cdot, t)\|_{W_q^1(\Omega)} + \|w(\cdot, t)\|_{W_q^1(\Omega)}) = \infty \quad \text{for all } q > n.$$

The following lemma gives a statement on ODE comparison.

**Lemma 2.2.** *Let  $a, b > 0$ . Assume that for some  $\hat{T} \in (0, \infty]$  and  $\hat{\tau} = \min\{1, \hat{T}/2\}$ , the nonnegative functions  $y \in C([0, \hat{T})) \cap C^1((0, \hat{T}))$ ,  $f \in L^1_{loc}([0, \hat{T}))$  and satisfy*

$$\begin{aligned} y'(t) + ay(t) &\leq f(t), \quad t \in (0, \hat{T}), \\ \int_t^{t+\hat{\tau}} f(s)ds &\leq b, \quad t \in (0, \hat{T} - \hat{\tau}). \end{aligned} \quad (2.2)$$

Then

$$y(t) \leq y(0) + 2b + b/a, \quad t \in (0, \hat{T}). \quad (2.3)$$

*Proof.* If  $\hat{T} > 2$ , then  $\hat{\tau} = 1$ . Thanks to [11, Lemma 3.4], we get (2.3).

If  $\hat{T} \leq 2$ , then  $\hat{T} = 2\hat{\tau}$ . For any  $t_0 \in [0, \hat{T})$  and  $t \in [t_0, t_0 + \hat{\tau}] \cap [0, \hat{T})$ , integrating (2.2) from  $t_0$  to  $t$ , there holds

$$y(t) \leq y(t_0) + \int_{t_0}^t f(s)ds \leq y(t_0) + b. \quad (2.4)$$

Hence,  $y(t) \leq y(0) + b$  for any  $t \in [0, \hat{\tau}]$ . Again by (2.4) and  $\hat{T} = 2\hat{\tau}$ , we have

$$y(t) \leq y(\hat{\tau}) + b \leq y(0) + 2b, \quad t \in (\hat{\tau}, \hat{T}).$$

Therefore, for any  $t \in (0, \hat{T})$ , there holds that  $y(t) \leq y(0) + 2b$ . This completes the proof.  $\square$

### 3 Proofs of Theorem 1.1 and Theorem 1.2

First of all, we introduce some notations. Let  $r_* = \|r(x, t)\|_{L^\infty(\Omega \times (0, \infty))}$  and

$$\begin{aligned} A &= 2\|u_0\|_\infty, \quad B = 2\|v_0\|_\infty, \quad Q = \max\{\|w_0\|_\infty, r_*/\mu\}, \\ p &= \min\{k \in \mathbb{N} : k > n/2\}, \quad \Sigma = \{\eta, \lambda, \mu, n, p, \Omega\}, \\ G_0 &= (A + B + 1)(\|w_0\|_{W^1_{2(p+1)}(\Omega)} + r_*), \quad H_0 = (A + B + 1)Q + \|w_0\|_{W^2_{p+1}(\Omega)} + r_*. \end{aligned}$$

Moreover, we define

$$T := \sup\{\tilde{t} \in (0, T_m) : \|u(\cdot, t)\|_\infty \leq A, \|v(\cdot, t)\|_\infty \leq B \text{ in } (0, \tilde{t})\}.$$

Clearly, by the continuity of the solution, we have  $T \in (0, T_m]$  and

$$\sup_{t \in (0, T)} \|u(\cdot, t)\|_\infty \leq A, \quad \sup_{t \in (0, T)} \|v(\cdot, t)\|_\infty \leq B. \quad (3.1)$$

We set from now on that  $\tau := \min\{1, T/2\}$ . The following lemma provides the  $L^{2(p+1)}$ -bound of  $\nabla w$  in  $(0, T)$ .

**Lemma 3.1.** *Let  $n \geq 2$ ,  $\eta_1 = \eta_2 = 0$ . Then there exists  $C = C(\Sigma) > 0$  such that*

$$\int_\Omega |\nabla w(\cdot, t)|^{2(p+1)} \leq CG_0^{2(p+1)}, \quad t \in (0, T). \quad (3.2)$$

*Proof.* According to the standard  $L^p$ - $L^q$  estimates for  $(e^{t\Delta})_{t \geq 0}$  ([18, Lemma 1.3 (ii)-(iii)]), we can find  $\lambda_1, C_1, C_2 > 0$  depending on  $\Sigma$  such that, for all  $t \in (0, T)$ ,

$$\begin{aligned} \|\nabla w(\cdot, t)\|_{2(p+1)} &\leq C_1 \|w_0\|_{W_{2(p+1)}^1(\Omega)} + C_1 \int_0^t \|\nabla e^{(t-s)\Delta} f(\cdot, s)\|_{2(p+1)} ds \\ &\leq C_1 \|w_0\|_{W_{2(p+1)}^1(\Omega)} + C_2 \int_0^t \left(1 + (t-s)^{-\frac{1}{2}}\right) e^{-\lambda_1(t-s)} \|f(\cdot, s)\|_{2(p+1)} ds, \end{aligned} \quad (3.3)$$

where  $f = -\lambda(u+v)w - \mu w + r$ . It is easy to see from (3.1) and (2.1) that, there is  $C_3 = C_3(\Sigma) > 0$  such that

$$\|f(\cdot, s)\|_{2(p+1)} \leq C_3[(A+B+1)Q + r_*], \quad \forall s \in (0, T).$$

Inserting this into (3.3) and using the Sobolev embedding theorem:  $W_{2(p+1)}^1(\Omega) \hookrightarrow L^\infty(\Omega)$  with  $p+1 > n/2 + 1$ , one can find  $C_4, C_5 > 0$  depending on  $\Sigma$  such that

$$\|\nabla w(\cdot, t)\|_{2(p+1)} \leq C_4[(A+B+1)Q + \|w_0\|_{W_{2(p+1)}^1(\Omega)} + r_*] \leq C_5 G_0, \quad t \in (0, T).$$

The proof is end.  $\square$

We proceed to find a space-time  $L^{p+1}$  bound for  $\Delta w$ .

**Lemma 3.2.** *Suppose that  $n \geq 2$  and  $\eta_1 = \eta_2 = 0$ . Then there exists  $C = C(\Sigma) > 0$  such that*

$$\int_t^{t+\tau} \int_\Omega |\Delta w|^{p+1} dx ds \leq C H_0^{p+1}, \quad t \in (0, T - \tau). \quad (3.4)$$

*Proof.* This proof is based on [9, Lemma 4.2] and [1, Lemma 4.3]. In view of the maximal Sobolev regularity properties of the Neumann heat semigroup  $(e^{t\Delta})_{t \geq 0}$  ([5]) and (3.1), we have

$$\int_0^{2\tau} \|w(\cdot, s)\|_{W_{p+1}^{p+1}(\Omega)}^{p+1} ds \leq C_1 \|w_0\|_{W_{p+1}^{p+1}(\Omega)}^{p+1} + C_1 \int_0^{2\tau} \|f(\cdot, s)\|_{p+1}^{p+1} ds \leq C_2 H_0^{p+1}, \quad (3.5)$$

where  $f = -\lambda(u+v)w - \mu w + r$  and  $C_i = C_i(\Sigma) > 0$  with  $i = 1, 2$ .

**Case I:**  $T > 2$ . In this case, it is easy to see from the definition of  $\tau$  that  $\tau = 1$ . Thus, (3.4) is equivalent to

$$\int_t^{t+1} \int_\Omega |\Delta w|^{p+1} dx ds \leq C H_0^{p+1}, \quad t \in (0, T - 1). \quad (3.6)$$

From (3.5) with  $\tau = 1$ , we know that (3.6) holds for  $t \in (0, 1]$ .

For  $t_0 \in (1, T - 1)$ , setting  $\sigma := t_0 - 1$  and hence  $\sigma \in (0, T - 2)$ . Let  $\rho \in C^\infty(\mathbb{R})$  be an increasing function satisfying

$$0 \leq \rho \leq 1 \quad \text{in } \mathbb{R}, \quad \rho \equiv 0 \quad \text{in } (-\infty, 0], \quad \rho \equiv 1 \quad \text{in } (1, \infty),$$

and define  $\rho_\sigma(t) = \rho(t - \sigma)$ . Clearly,  $\|\rho'\|_{C^1(\mathbb{R})} \leq C_3$  for some  $C_3 > 0$ .

Let  $\varphi$  be the unique classical solution of

$$\begin{cases} \varphi_t = \Delta \varphi, & x \in \Omega, \quad t > \sigma, \\ \partial_\nu \varphi = 0, & x \in \partial\Omega, \quad t > \sigma, \\ \varphi(x, \sigma) = w(x, \sigma), & x \in \Omega. \end{cases}$$

It is easy to deduce that  $\|\varphi(\cdot, t)\|_\infty \leq Q$  for  $t \in (\sigma, \infty)$ , and  $\phi := \rho_\sigma(t)\varphi$  solves

$$\begin{cases} \phi_t = \Delta \phi + \rho'_\sigma(t)\varphi, & x \in \Omega, \quad t > \sigma, \\ \partial_\nu \phi = 0, & x \in \partial\Omega, \quad t > \sigma, \\ \phi(x, \sigma) = 0, & x \in \Omega. \end{cases}$$

Thanks to the maximal Sobolev regularity properties of the Neumann heat semigroup  $(e^{t\Delta})_{t \geq 0}$  ([5]), there holds

$$\int_\sigma^{\sigma+2} \|\phi(\cdot, s)\|_{W_{p+1}^{2,p+1}(\Omega)}^{p+1} ds \leq C_4 \int_\sigma^{\sigma+2} \|\rho'_\sigma(s)\varphi(\cdot, s)\|_{p+1}^{p+1} ds \leq 2C_4|\Omega| \|\rho'\|_{C^1(\mathbb{R})}^{p+1} Q^{p+1} \leq C_5 Q^{p+1}, \quad (3.7)$$

where positive constants  $C_4 = C_4(n, p, \Omega)$  and  $C_5 = 2C_3C_4|\Omega|$ . Noticing that  $\phi(\cdot, t) = \varphi(\cdot, t)$  for  $t > \sigma + 1$ . Hence, we have from (3.7) that

$$\int_{\sigma+1}^{\sigma+2} \|\varphi(\cdot, s)\|_{W_{p+1}^{2,p+1}(\Omega)}^{p+1} ds \leq C_5 Q^{p+1}, \quad \sigma \in (0, T-2). \quad (3.8)$$

Let  $z(x, t) = w(x, t) - \varphi(x, t)$  for  $x \in \Omega$  and  $t \in [\sigma, T)$ , then  $z$  satisfies

$$\begin{cases} z_t = \Delta z + f(x, t), & x \in \Omega, \quad t \in (\sigma, T), \\ \partial_\nu z = 0, & x \in \partial\Omega, \quad t \in (\sigma, T), \\ z(x, \sigma) = 0, & x \in \Omega. \end{cases}$$

Again by the maximal Sobolev regularity properties of the Neumann heat semigroup  $(e^{t\Delta})_{t \geq 0}$  ([5]), one can find  $C_6, C_7 > 0$  depending on  $\Sigma$  such that

$$\int_\sigma^{\sigma+2} \|z(\cdot, s)\|_{W_{p+1}^{2,p+1}(\Omega)}^{p+1} ds \leq C_6 \int_\sigma^{\sigma+2} \|f(\cdot, s)\|_{p+1}^{p+1} ds \leq C_7 H_0^{p+1}. \quad (3.9)$$

Note that

$$\|w(\cdot, s)\|_{W_{p+1}^2(\Omega)} \leq \|z(\cdot, s)\|_{W_{p+1}^2(\Omega)} + \|\varphi(\cdot, s)\|_{W_{p+1}^2(\Omega)}, \quad \forall s \in (\sigma+1, \sigma+2).$$

This combined with (3.8) and (3.9) yields that

$$\int_{\sigma+1}^{\sigma+2} \|w(\cdot, s)\|_{W_{p+1}^{2,p+1}(\Omega)}^{p+1} ds \leq C_8 H_0^{p+1}, \quad \sigma \in (0, T-2),$$

and hence

$$\int_{t_0}^{t_0+1} \|w(\cdot, s)\|_{W_{p+1}^{2,p+1}(\Omega)}^{p+1} ds \leq C_8 H_0^{p+1}, \quad t_0 \in (1, T-1)$$

for the positive constant  $C_8 = C_8(\Sigma)$ . We thus obtain (3.6) for  $t \in (1, T-1)$  due to the arbitrariness of  $t_0$  in  $(1, T-1)$ . Recalling (3.5) with  $\tau = 1$ , we get (3.6).

**Case II:**  $T \leq 2$ . In this case  $\tau = T/2$ , i.e.,  $T = 2\tau$ . Due to (3.5), we obtain (3.4).  $\square$

Thanks to Lemmas 3.1, 3.2, we establish an uniform-in-time  $L^4$  regularity and a space-time  $L^6$  estimate for  $\nabla u$ .

**Lemma 3.3.** *Let  $n \geq 2$  and  $\eta_i = 0$  with  $i = 1, 2$ . Then there exists  $C = C(\Sigma) > 0$  such that*

$$\int_{\Omega} |\nabla u(\cdot, t)|^{2p} \leq C \|u_0\|_{W_{2p}^1(\Omega)}^{2p} [(\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1], \quad t \in (0, T). \quad (3.10)$$

Moreover, there is  $C' = C'(\chi, A, B, \Sigma) > 0$  such that

$$\int_t^{t+\tau} \int_{\Omega} |\nabla u|^{2(p+1)} dx ds \leq C', \quad t \in (0, T - \tau). \quad (3.11)$$

*Proof.* We first recall from [7, Lemma 2.2] that, for  $t \in (0, T_m)$ ,

$$\int_{\Omega} |\nabla u|^{2(p+1)} \leq 2(n + 4p^2) \|u\|_{\infty}^2 \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2, \quad (3.12)$$

and hence by (3.1), we have

$$\int_{\Omega} |\nabla u|^{2(p+1)} \leq 2(n + 4p^2) A^2 \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 =: k \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2, \quad t \in (0, T). \quad (3.13)$$

It follows from the first equation in (1.1) that

$$\begin{aligned} & \frac{1}{2p} \frac{d}{dt} \int_{\Omega} |\nabla u|^{2p} + \int_{\Omega} |\nabla u|^{2p} \\ &= \int_{\Omega} |\nabla u|^{2(p-1)} \nabla u \cdot \nabla u_t + \int_{\Omega} |\nabla u|^{2p} \\ &= \int_{\Omega} |\nabla u|^{2(p-1)} \nabla u \cdot \nabla (\Delta u - \chi \nabla \cdot (u \nabla w)) + \int_{\Omega} |\nabla u|^{2p} \\ &= \int_{\Omega} |\nabla u|^{2(p-1)} \nabla u \cdot \nabla \Delta u + \chi \int_{\Omega} \nabla \cdot (|\nabla u|^{2(p-1)} \nabla u) (\nabla \cdot (u \nabla w)) + \int_{\Omega} |\nabla u|^{2p} \\ &=: I(t) + J(t) + \int_{\Omega} |\nabla u|^{2p}, \quad t \in (0, T_m). \end{aligned} \quad (3.14)$$

In view of [16, Lemma 2.6 (ii)], there is  $C_1 = C_1(n, p, \Omega) > 0$  such that

$$\int_{\partial\Omega} |\nabla u|^{2(p-1)} \partial_{\nu} |\nabla u|^2 dS \leq (p-1) \int_{\Omega} |\nabla u|^{2(p-2)} |\nabla |\nabla u|^2|^2 + C_1 \int_{\Omega} |\nabla u|^{2p}, \quad t \in (0, T_m).$$

Hence, by direct computations, we have

$$\begin{aligned} I(t) &= \frac{1}{2} \int_{\Omega} |\nabla u|^{2(p-1)} \Delta |\nabla u|^2 - \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 \\ &= -\frac{1}{2} \int_{\Omega} \nabla |\nabla u|^{2(p-1)} \cdot \nabla |\nabla u|^2 + \frac{1}{2} \int_{\partial\Omega} |\nabla u|^{2(p-1)} \partial_{\nu} |\nabla u|^2 dS - \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 \\ &= -\frac{p-1}{2} \int_{\Omega} |\nabla u|^{2(p-2)} |\nabla |\nabla u|^2|^2 + \frac{1}{2} \int_{\partial\Omega} |\nabla u|^{2(p-1)} \partial_{\nu} |\nabla u|^2 dS - \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 \\ &\leq -\int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 + \frac{C_1}{2} \int_{\Omega} |\nabla u|^{2p}, \quad t \in (0, T_m). \end{aligned} \quad (3.15)$$

Next, we estimate  $J(t)$ . It is easy to deduce that

$$\begin{aligned} J(t) &= \chi \int_{\Omega} \left( \nabla |\nabla u|^{2(p-1)} \cdot \nabla u + |\nabla u|^{2(p-1)} \Delta u \right) (\nabla u \cdot \nabla w + u \Delta w) \\ &= \chi \int_{\Omega} \left( \nabla |\nabla u|^{2(p-1)} \cdot \nabla u \right) (\nabla u \cdot \nabla w) + \chi \int_{\Omega} u \Delta w \left( \nabla |\nabla u|^{2(p-1)} \cdot \nabla u \right) \end{aligned}$$



$$\begin{aligned}
& +\chi \int_{\Omega} |\nabla u|^{2(p-1)} \Delta u (\nabla u \cdot \nabla w) + \chi \int_{\Omega} u |\nabla u|^{2(p-1)} \Delta u \Delta w \\
& =: J_1(t) + J_2(t) + J_3(t) + J_4(t), \quad t \in (0, T_m).
\end{aligned} \tag{3.16}$$

Noticing that  $\nabla |\nabla u|^2 = 2D^2u \cdot \nabla u$ . Using Young's inequality:  $ab \leq |a|^q + |b|^{q'}$  with  $q > 1$  and  $q' = q/(q-1)$ , (3.1) and (3.13), the terms  $J_1(t), J_2(t)$  in the right hand side of (3.16) can be estimated as

$$\begin{aligned}
J_1(t) &= \chi(p-1) \int_{\Omega} \left( |\nabla u|^{2(p-2)} \nabla |\nabla u|^2 \cdot \nabla u \right) (\nabla u \cdot \nabla w) \\
&\leq \chi(p-1) \int_{\Omega} |\nabla u|^{2(p-1)} |\nabla w| |\nabla |\nabla u|^2| \\
&= 2\chi(p-1) \int_{\Omega} |\nabla u|^{2(p-1)} |\nabla w| |D^2u \cdot \nabla u| \\
&\leq 2\chi(p-1) \int_{\Omega} |\nabla u|^{2p-1} |\nabla w| |D^2u| \\
&\leq \frac{1}{8} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2u|^2 + 8\chi^2(p-1)^2 \int_{\Omega} |\nabla u|^{2p} |\nabla w|^2 \\
&\leq \frac{1}{8} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2u|^2 + \frac{1}{8k} \int_{\Omega} |\nabla u|^{2(p+1)} + 8^{2p+1} k^p [\chi(p-1)]^{2(p+1)} \int_{\Omega} |\nabla w|^{2(p+1)} \\
&\leq \frac{1}{4} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2u|^2 + 8^{2p+1} k^p [\chi(p-1)]^{2(p+1)} \int_{\Omega} |\nabla w|^{2(p+1)} \\
&= \frac{1}{4} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2u|^2 + C_2 A^{2p} \chi^{2(p+1)} \int_{\Omega} |\nabla w|^{2(p+1)}, \quad t \in (0, T),
\end{aligned} \tag{3.17}$$

where  $C_2 = 8^{2p+1} [2(n+4p^2)]^p (p-1)^{2(p+1)}$  due to the definition of  $k$ , and

$$\begin{aligned}
J_2(t) &= \chi(p-1) \int_{\Omega} u |\nabla u|^{2(p-2)} \Delta w (\nabla |\nabla u|^2 \cdot \nabla u) \\
&\leq \chi(p-1) \int_{\Omega} u |\Delta w| |\nabla u|^{2p-3} |\nabla |\nabla u|^2| \\
&\leq 2\chi(p-1) \int_{\Omega} u |\Delta w| |\nabla u|^{2(p-1)} |D^2u| \\
&\leq \frac{1}{8} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2u|^2 + 8[\chi A(p-1)]^2 \int_{\Omega} |\nabla u|^{2(p-1)} |\Delta w|^2 \\
&\leq \frac{1}{8} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2u|^2 + \frac{1}{8k} \int_{\Omega} |\nabla u|^{2(p+1)} + 8^p k^{(p-1)/2} [\chi A(p-1)]^{p+1} \int_{\Omega} |\Delta w|^{p+1} \\
&\leq \frac{1}{4} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2u|^2 + 8^p k^{(p-1)/2} [\chi A(p-1)]^{p+1} \int_{\Omega} |\Delta w|^{p+1} \\
&= \frac{1}{4} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2u|^2 + C_3 A^{2p} \chi^{p+1} \int_{\Omega} |\Delta w|^{p+1}, \quad t \in (0, T)
\end{aligned} \tag{3.18}$$

with  $C_3 = 8^p [2(n+4p^2)]^{(p-1)/2} (p-1)^{p+1}$ . In view of (3.1), (3.13) and the known inequality:  $|\Delta u| \leq \sqrt{n} |D^2u|$ , we estimate the last two terms in the right hand side of (3.16) as follows:

$$\begin{aligned}
J_3(t) &\leq \sqrt{n} \chi \int_{\Omega} |\nabla u|^{2p-1} |\nabla w| |D^2u| \\
&\leq \frac{1}{8} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2u|^2 + 2n \chi^2 \int_{\Omega} |\nabla u|^{2p} |\nabla w|^2
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{8} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 + \frac{1}{8k} \int_{\Omega} |\nabla u|^{2(p+1)} + (8k)^p (2n\chi^2)^{p+1} \int_{\Omega} |\nabla w|^{2(p+1)} \\
&\leq \frac{1}{4} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 + (8k)^p (2n\chi^2)^{p+1} \int_{\Omega} |\nabla w|^{2(p+1)} \\
&= \frac{1}{4} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 + C_4 A^{2p} \chi^{2(p+1)} \int_{\Omega} |\nabla w|^{2(p+1)}, \quad t \in (0, T)
\end{aligned} \tag{3.19}$$

with  $C_4 = [16(n + 4p^2)]^p (2n)^{p+1}$ , and

$$\begin{aligned}
J_4(t) &\leq \sqrt{n} A \chi \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u| |\Delta w| \\
&\leq \frac{1}{8} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 + 2n A^2 \chi^2 \int_{\Omega} |\nabla u|^{2(p-1)} |\Delta w|^2 \\
&\leq \frac{1}{8} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 + \frac{1}{16k} \int_{\Omega} |\nabla u|^{2(p+1)} + (16k)^{\frac{p-1}{2}} (2n A^2 \chi^2)^{\frac{p+1}{2}} \int_{\Omega} |\Delta w|^{p+1} \\
&\leq \frac{3}{16} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 + (16k)^{(p-1)/2} (2n A^2 \chi^2)^{\frac{p+1}{2}} \int_{\Omega} |\Delta w|^{p+1} \\
&= \frac{3}{16} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 + C_5 A^{2p} \chi^{p+1} \int_{\Omega} |\Delta w|^{p+1}, \quad t \in (0, T),
\end{aligned} \tag{3.20}$$

where  $C_5 = [32(n + 4p^2)]^{(p-1)/2} (2n)^{\frac{p+1}{2}}$ . Inserting (3.17)-(3.20) into (3.16) we find that

$$J(t) \leq \frac{15}{16} \int_{\Omega} |\nabla u|^2 |D^2 u|^2 + C' A^{2p} \chi^{2(p+1)} \int_{\Omega} |\nabla w|^{2(p+1)} + C'' A^{2p} \chi^{p+1} \int_{\Omega} |\Delta w|^{p+1}, \quad t \in (0, T)$$

with  $C' = C_2 + C_4$ ,  $C'' = C_3 + C_5$ . Plugging this and (3.15) into (3.14) yields that, for  $t \in (0, T)$ ,

$$\begin{aligned}
&\frac{1}{2p} \frac{d}{dt} \int_{\Omega} |\nabla u|^{2p} + \int_{\Omega} |\nabla u|^{2p} + \frac{1}{16} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 \\
&\leq C' A^{2p} \chi^{2(p+1)} \int_{\Omega} |\nabla w|^{2(p+1)} + C'' A^{2p} \chi^{p+1} \int_{\Omega} |\Delta w|^{p+1} + \left( \frac{C_1}{2} + 1 \right) \int_{\Omega} |\nabla u|^{2p}.
\end{aligned} \tag{3.21}$$

By using Young's inequality and (3.13), one has

$$\begin{aligned}
\left( \frac{C_1}{2} + 1 \right) \int_{\Omega} |\nabla u|^{2p} &\leq \frac{1}{32k} \int_{\Omega} |\nabla u|^{2(p+1)} + (32k)^p \left( \frac{C_1}{2} + 1 \right)^{p+1} |\Omega| \\
&\leq \frac{1}{32} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 + (32k)^p \left( \frac{C_1}{2} + 1 \right)^{p+1} |\Omega| \\
&= \frac{1}{32} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 + C_6 A^{2p}, \quad t \in (0, T)
\end{aligned}$$

with  $C_6 = [64(n + 4p^2)]^p \left( \frac{C_1}{2} + 1 \right)^{p+1} |\Omega|$ . Inserting this into (3.21) gives that, for  $t \in (0, T)$ ,

$$\begin{aligned}
&\frac{1}{2p} \frac{d}{dt} \int_{\Omega} |\nabla u|^{2p} + \int_{\Omega} |\nabla u|^{2p} + \frac{1}{32} \int_{\Omega} |\nabla u|^{2(p-1)} |D^2 u|^2 \\
&\leq C' A^{2p} \chi^{2(p+1)} \int_{\Omega} |\nabla w|^{2(p+1)} + C'' A^{2p} \chi^{p+1} \int_{\Omega} |\Delta w|^{p+1} + C_6 A^{2p} \\
&\leq C_7 A^{2p} \left( \chi^{2(p+1)} \int_{\Omega} |\nabla w|^{2(p+1)} + \chi^{p+1} \int_{\Omega} |\Delta w|^{p+1} + 1 \right) =: g(t), \quad t \in (0, T),
\end{aligned} \tag{3.22}$$

where  $C_7 = \max\{C', C'', C_6\}$  depending only on  $n, p, \Omega$ . Applying (3.2) and (3.4), there exists  $C_8 = C_8(\Sigma) > 0$  such that

$$\int_t^{t+\tau} g(s)ds \leq C_8 A^{2p} [(\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1], \quad t \in (0, T - \tau).$$

In view of Lemma 2.2 with  $a = 1$ ,  $b = C_8 A^{2p} [(\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1]$ ,  $y(t) = \frac{1}{2p} \int_\Omega |\nabla u|^4$  and  $f(t) = g(t)$ , we find  $C_9, C_{10} > 0$  depending on  $\Sigma$  such that

$$\begin{aligned} \int_\Omega |\nabla u(\cdot, t)|^{2p} &\leq C_9 A^{2p} [(\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1] + C_9 \|\nabla u_0\|_{2p}^{2p} \\ &\leq C_{10} \|u_0\|_{W_{2p}^1(\Omega)}^{2p} [(\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1], \quad t \in (0, T), \end{aligned}$$

where we have used  $A = 2\|u_0\|_\infty$  and the embedding:  $W_{2p}^1(\Omega) \hookrightarrow L^\infty(\Omega)$  with  $p > n/2$  in the derivation of the last inequality. This establishes (3.10).

Integrating (3.22) from  $t$  to  $t + \tau$  with  $t \in (0, T - \tau)$ , and using (3.2), (3.4) and (3.10), we have

$$\int_t^{t+\tau} \int_\Omega |\nabla u|^{2(p-1)} |D^2 u|^2 dx ds \leq C^*, \quad t \in (0, T - \tau)$$

for some  $C^* > 0$  depending on  $\chi, A, B, \Sigma$ . This in conjunction with (3.13) implies

$$\int_t^{t+\tau} \int_\Omega |\nabla u|^{2(p+1)} dx ds \leq \frac{C^*}{2(n + 4p^2)A^2}, \quad t \in (0, T - \tau).$$

Hence, we finally get (3.11) and the proof is completed.  $\square$

**Remark 3.1.** From the proof of Lemma 3.3, with some minor modifications, we see that for  $\eta_1, \eta_2 \geq 0$  and  $m, l > 1$ , the inequality (3.22) holds as well provided  $u$  is uniformly bounded in time. And, the inequality (3.22) also plays an important role in the proofs of Theorems 1.3, 1.4.

Based on Lemma 3.1, we obtain uniform  $L^\infty$  boundedness for  $u$ .

**Lemma 3.4.** Let  $n \geq 2$ ,  $\eta_i = 0$  with  $i = 1, 2$ . Then there exists  $C = C(\Sigma) > 0$  such that

$$\|u(\cdot, t)\|_\infty \leq \|u_0\|_\infty + CA\chi G_0, \quad t \in (0, T).$$

*Proof.* By using the standard  $L^p$ - $L^q$  estimates for  $(e^{t\Delta})_{t \geq 0}$  ([4, Lemma 3.3]), there exist positive constants  $\lambda_1, C_1$  depending on  $n, \Omega$  such that

$$\begin{aligned} \|u(\cdot, t)\|_\infty &\leq \|u_0\|_\infty + \chi \int_0^t \|e^{(t-s)\Delta} \nabla \cdot (u \nabla w)\|_\infty ds \\ &\leq \|u_0\|_\infty + C_1 \chi \int_0^t \left(1 + (t-s)^{-\frac{1}{2} - \frac{n}{4(p+1)}}\right) e^{-\lambda_1(t-s)} \|u \nabla w\|_{2(p+1)} ds, \quad t \in (0, T). \end{aligned}$$

Thanks to (3.1) and Lemma 3.1, there is  $C_2 = C_2(\Sigma) > 0$  such that

$$\|u(\cdot, \sigma) \nabla w(\cdot, \sigma)\|_{2(p+1)} \leq C_2 A G_0, \quad \sigma \in (0, T).$$

This in conjunction with the fact that  $0 < \frac{1}{2} + \frac{n}{4(p+1)} < 1$  due to  $p+1 > \frac{n}{2} + 1$ , one can find  $C_3 > 0$  depending on  $\Sigma$  fulfilling

$$\|u(\cdot, t)\|_\infty \leq \|u_0\|_\infty + C_1 C_2 G_0 \chi \int_0^t \left(1 + (t-s)^{-\frac{1}{2} - \frac{n}{4(p+1)}}\right) e^{-\lambda_1(t-s)} ds \leq \|u_0\|_\infty + C_3 A \chi G_0$$

for all  $t \in (0, T)$ . The proof is finished.  $\square$

With Lemma 3.3 at hand, we derive an upper bound for  $v$  in  $(0, T)$ .

**Lemma 3.5.** *Let  $n \geq 2$ ,  $\eta_1 = \eta_2 = 0$ . Then there exists  $C = C(\Sigma) > 0$  such that, for  $t \in (0, T)$ ,*

$$\|v(\cdot, t)\|_\infty \leq \|v_0\|_\infty + C\xi B\|u_0\|_{W_{2p}^1(\Omega)} \left[ (\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1 \right]^{1/2p}.$$

*Proof.* Thanks to the standard  $L^p$ - $L^q$  estimates for  $(e^{t\Delta})_{t \geq 0}$  ([4, Lemma 3.3]), there exist  $\lambda_1, C_1 > 0$  depending on  $n, p$  such that

$$\begin{aligned} \|v(\cdot, t)\|_\infty &\leq \|v_0\|_\infty + \xi \int_0^t \|e^{(t-s)\Delta} \nabla \cdot (v \nabla u)\|_\infty ds \\ &\leq \|v_0\|_\infty + C_1 \xi \int_0^t \left( 1 + (t-s)^{-\frac{1}{2} - \frac{n}{4p}} \right) e^{-\lambda_1(t-s)} \|v \nabla u\|_{2p} ds, \quad t \in (0, T). \end{aligned}$$

In view of (3.1) and (3.10), there exists  $C_2 = C_2(\Sigma) > 0$  such that

$$\begin{aligned} \|v(\cdot, \sigma) \nabla u(\cdot, \sigma)\|_{2p} &\leq \|v(\cdot, \sigma)\|_\infty \|\nabla u(\cdot, \sigma)\|_{2p} \\ &\leq C_2 B \|u_0\|_{W_{2p}^1(\Omega)} \left[ (\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1 \right]^{1/2p}, \quad \sigma \in (0, T). \end{aligned}$$

This combined the fact that  $0 < \frac{1}{2} + \frac{n}{4p} < 1$  due to  $p > \frac{n}{2}$ , we find  $C_3 = C_3(\Sigma) > 0$  such that

$$\|v(\cdot, t)\|_\infty \leq \|v_0\|_\infty + C_3 \xi B \|u_0\|_{W_{2p}^1(\Omega)} \left[ (\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1 \right]^{1/2p}, \quad t \in (0, T).$$

The proof is end.  $\square$

We finally show the global existence and boundedness of the classical solution of (1.1) under the condition that the initial data or  $\chi, \xi$  are small enough.

**Lemma 3.6.** *Let  $n \geq 2$  and  $\eta_1 = \eta_2 = 0$ . There exists  $\kappa = \kappa(\Sigma) > 0$  such that, if  $\chi, \xi$  and  $(u_0, v_0, w_0)$  satisfy*

$$\chi \leq \frac{\kappa}{G_0}, \quad \text{and} \quad \xi \leq \frac{\kappa}{\|u_0\|_{W_{2p}^1(\Omega)} \left[ (\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1 \right]^{1/2p}}, \quad (3.23)$$

*then (1.1) admits a unique nonnegative global solution  $(u, v, w) \in [C(\bar{\Omega} \times [0, \infty)) \cap C^{2,1}(\bar{\Omega} \times (0, \infty))]^3$ , and*

$$\|u(\cdot, t)\|_\infty \leq A, \quad \|v(\cdot, t)\|_\infty \leq B, \quad \|w(\cdot, t)\|_\infty \leq Q, \quad t \in (0, \infty).$$

*Moreover, there exists  $C = C(\chi, \xi, A, B, \Sigma) > 0$  such that*

$$\|u(\cdot, t)\|_{W_{2p}^1(\Omega)} + \|v(\cdot, t)\|_{W_{2p}^1(\Omega)} + \|w(\cdot, t)\|_{W_{2p}^1(\Omega)} \leq C, \quad t \in (0, \infty). \quad (3.24)$$

*Proof.* We shall break this proof into three steps and use  $C_i$  to denote the general positive constants that may depend on  $\chi, \xi, \Sigma$  and the initial data.

*Step 1: The uniform boundedness of the solution in  $(0, T_m)$ .* From Lemma 3.4 and Lemma 3.5, for any  $\chi, \xi > 0$ , there exists  $K = K(\Sigma) > 0$  such that

$$\|u(\cdot, t)\|_\infty \leq \|u_0\|_\infty + K\chi A G_0, \quad t \in (0, T), \quad (3.25)$$

and

$$\|v(\cdot, t)\|_\infty \leq \|v_0\|_\infty + K\xi B \|u_0\|_{W_{2p}^1(\Omega)} \left[ (\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1 \right]^{1/2p}, \quad t \in (0, T). \quad (3.26)$$

If  $(u_0, v_0, w_0)$  and  $\chi, \xi$  satisfy

$$\chi \leq \frac{1}{4KG_0} \quad \text{and} \quad \xi \leq \frac{1}{4K\|u_0\|_{W_{2p}^1(\Omega)} \left[ (\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1 \right]^{1/2p}},$$

then, according to (3.25) and (3.26), we find

$$\|u(\cdot, t)\|_\infty \leq 3\|u_0\|_\infty/2 < A, \quad \|v(\cdot, t)\|_\infty \leq 3\|v_0\|_\infty/2 < B, \quad t \in (0, T).$$

Due to the continuity of  $u$  and  $v$ , there holds  $T = T_m$  and

$$\|u(\cdot, t)\|_\infty \leq A, \quad \|v(\cdot, t)\|_\infty \leq B, \quad t \in (0, T_m). \quad (3.27)$$

*Step 2: The uniform  $L^{2p}$ -boundedness of  $\nabla v$  in  $(0, T_m)$ .* Thanks to Lemmas 3.1 and 3.3 with  $T$  replaced by  $T_m$ , there exists  $C_1 > 0$  such that

$$\|\nabla w(\cdot, t)\|_{2(p+1)} \leq C_1, \quad t \in (0, T_m), \quad (3.28)$$

and

$$\|\nabla u(\cdot, t)\|_{2p} \leq C_1, \quad t \in (0, T_m); \quad \int_t^{t+\delta} \int_\Omega |\nabla u|^{2(p+1)} dx ds \leq C_1, \quad t \in (0, T_m - \delta), \quad (3.29)$$

where  $\delta = \min\{1, T_m/2\}$ . Moreover, from Lemma 3.2 with  $T$  replaced by  $T_m$ , we find  $C_2 > 0$  such that

$$\int_t^{t+\delta} \int_\Omega |\Delta w|^{p+1} dx ds \leq C_2, \quad t \in (0, T_m - \delta). \quad (3.30)$$

It is easy to see that  $u$  satisfies

$$\begin{cases} u_t = \Delta u + F(x, t), & x \in \Omega, \quad t \in (0, T_m), \\ \partial_\nu u = 0, & x \in \partial\Omega, \quad t \in (0, T_m), \\ u(x, 0) = u_0, & x \in \Omega, \end{cases}$$

where  $F(x, t) = \nabla u \cdot \nabla w + u\Delta w$ . In view of Young's inequality and the first inequality in (3.27), (3.28), the second inequality in (3.29) and (3.30), there is  $C_3 > 0$  fulfilling

$$\int_t^{t+\delta} \int_\Omega |F(x, s)|^{p+1} dx ds \leq C_3, \quad t \in (0, T_m - \delta).$$

Hence, similar to the derivation of Lemma 3.2, one can show that, there exists  $C_4 > 0$  such that

$$\int_t^{t+\delta} \int_\Omega |\Delta u|^{p+1} dx ds \leq C_4, \quad t \in (0, T_m - \delta). \quad (3.31)$$

In view of the  $L^\infty$  boundedness of  $v$  in (3.27), following the arguments in the proof of Lemma 3.3 (cf. (3.22)), for some positive constant  $C_5$ , there holds

$$\frac{d}{dt} \int_\Omega |\nabla v|^{2p} + \int_\Omega |\nabla v|^{2p} \leq C_5 \left( \int_\Omega |\nabla u|^{2(p+1)} + \int_\Omega |\Delta u|^{p+1} + 1 \right),$$

and then by using the second inequality in (3.29) and (3.31) and Lemma 2.2, we have

$$\int_{\Omega} |\nabla v(\cdot, t)|^{2p} \leq C_6, \quad t \in (0, T_m) \quad (3.32)$$

for some  $C_6 > 0$ .

*Step 3: The global existence and boundedness of the solution.* From (2.1), (3.27), (3.28), the first inequality in (3.29) and (3.32), there is  $C_7 > 0$  such that

$$\|u(\cdot, t)\|_{W_{2p}^1(\Omega)} + \|v(\cdot, t)\|_{W_{2p}^1(\Omega)} + \|w(\cdot, t)\|_{W_{2p}^1(\Omega)} \leq C_7, \quad t \in (0, T_m).$$

This enables us to deduce that  $T_m = \infty$  due to  $2p > n$  and Lemma 2.1. Moreover, (3.24) follows directly. This completes the proof.  $\square$

*Proof of Theorems 1.1 and 1.2.* Recalling the definitions of  $G_0, H_0$  and taking

$$\chi_0 = \frac{\kappa}{G_0}, \quad \text{and} \quad \xi_0 = \frac{\kappa}{\|u_0\|_{W_{2p}^1(\Omega)} [(\chi G_0)^{2(p+1)} + (\chi H_0)^{p+1} + 1]^{1/2p}}$$

where  $\kappa = \kappa(\Sigma) > 0$  was given in Lemma 3.6. On the one hand, for the fixed  $\chi, \xi$ , the small values of  $\|w_0\|_{W_{2(p+1)}^1(\Omega)} + r_*$  and  $\|u_0\|_{W_{2p}^1(\Omega)}$  ensure (3.23), then Theorem 1.1 follows from Lemma 3.6 directly. On the other hand, for the given initial data  $(u_0, v_0, w_0)$ , if  $\chi \leq \chi_0$  and  $\xi \leq \xi_0$ , then we can show Theorem 1.2 from Lemma 3.6.  $\square$

## 4 Proof of Theorem 1.3

Define

$$\tilde{T} := \sup \{T \in (0, T_m) : \|v(\cdot, t)\|_{\infty} \leq B \text{ for all } t \in (0, T)\}.$$

Evidently, by the continuity of the solution, we have  $\tilde{T} \in (0, T_m]$  and

$$\|v(\cdot, t)\|_{\infty} \leq B, \quad t \in (0, \tilde{T}). \quad (4.1)$$

For the later use, we denote  $\theta = \min\{1, \tilde{T}/2\}$  and

$$\Gamma = \{\lambda, \mu, r_*, \eta, \chi, \Omega, \|u_0\|_{W_4^1(\Omega)}, \|v_0\|_{\infty}, \|w_0\|_{W_3^2(\Omega)}\}.$$

We first assert the uniform  $L^{\infty}$  boundedness of  $u$  provided (4.1).

**Lemma 4.1.** *Let  $n = 2, \eta_1 > 0, m \geq 2$  and  $\eta_2 = 0$ . Then there is  $C = C(\Gamma) > 0$  such that, the solution of (1.1) satisfies*

$$\int_t^{t+\theta} \int_{\Omega} |\nabla u|^2 dx ds \leq C, \quad t \in (0, \tilde{T} - \theta). \quad (4.2)$$

*Proof.* In the proof, we use  $C_i$  to denote the positive constants which may depend on  $\Gamma$ . Integrating the first equation in (1.1) upon  $\Omega$  yields

$$\frac{d}{dt} \int_{\Omega} u = \eta_1 \int_{\Omega} u - \eta_1 \int_{\Omega} u^m, \quad t \in (0, T_m), \quad (4.3)$$

which implies

$$\int_{\Omega} u \leq \max\{\|u_0\|_1, |\Omega|\} =: C_1, \quad t \in (0, T_m). \quad (4.4)$$

Integrating (4.3) from  $t$  to  $t + \theta$  with  $t \in (0, T_m - \theta)$ , and using the  $L^1$  boundedness of  $u$ , we find

$$\int_t^{t+\theta} \int_{\Omega} u^m dx ds \leq C_1 + \frac{C_1}{\eta_1}, \quad t \in (0, T_m - \theta). \quad (4.5)$$

Due to  $m \geq 2$ , by use of Young's inequality, there is  $C_2 > 0$  such that

$$\int_t^{t+\theta} \int_{\Omega} u^2 dx ds \leq C_2, \quad t \in (0, T_m - \theta). \quad (4.6)$$

Let  $f(x, t) = -\lambda(u + v)w - \mu w + r$ . It follows from (4.1) and (4.6) that, there is  $C_3 > 0$  such that

$$\int_t^{t+\theta} \int_{\Omega} f^2 dx ds \leq C_3, \quad t \in (0, \tilde{T} - \theta).$$

As in the proof of Lemma 3.2, one can show that, there exists  $C_4 > 0$  such that

$$\int_t^{t+\theta} \int_{\Omega} |\Delta w|^2 dx ds \leq C_4, \quad t \in (0, \tilde{T} - \theta). \quad (4.7)$$

Testing the  $u$ -equation in (1.1) by  $u$ , there holds

$$\begin{aligned} \frac{1}{2} \int_{\Omega} u^2 + \int_{\Omega} |\nabla u|^2 &\leq \chi \int_{\Omega} u \nabla u \cdot \nabla w + \eta_1 \int_{\Omega} u^2 \\ &= \frac{\chi}{2} \int_{\Omega} \nabla u^2 \cdot \nabla w + \eta_1 \int_{\Omega} u^2 \\ &= -\frac{\chi}{2} \int_{\Omega} u^2 \Delta w + \eta_1 \int_{\Omega} u^2 \\ &\leq \frac{\chi}{2} \|u\|_4^2 \|\Delta w\|_2 + \eta_1 \int_{\Omega} u^2, \quad t \in (0, T_m). \end{aligned}$$

In view of the Gagliardo-Nirenberg inequality ([13]) and (4.4), there is  $C_5 > 0$  such that

$$\|u\|_4^2 \leq C_5 (\|\nabla u\|_2 \|u\|_2 + \|u\|_1^2) \leq C_5 (\|\nabla u\|_2 \|u\|_2 + C_1^2), \quad t \in (0, T_m).$$

Hence, for some  $C_6 > 0$ ,

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} u^2 + 2 \int_{\Omega} |\nabla u|^2 &\leq \chi C_5 (\|\nabla u\|_2 \|u\|_2 + C_1^2) \|\Delta w\|_2 + 2\eta_1 \int_{\Omega} u^2 \\ &\leq \|\nabla u\|_2^2 + C_6 (\|u\|_2^2 \|\Delta w\|_2^2 + \|u\|_2^2 + \|\Delta w\|_2^2 + 1), \quad t \in (0, T_m). \end{aligned} \quad (4.8)$$

By rearrangement, we have

$$z'(t) + \int_{\Omega} |\nabla u|^2 \leq C_6 \left( \int_{\Omega} u^2 dx + 1 \right) \left( \int_{\Omega} |\Delta w|^2 dx + 1 \right) := C_6 z(t) h(t), \quad t \in (0, \hat{T}), \quad (4.9)$$

where

$$z(t) = \int_{\Omega} |u(\cdot, t)|^2 + 1, \quad h(t) = \int_{\Omega} |\Delta w(\cdot, t)|^2 + 1.$$

For any  $0 \leq \tilde{t} \leq t < \tilde{T}$ , it follows from (4.9) that

$$z(t) \leq z(\tilde{t})e^{C_6 \int_{\tilde{t}}^t h(s)ds}. \quad (4.10)$$

Clearly, for  $t \in [0, \theta]$ , from (4.10) and (4.7) we have

$$z(t) \leq z(0)e^{C_6 \int_0^t h(s)ds} \leq (\|u_0\|_2^2 + 1)e^{C_6(C_4+1)} \quad (4.11)$$

Fix  $t \in (\theta, \tilde{T})$ . Due to (4.6), there exists  $t_0 \in [t - \theta, t]$  such that  $z(t_0) = \int_{\Omega} u^2(\cdot, t_0) + 1 \leq C_2 + 1$ . Thanks to (4.7) we have

$$\int_{t_0}^t h(s)ds = \int_{t_0}^t \left( \int_{\Omega} |\Delta w|^2 dx + 1 \right) ds \leq \int_{t-\theta}^t \left( \int_{\Omega} |\Delta w|^2 dx + 1 \right) ds \leq C_4 + 1.$$

Again by (4.10),

$$z(t) \leq z(t_0)e^{C_6 \int_{t_0}^t h(s)ds} \leq (C_2 + 1)e^{C_6(C_4+1)}, \quad \theta < t < \tilde{T}.$$

This combined with (4.11) implies that  $z(t) = \int_{\Omega} |u(\cdot, t)|^2 + 1 \leq C_7$  in  $(0, \tilde{T})$  with  $C_7 > 0$ . Moreover, it follows from (4.9) that

$$\frac{d}{dt} \int_{\Omega} u^2 + \int_{\Omega} |\nabla u|^2 \leq C_6 C_7 \left( \int_{\Omega} |\Delta w|^2 + 1 \right), \quad t \in (0, \tilde{T}),$$

which by an integration upon  $(t, t + \theta)$  for  $t \in (0, \tilde{T} - \theta)$  implies that, there is  $C_8 > 0$  such that

$$\int_t^{t+\theta} \int_{\Omega} |\nabla u|^2 dx ds \leq C_8, \quad t \in (0, \tilde{T} - \theta).$$

This implies (4.2) and the proof is end.  $\square$

**Lemma 4.2.** *Suppose that  $n = 2$ ,  $\eta_1 > 0$ ,  $m \geq 2$  and  $\eta_2 = 0$ . Then one can find  $C = C(\Gamma) > 0$  such that*

$$\int_{\Omega} |\nabla w(\cdot, t)|^4 \leq C, \quad t \in (0, \tilde{T}). \quad (4.12)$$

*Proof.* By direct calculations (cf. (3.14) with  $p = 2$ ), there holds

$$\begin{aligned} & \frac{1}{4} \frac{d}{dt} \int_{\Omega} |\nabla w|^4 + \mu \int_{\Omega} |\nabla w|^4 \\ &= \int_{\Omega} |\nabla w|^2 \nabla u \cdot \nabla w_t + \mu \int_{\Omega} |\nabla w|^4 \\ &= \int_{\Omega} |\nabla w|^2 \nabla w \cdot \nabla (\Delta w - \lambda(u + v)w - \mu w + r) + \mu \int_{\Omega} |\nabla w|^4 \\ &= \int_{\Omega} |\nabla w|^2 \nabla w \cdot \nabla \Delta w - \lambda \int_{\Omega} |\nabla w|^2 \nabla w \cdot \nabla (uw + vw) + \int_{\Omega} |\nabla w|^2 \nabla w \cdot \nabla r \\ &\leq \int_{\Omega} |\nabla w|^2 \nabla w \cdot \nabla \Delta w - \lambda \int_{\Omega} w |\nabla w|^2 \nabla u \cdot \nabla w - \lambda \int_{\Omega} w |\nabla w|^2 \nabla v \cdot \nabla w + \int_{\Omega} |\nabla w|^2 \nabla w \cdot \nabla r \\ &=: I_1(t) + I_2(t) + I_3(t) + I_4(t), \quad t \in (0, T_m). \end{aligned} \quad (4.13)$$

In view of [7, Lemma 2.2], we have

$$\int_{\Omega} |\nabla w|^6 \leq 36Q^2 \int_{\Omega} |\nabla w|^2 |D^2 w|^2 =: \tilde{k} \int_{\Omega} |\nabla w|^2 |D^2 w|^2, \quad t \in (0, T_m). \quad (4.14)$$



Making use of [16, Lemma 2.6 (ii)], there is  $C_1 > 0$  depending on  $n, p, \Omega$  such that

$$\int_{\partial\Omega} |\nabla w|^2 \partial_\nu |\nabla w|^2 dS \leq \frac{1}{2} \int_{\Omega} |\nabla |\nabla w|^2|^2 + C_1 \int_{\Omega} |\nabla w|^4, \quad t \in (0, T_m).$$

Hence, similar to the derivation of (3.15) with  $p = 2$ , we get

$$\begin{aligned} I_1(t) &= \frac{1}{2} \int_{\Omega} |\nabla w|^2 \Delta |\nabla w|^2 - \int_{\Omega} |\nabla w|^2 |D^2 w|^2 \\ &= -\frac{1}{2} \int_{\Omega} |\nabla |\nabla w|^2|^2 + \frac{1}{2} \int_{\partial\Omega} |\nabla w|^2 \partial_\nu |\nabla w|^2 dS - \int_{\Omega} |\nabla w|^2 |D^2 w|^2 \\ &\leq -\int_{\Omega} |\nabla w|^2 |D^2 w|^2 - \frac{1}{4} \int_{\Omega} |\nabla |\nabla w|^2|^2 + \frac{C_1}{2} \int_{\Omega} |\nabla w|^4, \quad t \in (0, T_m). \end{aligned}$$

Thus, applying Young's inequality and (4.14), there holds

$$\begin{aligned} I_1(t) &\leq -\int_{\Omega} |\nabla w|^2 |D^2 w|^2 - \frac{1}{4} \int_{\Omega} |\nabla |\nabla w|^2|^2 + \frac{1}{4\tilde{k}} \int_{\Omega} |\nabla w|^6 + 2\tilde{k}^2 C_1^3 |\Omega| \\ &\leq -\frac{3}{4} \int_{\Omega} |\nabla w|^2 |D^2 w|^2 - \frac{1}{4} \int_{\Omega} |\nabla |\nabla w|^2|^2 + 2\tilde{k}^2 C_1^3 |\Omega|, \quad t \in (0, T_m). \end{aligned} \quad (4.15)$$

By Young's inequality and (4.14), the second term in (4.13) can be estimated as:

$$\begin{aligned} I_2(t) &\leq \lambda Q \int_{\Omega} |\nabla w|^3 |\nabla u| \leq \frac{1}{4\tilde{k}} \int_{\Omega} |\nabla w|^6 + \tilde{k} \lambda^2 Q^2 \int_{\Omega} |\nabla u|^2 \\ &\leq \frac{1}{4} \int_{\Omega} |\nabla w|^2 |D^2 w|^2 + \tilde{k} \lambda^2 Q^2 \int_{\Omega} |\nabla u|^2, \quad t \in (0, T_m), \end{aligned} \quad (4.16)$$

and, the third term in (4.13) can be estimated as:

$$\begin{aligned} I_3(t) &= \lambda \int_{\Omega} v \nabla \cdot (w |\nabla w|^2 \nabla w) \\ &= \lambda \int_{\Omega} v |\nabla w|^4 + \lambda \int_{\Omega} v w \nabla |\nabla w|^2 \cdot \nabla w + \lambda \int_{\Omega} v w |\nabla w|^2 \Delta w \\ &\leq \lambda B \int_{\Omega} |\nabla w|^4 + \lambda B Q \int_{\Omega} |\nabla |\nabla w|^2| |\nabla w| + \lambda B Q \int_{\Omega} |\nabla w|^2 |\Delta w| \\ &\leq \frac{1}{8} \int_{\Omega} |\nabla |\nabla w|^2|^2 + \frac{1}{4\tilde{k}} \int_{\Omega} |\nabla w|^6 + \int_{\Omega} |\Delta w|^2 + C_2 \\ &\leq \frac{1}{8} \int_{\Omega} |\nabla |\nabla w|^2|^2 + \frac{1}{4} \int_{\Omega} |\nabla w|^2 |D^2 w|^2 + \int_{\Omega} |\Delta w|^2 + C_2, \quad t \in (0, \tilde{T}), \end{aligned} \quad (4.17)$$

where  $C_2 = C_2(\Gamma) > 0$  and we have used (4.1) in the derivation of (4.17). For the last term  $I_4$ , again by the Young inequality and (4.14), we find

$$\begin{aligned} I_4(t) &= -\int_{\Omega} r \nabla w \cdot \nabla |\nabla w|^2 - \int_{\Omega} r |\nabla w|^2 \Delta w \\ &\leq \frac{1}{8} \int_{\Omega} |\nabla |\nabla w|^2|^2 + 2r_*^2 \int_{\Omega} |\nabla w|^2 + \int_{\Omega} |\Delta w|^2 + \frac{r_*^2}{4} \int_{\Omega} |\nabla w|^4 \\ &\leq \frac{1}{8} \int_{\Omega} |\nabla |\nabla w|^2|^2 + \int_{\Omega} |\Delta w|^2 + \frac{1}{4\tilde{k}} \int_{\Omega} |\nabla w|^6 + C_3 \\ &\leq \frac{1}{8} \int_{\Omega} |\nabla |\nabla w|^2|^2 + \int_{\Omega} |\Delta w|^2 + \frac{1}{4} \int_{\Omega} |\nabla w|^2 |D^2 w|^2 + C_3, \quad t \in (0, T_m) \end{aligned} \quad (4.18)$$

for some  $C_3 = C_3(\Gamma) > 0$ . Plugging (4.15)-(4.18) into (4.13), one can find  $C_4 = C_4(\Gamma) > 0$  such that

$$\frac{1}{4} \frac{d}{dt} \int_{\Omega} |\nabla w|^4 + \mu \int_{\Omega} |\nabla w|^4 \leq \tilde{k} \lambda^2 Q^2 \int_{\Omega} |\nabla u|^2 + 2 \int_{\Omega} |\Delta w|^2 + C_4, \quad t \in (0, \tilde{T}).$$

In view of (4.2), (4.7) and Lemma 2.2, we get (4.12). This completes the proof.  $\square$

Now Lemma 4.2 enables us to establish the uniform  $L^\infty$  boundedness of  $u$ .

**Lemma 4.3.** *Let  $n = 2$ ,  $\eta_1 > 0$ ,  $m \geq 2$  and  $\eta_2 = 0$ . Then there exists  $C = C(\Gamma) > 0$  such that*

$$\|u(\cdot, t)\|_\infty \leq C, \quad t \in (0, \tilde{T}). \quad (4.19)$$

*Proof.* Denote  $H(T) = \sup_{t \in (0, T)} \|u(\cdot, t)\|_\infty < \infty$  for  $T \in (0, \tilde{T})$ . Making use of the standard  $L^p$ - $L^q$  estimates for  $(e^{t\Delta})_{t \geq 0}$  ([4, Lemma 3.3]), one can find  $\lambda_1, C_1 > 0$  depending on  $\Omega$  such that

$$\begin{aligned} \|u(\cdot, t)\|_\infty &\leq \|u_0\|_\infty + \chi \int_0^t \|e^{(t-s)\Delta} \nabla \cdot (u \nabla w)\|_\infty ds + \eta_1 \int_0^t \|e^{(t-s)\Delta} (u - u^m)\|_\infty ds \\ &\leq \|u_0\|_\infty + \chi \int_0^t \|e^{(t-s)\Delta} \nabla \cdot (u \nabla w)\|_\infty ds + \eta_1 \int_0^t \|e^{(t-s)\Delta} (u - u^m)_+\|_\infty ds \\ &\leq C_1 + C_1 \chi \int_0^t \left(1 + (t-s)^{-\frac{5}{6}}\right) e^{-\lambda_1(t-s)} \|u \nabla w\|_3 ds, \quad t \in (0, T). \end{aligned} \quad (4.20)$$

By (4.12), (4.4) and the definition of  $H(T)$ , there exists  $C_2 = C_2(\Gamma) > 0$  such that

$$\|u \nabla w\|_3 \leq \|u\|_{12} \|\nabla w\|_4 = \left( \int_{\Omega} u^{12} \right)^{1/12} \|\nabla w\|_4 \leq C_2 H(T)^{11/12}, \quad t \in (0, T).$$

Inserting this into (4.20) yields that, for some  $C_3 = C_3(\Gamma) > 0$ ,

$$\begin{aligned} \|u(\cdot, t)\|_\infty &\leq C_1 + C_1 C_2 \chi H(T)^{11/12} \int_0^t \left(1 + (t-s)^{-\frac{5}{6}}\right) e^{-\lambda_1(t-s)} ds, \\ &\leq C_3 + C_3 H(T)^{11/12}, \quad t \in (0, T), \end{aligned}$$

which implies

$$H(T) \leq C_3 + C_3 H(T)^{11/12}, \quad T \in (0, \tilde{T}).$$

Hence, thanks to the Young inequality, we have  $H(T) \leq C_3^{12} + 12C_3$  for all  $T \in (0, \tilde{T})$ . This combined with the definition of  $H(T)$  finishes the proof.  $\square$

In order to obtain the  $L^\infty$  estimate of  $v$  in  $(0, \tilde{T})$ , we shall need the  $L^4$  regularity of  $\nabla u$ .

**Lemma 4.4.** *Assume that  $n = 2$ ,  $\eta_1 > 0$ ,  $m \geq 2$  and  $\eta_2 = 0$ . Then there exists  $C = C(\Gamma) > 0$  such that*

$$\|\nabla u(\cdot, t)\|_4 \leq C, \quad t \in (0, \tilde{T}); \quad \int_t^{t+\theta} \int_{\Omega} |\nabla u|^6 dx ds \leq C, \quad t \in (0, \tilde{T} - \theta). \quad (4.21)$$

*Proof.* Thanks to (4.1) and (4.19), as in the proof of Lemmas 3.1 and 3.2, there exists  $C_1 = C_1(\Gamma) > 0$  such that

$$\int_{\Omega} |\nabla w(\cdot, t)|^6 \leq C_1, \quad t \in (0, \tilde{T}); \quad \int_t^{t+\theta} \int_{\Omega} |\Delta w|^3 dx ds \leq C_1, \quad t \in (0, \tilde{T} - \theta). \quad (4.22)$$

With (4.19) at hand, following the arguments in the proof of Lemma 3.3 with  $n = p = 2$ , we obtain (cf. (3.22))

$$\frac{d}{dt} \int_{\Omega} |\nabla u|^4 + \int_{\Omega} |\nabla u|^4 + \int_{\Omega} |\nabla u|^2 |D^2 u|^2 \leq C_2 \left( \int_{\Omega} |\nabla w|^6 + \int_{\Omega} |\Delta w|^3 + 1 \right), \quad t \in (0, \tilde{T}) \quad (4.23)$$

for some  $C_2 = C_2(\Gamma) > 0$ . Then, using (4.22) and Lemma 2.2, we get the first estimate in (4.21).

Integrating (4.23) from  $t$  to  $t + \theta$  for  $t \in (0, \tilde{T} - \theta)$  firstly, and then applying the first inequality of (4.21) and (4.22) secondly, there is  $C_3 = C_3(\Gamma) > 0$ ,

$$\int_t^{t+\theta} \int_{\Omega} |\nabla u|^2 |D^2 u|^2 dx ds \leq C_3, \quad t \in (0, \tilde{T} - \theta).$$

This in conjunction with (3.12) with  $n = p = 2$  and (4.19) implies the second inequality of (4.21).  $\square$

We are now in the position to get the  $L^\infty$  estimate of  $v$  in  $(0, \tilde{T})$ .

**Lemma 4.5.** *Let  $n = 2$ ,  $\eta_1 > 0$ ,  $m \geq 2$  and  $\eta_2 = 0$ . Then there is  $C = C(\Gamma) > 0$  such that*

$$\|v(\cdot, t)\|_\infty \leq \|v_0\|_\infty + C\xi, \quad t \in (0, \tilde{T}). \quad (4.24)$$

*Proof.* Again by the standard  $L^p$ - $L^q$  estimates for  $(e^{t\Delta})_{t \geq 0}$  ([4, Lemma 3.3]), there exist positive constants  $\lambda_1, C_1$  depending on  $\Omega$  such that

$$\begin{aligned} \|v(\cdot, t)\|_\infty &\leq \|v_0\|_\infty + \xi \int_0^t \|e^{(t-s)\Delta} \nabla \cdot (v \nabla u)\|_\infty ds \\ &\leq \|v_0\|_\infty + C_1 \xi \int_0^t \left(1 + (t-s)^{-\frac{3}{4}}\right) e^{-\lambda_1(t-s)} \|v \nabla u\|_4 ds, \quad t \in (0, T_m). \end{aligned}$$

In view of (4.1) and the first inequality of (4.21), there exists  $C_2 = C_2(\Gamma) > 0$  such that

$$\|v(\cdot, \sigma) \nabla u(\cdot, \sigma)\|_4 \leq \|v(\cdot, \sigma)\|_\infty \|\nabla u(\cdot, \sigma)\|_4 \leq C_2, \quad \sigma \in (0, \tilde{T}).$$

Hence, one can find  $C_3 = C_3(\Gamma)$  such that (4.24) holds.  $\square$

*Proof of Theorem 1.3.* From Lemma 4.5, we can see that, there is  $K_1 = K_1(\Gamma) > 0$  such that, for any  $\xi > 0$ ,

$$\|v(\cdot, t)\|_\infty \leq \|v_0\|_\infty + K_1(\Gamma)\xi, \quad t \in (0, \tilde{T}). \quad (4.25)$$

Noting that  $K_1(\Gamma)$  is independent of  $\xi$ . Hence, if

$$\xi \leq \frac{\|v_0\|_\infty}{2K_1(\Gamma)},$$

then we have by (4.25) that  $\|v(\cdot, t)\|_\infty \leq 3\|v_0\|_\infty/2 < B$  for all  $t \in (0, \tilde{T})$ . Therefore, it follows from the definition of  $\tilde{T}$  that  $\tilde{T} = T_m$ , and  $\|v(\cdot, t)\|_\infty \leq B$  for all  $t \in (0, T_m)$ . Moreover, Lemmas 4.1-4.5 holds with  $\tilde{T}$  replaced by  $T_m$ . As in the proof of Lemma 3.6, one can show that  $\|\nabla v(\cdot, t)\|_4 < \infty$  for any  $t \in (0, T_m)$ , and hence

$$\|u(\cdot, t)\|_{W_4^1(\Omega)} + \|v(\cdot, t)\|_{W_4^1(\Omega)} + \|w(\cdot, t)\|_{W_4^1(\Omega)} < \infty, \quad t \in (0, T_m).$$

Thus, we have  $T_m = \infty$  due to Lemma 2.1. The proof Theorem 1.3 is completed.  $\square$

## 5 Proof of Theorem 1.4

Let  $\delta = \min\{1, T_m/2\}$ . Similar to the derivation of (4.4) and (4.5), we get space-time  $L^m$  (res.  $L^l$ ) regularity for  $u$  (res.  $v$ ).

**Lemma 5.1.** *Let  $n = 2$ . Suppose that  $\eta_1, \eta_2 > 0$  and  $m, l$  satisfy (1.7). Then, there is  $C > 0$  such that*

$$\int_{\Omega} u \leq C, \quad \int_{\Omega} v \leq C, \quad t \in (0, T_m), \quad (5.1)$$

and

$$\int_t^{t+\delta} \int_{\Omega} u^m dx ds \leq C, \quad \int_t^{t+\delta} \int_{\Omega} v^l dx ds \leq C, \quad t \in (0, T_m - \delta). \quad (5.2)$$

Next we derive the space-time  $L^2$  estimate for  $\nabla u$ .

**Lemma 5.2.** *Let  $n = 2$ ,  $\eta_1, \eta_2 > 0$  and  $m, l$  satisfy (1.7). Then, one can find  $C > 0$  such that*

$$\int_t^{t+\delta} \int_{\Omega} |\nabla u|^2 dx ds \leq C, \quad t \in (0, T_m - \delta). \quad (5.3)$$

*Proof.* Noting that (5.2) holds with  $m, l \geq 2$ , there is  $C_1 > 0$  such that

$$\int_t^{t+\delta} \int_{\Omega} |f|^{\tilde{m}} dx ds \leq C_1, \quad t \in (0, T_m - \delta).$$

where  $f(x, t) = -\lambda(u+v)w - \mu w + r$  and  $\tilde{m} = \min\{m, l\} \geq 2$ . As in the proof of Lemma 3.2, there exists  $C_2 > 0$  such that

$$\int_t^{t+\delta} \int_{\Omega} |\Delta w|^{\tilde{m}} dx ds \leq C_2, \quad t \in (0, T_m - \delta). \quad (5.4)$$

Hence, making use of the Young inequality, we have

$$\int_t^{t+\delta} \int_{\Omega} |\Delta w|^2 dx ds \leq C_3, \quad t \in (0, T_m - \delta) \quad (5.5)$$

for some  $C_3 > 0$ . Due to  $m \geq 2$  and the estimation for  $u$  in (5.2), the Young inequality says that, for some  $C_4 > 0$ ,

$$\int_t^{t+\delta} \int_{\Omega} u^2 dx ds \leq C_4, \quad t \in (0, T_m - \delta). \quad (5.6)$$

With the  $L^1$ -estimation of  $u$  in (5.1), (5.5) and (5.6) at hand, following the line of the proof of Lemma 4.1, we get (5.3).  $\square$

The following lemma asserts the uniform-in-time  $L^4$  boundedness of  $\nabla w$ .

**Lemma 5.3.** *Let  $n = 2$ ,  $\eta_1, \eta_2 > 0$  and  $m, l$  satisfy (1.7). Then, one can find  $C > 0$  such that*

$$\int_{\Omega} |\nabla w(\cdot, t)|^4 \leq C, \quad t \in (0, T_m). \quad (5.7)$$

*Proof.* The arguments are same with the proof of Lemma 4.2 except the estimation of  $I_3$  in (4.17). By use of Young's inequality, (4.14) and (1.7), we find

$$\begin{aligned}
I_3(t) &= \lambda \int_{\Omega} v \nabla \cdot (w |\nabla w|^2 \nabla w) \\
&= \lambda \int_{\Omega} v |\nabla w|^4 + \lambda \int_{\Omega} v w \nabla |\nabla w|^2 \cdot \nabla w + \lambda \int_{\Omega} v w |\nabla w|^2 \Delta w \\
&\leq \lambda \int_{\Omega} v |\nabla w|^4 + \lambda Q \int_{\Omega} v |\nabla |\nabla w|^2| |\nabla w| + \lambda Q \int_{\Omega} v |\nabla w|^2 |\Delta w| \\
&\leq (12\tilde{k})^2 \lambda^3 \int_{\Omega} v^3 + \frac{1}{6\tilde{k}} \int_{\Omega} |\nabla w|^6 + \frac{1}{8} \int_{\Omega} |\nabla |\nabla w|^2|^2 + 2\lambda^2 Q^2 \int_{\Omega} v^2 |\nabla w|^2 \\
&\quad + (12\tilde{k})^{1/2} (\lambda Q)^{3/2} \int_{\Omega} v^{3/2} |\Delta w|^{3/2} \\
&\leq \left[ (12\tilde{k})^2 \lambda^3 + (12\tilde{k})^{1/2} 2\sqrt{2} \lambda^3 Q^3 \right] \int_{\Omega} v^3 + \frac{1}{4\tilde{k}} \int_{\Omega} |\nabla w|^6 + \frac{1}{8} \int_{\Omega} |\nabla |\nabla w|^2|^2 \\
&\quad + C_1 \int_{\Omega} v^{\frac{3\tilde{m}}{2\tilde{m}-3}} + C_1 \int_{\Omega} |\Delta w|^{\tilde{m}} \\
&\leq \frac{1}{8} \int_{\Omega} |\nabla |\nabla w|^2|^2 + \frac{1}{4} \int_{\Omega} |\nabla w|^2 |D^2 w|^2 + C_1 \int_{\Omega} |\Delta w|^{\tilde{m}} + C_2 \int_{\Omega} v^l + C_2, \quad t \in (0, T_m). \quad (5.8)
\end{aligned}$$

Inserting (4.15), (4.16), (5.8) and (4.18) into (4.13), one can find  $C_3 > 0$  such that

$$\frac{1}{4} \frac{d}{dt} \int_{\Omega} |\nabla w|^4 + \mu \int_{\Omega} |\nabla w|^4 \leq C_3 \left( \int_{\Omega} |\nabla u|^2 + \int_{\Omega} |\Delta w|^{\tilde{m}} + \int_{\Omega} |\Delta w|^2 + \int_{\Omega} v^l + 1 \right), \quad t \in (0, T_m).$$

Thanks to the second inequality in (5.2), (5.3), (5.4), (5.5) and Lemma 2.2, we get (5.7). This completes the proof.  $\square$

Noting that, we only use the uniform-in-time  $L^1$  regularity of  $u$  and  $L^4$  estimate of  $\nabla w$  in the proof of Lemma 4.3. Hence, by the similar arguments, we obtain the uniform-in-time  $L^\infty$  boundedness of  $u$ .

**Lemma 5.4.** *Assume that  $n = 2$ ,  $\eta_1, \eta_2 > 0$  and  $m, l$  satisfy (1.7). Then, there is  $C > 0$  such that*

$$\|u(\cdot, t)\|_{\infty} \leq C, \quad t \in (0, T_m). \quad (5.9)$$

*Proof.* Following the line in the proof of Lemma 4.3, using the  $L^1$ -estimate of  $u$  in (5.1) and (5.7), one can easily prove this lemma.  $\square$

The coming lemma provides the  $L^4$  estimate for  $\nabla u$ .

**Lemma 5.5.** *Assume that  $n = 2$ ,  $\eta_1, \eta_2 > 0$  and  $m, l$  satisfy (1.7). Then, there is  $C > 0$  such that*

$$\int_{\Omega} |\nabla u(\cdot, t)|^4 \leq C, \quad t \in (0, T_m), \quad (5.10)$$

and

$$\int_t^{t+\delta} \int_{\Omega} |\nabla u(\cdot, t)|^6 \leq C, \quad t \in (0, T_m - \delta). \quad (5.11)$$

*Proof.* By (5.9) and the second estimate in (5.2) with  $l \geq 3$  as well as Young's inequality, there is  $C_1 > 0$  such that

$$\int_t^{t+\delta} \int_{\Omega} |f(x, s)|^3 dx ds \leq C_1, \quad t \in (0, T_m - \delta).$$

where  $f = -\lambda(u + v)w - \mu w + r$ . As in the proof of Lemma 3.2, we find  $C_2 > 0$  such that

$$\int_t^{t+\delta} \int_{\Omega} |\Delta w|^3 dx ds \leq C_2, \quad t \in (0, T_m - \delta). \quad (5.12)$$

In view of the Gagliardo-Nirenberg inequality, there holds

$$\begin{aligned} \int_{\Omega} |\nabla w(\cdot, t)|^6 &= \|\nabla w(\cdot, t)\|_6^6 \leq C_3 (\|\Delta w(\cdot, t)\|_3^3 \|w(\cdot, t)\|_{\infty}^3 + \|w(\cdot, t)\|_{\infty}^6) \\ &\leq C_3 Q^3 \int_{\Omega} |\Delta w(\cdot, t)|^3 + C_3 Q^6, \quad t \in (0, T_m) \end{aligned}$$

for some  $C_3 > 0$ . This combined with (5.12) yields that, there is  $C_4 > 0$  such that

$$\int_t^{t+\delta} \int_{\Omega} |\nabla w|^6 dx ds \leq C_4, \quad t \in (0, T_m - \delta). \quad (5.13)$$

In view of (5.9), following the proof of Lemma 3.3 with  $n = p = 2$ , we get (cf. (3.22))

$$\begin{aligned} &\frac{d}{dt} \int_{\Omega} |\nabla u(\cdot, t)|^4 + \int_{\Omega} |\nabla u(\cdot, t)|^4 + \int_{\Omega} |\nabla u(\cdot, t)|^2 |D^2 u(\cdot, t)|^2 \\ &\leq C_5 \left( \int_{\Omega} |\nabla w(\cdot, t)|^6 + \int_{\Omega} |\Delta w(\cdot, t)|^3 + 1 \right) := G(t), \quad t \in (0, T_m) \end{aligned} \quad (5.14)$$

with some  $C_5 > 0$ . Due to (5.12) and (5.13), it is easy to see that, there exists  $C_6 > 0$  such that

$$\int_t^{t+\delta} G(s) ds \leq C_6, \quad t \in (0, T_m - \delta). \quad (5.15)$$

Making use of Lemma 2.2 with (5.14) and (5.15), we have

$$\int_{\Omega} |\nabla u(\cdot, t)|^4 \leq C_7, \quad t \in (0, T_m).$$

This shows (5.10). Moreover, integrating (5.14) from  $t$  to  $t + \delta$  for  $t \in (0, T_m - \delta)$  firstly and using (5.10), (5.15) secondly, it follows that, for some  $C_7 > 0$ ,

$$\int_t^{t+\delta} \int_{\Omega} |\nabla u|^2 |D^2 u|^2 dx ds \leq C_7, \quad t \in (0, T_m - \delta).$$

Finally, by (3.12) with  $n = p = 2$  and (5.9), we get (5.11).  $\square$

In light of (5.10), we show the uniform-in-time boundedness of  $v$ .

**Lemma 5.6.** *Let  $n = 2$ ,  $\eta_1, \eta_2 > 0$  and  $m, l$  satisfy (1.7). Then, there exists  $C > 0$  such that*

$$\|v(\cdot, t)\|_{\infty} \leq C, \quad t \in (0, T_m). \quad (5.16)$$

*Proof.* Thanks to uniform-in-time  $L^4$  estimate for  $\nabla u$  in (5.10) and the  $L^1$  regularity for  $v$  in (5.1), parallel to the arguments in the proof of Lemma 4.3, we easily get the desired result.  $\square$

*Proof of Theorem 1.4.* We first note that,  $u$  satisfies

$$\begin{cases} u_t = \Delta u + F(x, t), & x \in \Omega, \quad t \in (0, T_m), \\ \partial_\nu u = 0, & x \in \partial\Omega, \quad t \in (0, T_m), \\ u(x, 0) = u_0, & x \in \Omega, \end{cases}$$

where  $F(x, t) = \nabla u \cdot \nabla w + u\Delta w + \eta_1(u - u^m)$ . In view of Young's inequality and (5.9), (5.11), (5.12) and (5.13), there exists  $C_1 > 0$  such that  $\int_t^{t+\delta} \int_\Omega |F(x, s)|^3 dx ds \leq C_1$  for all  $t \in (0, T_m - \delta)$ . Similar to the proof of Lemma 3.2, we get that, for some  $C_2 > 0$ ,

$$\int_t^{t+\delta} \int_\Omega |\Delta u|^3 dx ds \leq C_2, \quad t \in (0, T_m - \delta). \quad (5.17)$$

With (5.16) at hand, by using the similar arguments in the proof of Lemma 3.3 with  $n = p = 2$  (cf. (3.22)), one can obtain that, for  $C_3 > 0$ ,

$$\frac{d}{dt} \int_\Omega |\nabla v|^4 + \int_\Omega |\nabla v|^4 \leq C_3 \left( \int_\Omega |\nabla u|^6 + \int_\Omega |\Delta u|^3 + 1 \right).$$

This in conjunction with (5.11) and (5.17) and Lemma 2.2, we have

$$\int_\Omega |\nabla v(\cdot, t)|^4 \leq C_4, \quad t \in (0, T_m) \quad (5.18)$$

for some  $C_4 > 0$ . From the second estimate in (2.1), (5.7), (5.9), (5.10), (5.16) and (5.18), there exists  $C_5 > 0$  such that

$$\|u(\cdot, t)\|_{W_4^1(\Omega)} + \|v(\cdot, t)\|_{W_4^1(\Omega)} + \|w(\cdot, t)\|_{W_4^1(\Omega)} \leq C_5, \quad t \in (0, T_m).$$

It follows from Lemma 2.1 that  $T_m = \infty$ . This completes the proof.  $\square$

## References

- [1] T. Black, *Global generalized solutions to a forager-exploiter model with superlinear degradation and their eventual regularity properties*. arXiv:1911.02321v2, 2019.
- [2] X. Cao, *Global bounded solutions of the higher-dimensional Keller-Segel system under smallness conditions in optimal spaces*, Discrete Contin. Dyn. Syst. Ser. A, 35(2015), 1891-1904.
- [3] L. Corrias and B. Perthame, *Asymptotic decay for the solutions of the parabolic-parabolic Keller-Segel chemotaxis system in critical spaces*, Math. Comput. Model., 47(2008), 755-764.
- [4] K. Fujie, A. Ito, M. Winkler and T. Yokota, *Stabilization in a chemotaxis model for tumor invasion*, Discrete Contin. Dyn. Syst., 36(2016), 151-169.
- [5] Y. Giga and H. Sohr, *Abstract  $L^p$  estimates for the Cauchy problem with applications to the Navier-Stokes equations in exterior domains*, J. Funct. Anal., 102(1991), 72-94.
- [6] H. Jin and Z. Wang, *Global stability of prey-taxis systems*, J. Differential Equations, 262(2017), 1257-1290.
- [7] J. Lankeit and Y.L. Wang, *Global existence, boundedness and stabilization in a high-dimensional chemotaxis system with consumption*, Discrete Contin. Dyn. Syst., 37(12)(2017), 6099-6121.

- [8] Y. Liu, *Global existence and boundedness of classical solutions to a forager-exploiter model with volume-filling effects*, Nonlinear Anal.: Real World Appl., 50(2019), 519-531.
- [9] Y. Lou and M. Winkler, *Global existence and uniform boundedness of smooth solutions to a cross-diffusion system with equal diffusion rates*, Comm. Partial Differential Equations, 40(10)(2015), 1905-1941.
- [10] K. Osaki, T. Tsujikawa, A. Yagi and M. Mimura, *Exponential attractor for a chemotaxis growth system of equations*, Nonlinear Anal., 51 (2002), 119-144.
- [11] C. Stinner, C. Surulescu and M. Winkler, *Global weak solutions in a PDE-ODE system modeling multiscale cancer cell invasion*, SIAM J. Math. Anal., 46(2014), 1969-2007.
- [12] N. Tania, B. Vanderlei, J.P. Heath and L. Edelstein-Keshet, *Role of social interactions in dynamic patterns of resource patches and forager aggregation*, Proc. Natl. Acad. Sci. USA, 109(2012), 11228-11233.
- [13] Y. Tao and Z. Wang, *Competing effects of attraction vs. repulsion in chemotaxis*, Math. Models Methods Appl. Sci., 23(1)(2012), 1-36.
- [14] Y. Tao and M. Winkler, *Large time behavior in a forager-exploiter model with different taxis strategies for two groups in search of food*, Math. Models Methods Appl. Sci., 29(11)(2019), 2151-2182.
- [15] J. Wang and M. Wang, *Boundedness and global stability of the two-predator and one-prey models with nonlinear prey-taxis*, Z. Angew. Math. Phys., 69(3)(2018), 63(1-24).
- [16] J. Wang and M. Wang, *Global solution of a diffusive predator-prey model with prey-taxis*, Comput. Math. Appl., 77(2019), 2676-2694.
- [17] J. Wang and M. Wang, *Boundedness in the higher-dimensional Keller-Segel model with signal-dependent motility and logistic growth*, J. Math. Phys., 60(2019), 011507.
- [18] M. Winkler, *Aggregation vs. global diffusive behavior in the higher-dimensional Keller-Segel model*, J. Differential Equations, 248(12)(2010), 2889-2905.
- [19] M. Winkler, *Global generalized solutions to a multi-dimensional doubly tactic resource consumption model accounting for social interactions*, Math. Models Methods Appl. Sci., 29(3)(2019), 373-418.
- [20] S. Wu, J. Shi and B. Wu, *Global existence of solutions and uniform persistence of a diffusive predator-prey model with prey-taxis*, J. Differential Equations, 260(2016), 5847-5874.
- [21] T. Xiang, *Sub-logistic source can prevent blowup in the 2D minimal Keller-Segel chemotaxis system*, J. Math. Phys., 59(2018), 081502.
- [22] T. Xiang and J. Zheng, *A new result for 2D boundedness of solutions to a chemotaxis-haptotaxis model with/without sub-logistic source*, Nonlinearity, 32(2019), 4890-4911.