

Constructions of Pairs of Orthogonal Latin Cubes*

Vladimir N. Potapov
Sobolev Institute of Mathematics, Novosibirsk, Russia
vpotapov@math.nsc.ru

Abstract

A pair of orthogonal latin cubes of order q is equivalent to an MDS code with distance 3 or to an $\text{OA}_1(3, 5, q)$ orthogonal array. We construct pairs of orthogonal latin cubes for a sequence of previously unknown orders $q_i = 16(18i - 1) + 4$ and $q'_i = 16(18i + 5) + 4$. The minimal new obtained parameters of orthogonal arrays are $\text{OA}_1(3, 5, 84)$.

Keywords— latin square, latin cube, MOLS, MDS code, block design, Steiner system, orthogonal array

MSC2010: 05B15, 94B05, 05B05

1 Introduction

A *latin square* of order q is a $q \times q$ array of q symbols where each symbol occurs exactly once in every row and in every column. A k -dimensional array satisfying the same condition is called a *latin k -cube*. Any 2-dimensional axis-aligned plane (face) of a latin k -cube of order q is a latin square of order q by definition. Two latin squares are *orthogonal* if, when they are superimposed, every ordered pair of symbols appears exactly once. For brevity, a pair of orthogonal latin squares is called POLS. If in a set of latin squares, any two latin squares are orthogonal then this set is called a system of Mutually Orthogonal Latin Squares (MOLS). Two latin k -cubes are *orthogonal* if any pair of corresponding 2-dimensional faces of these cubes is a POLS. Bose, Shrikhande and Parker [1] proved that for each positive integer q , $q \neq 2, 6$, there exists POLS of order q and POLS of orders 2 and 6 are not exist. As a corollary we obtain nonexistence of pairs of orthogonal latin k -cubes of orders 2, 6. A nonexistence of pairs of orthogonal latin k -cubes of orders q if $k > q - 1$ follows from the sphere-packing (Hamming) bound. But the complete spectrum of possible orders of pairs of orthogonal latin k -cubes remains unknown for any $k \geq 3$. Ten is the minimum order for which it is not known whether pairs of orthogonal latin 3-cubes exist. In this paper we construct pairs of orthogonal latin 3-cubes for a sequence of previously unknown orders $q_i = 16(18i - 1) + 4$ and $q'_i = 16(18i + 5) + 4$. New pairs of orthogonal latin 3-cubes are created as files. These files are available on the website <https://iee-dataport.org/open-access/graeco-latin-cubes>. The minimum new order for a pair of orthogonal latin cubes obtained by proposed construction is 84.

Let $Q_q = \{0, \dots, q - 1\}$. A subset C of Q_q^d is called an *MDS(t, d, q) code* (of order q , code distance $t + 1$ and length d) if $|C \cap \Gamma| = 1$ for each t -dimensional axis-aligned plane Γ . Ethier and Mullen [3] proved that *MDS($2, 2 + s, q$)* codes are equivalent to pairs of orthogonal latin s -cubes of order q . There are two well-known methods for constructing MDS codes. If q is a prime power,

* The research was carried out at the Sobolev Institute of Mathematics at the expense of the Russian Science Foundation 18-11-00136.

then we can consider Q_q as the Galois field $GF(q)$. MDS codes obtained as the solution of an appropriate system of linear equations over $GF(q)$ are known as Reed–Solomon codes. If there exists an $MDS(t, d, p_1)$ code and an $MDS(t, d, p_2)$ code, then we get an $MDS(t, d, p_1 p_2)$ code by a product construction (McNeish’s theorem). We represent a new construction of $MDS(2, 5, q)$ codes that is similar to Wilson’s construction for pairs of orthogonal latin squares with aligned subsquares (see, [2] and [5]).

The problem of existence of MDS codes with non-prime-power orders is connected to the problem of existence of Steiner block designs. By methods of random graph theory Keevash [6] and Glock et al. [4] proved that the natural divisibility conditions are sufficient for existence of Steiner system $S(t, k, n)$ apart from a finite number of exceptional n for given fixed t and k . It is not difficult to prove that any MDS code is equivalent to a transversal in an appropriate multipartite hypergraph (see [9]). Then the existence of $MDS(t, d, q)$ codes follows from [7] (Theorem 1.7) apart from a finite number of exceptional q for given fixed t and d . In the last section of this paper we propose a construction of pairs of orthogonal latin 3-cubes based on Steiner block designs.

Note that an $MDS(2, q+1, q)$ code (a pair of orthogonal $(q-1)$ -cubes) is an 1-error correcting perfect code. The existence of such codes is a well-known problem if q is not a prime power (see [8]).

2 Connection between MDS codes and orthogonal systems

An $OA_\lambda(s, d, q)$ orthogonal array is a $\lambda q^s \times d$ array whose entries are from Q_q such that in every subset of s columns of the array, every s -tuple from Q_q^s appears in exactly λ rows. Further we consider only orthogonal arrays with $\lambda = 1$. In this case every column of the orthogonal array is a function $f : Q_q^s \rightarrow Q_q$. A set of columns of an orthogonal array with $\lambda = 1$ is called an orthogonal system. In other words, a system consisting of d functions f_1, \dots, f_d , $f_i : Q_q^s \rightarrow Q_q$ ($d \geq s$) is *orthogonal* if for each subsystem f_{i_1}, \dots, f_{i_s} consisting of s functions it holds

$$\{(f_{i_1}(\bar{x}), \dots, f_{i_s}(\bar{x})) \mid \bar{x} \in Q_q^s\} = Q_q^s.$$

If the system remains orthogonal after substitution any constants for each subset of variables, then it is called *strong-orthogonal*. If the number of variables is two, then such system is a system of MOLS (see [3]). If $s = 3$, it is a set of Mutually Orthogonal Latin Cubes (MOLC). Ethier and Mullen [3] proved that MDS codes are equivalent to strong-orthogonal systems. Moreover, by a replacement of variables it is possible to obtain a strong-orthogonal system consisting of $d - s$ functions from any orthogonal system consisting of d functions over Q_q^s .

Proposition 1 *The following conditions are equivalent:*

- 1) *a system consisting of t functions f_1, \dots, f_t , $f_i : Q_q^s \rightarrow Q_q$ is strong-orthogonal;*
- 2) *the set $C = \{(x_1, \dots, x_s, f_1(\bar{x}), \dots, f_t(\bar{x})) : x_i \in Q_q\}$ is an $MDS(t, t + s, q)$ code;*
- 3) *the array consisting of all elements of C as rows is an $OA_1(s, t + s, q)$ orthogonal array.*

A projection (punctured code) of an $MDS(t, t + s, q)$ code onto a hyperplane is equal to a removal of one of the functions f_i . The punctured code is an $MDS(t - 1, t + s - 1, q)$ code by Proposition 1. Consequently, an existence of $MDS(2, 2 + s, q)$ code or a pair of orthogonal latin s -cubes of order q follows from an existence of an $MDS(t, t + s, q)$ code if $t \geq 2$.

Sometimes the terms “latin cube” and “ t mutually orthogonal latin cubes” is used for $OA_q(2, 4, q)$ and $OA_q(2, t + 3, q)$ orthogonal arrays respectively. It is easy to see that our definition of a system of MOLC is stronger.

3 Constructions of MDS codes

The Hamming distance ρ between two elements of Q_q^d is the number of positions at which the corresponding symbols are different. In this paper we use only the Hamming distance. The code distance of $C \subset Q_q^d$ is $\rho(C) = \min_{x \in C, y \in C, x \neq y} \rho(x, y)$. The distance between two subsets $A, B \subset Q_q^d$ is $\min_{x \in A, y \in B} \rho(x, y)$. The Singleton bound for the cardinality of a code $C \subset Q_q^d$ with distance $t + 1$ is $|C| \leq q^{d-t}$. MDS codes achieve equality in this bound.

Proposition 2 *A subset $C \subset Q_q^d$ with code distance $t + 1$ is an MDS code if and only if $|C| = q^{d-t}$.*

The Hamming bound for the cardinality of code $C \subset Q_q^d$ with distance 3 is $|C| \leq \frac{q^d}{1+(q-1)d}$. Then the inequalities $q^{d-2} \leq \frac{q^d}{1+(q-1)d}$ or $d \leq q + 1$ are a necessary condition for the existence of an $MDS(2, d, q)$ code. Consequently, an $MDS(2, 5, 3)$ code or a pair of orthogonal latin cubes of order 3 do not exist. Moreover, by puncturing codes we have a necessary condition $s \leq q - 1$ for the existence of an $MDS(t, t + s, q)$ if $t \geq 2$. For linear codes this condition $s \leq q - 1$ is in [8].

Let q be a prime power and let $Q_q = GF(q)$. A linear k -dimensional subspace $C \subset Q_q^d$ with distance t is called $[d, k, t]$ code over $GF(q)$. By Proposition 2 we see that any $[d, d - t, t + 1]$ code over $GF(q)$ is an $MDS(t, d, q)$ code. By using a well-known construction of a linear MDS code (see [8], Chapters 10,11, or [4], Theorem 9.1) by means of an appropriate parity-check matrix over $GF(q)$ we can conclude that the following proposition is true.

Proposition 3 *Let q be a prime power. Then for each integers $d \leq q + 1$ and ϱ , $3 \leq \varrho < d$, there exists a linear (over $GF(q)$) MDS code $C \subset Q_q^d$ with code distance ϱ .*

We will say that an $MDS(t, d, q)$ code M_0 is a *super* $MDS(t, d, q)$ code if there exist $MDS(t + 1, d, q)$ code M_1 and $MDS(t + 2, d, q)$ code M_2 such that $M_2 \subset M_1 \subset M_0$.

By removal of any row from a parity-check matrix of a linear MDS code with distance $t + 1$, we obtain a parity-check matrix of an MDS code with distance t that contains the original code. Thus Propositions 4 and 5 follow from Proposition 3.

Proposition 4 *Let q be a prime power. Then for each integers $d \leq q + 1$ and ϱ , $3 \leq \varrho < d - 2$, there exists a linear over $GF(q)$ super MDS code $C \subset Q_q^d$ with code distance ϱ .*

Proposition 5 *Let q be a prime power. Then for each integers $d \leq q + 1$ and ϱ , $3 \leq \varrho < d - 1$, there exists a linear over $GF(q)$ MDS code $C \subset Q_q^d$ with code distance ϱ that is an union of q disjoint linear over $GF(q)$ MDS code $C \subset Q_q^d$ with code distance $\varrho + 1$.*

The set $Q_{q_1 q_2}$ can be considered as the Cartesian product $Q_{q_1} \times Q_{q_2}$. Consequently, we can identify $Q_{q_1}^d \times Q_{q_2}^d$ and the hypercube $Q_{q_1 q_2}^d$. Thus if $C_1 \subset Q_{q_1}^d$ and $C_2 \subset Q_{q_2}^d$ then

$$C_1 \times C_2 = \{((x_1, y_1), (x_2, y_2), \dots, (x_d, y_d)) : (x_1, \dots, x_d) \in C_1, (y_1, \dots, y_d) \in C_2\} \subset Q_{q_1 q_2}^d.$$

Proposition 6 (McNeish) *Suppose M_1 is an (super) $MDS(t, d, q_1)$ code and M_2 is an (super) $MDS(t, d, q_2)$ code. Then $M_1 \times M_2$ is an (super) $MDS(t, d, q_1 q_2)$ code.*

By combining results of Propositions 3 and 6 we obtain that $MDS(2, 5, q)$ codes exist if $q = 2^{\delta_2} 3^{\delta_3} 5^{\delta_5} \dots$, where $\delta_2 \neq 1$ and $\delta_3 \neq 1$.

Let $A \subset Q_q$. Denote by π_A a function mapping from $Q_p \times Q_q$ to $Q_{p(q-|A|)+|A|}$ by the following rule: $\pi_A(x, y) = (x, y)$ if $y \notin A$, and $\pi_A(x, y) = y$ if $y \in A$. Let $C_1 \subset Q_p^d$ and $C_2 \subset Q_q^d$. Denote $C_1 \times_A C_2 = \{(\pi_A(z_1), \dots, \pi_A(z_d)) : \bar{z} \in C_1 \times C_2\}$. For any $C \subset Q_q^d$ we denote by $U_t(C)$ the t -neighborhood of C , i.e., $U_t(C) = \{x \in Q_q^d : \exists y \in C, \rho(x, y) \leq t\}$.

A set $D \subset Q_q^d$ is called an $MDS(t, d, q)$ with j - A -hole ($t+1 \leq j \leq d$) if

- 1) the code distance of D is equal to $t+1$;
- 2) $D \cap U_{d-j+t}(A^d) = \emptyset$;
- 3) $U_{d-j+t}(D) = Q_q^d \setminus A^d$;
- 4) $|D| = \sum_{k=0}^{j-t-1} \binom{d-t}{k} (q-|A|)^{d-t-k} |A|^k$.

For $t=2$ and $d=5$ we get that an $MDS(2, 5, q)$ code with 5- A -hole has cardinality $q^3 - |A|^3$ and an $MDS(2, 5, q)$ code with 4- A -hole has cardinality $(q-|A|)^3 + 3(q-|A|)^2|A|$.

Suppose that M is an $MDS(t, d, q)$ code, $a \in Q_q$, and $\bar{a} = (a, \dots, a) \in M$. It is easy to see that $M \setminus \{\bar{a}\}$ is an $MDS(t, d, q)$ with d - $\{a\}$ -hole. Let $A \subset Q_q$ and let M be an MDS code, $M \subset Q_q^d$. A subset $M \cap A^d$ is called a *subcode* of M if it is an MDS code in A^d with the same code distance as M . If $M \cap A^d$ is a subcode, then $M \setminus A^d$ is an $MDS(t, d, q)$ with d - A -hole.

Let us formulate a known construction of a POLS (see [2], Chapter 4) in introduced terms.

Proposition 7 *Suppose that*

- $M_1 \subset M$ is $MDS(3, 4, p)$ code and M is $MDS(2, 4, p)$ code,
- D is an $MDS(2, 4, q)$ code,
- E is an $MDS(2, 4, q_1 - q)$ code on alphabet A ,
- F is an $MDS(2, 4, q_1)$ code with 4- A -hole, where $|A| = q_1 - q$.

Then the set $C = E \cup (M_1 \times_A F) \cup ((M \setminus M_1) \times D)$ is an $MDS(2, 4, (p-1)q + q_1)$ code.

Consider an example of code C that is described in Proposition 7. An $MDS(2, 4, p)$ code is equivalent to a POLS. Determine $p = q_1 = 4$ and $q = 3$. Let M corresponds to the pair

$$\begin{array}{cccc} 0 & 1 & 2 & 3 \\ 3 & 2 & 1 & 0 \\ 1 & 0 & 3 & 2 \\ 2 & 3 & 0 & 1 \end{array} \quad \begin{array}{cccc} 0 & 1 & 2 & 3 \\ 2 & 3 & 0 & 1 \\ 3 & 2 & 1 & 0 \\ 1 & 0 & 3 & 2 \end{array}$$

and let M_1 corresponds to main diagonals of this squares. Suppose that D corresponds to the pair

$$\begin{array}{ccc} b & c & e \\ c & e & b \\ e & b & c \end{array} \quad \begin{array}{ccc} b & c & e \\ e & b & c \\ c & e & b \end{array}, F \text{ corresponds to the pair}$$

$$\begin{array}{cccc} a & b & c & e \\ c & e & a & b \\ b & a & e & c \\ e & c & b & . \end{array} \quad \begin{array}{cccc} b & a & e & c \\ a & b & c & e \\ c & e & a & b \\ e & c & b & . \end{array}$$

and $E = (a, a, a, a)$. Then the constructed code C is equivalent to the following POLS of order 13:

a	$0b$	$0c$	$1b$	$1c$	$1e$	$2b$	$2c$	$2e$	$3b$	$3c$	$3e$	$0e$
$0c$	$0e$	a	$1c$	$1e$	$1b$	$2c$	$2e$	$2b$	$3c$	$3e$	$3b$	$0b$
$0b$	a	$0e$	$1e$	$1b$	$1c$	$2e$	$2b$	$2c$	$3e$	$3b$	$3c$	$0c$
$3b$	$3c$	$3e$	a	$2b$	$2c$	$1b$	$1c$	$1e$	$0b$	$0c$	$0e$	$2e$
$3c$	$3e$	$3b$	$2c$	$2e$	a	$1c$	$1e$	$1b$	$0c$	$0e$	$0b$	$2b$
$3e$	$3b$	$3c$	$2b$	a	$2e$	$1e$	$1b$	$1c$	$0e$	$0b$	$0c$	$2c$
$1b$	$1c$	$1e$	$0b$	$0c$	$0e$	a	$3b$	$3c$	$2b$	$2c$	$2e$	$3e$
$1c$	$1e$	$1b$	$0c$	$0e$	$0b$	$3c$	$3e$	a	$2c$	$2e$	$2b$	$3b$
$1e$	$1b$	$1c$	$0e$	$0b$	$0c$	$3b$	a	$3e$	$2e$	$2b$	$2c$	$3c$
$2b$	$2c$	$2e$	$3b$	$3c$	$3e$	$0b$	$0c$	$0e$	a	$1b$	$1c$	$1e$
$2c$	$2e$	$2b$	$3c$	$3e$	$3b$	$0c$	$0e$	$0b$	$1c$	$1e$	a	$1b$
$2e$	$2b$	$2c$	$3e$	$3b$	$3c$	$0e$	$0b$	$0c$	$1b$	a	$1e$	$1c$
$0e$	$0c$	$0b$	$2e$	$2c$	$2b$	$3e$	$3c$	$3b$	$1e$	$1b$	$1c$	a
<hr/>												
$0b$	a	$0e$	$1b$	$1c$	$1e$	$2b$	$2c$	$2e$	$3b$	$3c$	$3e$	$0c$
a	$0b$	$0c$	$1e$	$1b$	$1c$	$2e$	$2b$	$2c$	$3e$	$3b$	$3c$	$0e$
$0c$	$0e$	a	$1c$	$1e$	$1b$	$2c$	$2e$	$2b$	$3c$	$3e$	$3b$	$0b$
$2b$	$2c$	$2e$	$3b$	a	$3e$	$0b$	$0c$	$0e$	$1b$	$1c$	$1e$	$3c$
$2e$	$2b$	$2c$	a	$3b$	$3c$	$0e$	$0b$	$0c$	$1e$	$1b$	$1c$	$3e$
$2c$	$2e$	$2b$	$3c$	$3e$	a	$0c$	$0e$	$0b$	$1c$	$1e$	$1b$	$3b$
$3b$	$3c$	$3e$	$2b$	$2c$	$2e$	$1b$	a	$1e$	$0b$	$0c$	$0e$	$1c$
$3e$	$3b$	$3c$	$2e$	$2b$	$2c$	a	$1b$	$1c$	$0e$	$0b$	$0c$	$1e$
$3c$	$3e$	$3b$	$2c$	$2e$	$2b$	$1c$	$1e$	a	$0c$	$0e$	$0b$	$1b$
$1b$	$1c$	$1e$	$0b$	$0c$	$0e$	$3b$	$3c$	$3e$	$2b$	a	$2e$	$2c$
$1e$	$1b$	$1c$	$0e$	$0b$	$0c$	$3e$	$3b$	$3c$	a	$2b$	$2c$	$2e$
$1c$	$1e$	$1b$	$0c$	$0e$	$0b$	$3c$	$3e$	$3b$	$2c$	$2e$	a	$2b$
$0e$	$0c$	$0b$	$3e$	$3c$	$3b$	$1e$	$1c$	$1b$	$2e$	$2b$	$2c$	a

Theorem 1 *Suppose that*

- $M_2 \subset M_1 \subset M$ is a super $MDS(2, 5, p)$ code,
- D is an $MDS(2, 5, q)$ code,
- E is an $MDS(2, 5, q_1 - q)$ code on alphabet A ,
- F is an $MDS(2, 5, q_1)$ code with 4- A -hole,
- G is an $MDS(2, 5, q_1)$ code with 5- A -hole, where $|A| = q_1 - q$.

Then the set $C = E \cup (M_2 \times_A G) \cup ((M_1 \setminus M_2) \times_A F) \cup ((M \setminus M_1) \times D)$ is an $MDS(2, 5, (p-1)q + q_1)$ code.

Proof. By the hypotheses of the theorem for any $y \in G$ there exist three $i \in \{1, \dots, 5\}$ such that $y_i \notin A$. Since code distance of M_2 equals 5, for any $x, x' \in M_2$ all coordinates are different. Consequently, if $(x, y) \neq (x', y')$ then $\pi_A(x, y) \neq \pi_A(x', y')$ for $x, x' \in M_2$ and $y, y' \in G$. Therefore $|M_2 \times_A G| = |M_2 \times G| = |M_2||G|$. By the same way we can prove that $|M_2 \times_A F| = |M_2||F|$ and $|M_1 \times_A F| = |M_1||F|$. Then it holds

$$|C| = |E| + |M_2||G| + (|M_1| - |M_2|)|F| + (|M| - |M_1|)|D| =$$

$$(q_1 - q)^3 + p(q_1^3 - (q_1 - q)^3) + (p^2 - p)(q^3 + 3q^2(q_1 - q)) + (p^3 - p^2)q^3 = (pq + q_1 - q)^3.$$

The code distance of $X \times Y$ is the minimum of the code distances of X and Y . If elements of Y contains not more than k symbols from A then $\rho(X \times_A Y) \geq \min(\rho(X) - k, \rho(Y))$. Hence the interior distances of the codes E , $M_2 \times_A G$, $(M_1 \setminus M_2) \times_A F$ and $(M \setminus M_1) \times D$ are not less than 3 by the hypotheses of the theorem. The distance between codes $(M \setminus M_1) \times D$ and E equals 5. The distance between $(M \setminus M_1) \times D$ and $(M_1 \setminus M_2) \times_A F$ (or $M_2 \times_A G$) is not less than the distance between $(M \setminus M_1) \times D$ and $(M_1 \setminus M_2) \times F$ (or $M_2 \times G$). This distance is not less than the distance between $M \setminus M_1$ and $M_1 \setminus M_2$ (or M_2), i.e., it is not less than the code distance of M .

We have that $U_2(E) \cap F = U_2(E) \cap G = \emptyset$ by the definition of a code with j - A -hole. Thus the distance between E and $(M_1 \setminus M_2) \times_A F$ (or $M_2 \times_A G$) is not less than the distance between E and F or G , i.e., it is not less than 3.

The distance between $M_1 \setminus M_2$ and M_2 is equal to 4. Take $(x_0, x_1, x_2, x_3, x_4)$ from $M_1 \setminus M_2$. Each element of $(x_0, x_1, x_2, x_3, x_4) \times_A F$ contains not more than one symbol from A . Consequently, the distance between $(x_0, x_1, x_2, x_3, x_4) \times_A F$ and $M_2 \times_A G$ is not less than $4 - 1 = 3$.

By the Singleton bound (Proposition 2) C is an MDS code. \blacktriangle

It is easy to see that the MDS code C constructed by using the theorem above contains subcodes of orders q and q_1 . These subcodes are $\bar{x} \times D$, where $\bar{x} \in M \setminus M_1$, and $E \cup (\bar{x} \times_A G)$, where $\bar{x} \in M_2$.

Proposition 8 *Let $k \leq p$ and $i \in \{1, \dots, k\}$. Suppose*

- *M is an MDS(2, 5, p) code that contains k disjoint MDS(3, 5, p) codes C_i ,*
- *D is an MDS(2, 5, q) code,*
- *F_i is an MDS(2, 5, q_1) code over alphabet $Q_q \cup A_i$ with 4- A_i -hole, where $|A_i| = q_1 - q$, $A_i \cap A_j = \emptyset$ if $i \neq j$.*

Then the set $S = (\bigcup_{i=1}^k C_i \times_{A_i} F_i) \cup ((M \setminus \bigcup_{i=1}^k C_i) \times D)$ is an MDS(2, 5, $(p - k)q + kq_1$) code with 4- B -hole, where $B = \bigcup_{i=1}^k A_i$.

Proof. By the hypotheses of the proposition for any $y \in F_i$ there exists $j_1, j_2 \in 1, \dots, 5$ such that $y_{j_1}, y_{j_2} \notin A_i$. Since code distance of C_i equals 4, any $x, x' \in C_i$ coincide in one coordinate at most. Consequently, if $(x, y) \neq (x', y')$ then $\pi_{A_i}(x, y) \neq \pi_{A_i}(x', y')$ for $x, x' \in C_i$ and $y, y' \in F_i$. Then $|C_i \times_{A_i} F_i| = |C_i||F_i| = |C_1||F_1|$. By direct calculation we obtain the following equalities:

$$|S| = k|C_1||F_1| + (|M| - k|C_1|)|D| = kp^2(q^3 + 3q^2(q_1 - q)) + (p^3 - kp^2)q^3 = (pq)^3 + 3(pq)^2k(q_1 - q).$$

The distance between $(M \setminus \bigcup_{i=1}^k C_i) \times D$ and $\bigcup_{i=1}^k (C_i \times_{A_i} F_i)$ is not less than the distance between $M \setminus (\bigcup_{i=1}^k C_i)$ and $\bigcup_{i=1}^k C_i$. Since $A_i \cap A_j = \emptyset$ if $i \neq j$, the distance between $C_i \times_{A_i} F_i$ and $C_j \times_{A_j} F_j$ is not less than the distance between C_i and C_j . For $i = 1, \dots, k$ we have $\rho(C_i \times_{A_i} F_i) \geq \min(\rho(C_i) - 1, \rho(F_i)) = 3$. The code distance of $(M \setminus \bigcup_{i=1}^k C_i) \times D$ are not less than the minimum of the code distances of D and M . Therefore, the code distance of S equals 3.

Let us prove that $S \cap U_3((\bigcup_{i=1}^k A_i)^5) = \emptyset$. By definition of 4- A_i -hole, each element of F_i contains not more than one symbol from A_i . So, each element of S contains not more than one symbol from $\bigcup_{i=1}^k A_i$.

Let us prove that $U_3(S) = (Q_q \cup (\bigcup_{i=1}^k A_i))^5 \setminus (\bigcup_{i=1}^k A_i)^5$. Consider any vector \overline{w} with 4 or less coordinates from $\bigcup_{i=1}^k A_i$. Without loss of generality, we take $\overline{w} = ((x_0, y_0), a_1, w_2, w_3, w_4)$, where $a_1 \in A_1$. Since F_1 is an $MDS(2, 5, q_1)$ code with 4- A_1 -hole, there is a vector $\overline{z} = (y_0, a_1, z_2, z_3, z_4) \in F_1$. Since C_1 is an $MDS(3, 5, p)$ code, there exists a vector $\overline{x} = (x_0, x_1, x_2, x_3, x_4) \in C_1$. Then the distance between vectors $\overline{x} \times_{A_1} \overline{z}$ and \overline{w} is equal to 3. \blacktriangle

Lemma 1 *There exists an $MDS(2, 5, 6)$ code with $4\text{-}\{a, b\}\text{-hole}$.*

The proof is by direct verification of the table below.

a	b	2	3	0	1	0	1	b	a	3	2
1	0	b	a	2	3	a	b	0	1	2	3
b	a	0	1	3	2	3	2	a	b	1	0
3	2	a	b	1	0	b	a	3	2	0	1
0	3	1	2	.	.	2	0	1	3	.	.
2	1	3	0	.	.	1	3	2	0	.	.

1	0	<i>a</i>	<i>b</i>	3	2	<i>b</i>	<i>a</i>	1	0	2	3
<i>b</i>	<i>a</i>	2	3	1	0	1	0	<i>a</i>	<i>b</i>	3	2
3	2	<i>b</i>	<i>a</i>	0	1	<i>a</i>	<i>b</i>	2	3	0	1
<i>a</i>	<i>b</i>	0	1	2	3	2	3	<i>b</i>	<i>a</i>	1	0
2	1	3	0	.	.	0	2	3	1	.	.
0	3	1	2	.	.	3	1	0	2	.	.

$$\begin{array}{cccccc} 2 & 3 & b & a & 1 & 0 \\ a & b & 1 & 0 & 3 & 2 \\ 0 & 1 & a & b & 2 & 3 \\ b & a & 3 & 2 & 0 & 1 \\ 3 & 0 & 2 & 1 & . & . \\ 1 & 2 & 0 & 3 & . & . \end{array} \qquad \begin{array}{cccccc} a & b & 3 & 2 & 1 & 0 \\ 3 & 2 & b & a & 0 & 1 \\ b & a & 0 & 1 & 3 & 2 \\ 0 & 1 & a & b & 2 & 3 \\ 1 & 3 & 2 & 0 & . & . \\ 2 & 0 & 1 & 3 & . & . \end{array}$$
$$\begin{array}{cccccc} b & a & 1 & 0 & 2 & 3 \\ 2 & 3 & a & b & 0 & 1 \\ a & b & 3 & 2 & 1 & 0 \\ 0 & 1 & b & a & 3 & 2 \\ 1 & 2 & 0 & 3 & . & . \\ 3 & 0 & 2 & 1 & . & . \end{array} \quad \begin{array}{cccccc} 2 & 3 & a & b & 0 & 1 \\ b & a & 2 & 3 & 1 & 0 \\ 1 & 0 & b & a & 2 & 3 \\ a & b & 1 & 0 & 3 & 2 \\ 3 & 1 & 0 & 2 & . & . \\ 0 & 2 & 3 & 1 & . & . \end{array}$$

0	2	3	1	.	.	1	2	0	3	.	.
3	1	0	2	.	.	2	1	3	0	.	.
1	3	2	0	.	.	0	3	1	2	.	.
2	0	1	3	.	.	3	0	2	1	.	.
.
.

3	1	0	2	.	.	3	0	2	1	.	.
0	2	3	1	.	.	0	3	1	2	.	.
2	0	1	3	.	.	2	1	3	0	.	.
1	3	2	0	.	.	1	2	0	3	.	.
.
.

Theorem 2 *If $q = 16(6s \pm 1) + 4$, then there exists an $MDS(2, 5, q)$ code.*

Proof. By Lemma 1 and Propositions 5 and 8 ($p = q = 4, k = 2, q_1 = 6$), there exists an $MDS(2, 5, 20)$ code with 4- A -hole, where $|A| = 4$. By Theorem 1 ($q_1 = 20, q = 16, k = 4$) we can obtain an $MDS(2, 5, 16p + 4)$ code if there exists a super $MDS(2, 5, p)$ code. Since any integer $p = 6s \pm 1$ is not divisible by 2 and 3, there exists a super $MDS(2, 5, p)$ code by Propositions 4 and 6. \blacktriangle

By Proposition 1 all $MDS(2, 5, q)$ codes are equivalent to pairs of orthogonal latin cubes of order q . If $6s - 1 = 18i - 1$ or $6s - 1 = 18i + 5$, then pairs of orthogonal latin cubes of order $q = 16(6s - 1) + 4$ were not previously known because in these cases q is divisible by 3 but it is not divisible by 9. Ten minimal new obtained orders (not only of type $q = 16(6s - 1) + 4$) are 84, 132, 276, 372, 516, 564, 660, 852, 948, 1140.

4 Connection between MDS codes and combinatorial designs

A *Steiner system* with parameters τ, d, q , $\tau \leq d$, written $S(\tau, d, q)$, is a set of d -element subsets of Q_q (called *blocks*) with the property that each τ -element subset of Q_q is contained in exactly one block.

Theorem 3 *If D_2 and D_3 are Steiner systems $S(2, 5, q)$ and $S(3, 5, q)$ respectively and $D_2 \subset D_3$, then there exists an $MDS(2, 5, q)$ code.*

Proof. Consider a block $X = \{x_1, x_2, x_3, x_4, x_5\} \in D_3 \setminus D_2$. Define a set $M_X = \{(x_{\tau 1}, x_{\tau 2}, x_{\tau 3}, x_{\tau 4}, x_{\tau 5}) \mid \tau \in \text{Alt}(5)\}$, where $\text{Alt}(5)$ is the alternating group.

By Proposition 3 there exists an $MDS(2, 5, 5)$ code that contains (a, a, a, a, a) for all $a \in Q_5$. Suppose that $X = \{x_1, x_2, x_3, x_4, x_5\} \in D_2$. Let us define an $MDS(2, 5, 5)$ code M_X over the alphabet X such that M_X contains $(x_i, x_i, x_i, x_i, x_i)$ for $i = 1, \dots, 5$. The intersections of pairs of such codes contain only elements of type (a, a, a, a, a) for $a \in Q_q$.

Let us to prove that $M = \bigcup_{X \in D_3} M_X$ is an $MDS(2, 5, q)$ code. The following holds:

$$|M| = q + |D_2|(5^3 - 5) + (|D_3| - |D_2|)|\text{Alt}(5)| =$$

$$q + 5 \cdot 24 \frac{q(q-1)}{4 \cdot 5} + 3 \cdot 4 \cdot 5 \left(\frac{q(q-1)(q-2)}{3 \cdot 4 \cdot 5} - \frac{q(q-1)}{4 \cdot 5} \right) = q^3.$$

Suppose that $X \in D_3 \setminus D_2$, $Y \in D_3$ and $X \neq Y$. The distance between codes M_X and M_Y is not less than 3 because $|X \cap Y| \leq 2$. Suppose $X, Y \in D_2$ and $X \neq Y$. Then $|X \cap Y| \leq 1$. If $x \in M_X$ is not a constant vector, then it contains not more than 2 equal symbols. If $x \in M_X$ and $y \in M_Y$ are not constant vectors, then $\rho(x, y) \geq 3$ by direct verification.

If $x, y \in M_X$ and $X \in D_2$, then $\rho(x, y) \geq 3$ by the definition of M_X . Any non-constant permutation from $\text{Alt}(5)$ permutes 3 or more elements. Therefore for $X \in D_3 \setminus D_2$ we obtain that $\rho(x, y) \geq 3$ for any distinct $x, y \in M_X$.

Thus we proved that the code distance of M is at least 3. So, M is an $MDS(2, 5, q)$ code by the Singleton bound (Proposition 2). \blacktriangle

The natural divisibility conditions for the existence of Steiner systems $S(2, 5, n)$ and $S(3, 5, n)$ simultaneously is that $n = 5$ or $41 \pmod{60}$. Steiner systems $S(3, 5, 41)$ are unknown. Steiner systems $S(2, 5, 65)$ and $S(3, 5, 65)$ exist. Systems $S(2, q + 1, q^3 + 1)$ are unitals and systems $S(3, q + 1, q^3 + 1)$ are spherical geometries if q is a prime power ($q = 4$ in this case). But it is unknown whether the system $S(3, 5, 65)$ contains the system $S(2, 5, 65)$. Keevash [6] and Glock et al. [4] proved that the natural divisibility conditions are sufficient for existence of Steiner system $S(t, k, n)$ (and inserted Steiner systems) apart from a finite number of exceptional n given fixed t and k . Therefore it is possible to use the theorem above for constructing $MDS(2, 5, q)$ codes if q is large enough.

5 Acknowledgments

My sincere thanks are due to D.S.Krotov who programmed the proposed method and calculated a series of new pairs orthogonal latin cubes (see <https://ieee-dataport.org/open-access/graeco-latin-cubes>).

References

- [1] R.C. Bose, S.S. Shrikhande and E.T. Parker, “Further results on the construction of mutually orthogonal latin squares and the falsity of Euler’s conjecture”, *Canad. J. Math.* **12** (1960), 189.
- [2] J. Denes and A.D. Keedwell, Latin squares: new development in the theory and applications. Annals of Discrete Mathematics, vol. 46, North-Holland, Amsterdam, 1991.
- [3] J.T. Ethier and G.L. Mullen, “Strong forms of orthogonality for sets of hypercubes”, *Discrete Math.* **312** (2012), no. 12-13, 2050–2061.
- [4] S. Glock, D. Kühn, A. Lo, and D. Osthus, The existence of designs via iterative absorption: hypergraph F -designs for arbitrary F , arXiv preprint arXiv:1611.06827v3[math.CO], 2020.
- [5] K. Heinrich and L. Zhu, “Existence of orthogonal Latin squares with aligned subsquares”, *Discrete Math.* **59** (1986), no. 1-2, 69–78.
- [6] P. Keevash, The existence of designs, arXiv preprint arXiv:1401.3665v3[math.CO], 2019.
- [7] P. Keevash, The existence of designs II, arXiv preprint arXiv:1802.05900[math.CO], 2018.
- [8] F.J. MacWilliams and N.J.A. Sloane, The theory of error-correcting codes. Elsevier/North-Holland, Amsterdam, 1977.
- [9] V.N. Potapov, “On the multidimensional permanent and q -ary designs”, *Siberian Electronic Mathematical Reports* **11** (2014), 451–456.