

Characterization and Generation of Absolutely Separable States

Saronath Halder, Shiladitya Mal, Aditi Sen(De)

Harish-Chandra Research Institute, HBNI, Chhatnag Road, Jhusi, Allahabad 211 019, India

We characterize the boundary of the convex compact set of absolutely separable states that cannot be transformed to entangled states by global unitary operators in $2 \otimes d$ Hilbert space. In particular, we show that the absolutely separable states of rank- $(2d - 1)$ are extreme points of such sets. Moreover, we prove the existence of full-rank extreme and boundary points for those Hilbert spaces. Properties of certain interior points are also explored. We further show that by exploring the boundary of the above set, it is possible to develop an algorithm to generate the absolutely separable states which stay outside the maximal ball. By exploring paradigmatic noise models, we find the amount of local noise which the input entangled states can sustain, so that the output states do not become absolutely separable. Interestingly, we report that with the decrease of entanglement of the pure input state, critical depolarizing noise value, transferring an entangled state to an absolutely separable one, increases, thereby showing advantages of sharing nonmaximally entangled states.

I. INTRODUCTION

Characterization of resourceful quantum states is important from the perspective of several quantum information processing tasks [1, 2]. These include quantum communication protocols like quantum state transfer [3, 4] using entangled states [5, 6], encoding of classical information into quantum states [7, 8], secure communication via entangled states [9–13], measurement-based quantum computation [14–16]. In a resource theory, along with the characterization of set of states according to certain tasks, understanding the set of operations, known as free operations, by which resourceful states cannot be created is also important. For example, in the theory of entanglement, the set of local operations and classical communication constitute the free operations by which only separable states can be produced. Therefore, characterising useless states in any paradigm can be essential to understand the free operations.

Two-qubit gates or joint unitary operators acting on two-qubit pure states can, in general, create entanglement in the systems [17–19]. For example, from the initial product state $|-\rangle|0\rangle$, with $|-\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$, two-qubit *CNOT* gate can create a maximally entangled state. However, it was shown that there are bipartite states from which it is not possible to generate entanglement by acting joint (global) unitary operations. They are called *absolutely separable states* [20, 21] (see also Refs. [22–27] in this regard). In a resource theory in which global unitary operators are free operations, the set of absolutely separable states are not useful states. However, it is important to understand the properties of such states from resource theoretic point of view.

Recently, it has been shown that for a given Hilbert space, the set of absolutely separable states form a convex and compact set [25]. Therefore, it is always possible to detect those states by constructing witness operators (for entanglement witnesses, see Refs. [28–30]) from separable ones which can be converted to entangled states under global unitary operators by applying

the *Hahn-Banach separation theorem* [31]. In this context, we note that to construct optimal witness operators, it is important to explore the boundary points of the set of absolutely separable states. Moreover, *Krein-Milman theorem* [32] states that a convex compact set corresponding to a finite dimensional vector space is equal to the convex hull of the extreme points of that set. Therefore, to understand a convex compact set, it is enough to know about the extreme points of that set.

The main objectives of the present work is twofold: (i) We consider the characterization of absolutely separable states by exploring the boundary of the set of absolutely separable states when the quantum system is associated with a $2 \otimes d$ Hilbert space. In particular, we show the states having rank- $(2d - 1)$ in $2 \otimes d$ always be extreme points of the set. On the other hand, the states having full-rank can be interior as well as extreme points of the above set, which can be on or outside of the maximal ball. By exploring the boundary of this set, it is also possible to develop an algorithm to generate the absolutely separable states which stay outside the *maximal ball* (it is the maximal ball around maximally mixed state in which all the states are separable) [33, 34]. (ii) We further search for noisy scenarios which result absolutely separable states. Finding such situations can be interesting from the perspective of experiments. Specifically, we find critical strengths of local noise for different prototypical noise models [1, 2], which lead to absolutely separable states. Since entangled states cannot be generated from absolutely separable states by applying global unitary operators, effects of local decoherence on state space induces irreversibility in the theory of entanglement.

This manuscript is arranged as follows: After a few basics, presented in Sec. II, we characterize the set of absolutely separable states in Sec. III. In Sec. IV, we address the question of irreversibility appeared due to the existence of absolutely separable states when an entangled state is affected by decoherence. Finally, we conclude in Sec. V.

II. PRELIMINARIES

Absolutely separable states are those which cannot be transformed to an entangled states under the action of global unitary operators. For any given Hilbert space, the identity operator is an example of absolutely separable operator. For a $2 \otimes d$ Hilbert space, there is a *necessary and sufficient* condition to check whether a separable state is absolutely separable [21, 22, 24]. In particular, it was shown that in a $2 \otimes d$ system, a state is absolutely separable if and only if

$$\lambda_1 - \lambda_{2d-1} - 2\sqrt{\lambda_{2d-2}\lambda_{2d}} \leq 0, \quad (1)$$

where $\lambda_1, \dots, \lambda_{2d}$ are the eigenvalues in decreasing order corresponding to a bipartite density matrix ρ_{AB} , associated with a $2 \otimes d$ Hilbert Space [21, 22, 24].

With increasing importance of the entangled states, the properties of states lying in the neighborhood of the maximally mixed states [35] were studied. It was shown that the largest ball of separable as well as absolutely separable states around maximally mixed state in a two-qubit system can be described by $\text{Tr}(\rho^2) \leq \frac{1}{3}$ [20, 33, 34]. This ball is known as *maximal ball*. In general, for a $d \otimes d$ system, the maximal ball can be described by $\text{Tr}(\rho^2) \leq \frac{1}{d^2-1}$ [33, 34]. However, it was found that there are absolutely separable states which reside outside the maximal ball [20]. So, to understand these states, it is important to construct such states.

The possible structures of witness operators which separates absolutely separable states from the separable states were explored in [25]. But to make these operators optimal, one may require to explore the boundary points of the set of absolutely separable states. So, understanding these boundary points is one of the main objectives, discussed in the succeeding section. In this regard, remember that a state is not an extreme point of a convex set if it can be written as a convex combination of two or more absolutely separable states. On the other hand, if an absolutely separable state does not allow any such decomposition, then it must be an extreme point of the set.

III. CHARACTERIZATION OF ABSOLUTELY SEPARABLE STATES

Let us now concentrate on any two-party state, ρ_{AB} in $2 \otimes d$ which is absolutely separable. It is known that the pure product states can always be transformed to entangled states under global unitary operations. From condition (1), it is also clear that rank-2 states cannot be absolutely separable states in $2 \otimes d$ [21, 22, 24]. Moreover, the condition given in (1), tells that the mixed states having rank $\leq (2d-2)$ cannot be absolutely separable in $2 \otimes d$. So, in $2 \otimes 2$, absolutely separable states can have rank 3 and 4. Since the set of absolutely separable states is a convex and compact set in any dimension

[25], it is possible to explore certain geometric properties of such a set, especially the structure of the extreme points of the set of absolutely separable states. Here, we say this set as **AS**. We prove the following proposition for absolutely separable states having lowest rank in $2 \otimes 2$:

Proposition 1. *All rank-3 absolutely separable states are extreme points of AS in $2 \otimes 2$.*

Proof. In $2 \otimes 2$, a state is absolutely separable if and only if $\lambda_1 - \lambda_3 - 2\sqrt{\lambda_2\lambda_4} \leq 0$, where $\lambda_1, \dots, \lambda_4$ are the eigenvalues of a given density matrix in decreasing order. For a rank-3 state, $\lambda_4 = 0$, which implies that a rank-3 state is absolutely separable iff $\lambda_1 \leq \lambda_3$. But we have assumed that $\lambda_1 \geq \lambda_3$. This indicates that a rank-3 state is absolutely separable if and only if $\lambda_1 = \lambda_2 = \lambda_3 = \frac{1}{3}$. Now, consider the spectral decomposition of a state, $\rho_{AB} = \frac{1}{3} |\psi_1\rangle \langle \psi_1| + \frac{1}{3} |\psi_2\rangle \langle \psi_2| + \frac{1}{3} |\psi_3\rangle \langle \psi_3|$, where $\{|\psi_i\rangle\}$, $i=1,2,3$ are orthogonal states. Obviously, this state is the only absolutely separable state in the 3-dimensional subspace spanned by $\{|\psi_1\rangle, |\psi_2\rangle, |\psi_3\rangle\}$. Therefore, this state cannot be written as a convex combination of two or more absolutely separable states, proving the fact that all rank-3 absolutely separable states are extreme points of **AS**. \square

Since, there is only one particular structure, possible for rank-3 absolutely separable states, any rank-3 state which does not capture such a structure must not be an absolutely separable state. Now, we mention here the following remark:

Remark 1. *The number of rank-3 extreme points of AS in $2 \otimes 2$ can be infinitely many and they resides on the maximal ball.*

This is because in a two-qubit Hilbert space, one can choose three orthogonal states in infinitely many possible ways and then, take an equal mixture of three orthogonal states, leaving the final state to be an extreme point of **AS**. Next, using the condition given in (1), the generalization of the above proposition for a $2 \otimes d$ system is presented:

Corollary 1. *All rank- $(2d-1)$ absolutely separable states are extreme points of AS in $2 \otimes d$.*

Proof. The proof is in the same line as Proposition 1. It uses the fact that any state of **AS**, having rank- $(2d-1)$, is the only state of **AS** supported in a proper subspace (of $2 \otimes d$ Hilbert space) spanned by a set of orthogonal states $\{|\psi_i\rangle\}_{i=1}^{2d-1}$. \square

The above state has a spectral decomposition: $\rho = \frac{1}{2d-1} (\sum_{i=1}^{2d-1} |\psi_i\rangle \langle \psi_i|)$, where $|\psi_i\rangle$ s are orthogonal states. Clearly, purity of the state ρ is given by $\text{Tr}[\rho^2] = \frac{1}{2d-1}$.

Remark 2. *All rank- $(2d-1)$ absolutely separable states in $2 \otimes d$ having purity $\frac{1}{2d-1}$ are included in the maximal ball*

characterised by $[\text{Tr}(\rho_{AB}^2) \leq \frac{1}{2d-1}]$ and they all lie on the surface of the ball.

Interestingly, there exist rank-4 absolutely separable states in $2 \otimes 2$ which can stay outside the maximal ball [20]. In this context, we mention that there is no known protocol to produce rank-4 absolutely separable states in $2 \otimes 2$ which reside outside the maximal ball. We now propose a prescription to produce rank-4 absolutely separable states which reside outside the maximal ball. This can be done by exploring the boundary of **AS**.

1. Take a rank-3 state in $2 \otimes 2$ which is not absolutely separable, i.e., such states must reside outside the maximal ball. From Proposition 1, it is clear that such states are of the form $\sum_{i=1}^3 p_i |\psi_i\rangle \langle \psi_i|$ with at least one p_i , not equal to the other p_i s.
2. Consider a pure state in $2 \otimes 2$ which is orthogonal to the previous state. Pure states are not included in the maximal ball and they cannot be absolutely separable either.
3. Suitable convex combination of these two states can produce rank-4 states in $2 \otimes 2$ which are absolutely separable. But we have to take the convex combination in such a way that the newly generated states reside outside the maximal ball for some choices of parameters.

Example. Let us now illustrate the recipe discussed above by an example. We consider a two-qubit rank-3 state $\rho_1 = \frac{1}{2} |00\rangle \langle 00| + \frac{1}{4} |01\rangle \langle 01| + \frac{1}{4} |10\rangle \langle 10|$, which is not an absolutely separable state. Also consider a pure state $\rho_2 = |11\rangle \langle 11|$ which cannot be absolutely separable. These two states are orthogonal to each other. So, any convex combination of them must be a rank-4 state. We now take convex combination of these two states $q\rho_1 + (1-q)\rho_2$ in a way that $q = \frac{16}{17}$. It can be checked that the newly prepared state is an absolutely separable states. Interestingly, if $q = \frac{16}{17}$, then $\text{Tr}[q\rho_1 + (1-q)\rho_2]^2 > \frac{1}{3}$, indicating the fact that the state resides outside the maximal ball.

Notice that in the above, when $q = \frac{16}{17}$, the newly prepared rank-4 state is just included into **AS**. Therefore, it is a boundary point of the set. But it is not known whether the state is an extreme point of **AS**. Note that the above protocol is quite easy to generalize for $2 \otimes d$. In that case, one has to start with a rank- $(2d-1)$ state which is not included in the maximal ball as well as in **AS** (can be found by Corollary 1). Then consider a pure state which is orthogonal to the previous state and the rest is as described in the above.

Let us now move to full-rank, i.e., rank-4 absolutely separable states in $2 \otimes 2$ and discuss different properties of a set consisting of such states. It is known that there exists a ball around the maximally mixed state, (\mathbb{I}/d^2) , in $d \otimes d$ and all the states, belonging to that ball, are

absolutely separable [33, 34]. So, the maximally mixed state is an interior point of **AS** in $d \otimes d$. Clearly, any state ρ (except the maximally mixed state) which is an interior point of **AS**, can be written as a convex combination of the maximally mixed state (\mathbb{I}/d^2) and another absolutely separable state σ , where $\rho \neq \sigma$, and both ρ and σ are not maximally mixed. Note that for different states ρ , the states σ can be different. However, the states which do not allow such a decomposition must be boundary points of **AS**. We now present the following observation:

Proposition 2. *In $2 \otimes 2$, any rank-4 absolutely separable state ρ which satisfy the condition $\lambda_1 - \lambda_3 = 2\sqrt{\lambda_2\lambda_4}$, cannot be written as a convex combination of the maximally mixed state $(\mathbb{I}/4)$ and another absolutely separable state σ , not maximally mixed, with $\rho \neq \sigma$, where λ_i s are the eigenvalues of ρ in decreasing order.*

Proof. We consider a spectral decomposition of the above state $\rho = \sum_{i=1}^4 \lambda_i |\psi_i\rangle \langle \psi_i|$. Now, consider a decomposition of ρ as the following:

$$\rho = \sum_{i=1}^4 \lambda_i |\psi_i\rangle \langle \psi_i| = (1-4\epsilon) \sum_{i=1}^4 \frac{(\lambda_i - \epsilon)}{1-4\epsilon} |\psi_i\rangle \langle \psi_i| + (4\epsilon) \sum_{i=1}^4 \frac{\epsilon}{4\epsilon} |\psi_i\rangle \langle \psi_i| = (1-4\epsilon)\sigma + (4\epsilon)\frac{\mathbb{I}}{4}, \quad (2)$$

where ϵ can be considered as a very small number and \mathbb{I} is the identity operator. Now, the question is whether σ is an absolutely separable state. The state σ is absolutely separable if and only if the eigenvalues of it $(\lambda_i - \epsilon)/(1-4\epsilon)$ (in the decreasing order) obey the condition (1). If it is the case then $\lambda_1 - \lambda_3 \leq 2\sqrt{(\lambda_2 - \epsilon)(\lambda_4 - \epsilon)}$. But this cannot be because we have assumed $\lambda_1 - \lambda_3 = 2\sqrt{\lambda_2\lambda_4}$. So, for any $\epsilon > 0$, $(\lambda_1 - \lambda_3) > 2\sqrt{(\lambda_2 - \epsilon)(\lambda_4 - \epsilon)}$, leading to the fact that σ cannot be absolutely separable. Thus, it is proved that if the eigenvalues of ρ (in the decreasing order) obey the condition $\lambda_1 - \lambda_3 = 2\sqrt{\lambda_2\lambda_4}$ then the state cannot be written as a convex combination of the maximally mixed state and another absolutely separable state. \square

In the above, it is clear that any state ρ whose eigenvalues (in the decreasing order) obey the condition $\lambda_1 - \lambda_3 = 2\sqrt{\lambda_2\lambda_4}$, are the boundary points of **AS**. Next, we describe an important property of certain interior points of the set.

Proposition 3. *In $2 \otimes 2$, if the eigenvalues (in the decreasing order) of a rank-4 absolutely separable state (except the maximally mixed state) satisfy the condition $\lambda_1 - \lambda_3 < 2\sqrt{\lambda_2\lambda_4}$ the state can be written as a convex combination of the maximally mixed state and a boundary point of **AS**.*

Proof. We consider the states other than the maximally mixed state. Suppose, the spectral decomposition of such a state is given by $\rho = \sum_{i=1}^4 \lambda_i |\psi_i\rangle \langle \psi_i|$. This state can be decomposed into the following form:

$$\rho = \sum_{i=1}^4 \lambda_i |\psi_i\rangle \langle \psi_i| = (1-4\epsilon) \sum_{i=1}^4 \frac{(\lambda_i - \epsilon)}{1-4\epsilon} |\psi_i\rangle \langle \psi_i| + (4\epsilon) \sum_{i=1}^4 \frac{\epsilon}{4\epsilon} |\psi_i\rangle \langle \psi_i| = (1-4\epsilon)\sigma + (4\epsilon)\frac{\mathbb{I}}{4}, \quad (3)$$

Our goal is to prove that the state σ in the above decomposition is a boundary point of **AS**. Because if it is the case then the absolutely separable state ρ can be written as the convex combination of the maximally mixed state and a boundary point of **AS**. Notice that the eigenvalues of σ are given by $(\lambda_i - \epsilon)/(1 - 4\epsilon)$. It is a boundary point of **AS** if $\lambda_1 - \lambda_3 = 2\sqrt{(\lambda_2 - \epsilon)(\lambda_4 - \epsilon)}$. We have assumed that $\lambda_1 - \lambda_3 < 2\sqrt{\lambda_2\lambda_4}$. Clearly, it is possible to consider $\epsilon \leq \lambda_4$, such that σ becomes a boundary point. The value of ϵ can be found by solving the quadratic equation $\lambda_1 - \lambda_3 = 2\sqrt{(\lambda_2 - \epsilon)(\lambda_4 - \epsilon)}$. This completes the proof. \square

Obviously, the states of Proposition 3 are interior points of **AS**. However, like Proposition 1, Proposition 3 can also be generalized for $2 \otimes d$ following the same argument given in the above proof. This leaves us the following corollary.

Corollary 2. *For a $2 \otimes d$ system if the eigenvalues (in decreasing order) of a given absolutely separable state (except the maximally mixed state) satisfy the condition $\lambda_1 - \lambda_{2d-1} < 2\sqrt{\lambda_{2d-2}\lambda_{2d}}$, the state can be written as a convex combination of the maximally mixed state and a boundary point of **AS**. Obviously, such a state must be an interior point of **AS**.*

Let us now address another important question: Are all boundary points of **AS** extreme points? In the following, we show that this is not the case.

Proposition 4. *In $2 \otimes 2$, there exist rank-4 absolutely separable states which are the boundary points of **AS** but they are not extreme points of the set.*

Proof. We simply construct a class of states, the eigenvalues (in the decreasing order) of which satisfy the condition (1) with equality and the constructed states allow some convex decomposition. Specifically, let us consider a two-qubit state $\sigma_1 = \sum_{i=1}^4 \lambda_i |\psi_i\rangle \langle \psi_i|$ (spectral decomposition), where λ_i s are in decreasing order and they satisfy the condition $\lambda_1 - \lambda_3 = 2\sqrt{\lambda_2\lambda_4}$. Similarly, we have another two-qubit state $\sigma_2 = \sum_{i=1}^4 \lambda'_i |\psi_i\rangle \langle \psi_i|$ (spectral decomposition), with λ'_i s being in decreasing order, satisfying the condition $\lambda'_1 - \lambda'_3 = 2\sqrt{\lambda'_2\lambda'_4}$. For any convex combination $x\sigma_1 + (1-x)\sigma_2$, $0 < x < 1$, the newly generated state can have the following eigenvalues: $\mu_i = x\lambda_i + (1-x)\lambda'_i$, for $i = 1, \dots, 4$. μ_i s are also in decreasing order. It can be shown that $\mu_1 - \mu_3 = 2\sqrt{\mu_2\mu_4}$ if $\lambda'_2/\lambda'_4 = \lambda_2/\lambda_4$. We assume $\lambda'_2/\lambda'_4 = \lambda_2/\lambda_4 = \kappa$. This along with the conditions $\sum_i \lambda_i = \sum_i \lambda'_i = 1$, it can be shown that $\lambda_1 = [1 - (1 + \kappa)\lambda_4 + 2\sqrt{\kappa\lambda_4}]/2$, $\lambda_2 = \kappa\lambda_4$, $\lambda_3 = [1 - (1 + \kappa)\lambda_4 - 2\sqrt{\kappa\lambda_4}]/2$. λ'_i s also satisfy similar relations. [Notice that for $i = 1, 2, 3$, λ_i s or λ'_i s are the function of λ_4 or λ'_4 respectively. Similarly, for $i = 1, 2, 3$, μ_i s are also the same function of μ_4 .] However, for proper choice of κ , λ_4 , λ'_4 , one can get λ_i s and

λ'_i s in decreasing order. Thus, one can generate absolutely separable states which satisfy the condition given in (1) with equality. Moreover, these states allow convex decomposition, implying the fact that such states are boundary points but not extreme points of **AS**. To constitute an example, one may consider $\kappa = 2.5$, $\lambda_4 = 0.1$, and $\lambda'_4 = 0.11$. \square

In the above context, we mention that if we assume for σ_1 , $\lambda_1 - \lambda_3 < 2\sqrt{\lambda_2\lambda_4}$ and for σ_2 , $\lambda'_1 - \lambda'_3 < 2\sqrt{\lambda'_2\lambda'_4}$, for any newly generated state, $\mu_1 - \mu_3$ must be less than $2\sqrt{\mu_2\mu_4}$. Nevertheless, which states are the rank-4 extreme points of **AS** in $2 \otimes 2$ is still an open problem but we are able to prove the existence of such states in the succeeding proposition.

Proposition 5. *In $2 \otimes 2$, there exist rank-4 absolutely separable states outside the maximal ball which are extreme points of **AS**.*

Proof. In Proposition 1, we have proved that all rank-3 absolute separable states are extreme points of **AS**. Moreover, they reside on the surface of maximal ball, defined by $\text{Tr}[\rho^2] \leq \frac{1}{3}$. We mention here that the maximal ball is a convex set, based on the fact that convex combination does not allow to increase purity. We know that there are rank-4 absolute separable states outside the maximal ball. Example of such a state is explicitly constructed after Proposition 1. Obviously, these rank-4 states cannot be written as a convex combination of rank-3 extreme points of **AS**. Since these states reside outside the maximal ball. Therefore, such a state is either an extreme point of **AS** or they can be written as a convex combination of rank-4 extreme points of **AS** outside the maximal ball. This completes the proof. \square

IV. INTERCONVERTABILITY BETWEEN ABSOLUTE SEPARABLE AND ENTANGLE STATE

Entanglement is a resource for different quantum information processing tasks. Since absolutely separable states cannot be transformed to entangled states under global unitary operations, their existence puts restriction on the state space. First, we discuss how auxiliary systems can help to overcome absolute separability and prescribe a method to identify operations on **AS** so that it becomes entangled. Secondly, we address how an entangled state get converted into separable or absolutely separable state through noisy channels.

A. Qubit assisted entanglement generation

It is easy to notice that given an absolutely separable state if one enlarges the local dimension, one can find joint unitary operator(s) corresponding to the higher dimensional Hilbert space, which can generate entanglement. In this regard, we present the following:

Observation 1. *If the absolutely separable state is of full-rank, there can be restriction on the form of the auxiliary qubit, where for a non-full rank absolutely separable state, any auxiliary qubit can serve the purpose.*

The first part is simple to establish. Suppose, for a two-qubit system, the maximally mixed state is given. Obviously, that state is an absolutely separable state. Now, if the second party enlarges the local dimension by considering an auxiliary qubit, then it must not be prepared in the maximally mixed state in order to generate entanglement by applying a global unitary operator. Because if the auxiliary qubit is prepared in a maximally mixed state, the overall state is again a maximally mixed state in $2 \otimes 4$ and thus, generating entanglement using any global unitary operator is not possible.

On the other hand, suppose, a non-full rank absolutely separable state is given in $2 \otimes 2$. Then an auxiliary qubit can be properly chosen if one wants to enlarge the local dimension, so that a separable state of rank lower than $2d - 1$ can be obtained, which by construction is not absolutely separable. So, using the final state, entanglement can be generated via a suitable global unitary operator. This follows from the fact that in $2 \otimes d$, a state which have rank less than $2d - 1$ cannot be absolutely separable.

From the above observation, we prescribe a possible way of creating entangled state from **AS** and derive Kraus operators for that kind of operations. Suppose global unitary operator, U , acts both on an absolutely separable state and an auxiliary state $|0\rangle\langle 0|$. Mathematically,

$$\Lambda(\rho_{AS}) = \text{Tr}_B[U(\rho_{AS} \otimes |0\rangle_B\langle 0|)U^\dagger] = \sum_{\mu} K_{\mu}\rho_{AS}K_{\mu}^\dagger \quad (4)$$

where $K_{\mu} = \langle \mu|U|0\rangle$ with $\sum K_{\mu}^\dagger K_{\mu} = \mathbb{I}$.

B. Identifying absolutely separable states in noisy environments

In realistic scenarios, due to imperfection in preparation procedure or interaction with the environment, the entangled state generated in the laboratory is not a pure one. This effect of noise can be modeled in various ways. Typically, a local noisy channel destroys entanglement, thereby creates a separable state [5] which may or may not be absolutely separable. We are interested to find the amount of noise required in the channel to produce absolutely separable states. Such a study can be important for two reasons – (1) in experiments, a state is always affected by noise, (2) finding out the range of noise parameter which makes the state absolutely separable, is significant to avoid them.

We deal with three paradigmatic noise models, namely depolarizing, amplitude damping, and phase damping channels which can affect an initial state differently, thereby, generate absolutely separable states

in independent ways. Before discussing the consequence of noisy channels on a given state, let us first fix the transformation of the input state that happens due to the interactions between the environment and the system. The depolarizing channel (DPC), Λ_{DPC} takes an arbitrary quantum state, ρ to $\rho' = \Lambda_{DPC}(\rho) = p\rho + \frac{(1-p)}{3} \sum_{i=x,y,z} \sigma_i \rho \sigma_i$ where σ_i ($i = x, y, z$)s represent the Pauli operators and $(1 - p)$ is the strength of the noise.

On the other hand, the amplitude damping channel (ADC), Λ_{ADC} acts asymmetrically on the states $|0\rangle$ and $|1\rangle$. In particular, it keeps $|0\rangle$ unchanged while it flips $|1\rangle$ to $|0\rangle$ with probability $(1 - p)$ and the corresponding Kraus operators which describe the effects of ADC are given by

$$K_1 = \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{p} \end{pmatrix}, \quad K_2 = \begin{pmatrix} 0 & \sqrt{1-p} \\ 0 & 0 \end{pmatrix}. \quad (5)$$

The input state ρ after sending through the ADC results in an output state, represented as

$$\Lambda_{ADC}(\rho) = \sum_{i=1}^2 K_i \rho K_i^\dagger. \quad (6)$$

Finally, we consider a scenario in which a qubit is sent through a phase damping channel (PDC), described by

$$\Lambda_{PDC}(\rho) = \sum_{i=1}^3 K_i \rho K_i^\dagger, \quad (7)$$

with the operators K_i , $i = 1, 2, 3$ are given by

$$K_1 = \sqrt{1-p} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad K_2 = \sqrt{p} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad (8)$$

$$K_3 = \sqrt{p} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

A noisy environment always takes an initial pure state, $|\psi\rangle = \cos \frac{x}{2} |00\rangle + e^{-i\phi} \sin \frac{x}{2} |11\rangle$ with $0 \leq x \leq \pi$ and $0 \leq \phi \leq 2\pi$ to a mixed one which may or may not be an entangled state. Note that the input state $|\psi\rangle$ is entangled for all values of x and ϕ except when $x = 0$ or π . We assume here that two independent and identical channels, $\Lambda^1(p) \otimes \Lambda^2(p)$, act on the input state where Λ^i can be either DPC or ADC or PDC and we are interested with the properties of output state.

1. Absolutely separable states via depolarizing channels

Let us start the investigation by studying the features of the output state obtained after sending the initial state via local depolarizing channels (also see Ref. [36] in this regard), i.e.,

$$|\psi\rangle \rightarrow \Lambda_{DPC}^1 \otimes \Lambda_{DPC}^2(|\psi\rangle\langle\psi|) = \rho'(p, x, \phi). \quad (9)$$

The characteristics of the resulting state can be summarized as the following:




FIG. 1. (Color online) Creation of absolutely separable states from a pure state $|\psi\rangle = \cos(x/2)|00\rangle + e^{-i\phi}\sin(x/2)|11\rangle$. Note that $|\psi\rangle$ is maximally entangled with $x = \pi/2$. It is sent through local depolarizing channels which produce separable as well as absolutely separable states with the variation of parameter p . Blue, green, red lines correspond to $x = \pi/2$, $x = \pi/6$, and $x = \pi/12$ respectively. Dotted lines represent the minimum eigenvalues of the partial transposed output states for different values of x while dashed ones are the condition for absolutely separable states, given in Eq. (1). In the figure, it is clear that there exists a range of p where the output state is separable but not absolutely separable when the input state is a pure nonmaximally entangled state.

1. The output state is a rank-4 state. As discussed in Sec. III, the full-rank states can in principle be an absolutely separable for certain choices of p , x and ϕ . For any fixed values of x , we find that there exists a critical value of noise above which the output state is absolutely separable.
2. Two eigenvalues of the output state are same and are independent of the input state parameter, x and ϕ , while the other two depends on x and p . For a fixed value of x , we observe that there exists a critical value of noise $1 - p_{abs}$, above which the state is absolutely separable (see Fig. 1).
3. The partial transposed output state leads to the condition that the state is entangled when $\sin x > [4(1+2p)(1-p)]/(4p-1)^2$ as depicted in Fig. 1.

Remember that we have taken here the strength of the noise as $(1-p)$ while in the above figure, the plots are made with respect to the parameter p . However, looking at the properties 2 and 3, we observe that when $x < \pi/2$, $(1-p_{abs})$ is strictly greater than $(1-p_{sep})$, below which the state is entangled. Interestingly, when $x = \pi/2$, i.e., the input state is maximally entangled, $(1-p_{abs}) = (1-p_{sep})$. The gap obtained between the threshold values of noise for separability and absolute separability when the input state is a nonmaximally entangled state, can be interesting. We know that nonmaximally entangled states are less useful than maximally entangled states in several quantum information protocols. Nonetheless, such disadvantages can

be compensated. Since in presence of certain amount of noise, separable states that are not absolutely separable are created which can be converted to entangled states under global unitary operations. This is shown in the Table I.

We also observe that if the entanglement of the input state decreases, the output state becomes separable quickly under the action of local DPCs. But lowering the input state entanglement, the range of the noise in which the output state remains only separable (not absolutely separable) increases. Specifically, for maximally entangled and nonmaximally entangled states, denoted by $|\psi_{max}\rangle$ and $|\psi_{nm}\rangle$ respectively, we obtain that

$$(1 - p_{sep})(|\psi_{max}\rangle) > (1 - p_{sep})(|\psi_{nm}\rangle)$$

while

$$(1 - p_{abs})(|\psi_{max}\rangle) < (1 - p_{abs})(|\psi_{nm}\rangle).$$

Ent. of input	$1 - p_{sep}$	$1 - p_{abs}$	$\Delta p = p_{sep} - p_{abs}$
0.7715	0.29133	0.36114	0.0698
0.33225	0.21413	0.426103	0.21197

TABLE I. Entanglement of the input state is measured in von Neumann entropy [37]. The table shows that with the decrease of entanglement of the input state, the critical noise making a state separable but not absolute separable increases.

Remark 3. A maximally entangled state sent through a global depolarizing channel results the Werner state which is separable as well as absolutely separable with $p \leq 1/3$ [38]. On the other hand, if nonmaximally entangled state is mixed with the white noise, the gap in the strength of noise between separability and absolute separability emerges like the local depolarizing channels, establishing the usefulness of nonmaximally entangled states over the maximally entangled ones.

2. Absolutely separable states via amplitude damping channels

Let us now move to the scenario where two local ADCs are acted on $|\psi\rangle$, resulting to an output state of rank-4. It turns out that the output state is separable when $\tan(x/2) \geq 1/(1-p)$. By using Eq. (1), we can find the condition on p and x for which the resulting state is absolutely separable. Here also the eigenvalues are independent of the phase of the initial state.

Unlike the depolarizing channels, we find that the final state is absolutely separable only when $x \geq 2.325$. The criteria for absolute separability is satisfied in the




FIG. 2. (Color online) Regions in (x, p) plane indicates separable and absolutely separable states when $|\psi\rangle$ is sent through local amplitude damping channels. The blue colored region is for separability while the yellowish region represents states that are absolutely separable.

neighborhood of $p = 0.5$. The range of p in which the state is absolutely separable increases with the increase of x and becomes maximum when $x = \pi$. For example, when $x = \pi$, the state is separable in the entire range of p although it is absolutely separable when $p \in [0.302, 0.6998]$. For the input state with $x = 2.4$, the state remains absolutely separable when $p \in [0.414, 0.586]$ and separable for $0 \leq p \leq 0.61$. Therefore, like depolarizing channels, there also exists a range of parameter p in which the state is separable, but not absolutely separable as depicted in Fig. 2. However, the absolute separability of the resulting state with respect to (fixed) noise in this case does not depend on the entanglement of the input state which is in a sharp contrast of DPC.

3. Absolutely separable states via phase damping channels

Phase damping channels make the pure rank-1 state to a rank-2 one and hence the output state produced after PDCs cannot be absolutely separable, as discussed in Sec. III. To obtain absolutely separable state by using PDCs, we either consider rank-2 state which can produce rank-3 state with equal mixing parameter or we can start with a rank-3 or a rank-4 state that can be absolutely separable.

To illustrate this feature, let us first consider a rank-3 state, given by

$$\rho_3 = \frac{1}{3}|\psi\rangle\langle\psi| + \frac{1}{3}|01\rangle\langle 01| + \frac{1}{3}|10\rangle\langle 10|, \quad (10)$$




FIG. 3. (Color online) Map of separable (blue) and absolutely separable (yellow) states for local PDC with Werner state as input. The abscissa and ordinate respectively represent the noise parameter, p and the mixing parameter, q , of the Werner state.

where $|\psi\rangle = \cos(x/2)|00\rangle + e^{-i\phi}\sin(x/2)|11\rangle$. By Proposition 1, it is an extreme point of AS. If one sends the state through local PDCs, it is easy to check that the state becomes rank-4 except $x = 0$ as well as $x = \pi$ and remains absolutely separable, i.e., it satisfies the condition (1) in a strict sense. Note that at $x = 0$ and $x = \pi$, the state remains unaffected by the PDCs.

As a second example, let us consider the initial state for the PDCs as the Werner state, given by $q|\phi^+\rangle\langle\phi^+| + (1-q)\frac{\mathbb{I}}{4}$ which is of rank-4. It is known that the state is separable as well as absolutely separable when $q \leq 1/3$. When the state is sent through two noisy PDC channels, entanglement gets destroyed and hence the state becomes separable even for $q > 1/3$. We find that the state becomes separable against q in a bigger range than the value of q below which it is absolutely separable, as shown in Fig. 3. Therefore, phase damping noise introduces a gap between separable and absolutely separable regions for Werner states.

V. DISCUSSION

Absolute separable states are those which cannot be made entangled by the action of global unitary operations. Therefore, from the resource theoretic perspective, characterization of the set of useless states aka absolute separable states is an important enterprise. In this work, we considered $2 \otimes d$ dimensional states and showed that absolutely separable states of rank- $(2d-1)$ are all extreme points of the set of such states. We proved that the states with full-rank satisfying strict ab-

solute separability condition are the interior points. We also showed that there exists full rank states which are the boundary and the extreme points of the set.

We showed a possible method to make absolutely separable states entangled by adding auxiliary system. We also considered the effect of local noise on pure as well as mixed input entangled states. We found the

range of noise parameter which can produce absolutely separable states. We also showed that after sending maximally entangled states via local depolarizing channel, threshold noise value producing separable and absolute separable states coincide while with the decrease of entanglement content of the input pure state, the gap between these two critical values increases.

-
- [1] J. Preskill, Quantum computation and quantum information , Lecture notes for physics 229 (1998).
- [2] M. A. Nielsen and I. L. Chuang, Quantum computation and quantum information , Cambridge University Press (2000).
- [3] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, *Phys. Rev. Lett.* **70**, 1895 (1993).
- [4] D. Bouwmeester, J.-W. Pan, K. Mattle, M. Eibl, H. Weinfurter, and A. Zeilinger, *Nature* **390**, 575 (1997).
- [5] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Rev. Mod. Phys.* **81**, 865 (2009).
- [6] O. Gühne and G. Tóth, *Phys. Rep.* **474**, 1 (2009).
- [7] C. H. Bennett and S. J. Wiesner, *Phys. Rev. Lett.* **69**, 2881 (1992).
- [8] K. Mattle, H. Weinfurter, P. G. Kwiat, and A. Zeilinger, *Phys. Rev. Lett.* **76**, 4656 (1996).
- [9] A. K. Ekert, *Phys. Rev. Lett.* **67**, 661 (1991).
- [10] K. Boström and T. Felbinger, *Phys. Rev. Lett.* **89**, 187902 (2002).
- [11] A. Beige, B.-G. Englert, C. Kurtsiefer, and H. Weinfurter, *Acta Phys. Pol. A* **101**, 357 (2002).
- [12] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, *Rev. Mod. Phys.* **74**, 145 (2002).
- [13] V. Scarani, H. Bechmann-Pasquinucci, N. J. Cerf, M. Dušek, N. Lütkenhaus, and M. Peev, *Rev. Mod. Phys.* **81**, 1301 (2009).
- [14] R. Raussendorf and H. J. Briegel, *Phys. Rev. Lett.* **86**, 5188 (2001).
- [15] M. Hein, J. Eisert, and H. J. Briegel, *Phys. Rev. A* **69**, 062311 (2004).
- [16] H. J. Briegel, D. E. Browne, W. Dür, R. Raussendorf, and M. Van den Nest, *Nat. Phys.* **5**, 19 (2009).
- [17] B. Kraus and J. I. Cirac, *Phys. Rev. A* **63**, 062309 (2001).
- [18] M. S. Leifer, L. Henderson, and N. Linden, *Phys. Rev. A* **67**, 012306 (2003).
- [19] A. Chefles, *Phys. Rev. A* **72**, 042332 (2005).
- [20] M. Kuś and K. Życzkowski, *Phys. Rev. A* **63**, 032307 (2001).
- [21] F. Verstraete, K. Audenaert, and B. De Moor, *Phys. Rev. A* **64**, 012316 (2001).
- [22] R. Hildebrand, *Phys. Rev. A* **76**, 052325 (2007).
- [23] P. B. Slater, *Journal of Geometry and Physics* **59**, 17 (2009).
- [24] N. Johnston, *Phys. Rev. A* **88**, 062330 (2013).
- [25] N. Ganguly, J. Chatterjee, and A. S. Majumdar, *Phys. Rev. A* **89**, 052304 (2014).
- [26] S. Arunachalam, N. Johnston, and V. Russo, *Quant. Inf. Comput.* **15**, 0694 (2015).
- [27] M. A. Jivulescu, N. Lupa, and I. Nechita, *Quant. Inf. Comput.* **15**, 1165 (2015).
- [28] M. Horodecki, P. Horodecki, and R. Horodecki, *Phys. Lett. A* **223**, 1 (1996).
- [29] M. Lewenstein, B. Kraus, J. I. Cirac, and P. Horodecki, *Phys. Rev. A* **62**, 052310 (2000).
- [30] B. M. Terhal, *Linear Algebra and its Applications* **323**, 61 (2001).
- [31] R. B. Holmes, *Geometric Functional Analysis and its Applications*, Springer (1975).
- [32] M. Krein and D. Milman, *Studia Mathematica* **9**, 133 (1940).
- [33] K. Życzkowski, P. Horodecki, A. Sanpera, and M. Lewenstein, *Phys. Rev. A* **58**, 883 (1998).
- [34] L. Gurvits and H. Barnum, *Phys. Rev. A* **66**, 062311 (2002).
- [35] S. L. Braunstein, C. M. Caves, R. Jozsa, N. Linden, S. Popescu, and R. Schack, *Phys. Rev. Lett.* **83**, 1054 (1999).
- [36] S. N. Filippov, K. Y. Magadov, and M. A. Jivulescu, *New J. of Phys.* **19**, 083010 (2017).
- [37] C. H. Bennett, S. Popescu, D. Rohrlich, J. A. Smolin, and A. V. Thapliyal, *Phys. Rev. A* **63**, 012307 (2000).
- [38] R. F. Werner, *Phys. Rev. A* **40**, 4277 (1989).