

Radial flow induced by inhomogeneous magnetic field in heavy ion collisions

M. Haddadi Moghaddam^{1,2}, B. Azadegan², A. F. Kord², and W. M. Alberico¹

¹ Department of Physics, University of Turin and INFN, Turin, Via P. Giuria 1, I-10125 Turin, Italy

² Department of Physics, Hakim Sabzevari University (HSU), P.O.Box 397, Sabzevar, Iran

Abstract. In this paper, we study the effects of an in-homogeneous magnetic field on the quark gluon plasma dynamics, within the Magneto-Hydrodynamic framework. We have investigated the effect of an inhomogeneous external magnetic field on the transverse expansion of in-viscid fluid created in high energy nuclear collisions. Transverse velocity and energy density are modified by the presence of the magnetic field. This effects can also influence the transverse momentum spectrum for particles at the freeze-out surface. In this context we obtained an interesting comparison with data extracted from heavy-ion collisions.

Keywords: Heavy ions collision, Magneto-hydrodynamic

1 Introduction

Collisions of two heavy nuclei at high energy produce a hot and dense fireball. Quarks and gluons could reach the deconfined state, called quark gluon plasma (QGP), in a very short time ($\sim 1\text{fm}/c$) after the initial hard parton collisions of nuclei. recently a wide range of studies has shown that relativistic heavy-ion collisions create huge magnetic field due to the relativistic motion of the colliding heavy ions carrying large positive electric charge (For more references and details refer to it [1]-[3].).

Recently, some efforts in numerical and analytical works have been made, based on the relativistic magneto-hydrodynamic (RMHD) setup, to describe high energy heavy ion collisions (See, for example, [4]-[11]). The aim of our work is to generalize the Bjorken model by considering an inhomogeneous external magnetic field acting on the medium. We show that the presence of the magnetic field leads to non-zero radial flow [9].

2 Relativistic magneto-hydrodynamic

We deal with the case of an ideal non-resistive plasma, with vanishing electric field in the local rest-frame ($e^\mu = 0$), which is embedded in an external magnetic field (b_μ) [12]-[13]. The energy momentum conservation equations read:

$$d_\mu(T_{pl}^{\mu\nu} + T_{em}^{\mu\nu}) = 0, \quad (1)$$

where

$$T_{pl}^{\mu\nu} = (\epsilon + P)u^\mu u^\nu + P g^{\mu\nu} \quad (2)$$

$$T_{em}^{\mu\nu} = b^2 u^\mu u^\nu + \frac{1}{2} b^2 g^{\mu\nu} - b^\mu b^\nu. \quad (3)$$

In the above $g_{\mu\nu}$ is the metric tensor, ϵ and P are the energy density and pressure, respectively. Moreover d_μ is the covariant derivative. And the four velocity is defined as

$$u_\mu = \gamma(1, \mathbf{v}), \quad \gamma = \frac{1}{\sqrt{1 - v^2}}$$

satisfying the condition $u^\mu u_\mu = -1$.

Canonically one takes projections of the equation $d_\mu(T_{pl}^{\mu\nu} + T_{em}^{\mu\nu}) = 0$ along the parallel and perpendicular directions to u_ν , which gives:

$$D(\epsilon + b^2/2) + (\epsilon + P + b^2)\Theta + u_\nu b^\mu (d_\mu b^\nu) = 0, \quad (4)$$

$$(\epsilon + P + b^2)Du^\alpha + \nabla^\alpha(P + \frac{1}{2}b^2) - d_\mu(b^\mu b^\alpha) - u^\alpha u_\nu d_\mu(b^\mu b^\nu) = 0. \quad (5)$$

Notice that α should be a spacelike index. Moreover

$$D = u^\mu d_\mu, \quad \Theta = d_\mu u^\mu, \quad \nabla^\alpha = \Delta^\alpha_\nu d^\nu. \quad (6)$$

2.1 Induced radial flow in B-field

We consider the medium expands both radially and along the beam axis, the only nonzero components of $u_\mu = (u_\tau, u_\perp, 0, 0)$ are u_τ , which describes the boost-invariant longitudinal expansion [14], and u_\perp , which describes the transverse expansion. And we suppose the external magnetic field to be located in transverse plane as $b_\mu = (0, 0, b_\phi, 0)$.

We now seek the perturbation solution in the presence of a weak external magnetic field pointing along the ϕ direction in an inviscid fluid with infinite electrical conductivity:

$$u_\mu = (1, \lambda^2 u_\perp, 0, 0), \quad b_\mu = (0, 0, \lambda b_\phi, 0), \quad b^2 \equiv b^\mu b_\mu \\ \epsilon = \epsilon_0(\tau) + \lambda^2 \epsilon_1(\tau, x_\perp), \quad \epsilon_0(\tau) = \frac{\epsilon_c}{\tau^{4/3}} \quad (7)$$

In such setup, the conservation equations (4-5) reduce to the following partial differential equations

$$u_\perp - \tau^2 \partial_\perp \left(\frac{u_\perp}{x_\perp} \right) - \tau^2 \partial_\perp^2 u_\perp - \tau \partial_\tau u_\perp + 3\tau^2 \partial_\tau^2 u_\perp \\ - \frac{3\tau^{7/3}}{x_\perp \epsilon_c} b_\phi^2 - \frac{3\tau^{7/3}}{4\epsilon_c} \partial_\perp b_\phi^2 - \frac{9\tau^{10/3}}{4x_\perp \epsilon_c} \partial_\tau b_\phi^2 - \frac{3\tau^{10/3}}{4\epsilon_c} \partial_\perp \partial_\tau b_\phi^2 = 0. \quad (8)$$

For non-vanishing b_ϕ we assume a space-time profile of the magnetic field in central collisions in the form:

$$b_\phi^2(\tau, x_\perp) = B_c^2 \tau^{-1} \sqrt{\alpha} x_\perp e^{-\alpha x_\perp^2}. \quad (9)$$

We see that the magnitude of b_ϕ is zero at $x_\perp = 0$. Finally, one can find the solutions for transverse velocity u_\perp and correspondingly modified energy density ϵ_1 ; for more details refer to [9].

3 Particle transverse momentum spectrum

From the local equilibrium hadron distribution the transverse spectrum is calculated via the Cooper-Frye formula in the freeze out surface

$$S = E \frac{d^3 N}{dp^3} = \frac{g_i}{2\pi^2} \int_0^{x_f} x_\perp \tau_f(x_\perp) dx_\perp \left[m_T K_1\left(\frac{m_T u_\tau}{T_f}\right) I_0\left(\frac{m_T u_\perp}{T_f}\right) + p_T R_f K_0\left(\frac{m_T u_\tau}{T_f}\right) I_1\left(\frac{m_T u_\perp}{T_f}\right) \right] \quad (10)$$

Where $\tau_f(x_\perp)$ is the solution of the $T(\tau_f, x_\perp) = T_f$ and the degeneracy is $g_i = 2$ for both the pions and the protons. The above integral over x_\perp on the freeze-out surface is evaluated numerically.

The spectrum Eq. (10) is illustrated in the following figures for three different values of the freeze out temperature (140, 150 and 160 MeV) and compared with experimental results obtained at PHENIX [15]. in central collisions. Our proton spectrum appear to underestimate the experimental data, except at low p_T , but their behavior with p_T has the correct trend of a monotonically decrease. The pion spectrum, instead, appears in fair agreement with the experimental results, which are very close to the theoretical curves. This is an indication that hadrons with different masses have different sensitivities to the underlying hydrodynamic flow and to the electromagnetic fields. Indeed, the difference between the charge-dependent flow of light pions and heavy protons might arise because the former are more affected by the weak magnetic field than the heavy protons [16].

For comparison, we also show the results obtained by Gubser [17], which appear to be more flat and typically overestimate the experiment. We also notice that, for the proton case, the highest value of the freeze out temperature we employed (as suggested, e.g. in Ref. [18]) slightly brings (for protons) the calculation closer to the experimental data; however it also shows a kind of saturation phenomenon and points to the need of including other effects not considered in the present work.

References

1. K. Tuchin, "Time and space dependence of the electromagnetic field in relativistic heavy-ion collisions," *Phys. Rev. C* **88** (2013) no.2, 024911.
2. H. Li, X-L. Sheng, Q. Wang, " Electromagnetic fields with electric and chiral magnetic conductivities in heavy ion collisions ", *Phys. Rev. C* **94**, 044903 (2016)
3. U. Gursoy, D. Kharzeev and K. Rajagopal, "Magnetohydrodynamics, charged currents and directed flow in heavy ion collisions," *Phys. Rev. C* **89** (2014) no.5, 054905
4. Shi Pu, and Di-Lun Yang, " Transverse flow induced by inhomogeneous magnetic fields in the Bjorken expansion ", *Phys. Rev. D* **93**, 054042 (2016).




Fig. 1. Particles transverse momentum spectrum from central Au-Au collisions: black, purple and red lines correspond to a freeze out temperature of 140, 150 and 160 MeV, respectively. Circles: PHENIX data [15].

5. V. Roy, S. Pu, L. Rezzolla, D. Rischke, "Analytic Bjorken flow in one-dimensional relativistic magnetohydrodynamics", *Physics Letters B*, Vol. 750, (2015).
6. S. Pu, V. Roy, L. Rezzolla and D. H. Rischke, "Bjorken flow in one-dimensional relativistic magnetohydrodynamics with magnetization", *Phys. Rev. D* 93, 074022 (2016).
7. G. Inghirami, L. Del Zanna, A. Beraudo, M. Haddadi Moghaddam, F. Becattini, M. Bleicher, "Numerical magneto-hydrodynamics for relativistic nuclear collisions", *Eur. Phys. J. C* (2016) 76:659.
8. M. H. Moghaddam, B. Azadegan, A. F. Kord, W. M. Alberico, "Non-relativistic approximate numerical ideal-magneto-hydrodynamics of (1+1D) transverse flow in Bjorken scenario", *Eur. Phys. J. C* (2018) 78:255.
9. M. H. Moghaddam, B. Azadegan, A. F. Kord, W. M. Alberico, "Transverse expansion of hot magnetized Bjorken flow in heavy ion collisions", *Eur. Phys. J. C* (2019) 79:619.
10. V. Roy, S. Pu, L. Rezzolla and D.H. Rischke, "Effects of intense magnetic fields on reduced-MHD evolution in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions", *Phys. Rev. C* 96, 054909 (2017).
11. A. Das, S.S. Dave, P.S. Saumia and A. M. Srivastava, "Effects of magnetic field on the plasma evolution in relativistic heavy-ion collisions", *Phys. Rev. C* 96, 034902 (2017).
12. J. Goedbloed, R. Keppens, S. Poedts "Advanced Magnetohydrodynamics with Applications to Laboratory and Astrophysical Plasmas", Cambridge University Press, 2010.
13. A. M. Anile, "Relativistic fluids and magneto-fluids", Cambridge University Press (1989).
14. J. D. Bjorken, *Phys. Rev. D* 27, 140 (1983).
15. K. Adcox et al. (PHENIX Collaboration), "Formation of dense partonic matter in relativistic nucleus-nucleus collisions at RHIC: Experimental evaluation by the PHENIX Collaboration" *Nucl. Phys. A* 757, 184 (2005)
16. U. Gursoy, D. Kharzeev, E. Marcus, K. Rajagopal and C. Shen, "charge dependent flow induced by magnetic and electric field in heavy ion collisions," *Phys. Rev. C* **98**, 055201 (2018).
17. S. S. Gubser, "Symmetry constraints on generalizations of Bjorken flow", *Phys. Rev. D* 82, 085027 (2010).
18. C. Ratti, R. Bellwied, J. Noronha-Hostler, P. Parotto, I. Portillo Vazquez and J. M. Stafford, arXiv:1805.00088 [hep-ph].