

# NEW INEQUALITIES FOR THE FUNCTION $y = t \ln t$

MARKO KOSTIĆ

**ABSTRACT.** The main aim of this note, which can be viewed as a certain addendum to the paper [1], is to propose several new inequalities for the function  $y = t \ln t$ . We consider the local behaviour of this function near the point  $t = 1$ , as well as the global behaviour of this function on the intervals  $[1, \infty)$  and  $(0, 1]$ .

*AMS Mathematics Subject Classification* (2010): 26D20, 26D07, 33B30.

*Key words and phrases:* Logarithmic inequalities, differential calculus.

## 1. INTRODUCTION

The reading of paper [1] by C. Chesneau and Y. J. Bagul has strongly influenced us to write this note. In Theorem 2.1, we give new abstract local bounds for the function  $y = t \ln t$  near the point  $t = 1$ . The obtained inequalities can be used to improve the main results of paper [1], Proposition 1 and Proposition 2. We also present an interesting result with regards to these propositions, which claims that there is no rational real function which intermediates the functions  $\ln(1+x)$  and  $f(x)/\sqrt{x+1}$  for  $x \geq 0$  ( $x \in (-1, 0]$ ); here and hereafter,

$$f(x) := \pi + \frac{1}{2}(4 + \pi)x - 2(x + 2) \arctan \sqrt{x + 1}, \quad x \geq -1.$$

The following inequalities are well known (see also [3, Problem 3.6.19, p. 274] and [4]):

$$(1.1) \quad \ln(1+x) \leq \frac{x}{\sqrt{x+1}}, \quad x \geq 0, \quad \ln(1+x) \leq \frac{x(2+x)}{2(1+x)}, \quad x \geq 0,$$

$$(1.2) \quad \ln(1+x) \leq \frac{x(6+x)}{2(3+2x)}, \quad x \geq 0 \quad \text{and} \quad \ln(1+x) \leq \frac{(x+2)[(x+1)^3 - 1]}{3(1+x)[(x+1)^2 + 1]}, \quad x \geq 0.$$

Taken together, the first inequality in (1.1) and the second inequality in (1.2) are known in the existing literature as Karamata's inequality [2]. As clarified in [1], all these inequalities are weaker than the inequality:

$$(1.3) \quad \ln(1+x) \leq \frac{f(x)}{\sqrt{x+1}}, \quad x \geq 0.$$

This inequality has been proved in [1, Proposition 1]. In [1, Proposition 2], the authors have proved that

$$(1.4) \quad \ln(1+x) \geq \frac{f(x)}{\sqrt{x+1}}, \quad x \in (-1, 0],$$

as well.

Our approach leans heavily on the use of substitution  $t = \sqrt{x+1}$ . Then the inequalities (1.3) and (1.4) become

$$2 \ln t \leq \frac{f(t^2-1)}{t}, \quad t \geq 1 \quad \text{and} \quad 2 \ln t \geq \frac{f(t^2-1)}{t}, \quad t \in (0, 1],$$

i.e.,

$$(1.5) \quad 2t \ln t \leq f(t^2-1), \quad t \geq 1 \quad \text{and} \quad 2t \ln t \geq f(t^2-1), \quad t \in (0, 1].$$

We can prove (1.5) in the following way. Notice that

$$\left[ \ln t - \left( \frac{1}{2}(4+\pi)t - 2t \arctan t - 2 \right) \right]''(t) = -(t^2-1)^2 t^{-2} (t^2+1)^{-2}, \quad t > 0.$$

Using an elementary argumentation, this estimate implies

$$\ln t \leq \frac{1}{2}(4+\pi)t - 2t \arctan t - 2, \quad t > 0.$$

Define  $R(t) := 2t \ln t - f(t^2-1)$ ,  $t > 0$ . Since  $R'(t) = 2(1+\ln t) - (4+\pi)t + 4t \arctan t + 2$ ,  $t > 0$ , the previous inequality yields  $R'(t) \leq 0$ ,  $t > 0$  and (1.5). Moreover, by taking the limit of function  $R(\cdot)$  as  $t \rightarrow 0+$ , we get that  $2t \ln t - f(t^2-1) \in (2 - (\pi/2), 0]$  for  $t \in (0, 1]$ .

In this paper, we will first generalize the inequalities in (1.5) by considering the local behaviour of the function  $y = t \ln t$  near the point  $t = 1$ . We will use the following simple lemmata, which is known from the elementary courses of mathematical analysis:

**Lemma 1.1.** *Suppose  $t_0 \in \mathbb{R}$ ,  $a > 0$ ,  $n \in \mathbb{N}$  and function  $f : (t_0 - a, t_0 + a) \rightarrow \mathbb{R}$  is  $2n$ -times differentiable. If  $f^{(i)}(t_0) = 0$  for all  $i = 1, \dots, 2n-1$  and  $f^{(2n)}(t_0) > 0$  ( $f^{(2n)}(t_0) < 0$ ), then the function  $y = f(t)$  has a local minimum (maximum) at  $t = t_0$ .*

**Lemma 1.2.** *We have*

$$(\arctan x)^{(n)} = \frac{(-1)^{n-1}(n-1)!}{(1+x^2)^{n/2}} \sin(n\pi/2 - n \arctan x), \quad x \in \mathbb{R}, \quad n \in \mathbb{N}.$$

After that, we will prove the following result with regards to [1, Proposition 1, Proposition 2]:

**Theorem 1.3.** (i) *There do not exist real polynomials  $P(\cdot)$  and  $Q(\cdot)$  such that  $Q(x) \neq 0$  for  $x \geq 0$  and*

$$(1.6) \quad \ln(1+x) \leq \frac{P(x)}{Q(x)} \leq \frac{f(x)}{\sqrt{x+1}}, \quad x \geq 0.$$

(ii) *There do not exist real polynomials  $P(\cdot)$  and  $Q(\cdot)$  such that  $Q(x) \neq 0$  for  $x \in (-1, 0]$  and*

$$(1.7) \quad \ln(1+x) \geq \frac{P(x)}{Q(x)} \geq \frac{f(x)}{\sqrt{x+1}}, \quad x \in (-1, 0].$$

## 2. THE MAIN RESULTS AND THEIR PROOFS

We start this section by stating the following result:

**Theorem 2.1.** *Suppose that  $a \in (0, 1)$ ,  $P : (1 - a, 1 + a) \rightarrow \mathbb{R}$  is a function and  $P(1) = 0$ . Then the following holds:*

- (i) *If  $P'(1) \geq 2$  and there exists an odd natural number  $n$  such that  $P(\cdot)$  is  $(n + 2)$ -times differentiable,  $P^{(n+2)}(1) + 2(-1)^{n+1}n! > 0$  and*

$$P^{(j)}(1) + 2(-1)^{j+1}(j - 2)! = 0 \quad \text{for all } j = 2, 3, \dots, n + 1,$$

*then there exists a real number  $\zeta \in (0, a]$  such that*

$$(2.1) \quad 2t \ln t \leq P(t), \quad t \in [1, 1 + \zeta] \quad \text{and} \quad 2t \ln t \geq P(t), \quad t \in [1 - \zeta, 1].$$

- (ii) *Assume that there exists an even natural number  $n \geq 6$  such that  $P(\cdot)$  is  $(n + 1)$ -times differentiable,  $P^{(n+1)}(1) + 2(-1)^n(n - 1)! > 0$  and*

$$P^{(j)}(1) + 2(-1)^{j+1}(j - 2)! = 0 \quad \text{for all } j = 1, 2, \dots, n.$$

*Then there exists a real number  $\eta \in (0, a]$  such that*

$$(2.2) \quad \begin{aligned} 2t \ln t \leq P(t) \leq f(t^2 - 1), \quad t \in [1, 1 + \eta] \\ \text{and} \quad 2t \ln t \geq P(t) \geq f(t^2 - 1), \quad t \in [1 - \eta, 1]. \end{aligned}$$

- (iii) *Assume that there exists an even natural number  $n \geq 6$  such that  $P(\cdot)$  is  $(n + 1)$ -times differentiable,*

$$(2.3) \quad P^{(j)}(1) + 2(-1)^{j+1}(j - 2)! = 0 \quad \text{for all } j = 1, 2, 3, 4,$$

$$P^{(n+1)}(1) + 4 \left[ \frac{(-1)^n n!}{2^{(n+1)/2}} \sin((n + 1)\pi/4) + \frac{(-1)^{n+1} n!}{2^{n/2}} \sin(n\pi/4) \right] < 0$$

*and, for every  $j = 5, 6, \dots, n$ ,*

$$(2.4) \quad P^{(j)}(1) + 4 \left[ \frac{(-1)^{j-1}(j - 1)!}{2^{j/2}} \sin(j\pi/4) + \frac{(-1)^j(j - 1)!}{2^{(j-1)/2}} \sin((j - 1)\pi/4) \right] = 0.$$

*Then there exists a real number  $\eta \in (0, a]$  such that (2.2) holds.*

- (iv) *If  $P(\cdot)$  is five times differentiable, (2.3) holds and  $P^{(v)}(1) \in (-12, -8)$ , then there exists a real number  $\eta \in (0, a]$  such that (2.2) holds.*

*Proof.* Define  $G(t) := P(t) - 2t \ln t$ ,  $t > 0$ . Then, for every real number  $t > 0$ , we have  $G'(t) = P'(t) - 2(1 + \ln t)$ ,  $G''(t) = P''(t) - (2/t)$  and  $G^{(n)}(t) = P^{(n)}(t) + 2(-1)^{n+1}(n - 2)! \cdot t^{1-n}$ ,  $n \geq 3$ . The assumptions made in (i) imply that  $G'(1) \geq 0$ ,  $(G')^{(j)}(1) = 0$  for  $1 \leq j \leq n$  and  $(G')^{(n+1)}(1) > 0$ . Applying Lemma 1.1, we get that the function  $t \mapsto G'(t)$  has a local minimum at  $t = 1$ . Since  $G'(1) \geq 0$ , we get that the function  $t \mapsto G'(t)$  is non-negative in an open neighborhood of point  $t = 1$ , so that the mapping  $t \mapsto G(t)$  is increasing in an open neighborhood of point  $t = 1$ . This finishes the proof of (i). For the proof of (ii), define  $Q(t) := P(t) - f(t^2 - 1)$ ,  $t > 0$ . Then a simple computation yields that, for every real number  $t > 0$ , we have  $Q'(t) = P'(t) - (4 + \pi)t + 4t \arctan t + 2$  and  $Q''(t) = P''(t) - (4 + \pi) + 4 \arctan t + \frac{4t}{t^2 + 1}$ .

Using Leibniz rule and Lemma 1.2, for every real number  $t > 0$  and for every natural number  $n \geq 3$ , we can show that

$$\begin{aligned} Q^{(n)}(t) &= P^{(n)}(t) + 4[\cdot \arctan \cdot]^{(n-1)}(t) \\ &= P^{(n)}(t) + 4 \left[ t \frac{(-1)^{n-1}(n-1)!}{(1+t^2)^{n/2}} \sin(n\pi/2 - n \arctan t) \right. \\ &\quad \left. + \frac{(-1)^n(n-1)!}{(1+t^2)^{(n-1)/2}} \sin((n-1)\pi/2 - (n-1) \arctan t) \right]. \end{aligned}$$

Arguing as in the proof of (i), we have that  $(Q')^{(j)}(1) = 0$  for  $j = 0, 1, 2, 3$  and  $(Q')^{(4)}(1) < 0$ ; hence, the function  $t \mapsto Q'(t)$  has a local maximum at  $t = 1$  and the mapping  $t \mapsto Q(t)$  is decreasing in an open neighborhood of point  $t = 1$ . Similarly,  $(G')^{(j)}(1) = 0$  for  $j = 0, 1, 2, \dots, n-1$  and  $(G')^{(n)}(1) > 0$ ; hence, the function  $t \mapsto G'(t)$  has a local minimum at  $t = 1$  and the mapping  $t \mapsto G(t)$  is increasing in an open neighborhood of point  $t = 1$ . This completes the proof of (ii). The proof of (iii) can be deduced similarly, by interchanging the roles of  $G(t)$  and  $Q(t)$ . If the assumptions of (iv) holds, then we can apply Lemma 1.1, with  $n = 2$ , in order to see that the function  $t \mapsto G'(t)$  has a local minimum at  $t = 1$ , as well as the function  $t \mapsto G'(t)$  is non-negative in an open neighborhood of point  $t = 1$ ; hence, the mapping  $t \mapsto G(t)$  is increasing in an open neighborhood of point  $t = 1$ . Similarly, we can show that the mapping  $t \mapsto Q(t)$  is decreasing in an open neighborhood of point  $t = 1$ . The proof of the theorem is thereby complete.  $\square$

*Remark 2.2.* Define  $H(t) := f(t^2 - 1)$ ,  $t \in \mathbb{R}$ . Concerning the conditions used in Theorem 2.1, it is worth noting that the function  $H(\cdot)$  satisfies  $H(1) = 0$ ,  $H'(1) = H''(1) = 2$ ,  $H'''(1) = -2$ ,  $H^{(iv)}(1) = 4$  and  $H^{(v)}(1) = -8$ . This implies that the values of terms appearing at the right hand sides of (2.3) and (2.4) coincide for  $j = 1, 2, 3, 4$  and differ for  $j = 5$  (observe that  $G^{(v)}(1) = P^{(v)}(1) + 12$ ).

*Remark 2.3.* The parts (ii)-(iv) of Theorem 2.1 ensure the existence of a large class of elementary functions for which we can further refine the inequalities in (1.5) locally around the point  $t = 1$ . Compared with the function  $H(\cdot)$ , the most simplest example of function which provides a better estimate describing the local behaviour of function  $y = t \ln t$  around the point  $t = 1$  is given by the function  $t \mapsto H(t) - \epsilon(t-1)^5$ ,  $t > 0$ , where  $\epsilon \in (0, 1/30)$ .

Concerning the global behaviour of function  $y = t \ln t$ ,  $t > 0$ , it is clear that the inequalities in (1.5) give some very uninteresting estimates with regards to the asymptotic behaviour of function  $y = t \ln t$  when  $t \rightarrow +\infty$  or  $t \rightarrow 0+$ ; on the other hand, the importance of estimate (1.5) lies in the fact that it gives some bounds for the behaviour of function  $y = t \ln t$  on any compact interval  $[a, b]$ , where  $0 < a < 1 < b$ . It is clear that there exists a large class of infinitely differentiable functions  $P : (0, \infty) \rightarrow \mathbb{R}$  such that

$$(2.5) \quad \begin{aligned} 2t \ln t &\leq P(t) \leq f(t^2 - 1), \quad t \geq 1 \\ \text{and } 2t \ln t &\geq P(t) \geq f(t^2 - 1), \quad t \in (0, 1]. \end{aligned}$$

Finding new elementary functions  $P(\cdot)$  for which the equation (2.5) holds is without scope of this paper.

We close the paper by giving the proof of Theorem 1.3:

**Proof of Theorem 1.3.** Suppose that (1.6) holds for some real polynomials  $P(\cdot)$  and  $Q(\cdot)$  such that  $Q(x) \neq 0$  for  $x \geq 0$ . Without loss of generality, we may assume that  $Q(x) > 0$ ,  $x \geq 0$ . Using the substitution  $t = \sqrt{x+1}$ , we get that

$$2 \ln t \leq \frac{P(t^2 - 1)}{Q(t^2 - 1)} \leq \frac{f(t^2 - 1)}{t}, \quad t \geq 1.$$

If  $P(t) = \sum_{j=0}^n a_j t^j$  and  $Q(t) = \sum_{j=0}^m b_j t^j$  for some non-negative integers  $m, n$  and some real numbers  $a_j, b_j$  ( $a_n b_m \neq 0$ ; clearly, we cannot have  $P(x) \equiv 0$ ), we get

$$(2.6) \quad t \sum_{j=0}^n a_j (t^2 - 1)^j \leq f(t^2 - 1) \sum_{j=0}^m b_j (t^2 - 1)^j, \quad t \geq 1$$

and

$$(2.7) \quad \sum_{j=0}^n a_j (t^2 - 1)^j \geq 2 \ln t \sum_{j=0}^m b_j (t^2 - 1)^j, \quad t \geq 1.$$

Since  $f(t^2 - 1) \sim (2 - (\pi/2))t^2$ ,  $t \rightarrow +\infty$ , the estimate (2.6) implies  $n \leq m$ . The positivity of polynomial  $Q(\cdot)$  on the non-negative real axis implies  $b_m > 0$  so that (2.7) gives  $a_n > 0$ . Considering the asymptotic behaviour of terms appearing in (2.7), we get that the inequality  $n < m$  cannot be satisfied so that  $m = n$ . Dividing the both sides of (2.7) with  $t^{2n}$  and letting  $t \rightarrow +\infty$  in the obtained expression, we get that  $a_n/2b_n \geq +\infty$ , which is a contradiction. This completes the proof of (i). To prove (ii), suppose that the estimates

$$\ln(1+x) \geq \frac{P_0(x)}{Q_0(x)} \geq \frac{f(x)}{\sqrt{x+1}}, \quad x \in (-1, 0]$$

hold for some real polynomials  $P_0(\cdot)$  and  $Q_0(\cdot)$  such that  $Q_0(x) \neq 0$  for  $x \in (-1, 0]$ . Then (1.7) holds for some real polynomials  $P(\cdot)$  and  $Q(\cdot)$  such that  $Q(x) > 0$  for  $x \in (-1, 0]$ . Letting  $x \rightarrow -1-$  in (1.7), we get that  $Q(-1) = 0$ . If  $P(x) = \sum_{j=0}^n a_j x^j$  and  $Q(x) = \sum_{j=0}^m b_j x^j$  for some non-negative integers  $m, n$  and some real numbers  $a_j, b_j$  ( $a_n b_m \neq 0$ ; again, we cannot have  $P(x) \equiv 0$ ), this implies

$$(2.8) \quad \ln(1+x) \cdot \sum_{j=0}^m b_j x^j \geq \sum_{j=0}^n a_j x^j \geq \frac{f(x)}{\sqrt{x+1}} \sum_{j=0}^m b_j x^j, \quad x \in (-1, 0].$$

Letting  $x \rightarrow 0-$  in this expression, we get that  $a_0 = 0$  so that  $n \geq 1$  and  $x|P(x)$ . Define  $P_1(x) := P(x)/x$  and  $Q_1(x) := Q(x)/(x+1)$ . Then  $P_1(x)$  and  $Q_1(x)$  are real polynomials,  $Q_1(x) > 0$  for  $x \in (-1, 0]$  and after multiplication with  $\frac{x+1}{xQ(x)} \leq 0$  the estimate (2.8) implies

$$(2.9) \quad \frac{x+1}{x} \ln(1+x) \leq \frac{P_1(x)}{Q_1(x)} \leq \sqrt{x+1} \frac{f(x)}{x}, \quad x \in (-1, 0).$$

Letting  $x \rightarrow -1-$  in this expression, we get that  $\lim_{x \rightarrow -1-} \frac{P_1(x)}{Q_1(x)} = 0$ , which implies  $P_1(-1) = 0$ . Since  $P_1(x)$  is a non-zero polynomial, we get that  $x+1|P_1(x)$ . Multiplying the equation (2.9) with  $\frac{x}{x+1} \leq 0$ , we get

$$\ln(1+x) \geq \frac{P_1(x)}{Q_1(x)} \geq \frac{f(x)}{\sqrt{x+1}}, \quad x \in (-1, 0).$$

Letting  $x \rightarrow 0-$ , we get

$$\ln(1+x) \geq \frac{P_1(x)}{Q_1(x)} \geq \frac{f(x)}{\sqrt{x+1}}, \quad x \in (-1, 0].$$

Repeating this procedure, we get that for every natural number  $k$  we have  $(x+1)^k |Q(x)$ , which is a contradiction.

#### REFERENCES

- [1] C. Chesneau and Y. J. Bagul, *New sharp bounds for the logarithmic function*, Electronic J. Math. Anal. Appl. **8(1)** (2020), 140–145.
- [2] J. Karamata, *Sur quelques problèmes posés par Ramanujan*, J. Indian Math. Soc. (N.S.) **24** (1960), 343–365.
- [3] D. S. Mitrinović, *Analytic Inequalities*, Springer, Berlin, 1970.
- [4] F. Topsoe, *Some bounds for the logarithmic function*, In: Cho YJ, Kim JK, Dragomir SS Editors, Inequality theory and applications 4, New York: Nova Science Publishers, 137, 2007.

FACULTY OF TECHNICAL SCIENCES, UNIVERSITY OF NOVI SAD, TRG D. OBRADOVIĆA 6, 21125 NOVI SAD, SERBIA

*E-mail address:* marco.s@verat.net