# ON AUTOMATIC CONTINUITY OF THE DERIVATIONS ON MODULE EXTENSION OF BANACH ALGEBRAS

### H. FARHADI

ABSTRACT. In this article we shall focus on the derivations on module extension of Banach algebras and determine the general structure of them. Then we obtain some results concerning the automatic continuity of these mappings.

## 1. Introduction

Let  $\mathcal{A}$  be a Banach algebra (over the complex field  $\mathbb{C}$ ), and  $\mathcal{U}$  be a Banach  $\mathcal{A}$ bimodule. A linear mapping  $D: A \to \mathcal{U}$  is called a derivation if D(ab) = aD(b) +D(a)b holds for all  $a, b \in \mathcal{A}$ . If  $\mathcal{U} = \mathcal{A}$ , then D is said to be a derivation on Banach algebra  $\mathcal{A}$ . For any  $u \in \mathcal{U}$ , the mapping  $\mathrm{id}_u : \mathcal{A} \to \mathcal{U}$  given by  $\mathrm{id}_u(a) = au - ua$ is a continuous derivation called *inner derivation* induced by u. Derivations and their various properties are significant subjects in the study of Banach algebras. One of the most important problems related to these mappings is this question; Under what conditions is a derivation  $D: \mathcal{A} \to \mathcal{U}$  continuous? This question lies in the theory of automatic continuity which is an important subject in mathematical analysis and has attracted the attention of researchers during the last few decades. In this theory we are looking for conditions under which one can conclude that a linear mapping between two Banach algebras (or two Banach spaces, in general) is necessarily continuous. For more information, the reader may refer to [3] which is a detailed source in this context. Here we mention the most important established results in this regard. A celebrated theorem due to Johnson and Sinclair [17] states that every derivation on a semisimple Banach algebra is continuous. Later on, Ringrose in [21] showed that every derivation from a  $C^*$ -algebra  $\mathcal{A}$  into Banach  $\mathcal{A}$ -bimodule  $\mathcal{U}$  is continuous. In [2], Christensen proved that every derivation from nest algebra on Hilbert space  $\mathcal{H}$  into  $\mathbb{B}(\mathcal{H})$  is continuous. Additionally, some results on automatic continuity of the derivations on prime Banach algebras have been established by Villena in [26] and [27]. See also [4, 15] where the continuity of the derivations are studied on certain rpoducts of Banach algebras. Some results on the derivations and Jordan derivations on trivial extension and triangular Banach algebras have been established by in a number of papers; see [4–14]

Let  $\mathcal{A}$  be a Banach algebra and  $\mathcal{U}$  be a Banach  $\mathcal{A}$ -bimodule. The module extension or trivial extension Banach algebra associated to  $\mathcal{A}$  and  $\mathcal{U}$ , denoted by  $T(\mathcal{A},\mathcal{U})$ , is the  $\ell^1$ -direct sum  $\mathcal{A} \oplus \mathcal{U}$  equipped with the algebra multiplication given by

$$(a, u)(b, v) = (ab, av + ub)$$
  $(a, b \in \mathcal{A}, u, v \in \mathcal{U}).$ 

MSC(2010): 16E40; 46H40; 46H25.

Keywords: Module extension of Banach algebras, Derivation, Automatic continuity.

H. FARHADI

and the norm

2

$$||(a, u)|| = ||a|| + ||u||$$
  $(a \in \mathcal{A}, u \in \mathcal{U}).$ 

Many authors have studied this class of Banach algebras from different points of view and investigated various properties of these spaces; see for example [20, 28].

At the end of this section we introduce some notations and expressions used in the paper. Let  $\mathcal{A}$  be a Banach algebra and  $\mathcal{U}, \mathcal{V}$  be Banach  $\mathcal{A}$ -bimodules. The annihilator of  $\mathcal{U}$  over  $\mathcal{A}$  is defined as

$$ann_{\mathcal{A}}\mathcal{U} = \{ a \in \mathcal{A} \mid a\mathcal{U} = \mathcal{U}a = (0) \}.$$

Recall that a linear mapping  $\phi: \mathcal{U} \to \mathcal{V}$  is said to be a left  $\mathcal{A}$ -module homomorphism if  $\phi(au) = a\phi(u)$  whenever  $a \in \mathcal{A}$  and  $u \in \mathcal{U}$  and it is a right  $\mathcal{A}$ -module homomorphism if  $\phi(ua) = \phi(u)a$  ( $a \in \mathcal{A}, u \in \mathcal{U}$ ).  $\phi$  is called  $\mathcal{A}$ -module homomorphism, if  $\phi$  is both left and right  $\mathcal{A}$ -module homomorphism. The set of all continuous  $\mathcal{A}$ -module homomorphisms from  $\mathcal{U}$  into  $\mathcal{V}$  is denoted by  $Hom_{\mathcal{A}}(\mathcal{U}, \mathcal{V})$ .

## 2. Continuity of the derivations

Throughout this section  $\mathcal{A}, \mathcal{U}$  are Banach algebras and  $T(\mathcal{A}, \mathcal{U})$  denotes the corresponding module extension Banach algebra. We commence with the following proposition characterizing the derivations on  $T(\mathcal{A}, \mathcal{U})$ .

**Proposition 2.1.** Let  $D: T(A, \mathcal{U}) \to T(A, \mathcal{U})$  be a linear mapping such that

$$D((a, u)) = (\delta_1(a) + \tau_1(u), \delta_2(a) + \tau_2(u)) \qquad ((a, u) \in \mathcal{A} \times \mathcal{U})$$

in which  $\delta_1: \mathcal{A} \to \mathcal{A}$ ,  $\delta_2: \mathcal{A} \to \mathcal{U}$ ,  $\tau_1: \mathcal{U} \to \mathcal{A}$  and  $\tau_2: \mathcal{U} \to \mathcal{U}$  are linear mappings. Then the following are equivalent.

- (i) D is a derivation.
- (ii)  $\delta_1: \mathcal{A} \to \mathcal{A}, \ \delta_2: \mathcal{A} \to \mathcal{U}$  are derivations,  $\tau_2: \mathcal{U} \to \mathcal{U}$  is a linear mapping satisfying

$$\tau_2(au) = a\tau_2(u) + \delta_2(a)u$$
 and  $\tau_2(ua) = \tau_2(u)a + u\delta_2(a)$   $(a \in \mathcal{A}, u \in \mathcal{U}).$ 

 $\tau_1: \mathcal{U} \to \mathcal{A}$  is an  $\mathcal{A}$ -bimodule homomorphism for which  $u\tau_1(v) + \tau_1(u)v = 0$   $(u, v \in \mathcal{U})$ .

As a consequence, any derivation  $D: T(\mathcal{A}, \mathcal{U}) \to T(\mathcal{A}, \mathcal{U})$  can be written as  $D = D_1 + D_2$  where  $D_1((a, u)) = (\delta_1(a) + \tau_1(u), \tau_2(u))$  and  $D_2((a, u)) = (0, \delta_2(a))$  are derivations on  $T(\mathcal{A}, \mathcal{U})$ . Moreover, D is an inner derivation if and only if  $\delta_1, \delta_2$  are inner derivations and  $\tau_1 = 0$ .

*Proof.* The proof follows from a straightforward verification.

**Remark 2.2.** Note that one can show that the linear mapping  $\delta: \mathcal{A} \to \mathcal{U}$  is a derivation if and only if  $D: T(\mathcal{A}, \mathcal{U}) \to T(\mathcal{A}, \mathcal{U})$  given by  $D((a, u)) = (0, \delta(a))$  is a derivation. So if any derivation on  $T(\mathcal{A}, \mathcal{U})$  is continuous, then any derivation  $\delta: \mathcal{A} \to \mathcal{U}$  is continuous as well.

In the next proposition we give some sufficient conditions for derivations on  $\mathcal{A}$  be continuous.

**Proposition 2.3.** Suppose that there are A-bimodule homomorphisms  $\phi : A \to \mathcal{U}$  and  $\psi : \mathcal{U} \to A$  such that  $\phi o \psi = I_{\mathcal{U}}$  ( $I_{\mathcal{U}}$  is the identity mapping on  $\mathcal{U}$ ). If every derivation on  $T(A,\mathcal{U})$  is continuous, then every derivation on A is continuous.

*Proof.* Let  $\delta$  be a derivation on  $\mathcal{A}$ . Define the map  $\tau: \mathcal{U} \to \mathcal{U}$  by  $\tau = \phi o \delta o \psi$ . Then for every  $a \in \mathcal{A}, x \in \mathcal{U}$ ,

$$\tau(ax) = a\tau(x) + \delta(a)x$$
 and  $\tau(xa) = \tau(x)a + x\delta(a)$ .

So the map  $D: T(\mathcal{A}, \mathcal{U}) \to T(\mathcal{A}, \mathcal{U})$  defined by  $D((a, x)) = (\delta(a), \tau(x))$  is a derivation which is continuous by the hypothesis. Thus  $\delta$  is continuous.

**Remark 2.4.** If  $\mathcal{A}$  is a Banach algebra and  $\mathcal{I}$  is a closed ideal on it, then  $\frac{\mathcal{A}}{\mathcal{I}}$  is a Banach  $\mathcal{A}$ -bimodule and so we can consider  $T(\mathcal{A}, \frac{\mathcal{A}}{\mathcal{I}})$ . Suppose that  $\mathcal{A}$  possesses a bounded right (or left) approximate identity and every derivation on  $\mathcal{A}$  and every derivation from  $\mathcal{A}$  to  $\frac{\mathcal{A}}{\mathcal{I}}$  is continuous. Let D be a derivation on  $T(\mathcal{A}, \frac{\mathcal{A}}{\mathcal{I}})$  which has a structure as in Proposition 2.1. Then  $\tau_1: \frac{\mathcal{A}}{\mathcal{I}} \to \mathcal{A}$  is an  $\mathcal{A}$ -bimodule homomorphism. Since  $\mathcal{A}$  has a bounded right(left) approximate identity, then so does  $\frac{\mathcal{A}}{\mathcal{I}}$ . Hence  $\tau_1$  is continuous implying that D is continuous. So in this case any derivation on  $T(\mathcal{A}, \frac{\mathcal{A}}{\mathcal{T}})$  is continuous.

**Remark 2.5.** If  $\mathcal{I}$  is a closed ideal in a Banach algebra  $\mathcal{A}$  and  $\delta: \mathcal{A} \to \mathcal{A}$  is a derivation such that  $\delta(\mathcal{I}) \subseteq \mathcal{I}$ , then the mapping  $\tau: \frac{\mathcal{A}}{\mathcal{I}} \to \frac{\mathcal{A}}{\mathcal{I}}$  defined by  $\tau(a+\mathcal{I}) = \delta(a) + \mathcal{I}$  is well-defined and linear and

$$\tau(a(u+\mathcal{I})) = a\tau(u+\mathcal{I}) + \delta(a)(u+\mathcal{I}) \quad , \quad \tau((u+\mathcal{I})a) = \tau(u+\mathcal{I})a + (u+\mathcal{I})\delta(a).$$

Therefore the mapping D on  $T(\mathcal{A}, \frac{\mathcal{A}}{\mathcal{I}})$  defined by  $D((a, u)) = (\delta(a), \tau(u + \mathcal{I}))$  is a derivation. So if every derivation on  $T(\mathcal{A}, \frac{\mathcal{A}}{\mathcal{I}})$  is continuous, then every derivation  $\delta: \mathcal{A} \to \mathcal{A}$  with  $\delta(\mathcal{I}) \subseteq \mathcal{I}$  is continuous.

In Remarks 2.4 and 2.5, if we let  $\mathcal{I} = (0)$ , then we have the following proposition.

**Proposition 2.6.** Let A be a Banach algebra.

- (i) If A has a right (left) approximate identity and every derivation on A is continuous, then any derivation on T(A, A) is continuous.
- (ii) If every derivation on T(A, A) is continuous, then any derivation on A is continuous.

**Proposition 2.7.** Let  $\mathcal{A}$  be a semisimple Banach algebra which has a bounded approximate identity and  $\mathcal{U}$  be a Banach  $\mathcal{A}$ -bimodule with  $\operatorname{ann}_{\mathcal{A}}\mathcal{U} = (0)$ . Suppose that there exists a surjective left  $\mathcal{A}$ -module homomorphism  $\phi: \mathcal{A} \to \mathcal{U}$  and every derivation from  $\mathcal{A}$  into  $\mathcal{U}$  is continuous, then every derivation on  $T(\mathcal{A}, \mathcal{U})$  is continuous.

*Proof.* Let D be a derivation on  $T(\mathcal{A}, \mathcal{U})$  which has a structure as in Proposition 2.1. Since  $\mathcal{A}$  is semisimple, every derivation from  $\mathcal{A}$  into  $\mathcal{A}$  is continuous. Now by Proposition 2.1 and Remark 2.4, it follows that every derivation on  $T(\mathcal{A}, \mathcal{U})$  is continuous.

Ringrose [21] proved that every derivation from a  $C^*$ -algebra into a Banach bimodule is continuous. According to this result we give the following example satisfying the conditions of the above proposition.

**Example 2.8.** Let  $\mathcal{A}$  be a  $C^*$ -algebra and  $\mathcal{U}$  be a Banach  $\mathcal{A}$ -bimodule with  $ann_{\mathcal{A}}\mathcal{U}=(0)$ . Suppose that there exists a surjective left  $\mathcal{A}$ -module homomorphism  $\phi: \mathcal{A} \to \mathcal{U}$ . Hence  $\mathcal{A}$  is a semisimple Banach algebra with a bounded approximate identity. So by [21] and Proposition 2.7, any derivation on  $T(\mathcal{A},\mathcal{U})$  is continuous.

H. FARHADI

4

An non-trivial element p in an algebra  $\mathcal{A}$  is called an *idempotent* if  $p^2 = p$ .

**Proposition 2.9.** Let A be a prime Banach algebra with a non-trivial idempotent p (i.e.  $p \neq 0$ ) such that Ap is finite dimensional. Then every derivation on A is continuous.

*Proof.* Let  $\mathcal{U} = \mathcal{A}p$ . Then  $\mathcal{U}$  is a closed left ideal in  $\mathcal{A}$ . By the following make  $\mathcal{U}$  into a Banach  $\mathcal{A}$ -bimodule:

$$xa = 0$$
  $(x \in \mathcal{A}p, a \in \mathcal{A}),$ 

and the left multiplication is the usual multiplication of  $\mathcal{A}$ . So we can consider  $T(\mathcal{A},\mathcal{U})$  in this case. Since  $\mathcal{A}$  is prime, it follows that  $ann_{\mathcal{A}}\mathcal{U} = (0)$ . Let  $\delta : \mathcal{A} \to \mathcal{A}$  be a derivation. Define the mapping  $\tau : \mathcal{U} \to \mathcal{U}$  by  $\tau(ap) = \delta(ap)p$   $(a \in \mathcal{A})$ . The mapping  $\tau$  is well-defined and linear. Also

$$\tau(ax) = a\tau(x) + \delta(a)x$$
 and  $\tau(xa) = \tau(x)a + x\delta(a)$   $(a \in \mathcal{A}, x \in \mathcal{U}).$ 

Since  $\mathcal{U}$  is finite dimensional, it follows that  $\tau$  is continuous. By Proposition 2.1, the mapping  $D: T(\mathcal{A},\mathcal{U}) \to T(\mathcal{A},\mathcal{U})$  defined by  $D((a,x)) = (\delta(a),\tau(x))$  is a derivation. Now for D, the conditions of Proposition 2.1 are all satisfied and so D is continuous. Therefore  $\delta$  is continuous.

#### References

- [1] W. G. Bade and P. C. Curtis, *The continuity of derivation of Banach algebras*, J. Functional analysis **16** (1974), 372–378.
- [2] E. Christensen, Derivations of nest algebras, Math. Ann. 229 (1977), 155-161.
- [3] H. G. Dales, Banach Algebras and Automatic Continuity. In: London Math. Soc. Monographs. Oxford Univ. Press, Oxford 2000.
- [4] G. H. Esslamzadeh and H. Ghahramani, Existence, automatic continuity and invariant submodules of generalized derivations on modules, Aequationes mathematicae 84 (2012), 200– 185.
- [5] H. Ghahramani, Jordan derivation on trivial extension, Bulletin of the Iranian Mathematical Society 39 (2013), 645-635.
- [6] H. Ghahramani, Additive maps on some operator algebras behaving like (α, β)-derivations or generalized (α, β)-derivations at zero-product elements, Acta Mathematica Scientia 34B (2014), 1300–1287.
- [7] H. Ghahramani, On derivations and Jordan derivations through zero products, Operators and Matrices 8 (2014), 771–759.
- [8] H. Ghahramani, Characterizing jordan derivations of matrix rings through zero products, Mathematica Slovaca 65 (2015), 1277-1290.
- [9] H. Ghahramani, Jordan derivations on block upper triangular matrix algebras, Operators and Matrices 9 (2015), 181-188.
- [10] H. Ghahramani, On centralizers of Banach algebras, bulletin of the malaysian mathematical sciences society (1)38 (2015), 155-164.
- [11] H. Farhadi, H. Ghahramani, Derivations on semidirect products of Banach algebras, arXiv preprint arXiv:1606.07909.
- [12] H. Ghahramani, Characterizing jordan maps on triangular rings through commutative zero products, Mediterranean Journal of Mathematics 15 (2018), 38-48.
- [13] H. Ghahramani, Linear maps on group algebras determined by the action of the derivations or anti-derivations on a set of orthogonal elements, Results in Mathematics 73 (4) (2018), 133.
- [14] H. Farhadi, H. Ghahramani, Some notes on semidirect products of Banach algebras, Results in Mathematics 74 (3) (2019), 107-117.
- [15] H. Farhadi, E.Ghaderi, H. Ghahramani, Automatic continuity of some linear mappings from certain products of Banach algebras, arXiv preprint arXiv:1902.10081.
- [16] N. Grønbæk, Commutative Banach algebras, module derivations and semigroups, J. London Math. Soc. 40 (2) (1989) 137–157.

- [17] B. E. Johnson and A. M. Sinclair, Continuity of derivations and a problem of Kaplansky, Amer. J. of Math. 90 (1968), 1067–1073.
- [18] R. V. Kadison, Derivations of operator algebras, Ann. Math. 83 (1966), 280-293.
- [19] A. T. Lau, Analysis on a class of Banach algebras with application to harmonic analysis on locally compact groups and semigroups, Fund. Math. 118 (1983), 161–175.
- [20] A. R. Medghalchi and H. Pourmahmood-Aghababa, The first cohomology group of module extension Banach algebras, Rocky mountain J. Math. 41 (2011), 1639–1651.
- [21] J. R. Ringrose, Automatic continuity of derivations of operator algebras, J. London Math. Soc. 5 (1972), 432–438.
- [22] S. Sakai, Derivations of  $W^*$ -algebras, Ann. Math. 83 (1966), 273–279.
- [23] A. M. Sinclair, Continuous derivations on Banach algebras, Proc. Amer. Math. Soc. 20 (1969), 166–170.
- [24] I. M. Singer and J. Wermer, Derivations on commutative normed algebras, Math. Ann. 129 (1955), 260–264.
- [25] A. R. Villena, Automatic continuity in associative and nonassociative context, Irish Math. Soc. Bulletin 46 (2001), 43–76.
- [26] A. R. Villena, Derivations with a hereditary domain, J. London Math. Soc. 57 (1998), 469–477
- [27] A. R. Villena, Derivations with a hereditary domain, II, Studia Math. 130 (1998), 275–291.
- [28] Y. Zhang, Weak amenability of module extensions of Banach algebras, Trans. Amer. Math. Soc. 354 (2002), 4131–4151.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KURDISTAN, P. O. BOX 416, SANANDAJ, IRAN. E-mail address: h.farhadi@sci.uok.ac.ir