

A heavy neutral gauge boson near the Z boson mass pole via third generation fermions at the LHC

Mohammad Abdullah, Mykhailo Dalchenko, Teruki Kamon, Denis Rathjens, and Adrian Thompson

Mitchell Institute for Fundamental Physics and Astronomy, Department of Physics and Astronomy, Texas A&M University, College Station, TX 77843

We explore the physics of a new neutral gauge boson, (Z'), coupling to only third-generation particles with a mass near the electroweak gauge boson mass poles. A Z' boson produced by top quarks and decaying to tau leptons is considered. With a simple search strategy inspired by existing analyses of the standard model gauge boson production in association with top quarks, we show that the Large Hadron Collider has good exclusionary power over the model parameter space of the Z' boson even at the advent of the high-luminosity era. It is shown that the $t\bar{t}Z'$ process allows one to place limits on right-handed top couplings with a Z' boson that preferentially couples to third generation fermions, which are at present very weakly constrained.

I. Introduction

A new neutral massive vector gauge boson, designated as Z' , is considered to be a relatively low-hanging fruit for resonance searches at the CERN Large Hadron Collider (LHC). Due to the numerous allowed combinations of couplings to standard model (SM) fermions, however, there remain production and decay modes that are yet to be thoroughly explored. Of particular interest, in light of the flavor hierarchy in the SM, is a Z' that couples preferentially to the third generation fermions [1–7].

So far, the experimental coverage of third generation couplings in flavor non-universal Z' and their associated resonances has been mainly limited to heavy masses; $m_{Z'} \gtrsim m_t$ [8–12]. Recently, searches for light $b\bar{b}$ [13] resonances and $\tau^-\tau^+$ [14] resonances in association with b -jets have been performed, setting limits for the first time on these signatures. In addition to the inherent difficulties associated with third generation final states, resonances in the low mass regime are more challenging, especially in the vicinity of the electroweak gauge boson mass poles. Since the present limits are modest and ready to be improved over the upcoming run cycle of the High Luminosity (HL)-LHC, the following question is open: is there any new and exclusively third generation physics around the scale of the electroweak gauge boson masses?

To this end, and in contrast with existing studies, we propose a search for a Z' that is produced through top quarks and decays primarily to tau leptons. The negligibly low presence of top quarks in the proton's parton

distribution functions (PDFs) at the LHC energy range leads us to consider Z' production either through gluon splitting followed by top fusion, or top production followed by Z' emission. In both production mechanisms, the resonant ditau is supplemented with a top quark pair which, convolved with reliable top tagging, provides an additional handle in discriminating the signal from the background. Relevant studies can be found in Refs. [10, 15–31].

For the pragmatically oriented reader, this search is interesting in its own right in how it fits into the landscape of resonance searches at colliders. From a model building stand point, however, a model with only such interactions and particle content is not anomaly free and needs to be supplemented with additional physics. There are numerous ways to do so (see for example [32]) and they each have the potential of altering our limits or introducing additional constraints in different ways. Rather than restricting ourselves to one such specific model, we will opt for a minimal setup where the Z' couples only to right handed top quarks and right handed tau leptons, since this coupling scheme is not strongly constrained by existing experiments. We will provide the reader with the mathematical formalism needed to translate our results to a wider class of models.

This paper is organized as follows: in Section II we present our simplified model, in Section III we detail our search strategy, in Section IV we present our findings, and in Section V we conclude.

II. The Model

We consider a toy model that extends the SM Lagrangian with an additional $U(1)'$ gauge symmetry in the 3rd generation of the SM fermions and introduces a new gauge vector boson Z' . The new physics couplings are given as follows:

$$\begin{aligned} \mathcal{L} \supset Z'^\mu & \left[g_\tau \bar{\tau} \gamma_\mu (c_{\tau L} P_L + c_{\tau R} P_R) \tau + g_\nu \bar{\nu}_\tau \gamma_\mu (c_{\nu L} P_L) \nu_\tau \right. \\ & + g_t \bar{t} \gamma_\mu (c_{tL} P_L + c_{tR} P_R) t + g_b \bar{b} \gamma_\mu (c_{bL} P_L + c_{bR} P_R) b \left. \right] \\ & + \text{h.c.}, \end{aligned} \quad (1)$$

where g_τ , g_ν , g_t and g_b are new physics couplings, and c_{fX} ($X = L, R$) are the strengths of left handed and right handed couplings for fermion f . For the case we are interested in we set $c_{tR} = c_{\tau R} = 1$ and all other coefficients c_{fX} to zero. For the purpose of rescaling the signal, $\sigma(pp \rightarrow t\bar{t}Z') \times BR(Z' \rightarrow \tau^+\tau^-)$, it is useful to know the decay width of Z' which, in the massless fermion approximation, is given by:

$$\Gamma(Z' \rightarrow f\bar{f}) = \frac{1}{24\pi} c_f g_f^2 m_{Z'} (c_{fL}^2 + c_{fR}^2), \quad (2)$$

where $m_{Z'}$ is the Z' mass, $c_f = 1$ for leptons, and $c_f = 3$ for quarks. Note that, as written, the expression is valid for general values of c_L and c_R , not just for 0 and 1.

If a bottom quark coupling is included, then the dominant constraint for a light Z' boson comes from the added partial width to the bottomonium decay $\Upsilon(1S) \rightarrow \tau\tau$. The latest central value of

$$R_{\tau\mu} = \frac{\Gamma_{\Upsilon(1S) \rightarrow \tau\tau}}{\Gamma_{\Upsilon(1S) \rightarrow \mu\mu}} = 1.005 \pm 0.013(\text{stat}) \pm 0.022(\text{syst})$$

was reported by the BaBar Collaboration [33]. The SM prediction including one-loop QED diagrams and lepton mass effects is $R_{\tau\mu}^{SM} = 0.9924 \pm O(10^{-5})$ [34]. For the couplings in Equation 1, $R_{\tau\mu}$ is modified to

$$\begin{aligned} R_{\tau\mu} = R_{\tau\mu}^{SM} & \left[1 + (c_{bL} + c_{bR})(c_{\tau L} + c_{\tau R}) \frac{3g_q g_\tau m_\Upsilon^2}{8\pi\alpha(m_{Z'}^2 - m_\Upsilon^2)} \right. \\ & \left. + O\left(\frac{g_q^2 g_\tau^2 m_\Upsilon^4}{m_{Z'}^4}\right) \right] \end{aligned} \quad (3)$$

where α is the fine-structure constant and $m_\Upsilon \approx 9.46$ GeV. If the Z' contributes maximally to the upper- 1σ $R_{\tau\mu}$ value from the BaBar experiment (also including the second-order contribution), Equation 3 translates to the following bound:

$$(c_{bL} + c_{bR})(c_{\tau L} + c_{\tau R}) \left(\frac{g_q g_\tau}{0.01} \right) \left(\frac{40 \text{ GeV}}{m_{Z'}} \right)^2 \lesssim 3.29 \quad (4)$$

The limits placed by the CMS experiment on a pseudoscalar A produced by $b\bar{b}$ fusion and decaying into tau

leptons [14] can be recast as limits on a vector Z' , but at present they are not competitive with the limit from the BaBar experiment for light masses.

Note that if the Z' couples only to the right-handed projections of the third generation quarks, then the top and bottom couplings may be varied independently of each other in certain ultraviolet completions. The $\Upsilon(1S)$ decay bounds are presented here assuming the couplings are equal.

We now proceed to detail our search strategy for the $t\bar{t}Z'(\tau\tau)$ final state. The complementary search for $b\bar{b}Z'(\tau\tau)$ had been thoroughly investigated in the context of a $(B-L)_3$ model [7].

III. Search Strategy

For a quark-produced Z' decaying to tau leptons, QCD and $t\bar{t}$ are the first SM backgrounds of consideration; they can easily dwarf the signal process with their $\gtrsim 1000$ pb cross-section at $\sqrt{s} = 13$ TeV. If hadronically-decaying tau leptons τ_h are within the event selection, the τ_h fake rate for jets of about 0.5–1% [35] has substantial effect for such large cross-sections. However, the presence of light leptons ($\ell = e, \mu$) with $p_T \gtrsim 25$ GeV in the event selection can help mitigate those backgrounds.

The $b\bar{b}Z'$ production channel has been thoroughly investigated by Elahi & Martin [7] (from this point on referred to as EM19) using purely leptonic τ decay, and while it benefits from large statistics, it still suffers from these backgrounds as well as theoretical uncertainties in the proton's bottom-quark PDF. Production of $t\bar{t}Z'$, driven by gluon-splitting into top quarks, trades large statistics for smaller PDF uncertainties and the opportunity to use single- or multi-lepton triggers on additional leptons from the $t \rightarrow Wb$ decays. We take advantage of the higher lepton multiplicity available and consider semileptonic W decays from $t\bar{t} \rightarrow b\bar{b}W^+W^-$ and semileptonic $\tau\tau$ decays, for $\tau_\ell \rightarrow e\nu_\tau\nu_e$ or $\tau_\ell \rightarrow \mu\nu_\tau\nu_\mu$;

$$b\bar{b}(W \rightarrow \ell\nu_\ell)(W \rightarrow \bar{q}q)\tau_h\tau_\ell \quad (5)$$

The three-lepton multiplicity of this selection is sufficient to almost entirely eliminate $t\bar{t}$ as a background, so we neglect it for the analysis. One could also consider $\tau_\ell\tau_\ell$ or $\tau_h\tau_h$, but we find that the former does not yield good fidelity in the $m_{Z'}$ reconstruction and the latter does not sufficiently reduce the SM $t\bar{t}$ background. An important aspect of including a τ_h is that the p_T threshold for τ_h triggers needed to sustain high efficiency selection at CMS and ATLAS ($p_T(\tau_h) \gtrsim 30$ GeV) is relatively higher than the triggers for $\ell = e, \mu$ [36–38]. This is a major factor in the reduction of QCD and $t\bar{t}$ backgrounds, but effectively restricts us to $m_{Z'} \gtrsim 30$ GeV. It is in this way that $b\bar{b}Z'$ and $t\bar{t}Z'$ searches may be complementary probes.

The remaining irreducible backgrounds in this search are $t\bar{t}Z$, $t\bar{t}W^\pm$, $t\bar{t}H$, and $t\bar{t}\gamma^*$. Since we consider light




Figure 1: Representative Feynman diagrams for Z' production via $t\bar{t}$ fusion (left) and $t\bar{t}$ associated production (right).

$m_{Z'}$ near the Z pole, the event content and relevant backgrounds may be very similar to that of a SM vector boson produced in association with top quarks. One can utilize some of the same strategies used in $t\bar{t}W$ or $t\bar{t}Z$ searches [39–42]. From these analyses, we can motivate the choice of a three-lepton selection $\tau_h^\pm \ell^\mp \ell^\mp$ with two same-sign (SS) light leptons that are opposite in sign to the hadronically decaying tau lepton. The SS requirement serves to filter out opposite-sign (OS) signatures from SM Z decays to light leptons. Also, for $m_{Z'} \lesssim 50$ GeV, the $\tau_h\text{-}\tau_\ell$ system can be very proximate in $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2}$ (where η is the pseudorapidity and ϕ is the transverse azimuthal angle of the detector geometry) and merge into a single τ_h object during reconstruction. In these cases, a leptonically decaying associated W , together with the merged τ_h object, could fake a semileptonic $\tau_h\tau_\ell$; the 3-lepton requirement greatly reduces this form of contamination. In order to choose the best light-lepton partner to the hadronic tau, we pick the one with the smallest $\Delta R(\tau_h^\pm, \ell^\mp)$ to disfavor isotropically distributed $\tau_h - W$ pairs.

On the hadronic side of the selection, we require at least three jets in the event and at least one of them b -tagged; since the b -tagging efficiency is $\sim 60\%$ [43] and the correct tag of the signal process is mostly dependent on the lepton selection and overall jet multiplicity, the single b requirement helps increase acceptance without introducing sizable contamination.

Lastly, to ensure real sources of missing transverse energy (\cancel{E}_T) in the event from $W \rightarrow \ell\nu$ decays, we require $\cancel{E}_T > 30$ GeV. The kinematic cuts on b jets, light jets, the τ_h , the light (e/μ) leptons, and the \cancel{E}_T in each event are summarized in Table I.

We use `FeynRules` to generate a model file [44] [45] and `MadGraph5 V2.6.4` [46] to generate signal and background samples. `Pythia 8.2` [47] is used for parton showering and `DELPHES 3.4` [48] to model the detector response. We use a default `DELPHES` card modified to use a jet-finder cone radius of $\Delta R = 0.4$ and muon iso-

	$ \eta \leq$	$p_T \geq$	Multiplicity
$\ell^\pm = e(\mu)$	2.1(2.4)	26(23)	2 SS
τ_h^\mp	2.4	30	1
b -jets	2.4	20	≥ 1
Light jets	2.4	30	$\geq \text{Max}(3-j_b, 0)$
\cancel{E}_T	-	30	-

TABLE I: Basic selection criteria. j_b is the number of selected b -jets.

lation cone of $\Delta R = 0.3$ restricting the summed p_T of non-candidate tracks within the cone to less than 10% of the candidate muon p_T . For simplicity, we simulate signal and background at leading order (LO) and apply the “ k -factor” multipliers that account for the next-to-leading order (NLO) corrections for these processes [49]. These k -factors correspond to an increase of 22% in the $t\bar{t}W^\pm$ cross-section and an overall increase of 26% in the combined $t\bar{t}Z/H/\gamma^*$ cross-section. For the signal process, we approximate the k -factor as that of the $t\bar{t}Z$ process (+23%).

IV. Results

The mass spectra generated after the selection procedure described in Section III is shown in Figure 2. The usual invariant mass $m(\tau_h^\pm, \ell^\mp)$ is found to have the best shape determination and signal-to-background yield in comparison to various transverse-mass constructions such as $m_T(\tau_h^\pm, \ell^\mp)$ or $m_T(\tau_h^\pm, \ell^\mp, \cancel{E}_T)$.




Figure 2: Mass spectra for $t\bar{t}Z'$ in the semileptonic ditau channel before the k -factor NLO enhancement is applied. The signal samples were generated with couplings $g_t = 0.1$ and $g_\tau = 0.5$. Crystalball fits are shown as solid lines and the $t\bar{t}Z/H/\gamma^*$ and $t\bar{t}W$ backgrounds are stacked.

To evaluate the projected sensitivity to this model at benchmark luminosities (300 and 3000 fb⁻¹), we construct a parametrically defined log-likelihood function of g_q and $m_{Z'}$, holding g_τ fixed without loss of generality. To do this, we first perform fits to the signal and background mass spectra and use them to model the asymptotic form of the mass spectrum for each point in model parameter space. *Crystalball* distributions (constituting a power-law tail attached to a gaussian core) are used to fit the signal shape for six choices of $m_{Z'}$ between 40 and 150 GeV as well as the combined background shape; see Figure 2 as an example. Linear fits on the shape parameters as a function of mass allow us to interpolate smoothly between mass points. Additionally, to stabilize the interpolation, the power law order parameter n of the Crystalball was fixed to natural numbers excluding zero, and for these choices the best set of fits was found at $n = 1$. The normalization as a function of $m_{Z'}$ was parameterized using a fit to $\propto m_{Z'}^{-1}$. Changes in the coupling g_t also shift the normalization up and down by a factor of g_t^2 following from the factor of $g_q^2 g_\tau^2$ in the $gg \rightarrow t\bar{t}Z'(\tau^+\tau^-)$ cross section. The Crystalball parameter fits are shown in Figure 3.




Figure 3: Crystalball shape parameters from left to right: μ (mean), σ (width), and α (gaussian-to-power-law transition parameter). The power law order parameter is fixed at $n = 1$. On the far right we fit the integral over the invariant mass spectrum N for couplings $g_q = 0.1$, $g_\tau = 0.5$ at 3000 fb⁻¹.

We define a log-likelihood function (Equation 6) over the asymptotic binned events $N_{s+b}(g_q, m_{Z'})$ tested against the background-only hypothesis $N_b \sim (t\bar{t}W + t\bar{t}Z'/\gamma^*/H)$. We calculate the expected number of events by integrating the signal and background fits on $M(\tau_h^\pm, \ell^\mp)$ between bin edges. We take nuisance parameters σ_i as the Poisson error $\sqrt{N_b^i}$ in each bin.

$$\mathcal{L}(g, m_{Z'}) = - \sum_{\text{bins } i} \left(\frac{(N_{s+b}^i(g, m_{Z'}) - N_b^i)^2}{2\sigma_i^2} + \frac{1}{2} \ln(2\pi\sigma_i) \right) \quad (6)$$

To perform the likelihood scan, we use the **MultiNest** bayesian inference package [50] using flat priors on g_t and $m_{Z'}$ to determine the credible regions in $(m_{Z'}, g_q)$ space. **MultiNest** evaluated the posterior distributions using an evidence tolerance of 0.1 and a sample efficiency of 0.3.

The resulting 95% ($\sim 2\sigma$) exclusion on g_t and $m_{Z'}$ is shown in Figure 4 for 150, 300, and 3000 fb⁻¹ of inte-




Figure 4: Projected limits on $m_{Z'}$ versus g_t with fixed $g_\tau = 0.5$ for the minimal framework where the Z' boson couples only to right-handed top quarks and right handed tau leptons. The dark blue, blue, and light blue regions represent the projected 95% exclusion from the LHC at 150, 300, and 3000 fb⁻¹ of integrated luminosity, respectively. The limit from $R_{\mu\tau}$ in Υ decays (shown in dashed line) is valid only to models in which $g_t = g_b$.




Figure 5: Projected limits on $m_{Z'}$ versus the total cross-section times branching fraction $\sigma(pp \rightarrow Z') \times BR(Z' \rightarrow \tau\tau)$ for the minimal framework where the Z' boson couples only to right-handed top quarks and right handed tau leptons. The dark blue, blue, and light blue regions represent the projected 95% exclusion from the LHC at 150, 300, and 3000 fb⁻¹ of integrated luminosity, respectively.

grated luminosity. This limit contrasts nicely with the EM19 analysis in that we are limited to $m_{Z'} \gtrsim 30$ GeV but can achieve stable performance across masses greater than 80 GeV where EM19's analysis begins to lose performance. We remind the reader that how our results and EM19's fit together depends on the relationships between the various couplings in the full model. We also perform a scan over the cross-section times branching ratio $\sigma(pp \rightarrow Z') \times BR(Z' \rightarrow \tau\tau)$, shown in Figure 5. For this limit calculation we scale the signal shape according to $\left(\frac{\sigma \times BR(Z' \rightarrow \tau\tau)}{\text{pb}} \right) \times \left(\frac{L}{\text{pb}^{-1}} \right) \times \epsilon(m)$ where L is the luminosity and $\epsilon(m)$ is the signal efficiency param-

eterized by the Z' mass. The increased sensitivity with mass in Figure 5 is therefore purely due to the selection efficiency; this serves as a model-independent limit on the $t\bar{t}\tau_h\tau_\ell$ signature.

V. Conclusion

We have investigated the phenomenology of the Z' boson decaying to tau leptons with exclusive third-generation couplings in the mass range $m_{Z'} \in [30, 200]$ GeV produced in association with top quarks in the high-luminosity era of the CERN LHC. Although our focus was placed on the $\tau_h\tau_\ell$ channel in order to suppress QCD and $t\bar{t}$ backgrounds using light lepton triggers, a study of how the $\tau_h\tau_h$ channel is worth pursuing. It should be noted that for right-handed top couplings, the Z' bosons are almost entirely unconstrained by existing experiments, opening the opportunity to make new exclusions of right-chiral models that exceed any existing con-

straints, using luminosities available by the early runs of the HL-LHC. Finally, the search strategy presented here should be complementary to one aiming to discover the Z' boson through $b\bar{b}$ production and decay to tau leptons.

Acknowledgements

The authors are grateful to B. Dutta for the motivating discussions and guidance in the model selection and search strategy. We also acknowledge the Texas A&M University Brazos HPC cluster that contributed to the research reported here. M.A., R.E., T.K. and D.R. are supported in part by the DOE grant de-sc0010813. M.D. and A.T. thank the Mitchell Institute for Fundamental Physics and Astronomy for support. T.K. is also supported in part by the Qatar National Research Fund under project NPRP 9-328-1-066.

-
- [1] B. Holdom, “Hints of new flavor physics at LEP?,” *Phys. Lett. B.* **339** (1994) 114, [arXiv:hep-ph/9407311 \[hep-ph\]](#).
- [2] P. H. Frampton, M. B. Wise, and B. D. Wright, “An Elusive Z-prime coupled to beauty,” *Phys. Rev. D.* **54** (1996) 5820–5823, [arXiv:hep-ph/9604260 \[hep-ph\]](#).
- [3] D. J. Muller and S. Nandi, “Top flavor: A Separate SU(2) for the third family,” *Phys. Lett. B.* **383** (1996) 345–350, [arXiv:hep-ph/9602390 \[hep-ph\]](#).
- [4] A. A. Andrianov, P. Osland, A. A. Pankov, N. V. Romanenko, and J. Sirkka, “On the phenomenology of a Z' coupling only to third family fermions,” *Phys. Rev. D.* **58** (1998) 075001, [arXiv:hep-ph/9804389 \[hep-ph\]](#).
- [5] K. S. Babu, A. Friedland, P. A. N. Machado, and I. Mocioiu, “Flavor Gauge Models Below the Fermi Scale,” *JHEP* **12** (2017) 096, [arXiv:1705.01822 \[hep-ph\]](#).
- [6] A. Kamada, M. Yamada, and T. T. Yanagida, “Self-interacting dark matter with a vector mediator: kinetic mixing with the $U(1)_{(B-L)_3}$ gauge boson,” *JHEP* **03** (2019) 021, [arXiv:1811.02567 \[hep-ph\]](#).
- [7] F. Elahi and A. Martin, “LHC constraints on a $(B-L)_3$ gauge boson,” *Phys. Rev. D.* **100** (2019) 035016, [arXiv:1905.10106 \[hep-ph\]](#).
- [8] ATLAS Collaboration, G. Aad *et al.*, “A search for high-mass resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector,” *JHEP* **07** (2015) 157, [arXiv:1502.07177 \[hep-ex\]](#).
- [9] CMS Collaboration, V. Khachatryan *et al.*, “Search for heavy resonances decaying to tau lepton pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **02** (2017) 048, [arXiv:1611.06594 \[hep-ex\]](#).
- [10] CMS Collaboration, A. M. Sirunyan *et al.*, “Search for $t\bar{t}$ resonances in highly boosted lepton+jets and fully hadronic final states in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **07** (2017) 001, [arXiv:1704.03366 \[hep-ex\]](#).
- [11] CMS Collaboration, J. Haller *et al.*, “Searches for new resonances with couplings to third generation quarks with the CMS detector,” *PoS EPS-HEP2017* (2017) 289.
- [12] ATLAS Collaboration, M. Aaboud *et al.*, “Search for additional heavy neutral higgs and gauge bosons in the ditau final state produced in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13$ tev with the atlas detector,” *JHEP* **01** (2018) 055, [arXiv:1709.07242 \[hep-ex\]](#). [http://dx.doi.org/10.1007/JHEP01\(2018\)055](http://dx.doi.org/10.1007/JHEP01(2018)055).
- [13] CMS Collaboration, A. M. Sirunyan *et al.*, “Search for low-mass resonances decaying into bottom quark-antiquark pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *Phys. Rev. D.* **99** (2019) 012005, [arXiv:1810.11822 \[hep-ex\]](#).
- [14] CMS Collaboration, A. M. Sirunyan *et al.*, “Search for a low-mass $\tau^+\tau^-$ resonance in association with a bottom quark in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **05** (2019) 210, [arXiv:1903.10228 \[hep-ex\]](#).
- [15] C. T. Hill, “Topcolor: Top quark condensation in a gauge extension of the standard model,” *Phys. Lett. B.* **266** (1991) 419.
- [16] C. T. Hill and S. J. Parke, “Top production: Sensitivity to new physics,” *Phys. Rev. D.* **49** (1994) 4454–4462, [arXiv:hep-ph/9312324 \[hep-ph\]](#).
- [17] C. T. Hill, “Topcolor assisted technicolor,” *Phys. Lett. B.* **345** (1995) 483, [arXiv:hep-ph/9411426 \[hep-ph\]](#).
- [18] R. M. Harris and S. Jain, “Cross Sections for Leptophobic Topcolor Z' Decaying to Top-Antitop,” *Eur. Phys. J. C.* **72** (2012) 2072, [arXiv:1112.4928 \[hep-ph\]](#).
- [19] J. L. Rosner, “Prominent decay modes of a leptophobic Z' ,” *Phys. Lett. B.* **387** (1996) 113, [arXiv:hep-ph/9607207 \[hep-ph\]](#).
- [20] K. R. Lynch, E. H. Simmons, M. Narain, and S. Mrenna, “Finding Z' bosons coupled preferentially to

- the third family at LEP and the Tevatron,” *Phys. Rev. D.* **63** (2001) 035006, [arXiv:hep-ph/0007286](https://arxiv.org/abs/hep-ph/0007286) [[hep-ph](#)].
- [21] M. Carena, A. Daleo, B. A. Dobrescu, and T. M. P. Tait, “ Z' gauge bosons at the Tevatron,” *Phys. Rev. D.* **70** (2004) 093009, [arXiv:hep-ph/0408098](https://arxiv.org/abs/hep-ph/0408098) [[hep-ph](#)].
- [22] D. Choudhury, R. M. Godbole, R. K. Singh, and K. Wagh, “Top production at the Tevatron/LHC and nonstandard, strongly interacting spin one particles,” *Phys. Lett. B.* **657** (2007) 69, [arXiv:0705.1499](https://arxiv.org/abs/0705.1499) [[hep-ph](#)].
- [23] CMS Collaboration, V. Khachatryan *et al.*, “Search for resonant $t\bar{t}$ production in proton-proton collisions at $\sqrt{s} = 8$ TeV,” *Phys. Rev. D.* **93** (2016) 012001, [arXiv:1506.03062](https://arxiv.org/abs/1506.03062) [[hep-ex](#)].
- [24] CMS Collaboration, “Search for $t\bar{t}$ resonances at the HL-LHC and HE-LHC with the Phase-2 CMS detector,” Tech. Rep. CMS-PAS-FTR-18-009, CERN, Geneva, 2018. <http://cds.cern.ch/record/2649032>.
- [25] ATLAS Collaboration, M. Aaboud *et al.*, “Search for heavy particles decaying into top-quark pairs using lepton-plus-jets events in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector,” *Eur. Phys. J. C.* **78** (2018) 565, [arXiv:1804.10823](https://arxiv.org/abs/1804.10823) [[hep-ex](#)].
- [26] CMS Collaboration, A. M. Sirunyan *et al.*, “Measurement of the jet mass in highly boosted $t\bar{t}$ events from pp collisions at $\sqrt{s} = 8$ TeV,” *Eur. Phys. J. C.* **77** (2017) 467, [arXiv:1703.06330](https://arxiv.org/abs/1703.06330) [[hep-ex](#)].
- [27] L. Cerrito *et al.*, “Discovering and profiling Z' bosons using asymmetry observables in top quark pair production with the lepton-plus-jets final state at the LHC,” [arXiv:1609.05540](https://arxiv.org/abs/1609.05540) [[hep-ph](#)].
- [28] C. Arina *et al.*, “A comprehensive approach to dark matter studies: exploration of simplified top-philic models,” *JHEP* **11** (2016) 111, [arXiv:1605.09242](https://arxiv.org/abs/1605.09242) [[hep-ph](#)].
- [29] K. Pedersen and Z. Sullivan, “ μ_x boosted-bottom-jet tagging and Z boson searches,” *Phys. Rev. D.* **93** (2016) 014014, [arXiv:1511.05990](https://arxiv.org/abs/1511.05990) [[hep-ph](#)].
- [30] P. J. Fox, I. Low, and Y. Zhang, “Top-philic Z' forces at the LHC,” *JHEP* **03** (2018) 074, [arXiv:1801.03505](https://arxiv.org/abs/1801.03505) [[hep-ph](#)].
- [31] E. Alvarez, R. M. S. Seoane, and A. Juste, “Four-top as probe of light top-philic New Physics,” [arXiv:1910.09581](https://arxiv.org/abs/1910.09581) [[hep-ph](#)].
- [32] S. M. Boucenna, A. Celis, J. Fuentes-Martin, A. Vicente, and J. Virto, “Phenomenology of an $SU(2) \times SU(2) \times U(1)$ model with lepton-flavour non-universality,” *JHEP* **12** (2016) 059, [arXiv:1608.01349](https://arxiv.org/abs/1608.01349) [[hep-ph](#)].
- [33] BABAR Collaboration, del Amo Sanchez *et al.*, “Test of Lepton Universality in $\Upsilon(1S)$ Decays at BABAR,” *Phys. Rev. Lett.* **104** (2010) 191801. <https://link.aps.org/doi/10.1103/PhysRevLett.104.191801>.
- [34] D. Aloni, A. Efrati, Y. Grossman, and Y. Nir, “ Υ and ψ leptonic decays as probes of solutions to the $R_D^{(*)}$ puzzle,” *JHEP* **06** (2017) 019, [arXiv:1702.07356](https://arxiv.org/abs/1702.07356) [[hep-ph](#)].
- [35] CMS Collaboration, A. M. Sirunyan *et al.*, “Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV,” *JINST* **13** (2018) P10005, [arXiv:1809.02816](https://arxiv.org/abs/1809.02816) [[hep-ex](#)].
- [36] CMS Collaboration, “Level-1 E/Gamma trigger performance in Run II,” <https://cds.cern.ch/record/2682202>.
- [37] CMS Collaboration, “Level-1 Muon Trigger Performance,” <https://cds.cern.ch/record/2629852>.
- [38] CMS Collaboration, “Level-1 τ trigger performance in 2017 data,” <https://cds.cern.ch/record/2305547>.
- [39] ATLAS Collaboration, M. Aaboud *et al.*, “Measurement of the $t\bar{t}Z$ and $t\bar{t}W$ cross sections in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector,” *Phys. Rev. D* **99** (2019) 072009. <https://link.aps.org/doi/10.1103/PhysRevD.99.072009>.
- [40] CMS Collaboration, A. M. Sirunyan *et al.*, “Evidence for associated production of a Higgs boson with a top quark pair in final states with electrons, muons, and hadronically decaying τ leptons at $\sqrt{s} = 13$ TeV,” *JHEP* **08** (2018) 066, [arXiv:1803.05485](https://arxiv.org/abs/1803.05485) [[hep-ex](#)].
- [41] CMS Collaboration, A. Sirunyan *et al.*, “Measurement of the cross section for top quark pair production in association with a W or Z boson in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **08** (2018) 11, [arXiv:1711.02547](https://arxiv.org/abs/1711.02547) [[hep-ex](#)]. [https://doi.org/10.1007/JHEP08\(2018\)011](https://doi.org/10.1007/JHEP08(2018)011).
- [42] CMS Collaboration, A. M. Sirunyan *et al.*, “Measurement of top quark pair production in association with a Z boson in proton-proton collisions at $\sqrt{s} = 13$ TeV,” [arXiv:1907.11270](https://arxiv.org/abs/1907.11270) [[hep-ex](#)].
- [43] CMS Collaboration, A. M. Sirunyan *et al.*, “Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV,” *JINST* **13** (2018) P05011, [arXiv:1712.07158](https://arxiv.org/abs/1712.07158) [[physics.ins-det](#)].
- [44] N. D. Christensen and C. Duhr, “FeynRules - Feynman rules made easy,” *Comput. Phys. Commun.* **180** (2009) 1614–1641, [arXiv:0806.4194](https://arxiv.org/abs/0806.4194) [[hep-ph](#)].
- [45] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr, and B. Fuks, “FeynRules 2.0 - A complete toolbox for tree-level phenomenology,” *Comput. Phys. Commun.* **185** (2014) 2250, [arXiv:1310.1921](https://arxiv.org/abs/1310.1921) [[hep-ph](#)].
- [46] J. Alwall *et al.*, “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations,” *JHEP* **07** (2014) 079, [arXiv:1405.0301](https://arxiv.org/abs/1405.0301) [[hep-ph](#)].
- [47] T. Sjostrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, “An Introduction to PYTHIA 8.2,” *Comput. Phys. Commun.* **191** (2015) 159, [arXiv:1410.3012](https://arxiv.org/abs/1410.3012) [[hep-ph](#)].
- [48] DELPHES 3 Collaboration, J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaitre, A. Mertens, and M. Selvaggi, “DELPHES 3, A modular framework for fast simulation of a generic collider experiment,” *JHEP* **02** (2014) 057, [arXiv:1307.6346](https://arxiv.org/abs/1307.6346) [[hep-ex](#)].
- [49] F. Maltoni, D. Pagani, and I. Tsinikos, “Associated production of a top-quark pair with vector bosons at NLO in QCD: impact on $t\bar{t}H$ searches at the LHC,” *JHEP* **02** (2016) 113, [arXiv:1507.05640](https://arxiv.org/abs/1507.05640) [[hep-ph](#)].
- [50] F. Feroz, M. P. Hobson, and M. Bridges, “MultiNest: an efficient and robust Bayesian inference tool for cosmology and particle physics,” *Mon. Not. Roy. Astron. Soc.* **398** (2009) 1601, [arXiv:0809.3437](https://arxiv.org/abs/0809.3437) [[astro-ph](#)].