A new characterization of simple K_3 -groups using same-order type

Igor Lima and Josyane Pereira

ABSTRACT. Let G be a group, define an equivalence relation \sim as below:

$$\forall g, h \in G, g \sim h \iff |g| = |h|$$

the set of sizes of equivalence classes with respect to this relation is called the same-order type of G and denoted by $\alpha(G)$. And G is said a α_n -group if $|\alpha(G)| = n$. Let $\pi(G)$ be the set of prime divisors of the order of G. A simple group of G is called a simple K_n -group if $|\pi(G)| = n$. We give a new characterization of simple K_3 -groups using same-order type. Indeed we prove that a nonabelian simple group G has same-order type $\{r, m, n, k, l\}$ if and only if $G \cong PSL(2,q)$, with q=7,8 or 9. This result generalizes the main results in [4], [6] and [8]. Moreover based on the main result in [8] we have the natural question: Let S be a nonabelian simple α_n -group and G a α_n -group such that |S| = |G|. Then $S \cong G$. In this paper with a counterexample we give a negative answer to this question.

1. Introduction and Preliminaries

In this paper all the groups we consider are finite.

Let G a group and $\pi_e(G)$ be the set of element orders of G. Let $t \in \pi_e(G)$ and s_t be the number of elements of order t in G. Let $nse(G) = \{s_t | t \in \pi_e(G)\}$ the set of sizes of elements with the same order in G. Some authors have studied the influence of nse(G) on the structure of G (see [1], [5], [8] and [9]). For instance G. Shen in [6] proved that G is a continuation in [4] was proved that if G is a group such that G is a continuation in [4] was proved that if G is a group such that G is the only group such that G is a group of that G is the only group such that G is a group such that G is the only group such that G is a group of that G is the only group such that G is a group of that G is the only group such that G is a group of that G is a group of that G is the only group such that G is a group of that G is an onabelian simple group G has same-order type G is an only if $G \cong G$ (see Th. 1.1 [8]).

Let G be a group, in [8] was defined an equivalence relation \sim as below:

$$\forall g, h \in G, g \sim h \iff |g| = |h|$$

the set of sizes of equivalence classes with respect to this relation is called the same-order type of G and denoted by $\alpha(G)$. And G is said a α_n -group if $|\alpha(G)| = n$. Note that $\alpha(G)$ is equal to the set of sizes of elements with the same order in G, hence $|nse(G)| = |\alpha(G)|$.

We give a new characterization of PSL(2,7), PSL(2,8) and PSL(2,9) using same-order type.

1

²⁰¹⁰ Mathematics Subject Classification. 20D60, 20D66.

Key words and phrases. Element order, Same-order type, Characterization, Simple group.

The first author was supported by DPI/UnB, Brazil.

The second author was supported by CNPq.

THEOREM 1.1. Let G be a simple K_3 -group with same-order type $\{r, m, n, k, l\}$. Then $G \cong PSL(2,7), PSL(2,8)$ or PSL(2,9).

This result generalizes the main results in [4], [6] and [8]. Combination the main results in [4] and [6] with Theorem 1.1 we have the following result

COROLLARY 1.2. A simple K_3 -group G has same-order type $\{r, m, n, k, l\}$ if and only if $G \cong PSL(2,7), PSL(2,8)$ or PSL(2,9).

We see easily that the only α_1 -groups are 1 and a cyclic group of order 2. In [6] R. Shen characterized α_2 -groups as nilpotent groups and α_3 -groups as solvable groups. Moreover Taghvasani-Zarrin (see Th. 1.1 in [8]) showed that the only nonabelian simple α_4 -group is the A_5 . As noted in [4] and [8] finite groups G cannot be determined by nse(G). Indeed in 1987 Thompson gave a first example as follows: Let $G_1 = (C_2 \times C_2 \times C_2 \times C_2) \times A_7$ and $G_2 = PSL(3,4) \times C_2$ be the maximal subgroups of Mathieu group M_{23} . Then $nse(G_1) = nse(G_2)$, but $G_1 \ncong G_2$.

Motived by the main result in [8] about a new characterization of A_5 using same-order type, we have the natural question

QUESTION 1.3. Let S be a nonabelian simple α_n -group and G a α_n -group such that |S| = |G|. Then $S \cong G$.

We give a negative answer to this question in the last section.

2. Proof of Theorem 1.1

We need of one preliminary result to prove the main Theorem. The following result is a property very interesting of simple groups (see Lemma 2.7 in [8]).

LEMMA 2.1. Let G be a nonabelian simple group. Then there exist two odd prime divisors p and q of the order of G such that $s_p \neq s_q$.

In fact if *G* is a nonabelian simple group then there exist two odd prime divisors *p* and *q* of the order of *G* such that $\{1, s_2, s_p, s_q\} \subseteq \alpha(G)$ (see Corollary 2.8 in [8]).

We are now ready to conclude the proof of main Theorem.

Proof of Theorem 1.1: As G is a nonabelian simple group, it follows that $s_2 > 1$, w.l.g. r = 1 and $s_2 = m$. From Lemma 2.1 there exist odd prime divisors p and q of the order of G such that $n = s_p \neq s_q = k$, hence $\pi(G) = \{2, p, q\}$ because G is a simple K_3 -group. Therefore $\{1, s_2, s_p, s_q\} \subseteq \alpha(G) = \{r, m, n, k, l\}$. So there exist a divisor $t \notin \pi(G)$ of order of G such that $s_t = l$. It's well known that the only nonabelian simple groups of order divisible by exactly three primes are the following eight groups: PSL(2,q), where $q \in \{5,7,8,9,17\}$, PSL(3,3), PSU(3,3) and PSU(4,2), see Th. 1 and Th. 2 in [3]. Now we arguing as in the proof of Th. 1.1 in [8] and a GAP check yields that $|\alpha(PSL(2,7))| = 5$, $|\alpha(PSL(2,8))| = 5$, $|\alpha(PSL(2,9))| = 5$ and all others groups are α_n -groups with $n \ge 6$ (except A_5 since $|\alpha(A_5)| = 4$). The result is follows.

3. A counterexample to a Question 1.3

Now we give a counterexample to the Question 1.3. Firstly we observed that by the main Theorem in [6], we have that $\alpha(PSL(2,7))$ is uniquely determined and we have that $\alpha(PSL(2,7)) = \{1,21,56,42,48\}$ hence PSL(2,7) is a α_5 -group. Let $G = Q_8 \times (C_7 \rtimes C_3)$, where Q_8 is the quaternion group of order 8. As |G| = 168 and G is a soluble group then is sufficient to prove that $|\alpha(G)| = 5$. Indeed the only 2-Sylow subgroup Q_8 is a normal

subgroup of G and using Sylow's Theorem it follows that $s_2 = 8$. Note that a 7-Sylow subgroup of G is isomorphic to C_7 and is a normal subgroup of $Q_8 \cdot C_7$ and $C_7 \rtimes C_3$, hence the normalizer N of C_7 has same order of G. Again from Sylow's Theorem we have that C_7 is a normal subgroup of G and $s_7 = 56$. As the number of 3-Sylow subgroup of G is 7, then $s_3 = 14$. The number of elements of G of order 2, 4 are respectively 1 and 6, hence $s_2 = 1$, $s_4 = 6$ and consequently $s_6 = 14$, $s_{12} = 84$, $s_{14} = 6$ and $s_{28} = 36$ (because of direct product in the structure of G). Therefore $G(G) = \{1, 1, 14, 6, 14, 6, 84, 6, 36\}$ and G is a G-group. Clearly |PSL(2,7)| = 168 = |G| but $PSL(2,7) \ncong G$.

We can obtain others groups G with the computational group theory system GAP [2]: $G = C_7 \times (Q_8 \rtimes C_3)$ or $G = C_2 \times ((C_{14} \times C_2) \rtimes C_3)$. These groups are also counterexamples to the Question 1.3.

Acknowledgments: We thank L. J. Taghvasani for pointing out a mistake of ours and talking to us about Question 1.3 in the olderst version of [8].

References

- [1] C. S. Anabanti. A counterexample to Zarrin's Conjecture on sizes of finite nonabelian simple groups in relation to involution sizes. *Archiv der Mathematik*, **112** (3) (2019), 225–226.
- [2] The GAP Group (2019). GAP Groups, Algorithms, and Programming, Version 4.10.2. https://www.gapsystem.org
- [3] M. Herzog. On finite simple groups of order divisible by three primes only. J. Algebra, 10 (1968), 383–388.
- [4] M. Khatami, B. Khosravi, Z. Akhlaghi. A new characterization for some linear groups. *Monatsh. Math.*, 163 (2011), 39-50.
- [5] I. A. Malinowska. Finite groups with few normalizers or involutions. Archiv der Mathematik, 112 (3) (2019), 459–465.
- [6] R. Shen. On groups with given same-order type. Comm. Algebra, 40 (2012), 2140-2150.
- [7] R. Shen, C. Shao, Q. Jiang, W. Mazurov. A new characterization A5, Monatsh. Math., 160 (2010), 337-341.
- [8] L. J. Taghvasani and M. Zarrin. A characterization of A5 by its same-order type, Monatsh. Math., 182 (2017), 731-736.
- [9] M. Zarrin. A counterexample to Herzog's Conjecture on the number of involutions. *Arch. Math.*, 111 (2018), 349–351.

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DE BRASÍLIA, BRASILIA-DF, 70910-900 BRAZIL *Email address*: (Lima) igor.matematico@gmail.com, (Pereira) josyanedsp@gmail.com