

**UNIQUE SOLVABILITY OF ELLIPTIC PROBLEMS ASSOCIATED
WITH TWO-PHASE INCOMPRESSIBLE FLOWS IN
UNBOUNDED DOMAINS**

HIROKAZU SAITO AND XIN ZHANG

ABSTRACT. This paper shows the unique solvability of elliptic problems associated with two-phase incompressible flows, which are governed by the two-phase Navier-Stokes equations, in unbounded domains such as the whole space separated by a compact interface and the whole space separated by a non-compact interface. As a by-product of the unique solvability of elliptic problems, we obtain the Helmholtz-Weyl decomposition for two-phase incompressible flows.

1. INTRODUCTION AND MAIN RESULTS

1.1. Introduction. Let Ω_+ be a bounded domain in the N -dimensional Euclidean space \mathbf{R}^N , $N \geq 2$, with boundary Σ , and let $\Omega_- = \mathbf{R}^N \setminus (\Omega_+ \cup \Sigma)$. Let us define $\rho = \rho_+ \mathbb{1}_{\Omega_+} + \rho_- \mathbb{1}_{\Omega_-}$ for positive constants ρ_{\pm} , where $\mathbb{1}_A$ is the indicator function of $A \subset \mathbf{R}^N$. We set for an open set G of \mathbf{R}^N and for $q \in (1, \infty)$

$$\widehat{H}_q^1(G) = \{f \in L_{q,\text{loc}}(\overline{G}) : \nabla f \in L_q(G)^N\},$$

and define a solenoidal space $J_q(\mathbf{R}^N \setminus \Sigma)$ and a space $G_q(\mathbf{R}^N \setminus \Sigma)$ as follows:

$$J_q(\mathbf{R}^N \setminus \Sigma) = \{\mathbf{u} \in L_q(\mathbf{R}^N \setminus \Sigma)^N : (\mathbf{u}, \nabla \varphi)_{\mathbf{R}^N \setminus \Sigma} = 0 \text{ for any } \varphi \in \widehat{H}_{q'}^1(\mathbf{R}^N)\},$$

$$G_q(\mathbf{R}^N \setminus \Sigma) = \{\mathbf{v} \in L_q(\mathbf{R}^N \setminus \Sigma)^N : \mathbf{v} = \rho^{-1} \nabla \psi \text{ for some } \psi \in \widehat{H}_q^1(\mathbf{R}^N)\},$$

where $q' = q/(q-1)$ and

$$(\mathbf{u}, \nabla \varphi)_{\mathbf{R}^N \setminus \Sigma} = \int_{\mathbf{R}^N \setminus \Sigma} \mathbf{u}(x) \cdot \nabla \varphi(x) dx.$$

Here the central dot denotes the scalar product of \mathbf{R}^N .

Let $\mathbf{f} \in L_q(\mathbf{R}^N \setminus \Sigma)^N$. One of the purposes of this paper is to show the unique solvability of the following weak elliptic problem: Find $u \in \widehat{H}_q^1(\mathbf{R}^N)$ such that

$$(1.1) \quad (\rho^{-1} \nabla u, \nabla \varphi)_{\mathbf{R}^N \setminus \Sigma} = (\mathbf{f}, \nabla \varphi)_{\mathbf{R}^N \setminus \Sigma} \quad \text{for any } \varphi \in \widehat{H}_{q'}^1(\mathbf{R}^N).$$

This weak elliptic problem arises from the study of two-phase incompressible flows governed by the two-phase Navier-Stokes equations. The momentum equation of the two-phase Navier-Stokes equations is linearized as

$$\rho \partial_t \mathbf{u} - \mu \Delta \mathbf{u} + \nabla \mathbf{p} = \mathbf{g} \quad \text{in } \mathbf{R}^N \setminus \Sigma,$$

2010 *Mathematics Subject Classification.* Primary: 35Q30; Secondary: 76D07.

Key words and phrases. Elliptic problem; Unbounded domain; Two-phase incompressible flow; Helmholtz-Weyl decomposition.

The work of the first author was supported by JSPS KAKENHI Grant Number JP17K14224, and the second author was supported by the Top Global University Project.

where \mathbf{g} is a given function and $\mu = \mu_+ \mathbb{1}_{\Omega_+} + \mu_- \mathbb{1}_{\Omega_-}$ for positive constants μ_{\pm} describing the viscosity coefficients, and then the unique solvability of (1.1) enables us to eliminate the pressure \mathbf{p} from the linearized equation. This elimination of pressure plays an important role in applications such as the generation of analytic C_0 -semigroups, the maximal regularity, and the local and global solvability of the two-phase Navier-Stokes equations (cf. [17], [13], and [20]). Another very important application of the unique solvability of (1.1) is a two-phase version of the Helmholtz-Weyl decomposition as follows:

$$(1.2) \quad L_q(\mathbf{R}^N \setminus \Sigma)^N = J_q(\mathbf{R}^N \setminus \Sigma) \oplus G_q(\mathbf{R}^N \setminus \Sigma),$$

where \oplus denotes the direct sum. Note that this decomposition is equivalent to the unique solvability of (1.1).

Prüss and Simonett [17, Proposition 8.6.2] proved the unique solvability of a weak elliptic problem associated with two-phase incompressible flows in the case where Ω_{\pm} are both bounded domains, while the case of unbounded domains is not very well known to the best of our knowledge. This motivates us to study the unique solvability of (1.1). Furthermore, as examples of unbounded domains with non-compact interface Σ , we also treat the whole space with a flat interface and the whole space with a bent interface in the present paper. The former is in Subsection 3.4 below, while the latter is in Subsection 4.3 below.

At this point, we introduce a short history of one-phase case for the unique solvability of weak elliptic problems and the Helmholtz-Weyl decomposition.

We first introduce the classical weak Neumann problem: Find $u \in \widehat{H}_q^1(D)$ such that

$$(1.3) \quad (\nabla u, \nabla \varphi)_D = (\mathbf{f}, \nabla \varphi)_D \quad \text{for any } \varphi \in \widehat{H}_q^1(D),$$

where $\mathbf{f} \in L_q(D)^N$ and D is a domain in \mathbf{R}^N . It is well known that the unique solvability of (1.3) is equivalent to the following Helmholtz-Weyl decomposition:

$$(1.4) \quad L_q(D)^N = L_{q,\sigma}(D) \oplus G_q(D),$$

where $L_{q,\sigma}(D)$ and $G_q(D)$ are given by

$$\begin{aligned} L_{q,\sigma}(D) &= \overline{C_{0,\sigma}^{\infty}(D)}^{\|\cdot\|_{L_q(D)}}, \quad C_{0,\sigma}^{\infty}(D) = \{\mathbf{u} \in C_0^{\infty}(D)^N : \operatorname{div} \mathbf{u} = 0 \text{ in } D\}, \\ G_q(D) &= \{\mathbf{v} \in L_q(D)^N : \mathbf{v} = \nabla \psi \text{ for some } \psi \in \widehat{H}_q^1(D)\}. \end{aligned}$$

The investigation of (1.4) (or (1.3)) can be traced back to Weyl [31]. Although (1.4) holds for any domain D in \mathbf{R}^N when $q = 2$ (cf. e.g. [28]), the general L_q -framework is more involved. According to [29, 10, 15, 26, 8, 16, 9, 1, 12], we can conclude that (1.4) is valid for any $q \in (1, \infty)$ whenever D is \mathbf{R}^N itself, the half space, a bounded or an exterior domain in \mathbf{R}^N with smooth boundary, a perturbed half space, a flat layer, an aperture domain, or a bounded convex domain. One also knows in [5] that (1.4) holds only when $3/2 - \varepsilon < q < 3 + \varepsilon$ for some $\varepsilon = \varepsilon(D) > 0$, assuming D is a bounded Lipschitz domain in \mathbf{R}^N . Note that (1.4) may fail for some unbounded domain and $q \in (1, \infty)$, which is pointed out in [14]. One however has a chance to obtain (1.4) for any $q \in (1, \infty)$ and for general unbounded domains, called uniform C^1 domains, by introducing mixed L_q -spaces due to [6, 7].

The classical (1.4) is widely used for one-phase problems with non-slip boundary condition. On the other hand, to handle one-phase incompressible flows with a free

surface, we make use of the weak Dirichlet problem as follows: Let Γ be a connected component of the boundary of D and

$$\widehat{H}_{q,\Gamma}^1(D) = \{u \in \widehat{H}_q^1(D) : u = 0 \text{ on } \Gamma\} \quad (1 < q < \infty).$$

One then says that the weak Dirichlet problem is uniquely solvable in $\widehat{H}_{q,\Gamma}^1(D)$ if and only if for any $\mathbf{f} \in L_q(D)^N$ there is a unique solution $u \in \widehat{H}_{q,\Gamma}^1(D)$ to

$$(\nabla u, \nabla \varphi)_D = (\mathbf{f}, \nabla \varphi)_D \quad \text{for any } \varphi \in \widehat{H}_{q,\Gamma}^1(D)$$

and there holds the estimate: $\|\nabla u\|_{L_q(D)} \leq C\|\mathbf{f}\|_{L_q(D)}$ for some positive constant C independent of u , \mathbf{f} , and φ . In [27, 23, 17], it is proved that the weak Dirichlet problem is uniquely solvable in $\widehat{H}_{q,\Gamma}^1(D)$ when D is \mathbf{R}^N , the half space, a bounded or an exterior domain with smooth boundary (cf. also [2, 18]).

Finally, we introduce the strong elliptic problem associated with (1.1). For an open set G of \mathbf{R}^N and $q \in (1, \infty)$, one sets

$$\widehat{H}_q^2(G) = \{f \in L_{1,\text{loc}}(G) : \nabla f \in \widehat{H}_q^1(G)^N\}.$$

The strong elliptic problem is then stated as follows: Find $v_\pm \in \widehat{H}_q^1(\Omega_\pm) \cap \widehat{H}_q^2(\Omega_\pm)$ such that

$$(1.5) \quad \begin{cases} \Delta v_\pm = \text{div } \mathbf{f}_\pm & \text{in } \Omega_\pm, \\ \rho_+ v_+ = \rho_- v_- & \text{on } \Sigma, \\ \mathbf{n} \cdot \nabla(v_+ - v_-) = \mathbf{n} \cdot (\mathbf{f}_+ - \mathbf{f}_-) & \text{on } \Sigma, \end{cases}$$

where \mathbf{n} is a unit normal vector on Σ pointing from Ω_+ into Ω_- . Throughout this paper, \mathbf{n} is seen as an N -vector of function defined on \mathbf{R}^N (cf. [21, Corollary A.3] and Assumption 1.1 below). In this paper, we first prove the unique solvability of (1.5), and then we prove the unique solvability of (1.1) by using the result of the strong elliptic problem. This approach is also applied to the problems with non-compact interfaces in Sections 3 and 4 below.

Notation. Let G be an open set in \mathbf{R}^N , and let $u = u(x)$ and $\mathbf{v} = \mathbf{v}(x) = (v_1(x), \dots, v_N(x))^{\text{T}}$ be respectively a scalar-valued function on G and a vector-valued function on G , where $x = (x_1, \dots, x_N)$. Then, for $\partial_j = \partial/\partial x_j$,

$$\nabla u = (\partial_1 u, \dots, \partial_N u)^{\text{T}}, \quad \nabla \mathbf{v} = \{\partial_j v_k : j, k = 1, \dots, N\},$$

and also $\text{div } \mathbf{v} = \sum_{j=1}^N \partial_j v_j$. Furthermore, for $\mathbf{u} = \mathbf{u}(x) = (u_1(x), \dots, u_N(x))^{\text{T}}$ and $v = v(x)$ defined on G ,

$$(u, v)_G = \int_G u(x)v(x) dx, \quad (\mathbf{u}, \mathbf{v})_G = \int_G \mathbf{u}(x) \cdot \mathbf{v}(x) dx = \sum_{j=1}^N \int_G u_j(x)v_j(x) dx.$$

Let X be a Banach space. Then X^m , $m \geq 2$, denotes the m -product space of X , while the norm of X^m is usually denoted by $\|\cdot\|_X$ instead of $\|\cdot\|_{X^m}$ for the sake of simplicity. For another Banach space Y , $\mathcal{L}(X, Y)$ stands for the Banach space of all bounded linear operators from X to Y . In addition, $\mathcal{L}(X) = \mathcal{L}(X, X)$.

Let $p \geq 1$ or $p = \infty$, and let $q \in (1, \infty)$. The Lebesgue spaces on G are denoted by $L_p(G)$ with norm $\|\cdot\|_{L_p(G)}$, while the Sobolev spaces on G are denoted by

¹ \mathbf{M}^{T} denotes the transpose of \mathbf{M} .

$H_p^n(G)$, $n \in \mathbf{N}$, with norm $\|\cdot\|_{H_p^n(G)}$. Set $H_p^0(G) = L_p(G)$ and $\mathbf{N}_0 = \mathbf{N} \cup \{0\}$. For any multi-index $\alpha = (\alpha_1, \dots, \alpha_N) \in \mathbf{N}_0^N$,

$$\partial^\alpha u = \partial_x^\alpha u = \frac{\partial^{|\alpha|} u(x)}{\partial x_1^{\alpha_1} \dots \partial x_N^{\alpha_N}} \quad \text{with } |\alpha| = \alpha_1 + \dots + \alpha_N.$$

Let $s \in (0, \infty) \setminus \mathbf{N}$, and then $[s]$ stands for the largest integer less than s , while $W_q^s(G) = \{f \in L_q(G) : \|f\|_{W_q^s(G)} < \infty\}$ with

$$\|f\|_{W_q^s(G)} = \|f\|_{H_q^{[s]}(G)} + \sum_{|\alpha|=[s]} \left(\int_G \int_G \frac{|\partial^\alpha f(x) - \partial^\alpha f(y)|^q}{|x-y|^{N+(s-[s])q}} dx dy \right)^{1/q}.$$

In addition, $C_0^\infty(G)$ stands for the set of all functions in $C^\infty(G)$ whose supports are compact subsets of G , while for a domain D of \mathbf{R}^N

$$E_q(D) = \{\mathbf{f} \in L_q(D)^N : \operatorname{div} \mathbf{f} \in L_q(D)\}$$

endowed with $\|\mathbf{f}\|_{E_q(D)} = \|\mathbf{f}\|_{L_q(D)} + \|\operatorname{div} \mathbf{f}\|_{L_q(D)}$.

1.2. Main results. We first introduce assumptions for Ω_\pm .

Assumption 1.1. (a) r is a real number satisfying $r > N$.

(b) Ω_+ is a bounded domain of \mathbf{R}^N with boundary Σ of class $W_r^{2-1/r}$.

(c) $\Omega_- = \mathbf{R}^N \setminus (\Omega_+ \cup \Sigma)$.

Now we state our main result for the strong elliptic problem (1.5).

Theorem 1.2. *Suppose that Assumption 1.1 holds and ρ_\pm are positive constants. Let $q \in (1, \infty)$ and $q' = q/(q-1)$ with $\max(q, q') \leq r$.*

(1) **Existence.** *Let $\mathbf{f}_\pm \in E_q(\Omega_\pm)$ with $\mathbf{n} \cdot \mathbf{f}_\pm \in H_q^1(\Omega_\pm)$. Then the strong elliptic problem (1.5) admits solutions $v_\pm \in \widehat{H}_q^1(\Omega_\pm) \cap \widehat{H}_q^2(\Omega_\pm)$ satisfying*

$$(1.6) \quad \|\nabla^2 v_\pm\|_{L_q(\Omega_\pm)} \leq C \sum_{\mathbf{s} \in \{+, -\}} \left(\|\mathbf{f}_\mathbf{s}\|_{E_q(\Omega_\mathbf{s})} + \|\mathbf{n} \cdot \mathbf{f}_\mathbf{s}\|_{H_q^1(\Omega_\mathbf{s})} \right),$$

$$(1.7) \quad \|\nabla v_\pm\|_{L_q(\Omega_\pm)} \leq C \sum_{\mathbf{s} \in \{+, -\}} \|\mathbf{f}_\mathbf{s}\|_{L_q(\Omega_\mathbf{s})},$$

with some positive constant $C = C(N, q, r, \rho_+, \rho_-)$. Additionally, if $\mathbf{n} \cdot (\mathbf{f}_+ - \mathbf{f}_-) = 0$ on Σ , then v_\pm satisfy

$$(1.8) \quad \|\nabla^2 v_\pm\|_{L_q(\Omega_\pm)} \leq C \sum_{\mathbf{s} \in \{+, -\}} \|\mathbf{f}_\mathbf{s}\|_{E_q(\Omega_\mathbf{s})}$$

for some positive constant $C = C(N, q, r, \rho_+, \rho_-)$.

(2) **Uniqueness.** *If $v_\pm \in \widehat{H}_q^1(\Omega_\pm) \cap \widehat{H}_q^2(\Omega_\pm)$ satisfies*

$$\Delta v_\pm = 0 \text{ in } \Omega_\pm, \quad \rho_+ v_+ = \rho_- v_- \text{ and } \mathbf{n} \cdot \nabla(v_+ - v_-) = 0 \text{ on } \Sigma,$$

then $v_\pm = \rho_\pm^{-1} c$ for some constant c .

For the weak elliptic problem (1.1), our main result reads as

Theorem 1.3. *Suppose that Assumption 1.1 holds and $\rho = \rho_+ \mathbf{1}_{\Omega_+} + \rho_- \mathbf{1}_{\Omega_-}$ for positive constants ρ_\pm . Let $q \in (1, \infty)$ and $q' = q/(q-1)$ with $\max(q, q') \leq r$.*

- (1) **Existence.** Let $\mathbf{f} \in L_q(\mathbf{R}^N \setminus \Sigma)^N$. Then the weak elliptic problem (1.1) admits a solution $u \in \widehat{H}_q^1(\mathbf{R}^N)$ satisfying the estimate: $\|\nabla u\|_{L_q(\mathbf{R}^N)} \leq C\|\mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \Sigma)}$ for some positive constant $C = C(N, q, r, \rho_+, \rho_-)$.
- (2) **Uniqueness.** If $u \in \widehat{H}_q^1(\mathbf{R}^N)$ satisfies

$$(\rho^{-1}\nabla u, \nabla \varphi)_{\mathbf{R}^N \setminus \Sigma} = 0 \quad \text{for any } \varphi \in \widehat{H}_q^1(\mathbf{R}^N),$$

then $u = c$ for some constant c .

Furthermore, we have by Theorem 1.3 the two-phase version of the Helmholtz-Weyl decomposition as follows:

Theorem 1.4. *Suppose that the same assumption as in Theorem 1.3 holds. Then the decomposition (1.2) holds.*

This paper is organized as follows: The next section introduces some function spaces and lemmas, which are used in Section 3. Section 3 treats strong elliptic problems with and without resolvent parameter λ and a weak elliptic problem in the whole space with a flat interface. Section 4 treats strong and weak elliptic problems similar to Section 3 in the whole space with a bent interface, and proves the unique solvability of the problems by using results obtained in Section 3. In Section 5, we first introduce the unique solvability of a strong elliptic problem with resolvent parameter λ in a bounded domain, which is proved by the standard localization technique together with a result given in Section 4. Next, we prove the unique solvability of the strong elliptic problem without λ by using the result with λ and the Riesz-Schauder theory. Section 6 proves our main results as stated above, i.e. Theorems 1.2 and 1.3, by the main result of Section 5 with a cut-off technique.

2. PRELIMINARIES

Let us define

$$\begin{aligned} F_q(\mathbf{R}_\pm^N) &= \{\mathbf{f}_\pm = (f_{\pm 1}, \dots, f_{\pm N})^\top \in E_q(\mathbf{R}_\pm^N) : f_{\pm N} \in H_q^1(\mathbf{R}_\pm^N)\}, \\ \|\mathbf{f}_\pm\|_{F_q(\mathbf{R}_\pm^N)} &= \|\mathbf{f}_\pm\|_{E_q(\mathbf{R}_\pm^N)} + \|f_{\pm N}\|_{H_q^1(\mathbf{R}_\pm^N)}, \end{aligned}$$

where \mathbf{R}_\pm^N are half spaces given by

$$\mathbf{R}_\pm^N = \{x = (x', x_N) : x' = (x_1, \dots, x_{N-1}) \in \mathbf{R}^{N-1}, \pm x_N > 0\}.$$

The following lemma is proved in [22] (cf. also [11, Theorem III.2.1]).

Lemma 2.1. *Let $q \in (1, \infty)$. Then, for any $\mathbf{f}_\pm \in F_q(\mathbf{R}_\pm^N)$, there exists a sequence $\{\mathbf{f}_\pm^{(j)}\}_{j=1}^\infty \subset C_0^\infty(\mathbf{R}^N)^N$ such that $\lim_{j \rightarrow \infty} \|\mathbf{f}_\pm^{(j)} - \mathbf{f}_\pm\|_{F_q(\mathbf{R}_\pm^N)} = 0$.*

For positive constants ρ_\pm , we set

$$(2.1) \quad A_\pm = \sqrt{\rho_\pm \lambda + |\xi'|^2}, \quad (\xi', \lambda) \in \mathbf{R}^{N-1} \times (\mathbf{C} \setminus (-\infty, 0]),$$

where we have chosen a branch cut along the negative real axis and a branch of the square root so that $\Re \sqrt{z} > 0$ for $z \in \mathbf{C} \setminus (-\infty, 0]$. In addition, we set

$$\Sigma_{\sigma, \lambda_0} = \{z \in \mathbf{C} : |\arg z| < \pi - \sigma, |\lambda| > \lambda_0\} \quad (0 < \sigma < \pi/2, \lambda_0 \geq 0).$$

Then we have

Lemma 2.2. *Let $s \in \mathbf{R}$, $\alpha' \in \mathbf{N}_0^{N-1}$, and $\xi' \in \mathbf{R}^{N-1}$*

(1) Let $\sigma \in (0, \pi/2)$ and $\lambda \in \Sigma_{\sigma,0}$. Then

$$\begin{aligned} |\partial_{\xi'}^{\alpha'} A_{\pm}^s| &\leq C(|\lambda|^{1/2} + |\xi'|)^{s-|\alpha'|} \quad (\xi' \in \mathbf{R}^{N-1} \setminus \{0\}), \\ |\partial_{\xi'}^{\alpha'} (\rho_+ A_- + \rho_- A_+)^s| &\leq C(|\lambda|^{1/2} + |\xi'|)^{s-|\alpha'|} \quad (\xi' \in \mathbf{R}^{N-1} \setminus \{0\}), \end{aligned}$$

where $C = C(N, s, \alpha', \sigma, \rho_+, \rho_-)$ is a positive constant.

(2) There exists a positive constant $C = C(N, s, \alpha')$ such that

$$|\partial_{\xi'}^{\alpha'} |\xi'|^s| \leq C|\xi'|^{s-|\alpha'|}.$$

Proof. The proof is similar to [25, Lemma 5.2], so that the detailed proof may be omitted. \square

For $a = (a_1, \dots, a_{N-1}, a_N)$, we set $a' = (a_1, \dots, a_{N-1})$. Then the partial Fourier transform of $f = f(x)$, $x = (x_1, \dots, x_N)$, and its inverse transform are respectively defined by

$$\begin{aligned} (2.2) \quad \widehat{f}(x_N) &= \widehat{f}(\xi', x_N) = \int_{\mathbf{R}^{N-1}} e^{-ix' \cdot \xi'} f(x', x_N) dx', \\ \mathcal{F}_{\xi'}^{-1}[\widehat{f}(\xi', x_N)](x') &= \frac{1}{(2\pi)^{N-1}} \int_{\mathbf{R}^{N-1}} e^{ix' \cdot \xi'} \widehat{f}(\xi', x_N) d\xi'. \end{aligned}$$

The following two lemmas are proved in [25, Lemma 5.4].

Lemma 2.3. Let $q \in (1, \infty)$ and $\sigma \in (0, \pi/2)$. Assume that $k(\xi', \lambda)$ and $\ell(\xi', \lambda)$ are defined on $(\mathbf{R}^{N-1} \setminus \{0\}) \times \Sigma_{\sigma,0}$, which are many times differentiable with respect to ξ' , and satisfy for any multi-index $\alpha' \in \mathbf{N}_0^{N-1}$ and $(\xi', \lambda) \in (\mathbf{R}^{N-1} \setminus \{0\}) \times \Sigma_{\sigma,0}$

$$|\partial_{\xi'}^{\alpha'} k(\xi', \lambda)| \leq c_1(\alpha')(|\lambda|^{1/2} + |\xi'|)^{-|\alpha'|}, \quad |\partial_{\xi'}^{\alpha'} \ell(\xi', \lambda)| \leq c_1(\alpha')|\xi'|^{-|\alpha'|},$$

where $c_1(\alpha')$ is a positive constant independent of ξ' and λ . Furthermore, define the operators $K_j(\lambda)$ and $L_j(\lambda)$ ($j = 1, 2, \lambda \in \Sigma_{\sigma,0}$) by the formulas:

$$\begin{aligned} [K_1(\lambda)f_{\pm}](x) &= \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[k(\xi', \lambda) \lambda^{1/2} e^{-(Ax_N + By_N)} \widehat{f}_{\pm}(\xi', \pm y_N) \right] (x') dy_N, \\ [L_1(\lambda)f_{\pm}](x) &= \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\ell(\xi', \lambda) |\xi'| e^{-(Ax_N + By_N)} \widehat{f}_{\pm}(\xi', \pm y_N) \right] (x') dy_N, \end{aligned}$$

where $A, B \in \{A_+, A_-\}$ and $x = (x', x_N) \in \mathbf{R}_+^N$;

$$\begin{aligned} [K_2(\lambda)f_{\pm}](x) &= \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[k(\xi', \lambda) \lambda^{1/2} e^{-(-Ax_N + By_N)} \widehat{f}_{\pm}(\xi', \pm y_N) \right] (x') dy_N, \\ [L_2(\lambda)f_{\pm}](x) &= \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\ell(\xi', \lambda) |\xi'| e^{-(-Ax_N + By_N)} \widehat{f}_{\pm}(\xi', \pm y_N) \right] (x') dy_N, \end{aligned}$$

where $A, B \in \{A_+, A_-\}$ and $x = (x', x_N) \in \mathbf{R}_-^N$. Then,

$$K_1(\lambda), L_1(\lambda) \in \mathcal{L}(L_q(\mathbf{R}_+^N), L_q(\mathbf{R}_+^N)), \quad K_2(\lambda), L_2(\lambda) \in \mathcal{L}(L_q(\mathbf{R}_+^N), L_q(\mathbf{R}_-^N)),$$

and also their operator norms do not exceed some positive constant C depending on $N, q, \sigma, \rho_+, \rho_-$, and $\max\{c_1(\alpha') : |\alpha'| \leq N+1\}$, but independent of λ .

Lemma 2.4. Let $q \in (1, \infty)$. Assume that $m(\xi')$ is defined on $\mathbf{R}^{N-1} \setminus \{0\}$, which is many times differentiable with respect to ξ' , and satisfies for any multi-index $\alpha' \in \mathbf{N}_0^{N-1}$ and $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$

$$|\partial_{\xi'}^{\alpha'} m(\xi')| \leq c_2(\alpha')|\xi'|^{-|\alpha'|},$$

where $c_2(\alpha')$ is a positive constant independent of ξ' . Furthermore, define the operators M_j ($j = 1, 2$) by the formulas:

$$[M_1 f_{\pm}](x) = \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[m(\xi') |\xi'| e^{-|\xi'| (x_N + y_N)} \widehat{f}_{\pm}(\xi', \pm y_N) \right] (x') dy_N$$

for $x = (x', x_N) \in \mathbf{R}_+^N$;

$$[M_2 f_{\pm}](x) = \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[m(\xi') |\xi'| e^{-|\xi'| (-x_N + y_N)} \widehat{f}_{\pm}(\xi', \pm y_N) \right] (x') dy_N$$

for $x = (x', x_N) \in \mathbf{R}_-^N$. Then,

$$M_1 \in \mathcal{L}(L_q(\mathbf{R}_\pm^N), L_q(\mathbf{R}_\pm^N)), \quad M_2 \in \mathcal{L}(L_q(\mathbf{R}_\pm^N), L_q(\mathbf{R}_\mp^N)),$$

and also their operator norms do not exceed some positive constant C depending on N, q , and $\max\{c_2(\alpha') : |\alpha'| \leq N + 1\}$.

We next prove

Lemma 2.5. *Let $q \in (1, \infty)$ and $\sigma \in (0, \pi/2)$. Assume that $k(\xi', \lambda)$ and $\ell(\xi', \lambda)$ are defined on $(\mathbf{R}^{N-1} \times \{0\}) \times \Sigma_{\sigma, 0}$, which are many times differentiable with respect to ξ' , and satisfy for any multi-index $\alpha' \in \mathbf{N}_0^{N-1}$ and $(\xi', \lambda) \in (\mathbf{R}^{N-1} \setminus \{0\}) \times \Sigma_{\sigma, 0}$*

$$|\partial_{\xi'}^{\alpha'} k(\xi', \lambda)| \leq c_3(\alpha') (|\lambda|^{1/2} + |\xi'|)^{-|\alpha'|}, \quad |\partial_{\xi'}^{\alpha'} \ell(\xi', \lambda)| \leq c_3(\alpha') (|\lambda|^{1/2} + |\xi'|)^{-1-|\alpha'|}$$

where $c_3(\alpha')$ is a positive constant independent of ξ' and λ . Furthermore, define the operators $K_j(\lambda)$ and $L_j(\lambda)$ ($j = 1, 2, \lambda \in \Sigma_{\sigma, 0}$) by the formulas:

$$[K_1(\lambda) f_{\pm}](x) = \mathcal{F}_{\xi'}^{-1} \left[k(\xi', \lambda) e^{-Ax_N} \widehat{f}_{\pm}(\xi', 0) \right] (x'),$$

$$[L_1(\lambda) f_{\pm}](x) = \mathcal{F}_{\xi'}^{-1} \left[\ell(\xi', \lambda) e^{-Ax_N} \widehat{f}_{\pm}(\xi', 0) \right] (x'),$$

where $A \in \{A_+, A_-\}$ and $x = (x', x_N) \in \mathbf{R}_\pm^N$;

$$[K_2(\lambda) f_{\pm}](x) = \mathcal{F}_{\xi'}^{-1} \left[k(\xi', \lambda) e^{Ax_N} \widehat{f}_{\pm}(\xi', 0) \right] (x'),$$

$$[L_2(\lambda) f_{\pm}](x) = \mathcal{F}_{\xi'}^{-1} \left[\ell(\xi', \lambda) e^{Ax_N} \widehat{f}_{\pm}(\xi', 0) \right] (x'),$$

where $A \in \{A_+, A_-\}$ and $x = (x', x_N) \in \mathbf{R}_\pm^N$. Then the following assertions hold.

(1) $K_1(\lambda) \in \mathcal{L}(H_q^1(\mathbf{R}_\pm^N), H_q^1(\mathbf{R}_\pm^N))$ with

$$(2.3) \quad \|(\lambda^{1/2} K_1(\lambda) f_{\pm}, \nabla K_1(\lambda) f_{\pm})\|_{L_q(\mathbf{R}_\pm^N)} \leq C \|(\lambda^{1/2} f_{\pm}, \nabla f_{\pm})\|_{L_q(\mathbf{R}_\pm^N)},$$

while $K_2(\lambda) \in \mathcal{L}(H_q^1(\mathbf{R}_\pm^N), H_q^1(\mathbf{R}_\mp^N))$ with

$$(2.4) \quad \|(\lambda^{1/2} K_2(\lambda) f_{\pm}, \nabla K_2(\lambda) f_{\pm})\|_{L_q(\mathbf{R}_\mp^N)} \leq C \|(\lambda^{1/2} f_{\pm}, \nabla f_{\pm})\|_{L_q(\mathbf{R}_\pm^N)}.$$

In addition, $K_1(\lambda) \in \mathcal{L}(H_q^2(\mathbf{R}_\pm^N), H_q^2(\mathbf{R}_\pm^N))$ with

$$(2.5) \quad \|(\lambda K_1(\lambda) f_{\pm}, \lambda^{1/2} \nabla K_1(\lambda) f_{\pm}, \nabla^2 K_1(\lambda) f_{\pm})\|_{L_q(\mathbf{R}_\pm^N)} \\ \leq C \|(\lambda f_{\pm}, \lambda^{1/2} \nabla f_{\pm}, \nabla^2 f_{\pm})\|_{L_q(\mathbf{R}_\pm^N)},$$

while $K_2(\lambda) \in \mathcal{L}(H_q^2(\mathbf{R}_\pm^N), H_q^2(\mathbf{R}_\mp^N))$ with

$$(2.6) \quad \|(\lambda K_2(\lambda) f_{\pm}, \lambda^{1/2} \nabla K_2(\lambda) f_{\pm}, \nabla^2 K_2(\lambda) f_{\pm})\|_{L_q(\mathbf{R}_\mp^N)} \\ \leq C \|(\lambda f_{\pm}, \lambda^{1/2} \nabla f_{\pm}, \nabla^2 f_{\pm})\|_{L_q(\mathbf{R}_\pm^N)}.$$

(2) $L_1(\lambda) \in \mathcal{L}(H_q^1(\mathbf{R}_\pm^N), H_q^2(\mathbf{R}_\pm^N))$ with

$$(2.7) \quad \begin{aligned} & \|(\lambda L_1(\lambda)f_\pm, \lambda^{1/2}\nabla L_1(\lambda)f_\pm, \nabla^2 L_1(\lambda)f_\pm)\|_{L_q(\mathbf{R}_\pm^N)} \\ & \leq C\|(\lambda^{1/2}f_\pm, \nabla f_\pm)\|_{L_q(\mathbf{R}_\pm^N)}, \end{aligned}$$

$$(2.8) \quad \begin{aligned} & \|(\lambda^{1/2}L_1(\lambda)f_\pm, \nabla L_1(\lambda)f_\pm)\|_{L_q(\mathbf{R}_\pm^N)} \\ & \leq C\left(\|f_\pm\|_{L_q(\mathbf{R}_\pm^N)} + |\lambda|^{-1/2}\|\nabla f_\pm\|_{L_q(\mathbf{R}_\pm^N)}\right), \end{aligned}$$

while $L_2(\lambda) \in \mathcal{L}(H_q^1(\mathbf{R}_\pm^N), H_q^2(\mathbf{R}_\pm^N))$ with

$$(2.9) \quad \begin{aligned} & \|(\lambda L_2(\lambda)f_\pm, \lambda^{1/2}\nabla L_2(\lambda)f_\pm, \nabla^2 L_2(\lambda)f_\pm)\|_{L_q(\mathbf{R}_\pm^N)} \\ & \leq C\|(\lambda^{1/2}f_\pm, \nabla f_\pm)\|_{L_q(\mathbf{R}_\pm^N)}, \end{aligned}$$

$$(2.10) \quad \begin{aligned} & \|(\lambda^{1/2}L_2(\lambda)f_\pm, \nabla L_2(\lambda)f_\pm)\|_{L_q(\mathbf{R}_\pm^N)} \\ & \leq C\left(\|f_\pm\|_{L_q(\mathbf{R}_\pm^N)} + |\lambda|^{-1/2}\|\nabla f_\pm\|_{L_q(\mathbf{R}_\pm^N)}\right). \end{aligned}$$

Here C is a positive constant depending on N , q , σ , ρ_+ , ρ_- , and $\max\{c_3(\alpha') : |\alpha'| \leq N+1\}$, but independent of λ .

Proof. (1). First, we prove (2.3). Let $x \in \mathbf{R}_\pm^N$, and then note that

$$(2.11) \quad \begin{aligned} e^{-Ax_N} \widehat{f}_\pm(\xi', 0) &= -\int_0^\infty \frac{\partial}{\partial y_N} \left(e^{-A(x_N+y_N)} \widehat{f}_\pm(\xi', \pm y_N) \right) dy_N \\ &= \int_0^\infty A e^{-A(x_N+y_N)} \widehat{f}_\pm(\xi', \pm y_N) dy_N \\ &\mp \int_0^\infty e^{-A(x_N+y_N)} \widehat{\partial_N f}_\pm(\xi', \pm y_N) dy_N. \end{aligned}$$

By these relations, we have

$$\begin{aligned} [K_1(\lambda)f_\pm](x) &= \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[k(\xi', \lambda) A e^{-A(x_N+y_N)} \widehat{f}_\pm(\xi', \pm y_N) \right] (x') dy_N \\ &\mp \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[k(\xi', \lambda) e^{-A(x_N+y_N)} \widehat{\partial_N f}_\pm(\xi', \pm y_N) \right] (x') dy_N \\ &=: I_\pm \mp J_\pm. \end{aligned}$$

Let $R(A) = \rho_+$ when $A = A_+$ and $R(A) = \rho_-$ when $A = A_-$, which gives us the following formulas:

$$(2.12) \quad A = \frac{A^2}{A} = \frac{1}{A} \left(R(A)\lambda - \sum_{j=1}^{N-1} (i\xi_j)^2 \right), \quad 1 = \frac{A^2}{A^2} = \frac{1}{A^2} \left(R(A)\lambda - \sum_{j=1}^{N-1} (i\xi_j)^2 \right).$$

By using these formulas, we can write I_\pm and J_\pm as

$$\begin{aligned} I_\pm &= \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\frac{kR(A)}{A} \lambda^{1/2} e^{-A(x_N+y_N)} \widehat{\lambda^{1/2} f}_\pm(\xi', \pm y_N) \right] (x') dy_N \\ &\quad - \sum_{j=1}^{N-1} \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\frac{ki\xi_j}{A|\xi'|} |\xi'| e^{-A(x_N+y_N)} \widehat{\partial_j f}_\pm(\xi', \pm y_N) \right] (x') dy_N, \end{aligned}$$

$$\begin{aligned}
 J_{\pm} &= \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{kR(A)\lambda^{1/2}}{A^2} \lambda^{1/2} e^{-A(x_N+y_N)} \widehat{\partial_N f_{\pm}}(\xi', \pm y_N) \right] (x') dy_N \\
 &\quad - \sum_{j=1}^{N-1} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{k(i\xi_j)^2}{A^2|\xi'|} |\xi'| e^{-A(x_N+y_N)} \widehat{\partial_N f_{\pm}}(\xi', \pm y_N) \right] (x') dy_N,
 \end{aligned}$$

where $k = k(\xi', \lambda)$. Set $\Xi = (i\xi_1, \dots, i\xi_{N-1}, -A)^T$, and thus

$$\begin{aligned}
 &(\lambda^{1/2}, \nabla) I_{\pm} \\
 &= \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{(\lambda^{1/2}, \Xi)kR(A)}{A} \lambda^{1/2} e^{-A(x_N+y_N)} \widehat{\lambda^{1/2} f_{\pm}}(\xi', \pm y_N) \right] (x') dy_N \\
 &\quad - \sum_{j=1}^{N-1} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{(\lambda^{1/2}, \Xi)k i \xi_j}{A|\xi'|} |\xi'| e^{-A(x_N+y_N)} \widehat{\partial_j f_{\pm}}(\xi', \pm y_N) \right] (x') dy_N, \\
 &(\lambda^{1/2}, \nabla) J_{\pm} \\
 &= \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{(\lambda^{1/2}, \Xi)kR(A)\lambda^{1/2}}{A^2} \lambda^{1/2} e^{-A(x_N+y_N)} \widehat{\partial_N f_{\pm}}(\xi', \pm y_N) dy_N \right] (x') dy_N \\
 &\quad - \sum_{j=1}^{N-1} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{(\lambda^{1/2}, \Xi)k(i\xi_j)^2}{A^2|\xi'|} |\xi'| e^{-A(x_N+y_N)} \widehat{\partial_N f_{\pm}}(\xi', \pm y_N) dy_N \right] (x') dy_N.
 \end{aligned}$$

By Lemma 2.2 and Leibniz's formula, we have for any multi-index $\alpha' \in \mathbf{N}_0^{N-1}$

$$\begin{aligned}
 \left| \partial_{\xi'}^{\alpha'} \left\{ \frac{(\lambda^{1/2}, \Xi)k(\xi', \lambda)R(A)}{A} \right\} \right| &\leq C|\xi'|^{-|\alpha'|} \quad (\xi' \in \mathbf{R}^{N-1} \setminus \{0\}), \\
 \left| \partial_{\xi'}^{\alpha'} \left\{ \frac{(\lambda^{1/2}, \Xi)k(\xi', \lambda) i \xi_j}{A|\xi'|} \right\} \right| &\leq C|\xi'|^{-|\alpha'|} \quad (\xi' \in \mathbf{R}^{N-1} \setminus \{0\}), \\
 \left| \partial_{\xi'}^{\alpha'} \left\{ \frac{(\lambda^{1/2}, \Xi)k(\xi', \lambda)R(A)\lambda^{1/2}}{A^2} \right\} \right| &\leq C|\xi'|^{-|\alpha'|} \quad (\xi' \in \mathbf{R}^{N-1} \setminus \{0\}), \\
 \left| \partial_{\xi'}^{\alpha'} \left\{ \frac{(\lambda^{1/2}, \Xi)k(\xi', \lambda) i \xi_j}{A^2|\xi'|} \right\} \right| &\leq C|\xi'|^{-|\alpha'|} \quad (\xi' \in \mathbf{R}^{N-1} \setminus \{0\}),
 \end{aligned}$$

with some positive constant $C = C(N, \alpha', \sigma, \rho_+, \rho_-)$ independent of ξ' and λ . Combining these properties with Lemma 2.3 furnishes

$$\begin{aligned}
 \|(\lambda^{1/2} I_{\pm}, \nabla I_{\pm})\|_{L_q(\mathbf{R}_+^N)} &\leq C \|(\lambda^{1/2} f_{\pm}, \nabla f_{\pm})\|_{L_q(\mathbf{R}_{\pm}^N)}, \\
 \|(\lambda^{1/2} J_{\pm}, \nabla J_{\pm})\|_{L_q(\mathbf{R}_+^N)} &\leq C \|(\lambda^{1/2} f_{\pm}, \nabla f_{\pm})\|_{L_q(\mathbf{R}_{\pm}^N)},
 \end{aligned}$$

which implies (2.3) holds. Analogously, we can prove (2.4)-(2.6). This completes the proof of (1).

(2). The estimates (2.7) and (2.9) follow respectively from (2.3) and (2.4). In addition, (2.8) and (2.10) follow respectively from (2.7) and (2.9). This completes the proof of (2). \square

Let $\text{sign}(a)$ be the sign function of a , that is, $\text{sign}(a) = 1$ when $a > 0$, $\text{sign}(a) = -1$ when $a < 0$, and $\text{sign}(a) = 0$ when $a = 0$. Then we have

Lemma 2.6. *Let $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$ and $\varepsilon > 0$. Then, for any $\lambda \in \mathbf{C} \setminus (-\infty, 0]$,*

$$(2.13) \quad \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-\varepsilon|\xi|^2} \frac{i\xi_N e^{i a \xi_N}}{\rho_{\pm} \lambda + |\xi|^2} d\xi_N = -\frac{\text{sign}(a)}{2} e^{\varepsilon \rho_{\pm} \lambda} e^{-A_{\pm} |\lambda|},$$

where $\xi = (\xi', \xi_N)$. In addition,

$$(2.14) \quad \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-\varepsilon|\xi|^2} \frac{i\xi_N e^{ia\xi_N}}{|\xi|^2} d\xi_N = -\frac{\text{sign}(a)}{2} e^{-|\xi'| |a|}.$$

Proof. These formulas follow from the residue theorem immediately, so that the detailed proof may be omitted. \square

3. PROBLEMS IN THE WHOLE SPACE WITH FLAT INTERFACE

Let us introduce the flat interface:

$$\mathbf{R}_0^N = \{x = (x', x_N) : x' = (x_1, \dots, x_{N-1}) \in \mathbf{R}^{N-1}, x_N = 0\}.$$

This section mainly considers two strong elliptic problems as follows:

$$(3.1) \quad \begin{cases} \rho_{\pm} \lambda v_{\pm} - \Delta v_{\pm} = -\text{div } \mathbf{f}_{\pm} + g_{\pm} & \text{in } \mathbf{R}_{\pm}^N, \\ \rho_+ v_+ = \rho_- v_- & \text{on } \mathbf{R}_0^N, \\ \partial_N v_+ - \partial_N v_- = f_{+N} - f_{-N} + h_+ - h_- & \text{on } \mathbf{R}_0^N, \end{cases}$$

where λ is the resolvent parameter varying in $\Sigma_{\sigma,0}$ ($0 < \sigma < \pi/2$), and

$$(3.2) \quad \begin{cases} \Delta v_{\pm} = \text{div } \mathbf{f}_{\pm} & \text{in } \mathbf{R}_{\pm}^N, \\ \rho_+ v_+ = \rho_- v_- & \text{on } \mathbf{R}_0^N, \\ \partial_N v_+ - \partial_N v_- = f_{+N} - f_{-N} & \text{on } \mathbf{R}_0^N. \end{cases}$$

Throughout this section, we assume that ρ_{\pm} are positive constants. Concerning (3.1) and (3.2), we prove the following theorems.

Theorem 3.1. *Let $\sigma \in (0, \pi/2)$ and $q \in (1, \infty)$. Then, for any*

$$\mathbf{f}_{\pm} = (f_{\pm 1}, \dots, f_{\pm N})^{\top} \in F_q(\mathbf{R}_{\pm}^N), \quad g_{\pm} \in L_q(\mathbf{R}_{\pm}^N), \quad h_{\pm} \in H_q^1(\mathbf{R}_{\pm}^N)$$

and for any $\lambda \in \Sigma_{\sigma,0}$, the strong elliptic problem (3.1) admits unique solutions $v_{\pm} \in H_q^2(\mathbf{R}_{\pm}^N)$. In addition, the solutions v_{\pm} satisfy

$$(3.3) \quad \begin{aligned} & \sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda v_{\mathfrak{s}}, \lambda^{1/2} \nabla v_{\mathfrak{s}}, \nabla^2 v_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} \\ & \leq C_1 \sum_{\mathfrak{s} \in \{+, -\}} \|(\text{div } \mathbf{f}_{\mathfrak{s}}, g_{\mathfrak{s}}, \lambda^{1/2} f_{\mathfrak{s}N}, \nabla f_{\mathfrak{s}N}, \lambda^{1/2} h_{\mathfrak{s}}, \nabla h_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)}, \end{aligned}$$

and also

$$(3.4) \quad \begin{aligned} & \sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda^{1/2} v_{\mathfrak{s}}, \nabla v_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} \leq C_1 \sum_{\mathfrak{s} \in \{+, -\}} \left(\|\mathbf{f}_{\mathfrak{s}}\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} \right. \\ & \quad \left. + |\lambda|^{-1/2} \|g_{\mathfrak{s}}\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} + \|h_{\mathfrak{s}}\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} + |\lambda|^{-1/2} \|\nabla h_{\mathfrak{s}}\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} \right), \end{aligned}$$

where $C_1 = C_1(N, q, \sigma, \rho_+, \rho_-)$ is a positive constant independent of λ . Additionally, if $f_{+N} - f_{-N} = 0$ on \mathbf{R}_0^N , then v_{\pm} satisfy

$$(3.5) \quad \begin{aligned} & \sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda v_{\mathfrak{s}}, \lambda^{1/2} \nabla v_{\mathfrak{s}}, \nabla^2 v_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} \\ & \leq C_1 \sum_{\mathfrak{s} \in \{+, -\}} \|(\text{div } \mathbf{f}_{\mathfrak{s}}, g_{\mathfrak{s}}, \lambda^{1/2} h_{\mathfrak{s}}, \nabla h_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)}. \end{aligned}$$

Theorem 3.2. *Let $q \in (1, \infty)$.*

- (1) **Existence.** Let $\mathbf{f}_\pm = (f_{\pm 1}, \dots, f_{\pm N})^\top \in F_q(\mathbf{R}_\pm^N)$. Then the strong problem (3.2) admits solutions $v_\pm \in \widehat{H}_q^1(\mathbf{R}_\pm^N) \cap \widehat{H}_q^2(\mathbf{R}_\pm^N)$ satisfying

$$(3.6) \quad \sum_{s \in \{+, -\}} \|\nabla^2 v_s\|_{L_q(\mathbf{R}_s^N)} \leq C_2 \sum_{s \in \{+, -\}} \|(\operatorname{div} \mathbf{f}_s, \nabla f_{sN})\|_{L_q(\mathbf{R}_s^N)},$$

$$(3.7) \quad \sum_{s \in \{+, -\}} \|\nabla v_s\|_{L_q(\mathbf{R}_s^N)} \leq C_2 \sum_{s \in \{+, -\}} \|\mathbf{f}_s\|_{L_q(\mathbf{R}_s^N)},$$

with some positive constant $C_2 = C_2(N, q, \rho_+, \rho_-)$. Additionally, if $f_{+N} - f_{-N} = 0$ on \mathbf{R}_0^N , then v_\pm satisfy

$$(3.8) \quad \sum_{s \in \{+, -\}} \|\nabla^2 v_s\|_{L_q(\mathbf{R}_s^N)} \leq C_2 \sum_{s \in \{+, -\}} \|\operatorname{div} \mathbf{f}_s\|_{L_q(\mathbf{R}_s^N)}.$$

- (2) **Uniqueness.** If $v_\pm \in \widehat{H}_q^1(\mathbf{R}_\pm^N) \cap \widehat{H}_q^2(\mathbf{R}_\pm^N)$ satisfy

$$(3.9) \quad \Delta v_\pm = 0 \text{ in } \mathbf{R}_\pm^N, \quad \rho_+ v_+ = \rho_- v_- \text{ and } \partial_N v_+ - \partial_N v_- = 0 \text{ on } \mathbf{R}_0^N$$

then $v_\pm = \rho_\pm^{-1} c$ for some constant c .

3.1. Auxiliary problems. We start with

$$(3.10) \quad \rho_\pm \lambda U_{\pm, \lambda} - \Delta U_{\pm, \lambda} = -\operatorname{div} \mathbf{f}_\pm + g_\pm \quad \text{in } \mathbf{R}_\pm^N.$$

For this problem, we prove

Lemma 3.3. Let $\sigma \in (0, \pi/2)$ and $q \in (1, \infty)$. Then, for any $\lambda \in \Sigma_{\sigma, 0}$, $\mathbf{f}_\pm = (f_{\pm 1}, \dots, f_{\pm N})^\top \in C_0^\infty(\mathbf{R}^N)^N$, and $g_\pm \in C_0^\infty(\mathbf{R}_\pm^N)$, there exist $U_{\pm, \lambda} \in H_q^2(\mathbf{R}^N)$, satisfying (3.10), such that the following assertions hold.

- (1) There hold the estimates:

$$\begin{aligned} \|(\lambda U_{\pm, \lambda}, \lambda^{1/2} \nabla U_{\pm, \lambda}, \nabla^2 U_{\pm, \lambda})\|_{L_q(\mathbf{R}^N)} &\leq C \|(\operatorname{div} \mathbf{f}_\pm, g_\pm)\|_{L_q(\mathbf{R}_\pm^N)}, \\ \|(\lambda^{1/2} U_{\pm, \lambda}, \nabla U_{\pm, \lambda})\|_{L_q(\mathbf{R}^N)} &\leq C \left(\|\mathbf{f}_\pm\|_{L_q(\mathbf{R}_\pm^N)} + |\lambda|^{-1/2} \|g_\pm\|_{L_q(\mathbf{R}_\pm^N)} \right), \end{aligned}$$

with a positive constant $C = C(N, q, \sigma, \rho_+, \rho_-)$ independent of λ .

- (2) The traces of $\partial_N U_{\pm, \lambda}$ on \mathbf{R}_0^N are given by

$$\begin{aligned} (\partial_N U_{\pm, \lambda})(x', 0) &= f_{\pm N}(x', 0) \mp \sum_{j=1}^{N-1} \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[i \xi_j e^{-A_\pm y_N} \widehat{f}_{\pm j}(\xi', \pm y_N) \right] (x') dy_N \\ &\quad - \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[A_\pm e^{-A_\pm y_N} \widehat{f}_{\pm N}(\xi', \pm y_N) \right] (x') dy_N, \end{aligned}$$

where A_\pm are defined as (2.1) and the symbols $\mathcal{F}_{\xi'}^{-1}$ and $\widehat{\cdot}$ are defined as (2.2).

Proof. Let us introduce the Fourier transform of $u = u(x)$ on \mathbf{R}^N and the inverse Fourier transform of $v = v(\xi)$ on \mathbf{R}^N as follows:

$$\mathcal{F}[u](\xi) = \int_{\mathbf{R}^N} e^{-ix \cdot \xi} u(x) dx, \quad \mathcal{F}_\xi^{-1}[v](x) = \frac{1}{(2\pi)^N} \int_{\mathbf{R}^N} e^{ix \cdot \xi} v(\xi) d\xi.$$

Let $u_\pm = u_\pm(x', x_N)$ be functions defined on \mathbf{R}_\pm^N , and define the odd extensions $E_\pm^o u_\pm$ of u_\pm and the even extensions $E_\pm^e u_\pm$ of u_\pm as follows:

$$(E_+^o u_+)(x) = \begin{cases} u_+(x', x_N) & (x_N > 0), \\ -u_+(x', -x_N) & (x_N < 0), \end{cases}$$

$$\begin{aligned} (E_-^o u_-)(x) &= \begin{cases} -u_-(x', -x_N) & (x_N > 0), \\ u_-(x', x_N) & (x_N < 0), \end{cases} \\ (E_+^e u_+)(x) &= \begin{cases} u_+(x', x_N) & (x_N > 0), \\ u_+(x', -x_N) & (x_N < 0), \end{cases} \\ (E_-^e u_-)(x) &= \begin{cases} u_-(x', -x_N) & (x_N > 0), \\ u_-(x', x_N) & (x_N < 0). \end{cases} \end{aligned}$$

In addition, we set

$$(3.11) \quad \mathbf{F}_\pm = (F_{\pm 1}, \dots, F_{\pm N-1}, F_{\pm N})^\top = (E_\pm^o f_{\pm 1}, \dots, E_\pm^o f_{\pm N-1}, E_\pm^e f_{\pm N})^\top.$$

It then holds that

$$\partial_j F_{\pm j} = E_\pm^o \partial_j f_{\pm j} \quad (j = 1, \dots, N-1), \quad \partial_N F_{\pm N} = E_\pm^o \partial_N f_{\pm N},$$

which implies

$$(3.12) \quad \operatorname{div} \mathbf{F}_\pm = E_\pm^o z_\pm \quad \text{for } z_\pm = \operatorname{div} \mathbf{f}_\pm.$$

We thus see that

$$(3.13) \quad \mathcal{F}[\operatorname{div} \mathbf{F}_\pm](\xi) = i\xi \cdot \mathcal{F}[\mathbf{F}_\pm](\xi) = \mathcal{F}[E_\pm^o z_\pm](\xi).$$

Let us define for $\varepsilon > 0$

$$V_\pm^\varepsilon(x) = -\mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{i\xi \cdot \mathcal{F}[\mathbf{F}_\pm](\xi)}{\rho_\pm \lambda + |\xi|^2} \right] (x)$$

and

$$(3.14) \quad F_{\pm, \operatorname{div}}^\varepsilon = \mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \mathcal{F}[\operatorname{div} \mathbf{F}_\pm](\xi) \right] (x).$$

Then V_\pm^ε solve by (3.13)

$$(3.15) \quad \rho_\pm \lambda V_\pm^\varepsilon - \Delta V_\pm^\varepsilon = -F_{\pm, \operatorname{div}}^\varepsilon \quad \text{in } \mathbf{R}^N.$$

On the other hand, one has

$$\begin{aligned} \lambda^{1/2} V_\pm^\varepsilon(x) &= -\mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{\lambda^{1/2} i\xi \cdot \mathcal{F}[\mathbf{F}_\pm](\xi)}{\rho_\pm \lambda + |\xi|^2} \right] (x), \\ \partial_j V_\pm^\varepsilon(x) &= \mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{\xi_j \xi \cdot \mathcal{F}[\mathbf{F}_\pm](\xi)}{\rho_\pm \lambda + |\xi|^2} \right] (x) \quad (j = 1, \dots, N), \end{aligned}$$

and also by (3.13)

$$\begin{aligned} \lambda V_\pm^\varepsilon(x) &= -\mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{\lambda}{\rho_\pm \lambda + |\xi|^2} \mathcal{F}[\operatorname{div} \mathbf{F}_\pm](\xi) \right] (x), \\ \lambda^{1/2} \partial_j V_\pm^\varepsilon(x) &= -\mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{\lambda^{1/2} i\xi_j}{\rho_\pm \lambda + |\xi|^2} \mathcal{F}[\operatorname{div} \mathbf{F}_\pm](\xi) \right] (x) \quad (j = 1, \dots, N), \\ \partial_k \partial_l V_\pm^\varepsilon(x) &= \mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{\xi_k \xi_l}{\rho_\pm \lambda + |\xi|^2} \mathcal{F}[\operatorname{div} \mathbf{F}_\pm](\xi) \right] (x) \quad (k, l = 1, \dots, N). \end{aligned}$$

For $j, k, l = 1, \dots, N$ and for any multi-index $\alpha \in \mathbf{N}_0^N$,

$$\left| \partial_\xi^\alpha \left(\frac{\lambda}{\rho_\pm \lambda + |\xi|^2} \right) \right| \leq C |\xi|^{-|\alpha|} \quad (\xi \in \mathbf{R}^N \setminus \{0\}),$$

$$\begin{aligned} \left| \partial_\xi^\alpha \left(\frac{\lambda^{1/2} i \xi_j}{\rho_\pm \lambda + |\xi|^2} \right) \right| &\leq C |\xi|^{-|\alpha|} \quad (\xi \in \mathbf{R}^N \setminus \{0\}), \\ \left| \partial_\xi^\alpha \left(\frac{\xi_k \xi_l}{\rho_\pm \lambda + |\xi|^2} \right) \right| &\leq C |\xi|^{-|\alpha|} \quad (\xi \in \mathbf{R}^N \setminus \{0\}), \end{aligned}$$

with some positive constant $C = C(N, \alpha, \sigma)$ (cf. [25, Section 3]). Thus, applying the classical Fourier multiplier theorem to the above formulas of V_\pm^ε and setting

$$(3.16) \quad \mathbf{F}_\pm^\varepsilon = \mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \mathcal{F}[\mathbf{F}_\pm](\xi) \right] (x)$$

furnish the following estimates:

$$(3.17) \quad \begin{aligned} \|(\lambda^{1/2} V_\pm^\varepsilon, \nabla V_\pm^\varepsilon)\|_{L_q(\mathbf{R}^N)} &\leq C \|\mathbf{F}_\pm^\varepsilon\|_{L_q(\mathbf{R}^N)}, \\ \|(\lambda V_\pm^\varepsilon, \lambda^{1/2} \nabla V_\pm^\varepsilon, \nabla^2 V_\pm^\varepsilon)\|_{L_q(\mathbf{R}^N)} &\leq C \|F_{\pm, \text{div}}^\varepsilon\|_{L_q(\mathbf{R}^N)}, \end{aligned}$$

where C is a positive constant independent of ε . Analogously,

$$(3.18) \quad \begin{aligned} |\lambda| \|V_\pm^\varepsilon - V_\pm^{\varepsilon'}\|_{L_q(\mathbf{R}^N)} &\leq C \|F_{\pm, \text{div}}^\varepsilon - F_{\pm, \text{div}}^{\varepsilon'}\|_{L_q(\mathbf{R}^N)}, \\ |\lambda|^{1/2} \|\nabla(V_\pm^\varepsilon - V_\pm^{\varepsilon'})\|_{L_q(\mathbf{R}^N)} &\leq C \|F_{\pm, \text{div}}^\varepsilon - F_{\pm, \text{div}}^{\varepsilon'}\|_{L_q(\mathbf{R}^N)}, \\ \|\nabla^2(V_\pm^\varepsilon - V_\pm^{\varepsilon'})\|_{L_q(\mathbf{R}^N)} &\leq C \|F_{\pm, \text{div}}^\varepsilon - F_{\pm, \text{div}}^{\varepsilon'}\|_{L_q(\mathbf{R}^N)}, \end{aligned}$$

for $\varepsilon, \varepsilon' > 0$ and a positive constant C independent of ε and ε' .

Next, we compute the formulas of $\partial_N V_\pm^\varepsilon$ on \mathbf{R}_0^N . We have

$$\mathcal{F}[E_\pm^o z_\pm](\xi) = \int_0^\infty (\pm e^{-iy_N \xi_N} \mp e^{iy_N \xi_N}) \widehat{z}_\pm(\xi', \pm y_N) dy_N,$$

while by (3.13)

$$(3.19) \quad \partial_N V_\pm^\varepsilon = -\mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{i \xi_N}{\rho_\pm \lambda + |\xi|^2} \mathcal{F}[E_\pm^o z_\pm](\xi) \right] (x).$$

These formulas yield

$$\begin{aligned} \partial_N V_\pm^\varepsilon &= - \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\widehat{z}_\pm(\xi', \pm y_N) \right. \\ &\quad \cdot \left. \left(\frac{1}{2\pi} \int_{-\infty}^\infty e^{-\varepsilon|\xi|^2} \frac{i \xi_N}{\rho_\pm \lambda + |\xi|^2} (\pm e^{i(x_N - y_N)\xi_N} \mp e^{i(x_N + y_N)\xi_N}) d\xi_N \right) \right] (x') dy_N. \end{aligned}$$

Inserting (2.13) into the above formula of $\partial_N V_\pm^\varepsilon$ furnishes

$$(3.20) \quad (\partial_N V_\pm^\varepsilon)(x', 0) = \mp e^{\varepsilon \rho_\pm \lambda} \int_0^\infty \mathcal{F}_{\xi'}^{-1} [e^{-A \pm y_N} \widehat{z}_\pm(\xi', \pm y_N)] (x') dy_N.$$

Now we consider the limit: $\varepsilon \rightarrow 0$. By the well-known property of the heat kernel $\mathcal{F}_\xi^{-1}[e^{-\varepsilon|\xi|^2}](x)$, we have

$$\lim_{\varepsilon \rightarrow 0} \|\mathbf{F}_\pm^\varepsilon - \mathbf{F}_\pm\|_{L_q(\mathbf{R}^N)} = 0, \quad \lim_{\varepsilon \rightarrow 0} \|F_{\pm, \text{div}}^\varepsilon - \text{div } \mathbf{F}_\pm\|_{L_q(\mathbf{R}^N)} = 0.$$

Combining the last property with (3.18) shows that there exist $V_\pm \in H_q^2(\mathbf{R}^N)$ such that $\lim_{\varepsilon \rightarrow 0} \|V_\pm^\varepsilon - V_\pm\|_{H_q^2(\mathbf{R}^N)} = 0$. One then sees that V_\pm satisfy, by (3.12), (3.15), and (3.17), the equations:

$$\rho_\pm \lambda V_\pm - \Delta V_\pm = -\text{div } \mathbf{F}_\pm = -\text{div } \mathbf{f}_\pm \quad \text{in } \mathbf{R}_\pm^N$$

and the estimates:

$$(3.21) \quad \|(\lambda^{1/2} V_\pm, \nabla V_\pm)\|_{L_q(\mathbf{R}^N)} \leq C \|\mathbf{F}_\pm\|_{L_q(\mathbf{R}^N)} \leq C \|\mathbf{f}_\pm\|_{L_q(\mathbf{R}_\pm^N)},$$

$$\|(\lambda V_{\pm}, \lambda^{1/2} \nabla V_{\pm}, \nabla^2 V_{\pm})\|_{L_q(\mathbf{R}^N)} \leq C \|\operatorname{div} \mathbf{F}_{\pm}\|_{L_q(\mathbf{R}^N)} \leq C \|\operatorname{div} \mathbf{f}_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)},$$

while there holds by (3.20)

$$(\partial_N V_{\pm})(x', 0) = \mp \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} [e^{-A_{\pm} y_N} \widehat{z}_{\pm}(\xi', \pm y_N)](x') dy_N.$$

Combining these formulas with

$$\widehat{z}_{\pm}(\xi', \pm y_N) = \sum_{j=1}^{N-1} i \xi_j \widehat{f}_{\pm j}(\xi', \pm y_N) + (\partial_N \widehat{f}_{\pm N})(\xi', \pm y_N)$$

and with integration by parts furnishes

$$(3.22) \quad (\partial_N V_{\pm})(x', 0) = f_{\pm N}(x', 0) \\ \mp \sum_{j=1}^{N-1} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} [i \xi_j e^{-A_{\pm} y_N} \widehat{f}_{\pm j}(\xi', \pm y_N)](x') dy_N \\ - \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} [A_{\pm} e^{-A_{\pm} y_N} \widehat{f}_{\pm N}(\xi', \pm y_N)](x') dy_N.$$

Finally, we set

$$W_{\pm}(x) = \mathcal{F}_{\xi}^{-1} \left[\frac{\mathcal{F} [E_{\pm}^e g_{\pm}](\xi)}{\rho_{\pm} \lambda + |\xi|^2} \right] (x) \quad (x \in \mathbf{R}^N).$$

Then W_{\pm} satisfy the equations:

$$\rho_{\pm} \lambda W_{\pm} - \Delta W_{\pm} = E_{\pm}^e g_{\pm} = g_{\pm} \quad \text{in } \mathbf{R}_{\pm}^N,$$

and also satisfy, similarly to (3.17), the estimates:

$$(3.23) \quad \|(\lambda^{1/2} W_{\pm}, \nabla W_{\pm})\|_{L_q(\mathbf{R}^N)} \leq C |\lambda|^{-\frac{1}{2}} \|E_{\pm}^e g_{\pm}\|_{L_q(\mathbf{R}^N)} \leq C |\lambda|^{-\frac{1}{2}} \|g_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)}, \\ \|(\lambda W_{\pm}, \lambda^{1/2} \nabla W_{\pm}, \nabla^2 W_{\pm})\|_{L_q(\mathbf{R}^N)} \leq C \|E_{\pm}^e g_{\pm}\|_{L_q(\mathbf{R}^N)} \leq C \|g_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)},$$

where C is a positive constant independent of λ . In addition, we have²

$$(3.24) \quad \partial_N W_{\pm} = 0 \quad \text{on } \mathbf{R}_0^N.$$

Thus, setting $U_{\pm, \lambda} = V_{\pm} + W_{\pm}$, we see that $U_{\pm, \lambda}$ satisfy all of the properties required in Lemma 3.3 by (3.21)-(3.24). This completes the proof of Lemma 3.3. \square

Next, we consider the case $\lambda = 0$ and $g_{\pm} = 0$ of (3.10):

$$(3.25) \quad \Delta U_{\pm, 0} = \operatorname{div} \mathbf{f}_{\pm} \quad \text{in } \mathbf{R}_{\pm}^N.$$

Concerning these equations, we prove

Lemma 3.4. *Let $q \in (1, \infty)$. Then, for any $\mathbf{f}_{\pm} = (f_{\pm 1}, \dots, f_{\pm N})^{\top} \in C_0^{\infty}(\mathbf{R}^N)$, there exist $U_{\pm, 0} \in \widehat{H}_q^1(\mathbf{R}^N) \cap \widehat{H}_q^2(\mathbf{R}^N)$, satisfying (3.25), such that the following assertions hold.*

(1) *There hold the estimates:*

$$\|\nabla^2 U_{\pm, 0}\|_{L_q(\mathbf{R}^N)} \leq C \|\operatorname{div} \mathbf{f}_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)}, \quad \|\nabla U_{\pm, 0}\|_{L_q(\mathbf{R}^N)} \leq C \|\mathbf{f}_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)}, \\ \text{with a positive constant } C = C(N, q, \rho_+, \rho_-).$$

²These properties follow from the uniqueness of solutions to $\lambda u - \Delta u = f$ in \mathbf{R}^N , see e.g. [19, Subsection 2.4].

(2) The traces of $\partial_N U_{\pm,0}$ on \mathbf{R}_0^N are given by

$$\begin{aligned} (\partial_N U_{\pm,0})(x', 0) &= f_{\pm N}(x', 0) \mp \sum_{j=1}^{N-1} \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[i\xi_j e^{-|\xi'|y_N} \widehat{f}_{\pm j}(\xi', \pm y_N) \right] (x') dy_N \\ &\quad - \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[|\xi'| e^{-|\xi'|y_N} \widehat{f}_{\pm N}(\xi', \pm y_N) \right] (x') dy_N, \end{aligned}$$

where the symbols $\mathcal{F}_{\xi'}^{-1}$ and $\widehat{\cdot}$ are defined as (2.2).

Proof. Let \mathbf{F}_\pm , $F_{\pm,\text{div}}^\varepsilon$, and $\mathbf{F}_\pm^\varepsilon$ be given in (3.11), (3.14), and (3.16), respectively. Set

$$U_{\pm,0}^\varepsilon = -\mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{i\xi \cdot \mathcal{F}[\mathbf{F}_\pm](\xi)}{|\xi|^2} \right] (x) \quad (\varepsilon > 0).$$

We then have

$$\partial_j U_{\pm,0}^\varepsilon(x) = \mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{\xi_j \xi \cdot \mathcal{F}[\mathbf{F}_\pm](\xi)}{|\xi|^2} \right] (x) \quad (j = 1, \dots, N),$$

and also by (3.13)

$$\partial_k \partial_l U_{\pm,0}^\varepsilon(x) = \mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{\xi_k \xi_l}{|\xi|^2} \mathcal{F}[\text{div } \mathbf{F}_\pm](\xi) \right] (x) \quad (k, l = 1, \dots, N).$$

By the L_q boundedness of the Riesz operators,

$$\begin{aligned} \|\nabla U_{\pm,0}^\varepsilon\|_{L_q(\mathbf{R}^N)} &\leq C \|\mathbf{F}_\pm^\varepsilon\|_{L_q(\mathbf{R}^N)}, \quad \|\nabla^2 U_{\pm,0}^\varepsilon\|_{L_q(\mathbf{R}^N)} \leq C \|F_{\pm,\text{div}}^\varepsilon\|_{L_q(\mathbf{R}^N)}, \\ \|\nabla(U_{\pm,0}^\varepsilon - U_{\pm,0}^{\varepsilon'})\|_{L_q(\mathbf{R}^N)} &\leq C \|\mathbf{F}_\pm^\varepsilon - \mathbf{F}_\pm^{\varepsilon'}\|_{L_q(\mathbf{R}^N)}, \\ \|\nabla^2(U_{\pm,0}^\varepsilon - U_{\pm,0}^{\varepsilon'})\|_{L_q(\mathbf{R}^N)} &\leq C \|F_{\pm,\text{div}}^\varepsilon - F_{\pm,\text{div}}^{\varepsilon'}\|_{L_q(\mathbf{R}^N)}, \end{aligned}$$

where $\varepsilon, \varepsilon' > 0$ and C is a positive constant independent of ε and ε' . Thus, similarly to the proof of Lemma 3.3, we can construct $U_{\pm,0} \in \widehat{H}_q^1(\mathbf{R}^N) \cap \widehat{H}_q^2(\mathbf{R}^N)$ satisfying $\Delta U_{\pm,0} = \text{div } \mathbf{f}_\pm$ in \mathbf{R}_\pm^N and the estimates:

$$\|\nabla U_{\pm,0}\|_{L_q(\mathbf{R}^N)} \leq C \|\mathbf{f}_\pm\|_{L_q(\mathbf{R}_\pm^N)}, \quad \|\nabla^2 U_{\pm,0}\|_{L_q(\mathbf{R}^N)} \leq C \|\text{div } \mathbf{f}_\pm\|_{L_q(\mathbf{R}_\pm^N)}.$$

We also obtain the formulas of $\partial_N U_{\pm,0}$, stated in Lemma 3.4 (2), from (2.14) and

$$\partial_N U_{\pm,0}^\varepsilon = -\mathcal{F}_\xi^{-1} \left[e^{-\varepsilon|\xi|^2} \frac{i\xi_N}{|\xi|^2} \mathcal{F}[E_{\pm}^q z_\pm](\xi) \right] (x)$$

in the same manner that we have obtained the formulas of $\partial_N U_{\pm,\lambda}$, stated in Lemma 3.3 (2), from (2.13) and (3.19). This completes the proof of Lemma 3.4. \square

3.2. Proof of Theorem 3.1. We prove Theorem 3.1 in this subsection. In view of Lemma 2.1, it suffices to consider

$$\mathbf{f}_\pm = (f_{\pm 1}, \dots, f_{\pm N})^\top \in C_0^\infty(\mathbf{R}^N)^N, \quad g_\pm \in C_0^\infty(\mathbf{R}_\pm^N), \quad h_\pm \in C_0^\infty(\mathbf{R}^N).$$

Let $U_{\pm,\lambda}$ be solutions of (3.10) constructed in Lemma 3.3 for $\lambda \in \Sigma_{\sigma,0}$. Setting $w_\pm = U_{\pm,\lambda} + w_\pm$ in (3.1) yields

$$(3.26) \quad \begin{cases} \rho_\pm \lambda w_\pm - \Delta w_\pm = 0 & \text{in } \mathbf{R}_\pm^N, \\ \rho_+ w_+ - \rho_- w_- = g_1 & \text{on } \mathbf{R}_0^N, \\ \partial_N w_+ - \partial_N w_- = g_2 & \text{on } \mathbf{R}_0^N. \end{cases}$$

where

$$\begin{aligned} g_1 &= -(\rho_+ U_{+, \lambda} - \rho_- U_{-, \lambda}), \\ g_2 &= -(\partial_N U_{+, \lambda} - \partial_N U_{-, \lambda}) + f_{+N} - f_{-N} + h_+ - h_-. \end{aligned}$$

We apply the partial Fourier transform, given in (2.2), to (3.26) in order to obtain

$$\begin{cases} \rho_{\pm} \lambda \widehat{w}_{\pm}(x_N) - (\partial_N^2 - |\xi'|^2) \widehat{w}_{\pm}(x_N) = 0, & \pm x_N > 0, \\ \rho_+ \widehat{w}_+(0) - \rho_- \widehat{w}_-(0) = \widehat{g}_1(0), \\ \partial_N \widehat{w}_+(0) - \partial_N \widehat{w}_-(0) = \widehat{g}_2(0). \end{cases}$$

Solving these ordinary differential equations with respect to x_N , we have

$$\widehat{w}_{\pm}(x_N) = \left(\pm \frac{A_{\mp}}{\rho_+ A_- + \rho_- A_+} \widehat{g}_1(0) - \frac{\rho_{\mp}}{\rho_+ A_- + \rho_- A_+} \widehat{g}_2(0) \right) e^{-A_{\pm}(\pm x_N)}.$$

Thus, setting $w_{\pm} = \mathcal{F}_{\xi'}^{-1}[\widehat{w}_{\pm}(x_N)](x')$, we see that w_{\pm} are solutions to (3.26).

From now on, we estimate w_{\pm} . To this end, we decompose w_{\pm} as follows:

$$(3.27) \quad w_{\pm} = \pm w_{\pm}^1 - \rho_{\mp} (w_{\pm}^2 - w_{\pm}^3 + w_{\pm}^4),$$

where

$$\begin{aligned} w_{\pm}^1 &= \mathcal{F}_{\xi'}^{-1} \left[\frac{A_{\mp}}{\rho_+ A_- + \rho_- A_+} e^{-A_{\pm}(\pm x_N)} \widehat{g}_1(0) \right] (x'), \\ w_{\pm}^2 &= \mathcal{F}_{\xi'}^{-1} \left[\frac{1}{\rho_+ A_- + \rho_- A_+} e^{-A_{\pm}(\pm x_N)} \widehat{h}_+(0) \right] (x'), \\ w_{\pm}^3 &= \mathcal{F}_{\xi'}^{-1} \left[\frac{1}{\rho_+ A_- + \rho_- A_+} e^{-A_{\pm}(\pm x_N)} \widehat{h}_-(0) \right] (x'), \\ w_{\pm}^4 &= \mathcal{F}_{\xi'}^{-1} \left[\frac{1}{\rho_+ A_- + \rho_- A_+} e^{-A_{\pm}(\pm x_N)} \widehat{g}_3(0) \right] (x'), \end{aligned}$$

with

$$g_3 = g_2 - (h_+ - h_-) = -(\partial_N U_{+, \lambda} - \partial_N U_{-, \lambda}) + f_{+N} - f_{-N}.$$

By Lemma 2.2 and Leibniz's formula, we have for $A \in \{A_+, A_-, i\xi_1, \dots, i\xi_{N-1}\}$

$$(3.28) \quad \left| \partial_{\xi'}^{\alpha'} \left(\frac{A}{\rho_+ A_- + \rho_- A_+} \right) \right| \leq C (|\lambda|^{1/2} + |\xi'|)^{-|\alpha'|},$$

where $\xi' = (\xi_1, \dots, \xi_{N-1}) \in \mathbf{R}^{N-1} \setminus \{0\}$, $\lambda \in \Sigma_{\sigma, 0}$, and $C = C(N, \alpha', \sigma) > 0$. Thus, applying Lemma 2.5 to the above formula of w_{\pm}^1 yields

$$\begin{aligned} (3.29) \quad & \|(\lambda w_{\pm}^1, \lambda^{1/2} \nabla w_{\pm}^1, \nabla^2 w_{\pm}^1)\|_{L_q(\mathbf{R}_{\pm}^N)} \\ & \leq C \sum_{s \in \{+, -\}} \|(\lambda U_{s, \lambda}, \lambda^{1/2} \nabla U_{s, \lambda}, \nabla^2 U_{s, \lambda})\|_{L_q(\mathbf{R}_s^N)}, \\ & \|(\lambda^{1/2} w_{\pm}^1, \nabla w_{\pm}^1)\|_{L_q(\mathbf{R}_{\pm}^N)} \leq C \sum_{s \in \{+, -\}} \|(\lambda^{1/2} U_{s, \lambda}, \nabla U_{s, \lambda})\|_{L_q(\mathbf{R}_{\pm}^N)}. \end{aligned}$$

Analogously, we have from Lemmas 2.2 and 2.5

$$\begin{aligned} (3.30) \quad & \|(\lambda w_{\pm}^2, \lambda^{1/2} \nabla w_{\pm}^2, \nabla^2 w_{\pm}^2)\|_{L_q(\mathbf{R}_{\pm}^N)} \leq C \|(\lambda^{1/2} h_+, \nabla h_+)\|_{L_q(\mathbf{R}_{\pm}^N)}, \\ & \|(\lambda^{1/2} w_{\pm}^2, \nabla w_{\pm}^2)\|_{L_q(\mathbf{R}_{\pm}^N)} \leq C \left(\|h_+\|_{L_q(\mathbf{R}_{\pm}^N)} + |\lambda|^{-1/2} \|\nabla h_+\|_{L_q(\mathbf{R}_{\pm}^N)} \right), \end{aligned}$$

and also

$$(3.31) \quad \begin{aligned} \|(\lambda w_{\pm}^3, \lambda^{1/2} \nabla w_{\pm}^3, \nabla^2 w_{\pm}^3)\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C \|(\lambda^{1/2} h_{-}, \nabla h_{-})\|_{L_q(\mathbf{R}_{-}^N)}, \\ \|(\lambda^{1/2} w_{\pm}^3, \nabla w_{\pm}^3)\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C \left(\|h_{-}\|_{L_q(\mathbf{R}_{-}^N)} + |\lambda|^{-1/2} \|\nabla h_{-}\|_{L_q(\mathbf{R}_{-}^N)} \right). \end{aligned}$$

Furthermore, Lemmas 2.2 and 2.5 yield

$$(3.32) \quad \begin{aligned} &\|(\lambda w_{\pm}^4, \lambda^{1/2} \nabla w_{\pm}^4, \nabla^2 w_{\pm}^4)\|_{L_q(\mathbf{R}_{\pm}^N)} \\ &\leq C \sum_{s \in \{+, -\}} \|(\lambda^{1/2} \partial_N U_{s, \lambda}, \nabla \partial_N U_{s, \lambda}, \lambda^{1/2} f_{sN}, \nabla f_{sN})\|_{L_q(\mathbf{R}_s^N)}. \end{aligned}$$

On the other hand, if $f_{+N} - f_{-N} = 0$ on \mathbf{R}_{-}^N , then we have

$$(3.33) \quad \begin{aligned} &\|(\lambda w_{\pm}^4, \lambda^{1/2} \nabla w_{\pm}^4, \nabla^2 w_{\pm}^4)\|_{L_q(\mathbf{R}_{\pm}^N)} \\ &\leq C \sum_{s \in \{+, -\}} \|(\lambda^{1/2} \partial_N U_{s, \lambda}, \nabla \partial_N U_{s, \lambda})\|_{L_q(\mathbf{R}_s^N)}. \end{aligned}$$

Next, we estimate $\|(\lambda^{1/2} w_{\pm}^4, \nabla w_{\pm}^4)\|_{L_q(\mathbf{R}_{\pm}^N)}$. Note that by Lemma 3.3 (2)

$$\begin{aligned} \widehat{g}_3(0) &= \sum_{s \in \{+, -\}} \sum_{j=1}^{N-1} \int_0^{\infty} i \xi_j e^{-A_s y_N} \widehat{f}_{sj}(\xi', \mathbf{s}y_N) dy_N \\ &\quad + \sum_{s \in \{+, -\}} \mathbf{s} \int_0^{\infty} A_s e^{-A_s y_N} \widehat{f}_{sN}(\xi', \mathbf{s}y_N) dy_N. \end{aligned}$$

We thus obtain

$$\begin{aligned} w_{\pm}^4 &= \sum_{s \in \{+, -\}} \sum_{j=1}^{N-1} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{i \xi_j e^{-(A_{\pm}(\pm x_N) + A_s y_N)}}{\rho_+ A_- + \rho_- A_+} \widehat{f}_{sj}(\xi', \mathbf{s}y_N) \right] (x') dy_N \\ &\quad + \sum_{s \in \{+, -\}} \mathbf{s} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{A_s e^{-(A_{\pm}(\pm x_N) + A_s y_N)}}{\rho_+ A_- + \rho_- A_+} \widehat{f}_{sN}(\xi', \mathbf{s}y_N) \right] (x') dy_N. \end{aligned}$$

Combining these formulas with Lemma 2.3, (2.12), and (3.28) furnishes

$$\|(\lambda^{1/2} w_{\pm}^4, \nabla w_{\pm}^4)\|_{L_q(\mathbf{R}_{\pm}^N)} \leq C \left(\|\mathbf{f}_+\|_{L_q(\mathbf{R}_+^N)} + \|\mathbf{f}_-\|_{L_q(\mathbf{R}_-^N)} \right).$$

Recalling $v_{\pm} = U_{\pm, \lambda} + w_{\pm}$ and (3.27), we obtain solutions v_{\pm} of (3.1) satisfying (3.3)-(3.5) by the last estimate, (3.29)-(3.33), and Lemma 3.3.

Finally, we prove the uniqueness of solutions to (3.1). Assume that $v_{\pm} \in H_q^2(\mathbf{R}_{\pm}^N)$ satisfy the homogeneous equations, i.e.

$$\begin{cases} \rho_{\pm} \lambda v_{\pm} - \Delta v_{\pm} = 0 & \text{in } \mathbf{R}_{\pm}^N, \\ \rho_+ v_+ = \rho_- v_- & \text{on } \mathbf{R}_0^N, \\ \partial_N v_+ = \partial_N v_- & \text{on } \mathbf{R}_0^N. \end{cases}$$

Let $\mathbf{f}_{\pm} \in C_0^{\infty}(\mathbf{R}_{\pm}^N)^N$. Since we know the existence of solutions to (3.1) for these \mathbf{f}_{\pm} as already discussed above, we have $w_{\pm} \in H_{q'}^2(\mathbf{R}_{\pm}^N)$, $q' = q/(q-1)$, satisfying

$$\begin{cases} \rho_{\pm} \lambda w_{\pm} - \Delta w_{\pm} = -\operatorname{div} \mathbf{f}_{\pm} & \text{in } \mathbf{R}_{\pm}^N, \\ \rho_+ w_+ = \rho_- w_- & \text{on } \mathbf{R}_0^N, \\ \partial_N w_+ = \partial_N w_- & \text{on } \mathbf{R}_0^N. \end{cases}$$

It then holds by integration by parts that

$$\begin{aligned} & (\rho_+ \nabla v_+, \mathbf{f}_+)_{\mathbf{R}_+^N} + (\rho_- \nabla v_-, \mathbf{f}_-)_{\mathbf{R}_-^N} = -(\rho_+ v_+, \operatorname{div} \mathbf{f}_+)_{\mathbf{R}_+^N} - (\rho_- v_-, \operatorname{div} \mathbf{f}_-)_{\mathbf{R}_-^N} \\ & = (\rho_+ v_+, \rho_+ \lambda w_+ - \Delta w_+)_{\mathbf{R}_+^N} + (\rho_- v_-, \rho_- \lambda w_- - \Delta w_-)_{\mathbf{R}_-^N} \\ & = (\rho_+ \lambda v_+ - \Delta v_+, \rho_+ w_+)_{\mathbf{R}_+^N} + (\rho_- \lambda v_- - \Delta v_-, \rho_- w_-)_{\mathbf{R}_-^N} = 0, \end{aligned}$$

and thus $\rho_\pm v_\pm = c_\pm$ for some constants c_\pm . By the equation: $\rho_\pm \lambda v_\pm - \Delta v_\pm = 0$ in \mathbf{R}_\pm^N , we conclude $c_\pm = 0$ since $\lambda \neq 0$. This implies the uniqueness of solutions to (3.1), which completes the proof of Theorem 3.1.

3.3. Proof of Theorem 3.2. We prove Theorem 3.2 in this subsection. In view of Lemma 2.1, it suffices to consider

$$\mathbf{f}_\pm = (f_{\pm 1}, \dots, f_{\pm N})^\top \in C_0^\infty(\mathbf{R}^N)^N.$$

Let $U_{\pm,0}$ be the solutions of (3.25) constructed in Lemma 3.4, and set $v_\pm = U_{\pm,0} + w_\pm$ in (3.2). Then,

$$(3.34) \quad \begin{cases} \Delta w_\pm = 0 & \text{in } \mathbf{R}_\pm^N, \\ \rho_+ w_+ - \rho_- w_- = g_1 & \text{on } \mathbf{R}_0^N, \\ \partial_N w_+ - \partial_N w_- = g_2 & \text{on } \mathbf{R}_0^N, \end{cases}$$

where

$$g_1 = -(\rho_+ U_{+,0} - \rho_- U_{-,0}), \quad g_2 = -(\partial_N U_{+,0} - \partial_N U_{-,0}) + f_{+N} - f_{-N}.$$

We apply the partial Fourier transform, given in (2.2), to (3.34) in order to obtain

$$\begin{cases} (\partial_N^2 - |\xi'|^2) \widehat{w}_\pm(x_N) = 0, & \pm x_N > 0, \\ \rho_+ \widehat{w}_+(0) - \rho_- \widehat{w}_-(0) = \widehat{g}_1(0), \\ \partial_N \widehat{w}_+(0) - \partial_N \widehat{w}_-(0) = \widehat{g}_2(0). \end{cases}$$

Solving these ordinary differential equations with respect to x_N , we have

$$\widehat{w}_\pm(x_N) = \left(\frac{1}{\rho_+ + \rho_-} \right) \left(\pm \widehat{g}_1(0) - \rho_\mp \frac{\widehat{g}_2(0)}{|\xi'|} \right) e^{\mp |\xi'| x_N}.$$

Thus, setting $w_\pm = \mathcal{F}_{\xi'}^{-1}[\widehat{w}_\pm(x_N)](x')$, we see that w_\pm are solutions to (3.34).

From now on, we estimate w_\pm . To this end, we decompose w_\pm as follows:

$$(3.35) \quad w_\pm = \pm \left(\frac{1}{\rho_+ + \rho_-} \right) w_\pm^1 - \left(\frac{\rho_\mp}{\rho_+ + \rho_-} \right) w_\pm^2,$$

where

$$w_\pm^1 = \mathcal{F}_{\xi'}^{-1} \left[\widehat{g}_1(0) e^{\mp |\xi'| x_N} \right] (x'), \quad w_\pm^2 = \mathcal{F}_{\xi'}^{-1} \left[\frac{\widehat{g}_2(0)}{|\xi'|} e^{\mp |\xi'| x_N} \right] (x').$$

Let us consider $\|\nabla w_\pm^1\|_{L_q(\mathbf{R}_\pm^N)}$ and $\|\nabla^2 w_\pm^1\|_{L_q(\mathbf{R}_\pm^N)}$. Similarly to (2.11), one has

$$\begin{aligned} \widehat{g}_1(0) e^{\mp |\xi'| x_N} &= \int_0^\infty |\xi'| e^{-|\xi'|(\pm x_N + y_N)} \widehat{g}_1(\xi', \pm y_N) dy_N \\ &\mp \int_0^\infty e^{-|\xi'|(\pm x_N + y_N)} \widehat{\partial_N g_1}(\xi', \pm y_N) dy_N, \end{aligned}$$

which, combined with $|\xi'| = -\sum_{j=1}^{N-1} (i\xi_j)^2/|\xi'|$, furnishes

$$\begin{aligned} \widehat{g}_1(0)e^{\mp|\xi'|x_N} &= -\sum_{j=1}^{N-1} \int_0^\infty \frac{i\xi_j}{|\xi'|} e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_j g_1}(\xi', \pm y_N) dy_N \\ &\mp \int_0^\infty e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_N g_1}(\xi', \pm y_N) dy_N. \end{aligned}$$

Inserting these relations into w_\pm^1 yields

$$\begin{aligned} w_\pm^1 &= -\sum_{j=1}^{N-1} \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\frac{i\xi_j}{|\xi'|} e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_j g_1}(\xi', \pm y_N) \right] (x') dy_N \\ &\mp \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_N g_1}(\xi', \pm y_N) \right] (x') dy_N. \end{aligned}$$

Thus, for $k, l = 1, \dots, N-1$,

$$\begin{aligned} \partial_k w_\pm^1 &= \sum_{j=1}^{N-1} \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\frac{\xi_k \xi_j}{|\xi'|^2} |\xi'| e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_j g_1}(\xi', \pm y_N) \right] (x') dy_N \\ &\mp \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\frac{i\xi_k}{|\xi'|} |\xi'| e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_N g_1}(\xi', \pm y_N) \right] (x') dy_N, \\ \partial_N w_\pm^1 &= \pm \sum_{j=1}^{N-1} \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\frac{i\xi_j}{|\xi'|} |\xi'| e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_j g_1}(\xi', \pm y_N) \right] (x') dy_N \\ &+ \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[|\xi'| e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_N g_1}(\xi', \pm y_N) \right] (x') dy_N, \\ \partial_l \partial_k w_\pm^1 &= \sum_{j=1}^{N-1} \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\frac{\xi_k \xi_j}{|\xi'|^2} |\xi'| e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_l \partial_j g_1}(\xi', \pm y_N) \right] (x') dy_N \\ &\mp \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\frac{i\xi_k}{|\xi'|} |\xi'| e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_l \partial_N g_1}(\xi', \pm y_N) \right] (x') dy_N, \\ \partial_k \partial_N w_\pm^1 &= \partial_N \partial_k w_\pm^1 \\ &= \pm \sum_{j=1}^{N-1} \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\frac{i\xi_j}{|\xi'|} |\xi'| e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_k \partial_j g_1}(\xi', \pm y_N) \right] (x') dy_N \\ &+ \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[|\xi'| e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_k \partial_N g_1}(\xi', \pm y_N) \right] (x') dy_N. \end{aligned}$$

Analogously, for $\Delta' g_1 = \sum_{j=1}^{N-1} \partial_j^2 g_1$,

$$\begin{aligned} \partial_N^2 w_\pm^1 &= -\int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[|\xi'| e^{-|\xi'|(\pm x_N+y_N)} \widehat{\Delta' g_1}(\xi', \pm y_N) \right] (x') dy_N \\ &\pm \sum_{j=1}^{N-1} \int_0^\infty \mathcal{F}_{\xi'}^{-1} \left[\frac{i\xi_j}{|\xi'|} |\xi'| e^{-|\xi'|(\pm x_N+y_N)} \widehat{\partial_j \partial_N g_1}(\xi', \pm y_N) \right] (x') dy_N. \end{aligned}$$

It then follows from Lemmas 2.2 and 2.4 that

$$(3.36) \quad \|\nabla w_\pm^1\|_{L_q(\mathbf{R}_\pm^N)} \leq C \sum_{\mathfrak{s} \in \{+, -\}} \|\nabla U_{\mathfrak{s}, 0}\|_{L_q(\mathbf{R}_\mathfrak{s}^N)},$$

$$\|\nabla^2 w_{\pm}^1\|_{L_q(\mathbf{R}_{\pm}^N)} \leq C \sum_{\mathfrak{s} \in \{+, -\}} \|\nabla^2 U_{\mathfrak{s}, 0}\|_{L_q(\mathbf{R}_{\pm}^N)}.$$

We similarly see that

$$(3.37) \quad \|\nabla^2 w_{\pm}^2\|_{L_q(\mathbf{R}_{\pm}^N)} \leq C \sum_{\mathfrak{s} \in \{+, -\}} (\|\nabla \partial_N U_{\mathfrak{s}, 0}\|_{L_q(\mathbf{R}_{\pm}^N)} + \|\nabla f_{\mathfrak{s}N}\|_{L_q(\mathbf{R}_{\pm}^N)})$$

and that, if $f_{+N} - f_{-N} = 0$ on \mathbf{R}_0^N ,

$$(3.38) \quad \|\nabla^2 w_{\pm}^2\|_{L_q(\mathbf{R}_{\pm}^N)} \leq C \sum_{\mathfrak{s} \in \{+, -\}} \|\nabla \partial_N U_{\mathfrak{s}, 0}\|_{L_q(\mathbf{R}_{\pm}^N)}.$$

Next, we estimate $\|\nabla w_{\pm}^2\|_{L_q(\mathbf{R}_{\pm}^N)}$. By Lemma 3.4,

$$\widehat{g}_2(0) = \sum_{\mathfrak{s} \in \{+, -\}} \int_0^{\infty} e^{-|\xi'| y_N} \left(i\xi' \cdot \widehat{\mathbf{f}}_{\mathfrak{s}}'(\xi', \mathfrak{s}y_N) + \mathfrak{s}|\xi'| \widehat{f}_{\mathfrak{s}N}(\xi', \mathfrak{s}y_N) \right) dy_N,$$

which implies that

$$\begin{aligned} w_{\pm}^2 &= \sum_{\mathfrak{s} \in \{+, -\}} \left\{ \sum_{j=1}^{N-1} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{i\xi_j}{|\xi'|} e^{-|\xi'|(\pm x_N + y_N)} \widehat{f}_{\mathfrak{s}j}(\xi', \mathfrak{s}y_N) \right] (x') dy_N \right. \\ &\quad \left. + \mathfrak{s} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[e^{-|\xi'|(\pm x_N + y_N)} \widehat{f}_{\mathfrak{s}N}(\xi', \mathfrak{s}y_N) \right] (x') dy_N \right\}. \end{aligned}$$

Then, for $k = 1, \dots, N-1$,

$$\begin{aligned} \partial_k w_{\pm}^2 &= \sum_{\mathfrak{s} \in \{+, -\}} \left\{ - \sum_{j=1}^{N-1} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{\xi_k \xi_j}{|\xi'|^2} |\xi'| e^{-|\xi'|(\pm x_N + y_N)} \widehat{f}_{\mathfrak{s}j}(\xi', \mathfrak{s}y_N) \right] (x') dy_N \right. \\ &\quad \left. + \mathfrak{s} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{i\xi_k}{|\xi'|} |\xi'| e^{-|\xi'|(\pm x_N + y_N)} \widehat{f}_{\mathfrak{s}N}(\xi', \mathfrak{s}y_N) \right] (x') dy_N \right\}, \\ \partial_N w_{\pm}^2 &= \sum_{\mathfrak{s} \in \{+, -\}} \left\{ \mp \sum_{j=1}^{N-1} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[\frac{i\xi_j}{|\xi'|} |\xi'| e^{-|\xi'|(\pm x_N + y_N)} \widehat{f}_{\mathfrak{s}j}(\xi', \mathfrak{s}y_N) \right] (x') dy_N \right. \\ &\quad \left. \mp \mathfrak{s} \int_0^{\infty} \mathcal{F}_{\xi'}^{-1} \left[|\xi'| e^{-|\xi'|(\pm x_N + y_N)} \widehat{f}_{\mathfrak{s}N}(\xi', \mathfrak{s}y_N) \right] (x') dy_N \right\}. \end{aligned}$$

Combining these formulas with Lemmas 2.2 and 2.4 furnishes

$$\|\nabla w_{\pm}^2\|_{L_q(\mathbf{R}_{\pm}^N)} \leq C \left(\|\mathbf{f}_+\|_{L_q(\mathbf{R}_+^N)} + \|\mathbf{f}_-\|_{L_q(\mathbf{R}_-^N)} \right).$$

Recalling $v_{\pm} = U_{\pm, 0} + w_{\pm}$ and (3.35), we obtain solutions v_{\pm} of (3.2) satisfying (3.6)-(3.8) by the last estimate, (3.36)-(3.38), and Lemma 3.4.

Finally, we prove the uniqueness of solutions to (3.2). Let v_{\pm} satisfy (3.9) and $\mathbf{f}_{\pm} \in C_0^{\infty}(\mathbf{R}_{\pm}^N)^N$. By the discussion above, there exists $w_{\pm} \in \widehat{H}_{q'}^1(\mathbf{R}_{\pm}^N) \cap \widehat{H}_{q'}^2(\mathbf{R}_{\pm}^N)$ with $q' = q/(q-1)$ satisfying

$$\begin{cases} \Delta w_{\pm} = \operatorname{div} \mathbf{f}_{\pm} & \text{in } \mathbf{R}_{\pm}^N, \\ \rho_+ w_+ = \rho_- w_- & \text{on } \mathbf{R}_0^N, \\ \partial_N w_+ = \partial_N w_- & \text{on } \mathbf{R}_0^N. \end{cases}$$

At this point, we introduce the following lemma (cf. e.g. [11, Section II.6], [22], [23, Proof of Theorem A.4]).

Lemma 3.5. *Let $q \in (1, \infty)$ and set*

$$d_q(x) = \begin{cases} (1 + |x|^2)^{1/2} & \text{when } q \neq N, \\ (1 + |x|^2)^{1/2} \log(2 + |x|^2)^{1/2} & \text{when } q = N. \end{cases}$$

Then, for any $u \in \widehat{H}_q^1(\mathbf{R}^N)$, there exists a constant c_u such that

$$\left\| \frac{u - c_u}{d_q} \right\|_{L_q(\mathbf{R}^N)} \leq C \|\nabla u\|_{L_q(\mathbf{R}^N)},$$

where $C = C(N, q)$ is a positive constant independent of u and c_u .

Now we set

$$\mathbf{f} = \mathbf{f}_+ \mathbf{1}_{\mathbf{R}_+^N} + \mathbf{f}_- \mathbf{1}_{\mathbf{R}_-^N}, \quad z = \rho_+ z_+ \mathbf{1}_{\mathbf{R}_+^N} + \rho_- z_- \mathbf{1}_{\mathbf{R}_-^N} \text{ for } z \in \{v, w\}.$$

Then $v \in \widehat{H}_q^1(\mathbf{R}^N)$ with $\nabla v = \rho_+(\nabla v_+) \mathbf{1}_{\mathbf{R}_+^N} + \rho_-(\nabla v_-) \mathbf{1}_{\mathbf{R}_-^N}$ by $\rho_+ v_+ = \rho_- v_-$ on \mathbf{R}_0^N , while $w \in \widehat{H}_{q'}^1(\mathbf{R}^N)$ with $\nabla w = \rho_+(\nabla w_+) \mathbf{1}_{\mathbf{R}_+^N} + \rho_-(\nabla w_-) \mathbf{1}_{\mathbf{R}_-^N}$ by $\rho_+ w_+ = \rho_- w_-$ on \mathbf{R}_0^N . By Lemma 3.5, there exist constants c_v and c_w such that

$$\left\| \frac{v - c_v}{d_q} \right\|_{L_q(\mathbf{R}^N)} \leq C \|\nabla v\|_{L_q(\mathbf{R}^N)}, \quad \left\| \frac{w - c_w}{d_{q'}} \right\|_{L_{q'}(\mathbf{R}^N)} \leq C \|\nabla w\|_{L_{q'}(\mathbf{R}^N)}.$$

Let ψ_R be a cut-off function of Sobolev's type as follows³: For $R > 0$ large enough and for $\psi \in C_0^\infty(\mathbf{R})$ satisfying $\psi(t) = 1$ when $|t| \leq 1/2$ and $\psi(t) = 0$ when $|t| \geq 1$,

$$\psi_R(x) = \begin{cases} 1 & \text{when } |x| \leq e^{\sqrt[3]{\log R}}, \\ \psi\left(\frac{\log \log |x|}{\log \log R}\right) & \text{when } |x| \geq e^{\sqrt[3]{\log R}}. \end{cases}$$

Note that for $D_R = \{x \in \mathbf{R}^N : e^{\sqrt{\log R}} \leq |x| \leq R\}$

$$(3.39) \quad |\nabla \psi_R(x)| \leq \frac{C}{\log \log R} \frac{1}{|x| \log |x|}, \quad \text{supp } \nabla \psi_R \subset D_R,$$

where C is a positive constant independent of R .

Choose $R > 0$ large enough so that $\psi_R = 1$ on $\text{supp } \mathbf{f}$. It then holds that

$$\begin{aligned} (\nabla v, \mathbf{f})_{\mathbf{R}^N} &= (\nabla(v - c_v), \mathbf{f})_{\mathbf{R}^N} = -(v - c_v, \text{div } \mathbf{f})_{\mathbf{R}^N} = -(v - c_v, \psi_R \text{div } \mathbf{f})_{\mathbf{R}^N} \\ &= -(v - c_v, \psi_R \Delta w_+)_{\mathbf{R}_+^N} - (v - c_v, \psi_R \Delta w_-)_{\mathbf{R}_-^N} \\ &= -(v - c_v, \text{div}(\psi_R \nabla w_+) - \nabla \psi_R \cdot \nabla w_+)_{\mathbf{R}_+^N} \\ &\quad - (v - c_v, \text{div}(\psi_R \nabla w_-) - \nabla \psi_R \cdot \nabla w_-)_{\mathbf{R}_-^N} \\ &= (\rho_+ \nabla v_+, \psi_R \nabla w_+)_{\mathbf{R}_+^N} + (v - c_v, \nabla \psi_R \cdot \nabla w_+)_{\mathbf{R}_+^N} \\ &\quad + (\rho_- \nabla v_-, \psi_R \nabla w_-)_{\mathbf{R}_-^N} + (v - c_v, \nabla \psi_R \cdot \nabla w_-)_{\mathbf{R}_-^N}. \end{aligned}$$

On the other hand,

$$\begin{aligned} &(\rho_+ \nabla v_+, \psi_R \nabla w_+)_{\mathbf{R}_+^N} + (\rho_- \nabla v_-, \psi_R \nabla w_-)_{\mathbf{R}_-^N} \\ &= (\psi_R \nabla v_+, \nabla(\rho_+ w_+ - c_w))_{\mathbf{R}_+^N} + (\psi_R \nabla v_-, \nabla(\rho_- w_- - c_w))_{\mathbf{R}_-^N} \\ &= -(\text{div}(\psi_R \nabla v_+), \rho_+ w_+ - c_w)_{\mathbf{R}_+^N} - (\text{div}(\psi_R \nabla v_-), \rho_- w_- - c_w)_{\mathbf{R}_-^N} \end{aligned}$$

³ See e.g. [11, Proof of Theorem II.7.1].

$$= -(\nabla\psi_R \cdot \nabla v_+, w - c_w)_{\mathbf{R}_+^N} - (\nabla\psi_R \cdot \nabla v_-, w - c_w)_{\mathbf{R}_-^N},$$

and thus

$$(3.40) \quad (\nabla v, \mathbf{f})_{\mathbf{R}^N} = (v - c_v, \nabla\psi_R \cdot \nabla w_+)_{\mathbf{R}_+^N} + (v - c_v, \nabla\psi_R \cdot \nabla w_-)_{\mathbf{R}_-^N} \\ - (\nabla\psi_R \cdot \nabla v_+, w - c_w)_{\mathbf{R}_+^N} - (\nabla\psi_R \cdot \nabla v_-, w - c_w)_{\mathbf{R}_-^N}.$$

We here see that by Lemma 3.5 and (3.39)

$$\begin{aligned} & |(v - c_v, \nabla\psi_R \cdot \nabla w_{\pm})_{\mathbf{R}_{\pm}^N}| \\ & \leq \left\| \frac{v - c_v}{d_q} \right\|_{L_q(\mathbf{R}^N)} \|d_q \nabla\psi_R\|_{L_{\infty}(\mathbf{R}^N)} \|\nabla w_{\pm}\|_{L_{q'}(\mathbf{R}_{\pm}^N)} \\ & \leq \frac{C}{\log \log R} \|\nabla v\|_{L_q(\mathbf{R}^N)} \|\nabla w_{\pm}\|_{L_{q'}(\mathbf{R}_{\pm}^N)} \rightarrow 0 \quad \text{as } R \rightarrow \infty, \end{aligned}$$

and also

$$\lim_{R \rightarrow \infty} (\nabla\psi_R \cdot \nabla v_{\pm}, w - c_w)_{\mathbf{R}_{\pm}^N} = 0.$$

By these properties, letting $R \rightarrow \infty$ in (3.40) furnishes

$$(\rho_+ \nabla v_+, \mathbf{f}_+)_{\mathbf{R}_+^N} + (\rho_- \nabla v_-, \mathbf{f}_-)_{\mathbf{R}_-^N} = (\nabla v, \mathbf{f})_{\mathbf{R}^N} = 0,$$

which implies $\rho_{\pm} v_{\pm} = c_{\pm}$ for some constants c_{\pm} . Since $\rho_+ v_+ = \rho_- v_-$ on \mathbf{R}_0^N , we have $c_+ = c_-$. Thus setting $c = c_+ = c_-$ yields $v_{\pm} = \rho_{\pm}^{-1} c$, which implies the uniqueness of solutions to (3.2). This completes the proof of Theorem 3.2.

3.4. Weak elliptic problem with flat interface. For $\mathbf{f} \in L_q(\mathbf{R}^N \setminus \mathbf{R}_0^N)^N$ with $q \in (1, \infty)$, this subsection considers the unique solvability of the following weak elliptic problem: Find $u \in \widehat{H}_q^1(\mathbf{R}^N)$ such that

$$(3.41) \quad (\rho^{-1} \nabla u, \nabla \varphi)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} = (\mathbf{f}, \nabla \varphi)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} \quad \text{for any } \varphi \in \widehat{H}_{q'}^1(\mathbf{R}^N),$$

where $q' = q/(q-1)$ and $\rho = \rho_+ \mathbf{1}_{\mathbf{R}_+^N} + \rho_- \mathbf{1}_{\mathbf{R}_-^N}$ for positive constants ρ_{\pm} . More precisely, we prove

Theorem 3.6. *Let $q \in (1, \infty)$ and $q' = q/(q-1)$.*

(1) **Existence.** *Let $\mathbf{f} \in L_q(\mathbf{R}^N \setminus \mathbf{R}_0^N)^N$. Then the weak elliptic problem (3.41) admits a solution $u \in \widehat{H}_q^1(\mathbf{R}^N)$ satisfying*

$$(3.42) \quad \|\nabla u\|_{L_q(\mathbf{R}^N)} \leq 2C_2(\rho_+ + \rho_-) \|\mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \mathbf{R}_0^N)},$$

where C_2 is the same constant as in Theorem 3.2.

(2) **Uniqueness.** *If $u \in \widehat{H}_q^1(\mathbf{R}^N)$ satisfies*

$$(3.43) \quad (\rho^{-1} \nabla u, \nabla \varphi)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} = 0 \quad \text{for any } \varphi \in \widehat{H}_{q'}^1(\mathbf{R}^N),$$

then $u = c$ for some constant c .

Proof. (1). Since $C_0^{\infty}(\mathbf{R}^N)$ is dense in $L_q(\mathbf{R}^N \setminus \mathbf{R}_0^N)$, it suffices to consider $\mathbf{f} \in C_0^{\infty}(\mathbf{R}^N \setminus \mathbf{R}_0^N)^N$. By Theorem 3.2, we have $v_{\pm} \in \widehat{H}_q^1(\mathbf{R}_{\pm}^N) \cap \widehat{H}_q^2(\mathbf{R}_{\pm}^N)$ satisfying (3.2) and (3.7) with $\mathbf{f}_+ = \mathbf{f}$ and $\mathbf{f}_- = \mathbf{f}$, where we note that $f_{+N} - f_{-N} = 0$ on \mathbf{R}_0^N in this case. Setting $u_{\pm} = \rho_{\pm} v_{\pm}$ and $u = u_+ \mathbf{1}_{\mathbf{R}_+^N} + u_- \mathbf{1}_{\mathbf{R}_-^N}$ furnishes that

$$\begin{cases} \rho_{\pm}^{-1} \Delta u_{\pm} = \operatorname{div} \mathbf{f}_{\pm} & \text{in } \mathbf{R}_{\pm}^N, \\ u_+ = u_- & \text{on } \mathbf{R}_0^N, \\ \rho_+^{-1} \partial_N u_+ = \rho_-^{-1} \partial_N u_- & \text{on } \mathbf{R}_0^N, \end{cases}$$

and that $u \in \widehat{H}_q^1(\mathbf{R}^N)$ with $\nabla u = \rho_+(\nabla v_+) \mathbf{1}_{\mathbf{R}_+^N} + \rho_-(\nabla v_-) \mathbf{1}_{\mathbf{R}_-^N}$ by $\rho_+ v_+ = \rho_- v_-$ on \mathbf{R}_0^N , Combining the last property with (3.7) yields

$$\|\nabla u\|_{L_q(\mathbf{R}^N)} \leq C_2(\rho_+ + \rho_-) \sum_{s \in \{+, -\}} \|\mathbf{f}_s\|_{L_q(\mathbf{R}_s^N)} \leq 2C_2(\rho_+ + \rho_-) \|\mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \mathbf{R}_0^N)},$$

which implies that u satisfies (3.42).

Next, we prove that u is a solution to (3.41). Let $\varphi \in \widehat{H}_q^1(\mathbf{R}^N)$. Then there exists $\{\varphi_j\}_{j=1}^\infty \subset C_0^\infty(\mathbf{R}^N)$ such that $\lim_{j \rightarrow \infty} \|\nabla(\varphi_j - \varphi)\|_{L_{q'}(\mathbf{R}^N)} = 0$. Thus,

$$\begin{aligned} (\mathbf{f}, \nabla \varphi)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} &= \lim_{j \rightarrow \infty} (\mathbf{f}, \nabla \varphi_j)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} = - \lim_{j \rightarrow \infty} \left\{ (\operatorname{div} \mathbf{f}_+, \varphi_j)_{\mathbf{R}_+^N} + (\operatorname{div} \mathbf{f}_-, \varphi_j)_{\mathbf{R}_-^N} \right\} \\ &= - \lim_{j \rightarrow \infty} \left\{ (\rho_+^{-1} \Delta u_+, \varphi_j)_{\mathbf{R}_+^N} + (\rho_-^{-1} \Delta u_-, \varphi_j)_{\mathbf{R}_+^N} \right\} \\ &= \lim_{j \rightarrow \infty} \left\{ (\rho_+^{-1} \nabla u_+, \nabla \varphi_j)_{\mathbf{R}_+^N} + (\rho_-^{-1} \nabla u_-, \nabla \varphi_j)_{\mathbf{R}_+^N} \right\} \\ &= \lim_{j \rightarrow \infty} (\rho^{-1} \nabla u, \nabla \varphi_j)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} = (\rho^{-1} \nabla u, \nabla \varphi)_{\mathbf{R}^N \setminus \mathbf{R}_0^N}. \end{aligned}$$

This completes the proof of (1).

(2). Let $u \in \widehat{H}_q^1(\mathbf{R}^N)$ satisfy (3.43), and let $\mathbf{f} \in C_0^\infty(\mathbf{R}^N)^N$. By (1), we have $v \in \widehat{H}_{q'}^1(\mathbf{R}^N)$ satisfying

$$(\rho^{-1} \nabla v, \nabla \psi)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} = (\mathbf{f}, \nabla \psi)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} \quad \text{for any } \psi \in \widehat{H}_q^1(\mathbf{R}^N).$$

Choosing $\psi = u$ in this equality yields

$$(\mathbf{f}, \nabla u)_{\mathbf{R}^N} = (\mathbf{f}, \nabla u)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} = (\rho^{-1} \nabla v, \nabla u)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} = (\rho^{-1} \nabla u, \nabla v)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} = 0,$$

which implies $u = c$ for some constant c . This completes the proof of (2). \square

4. PROBLEMS IN THE WHOLE SPACE WITH BENT INTERFACE

Let $\Phi : \mathbf{R}_x^N \rightarrow \mathbf{R}_y^N$ be a bijection of class C^1 and Φ^{-1} the inverse mapping of Φ , where the subscripts x, y denote their variables. Let $(\nabla_x \Phi)(x) = \mathcal{A} + \mathbf{B}(x)$ and $(\nabla_y \Phi^{-1})(\Phi(x)) = \mathcal{A}_{-1} + \mathbf{B}_{-1}(x)$. Assume that $\mathcal{A}, \mathcal{A}_{-1}$ are orthonormal matrices with constant coefficients and $\det \mathcal{A} = \det \mathcal{A}_{-1} = 1$, and also assume that $\mathbf{B}(x), \mathbf{B}_{-1}(x)$ are matrix-valued functions of $H_r^1(\mathbf{R}^N)$, $r \in (N, \infty)$, satisfying

$$(4.1) \quad \begin{aligned} \|(\mathbf{B}, \mathbf{B}_{-1})\|_{L_\infty(\mathbf{R}^N)} &\leq M_1 \quad (0 < M_1 \leq 1/2), \\ \|(\nabla \mathbf{B}, \nabla \mathbf{B}_{-1})\|_{L_r(\mathbf{R}^N)} &\leq M_2 \quad (M_2 \geq 1). \end{aligned}$$

In what follows, we will choose M_1 small enough eventually.

Let $\widetilde{\Omega}_\pm = \Phi(\mathbf{R}_\pm^N)$, $\widetilde{\Sigma} = \Phi(\mathbf{R}_0^N)$, and $\widetilde{\mathbf{n}} = \widetilde{\mathbf{n}}(y)$ be the unit normal vector on $\widetilde{\Sigma}$ pointing from $\widetilde{\Omega}_+$ into $\widetilde{\Omega}_-$. Let \mathcal{A}_{ij} and $B_{ij}(x)$ be respectively the (i, j) -component of \mathcal{A}_{-1} and the (i, j) -component of $\mathbf{B}_{-1}(x)$. In addition, setting $\Phi^{-1} = (\Phi_{-1,1}, \dots, \Phi_{-1,N})$, we see that $\widetilde{\Sigma}$ is represented as $\Phi_{-1,N}(y) = 0$. This representation implies that

$$(4.2) \quad \widetilde{\mathbf{n}}(\Phi(x)) = - \frac{(\nabla_y \Phi_{-1,N})(\Phi(x))}{|(\nabla_y \Phi_{-1,N})(\Phi(x))|} = - \frac{(\mathcal{A}_{N1} + B_{N1}(x), \dots, \mathcal{A}_{NN} + B_{NN}(x))^T}{\sqrt{\sum_{j=1}^N (\mathcal{A}_{Nj} + B_{Nj}(x))^2}}$$

$$= \frac{(\mathcal{A}_{-1} + \mathbf{B}_{-1}(x))^{\top} \mathbf{n}_0}{|(\mathcal{A}_{-1} + \mathbf{B}_{-1}(x))^{\top} \mathbf{n}_0|}$$

for $\mathbf{n}_0 = (0, \dots, 0, -1)^{\top}$. Especially, $\tilde{\mathbf{n}}$ is defined on \mathbf{R}^N through the relation (4.2).

Remark 4.1. Let J be the Jacobian of Φ , i.e. $J = \det(\nabla\Phi)$, and let $d = |(\mathcal{A}_{-1} + \mathbf{B}_{-1}(x))^{\top} \mathbf{n}_0|$. Since M_1 is small enough, there are positive constants C_3 and C_4 , independent of M_1 , M_2 , and r , such that the following inequalities hold:

$$(4.3) \quad C_3 \leq J(x), d(x) \leq C_4 \quad (x \in \mathbf{R}^N), \quad \sup_{x \in \mathbf{R}^N} |1 - J(x)| \leq C_4 M_1,$$

$$\|\nabla J\|_{L_r(\mathbf{R}^N)} \leq C_4 M_2, \quad \|\nabla d\|_{L_r(\mathbf{R}^N)} \leq C_4 M_2.$$

This section mainly considers two strong elliptic problems as follows:

$$(4.4) \quad \begin{cases} \rho_{\pm} \lambda \tilde{v}_{\pm} - \Delta \tilde{v}_{\pm} = -\operatorname{div} \tilde{\mathbf{f}}_{\pm} + \tilde{g}_{\pm} & \text{in } \tilde{\Omega}_{\pm}, \\ \rho_{+} \tilde{v}_{+} = \rho_{-} \tilde{v}_{-} & \text{on } \tilde{\Sigma}, \\ \tilde{\mathbf{n}} \cdot \nabla(\tilde{v}_{+} - \tilde{v}_{-}) = \tilde{\mathbf{n}} \cdot (\tilde{\mathbf{f}}_{+} - \tilde{\mathbf{f}}_{-}) + \tilde{h}_{+} - \tilde{h}_{-} & \text{on } \tilde{\Sigma}, \end{cases}$$

and also

$$(4.5) \quad \begin{cases} \Delta \tilde{v}_{\pm} = \operatorname{div} \tilde{\mathbf{f}}_{\pm} & \text{in } \tilde{\Omega}_{\pm}, \\ \rho_{+} \tilde{v}_{+} = \rho_{-} \tilde{v}_{-} & \text{on } \tilde{\Sigma}, \\ \tilde{\mathbf{n}} \cdot \nabla(\tilde{v}_{+} - \tilde{v}_{-}) = \tilde{\mathbf{n}} \cdot (\tilde{\mathbf{f}}_{+} - \tilde{\mathbf{f}}_{-}) & \text{on } \tilde{\Sigma}. \end{cases}$$

Assume that ρ_{\pm} are positive constants throughout this section and set $F_q(\tilde{\Omega}_{\pm})$ as

$$F_q(\tilde{\Omega}_{\pm}) = \{\tilde{\mathbf{f}}_{\pm} \in E_q(\tilde{\Omega}_{\pm}) : \tilde{\mathbf{n}} \cdot \tilde{\mathbf{f}}_{\pm} \in H_q^1(\tilde{\Omega}_{\pm})\}.$$

Concerning (4.4) and (4.5), we prove the following theorems.

Theorem 4.2. *Let M_1 , M_2 , r , $\tilde{\Omega}_{\pm}$, and $\tilde{\Sigma}$ be as above, and let $\sigma \in (0, \pi/2)$ and $q \in (1, r]$. Then there exist $M_1 \in (0, 1/2)$ and $\lambda_1 \geq 1$ such that, for any*

$$\tilde{\mathbf{f}}_{\pm} \in F_q(\tilde{\Omega}_{\pm}), \quad \tilde{g}_{\pm} \in L_q(\tilde{\Omega}_{\pm}), \quad \tilde{h}_{\pm} \in H_q^1(\tilde{\Omega}_{\pm})$$

and for any $\lambda \in \Sigma_{\sigma, \lambda_1}$, the strong elliptic problem (4.4) admits unique solutions $\tilde{v}_{\pm} \in H_q^2(\tilde{\Omega}_{\pm})$. In addition, the solutions \tilde{v}_{\pm} satisfies

$$(4.6) \quad \begin{aligned} & \|(\lambda \tilde{v}_{\pm}, \lambda^{1/2} \nabla \tilde{v}_{\pm}, \nabla^2 \tilde{v}_{\pm})\|_{L_q(\tilde{\Omega}_{\pm})} \\ & \leq C \sum_{s \in \{+, -\}} \left\| \left(\operatorname{div} \tilde{\mathbf{f}}_s, \tilde{g}_s, \lambda^{1/2} (\tilde{\mathbf{n}} \cdot \tilde{\mathbf{f}}_s), \nabla (\tilde{\mathbf{n}} \cdot \tilde{\mathbf{f}}_s), \lambda^{1/2} \tilde{h}_s, \nabla \tilde{h}_s \right) \right\|_{L_q(\tilde{\Omega}_s)}, \end{aligned}$$

and also

$$(4.7) \quad \begin{aligned} \|(\lambda^{1/2} \tilde{v}_{\pm}, \nabla \tilde{v}_{\pm})\|_{L_q(\tilde{\Omega}_{\pm})} & \leq C \sum_{s \in \{+, -\}} \left(\|\tilde{\mathbf{f}}_s\|_{L_q(\tilde{\Omega}_s)} + |\lambda|^{-1/2} \|\tilde{g}_s\|_{L_q(\tilde{\Omega}_s)} \right. \\ & \quad \left. + \|\tilde{h}_s\|_{L_q(\tilde{\Omega}_s)} + |\lambda|^{-1/2} \|\nabla \tilde{h}_s\|_{L_q(\tilde{\Omega}_s)} \right), \end{aligned}$$

where $C = C(M_2, r, N, q, \sigma, \rho_{+}, \rho_{-})$ is a positive constant independent of M_1 , λ_1 , and λ . Additionally, if $\tilde{\mathbf{n}} \cdot (\tilde{\mathbf{f}}_{+} - \tilde{\mathbf{f}}_{-}) = 0$ on $\tilde{\Sigma}$, then \tilde{v}_{\pm} satisfy

$$(4.8) \quad \begin{aligned} & \|(\lambda \tilde{v}_{\pm}, \lambda^{1/2} \nabla \tilde{v}_{\pm}, \nabla^2 \tilde{v}_{\pm})\|_{L_q(\tilde{\Omega}_{\pm})} \\ & \leq C \sum_{s \in \{+, -\}} \left\| \left(\operatorname{div} \tilde{\mathbf{f}}_s, \tilde{g}_s, \lambda^{1/2} \tilde{h}_s, \nabla \tilde{h}_s \right) \right\|_{L_q(\tilde{\Omega}_s)}. \end{aligned}$$

Here, M_1 depends only on N, q, σ, ρ_+ , and ρ_- , while λ_1 depends only on $M_2, r, N, q, \sigma, \rho_+$, and ρ_- .

Theorem 4.3. *Let $M_1, M_2, r, \tilde{\Omega}_\pm$, and $\tilde{\Sigma}$ be as above, and let $q \in (1, \infty)$. Assume that $\max(q, q') \leq r$ for $q' = q/(q-1)$. Then there exists $M_1 \in (0, 1/2)$, depending only on N, q, ρ_+ , and ρ_- , such that the following assertions hold.*

(1) **Existence.** *Let $\tilde{\mathbf{f}}_\pm \in F_q(\tilde{\Omega}_\pm)$. Then the strong elliptic problem (4.5) admits solutions $\tilde{v}_\pm \in \hat{H}_q^1(\tilde{\Omega}_\pm) \cap \hat{H}_q^2(\tilde{\Omega}_\pm)$ satisfying*

$$(4.9) \quad \|\nabla^2 \tilde{v}_\pm\|_{L_q(\tilde{\Omega}_\pm)} \leq C \sum_{s \in \{+, -\}} \left(\|\tilde{\mathbf{f}}_s\|_{E_q(\tilde{\Omega}_s)} + \|\tilde{\mathbf{n}} \cdot \tilde{\mathbf{f}}_s\|_{H_q^1(\tilde{\Omega}_s)} \right),$$

$$(4.10) \quad \|\nabla \tilde{v}_\pm\|_{L_q(\tilde{\Omega}_\pm)} \leq C' \sum_{s \in \{+, -\}} \|\tilde{\mathbf{f}}_s\|_{L_q(\tilde{\Omega}_s)},$$

with positive constants $C = C(M_2, r, N, q, \rho_+, \rho_-)$ and $C' = C'(N, q, \rho_+, \rho_-)$ independent of M_1 . Additionally, if $\tilde{\mathbf{n}} \cdot (\tilde{\mathbf{f}}_+ - \tilde{\mathbf{f}}_-) = 0$ on $\tilde{\Sigma}$, then \tilde{v}_\pm satisfy

$$(4.11) \quad \|\nabla^2 \tilde{v}_\pm\|_{L_q(\tilde{\Omega}_\pm)} \leq C \sum_{s \in \{+, -\}} \|\tilde{\mathbf{f}}_s\|_{E_q(\tilde{\Omega}_s)}.$$

(2) **Uniqueness.** *If $\tilde{v}_\pm \in \hat{H}_q^1(\tilde{\Omega}_\pm) \cap \hat{H}_q^2(\tilde{\Omega}_\pm)$ satisfy*

$$\Delta \tilde{v}_\pm = 0 \text{ in } \tilde{\Omega}_\pm, \quad \rho_+ \tilde{v}_+ = \rho_- \tilde{v}_- \text{ and } \tilde{\mathbf{n}} \cdot \nabla(\tilde{v}_+ - \tilde{v}_-) = 0 \text{ on } \tilde{\Sigma},$$

then $\tilde{v}_\pm = \rho_\pm^{-1} c$ for some constant c .

At this point, we introduce the following lemma, see [23, Lemma 2.4], which play an important role in proving Theorems 4.2 and 4.3 in the following subsections.

Lemma 4.4. *Let $q \in (1, \infty)$ and $r \in (N, \infty)$. Assume that $q \leq r$. Then there exists a positive constant $C = C(N, q, r)$ such that, for any $\varepsilon > 0$, $a \in L_r(\mathbf{R}_+^N)$, and $b \in H_q^1(\mathbf{R}_+^N)$,*

$$\|ab\|_{L_q(\mathbf{R}_+^N)} \leq \varepsilon \|\nabla b\|_{L_q(\mathbf{R}_+^N)} + C \varepsilon^{-\frac{N}{r-N}} \|a\|_{L_r(\mathbf{R}_+^N)}^{\frac{r}{r-N}} \|b\|_{L_q(\mathbf{R}_+^N)}.$$

4.1. Proof of Theorem 4.2. To prove Theorem 4.2, we first reduce (4.4) to a problem in the whole space with the flat interface by the change of variables: $y = \Phi(x)$. Let $D_j = \partial/\partial y_j$ and $\partial_j = \partial/\partial x_j$. One then notes the following fundamental relations: For $j, k = 1, \dots, N$,

$$\begin{aligned} D_j &= \sum_{l=1}^N (\mathcal{A}_{lj} + B_{lj}(x)) \partial_l, \quad \nabla_y = (\mathcal{A}_{-1} + \mathbf{B}_{-1}(x))^\top \nabla_x, \\ D_j D_k &= \sum_{l,m=1}^N \mathcal{A}_{lj} \mathcal{A}_{mk} \partial_l \partial_m \\ &\quad + \sum_{l,m=1}^N (\mathcal{A}_{lj} B_{mk}(x) + \mathcal{A}_{mk} B_{lj}(x) + B_{lj}(x) B_{mk}(x)) \partial_l \partial_m \\ &\quad + \sum_{l,m=1}^N (\mathcal{A}_{lj} + B_{lj}(x)) (\partial_l B_{mk})(x) \partial_m. \end{aligned}$$

Let $\tilde{\mathbf{u}}$ be a vector function on $\mathbf{R}_y^N \setminus \tilde{\Sigma} = \tilde{\Omega}_+ \cup \tilde{\Omega}_-$, and let $\mathbf{u} = \tilde{\mathbf{u}} \circ \Phi$. Then, for any $\tilde{\varphi} \in C_0^\infty(\mathbf{R}_y^N \setminus \tilde{\Sigma})$ and for $\varphi = \tilde{\varphi} \circ \Phi$,

$$\begin{aligned} (\operatorname{div}_y \tilde{\mathbf{u}}, \tilde{\varphi})_{\mathbf{R}_y^N \setminus \tilde{\Sigma}} &= -(\tilde{\mathbf{u}}, \nabla_y \tilde{\varphi})_{\mathbf{R}_y^N \setminus \tilde{\Sigma}} = -(J\mathbf{u}, (\mathcal{A}_{-1} + \mathbf{B}_{-1})^\top \nabla_x \varphi)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} \\ &= (\operatorname{div}_x \{J(\mathcal{A}_{-1} + \mathbf{B}_{-1})\mathbf{u}\}, \varphi)_{\mathbf{R}^N \setminus \mathbf{R}_0^N} \\ &= ([J^{-1} \operatorname{div}_x \{J(\mathcal{A}_{-1} + \mathbf{B}_{-1})\mathbf{u}\}] \circ \Phi^{-1}, \tilde{\varphi})_{\mathbf{R}_y^N \setminus \tilde{\Sigma}}, \end{aligned}$$

which implies

$$(4.12) \quad (\operatorname{div}_y \tilde{\mathbf{u}}) \circ \Phi = J^{-1} \operatorname{div}_x \{J(\mathcal{A}_{-1} + \mathbf{B}_{-1})\mathbf{u}\}.$$

Analogously, we observe for a scalar function \tilde{u} on $\mathbf{R}_y^N \setminus \tilde{\Sigma}$ and $u = \tilde{u} \circ \Phi$ that

$$\begin{aligned} (\Delta_y \tilde{u}, \tilde{\varphi})_{\mathbf{R}_y^N \setminus \tilde{\Sigma}} &= (\nabla_y \tilde{u}, \nabla_y \tilde{\varphi})_{\mathbf{R}_y^N \setminus \tilde{\Sigma}} \\ &= (J(\mathcal{A}_{-1} + \mathbf{B}_{-1})^\top \nabla_x u, (\mathcal{A}_{-1} + \mathbf{B}_{-1})^\top \nabla_x \varphi)_{\mathbf{R}_x^N \setminus \mathbf{R}_0^N} \\ &= (\operatorname{div}_x \{J(\mathcal{A}_{-1} + \mathbf{B}_{-1})(\mathcal{A}_{-1} + \mathbf{B}_{-1})^\top \nabla_x u\}, \varphi)_{\mathbf{R}_x^N \setminus \mathbf{R}_0^N}, \end{aligned}$$

and thus

$$(4.13) \quad (\Delta_y \tilde{u}) \circ \Phi = J^{-1} \operatorname{div}_x \{J(\mathcal{A}_{-1} + \mathbf{B}_{-1})(\mathcal{A}_{-1} + \mathbf{B}_{-1})^\top \nabla_x u\}.$$

For simplicity, one sets

$$\mathbf{C}_{-1}(x) = \mathcal{A}_{-1} \mathbf{B}_{-1}(x)^\top + \mathbf{B}_{-1}(x) \mathcal{A}_{-1}^\top + \mathbf{B}_{-1}(x) \mathbf{B}_{-1}(x)^\top,$$

and then

$$(4.14) \quad (\mathcal{A}_{-1} + \mathbf{B}_{-1}(x))(\mathcal{A}_{-1} + \mathbf{B}_{-1}(x))^\top = \mathbf{I} + \mathbf{C}_{-1}(x).$$

Furthermore, let us define

$$v_\pm = \tilde{v}_\pm \circ \Phi, \quad \mathbf{f}_\pm = \tilde{\mathbf{f}}_\pm \circ \Phi, \quad g_\pm = J(\tilde{g}_\pm \circ \Phi), \quad h_\pm = Jd(\tilde{h}_\pm \circ \Phi).$$

Then, by (4.2) and (4.12)-(4.14), we obtain an equivalent system of (4.4) as follows:

$$(4.15) \quad \begin{cases} \rho_\pm \lambda v_\pm - \Delta v_\pm = -\operatorname{div}(\mathbf{F}_\pm + \mathcal{F}(v_\pm)) + \rho_\pm \mathcal{G}_\lambda(v_\pm) + g_\pm & \text{in } \mathbf{R}_\pm^N, \\ \rho_+ v_+ = \rho_- v_- & \text{on } \mathbf{R}_0^N, \\ \mathbf{n}_0 \cdot \nabla(v_+ - v_-) = \mathbf{n}_0 \cdot \{(\mathbf{F}_+ + \mathcal{F}_+(v_+)) - (\mathbf{F}_- + \mathcal{F}_-(v_-))\} \\ \quad + h_+ - h_- & \text{on } \mathbf{R}_0^N, \end{cases}$$

where we have set

$$(4.16) \quad \begin{aligned} \mathbf{F}_\pm &= J(\mathcal{A}_{-1} + \mathbf{B}_{-1})\mathbf{f}_\pm, \quad \mathcal{F}(v_\pm) = (1 - J)\nabla v_\pm - J\mathbf{C}_{-1}\nabla v_\pm, \\ \mathcal{G}_\lambda(v_\pm) &= \lambda(1 - J)v_\pm. \end{aligned}$$

Remark 4.5. (1) The above \mathcal{F} and \mathcal{G}_λ are linear mappings.

(2) By (4.1), there is a constant $C_5 > 0$, independent of M_1 , M_2 , and r , such that

$$(4.17) \quad \|\mathbf{C}_{-1}\|_{L^\infty(\mathbf{R}^N)} \leq C_5 M_1, \quad \|\nabla \mathbf{C}_{-1}\|_{L^r(\mathbf{R}^N)} \leq C_5 M_2.$$

In what follows, $F_{\pm N}$ stands for the N th component of \mathbf{F}_\pm , while $\mathcal{F}_N(v_\pm)$ the N th component of $\mathcal{F}(v_\pm)$. By (4.3) and (4.12),

$$(4.18) \quad \|\operatorname{div} \mathbf{F}_\pm\|_{L_q(\mathbf{R}_\pm^N)} \leq C_4^{1-1/q} \|\operatorname{div} \tilde{\mathbf{f}}_\pm\|_{L_q(\tilde{\Omega}_\pm)}.$$

In addition, since it holds by (4.2) that

$$(4.19) \quad F_{\pm N} = -\mathbf{n}_0 \cdot \{J(\mathcal{A}_{-1} + \mathbf{B}_{-1})\mathbf{f}_\pm\} = -Jd(\tilde{\mathbf{n}} \cdot \tilde{\mathbf{f}}_\pm),$$

one has by (4.3) and Lemma 4.4 with $\varepsilon = 1$

$$(4.20) \quad \|F_{\pm N}\|_{H_q^j(\mathbf{R}_{\pm}^N)} \leq C_{M_2, r} \|\tilde{\mathbf{n}} \cdot \tilde{\mathbf{f}}_{\pm}\|_{H_q^j(\tilde{\Omega}_{\pm})} \quad (j = 0, 1).$$

Here and subsequently, $C_{M_2, r}$ stands for generic positive constants depending on M_2 , N , q , and r , but independent of M_1 . Similarly to (4.18) and (4.20),

$$(4.21) \quad \begin{aligned} \|\mathbf{F}_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C_6 \|\tilde{\mathbf{f}}_{\pm}\|_{L_q(\tilde{\Omega}_{\pm})}, \quad \|g_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)} \leq C_6 \|\tilde{g}_{\pm}\|_{L_q(\tilde{\Omega}_{\pm})}, \\ \|h_{\pm}\|_{H_q^j(\mathbf{R}_{\pm}^N)} &\leq C_{M_2, r} \|\tilde{h}_{\pm}\|_{H_q^j(\tilde{\Omega}_{\pm})} \quad (j = 0, 1), \end{aligned}$$

where C_6 is a positive constant independent of M_1 , M_2 , and r . Concerning $\mathcal{F}(v_{\pm})$ and $\mathcal{G}_{\lambda}(v_{\pm})$, we have by Lemma 4.4, (4.3), and (4.17)

Lemma 4.6. *Let $q \in (1, \infty)$, $r \in (N, \infty)$, and $\varepsilon > 0$. Assume that $q \leq r$. Then there exist a positive constant $\alpha_{M_2, r, \varepsilon}$ depending on M_2 , r , and ε , but independent of M_1 , and a positive constant C_7 independent of M_1 , M_2 , r , and ε , such that for any $w_{\pm} \in H_q^2(\mathbf{R}_{\pm}^N)$*

$$\begin{aligned} &\|(\operatorname{div} \mathcal{F}(w_{\pm}), \nabla \mathcal{F}(w_{\pm}))\|_{L_q(\mathbf{R}_{\pm}^N)} \\ &\leq C_7(M_1 + \varepsilon) \|\nabla^2 w_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)} + \alpha_{M_2, r, \varepsilon} \|\nabla w_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)}, \end{aligned}$$

and also

$$\begin{aligned} \|\mathcal{F}(w_{\pm})\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C_7 M_1 \|\nabla w_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)}, \\ \|\rho_{\pm} \mathcal{G}_{\lambda}(w_{\pm})\|_{L_q(\mathbf{R}_{\pm}^N)} &\leq C_7 M_1 |\lambda| \|w_{\pm}\|_{L_q(\mathbf{R}_{\pm}^N)} \quad (\lambda \in \mathbf{C}). \end{aligned}$$

From now on, we solve (4.15) by using the contraction mapping principle together with Theorem 3.1. Let $\lambda_1 \geq 1$ and $\lambda \in \Sigma_{\sigma, \lambda_1}$ for some $\sigma \in (0, \pi/2)$. We set $X_{q, \lambda} = H_q^2(\mathbf{R}_+^N) \times H_q^2(\mathbf{R}_-^N)$ endowed with the norm:

$$\|(w_+, w_-)\|_{X_{q, \lambda}} = \sum_{s \in \{+, -\}} \|(\lambda w_s, \lambda^{1/2} \nabla w_s, \nabla^2 w_s)\|_{L_q(\mathbf{R}_s^N)},$$

and also for $R > 0$

$$X_{q, \lambda}(R) = \{(w_+, w_-) \in X_{q, \lambda} : \|(w_+, w_-)\|_{X_{q, \lambda}} \leq R\}.$$

The next lemma then follows from Lemma 4.6 immediately.

Lemma 4.7. *Let λ_1 and λ be as above. Let $q \in (1, \infty)$, $r \in (N, \infty)$, and $\varepsilon > 0$. Assume that $q \leq r$. Then, for any $w_{\pm} \in H_q^2(\mathbf{R}_{\pm}^N)$,*

$$\begin{aligned} \sum_{s \in \{+, -\}} \|\nabla \mathcal{F}(w_s)\|_{L_q(\mathbf{R}_s^N)} &\leq \left\{ C_7(M_1 + \varepsilon) + \alpha_{M_2, r, \varepsilon} \lambda_1^{-1/2} \right\} \|(w_+, w_-)\|_{X_{q, \lambda}}, \\ \sum_{s \in \{+, -\}} \|\lambda^{1/2} \mathcal{F}(w_s)\|_{L_q(\mathbf{R}_s^N)} &\leq C_7 M_1 \|(w_+, w_-)\|_{X_{q, \lambda}}, \\ \sum_{s \in \{+, -\}} \|\rho_s \mathcal{G}_{\lambda}(w_s)\|_{L_q(\mathbf{R}_s^N)} &\leq C_7 M_1 \|(w_+, w_-)\|_{X_{q, \lambda}}, \end{aligned}$$

and also

$$\begin{aligned} &\sum_{s \in \{+, -\}} \left\| \left(\mathcal{F}(w_s), |\lambda|^{-1/2} \rho_s \mathcal{G}_{\lambda}(w_s) \right) \right\|_{L_q(\mathbf{R}_s^N)} \\ &\leq C_7 M_1 \sum_{s \in \{+, -\}} \|(\lambda^{1/2} w_s, \nabla w_s)\|_{L_q(\mathbf{R}_s^N)}. \end{aligned}$$

Here C_7 and $\alpha_{M_2, r, \varepsilon}$ are the same positive constants as in Lemma 4.6.

Let us define K_λ and L_λ as follows:

$$K_\lambda = \begin{cases} \sum_{s \in \{+, -\}} \|(\operatorname{div} \mathbf{F}_s, g_s, \lambda^{1/2} F_{sN}, \nabla F_{sN}, \lambda^{1/2} h_s, \nabla h_s)\|_{L_q(\mathbf{R}_s^N)}, \\ \sum_{s \in \{+, -\}} \|(\operatorname{div} \mathbf{F}_s, g_s, \lambda^{1/2} h_s, \nabla h_s)\|_{L_q(\mathbf{R}_s^N)} \text{ when } \tilde{\mathbf{n}} \cdot (\tilde{\mathbf{f}}_+ - \tilde{\mathbf{f}}_-) = 0 \text{ on } \tilde{\Sigma}, \end{cases}$$

$$L_\lambda = \sum_{s \in \{+, -\}} \left\| \left(\mathbf{F}_s, |\lambda|^{-1/2} g_s, h_s, |\lambda|^{-1/2} \nabla h_s \right) \right\|_{L_q(\mathbf{R}_s^N)}.$$

Fix a positive number S such that $K_\lambda \leq S/2$. Let C_1 be the same positive constant as in Theorem 3.1, and let $(w_+, w_-) \in X_{q, \lambda}(C_1 S)$. We consider

$$(4.22) \quad \begin{cases} \rho_\pm \lambda z_\pm - \Delta z_\pm = -\operatorname{div}(\mathbf{F}_\pm + \mathcal{F}(w_\pm)) + \rho_\pm \mathcal{G}_\lambda(w_\pm) + g_\pm & \text{in } \mathbf{R}_\pm^N, \\ \rho_+ z_+ = \rho_- z_- & \text{on } \mathbf{R}_0^N, \\ \mathbf{n}_0 \cdot \nabla(z_+ - z_-) = \mathbf{n}_0 \cdot \{(\mathbf{F}_+ + \mathcal{F}(w_+)) - (\mathbf{F}_- + \mathcal{F}(w_-))\} \\ \quad + h_+ - h_- & \text{on } \mathbf{R}_0^N. \end{cases}$$

Noting $\mathbf{n}_0 \cdot (\mathbf{F}_+ - \mathbf{F}_-) = 0$ on \mathbf{R}_0^N when $\tilde{\mathbf{n}} \cdot (\tilde{\mathbf{f}}_+ - \tilde{\mathbf{f}}_-) = 0$ on $\tilde{\Sigma}$ by virtue of (4.19), one has by Theorem 3.1 unique solutions $z_\pm \in H_q^2(\mathbf{R}_\pm^N)$ of (4.22) satisfying

$$\begin{aligned} & \| (z_+, z_-) \|_{X_{q, \lambda}} \\ & \leq C_1 \left(K_\lambda + \sum_{s \in \{+, -\}} \|(\operatorname{div} \mathcal{F}(w_s), \rho_s \mathcal{G}_\lambda(w_s), \lambda^{1/2} \mathcal{F}_N(w_s), \nabla \mathcal{F}_N(w_s))\|_{L_q(\mathbf{R}_s^N)} \right). \end{aligned}$$

Combining this inequality with Lemma 4.7 furnishes an a priori estimate for the solutions of (4.22) as follows:

$$(4.23) \quad \begin{aligned} \| (z_+, z_-) \|_{X_{q, \lambda}} & \leq C_1 \left\{ K_\lambda + 4 \left(C_7(M_1 + \varepsilon) + \alpha_{M_2, r, \varepsilon} \lambda_1^{-1/2} \right) \| (w_+, w_-) \|_{X_{q, \lambda}} \right\} \\ & \leq C_1 \left\{ K_\lambda + 4C_1 S \left(C_7(M_1 + \varepsilon) + \alpha_{M_2, r, \varepsilon} \lambda_1^{-1/2} \right) \right\}. \end{aligned}$$

Choosing M_1 , ε small enough and λ_1 large enough so that

$$(4.24) \quad 4C_1 C_7(M_1 + \varepsilon) \leq \frac{1}{4}, \quad 4C_1 \alpha_{M_2, r, \varepsilon} \lambda_1^{-1/2} \leq \frac{1}{4},$$

we see from (4.23) that

$$\| (z_+, z_-) \|_{X_{q, \lambda}} \leq C_1 \left(\frac{S}{2} + \frac{S}{2} \right) = C_1 S.$$

Thus we can define a map $\Psi : X_{q, \lambda}(C_1 S) \rightarrow X_{q, \lambda}(C_1 S)$ by $\Psi(w_+, w_-) = (z_+, z_-)$. Analogously, recalling Remark 4.5 (1) and using the a priori estimate (4.23), we can show that Ψ is a contraction mapping on $X_{q, \lambda}(C_1 S)$, which furnishes that there exists a fixed point $(w_+^*, w_-^*) \in X_{q, \lambda}(C_1 S)$, i.e. $(w_+^*, w_-^*) = \Psi(w_+^*, w_-^*)$. Then $(v_+, v_-) = (w_+^*, w_-^*)$ is a solution to (4.15) and satisfies by the first inequality of (4.23), together with (4.24),

$$(4.25) \quad \| (v_+, v_-) \|_{X_{q, \lambda}} \leq 2C_1 K_\lambda.$$

In addition, by (3.4),

$$\begin{aligned} & \sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda^{1/2} v_{\mathfrak{s}}, \nabla v_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} \\ & \leq C_1 \left(L_\lambda + \sum_{\mathfrak{s} \in \{+, -\}} \left\| \left(\mathcal{F}(v_{\mathfrak{s}}), |\lambda|^{-1/2} \rho_{\mathfrak{s}} \mathcal{G}_\lambda(v_{\mathfrak{s}}) \right) \right\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} \right), \end{aligned}$$

which, combined with Lemma 4.7 and (4.24), furnishes

$$\begin{aligned} \sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda^{1/2} v_{\mathfrak{s}}, \nabla v_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} & \leq C_1 \left(L_\lambda + C_7 M_1 \sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda^{1/2} v_{\mathfrak{s}}, \nabla v_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} \right) \\ & \leq C_1 L_\lambda + \frac{1}{2} \sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda^{1/2} v_{\mathfrak{s}}, \nabla v_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)}. \end{aligned}$$

It therefore holds that

$$(4.26) \quad \sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda^{1/2} v_{\mathfrak{s}}, \nabla v_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)} \leq 2C_1 L_\lambda.$$

The uniqueness of solutions of (4.15) also follows from the a priori estimate (4.23) together with (4.24).

Finally, setting $\tilde{v}_\pm = v_\pm \circ \Phi^{-1}$, we observe that $\tilde{v}_\pm \in H_q^2(\tilde{\Omega}_\pm)$ are solutions to (4.4) and that by (4.1), (4.3), and Lemma 4.4 with $\varepsilon = 1$

$$\sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda \tilde{v}_{\mathfrak{s}}, \lambda^{1/2} \nabla \tilde{v}_{\mathfrak{s}}, \nabla^2 \tilde{v}_{\mathfrak{s}})\|_{L_q(\tilde{\Omega}_{\mathfrak{s}})} \leq C \| (v_+, v_-) \|_{X_{q,\lambda}}$$

for some positive constant $C = C(M_2, N, q, r)$ independent of M_1 , λ_1 , and λ . Combining this inequality with (4.25), together with (4.18)-(4.21), furnishes (4.6) and (4.8). On the other hand, since

$$\sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda^{1/2} \tilde{v}_{\mathfrak{s}}, \nabla \tilde{v}_{\mathfrak{s}})\|_{L_q(\tilde{\Omega}_{\mathfrak{s}})} \leq C \sum_{\mathfrak{s} \in \{+, -\}} \|(\lambda^{1/2} v_{\mathfrak{s}}, \nabla v_{\mathfrak{s}})\|_{L_q(\mathbf{R}_{\mathfrak{s}}^N)}$$

for some positive constant C independent of M_1 , M_2 , λ_1 , λ , and r , the required estimate (4.7) follows from (4.26) together with (4.21). The uniqueness of solutions of (4.4) follows from the uniqueness of solutions of the equivalent system (4.15). This completes the proof of the Theorem 4.2.

4.2. Proof of Theorem 4.3. The uniqueness of solutions of (4.5) follows from the existence of solutions of (4.5) similarly to the proof of Theorem 3.2 (2), so that we prove the existence of solutions of (4.5) by following the idea of one-phase flows introduced in [22] in what follows.

Setting $v_\pm = \tilde{v}_\pm \circ \Phi$ and $\mathbf{f} = \tilde{\mathbf{f}}_\pm \circ \Phi$ in (4.5), we have, similarly to the previous subsection,

$$(4.27) \quad \begin{cases} \Delta v_\pm = \operatorname{div}(\mathbf{F}_\pm + \mathcal{F}(v_\pm)) & \text{in } \mathbf{R}_\pm^N, \\ \rho_+ v_+ = \rho_- v_- & \text{on } \mathbf{R}_0^N, \\ \mathbf{n}_0 \cdot \nabla(v_+ - v_-) = \mathbf{n}_0 \cdot \{(\mathbf{F}_+ + \mathcal{F}(v_+)) - (\mathbf{F}_- + \mathcal{F}(v_-))\} & \text{on } \mathbf{R}_0^N, \end{cases}$$

where \mathbf{F}_\pm and $\mathcal{F}(v_\pm)$ are given in (4.16). By Theorem 3.2, one sees for $w_\pm \in \widehat{H}_q^1(\mathbf{R}_\pm^N) \cap \widehat{H}_q^2(\mathbf{R}_\pm^N)$ that there is a unique solution $z_\pm \in \widehat{H}_q^1(\mathbf{R}_\pm^N) \cap \widehat{H}_q^2(\mathbf{R}_\pm^N)$ to

$$(4.28) \quad \begin{cases} \Delta z_\pm = \operatorname{div}(\mathbf{F}_\pm + \mathcal{F}(w_\pm)) & \text{in } \mathbf{R}_\pm^N, \\ \rho_+ z_+ = \rho_- z_- & \text{on } \mathbf{R}_0^N, \\ \mathbf{n}_0 \cdot \nabla(z_+ - z_-) = \mathbf{n}_0 \cdot \{(\mathbf{F}_+ + \mathcal{F}(w_+)) - (\mathbf{F}_- + \mathcal{F}(w_-))\} & \text{on } \mathbf{R}_0^N. \end{cases}$$

In addition, by (3.6) and (3.7),

$$\begin{aligned} \sum_{s \in \{+, -\}} \|\nabla^2 z_s\|_{L_q(\mathbf{R}_s^N)} &\leq C_2 \sum_{s \in \{+, -\}} \|(\operatorname{div} \mathbf{F}_s, \nabla F_{sN}, \operatorname{div} \mathcal{F}(w_s), \nabla \mathcal{F}_N(w_s))\|_{L_q(\mathbf{R}_s^N)}, \\ \sum_{s \in \{+, -\}} \|\nabla z_s\|_{L_q(\mathbf{R}_s^N)} &\leq C_2 \sum_{s \in \{+, -\}} \|(\mathbf{F}_s, \mathcal{F}(w_s))\|_{L_q(\mathbf{R}_s^N)}, \end{aligned}$$

which, combined with Lemma 4.6, furnishes a priori estimates for the solutions of (4.28) as follows:

$$(4.29) \quad \sum_{s \in \{+, -\}} \|\nabla^2 z_s\|_{L_q(\mathbf{R}_s^N)} \leq C_2 \sum_{s \in \{+, -\}} \left(\|(\operatorname{div} \mathbf{F}_s, \nabla F_{sN})\|_{L_q(\mathbf{R}_s^N)} + C_7(M_1 + \varepsilon) \|\nabla^2 w_s\|_{L_q(\mathbf{R}_s^N)} + \alpha_{M_2, r, \varepsilon} \|\nabla w_s\|_{L_q(\mathbf{R}_s^N)} \right),$$

$$(4.30) \quad \sum_{s \in \{+, -\}} \|\nabla z_s\|_{L_q(\mathbf{R}_s^N)} \leq C_2 \sum_{s \in \{+, -\}} \left(\|\mathbf{F}_s\|_{L_q(\mathbf{R}_s^N)} + C_7 M_1 \|\nabla w_s\|_{L_q(\mathbf{R}_s^N)} \right).$$

Remark 4.8. If $\tilde{\mathbf{n}} \cdot (\tilde{\mathbf{f}}_+ - \tilde{\mathbf{f}}_-) = 0$ on $\tilde{\Sigma}$, then $\mathbf{n}_0 \cdot (\mathbf{F}_+ - \mathbf{F}_-) = 0$ on \mathbf{R}_0^N by (4.19). When $\tilde{\mathbf{n}} \cdot (\tilde{\mathbf{f}}_+ - \tilde{\mathbf{f}}_-) = 0$ on $\tilde{\Sigma}$, one thus has by (3.8)

$$\sum_{s \in \{+, -\}} \|\nabla^2 z_s\|_{L_q(\mathbf{R}_s^N)} \leq C_2 \sum_{s \in \{+, -\}} \|(\operatorname{div} \mathbf{F}_s, \operatorname{div} \mathcal{F}(w_s))\|_{L_q(\mathbf{R}_s^N)},$$

and then the a priori estimate (4.29) is replaced by

$$(4.31) \quad \sum_{s \in \{+, -\}} \|\nabla^2 z_s\|_{L_q(\mathbf{R}_s^N)} \leq C_2 \sum_{s \in \{+, -\}} \left(\|\operatorname{div} \mathbf{F}_s\|_{L_q(\mathbf{R}_s^N)} + C_7(M_1 + \varepsilon) \|\nabla^2 w_s\|_{L_q(\mathbf{R}_s^N)} + \alpha_{M_2, r, \varepsilon} \|\nabla w_s\|_{L_q(\mathbf{R}_s^N)} \right).$$

From now on, we prove the existence of solutions v_\pm to (4.27). Let $v_\pm^{(0)} = 0$ and $v_\pm^{(j)} \in \widehat{H}_q^1(\mathbf{R}_\pm^N) \cap \widehat{H}_q^2(\mathbf{R}_\pm^N)$, $j \geq 1$, be the unique solutions to

$$(4.32) \quad \begin{cases} \Delta v_\pm^{(j)} = \operatorname{div}(\mathbf{F}_\pm + \mathcal{F}(v_\pm^{(j-1)})) & \text{in } \mathbf{R}_\pm^N, \\ \rho_+ v_+^{(j)} = \rho_- v_-^{(j)} & \text{on } \mathbf{R}_0^N, \\ \mathbf{n}_0 \cdot \nabla(v_+^{(j)} - v_-^{(j)}) = \mathbf{n}_0 \cdot \{(\mathbf{F}_+ + \mathcal{F}(v_+^{(j-1)})) - (\mathbf{F}_- + \mathcal{F}(v_-^{(j-1)}))\} & \text{on } \mathbf{R}_0^N. \end{cases}$$

By (4.29) and (4.30), $v_\pm^{(j)}$ satisfy

$$(4.33) \quad \begin{aligned} \sum_{s \in \{+, -\}} \|\nabla^2 v_s^{(j)}\|_{L_q(\mathbf{R}_s^N)} &\leq C_2 \sum_{s \in \{+, -\}} \left(\|(\operatorname{div} \mathbf{F}_s, \nabla F_{sN})\|_{L_q(\mathbf{R}_s^N)} + C_7(M_1 + \varepsilon) \|\nabla^2 v_s^{(j-1)}\|_{L_q(\mathbf{R}_s^N)} + \alpha_{M_2, r, \varepsilon} \|\nabla v_s^{(j-1)}\|_{L_q(\mathbf{R}_s^N)} \right), \end{aligned}$$

$$\begin{aligned}
 (4.34) \quad & \sum_{s \in \{+, -\}} \|\nabla v_s^{(j)}\|_{L_q(\mathbf{R}_s^N)} \\
 & \leq C_2 \sum_{s \in \{+, -\}} \left(\|\mathbf{F}_s\|_{L_q(\mathbf{R}_s^N)} + C_7 M_1 \|\nabla v_s^{(j-1)}\|_{L_q(\mathbf{R}_s^N)} \right).
 \end{aligned}$$

Inductively, it follows from (4.34) and $v_\pm^{(0)} = 0$ that

$$(4.35) \quad \sum_{s \in \{+, -\}} \|\nabla v_s^{(j)}\|_{L_q(\mathbf{R}_s^N)} \leq X_j \sum_{s \in \{+, -\}} \|\mathbf{F}_s\|_{L_q(\mathbf{R}_s^N)}, \quad X_j = C_2 \sum_{k=0}^{j-1} (C_2 C_7 M_1)^k,$$

which, inserted into (4.33), furnishes

$$\begin{aligned}
 & \sum_{s \in \{+, -\}} \|\nabla^2 v_s^{(j)}\|_{L_q(\mathbf{R}_s^N)} \leq C_2 \sum_{s \in \{+, -\}} \left(\|(\operatorname{div} \mathbf{F}_s, \nabla F_{sN})\|_{L_q(\mathbf{R}_s^N)} \right. \\
 & \left. + C_7 (M_1 + \varepsilon) \|\nabla^2 v_s^{(j-1)}\|_{L_q(\mathbf{R}_s^N)} + \alpha_{M_2, r, \varepsilon} X_j \|\mathbf{F}_s\|_{L_q(\mathbf{R}_s^N)} \right).
 \end{aligned}$$

Inductively, it follows from this inequality and $v_\pm^{(0)} = 0$ that

$$\begin{aligned}
 (4.36) \quad & \sum_{s \in \{+, -\}} \|\nabla^2 v_s^{(j)}\|_{L_q(\mathbf{R}_s^N)} \leq Y_j \sum_{s \in \{+, -\}} \|(\operatorname{div} \mathbf{F}_s, \nabla F_{sN})\|_{L_q(\mathbf{R}_s^N)} \\
 & + Z_j \sum_{s \in \{+, -\}} \|\mathbf{F}_s\|_{L_q(\mathbf{R}_s^N)},
 \end{aligned}$$

where

$$Y_j = C_2 \sum_{k=0}^{j-1} (C_2 C_7 (M_1 + \varepsilon))^k, \quad Z_j = C_2 \alpha_{M_2, r, \varepsilon} \sum_{k=0}^{j-1} (C_2 C_7 (M_1 + \varepsilon))^k X_{j-k}.$$

Let $u^{(j)} = \rho_+ v_+^{(j)} \mathbf{1}_{\mathbf{R}_+^N} + \rho_- v_-^{(j)} \mathbf{1}_{\mathbf{R}_-^N}$. One then sees by $\rho_+ v_+^{(j)} = \rho_- v_-^{(j)}$ on \mathbf{R}_0^N that $u^{(j)} \in \widehat{H}_q^1(\mathbf{R}^N)$ with $\nabla u^{(j)} = \rho_+ (\nabla v_+^{(j)}) \mathbf{1}_{\mathbf{R}_+^N} + \rho_- (\nabla v_-^{(j)}) \mathbf{1}_{\mathbf{R}_-^N}$. On the other hand, recalling Remark 4.5 (1), we have by (4.30)

$$\sum_{s \in \{+, -\}} \|\nabla (v_s^{(j)} - v_s^{(j-1)})\|_{L_q(\mathbf{R}_s^N)} \leq C_2 C_7 M_1 \sum_{s \in \{+, -\}} \|\nabla (v_s^{(j-1)} - v_s^{(j-2)})\|_{L_q(\mathbf{R}_s^N)},$$

which, combined with

$$\begin{aligned}
 & \frac{1}{C_8} \sum_{s \in \{+, -\}} \|\nabla (v_s^{(j)} - v_s^{(j-1)})\|_{L_q(\mathbf{R}_s^N)} \\
 & \leq \|\nabla (u^{(j)} - u^{(j-1)})\|_{L_q(\mathbf{R}^N)} \leq C_8 \sum_{s \in \{+, -\}} \|\nabla (v_s^{(j)} - v_s^{(j-1)})\|_{L_q(\mathbf{R}_s^N)}
 \end{aligned}$$

for some positive constants $C_8 \geq 1$ depending only on q , ρ_+ , and ρ_- , furnishes

$$\|\nabla (u^{(j)} - u^{(j-1)})\|_{L_q(\mathbf{R}^N)} \leq C_2 C_7 (C_8)^2 M_1 \|\nabla (u^{(j-1)} - u^{(j-2)})\|_{L_q(\mathbf{R}^N)}.$$

Choose M_1 and ε small enough so that

$$C_2 C_7 (C_8)^2 M_1 \leq \frac{1}{2}, \quad C_2 C_7 (M_1 + \varepsilon) \leq \frac{1}{2}.$$

Then the above inequality for $u^{(j)} - u^{(j-1)}$ implies that $\{u^{(j)}\}_{j=1}^\infty$ is a Cauchy sequence in $\widehat{H}_q^1(\mathbf{R}^N)$, and thus there exists $u \in \widehat{H}_q^1(\mathbf{R}^N)$ such that

$$(4.37) \quad \lim_{j \rightarrow \infty} \|\nabla(u^{(j)} - u)\|_{L_q(\mathbf{R}^N)} = 0.$$

Furthermore, we see that

$$X_j \leq C_2 \sum_{k=0}^{\infty} 2^{-k} = 2C_2, \quad Y_j \leq 2C_2, \quad Z_j \leq C_2 \alpha_{M_2, r, \varepsilon} \cdot 2 \cdot 2C_2,$$

which, combined with (4.35) and (4.36), furnishes

$$(4.38) \quad \sum_{s \in \{+, -\}} \|\nabla v_s^{(j)}\|_{L_q(\mathbf{R}_s^N)} \leq 2C_2 \sum_{s \in \{+, -\}} \|\mathbf{F}_s\|_{L_q(\mathbf{R}_s^N)},$$

$$(4.39) \quad \sum_{s \in \{+, -\}} \|\nabla^2 v_s^{(j)}\|_{L_q(\mathbf{R}_s^N)} \leq 2C_2 K,$$

where we have set

$$K = \sum_{s \in \{+, -\}} \|(\operatorname{div} \mathbf{F}_s, \nabla F_{sN})\|_{L_q(\mathbf{R}_s^N)} + 2C_2 \alpha_{M_2, r, \varepsilon} \sum_{s \in \{+, -\}} \|\mathbf{F}_s\|_{L_q(\mathbf{R}_s^N)}.$$

Let u_+ and u_- be respectively the restriction of u on \mathbf{R}_+^N and the restriction of u on \mathbf{R}_-^N . It then holds that $u_\pm \in \widehat{H}_q^1(\mathbf{R}_\pm^N)$ and

$$(4.40) \quad \nabla u = (\nabla u_+) \mathbb{1}_{\mathbf{R}_+^N} + (\nabla u_-) \mathbb{1}_{\mathbf{R}_-^N}, \quad u_+ = u_- \text{ on } \mathbf{R}_0^N.$$

Let us define $v_\pm = \rho_\pm^{-1} u_\pm$. By the second property of (4.40),

$$(4.41) \quad \rho_+ v_+ = \rho_- v_- \text{ on } \mathbf{R}_0^N.$$

In addition, we see by the definition of $u^{(j)}$ and the first property of (4.40) that

$$\|\nabla(v_\pm^{(j)} - v_\pm)\|_{L_q(\mathbf{R}_\pm^N)} = \rho_\pm^{-1} \|\nabla(u^{(j)} - u)\|_{L_q(\mathbf{R}_\pm^N)}$$

which, combined with (4.37), furnishes

$$(4.42) \quad \lim_{j \rightarrow \infty} \|\nabla(v_\pm^{(j)} - v_\pm)\|_{L_q(\mathbf{R}_\pm^N)} = 0.$$

Taking the the limit: $j \rightarrow \infty$ in (4.38) thus implies that v_\pm satisfy

$$(4.43) \quad \sum_{s \in \{+, -\}} \|\nabla v_s\|_{L_q(\mathbf{R}_s^N)} \leq 2C_2 \sum_{s \in \{+, -\}} \|\mathbf{F}_s\|_{L_q(\mathbf{R}_s^N)},$$

Next, we consider the higher regularity of v_\pm . Let $k, l = 1, \dots, N$. By (4.39), there exist $v_{\pm, kl} \in L_q(\mathbf{R}_\pm^N)$ such that $\partial_k \partial_l v_\pm^{(j)} \rightarrow v_{\pm, kl}$ weakly in $L_q(\mathbf{R}_\pm^N)$ as $j \rightarrow \infty$. Then we can prove $\partial_k \partial_l v_\pm = v_{\pm, kl} \in L_q(\mathbf{R}_\pm^N)$ by using the convergence in distribution. Consequently, $\partial_k \partial_l v_\pm^{(j)} \rightarrow \partial_k \partial_l v_\pm$ weakly in $L_q(\mathbf{R}_\pm^N)$ as $j \rightarrow \infty$. It thus follows from (4.39) that

$$(4.44) \quad \|\partial_k \partial_l v_\pm\|_{L_q(\mathbf{R}_\pm^N)} \leq \liminf_{j \rightarrow \infty} \|\partial_k \partial_l v_\pm^{(j)}\|_{L_q(\mathbf{R}_\pm^N)} \leq 2C_2 K.$$

Finally, we prove that v_\pm satisfies (4.27) by the limit: $j \rightarrow \infty$ in (4.32). The first line of (4.27) is satisfied immediately, and the third line of (4.27) can be proved by Remark 4.5 (1), Lemma 4.6, and the inequality (cf. [4, Proposition 16.2]):

$$\|f_\pm\|_{L_q(\mathbf{R}_0^N)} \leq q^{1/q} \|f_\pm\|_{L_q(\mathbf{R}_\pm^N)}^{1-1/q} \|\nabla f_\pm\|_{L_q(\mathbf{R}_\pm^N)}^{1/q} \quad \text{for any } f_\pm \in H_q^1(\mathbf{R}_\pm^N),$$

together with (4.39), (4.42), and (4.44). In addition, the second line of (4.27) is already obtained in (4.41). Thus, setting $\tilde{v}_\pm = v_\pm \circ \Phi$, we see that \tilde{v}_\pm are solutions to (4.5) and that the required estimates (4.9) and (4.10) follow from (4.43) and (4.44) in the same manner as in the last part of Subsection 4.1. When $\tilde{\mathbf{n}} \cdot (\tilde{\mathbf{f}}_+ - \tilde{\mathbf{f}}_-) = 0$ on $\tilde{\Sigma}$, we can obtain (4.11) by using (4.31) instead of (4.29) in the above argument. This completes the proof of Theorem 4.3.

4.3. Weak elliptic problem with bent interface. Throughout this subsection, we assume that $r \in (N, \infty)$ and $\tilde{\Omega}_\pm, \tilde{\Sigma}$ are given in Theorem 4.3. For $\mathbf{f} \in L_q(\mathbf{R}^N \setminus \tilde{\Sigma})^N$ with $q \in (1, \infty)$, this subsection considers the unique solvability of the following weak elliptic problem: Find $u \in \hat{H}_q^1(\mathbf{R}^N)$ such that

$$(4.45) \quad (\rho^{-1} \nabla u, \nabla \varphi)_{\mathbf{R}^N \setminus \tilde{\Sigma}} = (\mathbf{f}, \nabla \varphi)_{\mathbf{R}^N \setminus \tilde{\Sigma}} \quad \text{for any } \varphi \in \hat{H}_{q'}^1(\mathbf{R}^N),$$

where $q' = q/(q-1)$ and $\rho = \rho_+ \mathbf{1}_{\tilde{\Omega}_+} + \rho_- \mathbf{1}_{\tilde{\Omega}_-}$ for positive constants ρ_\pm . In the same manner that we have obtained Theorem 3.6 from Theorem 3.2, we can obtain the following theorem from Theorem 4.3.

Theorem 4.9. *Let $r, \tilde{\Omega}_\pm$, and $\tilde{\Sigma}$ be as above, and let $q \in (1, \infty)$. Assume that $\max(q, q') \leq r$ for $q' = q/(q-1)$ and $\rho = \rho_+ \mathbf{1}_{\tilde{\Omega}_+} + \rho_- \mathbf{1}_{\tilde{\Omega}_-}$ for constants $\rho_\pm > 0$.*

(1) **Existence.** *Let $\mathbf{f} \in L_q(\mathbf{R}^N \setminus \tilde{\Sigma})^N$. Then the weak elliptic problem (4.45) admits a solution $u \in \hat{H}_q^1(\mathbf{R}^N)$ satisfying*

$$\|\nabla u\|_{L_q(\mathbf{R}^N)} \leq 2C'(\rho_+ + \rho_-)\|\mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \tilde{\Sigma})},$$

where C' is the same constant as in Theorem 4.3.

(2) **Uniqueness.** *If $u \in \hat{H}_q^1(\mathbf{R}^N)$ satisfies*

$$(\rho^{-1} \nabla u, \nabla \varphi)_{\mathbf{R}^N \setminus \tilde{\Sigma}} = 0 \quad \text{for any } \varphi \in \hat{H}_{q'}^1(\mathbf{R}^N),$$

then $u = c$ for some constant c .

5. STRONG PROBLEMS IN BOUNDED DOMAINS

Throughout this section, we assume

- Assumption 5.1.** (a) r is a real number satisfying $r > N$.
 (b) G is a bounded domain with boundary Γ of class $W_r^{2-1/r}$.
 (c) G_+ is a subdomain of G with boundary Σ of class $W_r^{2-1/r}$ satisfying $\Sigma \cap \Gamma = \emptyset$.
 (d) $G_- = G \setminus (G_+ \cup \Sigma)$.

Let \mathbf{n} be a unit normal vector on Σ pointing from G_+ into G_- , and set

$$F_q(G_\pm) = \{\mathbf{f}_\pm \in E_q(G_\pm) : \mathbf{n} \cdot \mathbf{f}_\pm \in H_q^1(G_\pm)\}.$$

In this section, we consider the following strong elliptic problem:

$$(5.1) \quad \begin{cases} \Delta v_\pm = \operatorname{div} \mathbf{f}_\pm & \text{in } G_\pm, \\ \rho_+ v_+ = \rho_- v_- & \text{on } \Sigma, \\ \mathbf{n} \cdot \nabla(v_+ - v_-) = \mathbf{n} \cdot (\mathbf{f}_+ - \mathbf{f}_-) & \text{on } \Sigma, \\ v_- = 0 & \text{on } \Gamma. \end{cases}$$

More precisely, we prove

Theorem 5.2. *Suppose that Assumption 5.1 holds and ρ_{\pm} are positive constants. Let $q \in (1, \infty)$ with $\max(q, q') \leq r$ for $q' = q/(q-1)$.*

(1) **Existence.** *Let $\mathbf{f}_{\pm} \in F_q(G_{\pm})$. Then the strong elliptic problem (5.1) admits solutions $v_{\pm} \in H_q^2(G_{\pm})$ satisfying*

$$(5.2) \quad \|\nabla^2 v_{\pm}\|_{L_q(G_{\pm})} \leq C \sum_{\mathfrak{s} \in \{+, -\}} \left(\|\mathbf{f}_{\mathfrak{s}}\|_{E_q(G_{\mathfrak{s}})} + \|\mathbf{n} \cdot \mathbf{f}_{\mathfrak{s}}\|_{H_q^1(G_{\mathfrak{s}})} \right),$$

$$(5.3) \quad \|v_{\pm}\|_{H_q^1(G_{\pm})} \leq C \sum_{\mathfrak{s} \in \{+, -\}} \|\mathbf{f}_{\mathfrak{s}}\|_{L_q(G_{\mathfrak{s}})},$$

with a positive constant $C = C(N, q, r, \rho_+, \rho_-)$. Additionally, if $\mathbf{n} \cdot (\mathbf{f}_+ - \mathbf{f}_-) = 0$ on Σ , then v_{\pm} satisfy

$$(5.4) \quad \|\nabla^2 v_{\pm}\|_{L_q(G_{\pm})} \leq C \sum_{\mathfrak{s} \in \{+, -\}} \|\mathbf{f}_{\mathfrak{s}}\|_{E_q(G_{\mathfrak{s}})}.$$

(2) **Uniqueness.** *If $v_{\pm} \in H_q^2(G_{\pm})$ satisfy*

$$\Delta v_{\pm} = 0 \text{ in } G_{\pm}; \quad \rho_+ v_+ = \rho_- v_-, \quad \mathbf{n} \cdot \nabla(v_+ - v_-) = 0 \text{ on } \Sigma; \quad v_- = 0 \text{ on } \Gamma,$$

then $v_{\pm} = 0$.

5.1. **Strong elliptic problem with λ in G_{\pm} .** To prove Theorem 5.2, we consider

$$(5.5) \quad \begin{cases} \rho_{\pm} \lambda v_{\pm} - \Delta v_{\pm} = -\operatorname{div} \mathbf{f}_{\pm} + g_{\pm} & \text{in } G_{\pm}, \\ \rho_+ v_+ = \rho_- v_- & \text{on } \Sigma, \\ \mathbf{n} \cdot \nabla(v_+ - v_-) = \mathbf{n} \cdot (\mathbf{f}_+ - \mathbf{f}_-) + h_+ - h_- & \text{on } \Sigma, \\ v_- = 0 & \text{on } \Gamma. \end{cases}$$

Note that [24, Appendix B] already studies the strong elliptic problem in \mathbf{R}_+^N with the Dirichlet boundary condition in the case where λ is taken into account, so that resolvent estimates of solutions in the bent half-space case are also available similarly to the proof of Theorem 4.2. Combining that result with Theorem 4.2 and the standard localization technique yields

Theorem 5.3. *Suppose that the same assumption as in Theorem 5.2 holds, and let $\sigma \in (0, \pi/2)$. Then there is a constant $\lambda_2 \geq 1$ such that, for any $\lambda \in \Sigma_{\sigma, \lambda_2}$,*

$$g_{\pm} \in L_q(G_{\pm}), \quad \mathbf{f}_{\pm} \in F_q(G_{\pm}), \quad h_{\pm} \in H_q^1(G_{\pm}),$$

the strong elliptic problem (5.5) admits unique solutions $v_{\pm} \in H_q^2(G_{\pm})$. In addition, the solutions v_{\pm} satisfy

$$\begin{aligned} & \|(\lambda v_{\pm}, \lambda^{1/2} \nabla v_{\pm}, \nabla^2 v_{\pm})\|_{L_q(G_{\pm})} \\ & \leq C \sum_{\mathfrak{s} \in \{+, -\}} \|(\operatorname{div} \mathbf{f}_{\mathfrak{s}}, g_{\mathfrak{s}}, \lambda^{1/2} (\mathbf{n} \cdot \mathbf{f}_{\mathfrak{s}}), \nabla (\mathbf{n} \cdot \mathbf{f}_{\mathfrak{s}}), \lambda^{1/2} h_{\mathfrak{s}}, \nabla h_{\mathfrak{s}})\|_{L_q(G_{\mathfrak{s}})}, \end{aligned}$$

and also

$$\begin{aligned} \|(\lambda^{1/2} v_{\pm}, \nabla v_{\pm})\|_{L_q(G_{\pm})} & \leq C \sum_{\mathfrak{s} \in \{+, -\}} \left(\|\mathbf{f}_{\mathfrak{s}}\|_{L_q(\Omega_{\mathfrak{s}})} + |\lambda|^{-1/2} \|g_{\mathfrak{s}}\|_{L_q(G_{\mathfrak{s}})} \right. \\ & \quad \left. + \|h_{\mathfrak{s}}\|_{L_q(G_{\mathfrak{s}})} + |\lambda|^{-1/2} \|\nabla h_{\mathfrak{s}}\|_{L_q(G_{\mathfrak{s}})} \right), \end{aligned}$$

where $C = C(N, q, r, \rho_+, \rho_-, \sigma)$ is a positive constant independent of λ . Additionally, if $\mathbf{n} \cdot (\mathbf{f}_+ - \mathbf{f}_-) = 0$ on Σ , then v_\pm satisfy

$$\begin{aligned} & \|(\lambda v_\pm, \lambda^{1/2} \nabla v_\pm, \nabla^2 v_\pm)\|_{L_q(G_\pm)} \\ & \leq C \sum_{\mathfrak{s} \in \{+, -\}} \|(\operatorname{div} \mathbf{f}_\mathfrak{s}, g_\mathfrak{s}, \lambda^{1/2} h_\mathfrak{s}, \nabla h_\mathfrak{s})\|_{L_q(G_\mathfrak{s})}. \end{aligned}$$

5.2. Proof of Theorem 5.2. We start with the following strong elliptic problem:

$$(5.6) \quad \begin{cases} -\Delta v_\pm = d_\pm & \text{in } G_\pm, \\ \rho_+ v_+ = \rho_- v_- & \text{on } \Sigma, \\ \mathbf{n} \cdot \nabla(v_+ - v_-) = 0 & \text{on } \Sigma, \\ v_- = 0 & \text{on } \Gamma. \end{cases}$$

Concerning this problem, we have

Theorem 5.4. *Suppose that the same assumption as in Theorem 5.2 holds. Then, for any $d_\pm \in L_q(G_\pm)$, the strong elliptic problem (5.6) admits unique solutions $v_\pm \in H_q^2(G_\pm)$, which satisfy*

$$\sum_{\mathfrak{s} \in \{+, -\}} \|v_\mathfrak{s}\|_{H_q^2(G_\mathfrak{s})} \leq C \sum_{\mathfrak{s} \in \{+, -\}} \|d_\mathfrak{s}\|_{L_q(G_\mathfrak{s})}$$

for some positive constant $C = C(N, q, r, \rho_+, \rho_-)$.

Proof. The proof is based on the Riesz-Schauder theory together with Theorem 5.3, so that the detailed proof may be omitted. \square

Now we prove Theorem 5.2. Let $\mu = 2\lambda_2$ for λ_2 introduced in Theorem 5.3. The Theorem 5.3 yields $w_\pm \in H_q^2(G_\pm)$ such that

$$\begin{cases} \rho_\pm \mu w_\pm - \Delta w_\pm = -\operatorname{div} \mathbf{f}_\pm & \text{in } G_\pm, \\ \rho_+ w_+ = \rho_- w_- & \text{on } \Sigma, \\ \mathbf{n} \cdot \nabla(w_+ - w_-) = \mathbf{n} \cdot (\mathbf{f}_+ - \mathbf{f}_-) & \text{on } \Sigma, \\ w_- = 0 & \text{on } \Gamma, \end{cases}$$

while Theorem 5.4 yields $z_\pm \in H_q^2(G_\pm)$ such that

$$\begin{cases} -\Delta z_\pm = \rho_\pm \mu w_\pm & \text{in } G_\pm, \\ \rho_+ z_+ = \rho_- z_- & \text{on } \Sigma, \\ \mathbf{n} \cdot \nabla(z_+ - z_-) = 0 & \text{on } \Sigma, \\ z_- = 0 & \text{on } \Gamma. \end{cases}$$

Thus $v_\pm = w_\pm + z_\pm$ become solutions to (5.1) and satisfy (5.2)-(5.4) by Theorems 5.3 and 5.4. The uniqueness of solutions of (5.1) is already proved in Theorem 5.4. This completes the proof of Theorem 5.2.

6. PROOF OF THEOREMS 1.2 AND 1.3

Throughout this section, we assume that Ω_\pm satisfy Assumption 1.1.

First, let us introduce the notation used in this section. For $L > 0$, we set

$$B_L = \{x \in \mathbf{R}^N : |x| < L\}, \quad S_L = \{x \in \mathbf{R}^N : |x| = L\}.$$

Fix $R > 0$ such that $\Omega_+ \cup \Sigma \subset B_R$. We then define G , Γ , and G_\pm as follows:

$$G = B_{4R}, \quad \Gamma = S_{4R}, \quad G_+ = \Omega_+, \quad G_- = B_{4R} \setminus (\Omega_+ \cup \Sigma).$$

Let φ , ψ_0 , and ψ_∞ be functions in $C^\infty(\mathbf{R}^N)$ satisfying $0 \leq \varphi, \psi_0, \psi_\infty \leq 1$ and

$$\begin{aligned} \varphi(x) &= \begin{cases} 1 & \text{for } x \in B_{2R}, \\ 0 & \text{for } x \in \mathbf{R}^N \setminus B_{3R}, \end{cases} \\ \psi_0(x) &= \begin{cases} 1 & \text{for } x \in B_{(3+1/3)R}, \\ 0 & \text{for } x \in \mathbf{R}^N \setminus B_{(3+2/3)R}, \end{cases} \\ \psi_\infty(x) &= \begin{cases} 0 & \text{for } x \in B_{(2-2/3)R}, \\ 1 & \text{for } x \in \mathbf{R}^N \setminus B_{(2-1/3)R}. \end{cases} \end{aligned}$$

In addition, we set $\varphi_0(x) = \varphi(x)$ and $\varphi_\infty(x) = 1 - \varphi(x)$. For $q \in (1, \infty)$ and $D \in \{\mathbf{R}^N, G\}$, a Banach space $E_q(D \setminus \Sigma)$ is defined by

$$\begin{aligned} E_q(D \setminus \Sigma) &= \{\mathbf{f} \in L_q(D \setminus \Sigma)^N : \mathbf{f}|_{D_+} \in E_q(D_+) \text{ and } \mathbf{f}|_{D_-} \in E_q(D_-)\}, \\ \|\mathbf{f}\|_{E_q(D \setminus \Sigma)} &= \|\mathbf{f}|_{D_+}\|_{E_q(D_+)} + \|\mathbf{f}|_{D_-}\|_{E_q(D_-)}, \end{aligned}$$

where $\mathbf{f}|_{D_\pm}$ stands for the restriction of \mathbf{f} on D_\pm , respectively. On the other hand,

$$\begin{aligned} F_q(D \setminus \Sigma) &= \{\mathbf{f} \in E_q(D \setminus \Sigma) : \mathbf{n} \cdot \mathbf{f}|_{D_+} \in H_q^1(D_+) \text{ and } \mathbf{n} \cdot \mathbf{f}|_{D_-} \in H_q^1(D_-)\}, \\ \|\mathbf{f}\|_{F_q(D \setminus \Sigma)} &= \|\mathbf{f}\|_{E_q(D \setminus \Sigma)} + \|\mathbf{n} \cdot \mathbf{f}|_{D_+}\|_{H_q^1(D_+)} + \|\mathbf{n} \cdot \mathbf{f}|_{D_-}\|_{H_q^1(D_-)}. \end{aligned}$$

Remark 6.1. One sees that $\mathbf{f} \in F_q(D \setminus \Sigma)$ is equivalent to $\mathbf{f} = \mathbf{f}_+ \mathbf{1}_{D_+} + \mathbf{f}_- \mathbf{1}_{D_-}$ for some $\mathbf{f}_\pm \in E_q(D_\pm)$ with $\mathbf{n} \cdot \mathbf{f}_\pm \in H_q^1(\Omega_\pm)$.

The aim of this section is to prove Theorems 1.2 and 1.3. To this end, it suffices to prove the existence of solutions of (1.5) satisfying (1.6)-(1.8).⁴ Instead of (1.5), we consider the following equivalent system of (1.5) for simplicity of notation:

$$(6.1) \quad \begin{cases} \Delta v = \operatorname{div} \mathbf{f} & \text{in } \mathbf{R}^N \setminus \Sigma, \\ \llbracket \rho v \rrbracket = 0 & \text{on } \Sigma, \\ \llbracket \mathbf{n} \cdot \nabla v \rrbracket = \llbracket \mathbf{n} \cdot \mathbf{f} \rrbracket & \text{on } \Sigma, \end{cases}$$

where $\rho = \rho_+ \mathbf{1}_{\Omega_+} + \rho_- \mathbf{1}_{\Omega_-}$ for positive constants ρ_\pm and

$$\llbracket f \rrbracket(x_0) = \lim_{x \rightarrow x_0, x \in \Omega_+} f(x) - \lim_{x \rightarrow x_0, x \in \Omega_-} f(x) \quad (x_0 \in \Sigma).$$

We here recall the assumption for $q \in (1, \infty)$.

Assumption 6.2. Let $q \in (1, \infty)$ and $\max(q, q') \leq r$ for $q' = q/(q-1)$.

As mentioned above, it suffices to prove in this section

Theorem 6.3. Suppose that Assumption 6.2 holds. Then, for any $\mathbf{f} \in F_q(\mathbf{R}^N \setminus \Sigma)$, the system (6.1) admits a solution $v \in \widehat{H}_q^1(\mathbf{R}^N \setminus \Sigma) \cap \widehat{H}_q^2(\mathbf{R}^N \setminus \Sigma)$ satisfying

$$(6.2) \quad \|\nabla^2 v\|_{L_q(\mathbf{R}^N \setminus \Sigma)} \leq C \|\mathbf{f}\|_{F_q(\mathbf{R}^N \setminus \Sigma)}, \quad \|\nabla v\|_{L_q(\mathbf{R}^N \setminus \Sigma)} \leq C \|\mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \Sigma)},$$

⁴ The uniqueness of solutions of (1.5) follows from the existence of solutions of (1.5) similarly to the proof of Theorem 3.2 (2), while Theorem 1.3 follows from Theorem 1.2 as Theorem 3.6 has followed from Theorem 3.2.

with some positive constant $C = C(N, q, r, \rho_+, \rho_-)$. Additionally, if $\mathbf{n} \cdot (\mathbf{f}_+ - \mathbf{f}_-) = 0$ on Σ , then v satisfies

$$(6.3) \quad \|\nabla^2 v\|_{L_q(\mathbf{R}^N \setminus \Sigma)} \leq C \|\mathbf{f}\|_{E_q(\mathbf{R}^N \setminus \Sigma)}$$

for some positive constant $C = C(R, N, q, r, \rho_+, \rho_-)$.

6.1. Solution operators. First, let us consider the following problem in the whole space:

$$(6.4) \quad \Delta V = \operatorname{div} \mathbf{F} \quad \text{in } \mathbf{R}^N.$$

Similarly to [24, Page 1700], we obtain

Lemma 6.4. *Let $q \in (1, \infty)$. Then there exists a linear operator*

$$\mathcal{S}_\infty : E_q(\mathbf{R}^N) \rightarrow \widehat{H}_q^1(\mathbf{R}^N) \cap \widehat{H}_q^2(\mathbf{R}^N)$$

such that $V = \mathcal{S}_\infty \mathbf{F}$ is a solution to (6.4). In addition,

$$\begin{aligned} \|\mathcal{S}_\infty \mathbf{F}\|_{L_q(G)} + \|\nabla \mathcal{S}_\infty \mathbf{F}\|_{L_q(\mathbf{R}^N)} &\leq C \|\mathbf{F}\|_{L_q(\mathbf{R}^N)}, \\ \|\nabla^2 \mathcal{S}_\infty \mathbf{F}\|_{L_q(\mathbf{R}^N)} &\leq C \|\operatorname{div} \mathbf{F}\|_{L_q(\mathbf{R}^N)}, \end{aligned}$$

with some positive constant $C = C(R, N, q)$.

We next consider the following problem in a bounded domain:

$$(6.5) \quad \begin{cases} \Delta V = \operatorname{div} \mathbf{F} & \text{in } G \setminus \Sigma, \\ [\rho V] = 0 & \text{on } \Sigma, \\ [\mathbf{n} \cdot \nabla V] = [\mathbf{n} \cdot \mathbf{F}] & \text{on } \Sigma, \\ V = 0 & \text{on } \Gamma. \end{cases}$$

Then, by Theorem 5.2, we have

Lemma 6.5. *Suppose that Assumption 6.2 holds. Then there is a linear operator*

$$\mathcal{S}_0 : F_q(G \setminus \Sigma) \rightarrow H_q^2(G \setminus \Sigma)$$

such that, for any $\mathbf{F} \in F_q(G \setminus \Sigma)$, $V = \mathcal{S}_0 \mathbf{F}$ is a solution to (6.5). In addition,

$$\|\nabla^2 \mathcal{S}_0 \mathbf{F}\|_{L_q(G \setminus \Sigma)} \leq C \|\mathbf{F}\|_{F_q(G \setminus \Sigma)}, \quad \|\mathcal{S}_0 \mathbf{F}\|_{H_q^1(G \setminus \Sigma)} \leq C \|\mathbf{F}\|_{L_q(G \setminus \Sigma)},$$

with some positive constant $C = C(N, q, r, \rho_+, \rho_-)$. Additionally, if $[\mathbf{n} \cdot \mathbf{F}] = 0$ on Σ , then it holds that

$$\|\nabla^2 \mathcal{S}_0 \mathbf{F}\|_{L_q(G \setminus \Sigma)} \leq C \|\mathbf{F}\|_{E_q(G \setminus \Sigma)}$$

for some positive constant $C = C(N, q, r, \rho_+, \rho_-)$.

Let us define an operator $\mathcal{S} : F_q(\mathbf{R}^N \setminus \Sigma) \rightarrow \widehat{H}_q^1(\mathbf{R}^N \setminus \Sigma) \cap \widehat{H}_q^2(\mathbf{R}^N \setminus \Sigma)$ by

$$(6.6) \quad \mathcal{S} \mathbf{f} = \psi_\infty \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) + \psi_0 \mathcal{S}_0(\varphi_0 \mathbf{f}) \quad (\mathbf{f} \in F_q(\mathbf{R}^N \setminus \Sigma)).$$

Note that $\varphi_\infty \mathbf{f} \in E_q(\mathbf{R}^N)$ and $\varphi_0 \mathbf{f} \in F_q(G \setminus \Sigma)$ when $\mathbf{f} \in F_q(\mathbf{R}^N \setminus \Sigma)$. By Lemmas 6.4 and 6.5, the above $\mathcal{S} \mathbf{f}$ satisfies the following estimates:

$$(6.7) \quad \|\nabla^2 \mathcal{S} \mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \Sigma)} \leq C \|\mathbf{f}\|_{F_q(\mathbf{R}^N \setminus \Sigma)}, \quad \|\nabla \mathcal{S} \mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \Sigma)} \leq C \|\mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \Sigma)}.$$

Additionally, if $\mathbf{n} \cdot (\mathbf{f}_+ - \mathbf{f}_-) = 0$ on Σ , then it holds that

$$(6.8) \quad \|\nabla^2 \mathcal{S} \mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \Sigma)} \leq C \|\mathbf{f}\|_{E_q(\mathbf{R}^N \setminus \Sigma)}.$$

At this point, we introduce a function space $\mathcal{L}_q(\mathbf{R}^N)$ for $q \in (1, \infty)$. Let $R_1 = (2 - 2/3)R$ and $R_2 = (3 + 2/3)R$, and then

$$\mathcal{L}_q(\mathbf{R}^N) = \{f \in L_q(\mathbf{R}^N) : \text{supp } f \subset D_{R_1, R_2}, (f, 1)_{\mathbf{R}^N} = 0\}$$

and $\|f\|_{\mathcal{L}_q(\mathbf{R}^N)} = \|f\|_{L_q(\mathbf{R}^N)}$, where $D_{R_1, R_2} = \{x \in \mathbf{R}^N : R_1 \leq |x| \leq R_2\}$.

Now we consider for $f \in \mathcal{L}_q(\mathbf{R}^N)$ the following problem in the whole space:

$$(6.9) \quad \Delta v = f \quad \text{in } \mathbf{R}^N.$$

For this problem, we have

Lemma 6.6. *Let $q \in (1, \infty)$. Then there exists a linear operator*

$$\tilde{\mathcal{T}}_\infty : \mathcal{L}_q(\mathbf{R}^N) \rightarrow \hat{H}_q^1(\mathbf{R}^N) \cap \hat{H}_q^2(\mathbf{R}^N)$$

such that, for any $f \in \mathcal{L}_q(\mathbf{R}^N)$, $v = \tilde{\mathcal{T}}_\infty f$ is a solution to (6.9). In addition,

$$\begin{aligned} \|\tilde{\mathcal{T}}_\infty f\|_{L_q(G)} + \sup_{|x| \geq 4R} |x|^{N-1} |[\tilde{\mathcal{T}}_\infty f](x)| &\leq C \|f\|_{\mathcal{L}_q(\mathbf{R}^N)}, \\ \|\nabla \tilde{\mathcal{T}}_\infty f\|_{L_q(\mathbf{R}^N)} + \sup_{|x| \geq 4R} |x|^N |\nabla[\tilde{\mathcal{T}}_\infty f](x)| &\leq C \|f\|_{\mathcal{L}_q(\mathbf{R}^N)}, \\ \|\nabla^2 \tilde{\mathcal{T}}_\infty f\|_{L_q(\mathbf{R}^N)} &\leq C \|f\|_{\mathcal{L}_q(\mathbf{R}^N)}, \end{aligned}$$

with some positive constant $C = C(R, N, q)$.

Proof. See e.g. [24, pp.1703–1704] and [30, Lemma 1]. □

Next, we consider the following problem in a bounded domain:

$$(6.10) \quad \begin{cases} \Delta v = f & \text{in } G \setminus \Sigma, \\ \llbracket \rho v \rrbracket = 0 & \text{on } \Sigma, \\ \llbracket \mathbf{n} \cdot \nabla v \rrbracket = 0 & \text{on } \Sigma, \\ v = 0 & \text{on } \Gamma. \end{cases}$$

By Theorem 5.4, we have

Lemma 6.7. *Suppose that Assumption 6.2 holds. Then there is a linear operator*

$$\mathcal{T}_0 : L_q(G \setminus \Sigma) \rightarrow H_q^2(G \setminus \Sigma)$$

such that, for any $f \in L_q(G \setminus \Sigma)$, $v = \mathcal{T}_0 f$ is a solution to (6.10). In addition,

$$\|\mathcal{T}_0 f\|_{H_q^2(G \setminus \Sigma)} \leq C \|f\|_{L_q(G \setminus \Sigma)}$$

for some positive constant $C = C(N, q, r, \rho_+, \rho_-)$.

Finally, we introduce $\mathcal{T} : \mathcal{L}_q(\mathbf{R}^N) \rightarrow \hat{H}_q^1(\mathbf{R}^N \setminus \Sigma) \cap \hat{H}_q^2(\mathbf{R}^N \setminus \Sigma)$ as follows: For $f \in \mathcal{L}_q(\mathbf{R}^N)$, we choose a constant c_f so that

$$(6.11) \quad \int_{D_{R_3, R_4}} (\tilde{\mathcal{T}}_\infty f + c_f) dx = 0,$$

where $D_{R_3, R_4} = \{x \in \mathbf{R}^N : R_3 \leq |x| \leq R_4\}$ for $R_3 = (3 + 1/3)R$ and $R_4 = (4 - 1/3)R$. Let us define

$$(6.12) \quad \mathcal{T}_\infty f = \tilde{\mathcal{T}}_\infty f + c_f, \quad \mathcal{T} f = \varphi_\infty \mathcal{T}_\infty f + \varphi_0 \mathcal{T}_0 f.$$

We then have by (6.11)

$$(6.13) \quad \int_{D_{R_3, R_4}} \mathcal{T}f \, dx = \int_{D_{R_3, R_4}} \mathcal{T}_\infty f \, dx = 0.$$

In addition, since $c_f = -|D_{R_3, R_4}|^{-1} \int_{D_{R_3, R_4}} \tilde{\mathcal{T}}_\infty f \, dx$, it holds by Lemma 6.6 that

$$|c_f| \leq \frac{1}{|D_{R_3, R_4}|} \int_{D_{R_3, R_4}} |\tilde{\mathcal{T}}_\infty f| \, dx \leq C \|\tilde{\mathcal{T}}_\infty f\|_{L_q(G)} \leq C \|f\|_{\mathcal{L}_q(\mathbf{R}^N)}.$$

Thus $\mathcal{T}f$ satisfies, together with Lemmas 6.6 and 6.7, the following estimate:

$$(6.14) \quad \|\nabla \mathcal{T}f\|_{L_q(\mathbf{R}^N \setminus \Sigma)} + \|\nabla^2 \mathcal{T}f\|_{L_q(\mathbf{R}^N \setminus \Sigma)} \leq C \|f\|_{\mathcal{L}_q(\mathbf{R}^N)}.$$

6.2. An auxiliary problem. In this subsection, we consider for $f \in \mathcal{L}_q(\mathbf{R}^N)$ the following auxiliary problem:

$$(6.15) \quad \begin{cases} \Delta v = f & \text{in } \mathbf{R}^N \setminus \Sigma, \\ \llbracket \rho v \rrbracket = 0 & \text{on } \Sigma, \\ \llbracket \mathbf{n} \cdot \nabla v \rrbracket = 0 & \text{on } \Sigma. \end{cases}$$

Concerning this system, we prove

Lemma 6.8. *Suppose that Assumption 6.2 holds. Let $q \in [2, \infty)$ additionally and $f \in \mathcal{L}_q(\mathbf{R}^N)$. Then (6.15) admits a solution $v \in \hat{H}_q^1(\mathbf{R}^N \setminus \Sigma) \cap \hat{H}_q^2(\mathbf{R}^N \setminus \Sigma)$ satisfying*

$$(6.16) \quad \|\nabla v\|_{L_q(\mathbf{R}^N \setminus \Sigma)} + \|\nabla^2 v\|_{L_q(\mathbf{R}^N \setminus \Sigma)} \leq C \|f\|_{\mathcal{L}_q(\mathbf{R}^N)}$$

for some positive constant $C = C(R, N, q, r, \rho_+, \rho_-)$.

Proof. Let \mathcal{T} be the operator defined as (6.12). Then, in $\mathbf{R}^N \setminus \Sigma$,

$$\begin{aligned} \Delta \mathcal{T}f &= \varphi_\infty \Delta \mathcal{T}_\infty f + 2\nabla \varphi_\infty \cdot \nabla \mathcal{T}_\infty f + (\Delta \varphi_\infty) \mathcal{T}_\infty f \\ &\quad + \varphi_0 \Delta \mathcal{T}_0 f + 2\nabla \varphi_0 \cdot \nabla \mathcal{T}_0 f + (\Delta \varphi_0) \mathcal{T}_0 f. \end{aligned}$$

Since

$$\varphi_\infty \Delta \mathcal{T}_\infty f = \varphi_\infty f, \quad \varphi_0 \Delta \mathcal{T}_0 f = \varphi_0 f, \quad \varphi_0 + \varphi_\infty = 1,$$

it holds that

$$\Delta \mathcal{T}f = f + \mathcal{G}f \quad \text{in } \mathbf{R}^N \setminus \Sigma,$$

where

$$(6.17) \quad \mathcal{G}f = 2\nabla \varphi_\infty \cdot \nabla \mathcal{T}_\infty f + (\Delta \varphi_\infty) \mathcal{T}_\infty f + 2\nabla \varphi_0 \cdot \nabla \mathcal{T}_0 f + (\Delta \varphi_0) \mathcal{T}_0 f.$$

Thus we have achieved

$$(6.18) \quad \begin{cases} \Delta \mathcal{T}f = f + \mathcal{G}f & \text{in } \mathbf{R}^N \setminus \Sigma, \\ \llbracket \rho \mathcal{T}f \rrbracket = 0 & \text{on } \Sigma, \\ \llbracket \mathbf{n} \cdot \nabla \mathcal{T}f \rrbracket = 0 & \text{on } \Sigma. \end{cases}$$

The following lemma is proved in the appendix A below.

Lemma 6.9. *Suppose that Assumption 6.2 holds.*

- (1) *Then \mathcal{G} is a compact operator on $\mathcal{L}_q(\mathbf{R}^N)$.*
- (2) *Let $q \in [2, \infty)$ additionally. Then $(I + \mathcal{G})^{-1}$ exists in $\mathcal{L}(\mathcal{L}_q(\mathbf{R}^N))$.*

Setting $v = \mathcal{T}(I + \mathcal{G})^{-1}f$ for $f \in \mathcal{L}_q(\mathbf{R}^N)$, we see that v is a solution to (6.15) by (6.18) and satisfies (6.16) by (6.14). This completes the proof of Lemma 6.8 \square

6.3. Proof of Theorem 6.3. Let \mathcal{S} be the operator defined as (6.6). Then we observe that in $\mathbf{R}^N \setminus \Sigma$

$$\begin{aligned} \Delta \mathbf{Sf} &= \psi_\infty \Delta \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) + 2\nabla \psi_\infty \cdot \nabla \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) + (\Delta \psi_\infty) \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) \\ &\quad + \psi_0 \Delta \mathcal{S}_0(\varphi_0 \mathbf{f}) + 2\nabla \psi_0 \cdot \nabla \mathcal{S}_0(\varphi_0 \mathbf{f}) + (\Delta \psi_0) \mathcal{S}_0(\varphi_0 \mathbf{f}). \end{aligned}$$

Note that in $\mathbf{R}^N \setminus \Sigma$

$$\psi_\infty \Delta \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) + \psi_0 \Delta \mathcal{S}_0(\varphi_0 \mathbf{f}) = \psi_\infty \operatorname{div}(\varphi_\infty \mathbf{f}) + \psi_0 \operatorname{div}(\varphi_0 \mathbf{f}) = \operatorname{div} \mathbf{f}.$$

This relation implies

$$\Delta \mathbf{Sf} = \operatorname{div} \mathbf{f} + \mathcal{Rf} \quad \text{in } \mathbf{R}^N \setminus \Sigma,$$

where

$$(6.19) \quad \begin{aligned} \mathcal{Rf} &= 2\nabla \psi_\infty \cdot \nabla \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) + (\Delta \psi_\infty) \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) \\ &\quad + 2\nabla \psi_0 \cdot \nabla \mathcal{S}_0(\varphi_0 \mathbf{f}) + (\Delta \psi_0) \mathcal{S}_0(\varphi_0 \mathbf{f}). \end{aligned}$$

On the other hand,

$$[[\rho \mathbf{Sf}]] = 0, \quad [[\mathbf{n} \cdot \nabla \mathbf{Sf}]] = [[\mathbf{n} \cdot \mathbf{f}]] \quad \text{on } \Sigma,$$

and also one has⁵ for $\mathbf{f} \in F_q(\mathbf{R}^N \setminus \Sigma)$

$$(6.20) \quad \mathcal{Rf} \in \mathcal{L}_q(\mathbf{R}^N), \quad \|\mathcal{Rf}\|_{\mathcal{L}_q(\mathbf{R}^N)} \leq C \|\mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \Sigma)}.$$

Next, we consider

$$(6.21) \quad \begin{cases} \Delta w = -\mathcal{Rf} & \text{in } \mathbf{R}^N \setminus \Sigma, \\ [[\rho w]] = 0 & \text{on } \Sigma \\ [[\mathbf{n} \cdot \nabla w]] = 0 & \text{on } \Sigma. \end{cases}$$

In the following, the discussion of w is divided into two cases.

Case 1: $q \in [2, \infty)$. By Lemma 6.8 and (6.20), the system (6.21) admits a solution $w \in \widehat{H}_q^1(\mathbf{R}^N \setminus \Sigma) \cap \widehat{H}_q^2(\mathbf{R}^N \setminus \Sigma)$ satisfying

$$(6.22) \quad \|\nabla w\|_{L_q(\mathbf{R}^N \setminus \Sigma)} + \|\nabla^2 w\|_{L_q(\mathbf{R}^N \setminus \Sigma)} \leq C \|\mathbf{f}\|_{L_q(\mathbf{R}^N \setminus \Sigma)}.$$

Thus $v = \mathbf{Sf} + w$ solves (6.1) and satisfies (6.2)-(6.3) by (6.7), (6.8), and (6.22). This completes the proof of Theorem 6.3 for $q \in [2, \infty)$.

Case 2: $q \in (1, 2)$. Since Theorem 6.3 is already proved for $q \in [2, \infty)$ in Case 1 as above, one can prove by the result of Case 1 the following lemma.

Lemma 6.10. *Suppose that Assumption 6.2 holds. Let $q \in (1, 2)$ additionally and $f \in \mathcal{L}_q(\mathbf{R}^N)$. Then (6.15) admits a solution $v \in \widehat{H}_q^1(\mathbf{R}^N \setminus \Sigma) \cap \widehat{H}_q^2(\mathbf{R}^N \setminus \Sigma)$ satisfying*

$$\|\nabla v\|_{L_q(\mathbf{R}^N \setminus \Sigma)} + \|\nabla^2 v\|_{L_q(\mathbf{R}^N \setminus \Sigma)} \leq C \|f\|_{L_q(\mathbf{R}^N)}$$

for some positive constant $C = C(R, N, q, r, \rho_+, \rho_-)$.

Proof. See the appendix C below. □

By Lemma 6.10 and (6.20), the system (6.21) admits a solution $w \in \widehat{H}_q^1(\mathbf{R}^N \setminus \Sigma) \cap \widehat{H}_q^2(\mathbf{R}^N \setminus \Sigma)$ satisfying (6.22). Thus $v = \mathbf{Sf} + w$ solves (6.1) and satisfies (6.2)-(6.3) by (6.7), (6.8), and (6.22). This completes the proof of Theorem 6.3 for $q \in (1, 2)$, which furnishes the proof of Theorem 6.3.

⁵The first property of (6.20) is proved in the appendix B below, while the second property of (6.20) follows from Lemmas 6.4 and 6.5 immediately.

A.

This appendix proves Lemma 6.9 for \mathcal{G} defined as (6.17). We start with

Lemma A.1. *For q satisfying Assumption 6.2, \mathcal{G} is a compact operator on $\mathcal{L}_q(\mathbf{R}^N)$.*

Proof. Step 1: $\mathcal{G}f \in \mathcal{L}_q(\mathbf{R}^N)$ for any $f \in \mathcal{L}_q(\mathbf{R}^N)$. It is clear that

$$(A.1) \quad \text{supp } \mathcal{G}f \subset D_{R_1, R_2}.$$

In addition, by $\nabla\varphi_\infty = -\nabla\varphi_0$ and $\Delta\varphi_\infty = -\Delta\varphi_0$,

$$(A.2) \quad \begin{aligned} \mathcal{G}f &= -2\nabla\varphi_0 \cdot \nabla\mathcal{T}_\infty f - (\Delta\varphi_0)\mathcal{T}_\infty f + 2\nabla\varphi_0 \cdot \nabla\mathcal{T}_0 f + (\Delta\varphi_0)\mathcal{T}_0 f \\ &= \varphi_0\Delta\mathcal{T}_\infty f - \text{div}((\nabla\varphi_0)\mathcal{T}_\infty f + \varphi_0\nabla\mathcal{T}_\infty f) \\ &\quad - \varphi_0\Delta\mathcal{T}_0 f + \text{div}((\nabla\varphi_0)\mathcal{T}_0 f + \varphi_0\nabla\mathcal{T}_0 f) \quad \text{in } \mathbf{R}^N \setminus \Sigma, \end{aligned}$$

which, combined with $\varphi_0\Delta\mathcal{T}_\infty f = \varphi_0 f$ and $\varphi_0\Delta\mathcal{T}_0 f = \varphi_0 f$ in $\mathbf{R}^N \setminus \Sigma$, furnishes

$$\begin{aligned} \mathcal{G}f &= -\text{div}((\nabla\varphi_0)\mathcal{T}_\infty f + \varphi_0\nabla\mathcal{T}_\infty f) \\ &\quad + \text{div}((\nabla\varphi_0)\mathcal{T}_0 f + \varphi_0\nabla\mathcal{T}_0 f) \quad \text{in } \mathbf{R}^N \setminus \Sigma. \end{aligned}$$

It then holds that

$$\begin{aligned} (\mathcal{G}f, 1)_{B_{4R} \setminus \Sigma} &= (\text{div}((\nabla\varphi_0)\mathcal{T}_\infty f + \varphi_0\nabla\mathcal{T}_\infty f), 1)_{B_{4R} \setminus \Sigma} \\ &\quad + (\text{div}((\nabla\varphi_0)\mathcal{T}_0 f), 1)_{B_{4R} \setminus \Sigma} + (\text{div}(\varphi_0\nabla\mathcal{T}_0 f), 1)_{B_{4R} \setminus \Sigma} \\ &=: I_1 + I_2 + I_3. \end{aligned}$$

By Gauss's divergence theorem, we see that

$$I_1 = I_2 = 0, \quad I_3 = \int_{\Sigma} [\mathbf{n} \cdot \nabla\mathcal{T}_0 f] d\sigma = 0,$$

where $d\sigma$ is the surface element of Σ . Hence $(\mathcal{G}f, 1)_{\mathbf{R}^N} = (\mathcal{G}f, 1)_{B_{4R} \setminus \Sigma} = 0$, which, combined with (A.1), implies $\mathcal{G}f \in \mathcal{L}_q(\mathbf{R}^N)$.

Step 2: \mathcal{G} is a compact operator on $\mathcal{L}_q(\mathbf{R}^N)$.

Let $\{f^{(j)}\}_{j=1}^\infty$ be a bounded sequence in $\mathcal{L}_q(\mathbf{R}^N)$, i.e. there exists a positive constant M such that $\|f^{(j)}\|_{\mathcal{L}_q(\mathbf{R}^N)} \leq M$ for any $j \in \mathbf{N}$. Note that by (6.17), Lemmas 6.6 and 6.7, and $\text{supp } \nabla\varphi_0, \text{supp } \nabla\varphi_\infty \subset \{x \in \mathbf{R}^N : 2R \leq |x| \leq 3R\}$

$$(A.3) \quad \mathcal{G} \in \mathcal{L}(\mathcal{L}_q(\mathbf{R}^N), H_q^1(\mathbf{R}^N)).$$

Since $H_q^1(\mathbf{R}^N)$ is compactly embedded into $L_q(D_{R_1, R_2})$ by the Rellich-Kondrachov theorem (cf. [3, Theorem 6.3]), \mathcal{G} can be regarded as a compact operator from $\mathcal{L}_q(\mathbf{R}^N)$ into $L_q(D_{R_1, R_2})$ by (A.3). Thus there exists $G_f \in L_q(D_{R_1, R_2})$ such that

$$(A.4) \quad \lim_{j \rightarrow \infty} \|\mathcal{G}f^{(j)} - G_f\|_{L_q(D_{R_1, R_2})} = 0,$$

up to some extraction. Let us define

$$\tilde{G}_f = \begin{cases} G_f & \text{in } D_{R_1, R_2}, \\ 0 & \text{in } \mathbf{R}^N \setminus D_{R_1, R_2}, \end{cases}$$

and then $\text{supp } \tilde{G}_f \subset D_{R_1, R_2}$. In addition, by (A.4),

$$(\tilde{G}_f, 1)_{\mathbf{R}^N} = (G_f, 1)_{D_{R_1, R_2}} = \lim_{j \rightarrow \infty} (\mathcal{G}f^{(j)}, 1)_{D_{R_1, R_2}}$$

which, combined with $\mathcal{G}f^{(j)} \in \mathcal{L}_q(\mathbf{R}^N)$ as was proved in Step 1, implies $(\tilde{G}_f, 1)_{\mathbf{R}^N} = 0$. Hence $\tilde{G}_f \in \mathcal{L}_q(\mathbf{R}^N)$. On the other hand, by (A.4) and $\mathcal{G}f^{(j)} \in \mathcal{L}_q(\mathbf{R}^N)$,

$$\lim_{j \rightarrow \infty} \|\mathcal{G}f^{(j)} - \tilde{G}_f\|_{\mathcal{L}_q(\mathbf{R}^N)} = \lim_{j \rightarrow \infty} \|\mathcal{G}f^{(j)} - G_f\|_{L_q(D_{R_1, R_2})} = 0,$$

and therefore \mathcal{G} is a compact operator on $\mathcal{L}_q(\mathbf{R}^N)$. This completes the proof. \square

Now we prove Lemma 6.9 (2). In view of Lemma A.1 and the Riesz-Schauder theory, it suffices to prove that the kernel of $I + \mathcal{G}$ is trivial in what follows. Let us begin with some property of \mathcal{T} defined in (6.12).

Lemma A.2. *Suppose that Assumption 6.2 holds, and let $f \in \mathcal{L}_q(\mathbf{R}^N)$ with $\mathcal{T}f = 0$. Then $f = 0$.*

Proof. By the assumption $\mathcal{T}f = 0$, we have

$$(A.5) \quad \mathcal{T}_\infty f = 0 \quad \text{when } |x| \geq 3R, \quad \mathcal{T}_0 f = 0 \quad \text{when } |x| \leq 2R.$$

Here we set

$$w = \begin{cases} \mathcal{T}_0 f & (x \in B_{4R} \setminus B_{(3/2)R}), \\ 0 & (x \in B_{(3/2)R}). \end{cases}$$

Then w satisfies by the second property of (A.5) and $\mathcal{T}_0 f = 0$ on Γ

$$(A.6) \quad \Delta w = f \quad \text{in } G, \quad w = 0 \quad \text{on } \Gamma.$$

On the other hand, $\mathcal{T}_\infty f$ also satisfies (A.6) by the first property of (A.5). The uniqueness of solutions of (A.6) thus implies $\mathcal{T}_0 f = \mathcal{T}_\infty f$ in G , which, combined with the assumption $\mathcal{T}f = 0$ and $\varphi_\infty = 1 - \varphi_0$, furnishes

$$0 = \mathcal{T}f = (1 - \varphi_0)\mathcal{T}_\infty f + \varphi_0 \mathcal{T}_0 f = \mathcal{T}_\infty f \quad \text{in } G.$$

Combining this property with (A.5) yields $\mathcal{T}_\infty f = 0$ in \mathbf{R}^N . Since

$$\Delta \mathcal{T}_\infty f = f \quad \text{in } \mathbf{R}^N,$$

one concludes $f = 0$. This completes the proof of the lemma. \square

Next we complete the proof of Lemma 6.9 (2) by the following result.

Lemma A.3. *Suppose that Assumption 6.2 holds. Let $q \in [2, \infty)$ additionally and $f \in \mathcal{L}_q(\mathbf{R}^N)$ with $(I + \mathcal{G})f = 0$. Then $f = 0$.*

Proof. Let $\omega(x)$ be an element of $C_0^\infty(\mathbf{R}^N)$ satisfying

$$\omega(x) = \begin{cases} 1 & (|x| \leq 1), \\ 0 & (|x| \geq 2), \end{cases}$$

and set $\omega_L(x) = \omega(x/L)$ for $L > 0$ large enough. Note that $\mathcal{T}f \in H_{q', \text{loc}}^2(\mathbf{R}^N \setminus \Sigma)$ for $q' = q/(q-1)$ by $q \in [2, \infty)$. Then, by the assumption $(I + \mathcal{G})f = 0$, (6.18), and integration by parts,

$$(A.7) \quad \begin{aligned} 0 &= ((I + \mathcal{G})f, \omega_L \rho \mathcal{T}f)_{\mathbf{R}^N \setminus \Sigma} = (\Delta \mathcal{T}f, \omega_L \rho \mathcal{T}f)_{\mathbf{R}^N \setminus \Sigma} \\ &= -(\nabla \mathcal{T}f, \omega_L \rho \nabla \mathcal{T}f)_{\mathbf{R}^N \setminus \Sigma} - (\nabla \mathcal{T}f, (\nabla \omega_L) \rho \mathcal{T}f)_{\mathbf{R}^N}. \end{aligned}$$

Since $L > 0$ is large enough and $\text{supp}(\nabla \omega)(\cdot/L) \subset D_{L, 2L} = \{x \in \mathbf{R}^N : L \leq |x| \leq 2L\}$, one sees that on $\text{supp}(\nabla \omega)(\cdot/L)$

$$\mathcal{T}f = \tilde{\mathcal{T}}_\infty f + c_f, \quad \nabla \mathcal{T}f = \nabla \tilde{\mathcal{T}}_\infty f, \quad \rho = \rho_-.$$

We thus observe that

$$\begin{aligned}
 |(\nabla \mathcal{T}f, (\nabla \omega_L) \rho \mathcal{T}f)_{\mathbf{R}^N}| &= \frac{\rho_-}{L} \left| \int_{\mathbf{R}^N} \nabla \tilde{\mathcal{T}}_\infty f \cdot (\nabla \omega) \left(\frac{x}{L} \right) (\tilde{\mathcal{T}}_\infty f + c_f) dx \right| \\
 &\leq \frac{\rho_-}{L} \int_{D_{L,2L}} |\nabla \tilde{\mathcal{T}}_\infty f| \left| (\nabla \omega) \left(\frac{x}{L} \right) \right| |\tilde{\mathcal{T}}_\infty f| dx \\
 &+ \frac{\rho_- |c_f|}{L} \int_{D_{L,2L}} |\nabla \tilde{\mathcal{T}}_\infty f| \left| (\nabla \omega) \left(\frac{x}{L} \right) \right| dx \\
 &=: I_1 + I_2.
 \end{aligned}$$

By Lemma 6.6,

$$\begin{aligned}
 I_1 &\leq \frac{C}{L} \left(\sup_{x \in \mathbf{R}^N} |\nabla \omega(x)| \right) \|f\|_{\mathcal{L}_q(\mathbf{R}^N)}^2 \int_{D_{L,2L}} |x|^{-(2N-1)} dx, \\
 I_2 &\leq \frac{C}{L} |c_f| \left(\sup_{x \in \mathbf{R}^N} |\nabla \omega(x)| \right) \|f\|_{\mathcal{L}_q(\mathbf{R}^N)} \int_{D_{L,2L}} |x|^{-N} dx.
 \end{aligned}$$

Hence $\lim_{L \rightarrow \infty} (\nabla \mathcal{T}f, (\nabla \omega_L) \rho \mathcal{T}f)_{\mathbf{R}^N} = 0$.

Now we take the limit: $L \rightarrow \infty$ in (A.7), i.e.

$$\lim_{L \rightarrow \infty} \int_{\mathbf{R}^N \setminus \Sigma} \rho(x) |[\nabla \mathcal{T}f](x)|^2 \omega \left(\frac{x}{L} \right) dx = 0.$$

Then the monotone convergence theorem yields

$$\int_{\mathbf{R}^N \setminus \Sigma} \rho(x) |[\nabla \mathcal{T}f](x)|^2 = 0.$$

Thus $\nabla \mathcal{T}f = 0$ in $\mathbf{R}^N \setminus \Sigma$, which implies that there are constants c_\pm so that $\rho \mathcal{T}f = c_\pm$ in Ω_\pm . Since $[\rho \mathcal{T}f] = 0$ on Σ , one has $c_+ = c_-$. On the other hand, it holds by (6.13) that

$$c_+ |D_{R_3, R_4}| = \int_{D_{R_3, R_4}} \rho \mathcal{T}f dx = \rho_- \int_{D_{R_3, R_4}} \mathcal{T}f dx = 0,$$

which implies $c_+ = c_- = 0$. Hence $\mathcal{T}f = 0$, which, combined with A.2, furnishes $f = 0$. This completes the proof of the lemma. \square

B.

In this appendix, we prove for $\mathcal{R}\mathbf{f}$ defined as (6.19)

Lemma B.1. *Suppose that Assumption 6.2 holds. Then $\mathcal{R}\mathbf{f} \in \mathcal{L}_q(\mathbf{R}^N)$ for any $\mathbf{f} \in F_q(\mathbf{R}^N \setminus \Sigma)$.*

Proof. Since $\text{supp } \mathcal{R}\mathbf{f} \subset D_{R_1, R_2}$, it suffices to verify that $(\mathcal{R}\mathbf{f}, 1)_{\mathbf{R}^N} = 0$ for $\mathbf{f} \in F_q(\mathbf{R}^N \setminus \Sigma)$ in what follows. Let $\psi = 1 - \psi_\infty$. Similarly to (A.2), we write

$$\begin{aligned}
 \mathcal{R}\mathbf{f} &= \psi \Delta \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) - \text{div}((\nabla \psi) \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) + \psi \nabla \mathcal{S}_\infty(\varphi_\infty \mathbf{f})) \\
 &\quad - \psi_0 \Delta \mathcal{S}_0(\varphi_0 \mathbf{f}) + \text{div}((\nabla \psi_0) \mathcal{S}_0(\varphi_0 \mathbf{f}) + \psi_0 \nabla \mathcal{S}_0(\varphi_0 \mathbf{f})) \quad \text{in } \mathbf{R}^N \setminus \Sigma,
 \end{aligned}$$

which, combined with the facts:

$$\psi \Delta \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) = \psi \text{div}(\varphi_\infty \mathbf{f}) = 0, \quad \psi_0 \Delta \mathcal{S}_0(\varphi_0 \mathbf{f}) = \psi_0 \text{div}(\varphi_0 \mathbf{f}) = \text{div}(\varphi_0 \mathbf{f}),$$

furnishes that

$$\mathcal{R}\mathbf{f} = -\text{div}(\varphi_0 \mathbf{f}) - \text{div}((\nabla \psi) \mathcal{S}_\infty(\varphi_\infty \mathbf{f}) + \psi \nabla \mathcal{S}_\infty(\varphi_\infty \mathbf{f}))$$

$$+ \operatorname{div}((\nabla\psi_0)\mathcal{S}_0(\varphi_0\mathbf{f}) + \psi_0\nabla\mathcal{S}_0(\varphi_0\mathbf{f})) \quad \text{in } \mathbf{R}^N \setminus \Sigma.$$

It then holds that

$$\begin{aligned} (\mathcal{R}\mathbf{f}, 1)_{B_{4R}\setminus\Sigma} &= -(\operatorname{div}(\varphi_0\mathbf{f}), 1)_{B_{4R}\setminus\Sigma} \\ &\quad - (\operatorname{div}((\nabla\psi)\mathcal{S}_\infty(\varphi_\infty\mathbf{f}) + \psi\nabla\mathcal{S}_\infty(\varphi_\infty\mathbf{f})), 1)_{B_{4R}\setminus\Sigma} \\ &\quad + (\operatorname{div}((\nabla\psi_0)\mathcal{S}_0(\varphi_0\mathbf{f})), 1)_{B_{4R}\setminus\Sigma} + (\operatorname{div}(\psi_0\nabla\mathcal{S}_0(\varphi_0\mathbf{f})), 1)_{B_{4R}\setminus\Sigma} \\ &=: I_1 + I_2 + I_3 + I_4. \end{aligned}$$

By Gauss's divergence theorem, we see that

$$\begin{aligned} I_1 &= - \int_{\Sigma} [\mathbf{n} \cdot \mathbf{f}] \, d\sigma, \quad I_2 = I_3 = 0, \\ I_4 &= \int_{\Sigma} \psi_0 [\mathbf{n} \cdot \nabla\mathcal{S}_0(\varphi_0\mathbf{f})] \, d\sigma = \int_{\Sigma} \varphi_0 [\mathbf{n} \cdot \mathbf{f}] \, d\sigma = \int_{\Sigma} [\mathbf{n} \cdot \mathbf{f}] \, d\sigma, \end{aligned}$$

where $d\sigma$ is the surface element of Σ . Hence $(\mathcal{R}\mathbf{f}, 1)_{\mathbf{R}^N} = (\mathcal{R}\mathbf{f}, 1)_{B_{4R}\setminus\Sigma} = 0$, which completes the proof. \square

C.

In this appendix, we prove Lemma 6.10. Similarly to Subsection 6.2 and the appendix A, it suffices to prove the following lemma.

Lemma C.1. *Suppose that Assumption 6.2 holds. Let $q \in (1, 2)$ additionally and $f \in \mathcal{L}_q(\mathbf{R}^N)$ with $(I + \mathcal{G})f = 0$. Then $f = 0$.*

Proof. Since $(I + \mathcal{G})f = 0$, we have by (6.18)

$$\begin{cases} \Delta\mathcal{T}f = 0 & \text{in } \mathbf{R}^N \setminus \Sigma, \\ \llbracket \rho\mathcal{T}f \rrbracket = 0 & \text{on } \Sigma, \\ \llbracket \mathbf{n} \cdot \nabla\mathcal{T}f \rrbracket = 0 & \text{on } \Sigma. \end{cases}$$

On the other hand, one sees by Theorem 6.3 that for any $\Phi \in C_0^\infty(\mathbf{R}^N \setminus \Sigma) \subset F_{q'}(\mathbf{R}^N \setminus \Sigma)$, $q' = q/(q-1) \in (2, \infty)$, there exists $w \in \widehat{H}_{q'}^1(\mathbf{R}^N \setminus \Sigma) \cap \widehat{H}_{q'}^2(\mathbf{R}^N \setminus \Sigma)$ such that

$$\begin{cases} \Delta w = \operatorname{div} \Phi & \text{in } \mathbf{R}^N \setminus \Sigma, \\ \llbracket \rho w \rrbracket = 0 & \text{on } \Sigma, \\ \llbracket \mathbf{n} \cdot \nabla w \rrbracket = 0 & \text{on } \Sigma. \end{cases}$$

Similarly to the proof of uniqueness in Subsection 3.3, it holds that $\rho\mathcal{T}f = c$ for some constant c . One then concludes $c = 0$ by (6.13), i.e. $\mathcal{T}f = 0$, which, combined with Lemma A.2 furnishes $f = 0$. This completes the proof of Lemma C.1. \square

REFERENCES

- [1] T. Abe and Y. Shibata. On a resolvent estimate of the Stokes equation on an infinite layer. II. $\lambda = 0$ case. *J. Math. Fluid Mech.*, 5(3):245–274, 2003.
- [2] H. Abels. Generalized Stokes resolvent equations in an infinite layer with mixed boundary conditions. *Math. Nachr.*, 279(4):351–367, 2006.
- [3] R.A. Adams and J.J.F. Fournier. *Sobolev spaces*, volume 140 of *Pure and Applied Mathematics*. Elsevier/Academic Press, Amsterdam, 2nd edition, 2003.
- [4] E. DiBenedetto. *Real Analysis*. Birkhäuser Advanced Texts: Basel Textbooks. Birkhäuser, second edition, 2016.
- [5] E. Fabes, O. Mendez, and M. Mitrea. Boundary layers on Sobolev-Besov spaces and Poisson's equation for the Laplacian in Lipschitz domains. *J. Funct. Anal.*, 159(2):323–368, 1998.

- [6] R. Farwig, H. Kozono, and H. Sohr. An L^q -approach to Stokes and Navier-Stokes equations in general domains. *Acta Math.*, 195:21–53, 2005.
- [7] R. Farwig, H. Kozono, and H. Sohr. On the Helmholtz decomposition in general unbounded domains. *Arch. Math. (Basel)*, 88(3):239–248, 2007.
- [8] R. Farwig and H. Sohr. Generalized resolvent estimates for the Stokes system in bounded and unbounded domains. *J. Math. Soc. Japan*, 46(4):607–643, 1994.
- [9] R. Farwig and H. Sohr. Helmholtz decomposition and Stokes resolvent system for aperture domains in L^q -spaces. *Analysis*, 16(1):1–26, 1996.
- [10] D. Fujiwara and H. Morimoto. An L_r -theorem of the Helmholtz decomposition of vector fields. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 24(3):685–700, 1977.
- [11] G.P. Galdi. *An Introduction to the Mathematical Theory of the Navier-Stokes Equations: Steady-State Problems*. Springer Monogr. Math. Springer, New York, second edition, 2011.
- [12] J. Geng and Z. Shen. The Neumann problem and Helmholtz decomposition in convex domains. *J. Funct. Anal.*, 259(8):2147–2164, 2010.
- [13] S. Maryani and H. Saito. On the \mathcal{R} -boundedness of solution operator families for two-phase Stokes resolvent equations. *Differential Integral Equations*, 30(1-2):1–52, 2017.
- [14] V. N. Maslennikova and M. E. Bogovskii. Elliptic boundary value problems in unbounded domains with noncompact and nonsmooth boundaries. *Rend. Sem. Mat. Fis. Milano*, 56:125–138 (1988), 1986.
- [15] T. Miyakawa. On nonstationary solutions of the Navier-Stokes equations in an exterior domain. *Hiroshima Math. J.*, 12(1):115–140, 1982.
- [16] T. Miyakawa. The Helmholtz decomposition of vector fields in some unbounded domains. *Math. J. Toyama Univ.*, 17:115–149, 1994.
- [17] J. Prüss and G. Simonett. *Moving interfaces and quasilinear parabolic evolution equations*, volume 105 of *Monographs in Mathematics*. Birkhäuser/Springer, [Cham], 2016.
- [18] H. Saito. Global solvability of the Navier-Stokes equations with a free surface in the maximal L_p - L_q regularity class. *J. Differential Equations*, 264(3):1475–1520, 2018.
- [19] H. Saito. Existence of \mathcal{R} -bounded solution operator families for a compressible fluid model of Korteweg type on the half-space. *preprint*, 2019. arXiv:1901.06461 [math.AP].
- [20] H. Saito, Y. Shibata, and X. Zhang. Some free boundary problem for two phase inhomogeneous incompressible flow. 2018. preprint, arXiv: 1811.02179.
- [21] K. Schade and Y. Shibata. On strong dynamics of compressible nematic liquid crystals. *SIAM J. Math. Anal.*, 47(5):3963–3992, 2015.
- [22] Y. Shibata. *Introduction to the Mathematical Theory of Fluid Mechanics*. in japanese (to appear).
- [23] Y. Shibata. Generalized resolvent estimates of the Stokes equations with first order boundary condition in a general domain. *J. Math. Fluid Mech.*, 15(1):1–40, 2013.
- [24] Y. Shibata. On the local wellposedness of free boundary problem for the Navier-Stokes equations in an exterior domain. *Commun. Pure Appl. Anal.*, 17(4):1681–1721, 2018.
- [25] Y. Shibata and S. Shimizu. On the maximal L_p - L_q regularity of the Stokes problem with first order boundary condition; model problems. *J. Math. Soc. Japan*, 64(2):561–626, 2012.
- [26] C. G. Simader and H. Sohr. A new approach to the Helmholtz decomposition and the Neumann problem in L^q -spaces for bounded and exterior domains. In *Mathematical problems relating to the Navier-Stokes equation*, volume 11 of *Ser. Adv. Math. Appl. Sci.*, pages 1–35. World Sci. Publ., River Edge, NJ, 1992.
- [27] C. G. Simader and H. Sohr. *The Dirichlet problem for the Laplacian in bounded and unbounded domains*, volume 360 of *Pitman Research Notes in Mathematics Series*. Longman, Harlow, 1996. A new approach to weak, strong and $(2+k)$ -solutions in Sobolev-type spaces.
- [28] H. Sohr. *The Navier-Stokes equations*. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 2001. An elementary functional analytic approach, [2013 reprint of the 2001 original] [MR1928881].
- [29] V. A. Solonnikov. Estimates fo the solutions of nonstationary Navier-Stokes equations. *J. Soviet Math.*, 8:467–529, 1977.
- [30] S. Szufła. On the Hammerstein integral equation with weakly singular kernel. *Funkcial. Ekvac.*, 34(2):279–285, 1991.
- [31] H. Weyl. The method of orthogonal projection in potential theory. *Duke Math. J.*, 7:411–444, 1940.

FACULTY OF INDUSTRIAL SCIENCE AND TECHNOLOGY, TOKYO UNIVERSITY OF SCIENCE, 102-1
TOMINO, OSHAMAMBE-CHO, YAMAKOSHI-GUN, HOKKAIDO 049-3514, JAPAN

E-mail address: `hsaito@rs.tus.ac.jp`

FACULTY OF SCIENCE AND ENGINEERING, WASEDA UNIVERSITY, OKUBO 3-4-1, SHINJUKU-KU,
TOKYO 169-8555, JAPAN

E-mail address: `xinzhang@aoni.waseda.jp`