

Higher order asymptotic expansion of solutions to abstract linear hyperbolic equations

Motohiro Sobajima^{*}

Abstract. The paper concerned with higher order asymptotic expansion of solutions to the Cauchy problem of abstract hyperbolic equations of the form $u'' + Au + u' = 0$ in a Hilbert space, where A is a nonnegative selfadjoint operator. The result says that by assuming the regularity of initial data, asymptotic profiles (of arbitrary order) are explicitly written by using the semigroup e^{-tA} generated by $-A$. To prove this, a kind of maximal regularity for e^{-tA} is used.

Mathematics Subject Classification (2010): Primary:35L90, 35B40, Secondary:34G10, 35E15.

Key words and phrases: Hyperbolic equations in Hilbert spaces, higher order expansion of solution.

1 Introduction

Let H be a Hilbert space over \mathbb{R} with the inner product (\cdot, \cdot) and the norm $\|\cdot\|$. In this paper we consider higher order asymptotic expansion of solutions to abstract hyperbolic equations of the form

$$\begin{cases} u''(t) + Au(t) + u'(t) = 0, & t > 0, \\ (u, u')(0) = (u_0, u_1), \end{cases} \quad (1.1)$$

where $u : [0, t) \rightarrow H$ is an unknown function and A is a nonnegative selfadjoint operator in H . We denote $D(A)$ as a domain of A . The pair $(u_0, u_1) \in D(A^{1/2}) \times H$ is given.

The problem (1.1) is motivated as the generalization of the damped wave equation

$$\begin{cases} \partial_t^2 u(x, t) - \Delta u(x, t) + \partial_t u(x, t) = 0, & (x, t) \in \mathbb{R}^N \times (0, \infty), \\ (u, u')(x, 0) = (u_0(x), u_1(x)). \end{cases} \quad (1.2)$$

with $(u_0, u_1) \in H^1(\mathbb{R}^N) \times L^2(\mathbb{R}^N)$. The abstract framework as (1.1) is firstly introduced by Ikehata–Nishihara [6].

The equation (1.2) has been considered as a model of the phenomenon of heat conduction with finite propagation property (see Cattaneo [2] and Vernotte [20]). Therefore the solution of (1.2) is expected to have a similar profile of solutions to the heat equation

$$\begin{cases} \partial_t v(x, t) - \Delta v(x, t) = 0, & (x, t) \in \mathbb{R}^N \times (0, \infty), \\ v(x, 0) = v_0(x). \end{cases} \quad (1.3)$$

Actually it is known that under the suitable assumption, the solution u of (1.2) behaves like the solution v of (1.3) with $v_0 = u_0 + u_1$. This phenomenon is now so-called diffusion phenomenon for the damped wave equations. There are many investigation dealing with diffusion phenomena for

^{*} Department of Mathematics, Faculty of Science and Technology, Tokyo University of Science, 2641 Yamazaki, Noda-shi, Chiba, 278-8510, Japan, E-mail: msobajima1984@gmail.com

(1.2) (see e.g., Hsiao–Liu [4], Nishihara [11, 12], Karch [9], Yang–Milani [22] including some generalized problems like quasilinear systems). In the case of the damped wave equation in the exterior domain, Fourier analysis does not work well, and therefore, energy methods via integration by parts are often used (see e.g., Ikehata–Matsuyama [7], Ikehata [5] and Ikehata–Saeki [8]).

Another example of the (1.1) is the damped beam equation

$$\begin{cases} \partial_t^2 u(x, t) + \partial_x^4 u(x, t) - \alpha \partial_x^2 u(x, t) + \partial_t u(x, t) = \partial_x f(\partial_x u(x, t)), & (x, t) \in \mathbb{R} \times (0, \infty), \\ (u, u')(x, 0) = (u_0(x), u_1(x)). \end{cases} \quad (1.4)$$

The global existence and asymptotic behavior of solutions to (1.4) are studied in Takeda–Yoshikawa [18, 17, 19] and Takeda [16]. In these papers, the asymptotic behavior of solutions of (1.4) with $f \equiv 0$ plays a crucial role to consider global existence of nonlinear problem. Actually, in [17] the asymptotic behavior of solutions of nonlinear problem can be found as solutions of a linear problem.

For the further analysis, as an improvement of diffusion phenomena, the problem of higher order asymptotic expansion should be naturally considered. In 2001, Orive–Zuazua–Pazoto [13], they considered the following general problem of (1.2):

$$\begin{cases} \rho(x) \partial_t^2 u(x, t) - \operatorname{div} \left(a(x) \nabla u(x, t) \right) + a_0 \rho(x) \partial_t u(x, t) = 0, & (x, t) \in \mathbb{R}^N \times (0, \infty), \\ (u, u')(x, 0) = (u_0(x), u_1(x)), \end{cases} \quad (1.5)$$

where $\rho(x)$, $\rho(x)^{-1}$ and $a(x) = (a_{jk}(x))_{jk}$ are bounded and spatially periodic and the matrix $(a_{jk})_{jk}$ is uniformly and positively determined. In [13], higher order asymptotic expansion of solutions to (1.5) via the use of Bloch wave decomposition which is valid for spatial periodic coefficients. Later, Takeda [16] gave higher order asymptotic expansion by using the usual heat semigroup $e^{t\Delta}$ for (1.2) via Fourier multiplier theory with asymptotic expansion of symbol of evolution operator with respect to the variable of Fourier spaces ξ . An asymptotic expansion for wave part can be found in a recent paper Michihisa [10].

Instead of the various results in the previous works, in this paper the interest is how to systematically determine the asymptotic profile of arbitrary order. To discuss this problem we then consider an abstract hyperbolic equation (1.1) which is included three situations (1.2) ($A = -\Delta$), (1.4) ($A = \frac{d^4}{dx^4} - \alpha \frac{d^2}{dx^2}$) and (1.5) ($Au = \rho^{-1} \operatorname{div}(a \nabla u)$).

Concerning the previous works of the abstract setting, Ikehata–Nishihara [6] introduced the problem (1.1) and observed the asymptotic behavior of solutions in the following way:

$$\|u(t) - e^{-tA}(u_0 + u_1)\| \leq C(1+t)^{-1} \left(\log(e+t) \right)^{1/2+\varepsilon}, \quad (u_0, u_1) \in D(A) \times D(A^{1/2}).$$

After that Chill–Haraux [3] discussed the same problem and succeeded in removing the logarithmic correction of the above inequality, which is conjectured in Ikehata–Nishihara [6]. Radu–Todorova–Yordanov [14] studied also diffusion phenomena with respect to stronger norms $\|\cdot\|_{D(A^k)}$ (a similar analysis for a linear hyperbolic equation in Hilbert spaces with time-dependent damping term $b(t)u'$ can be found in Yamazaki [21]). Radu–Todorova–Yordanov [15] discussed a higher order approximation of solutions to $Bu'' + Au + u' = 0$ (B is bounded, selfadjoint and positively definite); however, their framework is only valid for semigroup in metric measure spaces $L^2(\Omega, \mu)$ with an extra maximal L^p – L^q regularity.

The purpose of the present paper is to discuss higher order asymptotic expansion of solutions (1.1) by using the semigroup e^{-tA} generated by $-A$. Moreover, the main topic is to give a way how to construct higher order asymptotic profiles of arbitrary order.

To begin with, we state existence of solutions to the problem (1.1). The following proposition follows from the standard theory for C_0 -semigroup on product spaces (see e.g., Ikehata–Nishihara [6] and also Brezis [1, Section X] when $A = -\Delta$).

Proposition 1.1. *Let A be a nonnegative selfadjoint operator in H with domain $D(A)$. Then the operator*

$$\mathcal{A} = \begin{pmatrix} 0 & -1 \\ A & 1 \end{pmatrix},$$

in $\mathcal{H} = D(A^{1/2}) \times H$ with domain $D(\mathcal{A}) = D(A) \times D(A^{1/2})$ is quasi- m -accretive in \mathcal{H} . Namely, $-\mathcal{A}$ generates a C_0 -semigroup in \mathcal{H} .

In particular, the solution u of (1.1) uniquely exists in the following sense:

(i) *for $(u_0, u_1) \in D(A^{1/2}) \times D(H)$, one has $(u(t), u'(t)) = e^{-t\mathcal{A}}(u_0, u_1)$ which implies*

$$u \in C^2([0, \infty); [D(A^{1/2})]^*) \cap C^1([0, \infty); H) \cap C([0, \infty); D(A^{1/2})).$$

and the equation (1.1) is satisfied in $[D(A^{1/2})]^$ (the dual space of $D(A^{1/2})$), that is,*

$$\langle u''(t), \varphi \rangle_{[D(A^{1/2})]^*, D(A^{1/2})} + (A^{1/2}u(t), A^{1/2}\varphi) + (u'(t), \varphi) = 0, \quad \forall \varphi \in D(A^{1/2})$$

(ii) *for $(u_0, u_1) \in D(A^{\frac{n+1}{2}}) \times D(A^{\frac{n}{2}})$ ($n \in \mathbb{N}$), one has $(u(t), u'(t)) = e^{-t\mathcal{A}}(u_0, u_1)$ which implies*

$$u \in \bigcap_{k=0}^{n+1} C^k([0, \infty); D(A^{\frac{n-k}{2}}))$$

Now we are in a position to give the result for higher order asymptotic expansion for abstract hyperbolic equation (1.1), which is the main result of the present paper.

Theorem 1.2. *Assume that $(u_0, u_1) \in D(A^{m+1/2}) \times D(A^{m+1/2})$ for some $m \in \mathbb{N} \cup \{0\}$. Let u be a unique solution of (1.1). Define $v_0 = u_0 + u_1$ and for $\ell \in \mathbb{N} \cup \{0\}$,*

$$\begin{aligned} \bar{u}_0(t) &= e^{-tA}v_0, \\ \bar{u}_\ell(t) &= A^\ell \left(\sum_{j=0}^{\ell} \binom{2\ell-1}{\ell+j-1} \frac{(-tA)^j}{j!} e^{-tA}v_0 + \sum_{k=0}^{\ell-1} \binom{2\ell-1}{\ell+k} \frac{(-tA)^k}{k!} e^{-tA}u_1 \right). \end{aligned}$$

Then there exists a positive constant C such that for every $t \geq 0$,

$$\left\| u(t) - \sum_{\ell=0}^m \bar{u}_\ell(t) \right\| \leq C(1+t)^{-m-1/2}.$$

Remark 1.1. Especially, if 0 is an eigenvalue of A and φ_0 is the corresponding eigenvector, then $e^{-tA}\varphi_0 = \varphi_0$. Therefore there is no possibility to improve the upper bound (1.6) in the general setting. Since the asymptotic profiles $\{\bar{u}_\ell\}$ satisfy

$$\|\bar{u}_\ell(t)\| \leq C_\ell(1+t)^{-\ell}, \tag{1.6}$$

the assertion in Theorem 1.2 can be understood as a higher order asymptotic expansion of solution u of (1.1). Of course in the several situation such as $-\Delta$ in \mathbb{R}^N , we have $\|e^{-tA}(u_0 + u_1)\| \leq C(1+t)^{-\alpha}$ for some $\alpha > 0$. In this case we need some effort to determine especially the regularity of initial data. We will not touch the details of specialized cases.

Remark 1.2. If $m = 0$, then Theorem 1.2 is weaker than those of Ikehata–Nishihara [6] and also Chill–Haraux [3]. Since the main topic of our result is to give a higher order asymptotic profiles, the optimality of decay rates is not precisely discussed.

Remark 1.3. In the case (1.2), we choose $H = L^2(\mathbb{R}^N)$ and $A = -\Delta$ endowed with domain $D(A) = H^2(\mathbb{R}^N)$. Takeda [16] obtained the same asymptotic expansion with a different expression

$$\begin{aligned} \bar{u}_\ell(t) &= \frac{1}{2} \sum_{j=0}^{\ell} \alpha_{j,k} (-t)^j (-\Delta)^{j+\ell} e^{t\Delta} u_0 \\ &+ \sum_{0 \leq k_1+k_2 \leq \ell} \alpha_{\ell-k_1-k_2,k_1} \beta_{k_2} (-t)^{\ell-k_1-k_2} (-\Delta)^{2\ell-k_1-k_2} e^{t\Delta} \left(\frac{1}{2} u_0 + u_1 \right), \end{aligned}$$

where

$$\begin{aligned} \alpha_{j,k} &= \frac{1}{j!k!} \left. \frac{d^k}{dr^k} \phi_j \right|_{r=0}, \quad \phi_j(r) = \left(\frac{1}{2} + \sqrt{\frac{1}{4} - r} \right)^{-2j}, \\ \beta_k &= \frac{1}{k!} \left. \frac{d^k}{dr^k} \psi \right|_{r=0}, \quad \psi(r) = \frac{1}{2} \left(\frac{1}{4} - r \right)^{-1/2}. \end{aligned}$$

This result is valid for the whole space case \mathbb{R}^N and the effect of high-frequency part is clearly written. On the other hand, Theorem 1.2 asserts that the same asymptotic expansion is valid for every nonnegative selfadjoint operator in a Hilbert space.

Remark 1.4. In the case (1.4), we choose $H = L^2(\mathbb{R})$ and $A = \frac{d^4}{dx^4} - \alpha \frac{d^2}{dx^2}$ endowed with domain $D(A) = H^4(\mathbb{R})$. According to Theorem 1.2 with $m = 0$, we can see that

$$\|u(t) - e^{-tA}(u_0 + u_1)\| \leq C(1+t)^{-\frac{1}{2}}.$$

On the other hand, in Takeda–Yoshikawa [17], it is shown that the asymptotic behavior of the solution u is given by $e^{-\alpha t A_0}(u_0 + u_1)$, where $A_0 = -\frac{d^2}{dx^2}$. The observation in Theorem 1.2 does not have contradiction because of the following estimate

$$\left\| e^{-tA}(u_0 + u_1) - e^{-\alpha t A_0}(u_0 + u_1) \right\| \leq \frac{C}{t} \|t A_0 e^{-\alpha t A_0} A_0(u_0 + u_1)\|, \quad u_0 + u_1 \in H^2(\mathbb{R}).$$

However, if we choose the latter profile, then since the asymptotic expansion in Theorem 1.2 is written by the operator A ($\neq A_0$), the their difference should be carefully analysed.

Remark 1.5. In the case (1.5) (with bounded ρ, ρ^{-1}), we choose $H = L^2(\mathbb{R})$ but an inner product different from the usual one:

$$(f, g)_H = \int_{\mathbb{R}^N} f g \rho dx.$$

Then we use $Au = \rho(x)^{-1} \operatorname{div}(a(x) \nabla u)$ endowed with domain $D(A) = H^2(\mathbb{R}^N)$. Of course we can consider the periodic setting $\mathbb{T}^N = (\mathbb{R}/\mathbb{Z})^N$ by choosing Sobolev spaces of periodic functions. Combining the strategy of Bloch wave decomposition, we can also deduce a similar result in [13].

Let us describe the strategy for the construction of asymptotic profiles. It is well known that the solution u of (1.1) has the energy decreasing property

$$E(u, t) + 2 \int_0^t \|u'(s)\|^2 ds = E(u, 0) = \|u_1\|^2 + \|A^{1/2}u_0\|^2,$$

where the energy functional of the solution u of (1.1) is defined as

$$E(u; t) = \|u'(t)\|^2 + \|A^{1/2}u(t)\|^2.$$

According to the experiences, we expect that the first order asymptotic profile of u is $e^{-tA}(u_0 + u_1)$. On the other hand, if w is the solution of (1.1), then by a direct computation we have

$$(1+t)\|w'(t)\|^2 \leq (1+t)E(w, t) + \|w(t)\|^2 \leq C.$$

Combining this concept, we consider the following auxiliary problem

$$\begin{cases} U_1'' + AU_1 + U_1' = Ae^{tA}(u_0 + u_1), & t > 0, \\ (U_1, U_1')(0) = (0, -u_1). \end{cases} \quad (1.7)$$

After some computation, we have $u(t) = e^{-tA}(u_0 + u_1) + U_1'(t)$, which is the first decomposition. It should be noticed that the function U_1 is completely the same as the function Z in Ikehata–Nishihara [6], however, the treatment of U_1 is different from [6] in which Z is understood as the solution of $Z' + AZ = u'$, $Z(0) = 0$. The idea of (1.7) can be understood as another use of the modified Morawetz method which is written in Ikehata–Matsuyama [7].

Then analysing the asymptotic profile of the solution U_1 and proceeding the similar argument as before, we successively obtain the sequence U_ℓ and their corresponding hyperbolic problems similar to (1.7). Finally, to obtain the desired decay property of error terms (such as U_1'), we use a kind of maximal regularity for the Cauchy problem of parabolic equation $V' + AV' = 0$. This consideration suggests that the solution u of (1.1) can be decomposed by

$$u(t) = \sum_{\ell=0}^m \frac{d^\ell}{dt^\ell} V_\ell(t) + \frac{d^{m+1}}{dt^{m+1}} U_{m+1}(t).$$

The present paper is organized as follows. In Section 2, as a preliminary, we state and prove a kind of maximal regularity result the Cauchy problem of parabolic equation $V' + AV' = 0$, which we will use later. In Section 3, we provide a proof of the simplest case $m = 0$ in Theorem 1.2 to clarify how to show the diffusion phenomena. The first part of Section 4 provides a construction of a family of function which will be understood as higher order asymptotic profiles. Finally, at the rest of Section 4 we show higher order asymptotic expansion of solutions to (1.1) (Theorem 1.2 for $m \in \mathbb{N}$).

2 Preliminary result for the property of the semigroup e^{-tA}

In this section, we give recall a kind of maximal regularity result for the abstract (selfadjoint) semigroup e^{-tA} . For the reader's convenience, we provide a short proof.

Lemma 2.1. *If $f \in H$, then for every $n \in \mathbb{N} \cup \{0\}$ and $t \geq 0$,*

$$\frac{\|e^{-tA}f\|^2}{2} + \frac{2^n}{n!} \int_0^t s^n \|A^{\frac{n+1}{2}} e^{-sA} f\|^2 ds = \frac{\|f\|^2}{2}. \quad (2.1)$$

Moreover, if $f \in D(A^{n/2})$, then there exists a positive constant C such that

$$\int_0^t (1+s)^n \|A^{\frac{n+1}{2}} e^{-sA} f\|^2 ds \leq C(\|f\|^2 + \|A^{n/2} f\|^2).$$

Proof. If $f \in D(A^{n/2+1})$, then $w(t) = e^{-tA} f$ satisfies

$$\begin{aligned} \frac{d}{dt} \|A^{k/2} w(t)\|^2 &= -2(A^{k/2} w(t), A^{k/2} w'(t)) \\ &= -2(A^{k/2} w(t), A^{k/2+1} w(t)) \\ &= -2\|A^{\frac{k+1}{2}} w(t)\|^2. \end{aligned}$$

In view of the above computation, in particular, we have

$$\frac{d}{dt} [t^{k+1} \|A^{\frac{k+1}{2}} w(t)\|^2] = (k+1)t^k \|A^{\frac{k+1}{2}} w(t)\|^2 - 2t^{k+1} \|A^{\frac{k+2}{2}} w(t)\|^2$$

and therefore

$$\begin{aligned} \frac{d}{dt} \left[\sum_{k=0}^{n-1} \frac{(2t)^{k+1}}{(k+1)!} \|A^{\frac{k+1}{2}} w(t)\|^2 \right] &= 2 \sum_{k=0}^{n-1} \left(\frac{(2t)^k}{k!} \|A^{\frac{k+1}{2}} w(t)\|^2 - \frac{(2t)^{k+1}}{(k+1)!} \|A^{\frac{k+2}{2}} w(t)\|^2 \right) \\ &= \|A^{1/2} w(t)\|^2 - \frac{(2t)^n}{n!} \|A^{\frac{n+1}{2}} w(t)\|^2. \end{aligned}$$

Integrating it over $[0, t]$, we obtain (2.1). \square

3 The first asymptotics

In this section, we will show the strategy how to find a asymptotic profile of solutions in the simplest situation $m = 0$ in Theorem 1.2.

To begin with, we prove the following lemma.

Lemma 3.1. *Assume that $(u_0, u_1) \in \mathcal{H}$ and let u be a unique solution of (1.1). Then there exists a positive constant $C > 0$ such that for every $t \geq 0$,*

$$(1+t)E(u, t) + \|u(t)\|^2 + \int_0^t \left((1+s)\|u'(s)\|^2 + \|A^{1/2} u(s)\|^2 \right) ds \leq C(E(u, 0) + \|u_0\|^2).$$

Proof. By taking a suitable approximation of (u_0, u_1) , We may assume $(u_0, u_1) \in D(\mathcal{A})$ without loss of generality. Then we see by a direct computation with the equation (1.1) that

$$\begin{aligned} \frac{d}{dt} E(u; t) &= 2(u', u'') + 2(A^{1/2} u', A^{1/2} u) \\ &= 2(u', u'' + Au) \\ &= -2\|u'\|^2. \end{aligned}$$

This yields

$$\|u'(t)\|^2 + \|A^{1/2}u\|^2 + 2 \int_0^t \|u'(s)\|^2 ds = \|u_1\|^2 + \|A^{1/2}u_0\|^2.$$

On the other hand, using the equation (1.1), we also have the following two identities:

$$\begin{aligned} \frac{d}{dt} \left[(3+t)E(u; t) \right] &= \|u'\|^2 + \|A^{1/2}u\|^2 - 2(3+t)\|u'\|^2, \\ \frac{d}{dt} E_*(u; t) &= 2\|u'\|^2 + 2(u, u'' + u') \\ &= 2\|u'\|^2 - 2\|A^{1/2}u\|^2. \end{aligned}$$

Summing up the above identities, we deduce

$$\frac{d}{dt} \left[(3+t)E(u; t) + E_*(u; t) \right] = -(3+2t)\|u'\|^2 - \|A^{1/2}u\|^2. \quad (3.1)$$

Noting that the differentiated function in (3.1) can be rewritten as

$$(3+t)E(u; t) + E_*(u; t) = (1+t)E(u, t) + 2\|A^{1/2}u\|^2 + 2 \left\| u' + \frac{1}{2}u \right\|^2 + \frac{1}{2}\|u\|^2,$$

we obtain the desired inequality. \square

The next lemma enables us to divide the solution u of (1.1) into two factors: one reflects the diffusion phenomenon and the other reflects an effect of hyperbolicity. This treatment is the difference between the one in Ikehata–Nishihara [6] and ours.

Lemma 3.2. *Let $(u_0, u_1) \in D(A^{1/2}) \times D(A^{1/2})$ and let U_1 be the unique solution of the problem*

$$\begin{cases} U_1'' + AU_1 + U_1' = Ae^{tA}(u_0 + u_1), & t > 0, \\ (U_1, U_1')(0) = (0, -u_1). \end{cases} \quad (3.2)$$

Then the function $e^{-tA}(u_0 + u_1) + U_1'$ coincides with the solution of (1.1).

Proof. Take $(u_{0\varepsilon}, u_{1\varepsilon}) \in D(A^2) \times D(A^2)$ such that $u_{0\varepsilon} \rightarrow u_0$ and $u_{1\varepsilon} \rightarrow u_1$ in $D(A^{1/2})$ as $\varepsilon \rightarrow 0$ and consider

$$\begin{cases} U_{1\varepsilon}'' + AU_{1\varepsilon} + U_{1\varepsilon}' = Ae^{tA}(u_{0\varepsilon} + u_{1\varepsilon}), & t > 0, \\ (U_{1\varepsilon}, U_{1\varepsilon}')(0) = (0, -u_{1\varepsilon}). \end{cases} \quad (3.3)$$

Since $Ae^{tA}(u_{0\varepsilon} + u_{1\varepsilon}) \in C([0, \infty); D(A))$, this problem has a unique solution

$$U_{1\varepsilon} \in C^2([0, \infty); D(A^{1/2})) \cap C^1([0, \infty); D(A)) \cap C([0, \infty); D(A^{3/2})).$$

Moreover, by using Duhamel's principle we can see that the solution $U_{1\varepsilon}$ can be represented by

$$(U_{1\varepsilon}(t), U_{1\varepsilon}'(t)) = e^{-tA}(0, -u_{1\varepsilon}) + \int_0^t e^{-(t-s)A}(0, Ae^{-sA}(u_{0\varepsilon} + u_{1\varepsilon})) ds$$

which converges to the solution of (3.2) in \mathcal{H} as $\varepsilon \rightarrow 0$:

$$e^{-tA}(0, -u_1) + \int_0^t e^{-(t-s)A}(0, Ae^{sA}(u_0 + u_1)) ds = (U_1(t), U_1'(t)).$$

Setting $v_\varepsilon(t) = e^{-tA}(u_{0\varepsilon} + u_{1\varepsilon})$ and $w_\varepsilon(t) = v_\varepsilon(t) + U_{1\varepsilon}'(t)$, we have

$$w_\varepsilon(0) = u_{0\varepsilon} + u_{1\varepsilon} - u_{1\varepsilon} = u_{0\varepsilon}$$

and

$$\begin{aligned} w_\varepsilon'(t) &= v_\varepsilon'(t) + U_{1\varepsilon}''(t) \\ &= v_\varepsilon'(t) + Av_\varepsilon(t) - AU_{1\varepsilon}(t) - U_{1\varepsilon}'(t) \\ &= -AU_{1\varepsilon}(t) - U_{1\varepsilon}'(t). \end{aligned}$$

This yields

$$\lim_{t \rightarrow 0} w_\varepsilon'(t) = -AU_{1\varepsilon}(0) - U_{1\varepsilon}'(0) = u_{1\varepsilon}$$

and also $w_\varepsilon' \in C^1((0, \infty); \mathcal{H})$ with

$$\begin{aligned} w_\varepsilon''(t) &= -AU_{1\varepsilon}'(t) - (w_\varepsilon(t) - v_\varepsilon(t))' \\ &= -A(w_\varepsilon(t) - v_\varepsilon(t)) - w_\varepsilon'(t) + v_\varepsilon'(t) \\ &= -Aw_\varepsilon(t) - w_\varepsilon'(t). \end{aligned}$$

This implies $(w_\varepsilon(t), w_\varepsilon'(t)) = e^{-tA}(u_{0,\varepsilon}, u_{1\varepsilon})$ and therefore $(w_\varepsilon(t), w_\varepsilon'(t)) \rightarrow (u(t), u'(t))$ in \mathcal{H} as $\varepsilon \rightarrow 0$. Consequently, noting that $v_\varepsilon(t) \rightarrow e^{-tA}(u_0 + u_1)$ in $D(A^{1/2})$ as $\varepsilon \rightarrow 0$ we have

$$u(t) = \lim_{\varepsilon \rightarrow 0} w_\varepsilon(t) = e^{tA}(u_0 + u_1) + U_1'(t).$$

The proof is complete. \square

Observe that a decaying property of $e^{-tA}(u_0 + u_1)$ cannot be expected as mentioned in Remark 1.1. In contrast, the other factor U' decays like $(1+t)^{-1/2}$ under a suitable assumption. To prove the decay property we proceed a similar way in the proof of Lemma 3.1, however, we need Lemma 2.1 which is a kind of maximal regularity result for e^{-tA} .

Lemma 3.3. *If $(u_0, u_1) \in D(A^{1/2}) \times D(A^{1/2})$, then*

$$(1+t)E(U_1; t) + \|U_1(t)\|^2 \in L^\infty((0, \infty)), \quad (1+t)\|U_1'(t)\|^2 + \|A^{1/2}U_1(t)\|^2 \in L^1((0, \infty))$$

Proof. By using the same approximation as in the proof of Lemma 3.2, we assume may assume $U_1 \in C^2([0, \infty); D(A^{1/2})) \cap C^1([0, \infty); D(A)) \cap C([0, \infty); D(A^{3/2}))$. Here we deduce an energy inequality for U_1 . By the equation (3.2), we have

$$\begin{aligned} \frac{d}{dt} E_*(U_1; t) &= 2\|U_1'\|^2 + 2(U_1, -AU_1 + Ae^{-tA}(u_0 + u_1)) \\ &= 2\|U_1'\|^2 - 2\|A^{1/2}U_1\|^2 + 2(U_1, Ae^{-tA}(u_0 + u_1)) \\ &\leq 2\|U_1'\|^2 - \|A^{1/2}U_1\|^2 + \|A^{1/2}e^{-tA}(u_0 + u_1)\|^2 \end{aligned}$$

and

$$\begin{aligned} \frac{d}{dt} \left[(6+t)E(U_1; t) \right] &= \|U_1'\|^2 + \|A^{1/2}U_1\|^2 + 2(6+t)(U_1', -U_1' + Ae^{-tA}(u_0 + u_1)) \\ &\leq \|U_1'\|^2 + \|A^{1/2}U_1\|^2 - (6+t)\|U_1'\|^2 + (6+t)\|Ae^{-tA}(u_0 + u_1)\|^2 \end{aligned}$$

In view of Lemma 2.1 with $f = u_0 + u_1 \in D(A^{1/2})$, we have

$$\begin{aligned} \frac{d}{dt} \left[(6+t)E(U_1; t) + 2E_*(U_1; t) \right] &+ (1+t)\|U_1'\|^2 + \|A^{1/2}U_1\|^2 \\ &\leq (6+t)\|Ae^{-tA}(u_0 + u_1)\|^2 + 2\|A^{1/2}e^{-tA}(u_0 + u_1)\|^2 \end{aligned}$$

which is integrable with respect to $t \in (0, \infty)$. Integrating the above estimate over $[0, t]$, we obtain all desired estimates. \square

Combining Lemmas 3.1, 3.2 and 3.3, we obtain the following proposition which is the same as the assertion of Theorem 1.2 with $m = 0$.

Proposition 3.4. *Assume $(u_0, u_1) \in D(A^{1/2}) \times D(A^{1/2})$. Then there exists a positive constant $C > 0$ such that for every $t \geq 0$,*

$$\|u(t) - e^{-tA}(u_0 + u_1)\| \leq C(1+t)^{-\frac{1}{2}}.$$

4 Higher order asymptotic expansion

In this section, we prove Theorem 1.2 with $m \in \mathbb{N}$ which gives higher order asymptotic expansion of solutions to (1.1). The philosophy of the proof is essentially the same as the previous section.

4.1 Construction of family of functions for asymptotics

To consider higher order asymptotic expansions, we define the following functions.

Definition 4.1. For $m \in \mathbb{N}$ and $(u_0, u_1) \in \mathcal{H}$, define $v_0 = u_0 + u_1$ and

$$\begin{aligned} V_0^{(1)}(t) &= e^{-tA}v_0, \\ V_m^{(1)}(t) &= (-1)^m \sum_{j=1}^m \binom{m-1}{j-1} \frac{(-tA)^j}{j!} e^{-tA}v_0, \quad m \in \mathbb{N} \\ V_0^{(2)}(t) &= 0, \\ V_m^{(2)}(t) &= (-1)^m \sum_{k=0}^{m-1} \binom{m-1}{k} \frac{(-tA)^k}{k!} e^{-tA}u_1, \quad m \in \mathbb{N} \end{aligned}$$

and $V_m(t) = V_m^{(1)}(t) + V_m^{(2)}(t)$.

The family $\{V_m\}$ will provide higher order asymptotic profiles; this will be explained after stating Lemma 4.4 below.

The function V_m can be successively found by using the following structure.

Lemma 4.1. Assume $u_1 \in D(A^{1/2})$. Then for $m \in \mathbb{N} \cup \{0\}$, V_{m+1} is a unique solution of

$$\begin{cases} V'_{m+1} + AV_{m+1} = -V'_m, & t > 0, \\ V_{m+1}(0) = (-1)^{m+1}u_1. \end{cases}$$

Proof. We use the Duhamel principle. The solution of

$$\begin{cases} w'_{m+1} + Aw_{m+1} = -V'_m, & t > 0, \\ w_{m+1}(0) = (-1)^{m+1}u_1. \end{cases}$$

can be written by the formula

$$w_{m+1}(t) = e^{-tA} \left[(-1)^{m+1}u_1 \right] - \int_0^t e^{-(t-s)A} V'_m(s) ds.$$

For the case $m = 0$, we see from the semigroup property that

$$\begin{aligned} w_1(t) &= e^{-tA} \left[-u_1 \right] - \int_0^t e^{-(t-s)A} V'_0(s) ds \\ &= -e^{-tA}u_1 - \int_0^t e^{-(t-s)A} (-Ae^{sA}v_0) ds \\ &= -e^{-tA}u_1 + tAe^{-tA}v_0 \\ &= V_1^{(1)}(t) + V_1^{(2)}(t). \end{aligned}$$

This gives the assertion with $m = 0$. For the case $m \in \mathbb{N}$, observe that

$$\begin{aligned} &\frac{d}{dt} V_m^{(1)}(t) \\ &= (-1)^m \left[\sum_{j=1}^m \binom{m-1}{j-1} \frac{(-tA)^{j-1}}{(j-1)!} (-A)e^{-tA}v_0 + \sum_{j=1}^m \binom{m-1}{j-1} \frac{(-tA)^j}{j!} (-A)e^{-tA}v_0 \right] \\ &= (-1)^m \left[\sum_{j=1}^m \binom{m-1}{j-1} \frac{(-tA)^{j-1}}{(j-1)!} (-A)e^{-tA}v_0 + \sum_{j=2}^{m+1} \binom{m-1}{j-2} \frac{(-tA)^{j-1}}{(j-1)!} (-A)e^{-tA}v_0 \right] \\ &= (-1)^m \sum_{j=1}^{m+1} \binom{m}{j-1} \frac{(-tA)^{j-1}}{(j-1)!} (-A)e^{-tA}v_0. \end{aligned}$$

Therefore

$$\begin{aligned} \int_0^t e^{-(t-s)A} \left(\frac{d}{dt} V_m^{(1)}(s) \right) ds &= \int_0^t e^{-(t-s)A} \left((-1)^m \sum_{j=1}^{m+1} \binom{m}{j-1} \frac{(-sA)^{j-1}}{(j-1)!} (-A)e^{-sA}v_0 \right) ds \\ &= (-1)^m \sum_{j=1}^{m+1} \binom{m}{j-1} \left(\int_0^t \frac{s^{j-1}}{(j-1)!} ds \right) (-A)^j e^{-tA}v_0 \\ &= (-1)^m \sum_{j=1}^{m+1} \binom{m}{j-1} \frac{(-tA)^j}{j!} e^{-tA}v_0 \\ &= -V_{m+1}^{(1)}(t). \end{aligned}$$

Similarly, we have

$$\begin{aligned}
& \frac{d}{dt} V_m^{(2)}(t) \\
&= (-1)^m \left[\sum_{k=1}^{m-1} \binom{m-1}{k} \frac{(-tA)^{k-1}}{(k-1)!} (-A) e^{-tA} u_1 + \sum_{k=0}^{m-1} \binom{m-1}{k} \frac{(-tA)^k}{k!} (-A) e^{-tA} u_1 \right] \\
&= (-1)^m \left[\sum_{k=1}^{m-1} \binom{m-1}{k} \frac{(-tA)^{k-1}}{(k-1)!} (-A) e^{-tA} u_1 + \sum_{k=1}^m \binom{m-1}{k-1} \frac{(-tA)^{k-1}}{(k-1)!} (-A) e^{-tA} u_1 \right] \\
&= (-1)^m \sum_{k=1}^m \binom{m}{k} \frac{(-tA)^{k-1}}{(k-1)!} (-A) e^{-tA} u_1
\end{aligned}$$

and then

$$\begin{aligned}
\int_0^t e^{-(t-s)A} \left(\frac{d}{dt} V_m^{(2)}(s) \right) ds &= (-1)^m \sum_{k=1}^m \binom{m}{k} \frac{(-tA)^k}{k!} e^{-tA} u_1 \\
&= -V_{m+1}^{(1)}(t) + (-1)^{m+1} e^{-tA} u_1.
\end{aligned}$$

Combining the above two identities, we obtain the assertions for $m \in \mathbb{N}$. \square

We prepare the formula of higher order derivative of the functions V_m .

Lemma 4.2. *Let V_m be as in Definition 4.1. For every $m, \ell \in \mathbb{N}$, one has*

$$\frac{d^\ell}{dt^\ell} V_m(t) = (-A)^\ell V_{m,\ell}(t),$$

where

$$V_{m,\ell}(t) = (-1)^m \left[\sum_{j=0}^m \binom{\ell+m-1}{\ell+j-1} \frac{(-tA)^j}{j!} e^{-tA} v_0 + \sum_{k=0}^{m-1} \binom{\ell+m-1}{\ell+k} \frac{(-tA)^k}{k!} e^{-tA} u_1 \right].$$

Proof. In Lemma 4.1, the following inequalities are already proved:

$$\begin{aligned}
\frac{d}{dt} V_m^{(1)}(t) &= (-1)^m (-A) \left(\sum_{j=0}^m \binom{1+(m-1)}{1+(j-1)} \frac{(-tA)^j}{j!} e^{-tA} v_0 \right), \\
\frac{d}{dt} V_m^{(2)}(t) &= (-1)^m (-A) \left(\sum_{k=0}^{m-1} \binom{1+(m-1)}{1+k} \frac{(-tA)^k}{k!} e^{-tA} u_1 \right).
\end{aligned}$$

As the same manner as in these proofs, we can directly check the desired assertion. \square

The following lemma gives integrability and boundness of functions V_m with respect to $t \in (0, \infty)$.

Lemma 4.3. *Let $(u_0, u_1) \in D(A^{\ell+1/2}) \times D(A^{\ell+1/2})$. Then for every $m, \ell \in \mathbb{N} \cup \{0\}$*

$$(1+t)^{2\ell} \left\| \frac{d^\ell}{dt^\ell} V_m(t) \right\|^2 \in L^\infty((0, \infty)), \quad (4.1)$$

$$(1+t)^{2\ell+1} \left\| \frac{d^{\ell+1}}{dt^{\ell+1}} V_m(t) \right\|^2 + (1+t)^{2\ell} \left\| A^{\ell+1/2} V_{m,\ell+1}(t) \right\|^2 \in L^1((\infty)). \quad (4.2)$$

Proof. By Lemma 4.2 and the boundedness of operators $(tA)^n e^{-\frac{t}{2}A}$ for $n \in \mathbb{N}$, we have

$$\begin{aligned} (1+t)^{2\ell+1} \left\| \left(\frac{d}{dt} \right)^{\ell+1} V_m(t) \right\|^2 &\leq C(1+t)^{2\ell+1} \left(\left\| A^{\ell+1} e^{-\frac{t}{2}A} v_0 \right\|^2 + \left\| A^{\ell+1} e^{-\frac{t}{2}A} u_1 \right\|^2 \right), \\ (1+t)^{2\ell} \left\| A^{\ell+1/2} V_{m,\ell+1}(t) \right\|^2 &\leq C(1+t)^{2\ell} \left(\left\| A^{\ell+1/2} e^{-\frac{t}{2}A} v_0 \right\|^2 + \left\| A^{\ell+1/2} e^{-\frac{t}{2}A} u_1 \right\|^2 \right). \end{aligned}$$

Applying Lemma 2.1 with $n = 2\ell + 1$, we obtain the desired estimate. \square

4.2 Proof of higher order asymptotic expansion

Here we provide the relation between the solution u of (1.1) and the family V_m .

Lemma 4.4. Assume $(u_0, u_1) \in D(A^{1/2}) \times D(A^{1/2})$. Let $\{U_m\}_{m \in \mathbb{N}}$ be a weak solution of

$$\begin{cases} U''_{m+1} + AU_{m+1} + U'_{m+1} = -V'_m, & t > 0, \\ (U_{m+1}, U'_{m+1})(0) = (0, (-1)^{m+1}u_1). \end{cases} \quad (4.3)$$

Then $U_m = V_m + U'_{m+1}$.

Proof. The strategy of the proof is essentially the same as Lemma 3.2. By taking a suitable approximation of (u_0, u_1) , we can assume that U_m is smooth enough without loss of generality. Putting $w = V_m + U'_{m+1}$, we see from Lemma 4.1 that

$$w(0) = V_m(0) + U'_{m+1}(0) = (-1)^m u_1 + (-1)^{m+1} u_1 = 0$$

and

$$\begin{aligned} w'(t) &= V'_m(t) + U''_{m+1}(t) \\ &= -AU_{m+1}(t) - U'_{m+1}(t). \end{aligned}$$

This gives

$$\lim_{t \rightarrow 0} w'(t) = -AU_{m+1}(0) - U'_{m+1}(0) = (-1)^m u_1$$

and also

$$\begin{aligned} w''(t) &= -AU'_{m+1}(t) + (V_m(t) - w(t))' \\ &= -A(w(t) - V_m(t)) + V'_m(t) - w'(t) \\ &= -Aw(t) - w'(t) + (AV_m(t) + V'_m(t)) \\ &= -Aw(t) - w'(t) - V'_{m-1}(t), \end{aligned}$$

where we have used Lemma 4.1 at the last line. Noting the uniqueness of solution to (4.3), we obtain $w = U_m$. \square

Formally speaking, from the viewpoint of Lemmas 3.2 and 4.4 we can see that the solution u of (1.1) can be decomposed as

$$\begin{aligned}
u(t) &= V_0(t) + u(t) - V_0(t) \\
&= V_0(t) + U_1'(t) \\
&= V_0(t) + V_1(t)' + (V_1(t) - U_1(t))' \\
&= V_0(t) + V_1'(t) + U_2''(t) \\
&\vdots \\
&= \sum_{\ell=0}^m \frac{d^\ell}{dt^\ell} V_\ell(t) + \frac{d^{m+1}}{dt^{m+1}} U_{m+1}(t).
\end{aligned}$$

By virtue of (4.1) in Lemma 4.3, we already have for every $\ell = 0, \dots, m$,

$$\left\| \frac{d^\ell}{dt^\ell} V_\ell(t) \right\| \leq C(1+t)^{-\ell}.$$

To ensure that the above description is the asymptotic expansion, we finally prove the following proposition.

Proposition 4.5. *Assume that $(u_0, u_1) \in D(A^{m+1/2}) \times D(A^{m+1/2})$. Then*

$$\left\| \frac{d^{m+1}}{dt^{m+1}} U_{m+1}(t) \right\| \leq (1+t)^{-m-\frac{1}{2}}.$$

The first step of the proof of Proposition 4.5 is to show the following lemma.

Lemma 4.6. *If $(u_0, u_1) \in D(A^{1/2}) \times D(A^{1/2})$, then*

$$(1+t)E(U_{m+1}; t) + \|U_{m+1}(t)\|^2 \in L^\infty, \quad (1+t)\|U_{m+1}'(t)\|^2 + \|A^{1/2}U_{m+1}(t)\|^2 \in L^1$$

Proof. The strategy is similar to Lemma 3.3. In view of Lemma 4.4, we compute

$$\begin{aligned}
&\frac{d}{dt} \left[(6+t)E(U_{m+1}, t) + 2E_*(U_{m+1}, t) \right] \\
&= \|U_{m+1}'\|^2 + \|A^{1/2}U_{m+1}\|^2 + 2(6+t)(U_{m+1}', -U_{m+1}' - V_m') \\
&\quad + 4\|U_{m+1}'\|^2 + 4(U_{m+1}, -AU_{m+1} - V_m') \\
&= -(7+2t)\|U_{m+1}'\|^2 - 3\|A^{1/2}U_{m+1}\|^2 - 2(6+t)(U_{m+1}', V_m') + 4(U_{m+1}, AV_{m,1}) \\
&= -(1+t)\|U_{m+1}'\|^2 - \|A^{1/2}U_{m+1}\|^2 - (6+t)\|V_m'\|^2 + 2\|A^{1/2}V_{m,1}\|^2.
\end{aligned}$$

By applying Lemma 4.3 (4.2) with $\ell = 0$, the proof is complete. \square

To give extra decay property of $\frac{d^\ell}{dt^\ell} U_{m+1}$ via induction, we use the following lemma.

Lemma 4.7. *Assume $(\tilde{u}_0, \tilde{u}_1) \in D(A^{1/2}) \times D(A^{1/2})$ and $F \in C([0, \infty); D(A))$. Let w be the solution of*

$$\begin{cases} w'' + Aw + w' = AF, & t > 0, \\ (w, w')(0) = (\tilde{u}_0, \tilde{u}_1). \end{cases} \quad (4.4)$$

If w and F satisfy additional condition

$$(1+t)^{2\ell+1}\|AF(t)\|^2 + (1+t)^{2\ell}\|A^{1/2}F(t)\|^2 + (1+t)^{2\ell-1}\|w(t)\|^2 \in L^1((0, \infty)),$$

then

$$(1+t)^{2\ell+1}E(w, t) + (1+t)^{2\ell}\|w\|^2 \in L^\infty((0, \infty)), \quad (1+t)^{2\ell+1}\|w'(t)\|^2 \in L^1((0, \infty)).$$

Proof. Let $t_0 \geq 1$ be a constant determined later. Since we can use an approximation $w_\varepsilon(t) = J_\varepsilon w(t)$ with $J_\varepsilon = (I + \varepsilon A)^{-1}$, which satisfies

$$\begin{cases} w_\varepsilon'' + Aw_\varepsilon + w_\varepsilon' = J_\varepsilon AF, & t > 0, \\ (w, w')(0) = (J_\varepsilon \tilde{u}_0, J_\varepsilon \tilde{u}_1) \in D(A^{3/2}) \times D(A^{3/2}), \end{cases} \quad (4.5)$$

we may assume $w \in C^2([0, \infty); D(A^{1/2})) \cap C^1([0, \infty); D(A)) \cap C([0, \infty); D(A^{3/2}))$. By direct computation with the use of (4.4), we have

$$\begin{aligned} & \frac{d}{dt} \left[(t_0 + t)^{2\ell+1}E(w, t) + (\ell + 2)(t_0 + t)^{2\ell}E_*(w, t) \right] \\ &= (2\ell + 1)(t_0 + t)^{2\ell}E(w, t) + 2(t_0 + t)^{2\ell+1}(w', -w' + AF) \\ & \quad + 2(\ell + 2) \left[\ell(t_0 + t)^{2\ell-1}E_*(w, t) + (t_0 + t)^{2\ell}\|w'\|^2 + (t_0 + t)^{2\ell}(w, -Aw - AF) \right] \\ &\leq (t_0 + t)^{2\ell} \left[(4\ell + 5 + \frac{2\ell(\ell+1)}{t_0+t})\|w'\|^2 - 2\|A^{1/2}w\|^2 - (t_0 + t)\|w'\|^2 \right] \\ & \quad + (t_0 + t)^{2\ell+1}\|AF\|^2 + (\ell + 2)^2(t_0 + t)^{2\ell}\|A^{1/2}F\|^2 + 4\ell(\ell + 1)(t_0 + t)^{2\ell-1}\|w\|^2. \end{aligned}$$

Taking $t_0 = 6 + 4\ell + 2\ell(\ell + 1)$, we obtain the desired estimate. \square

Here we prove Proposition 4.5.

Proof of Proposition 4.5. By Lemma 4.6, we have

$$(1+t)\|U'_{m+1}(t)\|^2 \in L^1((0, \infty)).$$

Here $w = U'_{m+1}$ satisfies

$$\begin{cases} w'' + Aw + w' = -V_m'' = -A^2V_{m,2}, & t > 0, \\ (w, w')(0) = ((-1)^{m+1}u_1, -AU_{m+1}(0) - V_m'(0)) = ((-1)^{m+1}u_1, AV_{m,1}(0)) \end{cases} \quad (4.6)$$

in view of the compatibility condition. By virtue of Lemma 4.3 with $\ell = 1$, applying Lemma 4.7 with $\ell = 1$, we have

$$(1+t)^3\|U''_{m+1}(t)\|^2 = (1+t)^3\|w'(t)\|^2 \in L^1((0, \infty)).$$

Successively, we apply Lemma 4.7 with $w = \frac{d^k}{dt^k}U_{m+1}$ and $\ell = k + 1$ ($k = 1, \dots, m$). Then finally, from Lemma 4.3 we deduce

$$(1+t)^{2m+1} \left\| \frac{d^{m+1}}{dt^{m+1}}U_{m+1}(t) \right\|^2 \leq C(1+t)^{2m+1}E\left(\frac{d^m}{dt^m}U_{m+1}, t\right) \in L^\infty((0, \infty));$$

note that we need the regularity $(u_0, u_1) \in D(A^{m+1/2}) \times D(A^{m+1/2})$ to verify all deductions. The proof is complete. \square

To close the paper, we give a proof of Theorem 1.2.

Proof of Theorem 1.2. Observing the decomposition (at the beginning of this section) and Lemma 4.2, we have

$$u(t) = \sum_{\ell=0}^m (-A)^\ell V_{\ell,\ell}(t) + \frac{d^{m+1}}{dt^{m+1}} U_{m+1}(t).$$

Combining the definition of $V_{\ell,\ell}$ and Proposition 4.5, we obtain the desired assertion. \square

Acknowledgements

This work is partially supported by JSPS KAKENHI Grant Number JP18K134450.

References

- [1] H. Brezis, “Functional analysis, Sobolev spaces and partial differential equations,” Universitext. Springer, New York, 2011.
- [2] C. Cattaneo, *Sur une forme de l’équation de la chaleur éliminant le paradoxe d’une propagation instantanée*, C. R. Acad. Sci. **247** (1958), 431–433.
- [3] R. Chill, A. Haraux, *An optimal estimate for the difference of solutions of two abstract evolution equations*, J. Differential Equations **193** (2003), 385–395.
- [4] L. Hsiao, T.-P. Liu, *Convergence to nonlinear diffusion waves for solutions of a system of hyperbolic conservation laws with damping*, Comm. Math. Phys. **43** (1992), 599–605.
- [5] R. Ikehata, *Diffusion phenomenon for linear dissipative wave equations in an exterior domain*, J. Differential Equations **186** (2002), 633–651.
- [6] R. Ikehata, K. Nishihara, *Diffusion phenomenon for second order linear evolution equations*, Studia Math. **158** (2003), 153–161.
- [7] R. Ikehata, T. Matsuyama, *L^2 -behaviour of solutions to the linear heat and wave equations in exterior domains*, Sci. Math. Jpn. **55** (2002), 33–42.
- [8] A. Saeki, R. Ikehata, *Remarks on the decay rate for the energy of the dissipative linear wave equations in exterior domains*, SUT J. Math. **36** (2000), 267–277.
- [9] G. Karch, *Selfsimilar profiles in large time asymptotics of solutions to damped wave equations*, Studia Math. **143** (2000), 175–197.
- [10] H. Michihisa, *L^2 asymptotic profiles of solutions to linear damped wave equations*, <https://arxiv.org/abs/1710.04870>.
- [11] K. Nishihara, *Asymptotic behavior of solutions of quasilinear hyperbolic equations with linear damping*, J. Differential Equations **137** (1997), 384–395.
- [12] K. Nishihara, *L^p - L^q estimates of solutions to the damped wave equation in 3-dimensional space and their application*, Math. Z. **244** (2003), 631–649.

- [13] R. Orive, E. Zuazua, A.F. Pazoto, *Asymptotic expansion for damped wave equations with periodic coefficients*, Math. Models Methods Appl. Sci. **11** (2001), 1285–1310.
- [14] P. Radu, G. Todorova, B. Yordanov, *Diffusion phenomenon in Hilbert spaces and applications*, J. Differential Equations **250** (2011), 4200–4218.
- [15] P. Radu, G. Todorova, B. Yordanov, *The generalized diffusion phenomenon and applications*, SIAM J. Math. Anal. **48** (2016), 174–203.
- [16] H. Takeda, *Higher-order expansion of solutions for a damped wave equation*, Asymptot. Anal. **94** (2015), 1–31.
- [17] H. Takeda, S. Yoshikawa, *On the initial value problem of the semilinear beam equation with weak damping II: Asymptotic profiles*, J. Differential Equations **253** (2012), 3061–3080.
- [18] H. Takeda, S. Yoshikawa, *On the initial value problem of the semilinear beam equation with weak damping I: Smoothing effect*, J. Math. Anal. Appl. **401** (2013), 244–258.
- [19] H. Takeda, S. Yoshikawa, *On the decay property of solutions to the Cauchy problem of the semilinear beam equation with weak damping for large initial data*, Nonlinear dynamics in partial differential equations, 507–514, Adv. Stud. Pure Math., **64**, Math. Soc. Japan, Tokyo, 2015.
- [20] P. Vernotte, *Les paradoxes de la théorie continue de l'équation de la chaleur*, Comptes Rendus **246** (1958), 3154–3155.
- [21] T. Yamazaki, *Asymptotic behavior for abstract wave equations with decaying dissipation*, Adv. Differential Equations **11** (2006), 419–456.
- [22] H. Yang, A. Milani, *On the diffusion phenomenon of quasilinear hyperbolic waves*, Bull. Sci. Math. **124** (2000), 415–433.