

Nature of $X(3872)$ from the line shape

Xian-Wei Kang^{1,2,*}

¹ *Key Laboratory of Beam Technology of Ministry of Education, College of Nuclear Science and Technology, Beijing Normal University, Beijing 100875, China*

² *Beijing Radiation Center, Beijing 100875, China*

**E-mail: 11112018023@bnu.edu.cn*

J. A. Oller

Departamento de Física, Universidad de Murcia, E-30071 Murcia, Spain

E-mail: oller@um.es

Using a general parameterization of two-body scattering amplitude, we systematically analyze the corresponding data on $X(3872)$, more explicitly, the CDF data on inclusive $p\bar{p}$ scattering to $J/\psi\pi^+\pi^-$, and the Belle and BaBar data on B decays to $KJ/\psi\pi^+\pi^-$ and $KD\bar{D}^{*0}$ around the $D^0\bar{D}^{*0}$ threshold. We achieve a good reproduction of data and find that the $X(3872)$ can be interpreted as a bound and/or virtual state, or even higher-order (double or triple) virtual state pole. The latter point was not realized previously in the literature. As a result, the compositeness of the $X(3872)$ can vary largely from nearly 0 to 1. More higher-precision data is needed to discriminate its pole structure and nature.

Keywords: CDD pole; two-body scattering; effective range expansion; line shape, compositeness; bound and virtual state.

1. Introduction

A series of exotic hadron candidates, called XYZ particles because their properties are hard to accommodate or cannot be accounted for by conventional potential models, have been observed. A variety of theoretical interpretations have been proposed, e.g., kinematical effects, molecular states, quark-gluon hybrids, etc. We want to stress that, generally, different scenarios predict different constituents, and in practice, it may involve several mechanisms. In such circumstances the concept of compositeness is a valuable tool to obtain insight on the nature of the resonance. In most cases, those newly observed states stay close to threshold of hadron pairs^{1,2} and the effective range expansion (ERE) may be applicable. We have such a good example in Ref.³, where the scattering length and effective range of

$B\bar{B}^{(*)}$ are calculated by reproducing the pole location of Z_b states, and the molecular component of $B\bar{B}^{(*)}$ pairs are predicted. However, the ERE radius of convergence may be severely reduced in the presence of a Castillejo-Dalitz-Dyson (CDD) pole⁴.

2. Formalism and data analysis

Since the discovery of the $X(3872)$ by the Belle collaboration⁵, a voluminous amount of papers devoted to its study have appeared. Its quantum numbers have been well determined by now as $I^G J^{PC} = 0^+(1^{++})$ ⁶. Within the ERE analysis, Ref.⁷ assigns the $X(3872)$ as a pure $D\bar{D}^*$ molecule, while in Ref.⁸ the virtual-state nature is proposed.

A nearby CDD pole around threshold could spoil the applicability of ERE to study the $X(3872)$ ⁹. In the presence of a CDD pole, the scattering amplitude can be written as¹⁰

$$t(E) = \left(\frac{\lambda}{E - M_{\text{CDD}}} + \beta - ik(E) \right)^{-1}, \quad (1)$$

where the CDD pole lies at M_{CDD} , its residue is λ , β is the subtraction constant and $k(E) = \sqrt{2\mu(E + i\frac{\Gamma_*}{2})}$. Throughout the paper, energy E , M_{CDD} and pole position are defined with respect to the $D^0\bar{D}^{*0}$ threshold. The width of D^{*0} meson is $\Gamma_* \approx 65$ keV, and is taken into account in our formalism by employing a complex mass for the D^{*0} . This width may play an important role in a quantitative analysis, since the width of the $X(3872)$, Γ_X , is also very small (smaller than 1.2 MeV as determined by experiment⁶). Removing the extra zero due to the CDD pole, one ends with the function $d(E)$, which accounts for the final-state interactions,

$$d(E) = \frac{1}{1 + \frac{E - M_{\text{CDD}}}{\lambda}(\beta - ik)}. \quad (2)$$

Equation (1) can be verified in the case in which cross-channel effect¹¹ can be disregarded, so that the partial-wave amplitude only has a right-hand cut.

By comparing Eq. (1) with the ERE

$$t(E) = \frac{1}{-1/a + 1/2rk^2 - ik}, \quad (3)$$

one obtains

$$1/a = \frac{\lambda}{M_{\text{CDD}}}, \quad r = -\frac{\lambda}{\mu M_{\text{CDD}}^2}. \quad (4)$$

If M_{CDD} is far from threshold, r would have a natural size around 1 fm, and the two meson constituents dominate; otherwise when M_{CDD} is vanishing, so that the CDD pole lies very close to threshold, the compositeness will be a small number, which implies the non-molecule nature.

For the definition of compositeness (X) of a resonance, we adopt the one developed in Ref.¹², adapted to the non-relativistic near-threshold case in Ref.³. Denoting by g_k^2 the residue in the variable k of the partial-wave amplitude for a pole, either a bound state or resonance, one has that $X = -ig_k^2$ for the former, while $X = |g_k^2|$ for a resonance if $M_R > 0$ (with M_R the mass of the resonance measured from the threshold).

We are now in position to obtain the theoretical formula for the event distributions, which are also the experimentally measured observables. We define the signal function as

$$\frac{d\hat{M}}{dE} = \frac{\Gamma_X |d(E)|^2}{2\pi|\alpha|^2}, \quad (5)$$

with α introduced such that in the vicinity of the pole position $d(E) \simeq \alpha/(E - E_p)$. As done in Ref.⁷, the background term, in the energy region considered, is treated as a constant or a linear function. The event distribution $N(E)$ is then expressed by the incoherent sum of the signal part and the background. In all the cases, the overall normalization constants, the parameters characterizing the background and the parameters λ, β and M_{CDD} , which determine the signal function, are free parameters, fitted to data. In practice, however, there are too many free parameters, which prevents us to extract any definite conclusion by reproducing the experimental data. To reduce the numbers of free parameters, we discuss several interesting scenarios that drive to different nature and pole structures for the $X(3872)$ (which may also represent feasible situations for other near-threshold states). As a result the number of free parameters in $d(E)$ for each case is 1 or 2, so that one can obtain for every scenario stable fits to data. The cases considered are:

- Case 1: Using the ERE with only the scattering length, which predicts the $X(3872)$ as a pure composite bound state⁷.
- Case 2: Imposing that $t(E)$ (for $\Gamma_* = 0$) has a double-virtual pole state as the limiting zero-width case of a resonance, one can express λ and β from E_R and M_{CDD} . That is, there are two free parameters, denoted by Case 2.I (generating two virtual-state poles). Furthermore, if one imposes $M_{\text{CDD}} = -3E_R$ then a triple virtual state pole happens for $\Gamma_* = 0$, and this is denoted as Case 2.II.

Case 3: Taking into account the coupled channel effects and imposing independence of the pole position with respect to the threshold of the charged or neutral $D\bar{D}^*$ pairs, so that the self-energy contribution due to re-scattering is necessarily suppressed. Then, the compositeness is very small at the level of only a few percent, which implies the “elementary” (or “bare”) nature of this particle. Here, only E_R determines the shape of $t(E)$, but we encounter a quadratic equation with two solutions which give rise to Case 3.I and Case 3.II.

The results that follow by fitting simultaneously all the combined data are shown in Fig. 1. There one can see that all these curves describe data almost equally well. The resulting pole locations are given in Table 1, indicating the compositeness X for those cases in which we can calculate it, as discussed above.

3. Conclusion and discussion

We propose to use a more general parameterization of a two-body scattering amplitude around threshold. With this advanced tool, we derive the formula for the event distributions with the free parameters obtained by fitting to data, which is well reproduced. The pole structures are analyzed, and it turns out that the $X(3872)$ can be interpreted as either bound or virtual state, or even as an “elementary” state. That is, the current data based line shapes is not enough for determining the properties of the $X(3872)$. Let us notice especially the Case 2, which implies that the $X(3872)$ stems from a double- or triple-virtual state pole. More higher-precision data or other observables are highly desired to further discriminate between those scenarios.

References

1. L. Y. Dai, J. Haidenbauer and U. G. Meißner, Phys. Rev. D **96**, no. 11, 116001 (2017) [arXiv:1710.03142 [hep-ph]].
2. L. Y. Dai, M. Shi, G. Y. Tang and H. Q. Zheng, Phys. Rev. D **92**, 014020 (2015), [arXiv:1206.6911 [hep-ph]].
3. X. W. Kang, Z. H. Guo and J. A. Oller, Phys. Rev. D **94**, no. 1, 014012 (2016) [arXiv:1603.05546 [hep-ph]].
4. L. Castillejo, R. H. Dalitz and F. J. Dyson, Phys. Rev. **101**, 453 (1956).
5. S. K. Choi *et al.* [Belle Collaboration], Phys. Rev. Lett. **91**, 262001 (2003) [hep-ex/0309032].




Fig. 1. Fit results for the different event distributions as a function of energy. Panels a) and b) denote the mode $D^0 \bar{D}^{*0}$ from BaBar¹³ and Belle¹⁶, respectively. Panels c), d) and e) denote the mode $J/\psi \pi^+ \pi^-$ from BaBar¹⁴, Belle¹⁵, and CDF¹⁷ collaborations, respectively. The black solid lines correspond to case 1. The CDD pole contribution is included in the expression of $d(E)$, and the rest of lines refer to cases making use of this parameterization. Case 2.I is shown by the red dotted lines, and case 2.II by the brown dashed lines, while cases 3.I and 3.II are given by the blue dash-double-dotted and green dash-dotted lines, in order. Part of the lines overlap so much that it is difficult to distinguish between them in the scale of the plots.

6. C. Patrignani *et al.* (Particle Data Group), *Chin. Phys. C* **40**, 100001 (2016).
7. E. Braaten and J. Stapleton, *Phys. Rev. D* **81**, 014019 (2010) [arXiv:0907.3167 [hep-ph]].
8. C. Hanhart, Y. S. Kalashnikova, A. E. Kudryavtsev and A. V. Nefediev, *Phys. Rev. D* **76**, 034007 (2007) [arXiv:0704.0605 [hep-ph]].
9. C. Hanhart, Y. S. Kalashnikova and A. V. Nefediev, *Eur. Phys. J. A* **47**, 101 (2011) [arXiv:1106.1185 [hep-ph]].
10. X. W. Kang and J. A. Oller, *Eur. Phys. J. C* **77**, no. 6, 399 (2017) [arXiv:1612.08420 [hep-ph]].

Table 1. Fits for all the cases including the pole positions and compositeness (X).

Case	Pole position [MeV]	X
1	$-0.19^{+0.01}_{-0.01} - i 0.0325$	1.0
2.I	$-0.36^{+0.08}_{-0.10} - i 0.18^{+0.01}_{-0.02}$ $-0.70^{+0.11}_{-0.13} + i 0.17^{+0.02}_{-0.01}$	
2.II	$-0.33^{+0.04}_{-0.03} - i 0.31^{+0.02}_{-0.01}$ $-0.84^{+0.07}_{-0.05} + i 0.77^{+0.03}_{-0.04}$ $-1.67^{+0.10}_{-0.08} - i 0.49^{+0.02}_{-0.02}$	
3.I	$-0.50^{+0.04}_{-0.03}$ $-0.68^{+0.05}_{-0.03}$	$0.061^{+0.003}_{-0.002}$
3.II	$-0.51^{+0.03}_{-0.01}$ $-1.06^{+0.05}_{-0.02}$	$0.158^{+0.001}_{-0.001}$

11. J. A. Oller and E. Oset, Phys. Rev. D **60**, 074023 (1999) [hep-ph/9809337].
12. Z. H. Guo and J. A. Oller, Phys. Rev. D **93**, no. 9, 096001 (2016) [arXiv:1508.06400 [hep-ph]].
13. B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. D **77**, 011102 (2008) [arXiv:0708.1565 [hep-ex]].
14. B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. D **77**, 111101 (2008) [arXiv:0803.2838 [hep-ex]].
15. I. Adachi *et al.* [Belle Collaboration], arXiv:0809.1224 [hep-ex].
16. T. Aushev *et al.* [Belle Collaboration], Phys. Rev. D **81**, 031103 (2010) [arXiv:0810.0358 [hep-ex]].
17. T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **103**, 152001 (2009) [arXiv:0906.5218 [hep-ex]].