

# Exact solution of the 1D Dirac equation for the inverse-square-root potential $1/\sqrt{x}$

A.M. Ishkhanyan<sup>1,2</sup>

<sup>1</sup>Russian-Armenian University, Yerevan, 0051 Armenia

<sup>2</sup>Institute for Physical Research, NAS of Armenia, Ashtarak, 0203 Armenia

We introduce a new exactly solvable potential for the one-dimensional time-independent Dirac equation. This is the inverse-square-root potential  $1/\sqrt{x}$  which belongs to a biconfluent Heun class. We present the exact solution of the Dirac equation for several configurations of vector, pseudo-vector and scalar potentials. In each case, each of the two fundamental solutions that compose the general solution of the problem is written in terms of an irreducible linear combination of two Hermite functions of a scaled and shifted argument. The fundamental solutions can alternatively be written in terms of confluent hypergeometric functions. For each of the discussed field-configurations, we derive the exact equation for bound-state energy eigenvalues and construct an accurate approximation for the energy spectrum.

**PACS numbers:** 03.65.Ge Solutions of wave equations: bound states, 03.65.Pm Relativistic wave equations, 02.30.Gp Special functions

**Keywords:** Dirac equation; inverse-square-root potential; bound state; bi-confluent Heun equation; Hermite function

**1. Introduction.** Exact analytic solutions of the Dirac equation [1], which is a relativistic wave equation of fundamental importance in physics [2], are rare. In the one-dimensional case, apart from the piece-wise constant potentials and their generalizations involving the Dirac  $\delta$ -functions, one may mention the Coulomb, linear, and exponential potentials [3]

$$V = V_0 + \frac{V_1}{x}, \quad V_0 + V_1 x, \quad V_0 + V_1 e^{x/\sigma}, \quad (1)$$

for which the Dirac equation can be solved in terms of confluent hypergeometric functions, and the potential

$$V = V_0 + V_1 \tanh\left(\frac{x}{\sigma}\right) + V_2 \coth\left(\frac{x}{\sigma}\right), \quad (2)$$

for which the Dirac equation can be solved in terms of ordinary hypergeometric functions. Many potentials reported so far, e.g., the Dirac-oscillator [4,5], Woods-Saxon [6], and Hulthén [7] potentials are in fact particular cases of these four potentials that can be derived by (generally complex) specifications of the involved parameters [8]. We note that the first three potentials given by equation (1) are particular truncated cases of the classical Kratzer [9], harmonic oscillator [10], and Morse [11] potentials for the Schrödinger equation, while the fourth potential given by equation (2) in general (if  $V_1 V_2 \neq 0$  and  $V_1 \neq V_2$ ) does not belong

to a known class of Schrödinger potentials (however, the solution for this potential can be constructed using the solution of the Schrödinger equation for the classical Pöschl-Teller potential [12]; note that the solution for a truncated version of potential (2) with  $V_2 = 0$  can be written using the solution for the Eckart potential [13]).

In the present paper, we introduce a new exactly solvable potential – the inverse square root potential

$$V = V_0 + \frac{V_1}{\sqrt{x}}. \quad (3)$$

We show that the time-independent Dirac equation for this potential can be solved in terms of linear combinations of the Hermite or confluent hypergeometric functions [14] of a scaled and shifted argument.

We consider the stationary one-dimensional Dirac equation for a spin 1/2 particle of rest mass  $m$  and energy  $E$ :

$$H|\psi\rangle = E|\psi\rangle, \quad (4)$$

with Hamiltonian  $H = K + \Pi$ , where  $K$  is the kinetic energy operator, and the interaction operator  $\Pi$  stands for the potential of the external field. For the two-component spinor wave function  $\psi = (\psi_A, \psi_B)^T$ , the kinetic energy operator is written as [2]

$$K = \begin{pmatrix} mc^2 & -i\hbar \frac{d}{dx} \\ -i\hbar \frac{d}{dx} & -mc^2 \end{pmatrix}, \quad (5)$$

where  $c$  is the speed of light and  $\hbar$  is the reduced Planck's constant (we use dimensional variables since this is useful for construction of new solutions using the known ones – see below). With the general  $2 \times 2$  Hermitian potential matrix given by real functions  $V, U, W, S$ :

$$\Pi = V(x)\sigma_0 + U(x)\sigma_1 + W(x)\sigma_2 + S(x)\sigma_3 = \begin{pmatrix} V + S & U - iW \\ U + iW & V - S \end{pmatrix}, \quad (6)$$

where  $\sigma_0$  is the unit matrix and  $\sigma_{1,2,3}$  are the three Pauli matrices, equation (4) reads

$$\begin{pmatrix} V + S + mc^2 & -i\hbar \frac{d}{dx} + U - iW \\ -i\hbar \frac{d}{dx} + U + iW & V - S - mc^2 \end{pmatrix} \begin{pmatrix} \psi_A \\ \psi_B \end{pmatrix} = E \begin{pmatrix} \psi_A \\ \psi_B \end{pmatrix}. \quad (7)$$

Without affecting the physical results, one can eliminate the space component  $U$  by applying the phase transformation  $\psi \rightarrow e^{-i\varphi}\psi$  with  $c\hbar \cdot d\varphi/dx = U$ . Hence, without loss of the generality, we assume  $U = 0$ .

The resulting system of equations can be solved for several configurations of functions  $V, W, S$  of the form  $a + b/\sqrt{x}$ , e.g., for spin symmetric or pseudo-spin symmetric cases ( $S \mp V = \text{const}$  and  $W = \text{const}$ ) or for the case of primarily electrostatic interaction for which  $V = V(x)$ ,  $W, S = \text{const}$ . More general combinations are also possible.

**2. Solution for a basic field configuration.** All the mentioned cases are treated in the same manner. A basic field configuration we consider is

$$(V, W, S) = \left( V_0 + \frac{V_1}{\sqrt{x}}, W_0, S_0 + \frac{S_1}{\sqrt{x}} \right) \quad (8)$$

with arbitrary  $V_{0,1}, W_0, S_{0,1}$ . To solve the Dirac equation (7) for this configuration, we apply the Darboux transformation [15]

$$\psi_A = a_1 \frac{dw}{dx} + a_2 w, \quad \psi_B = b_1 \frac{dw}{dx} + b_2 w. \quad (9)$$

to reduce the system to a single second-order differential equation for a new dependent variable  $w(x)$ . This is achieved by putting

$$(a_1, b_1) = \left( \sqrt{V_1 - S_1}, \sqrt{V_1 + S_1} \right), \quad (10)$$

$$a_2 = \frac{ib_1 \left( E + mc^2 + S_0 - V_0 + (S_1 - V_1) F(x) \right) + a_1 W_0}{c\hbar}, \quad (11)$$

$$b_2 = -\frac{ia_1 \left( -E + mc^2 + S_0 + V_0 + (S_1 + V_1) F(x) \right) + b_1 W_0}{c\hbar}. \quad (12)$$

where  $F = 1/\sqrt{x}$ . The resulting equation reads

$$\frac{d^2 w}{dx^2} + \frac{\left( A + B F(x) + (V_1^2 - S_1^2) F(x)^2 - i c \hbar \sqrt{V_1 - S_1} \sqrt{V_1 + S_1} F'(x) \right)}{c^2 \hbar^2} w = 0, \quad (13)$$

where the prime denotes differentiation and

$$A = (E - V_0)^2 - (mc^2 + S_0)^2 - W_0^2, \quad (14)$$

$$B = -2 \left( (E - V_0) V_1 + (mc^2 + S_0) S_1 \right). \quad (15)$$

This reduction suggests two immediate observations.

First, it is seen that in the case of spin symmetry, that is when  $S - V = C_s = \text{const}$  so that  $S_1 - V_1 = 0$ , we have  $a_1 = 0$  and  $a_2 = \text{const}$ . Due to the linearity of the Dirac equation, the constant  $a_2$  can be omitted, hence, in this case  $\psi_A = w(x)$ , while  $\psi_B$  is a linear combination of  $w$  and  $w'$ . Similarly, in the pseudo-spin symmetry case  $S + V = C_p = \text{const}$ , that is when  $S_1 + V_1 = 0$ , we have  $b_1 = 0$  and  $b_2 = \text{const}$ , hence, in this case  $\psi_B = w(x)$  and  $\psi_A$  is a linear combination of  $w$  and  $w'$ .

Second, it is seen that in spin and pseudo-spin symmetry cases, when  $S_1^2 = V_1^2$ , the terms proportional to  $F(x)^2$  and  $F'(x)$  vanish in equation (13) and the equation is reduced to the Schrödinger equation for the inverse-square-root potential:

$$\frac{d^2 w}{dx^2} + \frac{1}{c^2 \hbar^2} \left( A + \frac{B}{\sqrt{x}} \right) w = 0 \quad (16)$$

with energy  $A / (2mc^2)$  and potential  $-B / (2mc^2 \sqrt{x})$ . The general solution of this equation is presented in [16]. Rewritten in terms of our parameters, one of the two independent fundamental solutions composing this general solution of the equation, for real  $A$  and  $B$ , can be presented as

$$w = e^{-y^2/2} \left( H_\nu(y) - \text{sgn}(AB) \sqrt{2\nu} H_{\nu-1}(y) \right), \quad (17)$$

where

$$y = \sqrt{\frac{\sqrt{-4A}}{c\hbar}} \left( \sqrt{x} + \frac{B}{2A} \right), \quad (18)$$

and

$$\nu = \frac{B^2}{4c\hbar(-A)^{3/2}}. \quad (19)$$

We note that a second independent fundamental solution can be constructed by the simple change  $c\hbar \rightarrow -c\hbar$  in equations (18)-(19).

Discussing the general case of field configuration (8) for arbitrary parameters  $V_{1,2}, W_0, S_{1,2}$ , a main result we report is that a fundamental solution of equation (13) is given as

$$w = e^{-y^2/2} \left( H_\nu(y) + g H_{\nu-1}(y) \right), \quad (20)$$

where

$$y = \sqrt{-2\alpha_2} \left( \sqrt{2x} + \frac{\alpha_1}{2\alpha_2} \right), \quad (21)$$

$$(\nu, g) = \left( -\frac{\alpha_1^2}{4\alpha_2} - \frac{V_1^2 - S_1^2}{2\alpha_2 c^2 \hbar^2}, \frac{\alpha_1}{\sqrt{-2\alpha_2}} - i \frac{\sqrt{V_1 - S_1} \sqrt{V_1 + S_1}}{\sqrt{-\alpha_2} c \hbar} \right), \quad (22)$$

and 
$$(\alpha_1, \alpha_2) = \left( \frac{-B}{c\hbar\sqrt{-2A}}, \frac{\sqrt{-A}}{2c\hbar} \right). \quad (23)$$

It can be checked that for  $S_1 = \pm V_1$  this solution reproduces equations (17)-(19). We note that, mathematically, the presented solution applies not only for real parameters but for arbitrary *complex* parameters  $V_{1,2}, W_0, S_{1,2}$  as well as for arbitrary complex variable  $x$ . This observation may be useful if one discusses the non-Hermitian generalization of the Dirac Hamiltonian (6).

The presented solution is derived by the reduction of equation (13) to the biconfluent Heun equation [18,19]

$$\frac{d^2u}{dz^2} + \left( \frac{\gamma}{z} + \delta + \varepsilon z \right) \frac{du}{dz} + \frac{\alpha z - q}{z} u = 0, \quad (24)$$

the solution of which can be expanded in terms of (generally non-integer order) Hermite functions of a scaled and shifted argument [20]:

$$u = \sum_{n=0}^{\infty} c_n H_{n+\gamma-\alpha/\varepsilon} \left( \sqrt{-\varepsilon/2} (z + \delta/\varepsilon) \right). \quad (25)$$

Following the approach of [21,22], we apply the transformation  $\psi = \varphi(z) u(z)$  with  $z = \sqrt{2x}$  and  $\varphi = e^{\alpha_1 z + \alpha_2 z^2}$  to show that for the inverse-square-root potential (3) the parameters of resulting biconfluent Heun equation satisfy the equations

$$\gamma = -1, \quad q^2 - \delta q + \alpha = 0 \quad (26)$$

(mathematically, this means that the regular singularity of equation (24) at  $z = 0$  is *apparent*). With this, it is shown that the series (25) terminates on the second term [20] thus resulting in a closed-form solution involving just two Hermite functions:

$$u = e^{\alpha_1 z + \alpha_2 z^2} \left( H_{\gamma-\alpha/\varepsilon} \left( \sqrt{\frac{-\varepsilon}{2}} \left( z + \frac{\delta}{\varepsilon} \right) \right) + \frac{\sqrt{-\varepsilon}(\delta - q)}{\sqrt{2\alpha}} H_{1+\gamma-\alpha/\varepsilon} \left( \sqrt{\frac{-\varepsilon}{2}} \left( z + \frac{\delta}{\varepsilon} \right) \right) \right). \quad (27)$$

Rewritten in terms of parameters of equation (13), this yields the solution (20)-(23).

As regards the *general* solution of equation (13), it can be readily checked that this solution can be written as

$$w_G = e^{-y^2/2} \left( \Phi + \frac{g}{2\nu} \frac{d\Phi}{dy} \right) \quad (28)$$

with 
$$\Phi = c_1 \cdot H_\nu(y) + c_2 \cdot F_1 \left( -\frac{\nu}{2}; \frac{1}{2}; y^2 \right), \quad (29)$$

where  $c_{1,2}$  are arbitrary constants and  ${}_1F_1$  is the Kummer confluent hypergeometric function. We conclude this section by noting that in general the involved Hermite and hypergeometric functions are not polynomials since in general the index  $\nu = \gamma - \alpha / \varepsilon$  is not an integer.

**3. Another field-configuration.** The solution of the Dirac equation for several other field-configurations is constructed in the same way – via reduction to the biconfluent Heun equation (24). For instance, for the spin symmetric configuration  $S - V = C_s = \text{const}$  and  $W = W(x)$  equations (7) reduce to

$$\frac{d^2\psi_A}{dx^2} + \frac{1}{c^2\hbar^2} \left( (E + mc^2 + C_s)(E - mc^2 - C_s - 2V) - c\hbar \frac{dW}{dx} - W^2 \right) \psi_A = 0, \quad (30)$$

$$\psi_B = \frac{i}{E + mc^2 + C_s} \left( W\psi_A - c\hbar \frac{d\psi_A}{dx} \right). \quad (31)$$

For the field-configuration

$$(V, W, S) = \left( \frac{V_1}{\sqrt{x}}, \frac{W_1}{\sqrt{x}}, C_s + \frac{V_1}{\sqrt{x}} \right), \quad (32)$$

(for simplicity, we put  $V_0 = W_0 = 0$ ), reducing equation (30) to the biconfluent Heun equation (24), we verify that the parameters of the resulting biconfluent equation fulfill equations (26). As a result, we arrive at a fundamental solution for  $\psi_A$  written as an irreducible linear combination of two Hermite functions:

$$\psi_A = e^{-y^2/2} (H_\nu(y) + gH_{\nu-1}(y)) \quad (33)$$

with 
$$y = \sqrt{-2\alpha_2} \left( \sqrt{2x} + \frac{\alpha_1}{2\alpha_2} \right) \quad (34)$$

and (compare with (22)-(23))

$$(\nu, g) = \left( -\frac{\alpha_1^2}{4\alpha_2} + \frac{W_1^2}{2\alpha_2 c^2 \hbar^2}, \frac{\alpha_1}{\sqrt{-2\alpha_2}} - \frac{W_1}{c\hbar\sqrt{-\alpha_2}} \right), \quad (35)$$

$$(\alpha_1, \alpha_2) = \left( \frac{\sqrt{2}V_1(mc^2 + C_s + E)}{c\hbar\sqrt{(mc^2 + C_s)^2 - E^2}}, \frac{\sqrt{(mc^2 + C_s)^2 - E^2}}{2c\hbar} \right). \quad (36)$$

For completeness, in this case the general solution of the Dirac equation involving two independent fundamental solutions can be written as

$$\psi_A = e^{-y^2/2} \left( \Phi + \frac{g}{2\nu} \frac{d\Phi}{dy} \right) \quad (37)$$

with

$$\Phi = c_1 \cdot H_\nu(y) + c_2 \cdot F_1 \left( -\frac{\nu}{2}; \frac{1}{2}; y^2 \right), \quad (38)$$

where  $c_{1,2}$  are arbitrary constants.

We note that for the pseudo-spin symmetry configuration  $S + V = C_p = \text{const}$  with  $V(x)$  and  $W(x)$  given by equation (32), a fundamental solution for  $\psi_B$  is constructed by the formal change  $(\psi_A, C_s) \rightarrow (\psi_B, C_p)$  and  $(V_1, W_1, E) \rightarrow (-V_1, -W_1, -E)$  in equations (33)-(36).

**4. Bound states.** The discussed field configurations may support bound states. These states are derived by demanding the wave function to vanish at the infinity and in the origin [23]. As an example, consider the spin symmetric field configuration

$$V = \frac{V_1}{\sqrt{x}}, \quad W = 0, \quad S = \frac{V_1}{\sqrt{x}}. \quad (39)$$

This is a specific configuration that belongs to both families (8) and (32). Both approaches work yielding the same result. If viewed as a particular case of the field configuration (8) with  $V_0 = W_0 = S_0 = 0$  and  $S_1 = V_1$ , we have (see equations (14),(15))

$$A = E^2 - m^2 c^4, \quad B = -2(E + mc^2)V_1, \quad (40)$$

and the general solution of the Dirac equation (7) is written as

$$\psi_A = c_1 w(c\hbar) + c_2 w(-c\hbar), \quad (41)$$

$$\psi_B = \frac{-i\hbar}{E + mc^2} \frac{d\psi_A}{dx}, \quad (42)$$

where  $c_{1,2}$  are arbitrary constants and the function  $w$  is given through parameters  $A$  and  $B$  by equations (17)-(19).

The requirement for the wave function to vanish in the origin leads to a linear relation between coefficients  $c_1$  and  $c_2$ , while the second condition of vanishing the wave function at the infinity, after matching the leading asymptotes of the involved Hermite functions [24], results in the following transcendental equation for the index  $\nu$ :

$$H_\nu(-\sqrt{2\nu}) - \text{sgn}(AB)\sqrt{2\nu}H_{\nu-1}(-\sqrt{2\nu}) = 0. \quad (43)$$

Given  $\nu$  depends on  $E$ , this is the exact equation for the energy spectrum. This type of spectrum equations that involve two Hermite functions are faced in several other physical situations (see, e.g., [16,25,26]). For  $\text{sgn}(AB) = -1$  this is exactly the equation encountered when solving the Schrödinger equation for the inverse-square-root potential [16]. It has been shown that the equation possesses a countable infinite set of discrete positive roots  $\nu_n$ ,  $n \in \mathbb{N}$ . Together with the definition (19) for  $\nu$  and equations (40) for parameters  $A$  and  $B$ , this set determines the bound-state energy eigenvalues. We note that all  $\nu_n$  are not integers so that the bound-state wave-functions are not polynomials.

The calculation lines are as follows. Substituting equations (40), into equation (19), one arrives at the following cubic equation for energy  $E_n$ :

$$c^2 \hbar^2 (E_n - mc^2)^3 \nu_n^2 + (E_n + mc^2) V_1^4 = 0. \quad (44)$$

The discriminant  $D = -4c^2 \hbar^2 \nu_n^2 V_1^8 (27m^2 c^6 \hbar^2 \nu_n^2 + V_1^4)$  of this equation is always negative, hence, the cubic has only one real root [14]. This root is conveniently written through the dimensionless parameter

$$\theta = \frac{V_1^2}{3^{3/2} m \hbar c^3 \nu_n}. \quad (45)$$

The result reads

$$E_n = mc^2 + 3mc^2 |\theta|^{2/3} \frac{(\sqrt{\theta^2 + 1} - 1)^{2/3} - |\theta|^{2/3}}{(\sqrt{\theta^2 + 1} - 1)^{1/3}}. \quad (46)$$

Given  $\nu_n$  are known via equation (43), this is the exact expression for the energy spectrum.

To approximately solve equation (43), we note that the arguments and indexes of the involved Hermite functions  $H_\nu(z)$  belong to the *left* transient layer for which  $z \approx -\sqrt{2\nu + 1}$  [14,24]. An appropriate Airy-function approximation for this layer, where the argument of the Hermite function is negative, is suggested, e.g., in [27]. Following the approach of [16], we divide equation (43) by  $\sqrt{2\nu} H_{\nu-1}(-\sqrt{2\nu})$  and apply this approximation to show that the equation has solutions only if  $\text{sgn}(AB) = -1$ . Furthermore, the resulting approximation for the latter case reads

$$F \equiv 1 + \frac{H_\nu(-\sqrt{2\nu})}{\sqrt{2\nu} H_{\nu-1}(-\sqrt{2\nu})} \approx f(\nu) \left( \sin(\pi(\nu + 1/6)) + \frac{D_0}{\nu^{2/3}} \sin(\pi(\nu - 1/6)) \right), \quad (47)$$



where the pre-factor  $f(\nu)$  is a non-oscillatory function which does not adopt zero and  $D_0$  is a constant given as

$$D_0 = \frac{\Gamma(1/3)}{12\sqrt[3]{3}\Gamma(2/3)} \approx 0.11. \quad (48)$$

This is a highly accurate approximation. The accuracy is demonstrated in Fig. 1, where the filled circles stand for the exact values and the solid curves present the approximation.




Fig. 1. Approximation (47) compared with the exact function  $F$ . Filled circles - exact result, solid curves - approximation. For  $\nu < 1/2$  the function does not possess roots.

Thus, the exact energy spectrum equation (43) is accurately approximated by the transcendental equation

$$\frac{\sin(\pi(\nu + 1/6))}{\sin(\pi(\nu - 1/6))} + \frac{D_0}{\nu^{2/3}} = 0. \quad (49)$$

Since  $D_0$  is a small constant and  $\nu > 1/2$  (the smallest root is  $\nu_1 \approx 0.86$  see Fig.1), the second term of this equation is always small, hence, it can be treated as a perturbation. This leads to a simple, yet, highly accurate approximation

$$\nu_n \approx n - \frac{1}{6} + \frac{\sqrt{3}D_0}{2\pi(n-1/6)^{2/3}} - \frac{\sqrt{3}D_0^2}{4\pi(n-1/6)^{4/3}}, \quad n = 1, 2, 3, \dots \quad (50)$$

The relative error of this approximation is less than  $10^{-4}$  for all orders  $n \geq 2$  and the absolute error exceeds  $10^{-4}$  only for the first root with  $n = 1$ .

For large  $n$  we have  $\nu_n \approx n - 1/6$ . This result corresponds to the semiclassical limit for the solution of the Schrödinger equation for the inverse-square-root potential (3). It corresponds to the Maslov index equal to  $-1/6$  [27,27]. This index can be calculated by the

formula (4.47) by Quigg and Rosner [27] taking into account that for potential (3) the effective angular momentum is zero.

With exact equation (46), keeping just the first term  $\nu_n \approx n - 1/6$  in equation (50), the energy eigenvalues are expanded in terms of the quantum number  $n$  as

$$E_n \approx mc^2 \left( 1 - \frac{2}{(6n-1)^{2/3}} \left( \frac{3V_1^2}{c^3 m \hbar} \right)^{2/3} + \frac{2/3}{(6n-1)^{4/3}} \left( \frac{3V_1^2}{c^3 m \hbar} \right)^{4/3} \right). \quad (51)$$

This provides a qualitatively good description of the whole spectrum if  $V_1^2 / (m \hbar c^3) \leq 1$ . The approximation is compared with the exact result for the first seven levels in Table 1.

$n$	1	2	3	4	5	6	7
$E_n$ (exact)	-0.07534	0.279904	0.438093	0.530429	0.592080	0.636685	0.670730
$E_n$	-0.08538	0.276806	0.436756	0.529710	0.591638	0.636390	0.670521

Table 1. Comparison of approximation (51) with the exact formula (46) for  $V_1^2 / (m \hbar c^3) = 1$ :

$$(m, \hbar, c, V_1) = (1, 1, 1, -1).$$

**5. Bound states for the case of electrostatic potential.** As a second example consider the bound states of a Dirac particle in the presence of only electrostatic potential:

$$V = \frac{V_1}{\sqrt{x}}, \quad W = S = 0. \quad (52)$$

This is another particular case of the field configuration (8). Here,  $V_0 = W_0 = S_0 = S_1 = 0$  so that (see equations (14),(15))

$$A = E^2 - m^2 c^4, \quad B = -2EV_1 \quad (53)$$

and the general solution of the Dirac equation (7) is written as

$$\psi_A = \frac{dw_G}{dx} + \frac{i}{c\hbar} \left( E + mc^2 - \frac{V_1}{\sqrt{x}} \right) w_G, \quad (54)$$

$$\psi_B = \frac{dw_G}{dx} + \frac{i}{c\hbar} \left( E - mc^2 - \frac{V_1}{\sqrt{x}} \right) w_G, \quad (55)$$

where the function  $w_G$  is the general solution (28),(29) and  $y, \nu, g$  are given by equations (21)-(23). After a straightforward simplification, we have the following explicit result

$$\psi_A = e^{-\frac{y^2}{2}} \left( (E + mc^2 - 2i\hbar\alpha_2) \Phi + \frac{g}{2\nu} (E + mc^2 + 2i\hbar\alpha_2) \frac{d\Phi}{dy} \right), \quad (56)$$

$$\psi_B = e^{-\frac{y^2}{2}} \left( (E - mc^2 - 2i\hbar\alpha_2)\Phi + \frac{g}{2\nu}(E - mc^2 + 2i\hbar\alpha_2)\frac{d\Phi}{dy} \right), \quad (57)$$

where

$$\Phi = c_1 \cdot H_\nu(y) + c_2 \cdot F_1\left(-\frac{\nu}{2}; \frac{1}{2}; y^2\right). \quad (58)$$

Proceeding essentially in the same manner as in the previous case, the requirement of vanishing the wave function at the infinity, by matching the leading asymptotes, gives a simple linear relation between  $c_1$  and  $c_2$ :

$$2^\nu \Gamma\left(\frac{\nu+1}{2}\right)c_1 + i^\nu \sqrt{\pi}c_2 = 0, \quad (59)$$

while the requirement of vanishing in the origin results in the transcendental equation

$$\operatorname{Re} \left( \frac{\sqrt{\pi} \left(\frac{i}{2}\right)^\nu}{\Gamma\left(\frac{\nu+1}{2}\right)} \right) = \operatorname{Re} \left( \frac{{}_1F_1\left(-\frac{\nu}{2}; \frac{1}{2}; \frac{2\nu E^2}{m^2 c^4}\right) + \frac{2\nu E}{mc^2} {}_1F_1\left(1 - \frac{\nu}{2}; \frac{3}{2}; \frac{2\nu E^2}{m^2 c^4}\right)}{H_\nu\left(-\frac{\sqrt{2\nu E}}{mc^2}\right) + \sqrt{2\nu} H_{\nu-1}\left(-\frac{\sqrt{2\nu E}}{mc^2}\right)} \right), \quad (60)$$

where  $\Gamma$  is the Euler gamma-function and

$$\nu = -\frac{m^2 c^3 V_1^2 / \hbar}{(m^2 c^4 - E^2)^{3/2}}. \quad (61)$$

This is the exact equation for the energy spectrum.

Though rather complicated, equation (60) admits considerable simplification through the standard approximations for the Hermite function  $H_\nu$  [24] and the Kummer function  ${}_1F_1$  [14] if  $\nu$  is large. This is achieved by noting that the left-hand side of the equation is (exactly) equal to  $f_0 \cos^2(\pi\nu/2)$ , where  $f_0 = \sqrt{\pi} 2^{\nu+1/2} / \sqrt{-\nu} \Gamma(-\nu/2)$ . Dividing then the equation by  $f_0$ , and applying the known approximations, we reveal that the right-hand side of the resulting equation is approximated as  $\cos^2(\pi\nu/2) + g_0 \sin(\pi\nu + \Delta)$ , where  $g_0$  is a function that does not adopt zero and  $\Delta = \Delta(\nu)$  is a small quantity, less than 1, that exponentially vanishes at large  $\nu$ . Hence, we arrive at the simple equation

$$\sin(\pi(\nu + \Delta)) \approx 0 \quad (62)$$

with a small  $|\Delta = \Delta(\nu)| < 1$ . Neglecting the  $\Delta$ -term, we have  $\nu \approx -n$ . More accurately, the term can be treated iteratively with the first-order result

$$\nu \approx -n - \Delta(-n), \quad n = 1, 2, 3, \dots \quad (63)$$

With this, from equation (61), we have the approximate spectrum

$$E_n \approx mc^2 \sqrt{1 - \left( \frac{V_1^2}{mc^3 \hbar} \right)^{2/3} \frac{1}{(n + \Delta)^{2/3}}}. \quad (64)$$

This is a rather accurate approximation for all orders if  $V_1^2 / (m\hbar c^3) \leq 1$ . Even if  $\Delta$  is neglected, the relative error is less than  $10^{-3}$  for all orders  $n > 3$ . A more accurate result can be achieved if  $\Delta$  is expanded in terms of powers of  $n^{-2/3}$ . The accuracy of this approach with the first two correction terms is shown in [Table 2](#).

$n$	1	2	3	4	5	6	7
$E_n$ (exact)	0.297679	0.618900	0.723684	0.777986	0.811903	0.835392	0.852768
$E_n$	0.297679	0.619399	0.723691	0.777942	0.811863	0.835362	0.852747

Table 2. Comparison of approximation (64) for  $\Delta = \Delta_1 / n^{-2/3} + \Delta_2 / n^{-4/3}$  with the exact result for  $V_1^2 / (m\hbar c^3) = 1$ :  $(m, \hbar, c, V_1) = (1, 1, 1, -1)$ ,  $\Delta_{1,2} = (0.0316, 0.1177)$ .

Comparing the presented spectrum with that for the solution of the Schrödinger equation for the inverse-square-root potential [16], we note that the main difference is that here the Maslov index  $\Delta$  goes to zero as the quantum number  $n$  goes to infinity (we recall that in the Schrödinger case the limit Maslov index is  $-1/6$ ).

## Acknowledgments

The work was supported by the Armenian Committee of Science (SCS Grants No. 18RF-139 and No. 18T-1C276) and the Russian-Armenian (Slavonic) University at the expense of the Ministry of Education and Science of the Russian Federation.

## References

1. P.A.M. Dirac, "The quantum theory of the electron", Proc. Roy. Soc (London) A **117**, 610-624 (1928).
2. W. Greiner, *Relativistic Quantum Mechanics. Wave equations* (Springer, Berlin, 2000).
3. V.G. Bagrov and D.M. Gitman, *The Dirac Equation and its Solutions*, (de Gruyter, Boston 2014)
4. P.A. Cook, "Relativistic harmonic oscillators with intrinsic spin structure", Lett. Nuovo Cimento **1**, 419-426 (1971)
5. M. Moshinsky and A. Szczepaniak, "The Dirac oscillator", J. Phys. A **22**, L817-L819 (1989).
6. P. Kennedy, "The Woods-Saxon potential in the Dirac equation", J. Phys. A **35**, 689-698 (2002).

7. J.Y. Guo, Y. Yu, and S.W. Jin, "Transmission resonance for a Dirac particle in a one-dimensional Hulthén potential", *Cent. Eur. J. Phys.* **7**, 168-174 (2009).
8. A.M. Ishkhanyan, "Schrödinger potentials solvable in terms of the general Heun functions", *Ann. Phys.* **388**, 456-471 (2018).
9. A. Kratzer, "Die ultraroten Rotationsspektren der Halogenwasserstoffe", *Z. Phys.* **3**, 289-307 (1920).
10. E. Schrödinger, "Quantisierung als Eigenwertproblem (Erste Mitteilung)", *Annalen der Physik* **76**, 361-376 (1926).
11. P.M. Morse, "Diatomic molecules according to the wave mechanics. II. Vibrational levels", *Phys. Rev.* **34**, 57-64 (1929).
12. G. Pöschl, E. Teller, "Bemerkungen zur Quantenmechanik des anharmonischen Oszillators", *Z. Phys.* **83**, 143-151 (1933).
13. C. Eckart, "The penetration of a potential barrier by electrons", *Phys. Rev.* **35**, 1303-1309 (1930).
14. F.W.J. Olver, D.W. Lozier, R.F. Boisvert, and C.W. Clark (eds.), *NIST Handbook of Mathematical Functions* (Cambridge University Press, New York, 2010).
15. G. Darboux, "Sur une proposition relative aux équations linéaires", *Compt. Rend. Acad. Sci. Paris* **94**, 1456-1459 (1882).
16. A.M. Ishkhanyan, "Exact solution of the Schrödinger equation for the inverse square root potential  $V_0 / \sqrt{x}$ ", *Eur. Phys. Lett.* **112**, 10006 (2015).
17. R.R. Hartmann and M.E. Portnoi, "Quasi-exact solution to the Dirac equation for the hyperbolic-secant potential", *Phys. Rev. A* **89**, 012101 (2014).
18. A. Ronveaux (ed.), *Heun's Differential Equations* (Oxford University Press, London, 1995).
19. S.Yu. Slavyanov and W. Lay, *Special functions* (Oxford University Press, Oxford, 2000).
20. T.A. Ishkhanyan and A.M. Ishkhanyan, "Solutions of the bi-confluent Heun equation in terms of the Hermite functions", *Ann. Phys.* **383**, 79-91 (2017).
21. A. Ishkhanyan and V. Krainov, "Discretization of Natanzon potentials", *Eur. Phys. J. Plus* **131**, 342 (2016).
22. A.M. Ishkhanyan, "Schrödinger potentials solvable in terms of the confluent Heun functions", *Theor. Math. Phys.* **188**, 980-993 (2016).
23. M. Znojil, "Comment on "Conditionally exactly soluble class of quantum potentials", *Phys. Rev. A* **61**, 066101 (2000).
24. G. Szegő, *Orthogonal Polynomials*, 4th ed. (Amer. Math. Soc., Providence, 1975).
25. A.S. de Castro, "Comment on "Fun and frustration with quarkonium in a 1+1 dimension," by R.S. Bhalerao and B. Ram, *Am. J. Phys.* **69**, 817-818 (2001)", *Am. J. Phys.* **70**, 450-451 (2002).
26. R.M. Cavalcanti, "Comment on "Fun and frustration with quarkonium in a 1+1 dimension," by R.S. Bhalerao and B. Ram, *Am. J. Phys.* **69**, 817-818 (2001)", *Am. J. Phys.* **70**, 451-452 (2002).
27. A.M. Ishkhanyan and V.P. Krainov, "Maslov index for power-law potentials", *JETP Lett.* **105**, 43-46 (2017).
28. C. Quigg and J.L. Rosner, "Quantum mechanics with applications to quarkonium", *Phys. Rep.* **56**, 167-235 (1979).