

A RECTIFIABILITY RESULT FOR FINITE-PERIMETER SETS IN CARNOT GROUPS

SEBASTIANO DON, ENRICO LE DONNE, TERHI MOISALA, AND DAVIDE VITTONÉ

ABSTRACT. In the setting of Carnot groups, we are concerned with the rectifiability problem for subsets that have finite sub-Riemannian perimeter. We introduce a new notion of rectifiability that is, possibly, weaker than the one introduced by Franchi, Serapioni, and Serra Cassano. Namely, we consider subsets Γ that, similarly to intrinsic Lipschitz graphs, have a cone property: there exists an open dilation-invariant subset C whose translations by elements in Γ don't intersect Γ . However, a priori the cone C may not have any horizontal directions in its interior. In every Carnot group, we prove that the reduced boundary of every finite-perimeter subset can be covered by countably many subsets that have such a cone property. The cones are related to the semigroups generated by the horizontal half-spaces determined by the normal directions. We further study the case when one can find horizontal directions in the interior of the cones, in which case we infer that finite-perimeter subsets are countably rectifiable with respect to intrinsic Lipschitz graphs. A sufficient condition for this to hold is the existence of a horizontal one-parameter subgroup that is not an abnormal curve. As an application, we verify that this property holds in every filiform group, of either first or second type.

1. INTRODUCTION

The celebrated rectifiability theorem by De Giorgi, see [DG54, DG55], states that the reduced boundary of a set of finite perimeter in the Euclidean space \mathbb{R}^n is C^1 -rectifiable, i.e., it can be covered, up to a negligible set with respect to the Hausdorff measure \mathcal{H}^{n-1} , by a countable union of C^1 hypersurfaces. The proof of this theorem relies on the fact that the blow-up of a set of finite perimeter at a point of its reduced boundary is a set with constant normal, and each constant-normal set in \mathbb{R}^n is a half-space. The importance of having sufficiently regular sets of finite perimeter is evident on many classical key problems in Geometric Measure Theory and underlines the relevance of the notion of rectifiability in this context. A wide impact can be, for example, detected in developing a sufficiently rich theory for functions of bounded variation; see e.g. the monographs [Fed69, GMS98, AFP00, Mag12, EG15].

In the more general context of metric measure spaces, the regularity of finite-perimeter sets and the structure of some suitable notions of their boundaries has been object of several studies in the last decades. We refer to this task as the *rectifiability problem*.

2010 *Mathematics Subject Classification.* 53C17, 22E25, 49Q15, 28A75.

Key words and phrases. Sets of finite perimeter, Carnot groups, rectifiability, filiform groups, outer cone property, semigroups of horizontal normal, intrinsic Lipschitz graphs, abnormal directions.

S.D., E.L.D. and T.M. have been partially supported by the Academy of Finland (grant 288501 “*Geometry of subRiemannian groups*”) and by the European Research Council (ERC Starting Grant 713998 GeoMeG “*Geometry of Metric Groups*”). T.M. was also supported by the Finnish Cultural Foundation (grant 00170709). D.V. is supported by the University of Padova STARS Project “Sub-Riemannian Geometry and Geometric Measure Theory Issues: Old and New” (SUGGESTION), and by GNAMPA of INdAM (Italy) project “Applicazioni della Teoria delle Correnti all’Analisi Reale e al Trasporto Ottimo”.

In the current paper, we study the rectifiability problem in the setting of Carnot groups. A Carnot group \mathbb{G} of step $s \in \mathbb{N}$ is a connected and simply connected Lie group whose Lie algebra \mathfrak{g} is stratified into s layers, i.e., it is linearly decomposed as $\mathfrak{g} = \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_s$ with

$$[\mathfrak{g}_1, \mathfrak{g}_i] = \mathfrak{g}_{i+1} \text{ for } i = 1, \dots, s-1, \quad \mathfrak{g}_s \neq \{0\} \quad \text{and} \quad [\mathfrak{g}_1, \mathfrak{g}_s] = \{0\}.$$

We refer the reader to [FS82, LD17] for an introduction to Carnot groups. Canonically, every Carnot group has a one-parameter family of dilations that we denote by $\{\delta_t : t \geq 0\}$. We further fix a homogeneous distance d on \mathbb{G} , which is unique up to biLipschitz equivalence. In this setting the notions of perimeter and constant-normal set can be naturally defined (see Definitions 2.1 and 2.3). The original result of De Giorgi has been generalized by Franchi, Serapioni and Serra Cassano first in Heisenberg groups [FSSC01] and then in all step-2 Carnot groups [FSSC03]. The same result holds in the so-called type \star Carnot groups, which are Carnot groups satisfying a suitable algebraic condition, see [Mar14], which generalizes the step-2 condition but may hold in arbitrary step. These are the only classes of Carnot groups in which the rectifiability problem has been solved in a satisfactory way, so far.

In general, only few partial results are known. In arbitrary groups, the first delicate issue concerns the blow-up analysis of the reduced boundary of a set of finite perimeter. By [FSSC01] every tangent set at a point of the reduced boundary is a set with constant horizontal normal, but this is not enough to prove that the tangent is a half-space, in general. Indeed, in the Engel group (the simplest Carnot group of step 3) there are examples of sets with constant horizontal normal that are not half-spaces; see [FSSC03, Example 3.2]. However, in the paper [AKLD09], the authors prove that, among all the possible blow-ups of a set of constant horizontal normal in a Carnot group, there is always a half-space.

Another issue is to understand what is the correct notion of rectifiable set in Carnot groups. Namely, at the moment it is not known which kind of rectifiability property one should expect for finite-perimeter sets in these spaces. It is well-known that in the Euclidean setting, three equivalent notions of codimension-1 rectifiability are available: the countable covering family can be composed by C^1 hypersurfaces, Lipschitz images of sets in \mathbb{R}^{n-1} or Lipschitz codimension-1 graphs; see e.g. [Mat75, Mat95] for an account of rectifiability theory in the Euclidean spaces. Actually, a very natural notion of rectifiability, via Lipschitz images of open subsets of Euclidean spaces, was given in the setting of metric spaces already by Federer in [Fed69] (and later by Ambrosio and Kirchheim in [AK00a]). Unfortunately, this notion does not fit in the geometric structure of a Carnot group since, according to this definition, already the Heisenberg group is purely unrectifiable (see [AK00b]).

Nonetheless, a definition of rectifiability using a suitable notion of Lipschitz graphs or C^1 hypersurfaces can still be fruitful in the setting of Carnot groups. For this purpose, Franchi Serapioni and Serra Cassano introduced the notions of intrinsic Lipschitz graphs (see Definition 1.1) and of intrinsic C^1 hypersurfaces. We know that the notion of rectifiability with respect to intrinsic C^1 hypersurfaces implies the one with intrinsic Lipschitz graphs; see e.g. [Vit12, Theorem 3.2]. This stronger notion of rectifiability was used in [FSSC01, FSSC03, Mar14]. More precisely, the authors proved that the reduced boundary of a set of finite perimeter in a type \star group \mathbb{G} can be covered, up to a set of \mathcal{H}^{Q-1} -measure zero, by a countable union of intrinsic C^1 hypersurfaces, where Q is the Hausdorff dimension of \mathbb{G} .

It is not known whether the possibly weaker notion of rectifiability, obtained by replacing the intrinsic C^1 hypersurfaces by intrinsic Lipschitz graphs, leads to an equivalent definition. In fact, the validity of an intrinsic Rademacher-type theorem is still an unsolved problem. One of the aim

of this paper is to discuss another form of rectifiability that may be a priori even weaker than the intrinsic Lipschitz rectifiability.

The notion of intrinsic Lipschitz graph appeared in different equivalent forms in [FSSC06, FSSC11, FS16] and we briefly recall it here (see also Section 5 for a more complete discussion) in a way that is suitable for our purposes.

Definition 1.1. Let \mathbb{G} be a Carnot group and let $\mathbb{W}, \mathbb{L} \subseteq \mathbb{G}$ be homogeneous subgroups of \mathbb{G} such that $\mathbb{G} = \mathbb{W} \cdot \mathbb{L}$ and $\mathbb{W} \cap \mathbb{L} = \{0\}$. We say that $\Sigma \subseteq \mathbb{G}$ is an (entire) *intrinsic Lipschitz graph* if there exists $\beta > 0$ such that

(i) for every $p \in \Sigma$ one has

$$(1) \quad \Sigma \cap p \cdot \bigcup_{\ell \in \mathbb{L} \setminus \{0\}} B(\ell, \beta d(0, \ell)) = \emptyset;$$

(ii) $\pi_{\mathbb{W}}(\Sigma) = \mathbb{W}$.

The set $\bigcup_{\ell \in \mathbb{L}} B(\ell, \beta d(0, \ell))$ is a homogeneous cone around \mathbb{L} of aperture given by parameter β , while $\pi_{\mathbb{W}}$ is the canonical projection associated with the decomposition $\mathbb{G} = \mathbb{W} \cdot \mathbb{L}$. The cones introduced above are not the ones that are often used in the literature of Carnot groups, but they produce the same notion of intrinsic Lipschitz graph (see e.g. [DMV19, Remark A.2])

One of the goals of the current paper is to analyze the geometry of sets that possess a “cone-behavior” that resembles property (1) in the following weaker sense.

Definition 1.2. A non-empty set $C \subseteq \mathbb{G}$ is said to be a *cone* if

$$\delta_r C = C, \quad \text{for any } r > 0.$$

Definition 1.3. Let \mathbb{G} be a Carnot group. We say that a set $\Gamma \subseteq \mathbb{G}$ satisfies an (*outer*) *cone property* if there exists a cone $C \subseteq \mathbb{G}$ such that

$$\Gamma \cap pC = \emptyset, \quad \text{for every } p \in \Gamma.$$

A dilation-invariant set with nonempty interior, which turns out to be related with sets of locally finite perimeter, is the following semigroup. This notion has been introduced in [BLD13, BLD19] in the study of constant-normal sets in Carnot groups. (We point out that in [BLD19] the authors are mainly interested in an *inner cone property* of sets as opposed to the outer cone property.)

Definition 1.4. Let \mathbb{G} be a Carnot group. For any $\nu \in \mathfrak{g}_1 \setminus \{0\}$, the *semigroup of horizontal normal* ν is defined by

$$S_\nu := S(\exp(\nu^\perp + \mathbb{R}^+ \nu)),$$

where for any $A \subseteq \mathbb{G}$, the set $S(A) := \bigcup_{k=1}^{\infty} A^k$ is the semigroup generated by A . Here ν^\perp denotes the orthogonal space to ν within \mathfrak{g}_1 with respect to some scalar product that we fixed on \mathfrak{g}_1 .

Semigroups with horizontal normal are cones and, by a standard argument of Geometric Control Theory (see [AS13]), they have non-empty interior. It can be also proved (see Proposition 3.1) that any semigroup of horizontal direction ν has ν as constant horizontal normal. We first point out that semigroups of horizontal normal ν are minimal in the following sense: S_ν is contained in every set with ν as constant horizontal normal and for which the identity element 0 of \mathbb{G} has positive density, see [BLD19, Corollary 2.31 or Theorem 2.37]. This property, together with the dilation-invariance of such semigroups, allows us to perform a fruitful blow-up procedure and get our main result; see Theorem 4.6.

Theorem 1.5. *Let \mathbb{G} be a Carnot group and let $E \subseteq \mathbb{G}$ be a set of locally finite perimeter. Then there exists a family $\{C_h : h \in \mathbb{N}\}$ of open cones in \mathbb{G} and a family $\{\Gamma_h : h \in \mathbb{N}\}$ of subsets of \mathbb{G} such that each Γ_h satisfies the C_h -cone property (as in Definition 1.3) and*

$$\mathcal{F}E = \bigcup_{h \in \mathbb{N}} \Gamma_h,$$

where $\mathcal{F}E$ denotes the reduced boundary of E .

Notice that the previous result is obtained without requiring any assumption on the Carnot group \mathbb{G} . The first natural question one may ask is when a set satisfying a cone property is also an intrinsic Lipschitz graph. We can point out a sufficient condition. Indeed, since each C_h in the theorem above comes as a small shrinking of some semigroup S_ν , it is enough that S_ν is such that there exists $X \in \mathfrak{g}_1 \setminus \{0\}$ with $\exp(X) \in \text{int}(S_\nu)$ (see Remark 5.1 for more details). In Section 5 we find some conditions on the group \mathbb{G} that are sufficient to conclude that the reduced boundary of every set of finite perimeter in \mathbb{G} is intrinsically Lipschitz rectifiable (see Definition 5.2). In particular, we notice that, whenever the group possesses a non-abnormal horizontal direction (see Definition 5.9), then each set Γ_h appearing in Theorem 1.5 can be chosen to be an intrinsic Lipschitz graph (see Proposition 5.3 and Corollary 5.7). The notions of normal and abnormal curve naturally appeared in Geometric Control Theory (see [Mon02]), and in the study of regularity of geodesics in sub-Riemannian manifolds (see [Vit14]). In the paper [LDMO⁺16], it is proved that the one-parameter subgroup generated by $X \in \mathfrak{g}_1$ is non-abnormal if and only if

$$(2) \quad \text{span}\{\text{ad}_X^k(\mathfrak{g}_1) : k = 0, \dots, s-1\} = \mathfrak{g},$$

where the adjoint of X is defined by $\text{ad}_X^0(Y) := Y$ and $\text{ad}_X^k(Y) := \text{ad}_X^{k-1}([X, Y])$, for every $k \geq 1$ and every $Y \in \mathfrak{g}$. This gives us a purely algebraic sufficient condition on the group \mathbb{G} for the intrinsic Lipschitz rectifiability of reduced boundaries of sets of finite perimeter.

Theorem 1.6. *Let \mathbb{G} be a Carnot group and assume there exists $X \in \mathfrak{g}_1$ such that (2) holds. Then, the reduced boundary of every set of finite perimeter in \mathbb{G} is intrinsically Lipschitz rectifiable. In particular, this result can be applied to all filiform Carnot groups.*

For the second part of Theorem 1.6 see Section 6. Among filiform groups, we can find the Engel group, which is the simplest Carnot group of step 3. We also point out that possessing a non-abnormal horizontal direction is stable under direct products and quotients. More precisely, if \mathbb{G}_1 and \mathbb{G}_2 are two Carnot groups that possess a non-abnormal horizontal direction, and N is a normal subgroup of \mathbb{G}_1 , then both $\mathbb{G}_1 \times \mathbb{G}_2$ and \mathbb{G}_1/N possess a non-abnormal horizontal direction (see Propositions 5.13 and 5.14).

On the negative side, we point out that for example in the free group $\mathbb{F}_{2,3}$ of rank 2 and step 3, also known as Cartan group, all horizontal directions are abnormal and

$$\exp(\mathfrak{f}_1) \cap \text{int}(S_\nu) = \emptyset, \quad \text{for every } \nu \in \mathfrak{f}_1 \setminus \{0\},$$

where \mathfrak{f}_1 denotes the horizontal layer of the Lie algebra of $\mathbb{F}_{2,3}$ (see [BLD19]).

As an example of application, we remark that in [DV19], to study some properties of functions of bounded variation, the authors consider sub-Riemannian structures in which sets of finite perimeter have reduced boundary that is intrinsically Lipschitz rectifiable.

The outline of the paper is the following. Section 2 is devoted to the basic notions and facts related to Carnot groups. Section 3 is devoted to studying properties of closed semigroups of horizontal direction that will be crucial for the main theorem. Section 4 contains the proof of Theorem 1.5. Section 5 contains the definitions of intrinsic Lipschitz graphs, intrinsic Lipschitz

rectifiability and the proof of Theorem 1.6 together with applications to some classes of Carnot groups. In Section 6 we introduce filiform groups and determine their abnormal lines and their graded Lie algebra automorphisms.

2. PRELIMINARIES

For an introduction to Carnot groups we refer the reader to [FS82, LD17], while for a theory of sets of finite perimeter in Carnot groups, we refer to [AKLD09]. In what follows, let \mathbb{G} be a Carnot group of dimension n , let $\mathfrak{g} = \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_s$ be its stratified Lie algebra and denote by $m := \dim \mathfrak{g}_1$ the rank of \mathbb{G} . We fix a scalar product $\langle \cdot, \cdot \rangle$ on the horizontal layer \mathfrak{g}_1 of \mathfrak{g} and a left-invariant Haar measure μ on \mathbb{G} . We endow \mathbb{G} with the usual Carnot-Carathéodory metric d and we denote by B_r the metric ball of radius r at the identity element of \mathbb{G} .

It is well-known that there exists a family $\{\delta_r \in \mathbb{G}^{\mathbb{G}} : r \geq 0\}$ such that $\delta_0 := 0$ and, for any $r > 0$, δ_r is a graded diagonalizable automorphism of \mathbb{G} satisfying the following properties. For any $r, s \geq 0$, $\delta_{rs} = \delta_r \circ \delta_s$ and, for every $x, y \in \mathbb{G}$ and $r \geq 0$

$$d(\delta_r(x), \delta_r(y)) = rd(x, y).$$

Definition 2.1. Let Ω be an open set in \mathbb{G} . We say that a measurable set $E \subseteq \mathbb{G}$ has *locally finite perimeter* in Ω , if, for every $Y \in \mathfrak{g}_1$, there exists a Radon measure, denoted by $Y\mathbb{1}_E$, on Ω such that

$$\int_{A \cap E} Y\varphi d\mu = - \int_A \varphi d(Y\mathbb{1}_E), \quad \text{for every open set } A \Subset \Omega \text{ and every } \varphi \in C_c^1(A).$$

We say that E has *finite perimeter* in Ω if E has locally finite perimeter in Ω and, for every basis (X_1, \dots, X_m) of \mathfrak{g}_1 , the total variation $|D\mathbb{1}_E|(\Omega)$ of the measure $D\mathbb{1}_E := (X_1\mathbb{1}_E, \dots, X_m\mathbb{1}_E)$ is finite.

Definition 2.2. Let $E \subseteq \mathbb{G}$ be a set with locally finite perimeter in \mathbb{G} . We define the *reduced boundary* $\mathcal{F}E$ of E to be the set of points $p \in \mathbb{G}$ such that $|D\mathbb{1}_E|(B(p, r)) > 0$ for all $r > 0$ and there exists

$$\lim_{r \rightarrow 0} \frac{D\mathbb{1}_E(B(p, r))}{|D\mathbb{1}_E|(B(p, r))} =: \nu_E(p)$$

with $|\nu_E(p)| = 1$.

We denote by $\mathbb{S}(\mathbb{G})$ the unit sphere of \mathbb{G} and by $\mathbb{S}(\mathfrak{g}_1)$ the unit sphere in \mathfrak{g}_1 .

Definition 2.3. Let $E \subseteq \mathbb{G}$ be a set of locally finite perimeter in \mathbb{G} and let $\nu \in \mathbb{S}(\mathfrak{g}_1)$. We say that E has ν as *constant horizontal normal* if $\nu\mathbb{1}_E \geq 0$ in the sense of distributions and, for every $X \in \nu^\perp := \{Y \in \mathfrak{g}_1 : \langle \nu, Y \rangle = 0\}$, one has $X\mathbb{1}_E = 0$.

3. CONES IN CARNOT GROUPS

In this section, we show that semigroups of horizontal normal ν represent the “minimal” sets having ν as constant horizontal normal (see Proposition 3.1 and Lemma 3.5). Moreover, by Lemma 3.2, they are attainable sets of halfspaces in \mathfrak{g}_1 . By a standard argument of Geometric Control Theory (see e.g. [AS13, Theorem 8.1]), this implies that every S_ν has non-empty interior.

An important result of this section is Proposition 3.7, which proves that the semigroups S_ν are continuous with respect to the horizontal direction ν . This fact, resumed in Remark 3.8, will be used in the proof of Theorem 4.5.

The following proposition has been proven in [BLD19, Proposition 2.29]. We, however, write its short proof for the sake of completeness.

Proposition 3.1. *Let \mathbb{G} be a Carnot group, let $\nu \in \mathbb{S}(g_1)$ and let S_ν be the semigroup with normal ν . Then S_ν has ν as constant horizontal normal.*

Proof. By definition, $\mathbb{1}_{S_\nu \cdot \exp(sX)} \leq \mathbb{1}_{S_\nu}$ for all $s \geq 0$ and $X \in \{\nu\} \cup \nu^\perp$. Let $\varphi \in C_c^\infty(\mathbb{G}; [0, +\infty])$ and denote by $\Phi_X(p, s) := p \exp(sX)$ the flow of X at time s starting from point $p \in \mathbb{G}$. Then

$$\begin{aligned} \int_{\mathbb{G}} \mathbb{1}_{S_\nu} X \varphi \, d\mu &= \int_{\mathbb{G}} \mathbb{1}_{S_\nu} \cdot \lim_{s \rightarrow 0} \frac{1}{s} (\varphi(\Phi_X(\cdot, s)) - \varphi) \, d\mu \\ &= \lim_{s \rightarrow 0} \frac{1}{s} \left(\int_{\mathbb{G}} \mathbb{1}_{S_\nu \cdot \exp(sX)} \varphi \, d\mu - \int_{\mathbb{G}} \mathbb{1}_{S_\nu} \varphi \, d\mu \right) \leq 0 \end{aligned}$$

and therefore

$$- \int_G \mathbb{1}_{S_\nu} X \varphi \, d\mu \geq 0.$$

Hence $\langle X \mathbb{1}_{S_\nu}, \cdot \rangle$ is a positive linear functional and by Riesz's Theorem $X \mathbb{1}_{S_\nu}$ is a Radon measure. Moreover, since for each $X \in \nu^\perp$ also $-X \in \nu^\perp$, we get that $X \mathbb{1}_{S_\nu} = 0$ for all $X \in \nu^\perp$. Consequently, S_ν has ν as constant horizontal normal. \square

Lemma 3.2. *Let \mathbb{G} be a Carnot group, $T > 0$, and let $\gamma : [0, T] \rightarrow \mathbb{G}$ be a horizontal curve such that $\gamma(0) = 0$. If $\langle \dot{\gamma}(t), \nu \rangle \geq 0$ for almost every t , then $\gamma(T) \in \overline{S_\nu}$.*

Proof. Fix $T > 0$, define $X_1 := \nu$ and let X_2, \dots, X_m be such that (X_1, X_2, \dots, X_m) is an orthonormal basis for \mathfrak{g}_1 . We fix $T > 0$. We can assume without loss of generality that $\dot{\gamma} \in L^\infty([0, T]; \mathfrak{g}_1)$. Let $u_1, \dots, u_m \in L^\infty([0, T])$ be such that

$$\dot{\gamma}(t) = \sum_{i=1}^m u_i(t) X_i(\gamma(t)),$$

for \mathcal{L}^1 -almost every $t \in [0, T]$. Then, by assumption, we know that $u_1(t) \geq 0$ for \mathcal{L}^1 -almost every $t \in [0, T]$. Consider piecewise constant sequences $(u_1^h), \dots, (u_m^h)$ in $L^\infty([0, T])$, with $h \in \mathbb{N}$, such that

$$u_i^h \rightarrow u_i \quad \text{in } L^1([0, T]), \quad \text{as } h \rightarrow \infty,$$

for any $i = 1, \dots, m$ and such that $u_1^h(t) \geq 0$ for \mathcal{L}^1 -almost every $t \in [0, T]$. We can also assume that $\sup_{h \in \mathbb{N}} \sum_{i=1}^m \|u_i^h\|_\infty \leq M$, for some $M > 0$. According to the definition of S_ν and since u_i^h are piecewise constant, the curves defined by

$$\begin{cases} \dot{\gamma}^h(t) = \sum_{i=1}^m u_i^h(t) X_i(\gamma^h(t)) & \text{for } \mathcal{L}^1\text{-a.e. } t \in [0, T], \\ \gamma^h(0) = 0, \end{cases}$$

are such that $\gamma^h(t) \in S_\nu$ for any $t \in [0, T]$. Since $d(\gamma^h(t), 0) \leq Mt$ for every $t \in [0, T]$, there exists a compact set $K \subseteq \mathbb{G}$ for which

$$\bigcup_{h \in \mathbb{N}} \gamma^h([0, T]) \cup \gamma([0, T]) \subseteq K.$$

We prove that

$$\lim_{h \rightarrow \infty} d(\gamma^h(T), \gamma(T)) = 0.$$

It is not restrictive to work in coordinates and compute

$$\begin{aligned} |\gamma^h(T) - \gamma(T)| &= \left| \int_0^T \sum_{i=1}^m (u_i^h(t) X_i(\gamma^h(t)) - u_i(t) X_i(\gamma(t))) dt \right| \\ &\leq \int_0^T \sum_{i=1}^m |u_i^h(t)| |X_i(\gamma^h(t)) - X_i(\gamma(t))| dt \\ &\quad + \int_0^T \sum_{i=1}^m |u_i^h(t) - u_i(t)| |X_i(\gamma(t))| dt. \end{aligned}$$

Notice that, by the choice of u_i^h , the term

$$\alpha_h(T) := \int_0^T \sum_{i=1}^m |u_i^h(t) - u_i(t)| |X_i(\gamma(t))| dt$$

is infinitesimal as $h \rightarrow \infty$, and that, by the smoothness of X_1, \dots, X_n we can find $C > 0$ depending on T and K such that

$$|\gamma^h(T) - \gamma(T)| \leq \alpha_h(T) + CM \int_0^T |\gamma^h(t) - \gamma(t)| dt.$$

We are then in a position to apply Grönwall Lemma to get

$$|\gamma^h(T) - \gamma(T)| \leq \alpha_h(T) e^{CMT},$$

and letting $h \rightarrow \infty$, we conclude the proof. \square

Before proving Lemma 3.5, we make the following remark.

Remark 3.3. Let \mathbb{G} be a Carnot group and let d_L and d_R be the left- and right-invariant Carnot-Carathéodory distances on \mathbb{G} built with respect to the same scalar product on \mathfrak{g}_1 . Then the inversion is an isometry between (\mathbb{G}, d_L) and (\mathbb{G}, d_R) . In particular, for every $p \in \mathbb{G}$ we have that $d_L(0, p) = d_L(p^{-1}, 0) = d_R(p, 0) = d_R(0, p)$.

Given $p \in \mathbb{G}$ and a measurable set $F \subseteq \mathbb{G}$, we define the *lower and upper densities* $\theta_*(p, F)$ and $\theta^*(p, F)$ of F at p letting

$$\theta_*(p, F) := \liminf_{r \rightarrow 0} \frac{\mu(F \cap B(p, r))}{\mu(B(p, r))} \quad \theta^*(p, F) := \limsup_{r \rightarrow 0} \frac{\mu(F \cap B(p, r))}{\mu(B(p, r))}.$$

The *measure theoretic boundary* $\partial^* F$ of F is defined by

$$\partial^* F := \{p \in \mathbb{G} : \theta_*(p, F) > 0 \text{ and } \theta^*(p, F) < 1\}.$$

The Lebesgue representative \tilde{F} of F is defined by

$$(3) \quad \tilde{F} := \{p \in \mathbb{G} : \theta_*(p, F) = 1\}.$$

Notice that, by the Lebesgue's theorem, we know that $\tilde{F} = F$ up to a set of μ -measure zero. The following proposition is proved in [BLD19, Proposition 3.6]

Proposition 3.4. *Let $F \subseteq \mathbb{G}$ be a set with ν as constant horizontal normal. Then*

$$\mathcal{F}\tilde{F} = \partial\tilde{F} = \partial^*\tilde{F}.$$

Lemma 3.5 below will be used in the proof of Lemma 4.2.

Lemma 3.5. *Let \mathbb{G} be a Carnot group, let $F \subseteq \mathbb{G}$ be a set with ν as constant horizontal normal and assume that $\theta^*(0, F) > 0$. Then $\mathbb{1}_{S_\nu} \leq \mathbb{1}_F$ μ -almost everywhere.*

Proof. Recall first that every left-invariant Haar measure of a Carnot group is right-invariant, being Carnot groups nilpotent and therefore unimodular.

Since F has ν as constant horizontal normal, by [BLD13, Lemma 3.1] we have that,

$$\mathbb{1}_{F \cdot p} \leq \mathbb{1}_F \quad \text{for every } p \in S_\nu, \quad \mu\text{-almost everywhere.}$$

If $B^L(p, r)$ and $B^R(p, r)$ denote, respectively, the metric balls built with respect to the left-invariant and right-invariant metrics, then for all $p \in S_\nu$

$$\begin{aligned} \mu(F \cap B^L(0, r)) &\stackrel{\text{Rem. 3.3}}{=} \mu(F \cap B^R(0, r)) = \mu(F \cdot p \cap B^R(p, r)) \\ &\leq \mu(F \cap B^R(p, r)). \end{aligned}$$

Since $\theta^*(0, F) > 0$, we deduce that for all $p \in S_\nu$,

$$(4) \quad \theta_F^R(p) := \limsup_{r \rightarrow 0} \frac{\mu(F \cap B^R(p, r))}{\mu(B^R(p, r))} > 0.$$

On the other hand, by Lebesgue Theorem we have that $\theta_F^R(p) \in \{0, 1\}$ for μ -almost every $p \in \mathbb{G}$. By (4) we then get that μ -almost every $p \in S_\nu$ is a Lebesgue point for F , hence for almost every $p \in S_\nu$ one also has $p \in F$, as required. \square

In the following lemma, we denote by $\mathbb{R}O(n)$ the set of $n \times n$ matrices A that can be written as

$$A = \lambda B,$$

for some $\lambda \in \mathbb{R}$ and some $B \in O(n)$.

Lemma 3.6. *Let $n \in \mathbb{N}$. Then the following facts hold.*

- (i) *For every $p \in \mathbb{R}^n$ and every neighborhood U of p in \mathbb{R}^n , there exists a neighborhood $M \subseteq \mathbb{R}O(n)$ of the identity matrix such that*

$$\bigcap_{\ell \in M} \ell U$$

is a neighborhood of p .

- (ii) *For every $p \in \mathbb{R}^n \setminus \{0\}$ and every open neighborhood $M \subseteq \mathbb{R}O(n)$ of the identity matrix, the set Mp is an open neighborhood of p .*

Proof. (i) Consider $r, R \in (0, +\infty)$ such that $B(p, 2r) \subseteq U$ and $B(p, r) \subseteq B(0, R)$. We consider in $\mathbb{R}O(n)$ the distance coming from the usual operator norm. We define $M := B(\text{Id}, r/R)^{-1} \subseteq \mathbb{R}O(n)$. Then, take any $\ell \in M$, so that $\|\ell^{-1} - \text{Id}\| < r/R$. Then we have that for all $x \in B(0, R)$

$$|\ell^{-1}(x) - x| \leq \|\ell^{-1} - \text{Id}\| |x| \leq \frac{r}{R} R = r.$$

Consequently, by triangle inequality if $x \in B(p, r)$, and so also $x \in B(0, R)$, we have that

$$|\ell^{-1}(x) - p| \leq |\ell^{-1}(x) - x| + |x - p| \leq 2r.$$

Therefore, we showed that $\ell^{-1}(B(p, r)) \subseteq B(p, 2r) \subseteq U$, that is, $B(p, r) \subseteq \ell U$. Hence, we infer that $B(p, r) \subseteq \bigcap_{\ell \in M} \ell U$.

(ii) Consider the scaled orthogonal transformations $\mathbb{R}O(n)$ acting continuously and transitively on $\mathbb{R}^n \setminus \{0\}$. Fix $p \in \mathbb{R}^n \setminus \{0\}$ and let us denote by $(\mathbb{R}O(n))_p$ the stabilizer subgroup of p . By [Hel01, Theorem 3.2], the mapping

$$\begin{aligned} \psi: \mathbb{R}O(n)/(\mathbb{R}O(n))_p &\rightarrow \mathbb{R}^n \setminus \{0\} \\ [\ell] &\mapsto \ell(p) \end{aligned}$$

is a well-defined homeomorphism, and hence the map $\ell \mapsto \ell(p)$ obtained as $\psi \circ \pi$ is open. \square

Proposition 3.7. *Let \mathbb{G} be a Carnot group. Then, for every $\nu \in \mathbb{S}(\mathfrak{g}_1)$, there exists an open neighborhood U of ν in $\mathbb{S}(\mathfrak{g}_1)$ such that*

$$\bigcap_{\mu \in U} S_\mu$$

has non-empty interior. Moreover, if $\nu \in \mathbb{S}(\mathfrak{g}_1)$ and $X \in \mathfrak{g}_1$ are such that $\exp(X) \in \text{int}(S_\nu)$, then one can choose U in such a way that $X \in \text{int}(\bigcap_{\mu \in U} S_\mu)$.

Proof. Denote by \mathbb{F} the free Carnot group of both the same rank and step of \mathbb{G} (for a definition of free Lie algebra, see e.g. [VSCC92, p. 45] or [Var84, p. 174]). Then, the canonical projection $\pi: \mathbb{F} \rightarrow \mathbb{G}$ induces an isometry between the horizontal layers of the Lie algebras $\pi_*: \mathfrak{f}_1 \rightarrow \mathfrak{g}_1^1$. Define $\tilde{\nu} \in \mathfrak{f}_1$ such that $\pi_*(\tilde{\nu}) = \nu$, and consider a non-empty open set $W \subseteq S_{\tilde{\nu}}$, which exists by [BLD19, Proposition 2.26]. The canonical action of $\mathbb{R}O(m)$ on \mathfrak{f}_1 can be extended linearly to a map in $\text{Aut}(\mathfrak{f})$, since in free groups all non-horizontal left-invariant vector fields can be written in a (essentially) unique way as commutators of horizontal ones. Considering the corresponding Lie group automorphisms, with abuse of notation, we then identify the actions $\mathbb{R}O(m) \times \mathfrak{f}_1 \rightarrow \mathfrak{f}_1$ and $\mathbb{R}O(m) \times \mathbb{F} \rightarrow \mathbb{F}$. Notice that these actions are continuous and open. By Lemma 3.6 there exists an open neighborhood M of the identity in $\mathbb{R}O(m)$ such that $M\tilde{\nu}$ is an open neighborhood of $\tilde{\nu}$ and such that

$$\bigcap_{\ell \in M} \ell W$$

has non-empty interior. Set now $U := \pi_*(M\tilde{\nu})$. Since π_* is open, U is open and for every $\mu \in U$ we can find $m_0 \in M$ such that $\mu = \pi_*(m_0\tilde{\nu})$. By properties of orthogonal matrices, notice that we also have $\pi_*((m_0\tilde{\nu})^\perp) = \mu^\perp$. Since π is open as well, the inclusions

$$\begin{aligned} (5) \quad \pi \left(\text{int} \bigcap_{\ell \in M} \ell W \right) &\subseteq \pi(m_0 W) \subseteq \pi(m_0 S_{\tilde{\nu}}) = \pi(m_0(S(\exp(\tilde{\nu}^\perp + \mathbb{R}^+\tilde{\nu})))) \\ &= \pi S(\exp((m_0\tilde{\nu})^\perp + \mathbb{R}^+m_0\nu)) = S(\exp(\mu^\perp + \mathbb{R}^+\mu)) = S_\mu \end{aligned}$$

conclude the first part of the proof.

Assume now that $\exp(X) \in \text{int}(S_\nu)$. Then one can repeat the argument of the previous part of the proof with the additional conditions that $\exp(X) \in \text{int}(W)$ and, by (i) of Lemma 3.6,

$$\exp(X) \in \text{int} \left(\bigcap_{\ell \in M} \ell W \right).$$

The proof is again finished by (5). \square

¹It is understood that the metric on $\mathfrak{g}_1(\mathbb{F})$ is the pull-back metric of the metric on $\mathfrak{g}_1(\mathbb{G})$

For any subset $A \subseteq \mathbb{S}(\mathbb{G})$, we define

$$(6) \quad C(A) := \{\delta_r(a) : a \in A, r > 0\}.$$

Notice that $C(A)$ is a cone.

Remark 3.8. As an immediate consequence of Proposition 3.7, we notice that there exist $N \in \mathbb{N}$ and a finite family of triples $(U_i, \Omega_i, K_i)_{i=1}^N$ such that

- (1) $(U_i)_{i=1}^N$ is an open covering of $\mathbb{S}(\mathfrak{g}_1)$;
- (2) for every $i = 1, \dots, N$, Ω_i is a nonempty open subset of $\mathbb{S}(\mathbb{G})$ such that

$$C(\Omega_i) \subseteq \bigcap_{\nu \in U_i} S_\nu;$$

- (3) for every $i = 1, \dots, N$, $K_i \subseteq \Omega_i$ is a compact set with nonempty interior in $\mathbb{S}(\mathbb{G})$.

Moreover, if for every $\nu \in \mathbb{S}(\mathfrak{g}_1)$, there exists $X \in \mathfrak{g}_1$ such that $\exp(X) \in \text{int}(S_\nu)$, then each compact set K_i can be chosen in such a way that there exists $X_i \in \mathfrak{g}_1$ with $\exp(X_i) \in \text{int}(C(K_i))$.

4. PROOF OF THEOREM 1.5

In this section we show that, in Carnot groups, the reduced boundary of sets of locally finite perimeter can be decomposed into countably many pieces satisfying a cone property. This is precisely stated in Theorem 4.6, whose proof and statement complete and combine the previous Lemmata 4.2, 4.3, and 4.5.

Remark 4.1. If a subset Γ of a Carnot group satisfies the C -cone property, for some cone C , then Γ also satisfies the C^{-1} -cone property. Indeed, assume by contradiction that there exist $p \in \Gamma$ and $q \in \Gamma \cap pC^{-1}$. Since $q \in pC^{-1}$, then $p \in qC$. But Γ satisfies a C -cone property and, since $q \in \Gamma$ we have $\Gamma \cap qC = \emptyset$. This is in contradiction with $p \in \Gamma \cap qC$.

Lemma 4.2. *Let \mathbb{G} be a Carnot group. If $E \subseteq \mathbb{G}$ is a set of locally finite perimeter in \mathbb{G} and $\varepsilon > 0$, then, for every $p \in \mathcal{F}E$, there exists $r_p > 0$ such that for any $r \in (0, r_p)$ we have*

$$\mu(B_r \cap S_{\nu_E(p)} \cap p^{-1}(\mathbb{G} \setminus E)) \leq \varepsilon r^Q$$

and

$$\mu(B_r \cap S_{\nu_E(p)}^{-1} \cap p^{-1}E) \leq \varepsilon r^Q.$$

Proof. Fix $p \in \mathcal{F}E$ and notice that E admits a tangent at p that has, by the same argument of [FSSC03, Theorem 3.1], $\nu_E(p)$ as constant horizontal normal. This means that, for any sequence $r_{h_k} \rightarrow 0$, we can find a set F with constant horizontal normal $\nu_F \equiv \nu_E(p)$ and a subsequence r_{h_k} such that

$$\mathbb{1}_{\delta_{r_{h_k}}^{-1}(p^{-1}E)} \rightarrow \mathbb{1}_F \text{ in } L_{\text{loc}}^1(\mathbb{G}), \quad \text{as } k \rightarrow +\infty.$$

It is not restrictive to assume that $\tilde{F} = F$, where \tilde{F} is as in (3). By Proposition 3.4, since $0 \in \mathcal{F}\tilde{F}$ then $0 \in \partial^* \tilde{F}$ and by Lemma 3.5, one has

$$\mathbb{1}_{\delta_{r_{h_k}}^{-1}(p^{-1}E) \cap S_{\nu_E(p)}} \rightarrow \mathbb{1}_{S_{\nu_E(p)}} \text{ in } L_{\text{loc}}^1(\mathbb{G}), \quad \text{as } k \rightarrow +\infty.$$

Consequently, we have

$$\mathbb{1}_{\delta_{r_{h_k}}^{-1}(p^{-1}E) \cap S_{\nu_E(p)}} \rightarrow \mathbb{1}_{S_{\nu_E(p)}} \text{ in } L_{\text{loc}}^1(\mathbb{G}), \quad \text{as } r \rightarrow 0.$$

This completes the proof of the first statement. The proof of the second inequality follows from the first one by replacing E with $\mathbb{G} \setminus E$ and recalling that $S_{-\nu} = S_\nu^{-1}$. \square

Regarding next lemma, recall the notation $C(\Omega)$ introduced in (6).

Lemma 4.3. *Let \mathbb{G} be a Carnot group, let $K \subseteq \Omega \subseteq \mathbb{S}(\mathbb{G})$ be such that K is compact and Ω is open. Then*

$$\eta(C(\Omega), C(K)) := \inf \{ \mu(C(\Omega) \cap B_2 \cap \xi C(\Omega)^{-1} \cap \xi B_2) : \xi \in K \} > 0.$$

Proof. Since K is a compact set, there exists $\delta \in (0, 1)$ such that

$$0 < \delta < d(K, \mathbb{G} \setminus C(\Omega)).$$

Hence for any $\xi \in K$ one has $\xi B_\delta \subseteq C(\Omega) \cap B_2$. Therefore

$$\begin{aligned} \mu(C(\Omega) \cap B_2 \cap \xi C(\Omega)^{-1} \cap \xi B_2) &\geq \mu(C(\Omega) \cap B_2 \cap \xi C(\Omega)^{-1} \cap \xi B_\delta) \\ &= \mu(\xi C(\Omega)^{-1} \cap \xi B_\delta) \\ &= \mu(C(\Omega)^{-1} \cap B_\delta), \end{aligned}$$

which is a positive lower bound independent of $\xi \in K$. \square

By combining Remark 3.8 and Lemmata 4.2 and 4.3 we get the following corollary.

Corollary 4.4. *Let \mathbb{G} be a Carnot group, let $(U_i, \Omega_i, K_i)_{i=1}^N$ be as in Remark 3.8, and let $E \subseteq \mathbb{G}$ be a set of locally finite perimeter. Then, for every $p \in \mathcal{F}E$, there exists $r_p > 0$ such that, if $\nu_E(p) \in U_i$ and $r \in (0, r_p)$, then*

$$(7) \quad \mu(B_{2r} \cap C_i \cap p^{-1}(\mathbb{G} \setminus E)) \leq \frac{\eta_i}{3} r^Q$$

and

$$(8) \quad \mu(B_{2r} \cap C_i^{-1} \cap p^{-1}E) \leq \frac{\eta_i}{3} r^Q,$$

where $C_i := C(\Omega_i)$ and $\eta_i := \eta(C_i, C(K_i)) = \inf \{ \mu(C_i \cap B_2 \cap \xi C_i^{-1} \cap \xi B_2) : \xi \in K_i \}$.

Lemma 4.5. *Let \mathbb{G} be a Carnot group and let E be a set of locally finite perimeter in \mathbb{G} . Consider a family $(U_i, \Omega_i, K_i)_{i=1}^N$ as in Remark 3.8 and, for every $p \in \mathcal{F}E$, define $r_p > 0$ as in Corollary 4.4. Then, the sets*

$$(9) \quad F_{i,\ell} := \{ p \in \mathcal{F}E : \nu_E(p) \in U_i, r_p > \frac{1}{\ell} \},$$

defined for $i \in \{1, \dots, N\}$ and $\ell \in \mathbb{N}$, satisfy

$$(10) \quad F_{i,\ell} \cap p B_{1/\ell} \cap p C(K_i) = \emptyset \quad \forall p \in F_{i,\ell}.$$

Proof. Denote for shortness $\tilde{C}_i := C(K_i)$. Suppose by contradiction that there exist $x \in F_{i,\ell}$ and $y \in F_{i,\ell} \cap x B_{1/\ell} \cap x \tilde{C}_i$ with $y \neq x$. Let η_i be as in Corollary 4.4 and define

$$r := d(x, y) \quad \text{and} \quad I := x C_i \cap y C_i^{-1} \cap x B_{2r} \cap y B_{2r}.$$

By construction, we have that

$$r = d(x, y) < \frac{1}{\ell} < \min\{r_x, r_y\}$$

and using the facts that $I \subseteq x C_i \cap x B_{2r}$ and $I \subseteq y C_i^{-1} \cap y B_{2r}$, by applying (7) and (8) we have

$$\mu(I \setminus E) \leq \mu(x C_i \cap x B_{2r} \cap (\mathbb{G} \setminus E)) \leq \frac{\eta_i}{3} r^Q$$

and

$$\mu(I \cap E) \leq \mu(y C_i^{-1} \cap y B_{2r} \cap E) \leq \frac{\eta_i}{3} r^Q.$$

This contradicts the fact that by Lemma 4.3, we have $\mu(I) \geq \eta_i r^Q$. \square

Theorem 4.6. *Let \mathbb{G} be a Carnot group and let $E \subseteq \mathbb{G}$ be a set of locally finite perimeter in \mathbb{G} . Then there exist a countable family $\{C_h : h \in \mathbb{N}\}$ of open cones in \mathbb{G} and a countable family $\{\Gamma_h : h \in \mathbb{N}\}$ of subsets of \mathbb{G} such that each Γ_h satisfies the C_h -cone property and*

$$\mathcal{F}E = \bigcup_{h \in \mathbb{N}} \Gamma_h.$$

Proof. It is enough to show that, for any $i, \ell \in \mathbb{N}$, the set $F_{i,\ell}$ defined in (9) can be covered by a countable union of sets satisfying a cone property. This is simply done by covering $F_{i,\ell}$ by a countable family of balls $\{B(p_j, 1/\ell) : p_j \in F_{i,\ell}, j \in \mathbb{N}\}$ and decomposing $F_{i,\ell} = \bigcup_{j \in \mathbb{N}} (F_{i,\ell} \cap B(p_j, 1/\ell))$. We now set

$$(11) \quad \Gamma_{i,\ell}^j := F_{i,\ell} \cap B(p_j, 1/\ell)$$

Then by (10), we have that, for every $i, j \in \mathbb{N}$, the set $\Gamma_{i,\ell}^j$ has the $C(\text{int}(K_i))$ -cone property, where K_i are the compact subsets of the sphere introduced in Remark 3.8. Up to relabeling the family $\{\Gamma_{i,\ell}^j : i = 1, \dots, N, \ell, j \in \mathbb{N}\}$ and renaming $C_i := C(\text{int}(K_i))$, we have then a countable family $\{C_h : h \in \mathbb{N}\}$ of open cones and a countable family $\{\Gamma_h : h \in \mathbb{N}\}$ of sets such that each Γ_h satisfies the C_h -cone property and $\mathcal{F}E = \bigcup_{h \in \mathbb{N}} \Gamma_h$. \square

Remark 4.7. Notice that the family of cones $\{C_h : h \in \mathbb{N}\}$ appearing in Theorem 4.6 is indeed finite. This comes from the construction $C_i = C(\text{int}(K_i))$, and the fact that, by Remark 3.8, the family $\{K_i : i = 1, \dots, N\}$ is finite.

5. INTRINSIC LIPSCHITZ GRAPHS

In this section we follow [FSSC06] to recall the notion of intrinsic Lipschitz graph in Carnot groups. The construction of an intrinsic Lipschitz graph requires the space to have a decomposition into complementary subgroups (see Definition 1.1). As observed in Remark 5.1 below, in certain cases Theorem 4.6 can be strengthened to deduce that the reduced boundary of sets of finite perimeter is intrinsically Lipschitz-rectifiable, i.e., it can be covered by a countable union of intrinsic Lipschitz graphs.

Since we only deal with the notion of codimension-one rectifiable sets, Definition 1.1 will be used only in case $\dim \mathbb{L} = 1$. In this situation, according to [Vit12, Proposition 3.4] (see also [FS16]), one can see that, up to a modification of the parameter $\beta > 0$, sets Σ satisfying point (i) of Definition 1.1 can be extended to sets $\Sigma \subseteq \tilde{\Sigma}$ that satisfy (ii).

Remark 5.1. Assume $\Gamma \subseteq \mathbb{G}$ is a set with the C -cone property such that C is open and there exists $X \in \mathfrak{g}_1 \setminus \{0\}$ with the property that $\exp(X) \in C$. Then Γ is an intrinsic Lipschitz graph. Indeed, according to Definition 1.1, we may choose $\mathbb{L} := \{\exp(tX) : t \in \mathbb{R}\}$ and $\mathbb{W} := \exp(X^\perp \oplus \mathfrak{g}_2 \oplus \dots \oplus \mathfrak{g}_s)$. Then, since C is open, there exists $\varepsilon > 0$ such that $B(\exp(X), \varepsilon) \subseteq C$ and with the choice $\beta = \varepsilon$, condition (1) is satisfied.

In particular, Lemma 4.5 shows that, if for some $i = 1, \dots, N$, the set K_i defined in Remark 3.8 is such that $\text{int}(K_i) \cap \exp(\mathfrak{g}_1) \neq \emptyset$, then, for every $j, \ell \in \mathbb{N}$, the set $\Gamma_{i,\ell}^j$ defined in (11) is an intrinsic Lipschitz graph.

Definition 5.2. Let \mathbb{G} be a Carnot group of homogeneous dimension Q and let $E \subseteq \mathbb{G}$. We say that E is *intrinsically Lipschitz rectifiable* if there exists a countable family $\{\Sigma_h : h \in \mathbb{N}\}$ of intrinsic

Lipschitz graphs such that

$$\mathcal{H}^{Q-1} \left(E \setminus \bigcup_{h \in \mathbb{N}} \Sigma_h \right) = 0.$$

An immediate consequence of Remark 5.1 and Theorem 4.6 is given by Corollary 5.3 below.

Corollary 5.3. *Let \mathbb{G} be a Carnot group and assume that for all $\nu \in \mathbb{S}(\mathfrak{g}_1)$ there exists $X \in \mathbb{S}(\mathfrak{g}_1)$ such that $\exp(X) \in \text{int}(S_\nu)$. Then the reduced boundary of every set of locally finite perimeter in \mathbb{G} is intrinsically Lipschitz rectifiable.*

To describe some conditions on the group that guarantee the validity of the assumptions of Corollary 5.3, we introduce the definition of end-point map.

Definition 5.4. Let \mathbb{G} be a Carnot group. The *end-point map* $\text{End}: L^\infty([0, 1]; \mathfrak{g}_1) \rightarrow \mathbb{G}$ is defined by letting

$$\text{End}(h) = \gamma(1),$$

where $\gamma: [0, 1] \rightarrow \mathbb{G}$ is the horizontal curve that is the unique solution of

$$\begin{cases} \dot{\gamma}(t) = h(t) \\ \gamma(0) = 0. \end{cases}$$

With abuse of notation we also write $\text{End}(\gamma)$ meaning $\text{End}(h)$ for the defining control h .

In what follows, we say that a map $F: M \rightarrow N$ between topological spaces M and N is *locally open at $p \in M$* if, for every neighborhood U of p , the set $F(U)$ is a neighborhood of $F(p)$. If \mathbb{G} is a Carnot group and $X \in \mathfrak{g}_1$ is a horizontal direction, we also say that the end-point map $\text{End}: L^\infty([0, 1]; \mathfrak{g}_1) \rightarrow \mathbb{G}$ is locally open at X , if it is locally open at $h(t) \equiv X$.

Before proving Lemma 5.6, we point out some topological properties of the semigroups S_ν .

Proposition 5.5. *Let \mathbb{G} be a Carnot group and let $\nu \in \mathfrak{g}_1 \setminus \{0\}$. Then $\text{int}(S_\nu) = \text{int}(\overline{S_\nu})$.*

Proof. Since $\text{int}(S_\nu)$ has the (inner) $\text{int}(S_\nu)$ -cone property, then by [BLD19, Lemma 2.35] we have that

$$(12) \quad \text{int}(S_\nu) = \text{int}(\overline{\text{int}(S_\nu)}).$$

On the other hand, by [BLD19, Lemma 2.26], we also have that

$$(13) \quad \overline{S_\nu} = \overline{\text{int}(S_\nu)}.$$

The result follows combining (12) and (13). \square

Lemma 5.6. *Let \mathbb{G} be a Carnot group, let $X \in \mathfrak{g}_1 \setminus \{0\}$ and assume that the end-point map $\text{End}: L^\infty([0, 1]; \mathfrak{g}_1) \rightarrow \mathbb{G}$ is locally open at X . Then, for every $\nu \in \mathbb{S}(\mathfrak{g}_1)$ satisfying $\langle \nu, X \rangle > 0$, we have $\exp(X) \in \text{int}(S_\nu)$.*

Proof. Let $\varepsilon := \langle \nu, X \rangle > 0$ and

$$B_\varepsilon(X) := \{v \in L^\infty([0, 1]; \mathfrak{g}_1) : \|X - v\|_\infty < \varepsilon\}.$$

Since End is open at X , by Proposition 5.5 it suffices to show that $\text{End}(B_\varepsilon(X)) \subseteq \overline{S_\nu}$. On the other hand, by Lemma 3.2, if $v \in L^\infty([0, 1]; \mathfrak{g}_1)$ satisfies $\langle v(t), \nu \rangle > 0$ for almost every $t \in [0, 1]$, then $\text{End}(v) \in \overline{S_\nu}$. The proof is then achieved by noticing that, for every $v \in B_\varepsilon(X)$, we have

$$\langle v(t), \nu \rangle = \langle \nu, X \rangle - \langle \nu, X - v(t) \rangle \geq \varepsilon - \|\nu\| \|X - v\|_\infty > 0$$

for almost every $t \in [0, 1]$. \square

Corollary 5.7. *Let \mathbb{G} be a Carnot group and assume there exists a basis $\{X_i : i = 1, \dots, m\}$ of \mathfrak{g}_1 such that the end-point map $\text{End} : L^\infty([0, 1]; \mathfrak{g}_1) \rightarrow \mathbb{G}$ is locally open at X_i , for every $i = 1, \dots, m$. Then the reduced boundary of every set of locally finite perimeter in \mathbb{G} is intrinsically Lipschitz rectifiable.*

Proof. It is enough to combine Corollary 5.3, Lemma 5.6 and the following fact. If $X \in \mathfrak{g}_1$ and $\exp(X) \in \text{int}(S_\nu)$, then $\exp(-X) \in \text{int}(S_{-\nu})$. \square

Remark 5.8. Every Carnot group possessing a spanning set of pliable or strongly pliable vectors in the sense of [JS17] and [SS18], respectively, has the property that the reduced boundary of any set of locally finite perimeter is intrinsically Lipschitz rectifiable.

We next give a sufficient condition that has an equivalent algebraic formulation, and that can be more easily verified. This will be used to deduce that, for example, Corollary 5.7 applies to *filiform groups* (see Section 6).

Definition 5.9. Let \mathbb{G} be a Carnot group and let $\gamma : [0, 1] \rightarrow \mathbb{G}$ be a horizontal curve. We say that γ is *non-abnormal* if $d\text{End}(\gamma)$ has full rank. We also say that a horizontal direction $X \in \mathfrak{g}_1 \setminus \{0\}$ is non-abnormal, if $t \mapsto \exp(tX)$ is non-abnormal.

As pointed out in [ABB19] and in [Mon02], the fact that a curve is abnormal does not depend on its parametrization and one can develop a theory considering the end-point map defined on any L^p space, $1 \leq p \leq \infty$. Moreover, the Volterra expansion (see e.g. [ABB19, Formula (6.9)]), allows to compute the differential of the End-point map with respect to any variation in L^p . In particular, if the differential of $\text{End} : L^2([0, T]; \mathbb{R}^m) \rightarrow \mathbb{G}$ has full-rank at $X \in \mathfrak{g}_1$, then also the differential of $\text{End} : L^\infty([0, T]; \mathbb{R}^m) \rightarrow \mathbb{G}$ has full rank at X . This observation allows us to consider the formula for the differential of the end-point map in L^2 developed by [LDMO⁺16].

Proposition 5.10. *Let \mathbb{G} be a Carnot group and let $X \in \mathfrak{g}_1 \setminus \{0\}$. Then the curve $t \mapsto \exp(tX)$ is non-abnormal if and only if*

$$(14) \quad \text{span}\{\text{ad}_X^k(\mathfrak{g}_1) : k = 0, \dots, s-1\} = \mathfrak{g}.$$

Proof. Denote by $\gamma(t) := \exp(tX)$. It is enough to notice that, by [LDMO⁺16, Proposition 2.3]

$$\text{Im}(d\text{End}(\gamma)) = dR_{\gamma(1)}\text{span}\{\text{Ad}_{\gamma(t)}\mathfrak{g}_1 : t \in (0, 1)\} = dR_{\gamma(1)}\text{span}\{\text{ad}_X^k\mathfrak{g}_1 : k \in \mathbb{N}\}. \quad \square$$

Remark 5.11. Condition (14) is clearly open in X . In particular, if \mathbb{G} contains a non-abnormal curve, then there exists an open set $U \subseteq \mathbb{S}(\mathfrak{g}_1)$ such that $U = -U$ and $t \mapsto \exp(tX')$ is non-abnormal for all $X' \in U$.

Corollary 5.12. *Let \mathbb{G} be a Carnot group and assume there exists a non-abnormal $X \in \mathfrak{g}_1 \setminus \{0\}$. Then the reduced boundary of any set of locally finite perimeter in \mathbb{G} is intrinsically Lipschitz rectifiable.*

Proof. Since $\gamma(t) := \exp(tX)$ is non-abnormal, then $d\text{End}(\gamma)$ has full rank and, in particular, End is locally open at X . The proof then follows by combining Corollary 5.7 and Remark 5.11. \square

Proposition 5.13. *Let \mathbb{G}_1 and \mathbb{G}_2 be two Carnot groups possessing non-zero non-abnormal horizontal directions. Then the Carnot group $\mathbb{G}_1 \times \mathbb{G}_2$ possesses a non-zero non-abnormal horizontal direction.*

Proof. Denote by $\mathfrak{g}(\mathbb{G}_i)$ the Lie algebra of \mathbb{G}_i and by $\mathfrak{g}_1(\mathbb{G}_i)$ the related horizontal layer, for $i = 1, 2$. Recall that the Lie bracket of the product algebra $\mathfrak{g}(\mathbb{G}_1) \times \mathfrak{g}(\mathbb{G}_2)$ is defined by

$$[(Y_1, Y_2), (Z_1, Z_2)] = ([Y_1, Z_1], [Y_2, Z_2]),$$

for every $Y_1, Z_1 \in \mathfrak{g}(\mathbb{G}_1)$ and every $Y_2, Z_2 \in \mathfrak{g}(\mathbb{G}_2)$. Then, by induction on k , one can check that

$$\text{ad}_{(Y_1, Y_2)}^k(Z_1, Z_2) = (\text{ad}_{Y_1}^k(Z_1), \text{ad}_{Y_2}^k(Z_2)),$$

for every $k \in \mathbb{N}$, every $Y_1, Z_1 \in \mathfrak{g}(\mathbb{G}_1)$ and every $Y_2, Z_2 \in \mathfrak{g}(\mathbb{G}_2)$.

Let $X_1 \in \mathfrak{g}_1(\mathbb{G}_1)$ and $X_2 \in \mathfrak{g}_1(\mathbb{G}_2)$ be non-zero non-abnormal directions for \mathbb{G}_1 and \mathbb{G}_2 , respectively. Then (X_1, X_2) is non-abnormal for $\mathbb{G}_1 \times \mathbb{G}_2$. To prove this it is enough to notice that for any $k \in \mathbb{N}$ one has

$$\text{ad}_{(X_1, X_2)}^k(\mathfrak{g}_1(\mathbb{G}_1) \times \mathfrak{g}_1(\mathbb{G}_2)) = \text{ad}_{X_1}^k(\mathfrak{g}_1(\mathbb{G}_1)) \times \text{ad}_{X_2}^k(\mathfrak{g}_1(\mathbb{G}_2)) \quad \square$$

Proposition 5.14. *Let \mathbb{G} be a Carnot group possessing a non-zero non-abnormal horizontal direction and assume that $N \trianglelefteq \mathbb{G}$ is a normal subgroup of \mathbb{G} . Then, the Carnot group \mathbb{G}/N possesses a non-zero non-abnormal horizontal direction.*

Proof. Let $X \in \mathfrak{g}_1 \setminus \{0\}$ be a non-abnormal direction for \mathbb{G} . Then, if $\pi : \mathbb{G} \rightarrow \mathbb{G}/N$ is the canonical projection, the push-forward vector π_*X is non-abnormal for \mathbb{G}/N . This is true by the fact that

$$\pi\mathfrak{g} = \text{span}\{\pi_*\text{ad}_X^k\mathfrak{g}_1 : k \in \mathbb{N}\} = \text{span}\{\text{ad}_{\pi_*X}^k\pi_*\mathfrak{g}_1 : k \in \mathbb{N}\},$$

and by noticing that $\pi_*\mathfrak{g}_1 = \mathfrak{g}_1(\mathbb{G}/N)$. If $\pi_*X \neq 0$, the proof is concluded. If $\pi_*X = 0$, then $0 \in \mathfrak{g}_1(\mathbb{G}/N)$ is non-abnormal and, since non-abnormality is an open and dilation-invariant condition, all the elements in $\mathfrak{g}_1(\mathbb{G}/N)$ are non-abnormal. \square

6. FILIFORM GROUPS

In this section we briefly introduce filiform groups and study their abnormal horizontal lines and automorphisms. As a corollary, we obtain that, in all filiform groups, the reduced boundary of any set of locally finite perimeter is intrinsically Lipschitz rectifiable.

Definition 6.1. We say that a Carnot group \mathbb{G} is a *filiform group* of step s if the stratification $\mathfrak{g} = \mathfrak{g}_1 \oplus \dots \oplus \mathfrak{g}_s$ of its Lie algebra satisfies $\dim \mathfrak{g}_1 = 2$ and $\dim \mathfrak{g}_i = 1$, for every $i = 2, \dots, s$.

Definition 6.2. Let \mathbb{G} be a filiform group of step s . If its Lie algebra \mathfrak{g} has a basis $\{X_0, \dots, X_s\}$ with the only nonzero bracket relations

- (i) $[X_0, X_i] = X_{i+1}$ for $i = 1, \dots, s-1$, then \mathbb{G} is said to be *of the first kind*.
- (ii) $[X_0, X_i] = X_{i+1}$ for $i = 1, \dots, s-2$ and $[X_i, X_{s-i}] = (-1)^i X_s$ for $i = 1, \dots, s-1$, then \mathbb{G} is said to be *of the second kind*.

In what follows, any basis of the Lie algebra $\{X_0, \dots, X_s\}$ satisfying relations (i) or (ii) will be called *filiform basis*. The classification of filiform groups below follows from [Ver70, Proposition 5].

Proposition 6.3 (Vergne). *Let \mathbb{G} be a filiform group of step s . If $s = 2n + 1$ for $n \geq 2$, then \mathbb{G} is either of the first kind or of the second kind. Otherwise \mathbb{G} is of the first kind.*

We point out that our choice of basis for the filiform groups of the second kind differs from the one of Vergne. Indeed, in [Ver70] the author considers a basis $\{Y_0, \dots, Y_s\}$ for which $[Y_0, Y_i] = Y_{i+1}$ and $[Y_i, Y_{s-i}] = (-1)^i Y_s$ for $i = 1, \dots, s-1$. One sees that the basis of Vergne has an extra nonzero bracket $[Y_0, Y_{s-1}] = Y_s$ and the basis in Definition 6.2 (ii) is obtained by choosing $X_0 = Y_0 + Y_1$

and $Y_i = X_i$ for $i = 1, \dots, s$. In addition to having fewer nontrivial bracket relations, our choice of the basis has the benefit that it is adapted to the abnormal lines, as we will see next.

Proposition 6.4. *Let \mathbb{G} be a filiform group and let $\{X_0, \dots, X_s\}$ be a filiform basis for \mathfrak{g} . Then the line $t \mapsto \exp(tX_1)$ is abnormal. If \mathbb{G} is of the first kind, then this is the only abnormal horizontal line of \mathbb{G} . Otherwise there exists exactly one other abnormal horizontal line, namely $t \mapsto \exp(tX_0)$.*

Proof. We begin by proving the following claim: if \mathbb{G} is a filiform group of step $s \geq 3$ and $X \in \mathfrak{g}_1$, then the line $t \mapsto \exp(tX)$ is abnormal if and only if $[X, X_i] = 0$ for some $i = 2, \dots, s-1$. Assume first that $[X, X_i] = 0$ for some $i = 2, \dots, s-1$. Then, since \mathfrak{g}_i is one-dimensional, $[X, \mathfrak{g}_i] = 0$ and in particular

$$\mathfrak{g}_i \cap \text{span}\{\text{ad}_X^k(\mathfrak{g}_1) : k = 0, \dots, s-1\} = \text{ad}_X^i(\mathfrak{g}_1) = [X, \text{ad}_X^{i-1}(\mathfrak{g}_1)] = 0.$$

Hence, by Proposition 5.10, the line $t \mapsto \exp(tX)$ is abnormal. To prove the other implication, suppose that, for each $i = 2, \dots, s-1$, one has $[X, X_i] \neq 0$. Since the first layer \mathfrak{g}_1 does not contain elements of the center, we also find $Y_1 \in \mathfrak{g}_1$ for which $[X, Y_1] \neq 0$. Observe now that, since $\mathfrak{g}_i = \text{span}([X, X_{i-1}])$, one has $\mathfrak{g}_i = \text{span}(\text{ad}_X^{i-1}(Y_1))$ for all $i = 2, \dots, s$. By Proposition 5.10, the claim is proved.

As a consequence of the claim, since any filiform basis satisfies $[X_1, X_2] = 0$, the line $t \mapsto \exp(tX_1)$ is abnormal. If \mathbb{G} is of the second kind, then the relation $[X_0, X_{s-1}] = 0$ shows that the line $t \mapsto \exp(tX_0)$ is abnormal as well. We now show that, if \mathbb{G} is of the first kind, the line $t \mapsto \exp(tX_1)$ is the only abnormal horizontal line of \mathbb{G} . Indeed, for every $a \in \mathbb{R}$, the element $X := X_0 + aX_1 \in \mathfrak{g}_1$ satisfies $[X, X_i] = X_{i+1}$ for all $i = 1, \dots, s-1$. By the previous observations, the line $t \mapsto \exp(tX)$ is non-abnormal.

We are left to prove that a filiform group cannot possess more than two abnormal horizontal lines. This fact would imply that $t \mapsto \exp(tX_0)$ and $t \mapsto \exp(tX_1)$ are the only horizontal abnormal lines of the filiform groups of the second kind. First, notice that a line $t \mapsto \exp(tX)$ is abnormal if and only if $X \in \bigcup_{i=2}^{s-1} \ker \phi_i$, where $\phi_i: \mathfrak{g}_1 \rightarrow \mathfrak{g}_{i+1}$ is defined by $\phi_i(X) = [X, X_i]$. According to Proposition 6.3, the algebra $\mathfrak{g}/\mathfrak{g}_s$ is filiform of step $s-1$ and therefore is of the first kind. By the previous part of this proof, the set $\bigcup_{i=2}^{s-2} \ker \phi_i \subseteq \mathfrak{g}/\mathfrak{g}_s$ is one-dimensional. Since ϕ_{s-1} is surjective, also $\dim \ker \phi_{s-1} \leq 1$ and we conclude that $\bigcup_{i=2}^{s-1} \ker \phi_i$ cannot contain more than two linearly independent lines. \square

Corollary 6.5. *Every filiform group has a horizontal non-abnormal line. In particular, Corollary 5.12 applies to all filiform groups.*

Remark 6.6. Applying [LDMO⁺16, Proposition 2.21], one can check that in filiform groups of the first kind the only horizontal injective abnormal curve from the origin is, up to reparametrization, the line $t \mapsto \exp(tX_1)$. On the other hand, in filiform groups of the second kind there are also horizontal abnormal curves that are not lines. For example, the curve defined by

$$t \mapsto \begin{cases} \exp(tX_1) & \text{for } t \in [0, 1], \\ \exp(X_1) \exp((t-1)X_0) & \text{for } t \in [1, 2] \end{cases}$$

is abnormal.

For the sake of completeness, we end by describing the graded Lie algebra automorphisms (i.e., the stratification preserving Lie automorphisms) of filiform Lie algebras. By the following two propositions, we observe that any linear bijection on the horizontal layer that fixes the abnormal lines extends uniquely to a Lie algebra automorphism.

Proposition 6.7. *Let \mathfrak{g} be a filiform Lie algebra of the first kind with a filiform basis $\{X_0, \dots, X_s\}$. The linear transformation on \mathfrak{g}_1 that in basis $\{X_0, X_1\}$ is given by the matrix*

$$\begin{pmatrix} a & 0 \\ c & b \end{pmatrix}$$

induces a (graded) Lie algebra automorphism for every $a, b \in \mathbb{R} \setminus \{0\}$ and $c \in \mathbb{R}$. Moreover, every graded Lie algebra automorphism of \mathfrak{g} is of this form.

Proof. Notice that the only horizontal vectors that commute with \mathfrak{g}_2 are those parallel to X_1 . Therefore, any $\psi \in \text{Aut}(\mathfrak{g})$ maps $\psi(X_1) = bX_1$ with $b \in \mathbb{R} \setminus \{0\}$. Then mapping $\psi(X_0) = aX_0 + cX_1$ defines a Lie algebra automorphism for any choice of $a, b \in \mathbb{R} \setminus \{0\}$ and $c \in \mathbb{R}$ by $\psi(X_i) := \text{ad}_{\psi(X_0)}^{i-1}(\psi(X_1)) = a^i b X_i$. \square

Proposition 6.8. *Let \mathfrak{g} be a filiform Lie algebra of the second kind with a filiform basis $\{X_0, \dots, X_s\}$. The linear transformation on \mathfrak{g}_1 that in basis $\{X_0, X_1\}$ is given by the matrix*

$$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$$

induces a (graded) Lie algebra automorphism for every $a, b \in \mathbb{R} \setminus \{0\}$. Moreover, every graded Lie algebra automorphism of \mathfrak{g} is of this form.

Proof. Similarly to the filiform groups of the first kind, here X_1 is the unique direction commuting with \mathfrak{g}_2 , and hence the line bX_1 , $b \in \mathbb{R}$, must be fixed by every $\psi \in \text{Aut}(\mathfrak{g})$. Since in addition a graded Lie algebra automorphism maps abnormal horizontal lines into abnormal horizontal lines, necessarily $\psi(X_0) = aX_0$ for some $a \in \mathbb{R} \setminus \{0\}$ according to Proposition 6.4. Let us verify that the linear map defined by $\psi(X_0) = aX_0$ and $\psi(X_1) = bX_1$ on \mathfrak{g}_1 induces a Lie algebra automorphism for all $a, b \in \mathbb{R} \setminus \{0\}$ by explicitly calculating the bracket relations. Indeed, the extension

$$\begin{aligned} \psi(X_i) &= \psi(\text{ad}_{X_0}^{i-1} X_1) := \text{ad}_{\psi(X_0)}^{i-1} \psi(X_1) = a^{i-1} b X_i \quad \forall i = 1, \dots, s-1, \\ \psi(X_s) &= \psi(-[X_1, X_{s-1}]) := -[\psi(X_1), \psi(X_{s-1})] = a^{s-2} b^2 X_s \end{aligned}$$

satisfies

$$[\psi(X_i), \psi(X_{s-i})] = [a^{i-1} b X_i, a^{s-i-1} b X_{s-i}] = a^{s-2} b^2 (-1)^i X_s = \psi([X_i, X_{s-i}]) \quad \forall i = 2, \dots, s-2$$

and all the other brackets are zero, as required. \square

Remark 6.9. In the literature, there has not been a clear motivation why the two types of filiform groups are not isomorphic: in [Ver70, page 93] Vergne left it to the reader, while in [Pan89, page 49] Pansu claimed, erroneously, that in filiform groups of the second kind the only graded automorphisms are the homotheties (i.e., composition of isometries and intrinsic dilations). First of all, one can check that, in dimension 4, there is only one filiform group, known as the *Engel group*. Starting from dimension 6, we provided two ways to distinguish the two classes. A first reason is that the spaces of graded automorphisms are different: in filiform groups of the first kind this class is 3 dimensional, while for the second kind this class is 2 dimensional. A second reason, which has a control-theoretic flavor, is that the two classes have different number of abnormal one-parameter subgroups: in filiform groups of the first kind there is only one, while for the second kind there are two. Recall that the stratification of a stratifiable group is unique up to isomorphisms, see [LD17, Proposition 2.17]. Hence, since we showed that the two classes are different as stratified groups, they are different as Lie groups.

Remark 6.10. We stress that no filiform group is Pansu-rigid, as instead stated by Pansu [Pan89, page 49]. Indeed, in both types of filiform groups, there are more graded automorphisms than just the homotheties, in which case we would have said, by definition, that the Carnot group is Pansu-rigid, see [LDOW14]. It is however true that, in filiform groups of the second kind, the only graded automorphisms that are unipotent are the dilations.

REFERENCES

- [ABB19] Andrei Agrachev, Davide Barilari, and Ugo Boscain. A comprehensive introduction to sub-Riemannian geometry, 2019.
- [AFP00] Luigi Ambrosio, Nicola Fusco, and Diego Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [AK00a] Luigi Ambrosio and Bernd Kirchheim. Currents in metric spaces. *Acta Math.*, 185(1):1–80, 2000.
- [AK00b] Luigi Ambrosio and Bernd Kirchheim. Rectifiable sets in metric and Banach spaces. *Math. Ann.*, 318(3):527–555, 2000.
- [AKLD09] Luigi Ambrosio, Bruce Kleiner, and Enrico Le Donne. Rectifiability of sets of finite perimeter in Carnot groups: existence of a tangent hyperplane. *J. Geom. Anal.*, 19(3):509–540, 2009.
- [AS13] Andrei A Agrachev and Yuri Sachkov. *Control theory from the geometric viewpoint*, volume 87. Springer Science & Business Media, 2013.
- [BLD13] Costante Bellettini and Enrico Le Donne. Regularity of sets with constant horizontal normal in the Engel group. *Comm. Anal. Geom.*, 21(3):469–507, 2013.
- [BLD19] Costante Bellettini and Enrico Le Donne. Sets with constant normal in Carnot groups: properties and examples. *ArXiv e-prints*, 2019.
- [DG54] Ennio De Giorgi. Su una teoria generale della misura $(r - 1)$ -dimensionale in uno spazio ad r dimensioni. *Ann. Mat. Pura Appl. (4)*, 36:191–213, 1954.
- [DG55] Ennio De Giorgi. Nuovi teoremi relativi alle misure $(r - 1)$ -dimensionali in uno spazio ad r dimensioni. *Ricerche Mat.*, 4:95–113, 1955.
- [DMV19] Sebastiano Don, Annalisa Massaccesi, and Davide Vittone. Rank-one theorem and subgraphs of BV functions in Carnot groups. *J. Funct. Anal.*, 276(3):687–715, 2019.
- [DV19] Sebastiano Don and Davide Vittone. Fine properties of functions with bounded variation in Carnot-Carathéodory spaces. *J. Math. Anal. Appl.*, 479(1):482–530, 2019.
- [EG15] Lawrence C. Evans and Ronald F. Gariepy. *Measure theory and fine properties of functions*. Textbooks in Mathematics. CRC Press, Boca Raton, FL, revised edition, 2015.
- [Fed69] Herbert Federer. *Geometric measure theory*. Die Grundlehren der mathematischen Wissenschaften, Band 153. Springer-Verlag New York Inc., New York, 1969.
- [FS82] Gerald B. Folland and Elias M. Stein. *Hardy spaces on homogeneous groups*, volume 28. Princeton University Press, 1982.
- [FS16] Bruno Franchi and Raul Paolo Serapioni. Intrinsic Lipschitz graphs within Carnot groups. *J. Geom. Anal.*, 26(3):1946–1994, 2016.
- [FSSC01] Bruno Franchi, Raul Serapioni, and Francesco Serra Cassano. Rectifiability and perimeter in the Heisenberg group. *Math. Ann.*, 321(3):479–531, 2001.
- [FSSC03] Bruno Franchi, Raul Serapioni, and Francesco Serra Cassano. On the structure of finite perimeter sets in step 2 Carnot groups. *The Journal of Geometric Analysis*, 13(3):421–466, 2003.
- [FSSC06] Bruno Franchi, Raul Serapioni, and Francesco Serra Cassano. Intrinsic Lipschitz graphs in Heisenberg groups. *J. Nonlinear Convex Anal.*, 7(3):423–441, 2006.
- [FSSC11] Bruno Franchi, Raul Serapioni, and Francesco Serra Cassano. Differentiability of intrinsic Lipschitz functions within Heisenberg groups. *J. Geom. Anal.*, 21(4):1044–1084, 2011.
- [GMS98] Mariano Giaquinta, Giuseppe Modica, and Jiří Souček. *Cartesian currents in the calculus of variations. I*, volume 37 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1998. Cartesian currents.
- [Hel01] Sigurdur Helgason. *Differential geometry, Lie groups, and symmetric spaces*, volume 34 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001. Corrected reprint of the 1978 original.

- [JS17] Nicolas Juillet and Mario Sigalotti. Pliability, or the Whitney extension theorem for curves in Carnot groups. *Anal. PDE*, 10(7):1637–1661, 2017.
- [LD17] Enrico Le Donne. A primer on Carnot groups: homogenous groups, Carnot-Carathéodory spaces, and regularity of their isometries. *Anal. Geom. Metr. Spaces*, 5:116–137, 2017.
- [LDMO⁺16] Enrico Le Donne, Richard Montgomery, Alessandro Ottazzi, Pierre Pansu, and Davide Vittone. Sard property for the endpoint map on some Carnot groups. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 33(6):1639–1666, 2016.
- [LDOW14] Enrico Le Donne, Alessandro Ottazzi, and Ben Warhurst. Ultrarigid tangents of sub-Riemannian nilpotent groups. *Ann. Inst. Fourier (Grenoble)*, 64(6):2265–2282, 2014.
- [Mag12] Francesco Maggi. *Sets of finite perimeter and geometric variational problems*, volume 135 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2012. An introduction to geometric measure theory.
- [Mar14] Marco Marchi. Regularity of sets with constant intrinsic normal in a class of Carnot groups. *Ann. Inst. Fourier (Grenoble)*, 64(2):429–455, 2014.
- [Mat75] Pertti Mattila. Hausdorff m regular and rectifiable sets in n -space. *Trans. Amer. Math. Soc.*, 205:263–274, 1975.
- [Mat95] Pertti Mattila. *Geometry of sets and measures in Euclidean spaces*, volume 44 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995. Fractals and rectifiability.
- [Mon02] Richard Montgomery. *A tour of subriemannian geometries, their geodesics and applications*, volume 91 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2002.
- [Pan89] Pierre Pansu. Métriques de Carnot-Carathéodory et quasiisométries des espaces symétriques de rang un. *Ann. of Math. (2)*, 129(1):1–60, 1989.
- [SS18] Ludovic Sacchelli and Mario Sigalotti. On the Whitney extension property for continuously differentiable horizontal curves in sub-Riemannian manifolds. *Calc. Var. Partial Differential Equations*, 57(2):Art. 59, 34, 2018.
- [Var84] Veeravalli S. Varadarajan. *Lie groups, Lie algebras, and their representations*, volume 102 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1984. Reprint of the 1974 edition.
- [Ver70] Michèle Vergne. Cohomologie des algèbres de Lie nilpotentes. Application à l'étude de la variété des algèbres de Lie nilpotentes. *Bull. Soc. Math. France*, 98:81–116, 1970.
- [Vit12] Davide Vittone. Lipschitz surfaces, perimeter and trace theorems for BV functions in Carnot-Carathéodory spaces. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 11(4):939–998, 2012.
- [Vit14] Davide Vittone. The regularity problem for sub-Riemannian geodesics. In *Geometric measure theory and real analysis*, volume 17 of *CRM Series*, pages 193–226. Ed. Norm., Pisa, 2014.
- [VSCC92] Nicholas Th. Varopoulos, Laurent Saloff-Coste, and Thierry Coulhon. *Analysis and geometry on groups*, volume 100 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1992.

SEBASTIANO DON: DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX (MAD), FI-40014, UNIVERSITY OF JYVÄSKYLÄ, FINLAND.

E-mail address: `sedon@jyu.fi`

ENRICO LE DONNE: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO B. PONTECORVO 5, 56127 PISA, ITALY, & UNIVERSITY OF JYVÄSKYLÄ, DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX (MAD), FI-40014, FINLAND.

E-mail address: `enrico.ledonne@unipi.it`

TERHI MOISALA: DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX (MAD), FI-40014, UNIVERSITY OF JYVÄSKYLÄ, FINLAND.

E-mail address: `tekamois@jyu.fi`

DAVIDE VITTONI: DIPARTIMENTO DI MATEMATICA TULLIO LEVI-CIVITA, UNIVERSITÀ DI PADOVA, VIA TRIESTE 63, ITALY.

E-mail address: `vittone@math.unipd.it`