

Standing Waves for Nonautonomous Klein-Gordon-Maxwell Systems

Monica Lazzo ^{*} Lorenzo Pisani ^{*}

Abstract

We study a Klein-Gordon-Maxwell system, in a bounded spatial domain, under Neumann boundary conditions on the electric potential. We allow a nonconstant coupling coefficient. For sufficiently small data, we find infinitely many standing waves.

Keywords: Klein-Gordon-Maxwell systems, standing waves, variational methods, Ljusternik-Schnirelmann theory

MSC 2010: 35J50, 35J57, 35Q40, 35Q60

1 Introduction

We are interested in the system of nonautonomous elliptic equations

$$\begin{aligned}\Delta u &= m^2 u - (\omega + q(x)\phi)^2 u && \text{in } \Omega, \\ \Delta \phi &= q(x) (\omega + q(x)\phi) u^2 && \text{in } \Omega.\end{aligned}\tag{1}$$

where Δ is the Laplace operator in \mathbb{R}^3 , $\Omega \subset \mathbb{R}^3$ is a bounded and smooth domain, $m, \omega \in \mathbb{R}$, $q \in L^6(\Omega) \setminus \{0\}$. We complement these equations with the boundary conditions

$$u = 0 \quad \text{on } \partial\Omega,\tag{2a}$$

$$\frac{\partial \phi}{\partial \mathbf{n}} = \alpha \quad \text{on } \partial\Omega,\tag{2b}$$

^{*}Dipartimento di Matematica, Università degli Studi di Bari Aldo Moro, via E. Orabona 4, 70125 Bari, Italy; e-mail: monica.lazzo@uniba.it, lorenzo.pisani@uniba.it

where \mathbf{n} is the unit outward normal vector to $\partial\Omega$ and $\alpha \in H^{1/2}(\partial\Omega)$.

We look for *nontrivial* solutions, by which we mean pairs $(u, \phi) \in H_0^1(\Omega) \times H^1(\Omega)$, satisfying (1)-(2) in the usual weak sense, with $u \neq 0$. Note that, if (u, ϕ) is a nontrivial solution, the pair $(-u, \phi)$ is a nontrivial solution as well.

System (1) arises in connection with the so-called Klein-Gordon-Maxwell equations, which model the interaction of a charged matter field with the electromagnetic field (\mathbf{E}, \mathbf{H}) . They are the Euler-Lagrange equations of the Lagrangian density

$$\begin{aligned} \mathcal{L}_{KGM} = & \frac{1}{2}(|(\partial_t + i q \phi) \psi|^2 - |(\nabla - i q \mathbf{A}) \psi|^2 - m^2 |\psi|^2) + \\ & + \frac{1}{8\pi}(|\nabla \phi + \partial_t \mathbf{A}|^2 - |\nabla \times \mathbf{A}|^2), \end{aligned}$$

where ψ is a complex-valued function representing the matter field, while ϕ and \mathbf{A} are the gauge potentials, related to the electromagnetic field via the equations $\mathbf{E} = -\nabla \phi - \partial_t \mathbf{A}$, $\mathbf{H} = \nabla \times \mathbf{A}$. For the derivation of the Lagrangian density, and details on the physical model, we refer to [4, 5, 15].

Confining attention to *standing waves*, in equilibrium with a purely electrostatic field, amounts to imposing $\psi(t, x) = e^{i\omega t} u(x)$, where u is a real-valued function and ω is a real number, $\mathbf{A} = 0$, and $\phi = \phi(x)$. With these choices, the Klein-Gordon-Maxwell equations considerably simplify and become (1). In the physical model, the boundary condition (2a) means that the matter field is confined to the region Ω , while (2b) amounts to prescribing the normal component of the electric field on $\partial\Omega$. Up to a sign, the surface integral $\int_{\partial\Omega} \alpha \, d\sigma$ represents the flux of the electric field through the boundary of Ω , and thus, the total charge contained in Ω .

Let us point out that in the physical model the coupling coefficient q is constant (see [4, Subsection 5.15]); nonconstant coefficients, however, are worth investigating from a mathematical viewpoint.

For a constant coupling coefficient q , existence results for nontrivial solutions to Problem (1)-(2) were obtained in [13]. In this case, an invariance property holds and solutions to (1)-(2), for arbitrary ω , correspond to solutions of the same system with $\omega = 0$ (that is, *static*

solutions). Thus, with no loss of generality, in [13] the authors confined their attention to static solutions. Their results were generalized to the nonautonomous case in [14], assuming that the coupling coefficient q vanishes at most on a set of measure zero; this restriction on the zero-level set of q was later removed in [16]. Note that, absent the invariance property, the existence of solutions to (1)-(2) with $\omega \neq 0$ does not follow from the results in [14, 16]. Investigating Problem (1)-(2) with $\omega \neq 0$ is precisely the goal of the present paper.

Theorem 1.1. *Assume $\int_{\partial\Omega} \alpha \, d\sigma \neq 0$. Suppose that $|\omega| \leq |m|$ and the function q satisfies the following condition:*

(Q) *there exists $q_0 \in (0, +\infty)$ such that $|\{x \in \Omega \mid 0 < |q(x)| < q_0\}| = 0$.*

Then, if $\|\alpha\|_{H^{1/2}(\partial\Omega)} \|q\|_{L^6(\Omega)}$ is sufficiently small, Problem (1)-(2) has a sequence $\{(u_n, \phi_n)\}$ of distinct nontrivial solutions with the following properties:

- (i) $u_0 \geq 0$ in Ω ;
- (ii) *every bounded subsequence $\{u_{k_n}\}$ satisfies $\|q u_{k_n}\|_{L^3(\Omega)} \rightarrow 0$ as $n \rightarrow \infty$.*

A function q that satisfies (Q) may vanish in Ω , even on a set of positive measure; however, where q does not vanish, it must be bounded away from zero. This condition appears in results on the closedness of the range of the multiplication operator $u \mapsto q u$ (see [18]).

Without assumption (Q), we find nontrivial solutions provided that ω varies in a smaller range.

Theorem 1.2. *Assume $\int_{\partial\Omega} \alpha \, d\sigma \neq 0$. Suppose that $|\omega| \leq |m|/\sqrt{2}$. Then the same conclusions as in Theorem 1.1 hold.*

Following the approach in [16], we apply Ljusternik-Schnirelmann theory to a functional J , defined in an open subset Λ_q of $H_0^1(\Omega)$, whose critical points correspond to nontrivial solutions to Problem (1)-(2). Compared to the functional considered in [16], here J contains several additional terms, which depend on ω . Assuming $|\omega| \leq |m|$, all but one of these additional terms can be easily dealt with and entail no major complications in the study of J . Under assumption (Q), the excep-

tional term (the third summand in (14) below) can be controlled in a uniform fashion (see Lemma 3.3). Without assumption (Q), uniform bounds on the exceptional term are not available (see Remark 3.4). However, J retains its main properties for smaller values of ω , as in Theorem 1.2.

If the data are as small as in Theorem 1.1, the condition $\int_{\partial\Omega} \alpha \, d\sigma \neq 0$ is necessary for the existence of nontrivial solutions, as in [13, 14, 16].

Theorem 1.3. *Assume that $|\omega| \leq |m|$ and $\|\alpha\|_{H^{1/2}(\partial\Omega)} \|q\|_{L^6(\Omega)}$ is as small as required in Theorem 1.1. If $\int_{\partial\Omega} \alpha \, d\sigma = 0$, then Problem (1)-(2) has no nontrivial solutions.*

The assumptions in Theorems 1.1-1.3 are consistent with the literature, albeit on problems in unbounded domains. Limitations on the range of ω already appeared in [19]; later, they were required in [3] and the subsequent stream of related papers. Conditions on the smallest of q (if α is fixed) were imposed, for instance, in [4, Theorem 104], in line with Coleman's conjecture in [11].

We conclude this section by mentioning some recent work, loosely related to our own. In addition to the papers cited in [16], we refer to [9, 7] for results on Klein-Gordon-Maxwell systems in \mathbb{R}^3 . We also refer to [17] for a variant of the system involving fractional operators; to [10, 12] for results on Klein-Gordon-Maxwell-Proca systems; to [6, 8] for Klein-Gordon systems coupled with Born-Infeld type equations.

The paper is organized as follows. In Section 2 we collect some preliminary results and introduce the set Λ_q . In Section 3 we define the functional J and decompose it into the sum of several components, which we analyze separately. In Section 4 we show that J satisfies the requirements in Ljusternik-Schnirelmann theory. Finally, in Section 5 we prove Theorems 1.1-1.3.

2 Preliminaries

Throughout the paper we will use the following notation:

- For any integrable function $f : \Omega \rightarrow \mathbb{R}$, \bar{f} is the average of f in Ω and $\|f\|_p$ is the usual norm in $L^p(\Omega)$ ($p \in [1, \infty]$);

- $H_0^1(\Omega)$ is endowed with the norm $\|\nabla f\|_2$;
- $H^1(\Omega)$ is endowed with the norm $\|f\| := \left(\|\nabla f\|_2^2 + |\bar{f}|^2\right)^{1/2}$;
- for $p \in (1, 6]$, σ_p is the smallest positive number such that $\|f\|_p \leq \sigma_p \|\nabla f\|_2$ for every $f \in H_0^1(\Omega)$;
- for $p \in (1, 6]$, τ_p is the smallest positive number such that $\|f\|_p \leq \tau_p \|f\|$ for every $f \in H^1(\Omega)$;
- $A := \int_{\partial\Omega} \alpha \, d\sigma$, $\|\alpha\|_{1/2} := \|\alpha\|_{H^{1/2}(\partial\Omega)}$.

2.1 Reduction to homogeneous boundary conditions

As in [16], we begin by turning Problem (1)-(2) into an equivalent problem with homogeneous boundary conditions in both variables.

Let $\chi \in H^2(\Omega)$ be the unique solution of

$$\Delta \chi = \frac{A}{|\Omega|} \quad \text{in } \Omega, \quad \frac{\partial \chi}{\partial \mathbf{n}} = \alpha \quad \text{on } \partial\Omega, \quad \int_{\Omega} \chi \, dx = 0. \quad (3)$$

Note that, by elliptic regularity theory and Sobolev's inequalities, there exists $\kappa \in (0, \infty)$ such that

$$\|\chi\|_{\infty} \leq \kappa \|\alpha\|_{1/2}. \quad (4)$$

With $\varphi := \phi - \chi$, Problem (1)-(2) is equivalent to

$$\begin{cases} \Delta u = m^2 u - (\omega + q(\varphi + \chi))^2 u & \text{in } \Omega, \\ \Delta \varphi = q(\omega + q(\varphi + \chi)) u^2 - \frac{A}{|\Omega|} & \text{in } \Omega, \\ u = \frac{\partial \varphi}{\partial \mathbf{n}} = 0 & \text{on } \partial\Omega. \end{cases} \quad (5)$$

Weak solutions of (5) correspond to critical points of the functional F defined in $H_0^1(\Omega) \times H^1(\Omega)$ by

$$F(u, \varphi) = \|\nabla u\|_2^2 + \int_{\Omega} (m^2 - (\omega + q(\varphi + \chi))^2) u^2 \, dx - \|\nabla \varphi\|_2^2 + 2A \bar{\varphi}.$$

Indeed, standard computations show that F is continuously differentiable in $H_0^1(\Omega) \times H^1(\Omega)$ with

$$\begin{aligned} \langle F'_u(u, \varphi), v \rangle &= 2 \int_{\Omega} \left(\nabla u \nabla v + (m^2 - (\omega + q(\varphi + \chi))^2) u v \right) dx, \\ \langle F'_\varphi(u, \varphi), \psi \rangle &= -2 \int_{\Omega} \left(\nabla \varphi \nabla \psi + (q(\omega + q(\varphi + \chi)) u^2 - \frac{A}{|\Omega|}) \psi \right) dx, \end{aligned}$$

for every $u, v \in H_0^1(\Omega)$ and $\varphi, \psi \in H^1(\Omega)$. Since F is unbounded from above and from below, even modulo compact perturbations, a straightforward application of well-known results in critical point theory is precluded.

We follow [3] and associate solutions to Problem (5) with critical points of a functional J that depends only on the variable u and falls within the scope of classical critical point theory. The main ingredient in the construction of J is solving for φ the second equation in (5). We will repeatedly apply the following result.

Proposition 2.1. *For $b \in L^3(\Omega) \setminus \{0\}$ and $\rho \in L^{6/5}(\Omega)$, the homogeneous Neumann problem associated with the equation*

$$-\Delta\varphi + b^2\varphi = \rho \quad (6)$$

has a unique solution $\mathcal{L}_b(\rho)$ in $H^1(\Omega)$. If ρ does not change sign in Ω , then $\rho\mathcal{L}_b(\rho) \geq 0$ in Ω . Furthermore, $\mathcal{L}_b(\rho)$ depends continuously on b and ρ .

Proof. See [16, Proposition 2.1, Remark 2.2, and Remark 2.3]. \square

Remark 2.2. Let $b \in L^3(\Omega) \setminus \{0\}$ and $h \in L^\infty(\Omega)$. Observe that, for any $\tau \in \mathbb{R}$, $\mathcal{L}_b(b^2(h + \tau)) = \mathcal{L}_b(b^2h) + \tau$. With $\tau = -\inf h$ and $\tau = -\sup h$, respectively, Proposition 2.1 implies $\mathcal{L}_b(b^2h) - \inf h \geq 0$ and $\mathcal{L}_b(b^2h) - \sup h \leq 0$ in Ω , whence $\inf h \leq \mathcal{L}_b(b^2h) \leq \sup h$.

In the construction of J , we will consider Equation (6) with $b = qu$; to ensure its solvability, we confine u within the set

$$\Lambda_q := \left\{ u \in H_0^1(\Omega) \mid qu \neq 0 \right\}.$$

The set Λ_q is the complement in $H_0^1(\Omega)$ of the kernel of the bounded and linear operator $u \in H_0^1(\Omega) \mapsto qu \in L^3(\Omega)$. If q vanishes at most on a set of measure zero, then $\Lambda_q = H_0^1(\Omega) \setminus \{0\}$. In general, Λ_q satisfies the following properties.

Proposition 2.3. [16, Proposition 2.4]

- (a) Λ_q is open in $H_0^1(\Omega)$ with $\partial\Lambda_q = \{u \in H_0^1(\Omega) \mid qu = 0\}$.
- (b) If $u \in H_0^1(\Omega)$ and $\text{dist}(u, \partial\Lambda_q) \rightarrow 0$, then $\|qu\|_3 \rightarrow 0$.
- (c) Λ_q contains subsets with arbitrarily large genus.

3 The constrained functional

For $u \in \Lambda_q$, let

$$\rho_u := \frac{A}{|\Omega|} - (qu)^2 \chi - \omega qu^2 \quad (7)$$

and consider Equation (6) with $b = qu$ and $\rho = \rho_u$. By Proposition 2.1, the associated homogeneous Neumann problem has a unique solution $\mathcal{L}_{qu}(\rho_u)$ in $H^1(\Omega)$.

Let us define the map $\Phi : \Lambda_q \longrightarrow H^1(\Omega)$ by letting $\Phi(u) := \mathcal{L}_{qu}(\rho_u)$, and the functional $J : \Lambda_q \longrightarrow \mathbb{R}$ by letting $J(u) := F(u, \Phi(u))$.

Proposition 3.1. (a) *The map Φ is continuously differentiable in Λ_q . The graph of Φ is the set $\{(u, \varphi) \in \Lambda_q \times H^1(\Omega) \mid F'_\varphi(u, \varphi) = 0\}$.*
 (b) *The functional J is continuously differentiable in Λ_q . Furthermore, $(u, \varphi) \in \Lambda_q \times H^1(\Omega)$ is a critical point of F if, and only if, u is a critical point of J and $\varphi = \Phi(u)$.*

Proof. For the proof of Part (a), see [16, Section 3]. Note that all the assertions remain true despite the additional term $-\omega qu^2$ appearing in the right-hand side of (7) when $\omega \neq 0$. Part (b) easily follows from Part (a). \square

On account of Proposition 3.1, nontrivial solutions to Problem (1)-(2) are in one-to-one correspondence with critical points of J in Λ_q .

3.1 Decomposition of J

To simplify the notation, let $\varphi_u := \Phi(u)$. Since φ_u solves the homogeneous Neumann problem associated with the equation

$$-\Delta \varphi + (qu)^2 \varphi = \frac{A}{|\Omega|} - (qu)^2 \chi - \omega qu^2,$$

we get

$$\|\nabla \varphi_u\|_2^2 = A \overline{\varphi}_u - \int_{\Omega} (qu)^2 \chi \varphi_u \, dx - \int_{\Omega} \omega qu^2 \varphi_u \, dx - \int_{\Omega} (qu)^2 \varphi_u^2 \, dx,$$

and thus,

$$\begin{aligned} J(u) = F(u, \varphi_u) &= \|\nabla u\|_2^2 + \int_{\Omega} (m^2 - (\omega + q\chi)^2) u^2 dx + \\ &\quad - \int_{\Omega} (qu)^2 \chi \varphi_u dx - \int_{\Omega} \omega q u^2 \varphi_u dx + A \bar{\varphi}_u. \end{aligned}$$

For every $u \in \Lambda_q$, let

$$\xi_u := -\mathcal{L}_{qu}((qu)^2 \chi), \quad \eta_u := \frac{A}{|\Omega|} \mathcal{L}_{qu}(1), \quad \theta_u := -\mathcal{L}_{qu}(qu^2).$$

Note that η_u , ξ_u , and θ_u satisfy the equations

$$-\Delta \xi_u + (qu)^2 \xi_u = -(qu)^2 \chi, \quad (8)$$

$$-\Delta \eta_u + (qu)^2 \eta_u = \frac{A}{|\Omega|}, \quad (9)$$

$$-\Delta \theta_u + (qu)^2 \theta_u = -qu^2, \quad (10)$$

respectively, with homogeneous Neumann boundary conditions, and

$$\varphi_u = \xi_u + \eta_u + \omega \theta_u. \quad (11)$$

We will write the functional J in terms of u , ξ_u , η_u , and θ_u . Observe that

$$\int_{\Omega} (qu)^2 \chi \theta_u dx = \int_{\Omega} qu^2 \xi_u dx, \quad A \bar{\xi}_u = - \int_{\Omega} (qu)^2 \chi \eta_u dx, \quad (12)$$

$$A \bar{\theta}_u = - \int_{\Omega} qu^2 \eta_u dx, \quad (13)$$

these equalities are easily obtained by multiplying each of the equations (8)-(10) by the solution of the remaining two equations. Taking (11)-(13) into account yields

$$J(u) = \tilde{J}(u) + A \bar{\eta}_u + 2\omega A \bar{\theta}_u \quad (14)$$

for every $u \in \Lambda_q$, with

$$\begin{aligned} \tilde{J}(u) &= \|\nabla u\|_2^2 + \int_{\Omega} (m^2 - \omega^2) u^2 dx - \int_{\Omega} 2\omega qu^2 (\chi + \xi_u) dx + \\ &\quad + 2A \bar{\xi}_u - \int_{\Omega} (qu)^2 \chi^2 dx - \int_{\Omega} (qu)^2 \chi \xi_u dx - \int_{\Omega} \omega^2 qu^2 \theta_u dx. \end{aligned}$$

3.2 Properties of ξ_u , η_u , and θ_u

Since $\xi_u := -\mathcal{L}_{qu}((qu)^2 \chi)$, by Remark 2.2 we have

$$\|\xi_u\|_\infty \leq \|\chi\|_\infty \quad (15)$$

for every $u \in \Lambda_q$.

Lemma 3.2. [16, Lemma 3.3]

- (a) For every $u \in \Lambda_q$, $A\eta_u \geq 0$ in Ω .
- (b) There exists $\gamma \in (0, \infty)$ such that $\|\nabla \eta_u\|_2 \leq \gamma \|qu\|_3^2 |\bar{\eta}_u|$ for every $u \in \Lambda_q$.
- (c) Suppose $A \neq 0$. If $u \in \Lambda_q$ and $\|qu\|_3 \rightarrow 0$, then $A\bar{\eta}_u \rightarrow \infty$.

Lemma 3.3. Suppose that assumption (Q) is satisfied. Then, for every $u \in \Lambda_q$, we have $\|\theta_u\|_\infty \leq 1/q_0$.

Proof. Fix $u \in \Lambda_q$. Define $h \in L^\infty(\Omega)$ by

$$h(x) := \begin{cases} 1/q(x) & \text{if } |q(x)| \geq q_0, \\ 0 & \text{otherwise.} \end{cases}$$

In view of assumption (Q), we have $q = q^2 h$ in Ω , hence $\theta_u := -\mathcal{L}_{qu}(qu^2) = -\mathcal{L}_{qu}((qu)^2 h)$. Then $\|\theta_u\|_\infty \leq \|h\|_\infty$, by Remark 2.2, and the conclusion readily follows. \square

Remark 3.4. If assumption (Q) is satisfied, the map $u \in \Lambda_q \mapsto 2\omega A\bar{\theta}_u$, which appears as the third summand in (14), is bounded from below. Without assumption (Q), this need not be the case.

For instance, suppose that $q \in C(\Omega) \cap L^6(\Omega) \setminus \{0\}$ does not satisfy (Q). Hence, either $\inf \{q(x) \mid q(x) > 0\} = 0$ or $\sup \{q(x) \mid q(x) < 0\} = 0$. In the former case, let $\{s_n\}$ be any unbounded increasing sequence. Up to a subsequence, the open set

$$\Omega_n^+ := \{x \in \Omega \mid s_n < 1/q(x) < s_{n+1}\}$$

is nonempty. Take $u_n \in C_0^\infty(\Omega_n^+) \setminus \{0\} \subset \Lambda_q$. Define $h_n : \Omega \rightarrow \mathbb{R}$ by

$$h_n(x) := \begin{cases} 1/q(x) & \text{if } x \in \Omega_n^+, \\ s_n & \text{otherwise;} \end{cases}$$

clearly, $s_n \leq h_n < s_{n+1}$ in Ω . Since $q u_n^2 = (q u_n)^2 h_n$ in Ω , we have $\theta_{u_n} := -\mathcal{L}_{qu_n}(q u_n^2) = -\mathcal{L}_{qu_n}((q u_n)^2 h_n)$. By Remark 2.2, we get $\theta_{u_n} \leq -s_n$ in Ω and thus, $\bar{\theta}_{u_n} \rightarrow -\infty$. Likewise, in the case $\sup\{q(x) \mid q(x) < 0\} = 0$, we find a sequence $\{u_n\} \subset \Lambda_q$ such that $\bar{\theta}_{u_n} \rightarrow \infty$. Therefore, depending on the sign of ω and A , the map $u \in \Lambda_q \mapsto 2\omega A \bar{\theta}_u$ may be unbounded from below.

Lemma 3.5. *For every $u \in \Lambda_q$, $\left|2\omega A \bar{\theta}_u\right| \leq \int_{\Omega} \omega^2 u^2 dx + A \bar{\eta}_u$.*

Proof. Fix $u \in \Lambda_q$. By (13),

$$\left|2\omega A \bar{\theta}_u\right| = \left|\int_{\Omega} 2\omega q u^2 \eta_u dx\right| \leq \int_{\Omega} \omega^2 u^2 dx + \int_{\Omega} (q u)^2 \eta_u^2 dx.$$

Multiplying (9) by η_u yields

$$\|\nabla \eta_u\|_2^2 + \int_{\Omega} (q u)^2 \eta_u^2 dx = A \bar{\eta}_u.$$

The conclusion readily follows. \square

Lemma 3.6. *Let $\{u_n\} \subset \Lambda_q$ be bounded.*

- (a) *Suppose that $\{\|q u_n\|_3\}$ is bounded away from 0. Then: up to a subsequence, $\{\eta_{u_n}\}$ and $\{\theta_{u_n}\}$ converge in $H^1(\Omega)$.*
- (b) *If $A \neq 0$ and $\|q u_n\|_3 \rightarrow 0$, then $A \bar{\eta}_{u_n} + 2\omega A \bar{\theta}_{u_n} \rightarrow \infty$.*

Proof. (a) Let $\{u_n\} \subset \Lambda_q$ be bounded. Up to a subsequence, $\{u_n\}$ has in $L^6(\Omega)$ a limit u . Since $q u_n \rightarrow q u$ in $L^3(\Omega)$ and $\{\|q u_n\|_3\}$ is bounded away from 0, we deduce $q u \neq 0$; moreover, $q u_n^2 \rightarrow q u^2$ in $L^{6/5}(\Omega)$. Recall that $\eta_{u_n} := \mathcal{L}_{qu_n}(A/|\Omega|)$ and $\theta_{u_n} := -\mathcal{L}_{qu_n}(q u_n^2)$. Thus, by Proposition 2.1, η_{u_n} and θ_{u_n} converge in $H^1(\Omega)$ to $\mathcal{L}_{qu}(A/|\Omega|)$ and $-\mathcal{L}_{qu}(q u^2)$, respectively.

(b) Preliminarily, fix $u \in \Lambda_q$ and note that, by Lemma 3.2(b),

$$\|\eta_u\|^2 = \|\nabla \eta_u\|_2^2 + |\bar{\eta}_u|^2 \leq \left(\gamma^2 \|q u\|_3^4 + 1\right) |\bar{\eta}_u|^2.$$

Thus, by (13), Hölder's inequality, and Sobolev's embedding theorem,

$$\begin{aligned} \left|\omega A \bar{\theta}_u\right| &= \left|\int_{\Omega} \omega q u^2 \eta_u dx\right| \leq |\omega| \|q u\|_3 \|u\|_3 \|\eta_u\|_3 \\ &\leq \sigma_3 \tau_3 |\omega| \|q u\|_3 \|\nabla u\|_2 \|\eta_u\| \\ &\leq \sigma_3 \tau_3 |\omega| \|q u\|_3 \|\nabla u\|_2 \left(\gamma^2 \|q u\|_3^4 + 1\right)^{1/2} |\bar{\eta}_u|. \end{aligned}$$

Therefore,

$$A \bar{\eta}_u + 2 \omega A \bar{\theta}_u \geq |A| |\bar{\eta}_u| - \left| 2 \omega A \bar{\theta}_u \right| \geq (|A| - N(u)) |\bar{\eta}_u|, \quad (16)$$

with

$$N(u) := 2 \sigma_3 \tau_3 |\omega| \|q u\|_3 \|\nabla u\|_2 \left(\gamma^2 \|q u\|_3^4 + 1 \right)^{1/2}.$$

Now assume that $\{u_n\} \subset \Lambda_q$ is bounded and $\|q u_n\|_3 \rightarrow 0$. Then $N(u_n) \rightarrow 0$, and the conclusion follows from (16) and Lemma 3.2(c). \square

3.3 Properties of \tilde{J}

Lemma 3.7. *There exist $C_1 \in \mathbb{R}$, $C_2 \in (0, \infty)$, and $C_3 \in [0, \infty)$, which depend on Ω , ω , and the norms $\|q\|_6$ and $\|\alpha\|_{1/2}$, such that*

$$\tilde{J}(u) \geq C_1 \|\nabla u\|_2^2 + \int_{\Omega} (m^2 - \omega^2) u^2 dx - 2 \kappa |A| \|\alpha\|_{1/2} \quad (17)$$

and

$$\tilde{J}(u) \leq (C_2 + C_3 \|\theta_u\|) \|\nabla u\|_2^2 + 2 \kappa |A| \|\alpha\|_{1/2} \quad (18)$$

for every $u \in \Lambda_q$.

Proof. Fix $u \in \Lambda_q$ and recall that

$$\begin{aligned} \tilde{J}(u) &= \|\nabla u\|_2^2 + \int_{\Omega} (m^2 - \omega^2) u^2 dx - \int_{\Omega} 2 \omega q u^2 (\chi + \xi_u) dx + \\ &\quad + 2 A \bar{\xi}_u - \int_{\Omega} (q u)^2 \chi^2 dx - \int_{\Omega} (q u)^2 \chi \xi_u dx - \int_{\Omega} \omega^2 q u^2 \theta_u dx. \end{aligned}$$

By Hölder's inequality, Sobolev's embedding theorem, (4), and (15),

$$\begin{aligned} \left| \int_{\Omega} q u^2 (\chi + \xi_u) dx \right| &\leq 2 \|\chi\|_{\infty} \|q\|_6 \|u\|_{12/5}^2 \\ &\leq 2 \kappa \sigma_{12/5}^2 \|\alpha\|_{1/2} \|q\|_6 \|\nabla u\|_2^2, \end{aligned} \quad (19)$$

$$\int_{\Omega} (q u)^2 \chi^2 dx \leq \|\chi\|_{\infty}^2 \|q\|_6^2 \|u\|_3^2 \leq \kappa^2 \sigma_3^2 \|\alpha\|_{1/2}^2 \|q\|_6^2 \|\nabla u\|_2^2, \quad (20)$$

and

$$|\bar{\xi}_u| \leq \kappa \|\alpha\|_{1/2}. \quad (21)$$

Multiplying (8) and (10) by ξ_u and θ_u , respectively, gives

$$-\int_{\Omega} (qu)^2 \chi \xi_u dx \geq 0, \quad -\int_{\Omega} \omega^2 q u^2 \theta_u dx \geq 0. \quad (22)$$

Taking (19)-(22) into account yields (17), with

$$C_1 := 1 - 4|\omega| \kappa \sigma_{12/5}^2 \|\alpha\|_{1/2} \|q\|_6 - \kappa^2 \sigma_3^2 \|\alpha\|_{1/2}^2 \|q\|_6^2. \quad (23)$$

To prove (18), in addition to (19)-(21) observe that

$$\left| \int_{\Omega} (qu)^2 \chi \xi_u dx \right| \leq \|\chi\|_{\infty}^2 \|q\|_6^2 \|u\|_3^2 \leq \kappa^2 \sigma_3^2 \|\alpha\|_{1/2}^2 \|q\|_6^2 \|\nabla u\|_2^2$$

and

$$\left| \int_{\Omega} q u^2 \theta_u dx \right| \leq \|q\|_6 \|u\|_3^2 \|\theta_u\|_6 \leq \sigma_3^2 \|q\|_6 \|\nabla u\|_2^2 \|\theta_u\|.$$

Thus, (18) follows with

$$C_2 := 1 + |m^2 - \omega^2| \sigma_2^2 + 4|\omega| \kappa \sigma_{12/5}^2 \|\alpha\|_{1/2} \|q\|_6 + 2\kappa^2 \sigma_3^2 \|\alpha\|_{1/2}^2 \|q\|_6^2$$

and $C_3 := \omega^2 \sigma_3^2 \tau_6 \|q\|_6$. \square

4 Properties of J

Throughout this section, we will assume $A \neq 0$. For ease of discussion, we will refer to Case 1, if $|\omega| \leq |m|$ and (Q) is satisfied, and to Case 2, if $|\omega| \leq |m|/\sqrt{2}$. In either case, we will assume that $\|\alpha\|_{1/2} \|q\|_6$ is so small that the constant C_1 , as defined in (23), is strictly positive.

Proposition 4.1. *The functional J is bounded from below and coercive in Λ_q .*

Proof. Fix $u \in \Lambda_q$. In Case 1, Lemma 3.3 applies and implies

$$A\bar{\eta}_u + 2\omega A\bar{\theta}_u \geq -\frac{2|A||\omega|}{q_0},$$

in view of Lemma 3.2(a). Therefore, (14) and (17) yield

$$J(u) \geq C_1 \|\nabla u\|_2^2 + \int_{\Omega} (m^2 - \omega^2) u^2 dx - 2\kappa |A| \|\alpha\|_{1/2} - \frac{2|A||\omega|}{q_0}.$$

In Case 2, Lemma 3.5 implies

$$A\bar{\eta}_u + 2\omega A\bar{\theta}_u \geq - \int_{\Omega} \omega^2 u^2 dx.$$

Therefore, (14) and (17) yield

$$\begin{aligned} J(u) &\geq C_1 \|\nabla u\|_2^2 + \int_{\Omega} (m^2 - \omega^2) u^2 dx - 2\kappa |A| \|\alpha\|_{1/2} - \int_{\Omega} \omega^2 u^2 dx \\ &= C_1 \|\nabla u\|_2^2 + \int_{\Omega} (m^2 - 2\omega^2) u^2 dx - 2\kappa |A| \|\alpha\|_{1/2}. \end{aligned}$$

With $C_1 \in (0, \infty)$, the conclusions readily follow in both cases. \square

Proposition 4.2.

- (a) *Every sequence $\{u_n\} \subset \Lambda_q$ such that $\|qu_n\|_3 \rightarrow 0$ has a subsequence $\{u_{k_n}\}$ such that $J(u_{k_n}) \rightarrow \infty$.*
- (b) *J has complete sublevels.*

Proof. (a) Let $\{u_n\} \subset \Lambda_q$ and assume $\|qu_n\|_3 \rightarrow 0$.

In Case 1, note that $A\bar{\eta}_{u_n} \rightarrow \infty$, by Lemma 3.2(c), whereas $\{\bar{\theta}_{u_n}\}$ and $\{\tilde{J}(u_n)\}$ are bounded from below, by Lemma 3.3 and (17). Since $J(u_n) = \tilde{J}(u_n) + A\bar{\eta}_{u_n} + 2\omega A\bar{\theta}_{u_n}$, we deduce $J(u_n) \rightarrow \infty$.

In Case 2, we consider two possibilities. If $\{\|\nabla u_n\|_2\}$ is unbounded, there exists a subsequence $\{u_{k_n}\}$ such that $\|\nabla u_{k_n}\|_2 \rightarrow \infty$; thus, $J(u_{k_n}) \rightarrow \infty$, for J is coercive by Proposition 4.1. If $\{\|\nabla u_n\|_2\}$ is bounded, then $A\bar{\eta}_{u_n} + 2\omega A\bar{\theta}_{u_n} \rightarrow \infty$, by Lemma 3.6(b), whereas $\{\tilde{J}(u_n)\}$ is bounded from below, by (17). Since $J(u_n) = \tilde{J}(u_n) + A\bar{\eta}_{u_n} + 2\omega A\bar{\theta}_{u_n}$, we deduce $J(u_n) \rightarrow \infty$.

(b) Suppose that $\{u_n\} \subset \Lambda_q$, $J(u_n) \leq c$, for some $c \in \mathbb{R}$, and $u_n \rightarrow u$ in $H_0^1(\Omega)$. By Part (a), the sequence $\{\|qu_n\|_3\}$ is bounded away from 0 and thus, by Proposition 2.3(b), $u \in \Lambda_q$. \square

Proposition 4.3. *For any $\{u_n\} \subset \Lambda_q$, the sequence $\{J(u_n)\}$ is unbounded if, and only if, either $\{u_n\}$ is unbounded or $\{\|qu_n\|_3\}$ is not bounded away from 0.*

Proof. The “if” part of the statement easily follows from Proposition 4.1 and Proposition 4.2. We will prove the “only if” part by

way of contradiction. Suppose that there exists a bounded sequence $\{u_n\} \subset \Lambda_q$ such that $\{\|q u_n\|_3\}$ is bounded away from 0 and $J(u_n) \rightarrow \infty$.

Up to a subsequence, $\{\eta_{u_n}\}$ and $\{\theta_{u_n}\}$ converge in $H^1(\Omega)$, in view of Lemma 3.6(a). This clearly implies that $\{\bar{\eta}_{u_n}\}$ and $\{\bar{\theta}_{u_n}\}$ are bounded and thus, $\{\tilde{J}(u_n)\}$ is bounded from above, by (18). Since $J(u_n) = \tilde{J}(u_n) + A\bar{\eta}_{u_n} + 2\omega A\bar{\theta}_{u_n}$, we deduce that $\{J(u_n)\}$ is bounded from above, a contradiction. \square

Proposition 4.4. *The functional J satisfies the Palais-Smale condition in Λ_q .*

Proof. Suppose that $\{u_n\} \subset \Lambda_q$ is a Palais-Smale sequence, that is, $\{J(u_n)\}$ is bounded and $J'(u_n) \rightarrow 0$; we have to show that, up to a subsequence, $\{u_n\}$ converges in Λ_q .

Since J is coercive, $\{u_n\}$ is bounded in $H_0^1(\Omega)$; up to a subsequence, it converges weakly to some $u \in H_0^1(\Omega)$. Observe that

$$\Delta u_n = -\frac{1}{2} J'(u_n) + m^2 u_n - (\omega + q(\eta_{u_n} + \xi_{u_n} + \omega \theta_{u_n} + \chi))^2 u_n. \quad (24)$$

The first two summands in the right-hand side of (24) are clearly bounded in $H^{-1}(\Omega)$; we will show that the same is true for the third summand.

Since $\{J(u_n)\}$ is bounded, Proposition 4.2 implies that $\{\|q u_n\|_3\}$ is bounded away from 0. Lemma 3.6(a) applies: up to a subsequence, $\{\eta_{u_n}\}$ and $\{\theta_{u_n}\}$ converge in $H^1(\Omega)$, and are therefore bounded in $L^6(\Omega)$. By (15), $\{\xi_{u_n} + \chi\}$ is bounded in $L^6(\Omega)$ as well. It follows that $\{(\eta_{u_n} + \xi_{u_n} + \omega \theta_{u_n} + \chi)^2\}$ is bounded in $L^3(\Omega)$, which in turn implies that $\{q^2(\eta_{u_n} + \xi_{u_n} + \omega \theta_{u_n} + \chi)^2 u_n\}$ is bounded in $L^{6/5}(\Omega)$, hence in $H^{-1}(\Omega)$.

On account of (24), the sequence $\{\Delta u_n\}$ is bounded in $H^{-1}(\Omega)$; the compactness of the inverse Laplace operator implies that, up to a subsequence, $\{u_n\}$ converges to u in $H_0^1(\Omega)$. By Proposition 4.2(b), $u \in \Lambda_q$. \square

5 Proof of the main results

Proof of Theorems 1.1 and 1.2. On account of the correspondence between critical points of J and nontrivial solutions to Problem (1)-(2), it suffices to prove that J has a sequence of critical points $\{u_n\} \subset \Lambda_q$ satisfying (i) and (ii). Observe that $J(u) = J(|u|)$ for every $u \in \Lambda_q$. This easily follows from the fact that $\Phi(u) = \Phi(|u|)$ for every $u \in \Lambda_q$, by the very definition of Φ .

Suppose that $A \neq 0$ and $\|\alpha\|_{1/2} \|q\|_6$ is so small that the constant C_1 , as defined in (23), is strictly positive. As we have shown in Section 4, under the assumptions in Theorems 1.1 and 1.2, the functional J is bounded from below, has complete sublevels, and satisfies the Palais-Smale condition in Λ_q . These properties readily imply that J attains its minimum at some $u_0 \in \Lambda_q$; by the observation above, we can assume $u_0 \geq 0$ in Ω .

By Proposition 2.3(c), the set Λ_q has infinite genus. Thus, Ljusternik-Schnirelmann Theory applies (see [20, Corollary 4.1] and [1, Remark 3.6]) and J has a sequence $\{u_n\}_{n \geq 1}$ of critical points in Λ_q . Standard arguments show that $J(u_n) \rightarrow \infty$ (see [2, Chapter 10]).

Let $\{v_n\}$ be a bounded subsequence of $\{u_n\}$. In view of Proposition 4.3, every subsequence of $\{v_n\}$ has a subsequence $\{v_{k_n}\}$ such that $\|q v_{k_n}\|_3 \rightarrow 0$; this proves that $\|q v_n\|_3 \rightarrow 0$. \square

Proof of Theorem 1.3. Assume that $|\omega| \leq |m|$ and $\|\alpha\|_{1/2} \|q\|_6$ is so small that the constant C_1 , as defined in (23), is strictly positive.

Suppose that (u, ϕ) is a solution to (1)-(2) with $A = 0$ and let $\varphi := \phi - \chi$. Then, (u, φ) is a solution to

$$\begin{cases} \Delta u = m^2 u - (\omega + q(\varphi + \chi))^2 u & \text{in } \Omega, \\ \Delta \varphi = q(\omega + q(\varphi + \chi)) u^2 & \text{in } \Omega, \\ u = \frac{\partial \varphi}{\partial \mathbf{n}} = 0 & \text{on } \partial \Omega. \end{cases} \quad (25)$$

Multiplying by u the first equation in (25) gives

$$\begin{aligned} 0 &= \|\nabla u\|_2^2 + \int_{\Omega} m^2 u^2 dx - \int_{\Omega} (\omega + q(\varphi + \chi))^2 u^2 dx \\ &= \|\nabla u\|_2^2 + \int_{\Omega} (m^2 - \omega^2) u^2 dx - \int_{\Omega} (qu)^2 \varphi^2 dx - \int_{\Omega} 2\omega q u^2 \chi dx + \\ &\quad - \int_{\Omega} (qu)^2 \chi^2 dx - \int_{\Omega} 2(qu)^2 \varphi \chi dx - \int_{\Omega} 2\omega q u^2 \varphi dx. \end{aligned}$$

Multiplying by φ the second equation in (25) gives

$$\|\nabla \varphi\|_2^2 + \int_{\Omega} (qu)^2 \varphi^2 dx = - \int_{\Omega} (qu)^2 \varphi \chi dx - \int_{\Omega} \omega q u^2 \varphi dx. \quad (26)$$

Substituting (26) into the preceding equality gives

$$\begin{aligned} 0 &= \|\nabla u\|_2^2 + \int_{\Omega} (m^2 - \omega^2) u^2 dx + \int_{\Omega} (qu)^2 \varphi^2 dx - \int_{\Omega} 2\omega q u^2 \chi dx + \\ &\quad - \int_{\Omega} (qu)^2 \chi^2 dx + 2\|\nabla \varphi\|_2^2. \end{aligned}$$

Neglecting the nonnegative terms, and recalling (19), (20), and the definition of C_1 , we obtain

$$\begin{aligned} 0 &\geq \|\nabla u\|_2^2 - \int_{\Omega} 2\omega q u^2 \chi dx - \int_{\Omega} (qu)^2 \chi^2 dx \\ &\geq \left[1 - 2|\omega| \kappa \sigma_{12/5}^2 \|\alpha\|_{1/2} \|q\|_6 - \kappa^2 \sigma_3^2 \|\alpha\|_{1/2}^2 \|q\|_6^2 \right] \|\nabla u\|_2^2 \\ &\geq C_1 \|\nabla u\|_2^2, \end{aligned}$$

which implies $u = 0$. \square

References

- [1] A. Ambrosetti and V. Coti Zelati, Periodic solutions of singular lagrangian systems, Progr. Nonlinear Differential Equations Appl. 10, Birkhäuser, Boston, 1993.
- [2] A. Ambrosetti and A. Malchiodi, Nonlinear analysis and semilinear elliptic problems. Cambridge Studies in Advanced Mathematics, 104. Cambridge University Press, Cambridge, 2007.
- [3] V. Benci and D. Fortunato, Solitary waves of the nonlinear Klein-Gordon equation coupled with the Maxwell equations, Rev. Math. Phys. **14** (2002), 409-420.

- [4] V. Benci and D. Fortunato, Variational methods in nonlinear field equations. Solitary waves, hylomorphic solitons and vortices, Springer, Cham, 2014.
- [5] D. Bleecker, Gauge theory and variational principles, Dover Publications, 2005.
- [6] D. Bonheure, P. d'Avenia and A. Pomponio, On the electrostatic Born-Infeld equation with extended charges, Comm. Math. Phys. **346** (2016), 877–906.
- [7] S. J. Chen and L. Li, Infinitely many solutions for Klein-Gordon-Maxwell system with potentials vanishing at infinity, Z. Anal. Anwend. **37** (2018), 3950.
- [8] S. J. Chen and S. Z. Song, The existence of multiple solutions for the KleinGordon equation with concave and convex nonlinearities coupled with BornInfeld theory on \mathbb{R}^3 , Nonlinear Anal. Real World Appl. **38** (2017), 78-95.
- [9] S. Chen and X. Tang, Improved results for Klein-Gordon-Maxwell systems with general nonlinearity, Discrete Contin. Dyn. Syst. **38** (2018), 2333-2348.
- [10] M. Clapp, M. Ghimenti and A. M. Micheletti, Boundary layers to a singularly perturbed Klein-Gordon-Maxwell-Proca system in a compact Riemannian manifold with boundary, Adv. Nonlinear Anal., DOI: <https://doi.org/10.1515/anona-2017-0039>
- [11] S. Coleman, Q-balls, Nuclear Phys. B **262** (1985), 263-283; Erratum **269** (1986), 744-745.
- [12] P. d'Avenia, J. Mederski and A. Pomponio, Vortex ground states for Klein-Gordon-Maxwell-Proca type systems, J. Math. Phys. **58** (2017), 041503, 19 pp.
- [13] P. d'Avenia, L. Pisani and G. Siciliano, Klein-Gordon-Maxwell systems in a bounded domain, Discrete Contin. Dyn. Syst. **26** (2010), 135-149.
- [14] P. d'Avenia, L. Pisani and G. Siciliano, Nonautonomous Klein-Gordon-Maxwell systems in a bounded domain, Adv. Nonlinear Anal. **3** (2014), suppl. 1 s37-s45.

- [15] B. Felsager, *Geometry, Particles and Fields*, Springer, 1998.
- [16] M. Lazzo and L. Pisani, Klein-Gordon-Maxwell systems with non-constant coupling coefficient, *Adv. Nonlinear Stud.* **18** (2018) 55-63.
- [17] O. H. Miyagaki, E. L. de Moura and R. Ruviano, Positive ground state solutions for quasiscritical the [sic] fractional KleinGordonMaxwell system with potential vanishing at infinity, *Complex Variables and Elliptic Equations*, DOI: <https://doi.org/10.1080/17476933.2018.1434625>
- [18] J. C. Ramos Fernández, Some properties of multiplication operators acting on Banach spaces of measurable functions, *Bol. Mat.* **23** (2016), 221-237.
- [19] W. A. Strauss, Existence of solitary waves in higher dimensions, *Comm. Math. Phys.* **55** (1977), 149162.
- [20] A. Szulkin, Ljusternik-Schnirelmann theory on C^1 -manifolds, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **5** (1988), 119-139.