

Compatibility of any pair of 2-outcome measurements characterizes the Choquet simplex

Yui Kuramochi

the date of receipt and acceptance should be inserted later

Abstract For a compact convex subset K of a locally convex Hausdorff space, a measurement on $A(K)$ is a finite family of positive elements in $A(K)$ normalized to the unit constant 1_K , where $A(K)$ denotes the set of continuous real affine functionals on K . It is proved that a compact convex set K is a Choquet simplex if and only if any pair of 2-outcome measurements are compatible, i.e. the measurements are given as the marginals of a single measurement. This generalizes the finite-dimensional result of [Plávala M 2016 Phys. Rev. A **94**, 042108] obtained in the context of the foundations of quantum theory.

Keywords Choquet simplex · Bauer simplex · general probabilistic theory · compatibility of measurements

Mathematics Subject Classification (2010) 46A55 · 46B40 · 81P16 · 81P15

1 Introduction and main results

Let K be a compact convex subset of a locally convex Hausdorff space over the reals \mathbb{R} and let $A(K)$ denote the set of continuous real affine functionals on K , where $A(K)$ is a Banach space under the supremum norm $\|f\| := \sup_{x \in K} |f(x)|$ ($f \in A(K)$). (Throughout the paper all the linear spaces are over \mathbb{R} .) A measurement on $A(K)$ is a finite sequence $(f_k)_{k=1}^m \in A(K)^m$ for some integer m such that $f_k \geq 0$ ($1 \leq k \leq m$) and $\sum_{k=1}^m f_k = 1_K$, where

This work was supported by Cross-ministerial Strategic Innovation Promotion Program (SIP) (Council for Science, Technology and Innovation (CSTI)).

Yui Kuramochi
Photon Science Center, Graduate School of Engineering, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan
Tel.: +81-03-5841-0859
E-mail: kuramochi@qi.t.u-tokyo.ac.jp

$1_K \equiv 1$ is the unit constant functional. A measurement belonging to $A(K)^m$ is called m -outcome. Two measurements $(f_k)_{k=1}^m$ and $(g_j)_{j=1}^n$ on $A(K)$ are called *compatible* (or *jointly measurable*) if there exists a family $(h_{k,j})_{1 \leq k \leq m, 1 \leq j \leq n}$ in $A(K)$ such that

$$h_{k,j} \geq 0, \quad f_k = \sum_{j'=1}^n h_{k,j'}, \quad g_j = \sum_{k'=1}^m h_{k',j} \quad (1 \leq k \leq m, 1 \leq j \leq n).$$

In [5] Plávala showed that for finite-dimensional K , K is a simplex if and only if any pair of 2-outcome measurements are compatible. In the physical context, a compact convex set K corresponds to a state space of a general physical system and Plávala's result indicates that incompatibility of measurements [3] characterizes non-classicality of a physical system. The proof in [5] depends on the notion of the maximal face and is not straightforwardly applicable to infinite-dimensional compact sets. The purpose of this paper is to generalize this result to an arbitrary compact convex set K (Theorems 1 and 2).

The first main result of this paper is the following theorem.

Theorem 1 *Let K be a compact convex subset of a locally convex Hausdorff space. Then K is a Choquet simplex if and only if any pair $(f_k)_{k=1}^2$ and $(g_j)_{j=1}^2$ of 2-outcome measurements on $A(K)$ are compatible.*

Here a compact convex set K is called a Choquet simplex if the Banach dual $A(K)^*$ equipped with the dual positive cone

$$A(K)_+^* = \{ \psi \in A(K)^* \mid \psi(f) \geq 0 (\forall f \in A(K)_+) \}$$

is a lattice, where $A(K)_+ := \{ f \in A(K) \mid f(x) \geq 0 (\forall x \in K) \}$.

A Banach space E is said to have a Banach predual E_* if E_* is a Banach space such that its Banach dual $(E_*)^*$ is isometrically isomorphic to E . We can and do take such E_* , if exists, as a closed linear subspace of the Banach dual E^* . The second main result of this paper is then the following theorem.

Theorem 2 *Let K be a compact convex subset of a locally convex Hausdorff space. Suppose that $A(K)$ has a Banach predual. Then the following conditions are equivalent.*

- (i) K is a Bauer simplex.
- (ii) K is a Choquet simplex.
- (iii) Any pair $(f_k)_{k=1}^2$ and $(g_j)_{j=1}^2$ of 2-outcome measurements on $A(K)$ are compatible.

Here K is a Bauer simplex if the set $\partial_e K$ of extremal points of K is compact and any point in K is the barycenter of a unique boundary measure [1].

The above theorems are rephrased in terms of order-unit Banach space by using the following one-to-one correspondence ([1], Sections II.1 and II.2). A pair (E, e) of an Archimedean ordered linear space E and an order unit $e \in E$ is called an order-unit Banach space if the order-unit norm $\| \cdot \|$ on E induced by e is complete. We write by E_+ the positive cone of E . For each

order-unit Banach space (E, e) its dual E^* is a base-normed ordered linear space with the positive cone $E_+^* := \{ \phi \in E^* \mid \phi(a) \geq 0 \ (a \in E_+) \}$ and the base $S(E) := \{ \phi \in E_+^* \mid \|\phi\| = \phi(e) = 1 \}$. Each element of $S(E)$ is called a state on E . $S(E)$ is a weakly* compact convex subset of E^* and the order-unit Banach space $(A(S(E)), 1_{S(E)})$ is shown to be order and isometrically isomorphic to (E, e) . Conversely for any compact convex subset K of a locally convex Hausdorff space, $(A(K), 1_K)$ is an order-unit Banach space and K is continuously affine isomorphic to the set $S(A(K))$ of states on $A(K)$.

We also define the (m -outcome) measurement on an order-unit Banach space (E, e) as a finite-sequence (belonging to E^m) of positive elements of E normalized to e . The compatibility of measurements is defined in the same way as that of measurements on $A(K)$. By this correspondence we can readily see that Theorems 1 and 2 are respectively equivalent to

Theorem 3 *Let (E, e) be an order-unit Banach space. Then E^* is a lattice if and only if any pair of 2-outcome measurements $(a_j)_{j=1}^2$ and $(b_k)_{k=1}^2$ on (E, e) are compatible.*

Theorem 4 *Let (E, e) be an order-unit Banach space. Suppose that E has a Banach predual E_* . Then the following conditions are equivalent.*

- (i) $S(E)$ is a Bauer simplex.
- (ii) E^* is a lattice.
- (iii) Any pair of 2-outcome measurements $(a_j)_{j=1}^2$ and $(b_k)_{k=1}^2$ on (E, e) are compatible.

In the rest of this paper, after introducing some preliminaries in Section 2, we prove in Section 3 the main results Theorems 3 and 4.

2 Preliminary

This section reviews necessary results of order-unit Banach spaces with Banach preduals and those of simplexes. The reader is referred to [1] and [2, 4] for the complete proofs of the facts given in this section.

Let (E, e) be an order-unit Banach space with a Banach predual E_* . Then the predual E_* endowed with the predual positive cone

$$E_{*+} := \{ \psi \in E_* \mid \psi(a) \geq 0 \ (\forall a \in E_+) \}$$

has the base

$$S_*(E) := \{ \psi \in E_{*+} \mid \psi(e) = 1 \} = E_* \cap S(E)$$

and the base norm on E_* induced by $S_*(E)$ coincides with the predual norm

$$\|\psi\| = \sup_{a \in E: \|a\| \leq 1} |\psi(a)| \quad (\psi \in E_*),$$

which is the norm on E^* restricted to E_* ([2], Theorem 6; [4]). Moreover the positive cone E_+ is weakly* closed (i.e. $\sigma(E, E_*)$ -closed) ([4], in the proof). Therefore by the bipolar theorem E_+ is the dual cone of E_{*+} , i.e.

$$E_+ = \{ a \in E \mid \psi(a) \geq 0 \ (\forall \psi \in E_{*+}) \}.$$

By the Banach-Alaoglu theorem, the unit ball of E is weakly* compact and hence so is any interval

$$[a, b] := \{ x \in E \mid a \leq x \leq b \} \quad (a, b \in E).$$

We have seen that a Banach predual E_* of an order-unit Banach space E is a base-normed Banach space and the positive cone of E is the dual cone of that of E_* . We can conversely show that the Banach dual $(E_*)^*$ of a base-normed Banach space E_* is an order-unit Banach space, where the positive cone of $(E_*)^*$ is the dual cone of that of E_* ([2]; [1], II.1.15). Thus we can start either from an order-unit Banach space E with a Banach predual or from a base-normed space E_* , and these definitions result in effectively the same concept. We exploit the former definition since the main Theorems 2 and 4 are stated in terms of this.

A net $(x_i)_{i \in I}$ in an ordered linear space E is called bounded if $x_i \leq a$ ($\forall i \in I$) for some $a \in E$ and increasing if $i \leq i'$ implies $x_i \leq x_{i'}$ ($i, i' \in I$). We denote the supremum of the subset $\{ x_i \mid i \in I \}$ of E by $\sup_{i \in I} x_i$ if it exists.

The following lemma will be used in the proof of Theorem 4.

Lemma 1 *Let (E, e) be an order-unit Banach space with a Banach predual E_* . Then any bounded increasing net $(x_i)_{i \in I}$ in E is weakly* convergent to the supremum $\sup_{i \in I} x_i$.*

Proof Take an element $a \in E$ such that $x_i \leq a$ ($i \in I$). For a fixed $i_0 \in I$ we have $x_i \in [x_{i_0}, a]$ eventually and by the compactness of $[x_{i_0}, a]$ there exists a subnet $(x_{i(j)})_{j \in J}$ weakly* converging to some $x \in [x_{i_0}, a]$. Since $x_i \leq x_{i(j)}$ eventually for each $i \in I$, the weak* closedness of the positive cone E_+ implies $x_i \leq x$ for all $i \in I$. Moreover if $x_i \leq b \in E$ ($i \in I$), then again by the weak* closedness of E_+ we have $x \leq b$. Therefore $x = \sup_{i \in I} x_i$. Take arbitrary $\psi \in E_{*+}$. Then $(\psi(x_i))_{i \in I}$ is an increasing net in \mathbb{R} and

$$\psi(x) \geq \sup_{i \in I} \psi(x_i) \geq \sup_{j \in J} \psi(x_{i(j)}) = \psi(x),$$

which implies $\psi(x_i) \uparrow \psi(x)$. Since E_* is the linear span of E_{*+} , this implies $x_i \xrightarrow{\text{weakly*}} x$. \square

Let (E, e) be an order-unit Banach space. We define the linear order \leq on the double dual space E^{**} corresponding to the double dual cone

$$E_+^{**} := \{ x'' \in E^{**} \mid \langle x'', \psi \rangle \geq 0 \ (\forall \psi \in E_+^*) \},$$

where we introduced the notation $\langle x'', \psi \rangle := x''(\psi)$ ($x'' \in E^{**}, \psi \in E^*$). As usual we regard E as a norm-closed subspace of E^{**} . Then by the bipolar

theorem we have $E_+^{**} \cap E = E_+$, which implies that the order on E^{**} restricted to E coincides with the original order on E . Moreover (E^{**}, e) is an order-unit Banach space with the Banach predual E^* and E_+^{**} is the weak* closure of E_+ by the bipolar theorem. Let us introduce the sets

$$\begin{aligned}\mathcal{E}(E) &:= \{x \in E \mid 0 \leq x \leq e\}, \\ \mathcal{E}(E^{**}) &:= \{x'' \in E^{**} \mid 0 \leq x'' \leq e\}.\end{aligned}$$

Each element of $\mathcal{E}(E)$ or $\mathcal{E}(E^{**})$ is called an *effect*.

Lemma 2 *Let (E, e) be an order-unit Banach space. Then $\mathcal{E}(E)$ is weakly* dense in $\mathcal{E}(E^{**})$.*

Proof Suppose that $\mathcal{E}(E)$ is not weakly* dense in $\mathcal{E}(E^{**})$. Then by the Hahn-Banach separation theorem, there exist $\psi \in E^*$ and $x'' \in \mathcal{E}(E^{**})$ such that

$$\sup_{x \in \mathcal{E}(E)} \langle x, \psi \rangle < \langle x'', \psi \rangle.$$

Since the unit ball of E coincides with

$$\{-e + 2x \mid x \in \mathcal{E}(E)\},$$

we have

$$\|\psi\| = \sup_{x \in \mathcal{E}(E)} \langle -e + 2x, \psi \rangle < \langle -e + 2x'', \psi \rangle \leq \| -e + 2x'' \| \|\psi\| \leq \|\psi\|,$$

which is a contradiction. \square

The following well-known characterizations of the Choquet and Bauer simplexes are also necessary.

Lemma 3 ([1], II.3.1 and II.3.11) *Let (E, e) be an order-unit Banach space. Then E^* is a lattice if and only if E has the Riesz decomposition property, i.e.*

$$\begin{aligned}0 \leq u \leq v_1 + v_2, \quad 0 \leq v_1, v_2 \\ \implies \exists u_1, u_2 \in E: 0 \leq u_j \leq v_j \ (j = 1, 2), \quad u = u_1 + u_2\end{aligned}$$

holds for any $u, v_1, v_2 \in E$.

Lemma 4 ([1], II.4.1) *Let (E, e) be an order-unit Banach space. Then $S(E)$ is a Bauer simplex if and only if E is a lattice.*

3 Proofs of Theorems 3 and 4

In this section, we first prove Theorem 4 and then Theorem 3 by reducing to Theorem 4.

Let (E, e) be an order-unit Banach space. For each effect $a \in \mathcal{E}(E)$ there corresponds a 2-outcome measurement $(a, e - a)$ on (E, e) . Two effects $a, b \in E$ are said to be compatible if the corresponding 2-outcome measurements $(a, e - a)$ and $(b, e - b)$ are compatible. The following lemma is immediate from the definition of the compatibility (cf. [5], Proposition 7).

Lemma 5 *Let (E, e) be an order-unit Banach space. Then effects $a, b \in \mathcal{E}(E)$ are compatible if and only if there exists $c \in E$ such that*

$$0, a + b - e \leq c \leq a, b.$$

We also introduce the following notion of orthogonality.

Definition 1 Positive elements a and b in an ordered linear space E is said to be *orthogonal* if

$$0 \leq c \leq a, b \implies c = 0$$

for any $c \in E$. It readily follows that for positive elements $a, b \in E$ and positive reals $\alpha, \beta > 0$, a and b are orthogonal if and only if αa and βb are orthogonal.

Then we have

Lemma 6 *Let (E, e) be an order-unit Banach space satisfying the condition (iii) of Theorem 4 and let $a, b \in E$ be orthogonal positive elements. Suppose that $a \neq 0$. Then for any $\psi \in S(E)$ with $\psi(a) = \|a\|$, it holds that $\psi(b) = 0$.*

Proof The statement is obvious when $b = 0$ and we assume $b \neq 0$. Then since the effects $\|a\|^{-1}a$ and $\|b\|^{-1}b$ are compatible, by Lemma 5 there exists $c \in E$ such that

$$0, \|a\|^{-1}a + \|b\|^{-1}b - e \leq c \leq \|a\|^{-1}a, \|b\|^{-1}b.$$

This implies

$$0 \leq \min(\|a\|, \|b\|)c \leq a, b$$

and hence the orthogonality of a and b implies $c = 0$. Therefore

$$\|a\|^{-1}a + \|b\|^{-1}b - e \leq 0.$$

Thus for any state $\psi \in S(E)$ with $\psi(a) = \|a\|$ we have

$$0 \leq \|b\|^{-1}\psi(b) \leq \psi(e - \|a\|^{-1}a) = 0,$$

which implies $\psi(b) = 0$. □

Proof of Theorem 4.

(i) \implies (ii) is obvious.

(ii) \implies (i). This can be proved in the same way as in Proposition II.3.2 of [1] by noting Lemma 4 and that the set

$$\{w \in E \mid x, y \leq w \leq z\} = [x, z] \cap [y, z]$$

for any $x, y, z \in E$ is weakly* compact.

(ii) \implies (iii). Assume (ii) and take arbitrary 2-outcome measurements $(a_1, a_2) = (a_1, e - a_1)$ and $(b_1, b_2) = (b_1, e - b_1)$ on (E, e) . Then by $a_1 \leq e = b_1 + b_2$ and the Riesz decomposition property of E , there exist elements $c_{1,1}, c_{1,2} \in E$ such that

$$0 \leq c_{1,j} \leq b_j \quad (j = 1, 2), \quad a_1 = c_{1,1} + c_{1,2}.$$

If we put $c_{2,j} := b_j - c_{1,j} \geq 0$, we have

$$\begin{aligned} b_j &= c_{1,j} + c_{2,j} \quad (j = 1, 2), \\ a_2 &= e - a_1 = b_1 + b_2 - c_{1,1} - c_{1,2} = c_{2,1} + c_{2,2}. \end{aligned}$$

Thus (a_1, a_2) and (b_1, b_2) are compatible.

(iii) \implies (ii). We assume (iii) and prove the Riesz decomposition property of E . Take elements $u, v_1, v_2 \in E$ satisfying

$$0 \leq u \leq v_1 + v_2, \quad 0 \leq v_1, v_2.$$

Let

$$A := \{(x_1, x_2) \in E \times E \mid 0 \leq x_j \leq v_j \quad (j = 1, 2), \quad x_1 + x_2 \leq u\},$$

which is non-empty by $(0, 0) \in A$, and define a partial order \leq on A by

$$(x_1, x_2) \leq (y_1, y_2) \stackrel{\text{def.}}{\iff} x_1 \leq y_1 \text{ and } x_2 \leq y_2 \quad ((x_1, x_2), (y_1, y_2) \in A).$$

Let D be a directed subset of A and for each $\delta \in D$ we write as $\delta = (x_1^\delta, x_2^\delta)$. Then for each $j = 1, 2$ the net $(x_j^\delta)_{\delta \in D}$ in E is increasing and upper bounded by v_j . Thus by Lemma 1

$$x_j^\delta \xrightarrow{\text{weakly}^*} \sup_{\delta \in D} x_j^\delta =: x_j \in E.$$

By the weak* closedness of E_+ , we have $(x_1, x_2) \in A$ and hence (x_1, x_2) is a supremum of D in A . Therefore by Zorn's lemma there exists an element $(u_1, u_2) \in A$ maximal with respect to the order \leq on A .

We next prove that $u - u_1 - u_2 \geq 0$ and $v_1 - u_1 \geq 0$ are orthogonal. Take $c \in E$ such that

$$0 \leq c \leq u - u_1 - u_2, \quad v_1 - u_1.$$

Then this implies $(u_1, u_2) \leq (u_1 + c, u_2) \in A$ and hence by the maximality of (u_1, u_2) we have $c = 0$. Therefore $u - u_1 - u_2$ and $v_1 - u_1$ are orthogonal. Similar proof also applies to the orthogonality of $u - u_1 - u_2$ and $v_2 - u_2$.

Now assume $u - u_1 - u_2 \neq 0$. We then take a state $\psi \in S(E)$ such that

$$\psi(u - u_1 - u_2) = \|u - u_1 - u_2\| > 0.$$

Then by Lemma 6 we have $\psi(v_1 - u_1) = \psi(v_2 - u_2) = 0$ since $u - u_1 - u_2$ and $v_j - u_j$ are orthogonal ($j = 1, 2$). Hence, by noting $u \leq v_1 + v_2$, we have

$$0 < \psi(u - u_1 - u_2) = \psi(u - v_1 - v_2) \leq 0,$$

which is a contradiction. Therefore $u = u_1 + u_2$, which proves the Riesz decomposition property of E . \square

Proof of Theorem 3. The proof of (ii) \implies (iii) in Theorem 4 also applies to the “only if” part of the statement. To establish the “if” part, we assume that any pair of 2-outcome measurements on (E, e) is compatible and show that E^* is a lattice.

We first prove that any 2-outcome measurements on (E^{**}, e) is compatible. Since each 2-outcome measurements corresponds to an effect by

$$\mathcal{E}(E^{**}) \ni x'' \mapsto (x'', e - x''),$$

we have only to prove that any effects $a'', b'' \in \mathcal{E}(E^{**})$ are compatible. By Lemma 2 there exist nets $(a_i)_{i \in I}$ and $(b_i)_{i \in I}$ in $\mathcal{E}(E)$ such that $a_i \xrightarrow{\text{weakly}^*} a''$ and $b_i \xrightarrow{\text{weakly}^*} b''$. Since the effects a_i and b_i are compatible for each $i \in I$, there exists $c_i \in E$ such that

$$0, a_i + b_i - e \leq c_i \leq a_i, b_i.$$

Since c_i is bounded, the Banach-Alaoglu theorem implies that there exists a subnet of $(c_i)_{i \in I}$ weakly* converging to some $c'' \in E^{**}$. By the weak* closedness of the positive cone E_+^{**} , we have

$$0, a'' + b'' - e \leq c'' \leq a'', b'',$$

which implies the compatibility of the effects a'' and b'' .

Now by Theorem 4, E^{**} is a lattice and hence an AM-space and by Corollary 1 of [2] E^* is a AL-space and so is a lattice, which completes the proof. \square

References

1. Alfsen, E.M.: Compact Convex Sets and Boundary Integrals. Springer (1971)
2. Ellis, A.J.: The duality of partially ordered normed linear spaces. J. London Math. Soc. **39**(1), 730–744 (1964).
3. Heinosaari, T., Miyadera, T., Ziman, M.: An invitation to quantum incompatibility. J. Phys. A: Math. Theor. **49**(12), 123001 (2016).
4. Olubummo, Y., Cook, T.A.: The Predual of an Order-Unit Banach Space. Int. J. Theor. Phys. **38**(12), 3301–3303 (1999).
5. Plávala, M.: All measurements in a probabilistic theory are compatible if and only if the state space is a simplex. Phys. Rev. A **94**, 042108 (2016).