

A refined trans-Planckian censorship conjecture

Rong-Gen Cai*

*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Sciences, Beijing 100190, China*

Shao-Jiang Wang†

*Tufts Institute of Cosmology, Department of Physics and Astronomy,
Tufts University, 574 Boston Avenue, Medford, Massachusetts 02155, USA*

We propose a refined version of trans-Planckian censorship conjecture (TCC), which could be elaborated from the strong scalar weak gravity conjecture combined with some entropy bounds. In particular, no fine-tuning on the inflation model-building is required in the refined TCC, and it automatically passes the tests from those stringy examples that support the original TCC. Furthermore, our refined TCC could be consistent with hilltop eternal inflation.

I. INTRODUCTION

Decades of explorations along string theory as a promising candidate for quantum gravity reveal us a landscape [1] of a huge number of low-energy effective theories, beyond which are conjectured mostly in the swampland [2] that admit no consistent completion into quantum gravity in the ultraviolet (UV). The string-inspired swampland criteria (see, for example, a review [3]) have been proposed to discriminate those seemingly innocent theories consistent at low-energy in the infrared (IR).

The difficulties of locating de Sitter (dS) vacua in string theory have inspired the swampland dS conjecture (SdSC) [4] and its refinements [5–10]. The original SdSC [4] simply forbids any dS local extrema, which is in direct tension with the standard model Higgs potential [11–15] (see also [16]) that has a dS local maximum. Therefore, the refined SdSC [5–8, 10, 17] was proposed to relax the constraint so that only dS local minima is forbidden. In particular, both requirements of no quantum broken dS [5, 18] and no eternal inflation [7] disfavour any stable dS local minima. Similarly, the trans-Planckian censorship conjecture (TCC) [19] was recently proposed to further relax the constraint that, all modes that exit the Hubble horizon at the end of inflation should have their physical length larger than the Planck length at the beginning of inflation, namely, $a_i/(a_f H_f) > 1/M_{\text{Pl}}$. This strongly constrains the duration of inflation on the elapsed e-folding number,

$$e^N < \frac{M_{\text{Pl}}}{H_f}. \quad (1)$$

Therefore, TCC further relaxes the swampland dS criteria by forbidding any stable dS local minima but sparing the life of metastable dS local minima. As an immediate result, inflation cannot be eternal both into the past [20] and future directions [5, 7, 18, 21].

When applied to the inflationary phenomenology [22], TCC sets an upper bound on the inflationary Hubble scale and corresponding tensor-to-scalar ratio by

$$H < e^{-N} M_{\text{Pl}} \approx \mathcal{O}(10) \text{ MeV}, \quad (2)$$

$$r \equiv \frac{2}{\pi^2 \mathcal{P}_{\mathcal{R}}} \left(\frac{H}{M_{\text{Pl}}} \right)^2 < 6.8 \times 10^{-33}, \quad (3)$$

where $N = (1/3) \ln(M_{\text{Pl}}/H_0) \approx 46.2$ for an instantaneous reheating history [22, 23] and $\mathcal{P}_{\mathcal{R}} \approx 2.1 \times 10^{-9}$ from Planck 2018 [24]. This imposes a severe fine-tuning problem on the inflationary model-building with initial condition on slow-roll parameter down to $\epsilon \approx 10^{-33}$, which could be relaxed by invoking non-instantaneous reheating history [23] at the price of ultra-low reheating temperature (see [25] for its implications on the mass of primordial black holes). Other trials of model-building within TCC include inflationary dark matter [26], warm inflation [27, 28], initial states [29, 30], non-standard post-inflationary history [31, 32], D-term hybrid inflation [33], multi-stage inflation [34–36], inflection-point inflation [37], negative running of spectral index [38]. Beyond the original statement of TCC, [39] proposed a generalized TCC to save k -inflation [40] from the problem that sub-Planckian fluctuations could exit Hubble horizon without violating the original TCC. Furthermore, [41] casts the doubt on TCC as an universal swampland criterion but proposes a probabilistic interpretation.

Although TCC requires all trans-Planckian quantum modes to remain their quantum nature by never exiting the Hubble horizon, the apparent physical consequences of having trans-Planckian fluctuations stretched out of the Hubble horizon are unclear (see, however, [42][43][44] for self-completeness of Einstein gravity), which should be elaborated with other more established arguments. One inspiration comes from some earlier variants of refined SdSC [5, 7, 18] that also prohibit long-lasting dS vacua. Therefore, although the refined SdSC in [8] is silent about the dS lifetime, the arguments they used to obtain the refined SdSC could shed light on the physical interpretation of TCC. Recall that the refined SdSC [8] was found following from the swampland distance conjecture (SDC) [45] combined with the Bousso entropy bound

* cairg@itp.ac.cn

† schwang@cosmos.phy.tufts.edu

[46] applied to an accelerating universe, which could also be used in [21] to derive an entropic quasi-dS instability time. In particular, TCC was recently elaborated from SDC [47] in the large field excursion limit, which could also be derived from some entropy bounds [48]. To elaborate TCC over the whole field space, we use a strong version [49] of scalar weak gravity conjecture (WGC) [50] combined with some entropy bounds, and then a weaker version of TCC is obtained to naturally evade the fine-tuning problem of initial conditions without turning to any exotic model-building.

The outline for this paper is as follows: In Sec. II, we review the scalar WGC and then derive a bound for the species number from both scalar WGC and its strong version. In Sec. III, using previous bound on species number, we propose a refined TCC from two separate arguments of UV-IR hierarchy and entropy bounds. In Sec. IV, the refined TCC is reformulated for eternal inflation. In Sec. V, we conclude the result and discuss the refined TCC implications for inflation without fine-tuning problem.

II. SCALAR WEAK GRAVITY CONJECTURE

The scalar WGC [50] implements the belief that the net force mediated by scalar fields should be larger than the gravitational force. In specific, one considers a complex scalar field φ with a field-dependent mass term $m^2(\phi)|\varphi|^2$, where a canonically normalized scalar field ϕ mediates the scalar force between φ particles via a trilinear coupling term $\partial_\phi m^2(\phi_0)\delta\phi|\varphi|^2$ when expanding $\phi = \phi_0 + \delta\phi$ around a vacuum expectation value (vev) ϕ_0 . By requiring the scalar force larger than the gravity,

$$F_{\text{scalar}} \equiv \frac{(\partial_\phi m)^2}{4\pi r^2} > \frac{m^2}{8\pi M_{\text{Pl}}^2 r^2} \equiv F_{\text{gravity}}, \quad (4)$$

the scalar WGC reads

$$2(\partial_\phi m)^2 > \frac{m^2}{M_{\text{Pl}}^2}, \quad (5)$$

Similar form also holds for fermion or massless multi-scalar fields ϕ^i with kinetic term $g_{ij}\partial\phi^i\partial\phi^j$, that is, there should have a state with mass m satisfying the bound

$$2g^{ij}(\partial_{\phi^i} m)(\partial_{\phi^j} m) > \frac{m^2}{M_{\text{Pl}}^2}. \quad (6)$$

In particular, as an explicit fulfillment of (5), SDC [45] allows for a large distance replacement (corresponding to weak coupling limit) over the moduli space without a potential (or field space in the effective theory with a potential [51, 52]) if a tower of light states with mass

$$m(\phi) \sim m(\phi_0)e^{-\alpha\frac{\Delta\phi}{M_{\text{Pl}}}} \quad (7)$$

descends from the UV with some $\mathcal{O}(1)$ constant $\alpha > 0$. Note that the scalar WGC is formulated for any field

value in field space, while the SDC holds most straightforwardly at large distances in field space. Therefore, our use of scalar WGC in replacement of SDC is a non-trivial generalization of the argument used in [21]. Nevertheless, we all arrive at the same dS lifetime as shown in the end.

A. A bound for species number

Since both scalar WGC and SDC could constrain the mass scale of an infinite tower of states for a scalar theory coupled to gravity, the effective description of the scalar theory might be jeopardized by including sufficient number of states. Therefore, there exist a bound on the number of states included, above which we are not in any weakly-coupled gravitational regime but a strong coupling regime, or equivalently, for given N_s particle states, gravity becomes strongly coupled above a dubbed species scale (conjecture) [53–60],

$$\Lambda_s \equiv \frac{M_{\text{Pl}}}{\sqrt{N_s}}. \quad (8)$$

Recall that, for example, the number of Kaluza-Klein (KK) modes N_s we can include in the lower d dimensional gravity theory before the true quantum gravity cutoff scale M_{Pl}^D of the higher D dimensional gravity theory is $N_s \sim M_{\text{Pl}}^D/m_{\text{KK}}$. Likewise, $N_s \sim \Lambda_s/m$ is expected in general (see, for example, [3] around Eq. (5.17)). Hence the species number reads

$$N_s \sim \left(\frac{M_{\text{Pl}}}{m(\phi)}\right)^{\frac{2}{3}}, \quad (9)$$

which, after the use of scalar WGC (5), becomes

$$(\partial_\phi \ln N_s)^2 \gtrsim \frac{2}{9}M_{\text{Pl}}^{-2}. \quad (10)$$

This could be integrated directly to give

$$\ln N_s \gtrsim \frac{\sqrt{2}}{3} \frac{|\Delta\phi|}{M_{\text{Pl}}}. \quad (11)$$

On the other hand, if ϕ also derives the background expansion with Hubble rate H , one could define the corresponding Hubble slow-roll parameter as

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2} = \frac{\dot{\phi}^2}{2M_{\text{Pl}}^2 H^2}, \quad (12)$$

where we use the Friedmann equation $M_{\text{Pl}}^2 \dot{H} = -\frac{1}{2}\dot{\phi}^2$ in the last step. The e-folding number during given field excursion in the non-eternal inflationary regime is well-known,

$$\begin{aligned} |\Delta N| &= \int_{t_i}^{t_f} H dt = \int_{\phi_i}^{\phi_f} \frac{H}{\dot{\phi}} d\phi = \int_{\phi_i}^{\phi_f} \frac{1}{\sqrt{2\epsilon_H}} \frac{|d\phi|}{M_{\text{Pl}}} \\ &\equiv \left\langle \frac{1}{\sqrt{\epsilon_H}} \right\rangle \frac{|\Delta\phi|}{\sqrt{2}M_{\text{Pl}}}, \end{aligned} \quad (13)$$

where in the second line we define the averaged value for the inverse square-root of Hubble slow-roll parameter. Therefore, Eq. (11) becomes

$$N_s > e^{\frac{2}{3}|\Delta N|/\langle\epsilon_H^{-1/2}\rangle} \equiv e^{\frac{2}{3}N_\epsilon}, \quad (14)$$

where the abbreviation $N_\epsilon \equiv |\Delta N|/\langle\epsilon_H^{-1/2}\rangle$ is introduced for convenience. The physical meaning of bound (14) is illuminating: for given inflationary potential in an effective theory valid below Λ_s , the e-folding number is bounded from above by (14), otherwise larger e-folding number leads to larger field excursion, and hence larger number of light particle states would be included to spoil the effectiveness of original theory unless lowering Λ_s accordingly.

B. A strong scalar weak gravity conjecture

However, there are some concerns [49] on the scalar WGC, for example, it does not apply to all scalar fields (e.g. the massless mediator ϕ itself), and it does not simultaneously accommodate SDC for both $\phi \rightarrow \pm\infty$. Furthermore, it is not consistent with axion-like particles and fifth-force constraints, and it might be in phenomenological tension [61] with the (refined) swampland dS conjecture [4–10]. As a result, a stronger version of scalar WGC was proposed in [49] for any canonically normalized real scalar ϕ coupled to gravity, the scalar potential $V(\phi)$ at any field value should meet

$$2(V''')^2 - V''V'''' \geq \frac{(V'')^2}{M_{\text{Pl}}^2}, \quad (15)$$

which, after writing $m^2 = V''$, becomes

$$2(\partial_\phi m^2)^2 - m^2(\partial_\phi^2 m^2) \geq \frac{(m^2)^2}{M_{\text{Pl}}^2}. \quad (16)$$

The above bound possesses clear physical meaning [62] that the first and second terms on the left-hand-side denote the attractive and repulsive scalar forces, respectively. Therefore, this strong version of scalar WGC simply states that the net scalar force should surpass the gravity force over the whole field space.

To see whether the bound (14) is still held in the strong version of scalar WGC, we could insert (9) into (16) and found

$$9(\partial_\phi \ln N_s)^2 + 3\partial_\phi^2 \ln N_s \geq M_{\text{Pl}}^{-2}, \quad (17)$$

which could be directly solved as

$$N_s \geq \cosh^{\frac{1}{3}} \frac{|\Delta\phi|}{M_{\text{Pl}}} > e^{\frac{\sqrt{2}}{3}N_\epsilon}. \quad (18)$$

Apart from a $\mathcal{O}(1)$ factor difference in the exponent, this is the same as Eq. (14), which could also be derived from applying SDC on (9) in specific form of $m(\phi)$ with $m(\phi_0) \sim M_{\text{Pl}}$. However, our derivation of (18) is quite generic in the sense that it follows for any scalar field at any field value, and does not rely on the slow-roll approximation.

III. REFINED TRANS-PLANCKIAN CENSORSHIP CONJECTURE

Armed with Eq. (18), a refined TCC could be targeted with following two separate arguments:

A. UV-IR hierarchy

The Hubble horizon as an IR cutoff scale should be bounded by the UV cutoff scale set by the species scale [47],

$$H < \Lambda_s \equiv \frac{M_{\text{Pl}}}{\sqrt{N_s}} < M_{\text{Pl}} e^{-\frac{\sqrt{2}}{6}N_\epsilon}, \quad (19)$$

with Eq. (18) used in the last step, from which a refined TCC could be directly read off,

$$e^{\frac{\sqrt{2}}{6}N_\epsilon} < \frac{M_{\text{Pl}}}{H}. \quad (20)$$

The original TCC (1) serves as a stronger statement than (20) since $N_\epsilon \ll |\Delta N|$ for slow-roll inflation with $\epsilon_H \ll 1$. Therefore, all those stingy examples that support original TCC also automatically hold for the refined TCC. The physical consequence of refined TCC could be understood in the following way: for given potential with expansion history $\langle\epsilon_H^{-1/2}\rangle$ fixed, the violation occurs for large enough $|\Delta N|$, which leads to large field excursion $|\Delta\phi|$. Hence the species number N_s would be so large that the UV scale set by species scale Λ_s could be even smaller than the IR scale set by Hubble horizon H , which is not allowed by the hierarchy of scales.

B. Entropy bounds

Consider a spherical region of size R consisting of N_s species number of particles as radiations at temperature T , the total energy and entropy scale as

$$E \sim N_s R^3 T^4, \quad S \sim N_s R^3 T^3. \quad (21)$$

The absence of black hole formation imposes $R > R_{\text{BH}} \sim E/M_{\text{Pl}}^2 \sim N_s R^3 T^4/M_{\text{Pl}}^2$, which leads to a maximum radius $R < M_{\text{Pl}}/(\sqrt{N_s}T^2)$, or equivalently a maximum entropy $S \sim N_s R^3 T^3 < M_{\text{Pl}}^3/(\sqrt{N_s}T^3)$. This maximum entropy should be bounded by the Bekenstein-Hawking entropy, $S_{\text{BH}} \sim M_{\text{Pl}}^2 R^2 < M_{\text{Pl}}^4/(N_s T^4)$, as

$$\frac{M_{\text{Pl}}^3}{\sqrt{N_s}T^3} < \frac{M_{\text{Pl}}^4}{N_s T^4}, \quad (22)$$

which in turn gives rise to a maximum temperature $T < M_{\text{Pl}}/\sqrt{N_s}$ [59, 63, 64] larger than the Hubble temperature,

$$\frac{H}{2\pi} < \frac{M_{\text{Pl}}}{\sqrt{N_s}} < M_{\text{Pl}} e^{-\frac{\sqrt{2}}{6}N_\epsilon}, \quad (23)$$

namely,

$$e^{\frac{\sqrt{2}}{6}N_\epsilon} < \frac{M_{\text{Pl}}}{H}. \quad (24)$$

Similar bound could also be obtained if the radiation entropy is bounded by the Hubble entropy [65, 66],

$$N_s R^3 T^3 < M_{\text{Pl}}^2 R^3 H. \quad (25)$$

Hence there is also a maximum temperature, $T < (M_{\text{Pl}}^2 H/N_s)^{1/3}$, which should be larger than the Hubble temperature,

$$H^3 < \frac{M_{\text{Pl}}^2 H}{\sqrt{N_s}} < M_{\text{Pl}}^2 H e^{-\frac{\sqrt{2}}{6}N_\epsilon}, \quad (26)$$

namely,

$$e^{\frac{\sqrt{2}}{12}N_\epsilon} < \frac{M_{\text{Pl}}}{H}. \quad (27)$$

Note that similar arguments of using entropy bounds are adopted in [48] to elaborate SDC.

The above arguments implicitly assume that all N_s species number of particles have their masses spectrum below the temperature. However, it is fairly possible that some of the particle masses are heavier than the temperature so that they have contribution to the total energy $E \sim k_B N_s T$ ($k_B \equiv 1$ hereafter) but no contribution to the total entropy, which should be bounded by the Bekenstein entropy [67–69] for any weakly gravitating matter fields,

$$S_{\text{matter}} \leq 2\pi ER \sim 2\pi N_s TR. \quad (28)$$

If the matter inside the sphere is gravitationally stable so that black hole does not form, the Bekenstein entropy should be further bounded by the Bekenstein-Hawking entropy,

$$2\pi N_s TR \leq M_{\text{Pl}}^2 R^2. \quad (29)$$

Therefore, there is a maximum temperature $T < M_{\text{Pl}}^2 R/(2\pi N_s)$, which should be larger than the Hubble temperature,

$$\frac{H}{2\pi} < \frac{M_{\text{Pl}}^2 R}{2\pi N_s} < \frac{M_{\text{Pl}}^2}{2\pi H} e^{-\frac{\sqrt{2}}{3}N_\epsilon}, \quad (30)$$

namely,

$$e^{\frac{\sqrt{2}}{6}N_\epsilon} < \frac{M_{\text{Pl}}}{H}. \quad (31)$$

The same bound could also be obtained by bounding the Bekenstein entropy with Hubble entropy ,

$$2\pi N_s TR \leq M_{\text{Pl}}^2 R^3 H. \quad (32)$$

Hence there is also a maximum temperature $T < M_{\text{Pl}}^2 HR^2/(2\pi N_s)$, which should be larger than the Hubble temperature,

$$\frac{H}{2\pi} < \frac{M_{\text{Pl}}^2 HR^2}{2\pi N_s} < \frac{M_{\text{Pl}}^2}{2\pi H} e^{-\frac{\sqrt{2}}{3}N_\epsilon}, \quad (33)$$

namely,

$$e^{\frac{\sqrt{2}}{6}N_\epsilon} < \frac{M_{\text{Pl}}}{H}. \quad (34)$$

Note that the entropy bound alone is not sufficient to guarantee the refined TCC, since the UV-IR hierarchies $H < T$ and $R < H^{-1}$ are still needed for the relevant derivation of inequalities.

In summary, with Eq. (18), the refined TCC is always implied by entropy bounds either from radiation entropy (21) or Bekenstein entropy (28), which are bounded by either Bekenstein-Hawking entropy (22)(29) or Hubble entropy (25)(32).

IV. ETERNAL INFLATION

There are three types of eternal inflation frequently discussed in the literatures: old eternal inflation (a trapped dS vacuum) [70, 71], stochastic eternal inflation [72–74] (a hilltop potential [75]), and topological eternal inflation [76–78] (an eternally inflating topological defeat from a hilltop potential [79]). Whether the eternal inflation is in the swampland is an important question currently under debate [7, 80–86]. The old eternal inflation is generically inconsistent with the original SdSC [80, 81], but the stochastic eternal inflation was argued to be marginally consistent with the refined SdSC [82], which, however, was questioned with perturbative [83] and entropic [84] arguments. Nevertheless, [7] speculated a possibility that some variants of SdSC should be regarded as approximate consequences of the absence of eternal inflation. Furthermore, some topological eternal inflation models could be constructed to evade the swampland criteria for an eternally inflating bubble wall [85] and domain wall on the brane [86].

Our refined TCC (20) is elaborated from the arguments of Eq. (11), (13) and (19). Note that Eq. (13) is restricted to the non-eternal inflationary regime where the field excursion is dominated by the classical rolling. For eternal inflation, the elapsed e-folding number could be arbitrarily large as long as the field excursion is limited in the range where the quantum fluctuations dominate over classical rolling. In this case, one could use (11) and (19) directly without (13) to render the general version of our refined TCC,

$$e^{\frac{\sqrt{2}}{6} \frac{|\Delta\phi|}{M_{\text{Pl}}}} < \frac{M_{\text{Pl}}}{H}, \quad (35)$$

which shares a similar form derived in [87] from SDC on inflation. To see whether the stochastic eternal inflation could survive above criterion, we use a quadratic hilltop potential

$$V(\phi) = V_0 - \frac{1}{2}m^2\phi^2, \quad m^2 > 0, \quad (36)$$

where the eternal inflation is allowed in the small field region $|\Delta\phi| < \phi_c$ when the averaged quantum random




FIG. 1. The parameter space for a quadratic hilltop eternal inflation out of swampland by our refined TCC enclosed by the red and blue solid lines. The slow-roll approximation is also checked with the blue and red dashed lines.

walks dominate over classical rolling during each Hubble time, $H/(2\pi) \equiv \langle \delta\phi \rangle_Q > \delta\phi_C \equiv |\phi|/H$. ϕ_c is therefore solved from

$$\mathcal{P}_\zeta(\phi_c) \equiv \left(\frac{H^2(\phi_c)}{2\pi\dot{\phi}_c} \right)^2 \approx \frac{V(\phi_c)}{24\pi^2 M_{\text{Pl}}^4 \epsilon_V(\phi_c)} = 1. \quad (37)$$

Our refined TCC (35) is ensured provided that

$$e^{\frac{\sqrt{2}}{6} \frac{\phi_c}{M_{\text{Pl}}}} < \frac{M_{\text{Pl}}}{H(\phi_c)}, \quad (38)$$

which could be solved numerically as shown in Fig. 1 with blue solid line. Furthermore, (38) automatically guarantees the slow-roll condition for ϵ_V (blue dashed lines),

$$\epsilon_V(\phi_c) < \frac{1}{8\pi^2} e^{-\frac{\sqrt{2}}{3} \frac{\phi_c}{M_{\text{Pl}}}} < 1. \quad (39)$$

The slow-roll condition $|\eta_V(\phi_c)| \ll 1$ for $\eta_V(\phi_c)$ (red dashed lines) also automatically guarantees $\eta_V > -3$ [7] so that the exponentially suppressed probability to stay within $|\Delta\phi| < \phi_c$ beats the exponential expansion of the universe. On the other hand, the perturbativity argument [83] further requires

$$\mathcal{P}_\zeta < \frac{1}{16[3(\eta_V - \epsilon_V) - \epsilon_V(\eta_V - \epsilon_V) - \epsilon_V\eta_V]^2}, \quad (40)$$

which is shown in Fig. 1 with red solid line. Therefore, there is a parameter space for the quadratic hilltop eternal inflation out of swampland by our refined TCC (35).

V. CONCLUSION AND DISCUSSIONS

In this paper, we proposed a refined TCC as a consequence of a strong version of scalar WGC combined with either UV-IR hierarchy or some entropy bounds,

$$e^{\frac{|\Delta\phi|}{M_{\text{Pl}}}} / \mathcal{O}(1) < \mathcal{O}(1) \frac{M_{\text{Pl}}}{H}, \quad (41)$$

which is consistent with the hilltop eternal inflation. For non-eternal inflation, it further reduces to

$$e^{N_\epsilon / \mathcal{O}(1)} < \mathcal{O}(1) \frac{M_{\text{Pl}}}{H_f}. \quad (42)$$

where the $\mathcal{O}(1)$ factor on the left-hand-side is associated to certain combination of spacetime dimension, which could be easily worked out from previous arguments, and the $\mathcal{O}(1)$ factor on the right-hand-side is already mentioned in a footnote in [19], which could be the sound speed in the generalized TCC in [39]. The presence of an averaged Hubble slow-roll parameter in the exponent $N_\epsilon \equiv N / \langle \epsilon_H^{-1/2} \rangle$ would prolong the dS lifetime by

$$\Delta t_{\text{dS}} < \frac{1}{H_f} \int_{t_i}^{t_f} H dt = \frac{1}{H_f} \ln \frac{a_f}{a_i} < \frac{\langle \epsilon_H^{-1/2} \rangle}{H_f} \ln \frac{M_{\text{Pl}}}{H_f}, \quad (43)$$

which resembles the form of the entropic quasi-de Sitter instability time [21] (such a logarithmic correction to the Hubble time also appears in the scrambling time [19, 88]). Now the constraint on the Hubble scale could be greatly relaxed, for instance, $N_\epsilon \equiv N / \langle \epsilon_H^{-1/2} \rangle \approx N/10 = 6$ for an illustrative $\epsilon_H \approx 0.01$ and e-folding number $N = 60$, to

$$H < e^{-N_\epsilon} M_{\text{Pl}} \approx 6 \times 10^{15} \text{ GeV}. \quad (44)$$

The same relaxation could also be achieved even for a near critical expansion with $\epsilon_H \approx 1$ due to the presence of a $\mathcal{O}(1)$ factor on the left-hand-side of (42), for example, as shown in (20),

$$H < e^{-\frac{\sqrt{2}}{6} N} M_{\text{Pl}} \approx 2 \times 10^{12} \text{ GeV}, \quad (45)$$

for the same illustrative $N = 60$. Therefore, our refined TCC resolves the fine-tuning problem of inflationary initial conditions without invoking any exotic model-building. Furthermore, since our refined TCC is a weaker version of original TCC, it would automatically be supported by those stringy examples that also hold for the original TCC.

ACKNOWLEDGMENTS

We thank Suddhasattwa Brahma, Mark Hertzberg, Ken Olum, Eran Palti, Fabrizio Rompineve, Shan-Ming Ruan, Alexander Vilenkin, and Masaki Yamada for helpful discussions and correspondences. R.G.C. was

supported by the National Natural Science Foundation of China Grants No. 11435006, No. 11647601, No. 11690022, No. 11821505 and No. 11851302, and by the Strategic Priority Research Program of CAS Grant

No. XDB23030100, and by the Key Research Program of Frontier Sciences of CAS. SJW is supported by the postdoctoral scholarship of Tufts University from NSF.

-
- [1] L. Susskind, “The Anthropic landscape of string theory,” [arXiv:hep-th/0302219](#) [[hep-th](#)].
- [2] C. Vafa, “The String landscape and the swampland,” [arXiv:hep-th/0509212](#) [[hep-th](#)].
- [3] E. Palti, “The Swampland: Introduction and Review,” *Fortsch. Phys.* **67** no. 6, (2019) 1900037, [arXiv:1903.06239](#) [[hep-th](#)].
- [4] G. Obied, H. Ooguri, L. Spodyneiko, and C. Vafa, “De Sitter Space and the Swampland,” [arXiv:1806.08362](#) [[hep-th](#)].
- [5] G. Dvali and C. Gomez, “On Exclusion of Positive Cosmological Constant,” *Fortsch. Phys.* **67** no. 1-2, (2019) 1800092, [arXiv:1806.10877](#) [[hep-th](#)].
- [6] D. Andriot, “On the de Sitter swampland criterion,” *Phys. Lett.* **B785** (2018) 570–573, [arXiv:1806.10999](#) [[hep-th](#)].
- [7] T. Rudelius, “Conditions for (No) Eternal Inflation,” *JCAP* **1908** (2019) 009, [arXiv:1905.05198](#) [[hep-th](#)].
- [8] H. Ooguri, E. Palti, G. Shiu, and C. Vafa, “Distance and de Sitter Conjectures on the Swampland,” [arXiv:1810.05506](#) [[hep-th](#)].
- [9] S. K. Garg and C. Krishnan, “Bounds on Slow Roll and the de Sitter Swampland,” *JHEP* **11** (2019) 075, [arXiv:1807.05193](#) [[hep-th](#)].
- [10] D. Andriot and C. Roupec, “Further refining the de Sitter swampland conjecture,” *Fortsch. Phys.* **67** no. 1-2, (2019) 1800105, [arXiv:1811.08889](#) [[hep-th](#)].
- [11] F. Denef, A. Hebecker, and T. Wrase, “de Sitter swampland conjecture and the Higgs potential,” *Phys. Rev.* **D98** no. 8, (2018) 086004, [arXiv:1807.06581](#) [[hep-th](#)].
- [12] M. Cicoli, S. De Alwis, A. Maharana, F. Muia, and F. Quevedo, “De Sitter vs Quintessence in String Theory,” *Fortsch. Phys.* **67** no. 1-2, (2019) 1800079, [arXiv:1808.08967](#) [[hep-th](#)].
- [13] H. Murayama, M. Yamazaki, and T. T. Yanagida, “Do We Live in the Swampland?,” [arXiv:1809.00478](#) [[hep-th](#)].
- [14] K. Choi, D. Chway, and C. S. Shin, “The dS swampland conjecture with the electroweak symmetry and QCD chiral symmetry breaking,” *JHEP* **11** (2018) 142, [arXiv:1809.01475](#) [[hep-th](#)].
- [15] K. Hamaguchi, M. Ibe, and T. Moroi, “The swampland conjecture and the Higgs expectation value,” *JHEP* **12** (2018) 023, [arXiv:1810.02095](#) [[hep-th](#)].
- [16] C. Han, S. Pi, and M. Sasaki, “Quintessence Saves Higgs Instability,” [arXiv:1809.05507](#) [[hep-ph](#)].
- [17] S. K. Garg, C. Krishnan, and M. Zaid Zaz, “Bounds on Slow Roll at the Boundary of the Landscape,” *JHEP* **03** (2019) 029, [arXiv:1810.09406](#) [[hep-th](#)].
- [18] G. Dvali, C. Gomez, and S. Zell, “Quantum Breaking Bound on de Sitter and Swampland,” *Fortsch. Phys.* **67** no. 1-2, (2019) 1800094, [arXiv:1810.11002](#) [[hep-th](#)].
- [19] A. Bedroya and C. Vafa, “Trans-Planckian Censorship and the Swampland,” [arXiv:1909.11063](#) [[hep-th](#)].
- [20] A. Borde, A. H. Guth, and A. Vilenkin, “Inflationary space-times are incomplete in past directions,” *Phys. Rev. Lett.* **90** (2003) 151301, [arXiv:gr-qc/0110012](#) [[gr-qc](#)].
- [21] M.-S. Seo, “The entropic quasi-de Sitter instability time from the distance conjecture,” [arXiv:1911.06441](#) [[hep-th](#)].
- [22] A. Bedroya, R. Brandenberger, M. Loverde, and C. Vafa, “Trans-Planckian Censorship and Inflationary Cosmology,” [arXiv:1909.11106](#) [[hep-th](#)].
- [23] S. Mizuno, S. Mukohyama, S. Pi, and Y.-L. Zhang, “Universal Upper Bound on the Inflationary Energy Scale from the Trans-Planckian Censorship Conjecture,” [arXiv:1910.02979](#) [[astro-ph.CO](#)].
- [24] **Planck** Collaboration, N. Aghanim et al., “Planck 2018 results. VI. Cosmological parameters,” [arXiv:1807.06209](#) [[astro-ph.CO](#)].
- [25] R.-G. Cai and S.-J. Wang, “Mass bound for primordial black hole from trans-Planckian censorship conjecture,” [arXiv:1910.07981](#) [[astro-ph.CO](#)].
- [26] T. Tenkanen, “Trans-Planckian Censorship, Inflation and Dark Matter,” [arXiv:1910.00521](#) [[astro-ph.CO](#)].
- [27] S. Das, “Distance, de Sitter and Trans-Planckian Censorship conjectures: the status quo of Warm Inflation,” [arXiv:1910.02147](#) [[hep-th](#)].
- [28] G. Goswami and C. Krishnan, “Swampland, Axions and Minimal Warm Inflation,” [arXiv:1911.00323](#) [[hep-th](#)].
- [29] Y. Cai and Y.-S. Piao, “Pre-inflation and Trans-Planckian Censorship,” [arXiv:1909.12719](#) [[gr-qc](#)].
- [30] S. Brahma, “Trans-Planckian censorship, inflation and excited initial states for perturbations,” [arXiv:1910.04741](#) [[hep-th](#)].
- [31] M. Dhuria and G. Goswami, “Trans-Planckian Censorship Conjecture and Non-thermal post-inflationary history,” [arXiv:1910.06233](#) [[astro-ph.CO](#)].
- [32] M. Torabian, “Non-Standard Cosmological Models and the trans-Planckian Censorship Conjecture,” [arXiv:1910.06867](#) [[hep-th](#)].
- [33] K. Schmitz, “Trans-Planckian Censorship and Inflation in Grand Unified Theories,” [arXiv:1910.08837](#) [[hep-ph](#)].
- [34] A. Berera and J. R. Calderón, “Trans-Planckian censorship and other swampland bothers addressed in warm inflation,” [arXiv:1910.10516](#) [[hep-ph](#)].
- [35] H.-H. Li, G. Ye, Y. Cai, and Y.-S. Piao, “Trans-Planckian censorship of multi-stage inflation and dark energy,” [arXiv:1911.06148](#) [[gr-qc](#)].
- [36] M. Torabian, “Breathing Comoving Hubble: Initial Condition and Eternity in view of the trans-Planckian Censorship Conjecture,” [arXiv:1911.12304](#) [[hep-th](#)].
- [37] N. Okada, D. Raut, and Q. Shafi, “Inflection-Point Inflation with Axion Dark Matter in light of

- Trans-Planckian Censorship Conjecture,” [arXiv:1910.14586 \[hep-ph\]](#).
- [38] K. Kadota, C. S. Shin, T. Terada, and G. Tumurtushaa, “Trans-Planckian censorship and single-field inflaton potential,” [arXiv:1910.09460 \[hep-th\]](#).
- [39] W.-C. Lin and W. H. Kinney, “Trans-Planckian Censorship and k -inflation,” [arXiv:1911.03736 \[gr-qc\]](#).
- [40] C. Armendariz-Picon, T. Damour, and V. F. Mukhanov, “ k - inflation,” *Phys. Lett.* **B458** (1999) 209–218, [arXiv:hep-th/9904075 \[hep-th\]](#).
- [41] R. Saito, S. Shirai, and M. Yamazaki, “Is Trans-Planckian Censorship a Swampland Conjecture?,” [arXiv:1911.10445 \[hep-th\]](#).
- [42] G. Dvali and C. Gomez, “Self-Completeness of Einstein Gravity,” [arXiv:1005.3497 \[hep-th\]](#).
- [43] G. Dvali, G. F. Giudice, C. Gomez, and A. Kehagias, “UV-Completion by Classicalization,” *JHEP* **08** (2011) 108, [arXiv:1010.1415 \[hep-ph\]](#).
- [44] G. Dvali, C. Gomez, R. S. Isermann, D. Lüst, and S. Stieberger, “Black hole formation and classicalization in ultra-Planckian $2 \rightarrow N$ scattering,” *Nucl. Phys.* **B893** (2015) 187–235, [arXiv:1409.7405 \[hep-th\]](#).
- [45] H. Ooguri and C. Vafa, “On the Geometry of the String Landscape and the Swampland,” *Nucl. Phys.* **B766** (2007) 21–33, [arXiv:hep-th/0605264 \[hep-th\]](#).
- [46] R. Bousso, “A Covariant entropy conjecture,” *JHEP* **07** (1999) 004, [arXiv:hep-th/9905177 \[hep-th\]](#).
- [47] S. Brahma, “The trans-Planckian censorship conjecture from the swampland distance conjecture,” [arXiv:1910.12352 \[hep-th\]](#).
- [48] A. Kehagias and A. Riotto, “A Note on the Trans-Planckian Censorship Conjecture,” [arXiv:1911.09050 \[hep-th\]](#).
- [49] E. Gonzalo and L. E. Ibáñez, “A Strong Scalar Weak Gravity Conjecture and Some Implications,” *JHEP* **08** (2019) 118, [arXiv:1903.08878 \[hep-th\]](#).
- [50] E. Palti, “The Weak Gravity Conjecture and Scalar Fields,” *JHEP* **08** (2017) 034, [arXiv:1705.04328 \[hep-th\]](#).
- [51] D. Klaeuer and E. Palti, “Super-Planckian Spatial Field Variations and Quantum Gravity,” *JHEP* **01** (2017) 088, [arXiv:1610.00010 \[hep-th\]](#).
- [52] F. Baume and E. Palti, “Backreacted Axion Field Ranges in String Theory,” *JHEP* **08** (2016) 043, [arXiv:1602.06517 \[hep-th\]](#).
- [53] G. Veneziano, “Large N bounds on, and compositeness limit of, gauge and gravitational interactions,” *JHEP* **06** (2002) 051, [arXiv:hep-th/0110129 \[hep-th\]](#).
- [54] G. R. Dvali, G. Gabadadze, M. Kolanovic, and F. Nitti, “Scales of gravity,” *Phys. Rev.* **D65** (2002) 024031, [arXiv:hep-th/0106058 \[hep-th\]](#).
- [55] N. Arkani-Hamed, S. Dimopoulos, and S. Kachru, “Predictive landscapes and new physics at a TeV,” [arXiv:hep-th/0501082 \[hep-th\]](#).
- [56] J. Distler and U. Varadarajan, “Random polynomials and the friendly landscape,” [arXiv:hep-th/0507090 \[hep-th\]](#).
- [57] S. Dimopoulos, S. Kachru, J. McGreevy, and J. G. Wacker, “N-flation,” *JCAP* **0808** (2008) 003, [arXiv:hep-th/0507205 \[hep-th\]](#).
- [58] G. Dvali, “Black Holes and Large N Species Solution to the Hierarchy Problem,” *Fortsch. Phys.* **58** (2010) 528–536, [arXiv:0706.2050 \[hep-th\]](#).
- [59] G. Dvali and M. Redi, “Black Hole Bound on the Number of Species and Quantum Gravity at LHC,” *Phys. Rev.* **D77** (2008) 045027, [arXiv:0710.4344 \[hep-th\]](#).
- [60] G. Dvali and D. Lust, “Evaporation of Microscopic Black Holes in String Theory and the Bound on Species,” *Fortsch. Phys.* **58** (2010) 505–527, [arXiv:0912.3167 \[hep-th\]](#).
- [61] S. Shirai and M. Yamazaki, “Is Gravity the Weakest Force?,” [arXiv:1904.10577 \[hep-th\]](#).
- [62] A. Kusenko, V. Takhistov, M. Yamada, and M. Yamazaki, “Fundamental Forces and Scalar Field Dynamics in the Early Universe,” [arXiv:1908.10930 \[hep-th\]](#).
- [63] R. Brustein and G. Veneziano, “A Causal entropy bound,” *Phys. Rev. Lett.* **84** (2000) 5695–5698, [arXiv:hep-th/9912055 \[hep-th\]](#).
- [64] R. Brustein, S. Foffa, and G. Veneziano, “CFT, holography, and causal entropy bound,” *Phys. Lett.* **B507** (2001) 270–276, [arXiv:hep-th/0101083 \[hep-th\]](#).
- [65] G. Veneziano, “Entropy bounds and string cosmology,” 1999. [arXiv:hep-th/9907012 \[hep-th\]](#). <http://weblib.cern.ch/abstract?CERN-TH-99-176>.
- [66] R. Brustein, “Cosmological entropy bounds,” *Lect. Notes Phys.* **737** (2008) 619–659, [arXiv:hep-th/0702108 \[hep-th\]](#).
- [67] J. D. Bekenstein, “Black holes and the second law,” *Lett. Nuovo Cim.* **4** (1972) 737–740.
- [68] J. D. Bekenstein, “Black holes and entropy,” *Phys. Rev.* **D7** (1973) 2333–2346.
- [69] J. D. Bekenstein, “Generalized second law of thermodynamics in black hole physics,” *Phys. Rev.* **D9** (1974) 3292–3300.
- [70] A. H. Guth, “The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems,” *Phys. Rev.* **D23** (1981) 347–356.
- [71] A. H. Guth and E. J. Weinberg, “Could the Universe Have Recovered from a Slow First Order Phase Transition?,” *Nucl. Phys.* **B212** (1983) 321–364.
- [72] A. A. Starobinsky, “STOCHASTIC DE SITTER (INFLATIONARY) STAGE IN THE EARLY UNIVERSE,” *Lect. Notes Phys.* **246** (1986) 107–126.
- [73] A. D. Linde, “ETERNAL CHAOTIC INFLATION,” *Mod. Phys. Lett.* **A1** (1986) 81.
- [74] M. Aryal and A. Vilenkin, “The Fractal Dimension of Inflationary Universe,” *Phys. Lett.* **B199** (1987) 351–357.
- [75] G. Barenboim, W.-I. Park, and W. H. Kinney, “Eternal Hilltop Inflation,” *JCAP* **1605** no. 05, (2016) 030, [arXiv:1601.08140 \[astro-ph.CO\]](#).
- [76] A. Vilenkin, “Topological inflation,” *Phys. Rev. Lett.* **72** (1994) 3137–3140, [arXiv:hep-th/9402085 \[hep-th\]](#).
- [77] A. D. Linde, “Monopoles as big as a universe,” *Phys. Lett.* **B327** (1994) 208–213, [arXiv:astro-ph/9402031 \[astro-ph\]](#).
- [78] A. D. Linde and D. A. Linde, “Topological defects as seeds for eternal inflation,” *Phys. Rev.* **D50** (1994) 2456–2468, [arXiv:hep-th/9402115 \[hep-th\]](#).
- [79] L. Boubekeur and D. H. Lyth, “Hilltop inflation,” *JCAP* **0507** (2005) 010, [arXiv:hep-ph/0502047 \[hep-ph\]](#).
- [80] H. Matsui and F. Takahashi, “Eternal Inflation and Swampland Conjectures,” *Phys. Rev.* **D99** no. 2, (2019) 023533, [arXiv:1807.11938 \[hep-th\]](#).

- [81] K. Dimopoulos, “Steep Eternal Inflation and the Swampland,” *Phys. Rev.* **D98** no. 12, (2018) 123516, [arXiv:1810.03438 \[gr-qc\]](#).
- [82] W. H. Kinney, “Eternal Inflation and the Refined Swampland Conjecture,” *Phys. Rev. Lett.* **122** no. 8, (2019) 081302, [arXiv:1811.11698 \[astro-ph.CO\]](#).
- [83] S. Brahma and S. Shandera, “Stochastic eternal inflation is in the swampland,” *JHEP* **11** (2019) 016, [arXiv:1904.10979 \[hep-th\]](#).
- [84] Z. Wang, R. Brandenberger, and L. Heisenberg, “Eternal Inflation, Entropy Bounds and the Swampland,” [arXiv:1907.08943 \[hep-th\]](#).
- [85] J. J. Blanco-Pillado, H. Deng, and A. Vilenkin, “Eternal Inflation in Swampy Landscapes,” [arXiv:1909.00068 \[gr-qc\]](#).
- [86] C.-M. Lin, “Topological Eternal Hilltop Inflation and the Swampland Criteria,” [arXiv:1912.00749 \[hep-th\]](#).
- [87] M. Scalisi and I. Valenzuela, “Swampland distance conjecture, inflation and α -attractors,” *JHEP* **08** (2019) 160, [arXiv:1812.07558 \[hep-th\]](#).
- [88] Y. Sekino and L. Susskind, “Fast Scramblers,” *JHEP* **10** (2008) 065, [arXiv:0808.2096 \[hep-th\]](#).