

A Rigorous Theory of Conditional Mean Embeddings

Ilja Klebanov¹Ingmar Schuster²T. J. Sullivan^{1,3,4}

January 7, 2022

Abstract: Conditional mean embeddings (CMEs) have proven themselves to be a powerful tool in many machine learning applications. They allow the efficient conditioning of probability distributions within the corresponding reproducing kernel Hilbert spaces (RKHSs) by providing a linear-algebraic relation for the kernel mean embeddings of the respective joint and conditional probability distributions. Both centred and uncentred covariance operators have been used to define CMEs in the existing literature. In this paper, we develop a mathematically rigorous theory for both variants, discuss the merits and problems of each, and significantly weaken the conditions for applicability of CMEs. In the course of this, we demonstrate a beautiful connection to Gaussian conditioning in Hilbert spaces.

Keywords: conditional mean embedding, kernel mean embedding, Gaussian measure, reproducing kernel Hilbert space.

2010 Mathematics Subject Classification: 46E22, 62J02, 28C20.

1. Introduction

Reproducing kernel Hilbert spaces (RKHSs) have long been popular tools in machine learning because of the powerful property — often called the “kernel trick” — that many problems posed in terms of the base set \mathcal{X} of the RKHS \mathcal{H} (e.g. classification into two or more classes) become linear-algebraic problems in \mathcal{H} under the embedding of \mathcal{X} into \mathcal{H} induced by the reproducing kernel $k: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$. This insight has been used to define the *kernel mean embedding* (KME; [Berlinet and Thomas-Agnan, 2004](#); [Smola et al., 2007](#)) $\mu_X \in \mathcal{H}$ of an \mathcal{X} -valued random variable X as the \mathcal{H} -valued mean of the embedded random variable $k(X, \cdot)$, and also the *conditional mean embedding* (CME; [Fukumizu et al., 2004a](#), [Song et al., 2009](#)), which seeks to perform conditioning of the original random variable X through application of the Gaussian conditioning formula (also known as the Kálmán update) to the embedded *non-Gaussian* random variable $k(X, \cdot)$. This article aims to provide rigorous mathematical foundations for this attractive but apparently naïve approach to conditional probability, and hence to Bayesian inference.

¹ Zuse Institute Berlin, Takustraße 7, 14195 Berlin, Germany. klebanov@zib.de, sullivan@zib.de

² Zalando SE, 11501 Berlin, Germany. ingmar.schuster@zalando.de

³ Freie Universität Berlin, Arnimallee 6, 14195 Berlin, Germany. t.j.sullivan@fu-berlin.de

⁴ Mathematics Institute and School of Engineering, The University of Warwick, Coventry, CV4 7AL, United Kingdom. t.j.sullivan@warwick.ac.uk




Figure 1.1: While conditioning of the probability distributions in the original spaces \mathcal{X}, \mathcal{Y} is a possibly complicated, non-linear problem, the corresponding formula for their kernel mean embeddings reduces to elementary linear algebra — a common guiding theme when working with reproducing kernel Hilbert spaces.

To be somewhat more precise — while deferring technical points such as topological considerations, existence and uniqueness of conditional distributions etc. to [Section 2](#) — let us fix two RKHSs \mathcal{H} and \mathcal{G} over \mathcal{X} and \mathcal{Y} respectively, with reproducing kernels k and ℓ and canonical feature maps $\varphi(x) := k(x, \cdot)$ and $\psi(y) := \ell(y, \cdot)$. Let X and Y be random variables taking values in \mathcal{X} and \mathcal{Y} respectively with joint distribution \mathbb{P}_{XY} on $\mathcal{X} \times \mathcal{Y}$. Let μ_X , μ_Y , and $\mu_{Y|X=x}$ denote the kernel mean embeddings (KMEs) of the marginal distributions \mathbb{P}_X of X , \mathbb{P}_Y of Y , and the conditional distribution $\mathbb{P}_{Y|X=x}$ of Y given $X = x$ given by

$$\mu_X := \mathbb{E}[\varphi(X)] \in \mathcal{H}, \quad \mu_Y := \mathbb{E}[\psi(Y)] \in \mathcal{G}, \quad \mu_{Y|X=x} := \mathbb{E}[\psi(Y)|X = x] \in \mathcal{G}. \quad (1.1)$$

The *conditional mean embedding* (CME) offers a way to perform conditioning of probability distributions on \mathcal{X} and \mathcal{Y} by means of linear algebra in the corresponding feature spaces \mathcal{H} and \mathcal{G} ([Figure 1.1](#)). In terms of the kernel covariance operator C_X and cross-covariance operator C_{YX} defined later in [\(2.3\)](#), if C_X is invertible and $\mathbb{E}[g(Y)|X = \cdot]$ is an element of \mathcal{H} whenever $g \in \mathcal{G}$, then the well-known formula for the CME ([Song et al., 2009](#), Theorem 4) is

$$\mu_{Y|X=x} = C_{YX}C_X^{-1}\varphi(x), \quad x \in \mathcal{X}. \quad (1.2)$$

(We emphasise here that the CME $\mu_{Y|X=x}$ is *defined* in [\(1.1\)](#) as the KME of $\mathbb{P}_{Y|X=x}$; the claim implicit in [\(1.2\)](#) is that $\mu_{Y|X=x}$ can be *realised* through simple linear algebra involving cross-covariance operators; cf. the discussion of [Park and Muandet \(2020\)](#).) Note that there are in fact two theories of CMEs, one working with *centred* covariance operators ([Fukumizu et al., 2004a; Song et al., 2009](#)) and the other with *uncentred* ones ([Fukumizu et al., 2013](#)). We will discuss both theories in detail, but let us focus for a moment on the centred case for which the above formula was originally derived.

In the trivial case where X and Y are independent, the CME should yield $\mu_{Y|X=x} = \mu_Y$. However, independence implies that $C_{YX} = 0$, and so [\(1.2\)](#) yields $\mu_{Y|X=x} = 0$, regardless of x . In order to understand what has gone wrong it is helpful to consider in turn the two cases in which the constant function $\mathbf{1}_{\mathcal{X}}: x \mapsto 1$ is, or is not, an element of \mathcal{H} .

- If $\mathbb{1}_{\mathcal{X}} \in \mathcal{H}$, then C_X cannot be injective, since $C_X \mathbb{1}_{\mathcal{X}} = 0$, and (1.2) is not applicable.
- If $\mathbb{1}_{\mathcal{X}} \notin \mathcal{H}$ and X and Y are independent, then the assumption $\mathbb{E}[g(Y)|X = \cdot] \in \mathcal{H}$ for $g \in \mathcal{G}$ cannot be fulfilled (except for those special elements $g \in \mathcal{G}$ for which $\mathbb{E}[g(Y)] = 0$ or if $\mathbb{E}[\ell(y, Y)] = 0$ for all $y \in \mathcal{Y}$, respectively), and (1.2) is again not applicable.

In summary, (1.2) is never applicable for independent random variables except in certain degenerate cases. Note that this problem does not occur in the case of uncentred operators, where ${}^u C_X$ (defined in (2.5)) is typically injective.

Therefore, this paper aims to provide a rigorous theory of CMEs that addresses not only the above-mentioned pathology but also substantially generalises the assumptions under which CME can be performed. We will treat both centred and uncentred (cross-)covariance operators, with particular emphasis on the centred case, and will also exhibit a connection to Gaussian conditioning in general Hilbert spaces.

- (1) The standard assumption $\mathbb{E}[g(Y)|X = \cdot] \in \mathcal{H}$ for CME is rather restrictive.¹ We show in Section 4 that this assumption can be significantly weakened in the case of centred kernel (cross-)covariance operators as defined in (2.3): only $\mathbb{E}[g(Y)|X = \cdot]$ shifted by some constant function needs to lie in \mathcal{H} (Assumption B). In this setting, the correct expression of the CME formula is

$$\mu_{Y|X=x} = \mu_Y + (C_X^\dagger C_{XY})^* (\varphi(x) - \mu_X) \quad \text{for } \mathbb{P}_X\text{-a.e. } x \in \mathcal{X}, \quad (1.3)$$

where A^* denotes the adjoint and A^\dagger the Moore–Penrose pseudo-inverse of a linear operator A . As a first sanity check, note that this formula indeed yields $\mu_{Y|X=x} = \mu_Y$ when X and Y are independent. Similarly, as shown in Section 5, for *uncentred* kernel (cross-)covariance operators ${}^u C_X$ and ${}^u C_{XY}$ as defined later in (2.5), the correct formulation of the CME is

$$\mu_{Y|X=x} = ({}^u C_X^\dagger {}^u C_{XY})^* \varphi(x) \quad \text{for } \mathbb{P}_X\text{-a.e. } x \in \mathcal{X}. \quad (1.4)$$

- (2) Furthermore, the assumption $\mathbb{E}[g(Y)|X = \cdot] \in \mathcal{H}$, $g \in \mathcal{G}$, is hard to check in most applications. To the best of our knowledge, the only verifiable condition that implies this assumption is given by Fukumizu et al. (2004b, Proposition 4). However, this condition is itself difficult to check². We will present weaker assumptions (Assumption B*) for the applicability of CMEs that hold whenever the kernel k is characteristic.³ Characteristic kernels are well studied (see e.g. Sriperumbudur et al. (2010)) and therefore provide a verifiable condition as desired.
- (3) The applicability of (1.2) requires the additional assumptions that C_X is injective and that $\varphi(x)$ lies in the range of C_X , which is also hard to verify in practice.⁴ We show that

¹Fukumizu et al. (2013) themselves write “Note, however, that the assumptions [...] may not hold in general; we can easily give counterexamples for the latter in the case of Gaussian kernels.”. More precisely, for a Gaussian kernel k on, say, $[0, 1]$ and independent random variables X and Y , $\mathbb{E}[g(Y)|X = \cdot]$ is a constant function for each $g \in \mathcal{G}$, which does not lie in the RKHS corresponding to k (unless it happens to be the zero function) by Steinwart et al. (2006, Corollary 5) or Steinwart and Christmann (2008, Corollary 4.44).

²The original condition of Fukumizu et al. (2004a, Proposition 4) was verifiable in certain situations, but the proposition itself turned out to be incorrect. The corrected condition in the erratum (Fukumizu et al., 2004b) seems to be much harder to check — at least, no explicit case is given in which it is easier to verify than $\mathbb{E}[g(Y)|X = \cdot] \in \mathcal{H}$ for each $g \in \mathcal{G}$.

³A kernel k is called *characteristic* (Fukumizu et al., 2008) if the kernel mean embedding is injective as a function from $\{\mathbb{Q} \mid \mathbb{Q} \text{ is a prob. meas. on } \mathcal{X} \text{ with } \int_{\mathcal{X}} \|\varphi(x)\|_{\mathcal{H}} d\mathbb{Q}(x) < \infty\}$ into \mathcal{H} ; naturally, the KME cannot be injective as a function from the space of random variables on \mathcal{X} to \mathcal{H} , since random variables with the same law embed to the same point of \mathcal{H} .

⁴Note that, typically, $\dim \mathcal{H} = \infty$, in which case the compact operator C_X cannot possibly be surjective. To verify that $\varphi(x) \in \text{ran } C_X$, one would need to compute a singular value decomposition $C_X = \sum_{n \in \mathbb{N}} \sigma_n h_n \otimes h_n$ of C_X and check the Picard condition $\sum_{n \in \mathbb{N}} \sigma_n^{-2} \langle \varphi(x), h_n \rangle_{\mathcal{H}}^2 < \infty$.

both assumptions can be avoided completely by replacing $C_{YX}C_X^{-1}$ in (1.2) by $(C_X^\dagger C_{XY})^*$ in (1.3) and $({}^u C_X^\dagger {}^u C_{XY})^*$ in (1.4), which turn out to be globally-defined and bounded operators under rather weak assumptions (Assumptions C and u C).

- (4) The experienced reader will also observe that, modulo the replacement of $C_{YX}C_X^{-1}$ by $(C_X^\dagger C_{XY})^*$, (1.3) is identical to the familiar Sherman–Morrison–Woodbury / Schur complement formula for conditional Gaussian distributions, a connection on which we will elaborate in detail in Section 7. We call particular attention to the fact that the random variable $(\psi(Y), \varphi(X))$, which has no reason to be normally distributed, behaves very much like a Gaussian random variable in terms of its conditional mean.

Remark 1.1. Note that we stated (1.3) and (1.4) only for \mathbb{P}_X -a.e. $x \in \mathcal{X}$. This is the best that one can generally hope for, since the regular conditional probability $\mathbb{P}_{Y|X=x}$ is uniquely determined only for \mathbb{P}_X -a.e. $x \in \mathcal{X}$ (Kallenberg, 2006, Theorem 5.3). The work on CMEs so far completely ignores the fact that conditioning (especially on events of the form $X = x$) is not trivial, requires certain assumptions and, in general, yields results only for \mathbb{P}_X -a.e. $x \in \mathcal{X}$. In particular, the condition on $\mathbb{E}[g(Y)|X = \cdot]$ to lie in \mathcal{H} is ill posed, since these functions are uniquely defined only \mathbb{P}_X -a.e., which in certain situations may be practically nowhere, and the same reasoning applies to the above-mentioned condition given by Fukumizu et al. (2004b, Proposition 4). The existence and almost sure uniqueness of the regular conditional probability distribution $\mathbb{P}_{Y|X=x}$ will be addressed in a precise manner in Section 2.

Remark 1.2. The focus of this paper is the validity of the non-regularised *population* formulation of the CME in terms of the covariance structure of the KME of the data-generating distribution \mathbb{P}_{XY} . The construction of valid CME formulae based on *empirical sample data* (i.e. finitely many draws from \mathbb{P}_{XY}) is vital in practice but is also much harder to analyse. We give some remarks on this setting in Appendix B.

The rest of the paper is structured as follows. Section 2 establishes the notation and problem setting, and motivates some of the assumptions that are made. Section 3 discusses several critical assumptions for the applicability of the theory of CMEs and the relations among them. Section 4 proceeds to build a rigorous theory of CMEs using centred covariance operators, with the main results being Theorems 4.3 and 4.4, whereas Section 5 does the same for uncentred covariance operators, with the main results being Theorems 5.3 and 5.4. Section 6 reviews the established theory for the conditioning of Gaussian measures on Hilbert spaces, and this is then used in Section 7 to rigorously connect the theory of CMEs to the conditioning of Gaussian measures, with the main result being Theorem 7.1. We give some closing remarks in Section 8. Appendix A contains various auxiliary technical results and Appendix B discusses the possible extension of our results to empirical estimation of CMEs.

2. Setup and Notation

Throughout this paper, when considering Hilbert-space valued random variables $U \in \mathcal{L}^2(\Omega, \Sigma, \mathbb{P}; \mathcal{G})$ and $V \in \mathcal{L}^2(\Omega, \Sigma, \mathbb{P}; \mathcal{H})$ defined over a probability space $(\Omega, \Sigma, \mathbb{P})$, the expected value $\mathbb{E}[U] := \int_\Omega U(\omega) d\mathbb{P}(\omega)$ is meant in the sense of a Bochner integral (Diestel and Uhl, 1977, Section II.2), as are the uncentred and centred cross-covariance operators

$${}^u\text{Cov}[U, V] := \mathbb{E}[U \otimes V] \quad \text{and} \quad \text{Cov}[U, V] := \mathbb{E}[(U - \mathbb{E}[U]) \otimes (V - \mathbb{E}[V])]$$

from \mathcal{H} into \mathcal{G} , where, for $h \in \mathcal{H}$ and $g \in \mathcal{G}$, the outer product $g \otimes h: \mathcal{H} \rightarrow \mathcal{G}$ is the rank-one linear operator $(g \otimes h)(h') := \langle h, h' \rangle_{\mathcal{H}} g$. Naturally, we write ${}^u\text{Cov}[U]$ and $\text{Cov}[U]$ for the

covariance operators ${}^u\text{Cov}[U, U]$ and $\text{Cov}[U, U]$ respectively, and all of the above reduces to the usual definitions in the scalar-valued case. Both the centred and uncentred covariance operators of a square-integrable random variable are self-adjoint and non-negative, and — in the separable Hilbert case that is our exclusive focus — also trace-class (see [Baker \(1973\)](#); [Sazonov \(1958\)](#) for the centred case; the uncentred case follows from [Gohberg and Kreĭn \(1969, Corollary 2.1\)](#)).

Our treatment of CMEs will operate under the following assumptions and notation:

- Assumption 2.1.** (a) $(\Omega, \Sigma, \mathbb{P})$ is a probability space, \mathcal{X} is a measurable space, and \mathcal{Y} is a Borel space.⁵
- (b) $k: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ and $\ell: \mathcal{Y} \times \mathcal{Y} \rightarrow \mathbb{R}$ are symmetric and positive definite kernels, such that $k(x, \cdot)$ and $\ell(y, \cdot)$ are Borel-measurable functions for each $x \in \mathcal{X}$ and $y \in \mathcal{Y}$.
- (c) $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathcal{H}})$ and $(\mathcal{G}, \langle \cdot, \cdot \rangle_{\mathcal{G}})$ are the corresponding RKHSs, which we assume to be separable. Indeed, according to [Owhadi and Scovel \(2017\)](#), if the base sets \mathcal{X} and \mathcal{Y} are separable absolute Borel spaces or analytic subsets of Polish spaces, then separability of \mathcal{H} and \mathcal{G} follows from the measurability of their respective kernels and feature maps.
- (d) The corresponding canonical feature maps are $\varphi: \mathcal{X} \rightarrow \mathcal{H}$, $\varphi(x) := k(x, \cdot)$, and $\psi: \mathcal{Y} \rightarrow \mathcal{G}$, $\psi(y) := \ell(y, \cdot)$ respectively. Note that they satisfy the “reproducing properties” $\langle h, \varphi(x) \rangle_{\mathcal{H}} = h(x)$, $\langle g, \psi(y) \rangle_{\mathcal{G}} = g(y)$ for $x \in \mathcal{X}, y \in \mathcal{Y}, h \in \mathcal{H}, g \in \mathcal{G}$ and that φ and ψ are Borel measurable in view of [Steinwart and Christmann \(2008, Lemma 4.25\)](#).
- (e) $X: \Omega \rightarrow \mathcal{X}$ and $Y: \Omega \rightarrow \mathcal{Y}$ are random variables with distributions \mathbb{P}_X and \mathbb{P}_Y and joint distribution \mathbb{P}_{XY} . [Assumption 2.1\(a\)](#) and [Kallenberg \(2006, Theorem 5.3\)](#) ensure the existence of a \mathbb{P}_X -a.e.-unique regular version of the conditional probability distribution $\mathbb{P}_{Y|X=x}$; the choice of a representative of $\mathbb{P}_{Y|X=x}$ has no impact on our results. We assume that

$$\mathbb{E}[\|\varphi(X)\|_{\mathcal{H}}^2 + \|\psi(Y)\|_{\mathcal{G}}^2] < \infty, \quad (2.1)$$

which also implies that $\mathcal{X}_Y := \{x \in \mathcal{X} \mid \mathbb{E}[\|\psi(Y)\|_{\mathcal{G}}^2 | X = x] < \infty\}$ has full \mathbb{P}_X measure.⁶ Hence, $\mathcal{H} \subseteq \mathcal{L}^2(\mathbb{P}_X)$, $\mathcal{G} \subseteq \mathcal{L}^2(\mathbb{P}_Y)$ and $\mathcal{G} \subseteq \mathcal{L}^2(\mathbb{P}_{Y|X=x})$ for $x \in \mathcal{X}_Y$ since, by the reproducing property and the Cauchy–Schwarz inequality,

$$\begin{aligned} \|h\|_{\mathcal{L}^2(\mathbb{P}_X)}^2 &= \int_{\mathcal{X}} |h(x)|^2 d\mathbb{P}_X(x) = \int_{\mathcal{X}} |\langle h, \varphi(x) \rangle_{\mathcal{H}}|^2 d\mathbb{P}_X(x) \\ &\leq \int_{\mathcal{X}} \|h\|_{\mathcal{H}}^2 \|\varphi(x)\|_{\mathcal{H}}^2 d\mathbb{P}_X(x) = \mathbb{E}[\|\varphi(X)\|_{\mathcal{H}}^2] \|h\|_{\mathcal{H}}^2 \end{aligned} \quad (2.2)$$

for all $h \in \mathcal{H}$, and similarly for $g \in \mathcal{G}$ and $\mathbb{P}_Y, \mathbb{P}_{Y|X=x}, x \in \mathcal{X}_Y$. It follows from (2.2) that the inclusions $\iota_{\varphi, \mathbb{P}_X}: \mathcal{H} \hookrightarrow \mathcal{L}^2(\mathbb{P}_X)$, $\iota_{\psi, \mathbb{P}_Y}: \mathcal{G} \hookrightarrow \mathcal{L}^2(\mathbb{P}_Y)$ are bounded linear operators and so is $\iota_{\psi, \mathbb{P}_{Y|X=x}}: \mathcal{G} \hookrightarrow \mathcal{L}^2(\mathbb{P}_{Y|X=x})$ for $x \in \mathcal{X}_Y$.

- (f) We further assume that, for all $h \in \mathcal{H}$, $h = 0$ \mathbb{P}_X -a.e. in \mathcal{X} if and only if $h = 0$, i.e. almost everywhere equality separates points in \mathcal{H} . This assumption clearly holds if k is continuous and the topological support of \mathbb{P}_X is all of \mathcal{X} .⁷ It ensures that we can view \mathcal{H} as a subspace of $L^2(\mathbb{P}_X)$ and write $f \in \mathcal{H}$ for functions $f \in L^2(\mathbb{P}_X)$ whenever there exists $h \in \mathcal{H}$ (which, by this assumption, is unique) such that $f = h$ \mathbb{P}_X -a.e.
- (g) Several derivations will rely on the Bochner space $L^2(\mathbb{P}_X; \mathcal{F})$, which is isometrically isomorphic to the Hilbert tensor product space $L^2(\mathbb{P}_X) \otimes \mathcal{F}$. Here, \mathcal{F} denotes another Hilbert

⁵A space \mathcal{Y} is called a *Borel space* if it is Borel isomorphic to a Borel subset of $[0, 1]$. In particular, \mathcal{Y} is a Borel space if it is Polish, i.e. if it is separable and completely metrisable; see [Kallenberg \(2006, Chapter 1\)](#).

⁶Otherwise, $\mathbb{E}[\|\psi(Y)\|_{\mathcal{G}}^2] = \mathbb{E}[\mathbb{E}[\|\psi(Y)\|_{\mathcal{G}}^2 | X]]$ could not be finite.

⁷If k is continuous, then so is every $h \in \mathcal{H}$ ([Saitoh and Sawano, 2016, Theorem 2.3](#)). So, if $h \in \mathcal{H}$ and $|h(x)| = \varepsilon > 0$ for some $x \in \mathcal{X}$, then $|h| > \varepsilon/2$ on some open neighbourhood of x . Thus, if $\text{supp}(\mathbb{P}_X) = \mathcal{X}$, then $h = 0$ \mathbb{P}_X -a.e. cannot hold.

space, which in our case will be equal to either \mathbb{R} or \mathcal{G} . Motivated by the discussion in [Section 1](#) and the fact that $\mathbb{C}\text{ov}[f(X), f(X)] = \mathbb{V}[f(X)] = 0$ if and only if f is \mathbb{P}_X -a.e. constant, we consider the quotient space $L^2_{\mathcal{C}}(\mathbb{P}_X; \mathcal{F}) := L^2(\mathbb{P}_X; \mathcal{F})/\mathcal{C}$,⁸

$$\mathcal{C} := \{f \in L^2(\mathbb{P}_X; \mathcal{F}) \mid \exists c \in \mathcal{F}: f(x) = c \text{ for } \mathbb{P}_X\text{-a.e. } x \in \mathcal{X}\},$$

$$\langle [f_1], [f_2] \rangle_{L^2_{\mathcal{C}}(\mathbb{P}_X; \mathcal{F})} := \langle f_1 - \mathbb{E}[f_1(X)], f_2 - \mathbb{E}[f_2(X)] \rangle_{L^2(\mathbb{P}_X; \mathcal{F})}.$$

Note that, in the case $\mathcal{F} = \mathbb{R}$, we obtain $\langle [f_1], [f_2] \rangle_{L^2_{\mathcal{C}}(\mathbb{P}_X; \mathbb{R})} = \mathbb{C}\text{ov}[f_1(X), f_2(X)]$, in which case we will abbreviate the space $L^2(\mathbb{P}_X; \mathbb{R})$ by $L^2(\mathbb{P}_X)$ or simply L^2 and the space $L^2_{\mathcal{C}}(\mathbb{P}_X; \mathbb{R})$ by $L^2_{\mathcal{C}}(\mathbb{P}_X)$ or simply $L^2_{\mathcal{C}}$. For any closed subspace $U \subseteq L^2(\mathbb{P}_X)$ we can view $U \otimes \mathcal{F}$ as a subspace of $L^2(\mathbb{P}_X; \mathcal{F})$ by the above isometry and identify $(U \otimes \mathcal{F})_{\mathcal{C}} := (U \otimes \mathcal{F})/((U \otimes \mathcal{F}) \cap \mathcal{C})$ with a subspace of $L^2_{\mathcal{C}}(\mathbb{P}_X; \mathcal{F})$. Note that, in the particular case $U \subseteq \mathcal{H}$ (with U closed in $L^2(\mathbb{P}_X)$), the construction of $U \otimes \mathcal{F}$ and $(U \otimes \mathcal{F})_{\mathcal{C}}$ treats U as a subspace of $L^2(\mathbb{P}_X)$ and ignores the existence of the RKHS norm $\|\cdot\|_{\mathcal{H}}$.

- (h) We use overlines and superscripts to denote topological closures, so that, for example, $\overline{\mathcal{H}_{\mathcal{C}}}^{L^2_{\mathcal{C}}}$ is the closure of $\mathcal{H}_{\mathcal{C}}$ with respect to the norm $\|\cdot\|_{L^2_{\mathcal{C}}}$ and $\overline{\mathcal{H}}^{L^2}$ is the closure of \mathcal{H} with respect to the norm $\|\cdot\|_{L^2}$.
- (i) Since φ and ψ are Borel measurable, $Z := (\psi(Y), \varphi(X))$ is a well-defined $\mathcal{G} \oplus \mathcal{H}$ -valued random variable; [\(2.1\)](#) ensures that Z has finite second moment, hence its mean $\mathbb{E}[Z]$ and covariance operator $\mathbb{C}\text{ov}[Z]$ are well defined, Sazonov's theorem implies that $\mathbb{C}\text{ov}[Z]$ has finite trace, and we obtain the following block structures:

$$\mu := \mathbb{E}\left[\begin{pmatrix} \psi(Y) \\ \varphi(X) \end{pmatrix}\right] = \begin{pmatrix} \mu_Y \\ \mu_X \end{pmatrix}, \quad C := \mathbb{C}\text{ov}\left[\begin{pmatrix} \psi(Y) \\ \varphi(X) \end{pmatrix}\right] = \begin{pmatrix} C_Y & C_{YX} \\ C_{XY} & C_X \end{pmatrix}, \quad (2.3)$$

where the components

$$\begin{aligned} \mu_Y &:= \mathbb{E}[\psi(Y)], & C_Y &:= \mathbb{C}\text{ov}[\psi(Y)], & C_{YX} &:= \mathbb{C}\text{ov}[\psi(Y), \varphi(X)], \\ \mu_X &:= \mathbb{E}[\varphi(X)], & C_{XY} &:= \mathbb{C}\text{ov}[\varphi(X), \psi(Y)], & C_X &:= \mathbb{C}\text{ov}[\varphi(X)] \end{aligned}$$

are called the *kernel mean embeddings* (KME) and *kernel (cross-)covariance operators*, respectively. Note that $C_{XY}^* = C_{YX}$ and that the reproducing properties translate to the KMEs and covariance operators as follows: for arbitrary $h, h' \in \mathcal{H}$ and $g \in \mathcal{G}$,

$$\begin{aligned} \langle h, \mu_X \rangle_{\mathcal{H}} &= \mathbb{E}[h(X)], \\ \langle h, C_X h' \rangle_{\mathcal{H}} &= \mathbb{C}\text{ov}[h(X), h'(X)], \\ \langle h, C_{XY} g \rangle_{\mathcal{H}} &= \mathbb{C}\text{ov}[h(X), g(Y)], \end{aligned}$$

and so on. We are further interested in the *conditional kernel mean embedding* and the *conditional kernel covariance operator* given by

$$\mu_{Y|X=x} = \mathbb{E}[\psi(Y)|X=x], \quad C_{Y|X=x} = \mathbb{C}\text{ov}[\psi(Y)|X=x], \quad x \in \mathcal{X}_Y. \quad (2.4)$$

We set $\mu_{Y|X=x} := 0$ on the \mathbb{P}_X -null set $\mathcal{X} \setminus \mathcal{X}_Y$. Similarly, $Z = (\psi(Y), \varphi(X))$ has the uncentred kernel covariance structure

$${}^u C := {}^u \mathbb{C}\text{ov}\left[\begin{pmatrix} \psi(Y) \\ \varphi(X) \end{pmatrix}\right] = \begin{pmatrix} {}^u C_Y & {}^u C_{YX} \\ {}^u C_{XY} & {}^u C_X \end{pmatrix}, \quad (2.5)$$

where ${}^u C_Y := {}^u \mathbb{C}\text{ov}[\psi(Y)]$ etc. Note that, for $f_1, f_2 \in L^2(\mathbb{P}_X)$, ${}^u \mathbb{C}\text{ov}(f_1(X), f_2(X)) = \langle f_1, f_2 \rangle_{L^2(\mathbb{P}_X)}$, and similarly for functions of Y .

⁸By the variational characterisation of the expected value $\mathbb{E}[Z]$ of a random variable $Z \in L^2(\mathbb{P}; \mathcal{F})$, $\mathbb{E}[Z] = \arg \min_{m \in \mathcal{F}} \mathbb{E}[\|Z - m\|_{\mathcal{F}}^2]$, the norm $\|\cdot\|_{L^2_{\mathcal{C}}(\mathbb{P}_X; \mathcal{F})}$ coincides with the norm $\|[f]\| = \inf_{m \in \mathcal{C}} \|f - m\|_{L^2(\mathbb{P}_X; \mathcal{F})}$ induced on $L^2_{\mathcal{C}}(\mathbb{P}_X; \mathcal{F})$ by the norm $\|\cdot\|_{L^2(\mathbb{P}_X; \mathcal{F})}$.

(j) For $g \in \mathcal{G}$ we let $f_g(x) := \mathbb{E}[g(Y)|X = x]$. More precisely,

$$f_g(x) := \begin{cases} \mathbb{E}[g(Y)|X = x], & \text{for } x \in \mathcal{X}_Y, \\ 0, & \text{otherwise.} \end{cases}$$

These functions f_g will be of particular importance since, for $g = \psi(y)$, $y \in \mathcal{Y}$ and $x \in \mathcal{X}$, we obtain $f_{\psi(y)}(x) = \mu_{Y|X=x}(y)$, our main object of interest (note that $\mu_{Y|X=x} \in \mathcal{G}$ for each $x \in \mathcal{X}$, and so its pointwise evaluation at $y \in \mathcal{Y}$ is meaningful). By (2.1), (2.2), and the law of total expectation, $f_g \in L^2(\mathbb{P}_X)$ for every $g \in \mathcal{G}$, since

$$\begin{aligned} \|f_g\|_{L^2(\mathbb{P}_X)} &= \mathbb{E}[f_g(X)^2] = \mathbb{E}[\mathbb{E}[g(Y)|X]^2] \leq \mathbb{E}[\mathbb{E}[g(Y)^2|X]] \\ &= \mathbb{E}[g(Y)^2] = \|g\|_{L^2(\mathbb{P}_Y)} < \infty. \end{aligned}$$

Further, another application of the law of total expectation yields

$$\mathbb{E}[f_g(X)] = \mathbb{E}[g(Y)], \quad \mathbb{E}[f_{\psi(y)}(X)] = \mu_Y(y). \quad (2.6)$$

(k) For a linear operator A between Hilbert spaces, A^\dagger denotes its Moore–Penrose pseudo-inverse, i.e. the unique extension of $A|_{(\ker A)^\perp}^{-1} : \text{ran } A \rightarrow (\ker A)^\perp$ to a linear operator A^\dagger defined on $\text{dom } A^\dagger := (\text{ran } A) \oplus (\text{ran } A)^\perp$ subject to the criterion that $\ker A^\dagger = (\text{ran } A)^\perp$. In general, $\text{dom } A^\dagger$ is a dense but proper subspace and A^\dagger is an unbounded operator; global definition and boundedness occur precisely when $\text{ran } A$ is closed; see e.g. Engl et al. (1996, Section 2.1).

Remark 2.2. Measurability of $k(x, \cdot)$ and $\ell(y, \cdot)$ together with the separability of \mathcal{H} and \mathcal{G} guarantee the measurability of φ and ψ (Steinwart and Christmann, 2008, Lemma 4.25). Separability of \mathcal{H} and \mathcal{G} is also needed for Gaussian conditioning (see Owhadi and Scovel (2018) and Section 6), for the existence of a countable orthonormal basis of \mathcal{H} , and to ensure that weak (Pettis) and strong (Bochner) measurability of Hilbert-valued random variables coincide.

3. The Crucial Assumptions for CMEs

This section discusses various versions of the assumption $f_g \in \mathcal{H}$ under which we are going to prove various versions of the CME formula (note that, by Assumption 2.1(f), their formulations are unambiguous).

Assumption A. For all $g \in \mathcal{G}$, $f_g \in \mathcal{H}$.

Assumption B. For all $g \in \mathcal{G}$ there exists a function $h_g \in \mathcal{H}$ and a constant $c_g \in \mathbb{R}$ such that $h_g = f_g - c_g \mathbb{P}_X$ -a.e. in \mathcal{X} .

Assumption C. For all $g \in \mathcal{G}$ there exists a function $h_g \in \mathcal{H}$ such that

$$\text{Cov}[h_g(X) - f_g(X), h(X)] = 0 \quad \text{for all } h \in \mathcal{H}.$$

In this case we denote $c_g := \mathbb{E}[f_g(X) - h_g(X)]$ (in conformity with Assumption B).

Assumption ^uC. For all $g \in \mathcal{G}$ there exists a function $h_g \in \mathcal{H}$ such that

$${}^u\text{Cov}[h_g(X) - f_g(X), h(X)] = \langle h_g - f_g, h \rangle_{L^2(\mathbb{P}_X)} = 0 \quad \text{for all } h \in \mathcal{H}.$$




Figure 3.1: A hierarchy of CME-related assumptions. Sufficient conditions for validity of the CME formula are indicated by solid boxes while the insufficient Assumptions \mathbf{C} and ${}^u\mathbf{C}$, indicated by dashed boxes, have several strong theoretical implications and Assumption \mathbf{C} has a beautiful connection to Gaussian conditioning (Theorem 7.3). Assumption \mathbf{B}^* is the most favorable one, since it is verifiable in practice, and, by Lemma A.4, in particular is fulfilled if the kernel is universal or even just characteristic (marked in green). The shaded boxes correspond to Theorems 4.3, 4.4, 5.3, and 5.4.

Remark 3.1. Note that $\mathbf{A} \implies \mathbf{B} \implies \mathbf{C}$, that $\mathbf{A} \implies {}^u\mathbf{C}$, that $\mathbf{C} \implies \mathbf{B}$ if $\mathcal{H}_C \subseteq L^2_C(\mathbb{P}_X)$ is dense, and that $\mathbf{C} \implies \mathbf{A}$ and ${}^u\mathbf{C} \implies \mathbf{A}$ if $\mathcal{H} \subseteq L^2(\mathbb{P}_X)$ is dense.

Unlike Assumption \mathbf{A} , Assumptions \mathbf{B} and \mathbf{C} do not require the unfavourable property $\mathbb{1}_X \in \mathcal{H}$ for independent random variables X and Y . Instead, this case reduces to the trivial condition $0 \in \mathcal{H}$. At the same time, the proofs of the key properties of CMEs are not affected by replacing Assumption \mathbf{A} with Assumption \mathbf{B} as long as we work with centred operators (see Theorems 4.1 and 4.3 below). Therefore, it is surprising that this modification has not been considered earlier, even though the issues with independent random variables have been observed before (Fukumizu et al., 2013). One reason might be that, instead of centred operators, researchers started using uncentred ones, for which such a modification is not feasible.

Assumption \mathbf{C} , on the other hand, is not strong enough for proving the main formula for CMEs (the last statement of Theorem 4.3). Clearly, this cannot be expected: If \mathcal{X} and \mathcal{G} are reasonably large, but \mathcal{H} is not rich enough, e.g. $\mathcal{H} = \{0\}$ or $\mathcal{H} = \text{span}\{\mathbb{1}_X\}$, then no map from \mathcal{H} to \mathcal{G} can cover sufficiently many kernel mean embeddings, in particular the embeddings of the conditional probability $\mathbb{P}_{Y|X=x}$ for various x (while Assumption \mathbf{C} is trivially fulfilled for these choices of \mathcal{H}). The weakness of Assumption \mathbf{C} lies in the fact that it only requires the vanishing of the orthogonal projection of $[h_g] - [f_g]$ onto \mathcal{H}_C . Only if \mathcal{H}_C is rich enough (e.g. if it is dense in L^2_C) can this condition have useful implications. A similar reasoning applies to Assumption ${}^u\mathbf{C}$.

While it is nice to have a weaker form of Assumption \mathbf{A} , the Assumptions \mathbf{A} , \mathbf{B} and \mathbf{C} remain hard to check in practice. Another condition, provided by Fukumizu et al. (2004b,

Proposition 4), is also hard to verify in most applications.² Since characteristic kernels are well studied in the literature, [Lemma A.4](#) gives hope for a verifiable condition for the applicability of CMEs: it states that \mathcal{H}_C is dense in $L_C^2(\mathbb{P}_X)$ whenever the kernel k is characteristic. So, if the denseness of \mathcal{H}_C in $L_C^2(\mathbb{P}_X)$ were sufficient for performing CMEs, then the condition that k be characteristic would be sufficient as well, thus providing a favorable criterion for the applicability of formula (1.3). A similar argumentation applies to the condition that \mathcal{H} is dense in $L^2(\mathbb{P}_X)$ and the condition that k is L^2 -universal.⁹ Unfortunately, neither condition implies [Assumption B](#). Therefore, we will consider the following slightly weaker versions of [Assumptions A](#) and [B](#), under which CMEs can be performed if one allows for certain finite-rank approximations of the (cross-)covariance operators:

Assumption A*. For all $g \in \mathcal{G}$, $f_g \in \overline{\mathcal{H}}^{L^2}$.

Assumption B*. For all $g \in \mathcal{G}$ there exists a function $h_g \in \overline{\mathcal{H}}^{L^2}$ and a constant $c_g \in \mathbb{R}$ such that $h_g = f_g - c_g \mathbb{P}_X$ -a.e. in \mathcal{X} .

Note that [Assumption C](#) and [Assumption ^uC](#) have no weaker versions, since they would become trivial if $h_g \in \mathcal{H}_C$ were replaced by $h_g \in \overline{\mathcal{H}_C}^{L^2}$ and $h_g \in \mathcal{H}$ by $h_g \in \overline{\mathcal{H}}^{L^2}$ respectively.

Remark 3.2. In terms of the spaces L_C^2 and \mathcal{H}_C , [Assumptions A–B*](#) can be reformulated as follows: For all $g \in \mathcal{G}$,

- (A) $f_g \in \mathcal{H}$;
- (B) $[f_g] \in \mathcal{H}_C$;
- (C) the orthogonal projection $P_{\overline{\mathcal{H}_C}^{L^2}}[f_g]$ of $[f_g]$ onto $\overline{\mathcal{H}_C}^{L^2}$ lies in \mathcal{H}_C ;
- (^uC) the orthogonal projection $P_{\overline{\mathcal{H}}^{L^2}}f_g$ of f_g onto $\overline{\mathcal{H}}^{L^2}$ lies in \mathcal{H} ;
- (A*) $f_g \in \overline{\mathcal{H}}^{L^2}$;
- (B*) $[f_g] \in \overline{\mathcal{H}_C}^{L^2}$.

In summary, we consider the hierarchy of assumptions illustrated in [Figure 3.1](#). The main contributions of this paper are rigorous proofs of three versions of the CME formula under various assumptions:

- [Theorem 4.3](#) uses [Assumption B](#) and centred operators.
- [Theorem 4.4](#) uses [Assumption B*](#) and finite-rank approximations of centred operators.
- [Theorem 5.3](#) uses [Assumption A](#) and uncentred operators.
- [Theorem 5.4](#) uses [Assumption A*](#) and finite-rank approximations of uncentred operators.

Note that the theorems for uncentred covariance operators require stronger assumptions than their centred counterparts and that [Theorem 5.4](#) provides weaker statements than its centred analogue [Theorem 4.4](#) (we show only the convergence in $L^2(\mathbb{P}_X; \mathcal{G})$, which does not guarantee convergence for \mathbb{P}_X -a.e. $x \in \mathcal{X}$).

4. Theory for Centred Operators

In this section we will formulate and prove two versions of the CME formula (1.3) — the original one under [Assumption B](#) and a weaker version involving finite-rank approximations $C_X^{(n)}, C_{XY}^{(n)}$ of the (cross-)covariance operators under [Assumption B*](#). The following theorem demonstrates

⁹A kernel k on \mathcal{X} is called *L^2 -universal* ([Sriperumbudur et al., 2011](#)) if it is Borel measurable and bounded and if \mathcal{H} is dense in $L^2(\mathbb{Q})$ for any probability measure \mathbb{Q} on \mathcal{X} . Any L^2 -universal kernel is characteristic ([Sriperumbudur et al., 2011](#)).

the importance of [Assumption C](#) (which follows from [Assumption B](#)). It implies that the range of C_{XY} is contained in that of C_X , making the operator $C_X^\dagger C_{XY}$ well defined. By [Theorem A.1](#) it is even a bounded operator, which is a non-trivial result requiring the application of the closed graph theorem.¹⁰

Similar considerations cannot be performed, in general, under [Assumption B*](#) alone: it can no longer be expected that $\text{ran } C_{XY} \subseteq \text{ran } C_X$, which is why we must introduce the above-mentioned finite-rank approximations in order to guarantee that $\text{ran } C_{XY}^{(n)} \subseteq \text{ran } C_X^{(n)}$.

In summary, [Assumption B](#) allows for the simple CME formula (1.3) by [Theorem 4.1](#), while under [Assumption B*](#) we must make a detour using certain approximations. Note that this distinction is very similar to the theory of Gaussian conditioning in Hilbert spaces introduced by [Owhadi and Scovel \(2018\)](#) and recapped in [Section 6](#) below, a connection that will be elaborated upon in detail in [Section 7](#).

Theorem 4.1. *Under [Assumption 2.1](#), the following statements are equivalent:*

- (i) [Assumption C](#) holds.
- (ii) For each $g \in \mathcal{G}$ there exists $h_g \in \mathcal{H}$ such that $C_X h_g = C_{XY} g$.
- (iii) $\text{ran } C_{XY} \subseteq \text{ran } C_X$.

Proof. Note that (iii) is just a reformulation of (ii), so we only have to prove (i) \iff (ii). Let $g \in \mathcal{G}$ and $h, h_g \in \mathcal{H}$. By [Lemma A.6](#), $\text{Cov}[h(X), f_g(X)] = \langle h, C_{XY} g \rangle_{\mathcal{H}}$, and so

$$\begin{aligned} \text{Cov}[h(X), h_g(X)] &= \text{Cov}[h(X), f_g(X)] \quad \forall h \in \mathcal{H} \iff \langle h, C_X h_g \rangle_{\mathcal{H}} = \langle h, C_{XY} g \rangle_{\mathcal{H}} \quad \forall h \in \mathcal{H} \\ &\iff C_X h_g = C_{XY} g. \end{aligned}$$

■

Note that [Assumption C](#) implies that $[h_g] \in \mathcal{H}_{\mathcal{C}}$ is the orthogonal projection of $[f_g] \in L_{\mathcal{C}}^2$ onto $\mathcal{H}_{\mathcal{C}}$ with respect to $\langle \cdot, \cdot \rangle_{L_{\mathcal{C}}^2}$ (see the reformulation of [Assumption C](#) in [Remark 3.2](#)). Therefore, there might be some ambiguity in the choice of $h_g \in \mathcal{H}$ if \mathcal{H} contains constant functions. However, there is a particular choice of h_g that always works:

Proposition 4.2. *Under [Assumption 2.1](#), if [Assumption B](#) or [Assumption C](#) holds, then h_g may be chosen as*

$$h_g = C_X^\dagger C_{XY} g. \tag{4.1}$$

More precisely, if [Assumption C](#) holds, then $\text{Cov}[(C_X^\dagger C_{XY} g)(X) - f_g(X), h(X)] = 0$ for all $h \in \mathcal{H}$ and $g \in \mathcal{G}$; and if [Assumption B](#) holds, or even just $f_g \in \mathcal{H}_{\mathcal{C}}$ for some $g \in \mathcal{G}$, then there exists a constant $c_g \in \mathbb{R}$ such that \mathbb{P}_X -almost everywhere $f_g = c_g + C_X^\dagger C_{XY} g$.

Proof. By [Theorem 4.1](#), (4.1) is well defined. Under [Assumption C](#), for all $g \in \mathcal{G}$ and $h \in \mathcal{H}$, and appealing to [Theorem 4.1](#) and [Lemma A.6](#),

$$\text{Cov}[h(X), (C_X^\dagger C_{XY} g)(X)] = \langle h, C_X C_X^\dagger C_{XY} g \rangle_{\mathcal{H}} = \langle h, C_{XY} g \rangle_{\mathcal{H}} = \text{Cov}[h(X), f_g(X)].$$

If $f_g \in \mathcal{H}_{\mathcal{C}}$ for some $g \in \mathcal{G}$, then there exists a function $h'_g \in \mathcal{H}$ and a constant $c'_g \in \mathbb{R}$ such that, \mathbb{P}_X -a.e. in \mathcal{X} , $h'_g = f_g - c'_g$. [Theorem 4.1](#) implies that $C_X h'_g = C_{XY} g$, and so [Lemma A.5](#) implies that $h'_g - C_X^\dagger C_{XY} g$ is constant \mathbb{P}_X -a.e. Hence, $f_g - C_X^\dagger C_{XY} g$ is constant \mathbb{P}_X -a.e. ■

We now give our first main result, the rigorous statement of the CME formula for centred (cross-)covariance operators. In fact, we give two results: a “weak” result (4.2) under [Assumption C](#) in which the CME, as a function on \mathcal{X} , holds only when tested against elements of \mathcal{H} in the $L^2(\mathbb{P}_X)$ inner product, and a “strong” almost-sure equality in \mathcal{G} (4.3) under [Assumption B](#).

¹⁰Furthermore, by [Lemma A.3](#), this operator is actually Hilbert–Schmidt when thought of as an operator taking values in the appropriate L^2 space rather than in the RKHS.

Theorem 4.3 (Centred CME). *Under Assumptions 2.1 and C, $C_X^\dagger C_{XY} : \mathcal{G} \rightarrow \mathcal{H}$ is a bounded¹⁰ operator and, for all $y \in \mathcal{Y}$ and $h \in \mathcal{H}$,*

$$\langle h, \mu_{Y|X=\cdot}(y) \rangle_{L^2(\mathbb{P}_X)} = \left\langle h, (\mu_Y + (C_X^\dagger C_{XY})^* (\varphi(\cdot) - \mu_X))(y) \right\rangle_{L^2(\mathbb{P}_X)}. \quad (4.2)$$

Suppose in addition that any of the following four conditions holds:

- (i) the kernel k is characteristic;
- (ii) \mathcal{H}_C is dense in $L_C^2(\mathbb{P}_X)$;
- (iii) Assumption B holds;
- (iv) $f_{\psi(y)} \in \mathcal{H}_C$ for each $y \in \mathcal{Y}$.

Then, for \mathbb{P}_X -a.e. $x \in \mathcal{X}$,

$$\mu_{Y|X=x} = \mu_Y + (C_X^\dagger C_{XY})^* (\varphi(x) - \mu_X). \quad (4.3)$$

Proof. Theorems 4.1 and A.1 imply that $C_X^\dagger C_{XY}$ is well defined and bounded¹⁰ and that, for each $g \in \mathcal{G}$, we may choose the function $h_g \in \mathcal{H}$ in Assumptions B and C to be $h_g = C_X^\dagger C_{XY} g$ (by Proposition 4.2). Now (2.6), Lemma A.7, and the definition of c_g (see Assumption C) yield that, for $x \in \mathcal{X}$ and $y \in \mathcal{Y}$,

$$h_{\psi(y)}(x) + c_{\psi(y)} = (\mu_Y + (C_X^\dagger C_{XY})^* (\varphi(x) - \mu_X))(y). \quad (4.4)$$

This yields (4.2) for each $h \in \mathcal{H}$ via

$$\begin{aligned} & \langle h, (\mu_{Y|X=\cdot} - \mu_Y - (C_X^\dagger C_{XY})^* (\varphi(\cdot) - \mu_X))(y) \rangle_{L^2(\mathbb{P}_X)} \\ &= \langle h, f_{\psi(y)} - h_{\psi(y)} - c_{\psi(y)} \rangle_{L^2(\mathbb{P}_X)} \\ &= \underbrace{\text{Cov}[h(X), (f_{\psi(y)} - h_{\psi(y)})(X)]}_{=0} + \mathbb{E}[h(X)] \underbrace{(\mathbb{E}[(f_{\psi(y)} - h_{\psi(y)})(X)] - c_{\psi(y)})}_{=0} = 0. \end{aligned}$$

If (i) or (ii) holds (note that, by Lemma A.4, (i) \implies (ii)), then (4.3) follows directly. If (iii) or (iv) holds (with $f_{\psi(y)} = h_{\psi(y)} + c_{\psi(y)}$, $h_{\psi(y)} \in \mathcal{H}$, $c_{\psi(y)} \in \mathbb{R}$), then (4.3) can be obtained from

$$\begin{aligned} \mu_{Y|X=x}(y) &= \mathbb{E}[\ell(y, Y)|X=x] = f_{\psi(y)}(x) \stackrel{(*)}{=} h_{\psi(y)}(x) + c_{\psi(y)} \\ &= (\mu_Y + (C_X^\dagger C_{XY})^* (\varphi(x) - \mu_X))(y), \end{aligned}$$

where all equalities hold for \mathbb{P}_X -a.e. $x \in \mathcal{X}$ and the last equality follows from (4.4) (note that we might be arguing with two different choices of $h_{\psi(y)}$, which we may assume to agree by Proposition 4.2). \blacksquare

Note that step (*) in the proof of Theorem 4.3 genuinely requires condition (iv) (which follows from Assumption B), and Assumption C alone does not suffice. Again we see that \mathcal{H} needs to be rich enough. The reason that we get (4.2) in terms of the inner product of $L^2(\mathbb{P}_X)$, and not its weaker version in $L_C^2(\mathbb{P}_X)$, is that we took care of the shifting constant $c_g := \mathbb{E}[f_g(X) - h_g(X)]$.

Motivated by the theory of Gaussian conditioning in Hilbert spaces (Owhadi and Scovel, 2018) presented in Section 6 and Theorem 6.2 in particular, we hope to generalise CMEs to the case where $\text{ran } C_{XY} \subseteq \text{ran } C_X$ (i.e., by Theorem 4.1, Assumption C) does not necessarily hold. As mentioned above, this will require us to work with certain finite-rank approximations of the operators C_X and C_{XY} . We are still going to need some assumption that guarantees that \mathcal{H} is rich enough to be able to perform the conditioning process in the RKHSs. For this purpose Assumption B will be replaced by its weaker version B*.

Theorem 4.4 (Centred CME under finite-rank approximation). *Let [Assumption 2.1](#) hold. Further, let $(h_n)_{n \in \mathbb{N}}$ be a complete orthonormal system of \mathcal{H} that is an eigenbasis of C_X , let $\mathcal{H}^{(n)} := \text{span}\{h_1, \dots, h_n\}$, let $\mathcal{F} := \mathcal{G} \oplus \mathcal{H}$, let $P^{(n)}: \mathcal{F} \rightarrow \mathcal{F}$ be the orthogonal projection onto $\mathcal{G} \oplus \mathcal{H}^{(n)}$, and let*

$$C := \begin{pmatrix} C_Y & C_{YX} \\ C_{XY} & C_X \end{pmatrix}, \quad C^{(n)} := P^{(n)} C P^{(n)} = \begin{pmatrix} C_Y & C_{YX}^{(n)} \\ C_{XY}^{(n)} & C_X^{(n)} \end{pmatrix}.$$

Then $\text{ran } C_{XY}^{(n)} \subseteq \text{ran } C_X^{(n)}$ and therefore $h_g^{(n)} := C_X^{(n)\dagger} C_{XY}^{(n)} g \in \mathcal{H}$ is well defined for each $g \in \mathcal{G}$. For each $y \in \mathcal{Y}$ and $h \in \mathcal{H}$,

$$\langle h, \mu_{Y|X=\cdot}(y) \rangle_{L^2(\mathbb{P}_X)} = \lim_{n \rightarrow \infty} \langle h, \mu^{(n)}(\cdot, y) \rangle_{L^2(\mathbb{P}_X)}, \quad (4.5)$$

where, for $x \in \mathcal{X}$ and $y \in \mathcal{Y}$,

$$\mu^{(n)}(x, y) := (\mu_Y + (C_X^{(n)\dagger} C_{XY}^{(n)})^* (\varphi(x) - \mu_X))(y).$$

Suppose in addition that any of the following four conditions holds:

- (i) the kernel k is characteristic;*
- (ii) \mathcal{H}_C is dense in $L_C^2(\mathbb{P}_X)$;*
- (iii) [Assumption B*](#) holds;*
- (iv) $f_{\psi(y)} \in \overline{\mathcal{H}_C}^{L_C^2}$ for each $y \in \mathcal{Y}$.*

Then, as $n \rightarrow \infty$,

$$\|\mu^{(n)}(X, \cdot) - \mu_{Y|X}\|_{L^2(\mathbb{P}; \mathcal{G})} \rightarrow 0, \quad \|\mu_{Y|X=x} - \mu^{(n)}(x, \cdot)\|_{\mathcal{G}} \rightarrow 0 \text{ for } \mathbb{P}_X\text{-a.e. } x \in \mathcal{X}. \quad (4.6)$$

Proof. Note that, since C is a trace-class operator, so is $C^{(n)}$. Furthermore, by [Baker \(1973, Theorem 1\)](#), $C_{XY}^{(n)} = (C_X^{(n)})^{1/2} V C_Y^{1/2}$ for some bounded operator $V: \mathcal{G} \rightarrow \mathcal{H}$. Since $C_X^{(n)}$ has finite rank, this implies that $\text{ran } C_{XY}^{(n)} \subseteq \text{ran } C_X^{(n)}$. Similarly to the proof of [Theorem 4.3](#), we define $c_g^{(n)} := \mathbb{E}[(f_g - h_g^{(n)})(X)]$ for $g \in \mathcal{G}, n \in \mathbb{N}$ and obtain by [\(2.6\)](#) and [Lemma A.7](#) for $x \in \mathcal{X}, y \in \mathcal{Y}$ and $n \in \mathbb{N}$ that

$$h_{\psi(y)}^{(n)}(x) + c_{\psi(y)}^{(n)} = \mu^{(n)}(x, y). \quad (4.7)$$

Identity [\(4.5\)](#) can be obtained similarly to [\(4.2\)](#) except that we also need to show that $\text{Cov}[h(X), f_g(X)] = \lim_{n \rightarrow \infty} \text{Cov}[h(X), h_g^{(n)}(X)]$ for all $h \in \mathcal{H}$, as proved in [Lemma A.9\(a\)](#).

To establish [\(4.6\)](#), we first note that, by [Lemma A.9\(b\)](#), for all $g \in \mathcal{G}$, $[h_g^{(n)}]$ is the L_C^2 -orthogonal projection of $[f_g]$ onto $\mathcal{H}_C^{(n)}$. Now let $y \in \mathcal{Y}$ and $U := \bigcup_{n \in \mathbb{N}} \mathcal{H}_C^{(n)}$. Note that, by [Lemma A.4](#), (i) \implies (ii) \implies (iii) \implies (iv), so let us assume (iv). Since $\overline{U}^{\mathcal{H}_C} = \mathcal{H}_C$ and $[f_{\psi(y)}] \in \overline{\mathcal{H}_C}^{L_C^2}$ by assumption, and since [\(2.2\)](#) implies that $\|\cdot\|_{\mathcal{H}}$ is a stronger norm than $\|\cdot\|_{L^2}$, we also have $[f_{\psi(y)}] \in \overline{U}^{L_C^2}$ and [Lemma A.8](#) implies

$$\|[h_{\psi(y)}^{(n)}] - [f_{\psi(y)}]\|_{L_C^2} \xrightarrow{n \rightarrow \infty} 0. \quad (4.8)$$

For $x \in \mathcal{X}$ and $n \in \mathbb{N}$ let $\mathbf{m}^{(n)}(x) := h_{\psi(\cdot)}^{(n)}(x) = (C_X^{(n)\dagger} C_{XY}^{(n)})^* \varphi(x) \in \mathcal{G}$ and $\mathbf{m}(x) := f_{\psi(\cdot)}(x) = \mu_{Y|X=x} \in \mathcal{G}$. Then $\mathbf{m}^{(n)}, \mathbf{m} \in L^2(\mathbb{P}_X; \mathcal{G})$ by [\(2.1\)](#), since

$$\begin{aligned} \|\mathbf{m}^{(n)}\|_{L^2(\mathbb{P}_X; \mathcal{G})}^2 &= \mathbb{E}[\|(C_X^{(n)\dagger} C_{XY}^{(n)})^* \varphi(X)\|_{\mathcal{G}}^2] \leq \|(C_X^{(n)\dagger} C_{XY}^{(n)})^*\| \mathbb{E}[\|\varphi(X)\|_{\mathcal{H}}^2] < \infty, \\ \|\mathbf{m}\|_{L^2(\mathbb{P}_X; \mathcal{G})}^2 &= \mathbb{E}[\|\mathbb{E}[\psi(Y)|X]\|_{\mathcal{G}}^2] \leq \mathbb{E}[\mathbb{E}[\|\psi(Y)\|_{\mathcal{G}}^2|X]] = \mathbb{E}[\|\psi(Y)\|_{\mathcal{G}}^2] < \infty. \end{aligned}$$

So far, we have shown that, for each $y \in \mathcal{Y}$,

- $([\mathbf{m}^{(n)}](\cdot))(y)$ is the $L^2_{\mathcal{G}}(\mathbb{P}_X)$ -orthogonal projection of $([\mathbf{m}](\cdot))(y)$ onto $\mathcal{H}_{\mathcal{G}}^{(n)}$;
- $([\mathbf{m}^{(n)}](\cdot))(y) \rightarrow ([\mathbf{m}](\cdot))(y)$ in $L^2_{\mathcal{G}}(\mathbb{P}_X)$ as $n \rightarrow \infty$.

Hence, by [Lemma A.10\(a\)](#) and [\(b\)](#),

$$\|(\mathbf{m}^{(n)}(X) - \mathbb{E}[\mathbf{m}^{(n)}(X)]) - (\mathbf{m}(X) - \mathbb{E}[\mathbf{m}(X)])\|_{L^2(\mathbb{P}; \mathcal{G})} \xrightarrow{n \rightarrow \infty} 0. \quad (4.9)$$

Therefore, by [\(4.7\)](#) and the definition of $c_g^{(n)}$, $\mu^{(n)}(X, \cdot)$ converges to $\mu_{Y|X} = f_{\psi(\cdot)}(X) = \mathbf{m}(X)$ in $L^p(\mathbb{P}; \mathcal{G})$ for $p = 2$ and, since \mathbb{P} is a finite measure, also for $p = 1$. By [Lemma A.11](#), $(\mu^{(n)}(X, \cdot))_{n \in \mathbb{N}}$ is a martingale, and so [Diestel and Uhl \(1977, Theorem V.2.8\)](#) implies that this convergence even holds a.e., i.e. $\mu^{(n)}(x, \cdot)$ converges in \mathcal{G} to $\mu_{Y|X=x}$ for \mathbb{P}_X -a.e. $x \in \mathcal{X}$. ■

5. Theory for Uncentred Operators

Beginning with the work of [Song et al. \(2010a,b\)](#), uncentred (cross-)covariance operators became more commonly used than centred ones. This section shows how similar results to those of [Section 4](#) can be obtained for uncentred operators. Roughly speaking, the same conclusions can be made as in [Theorem 4.3](#) but under [Assumption A](#) in place of [B](#), while only weaker statements than in [Theorem 4.4](#) can be obtained in [Theorem 5.4](#) (no \mathbb{P}_X -a.e. convergence, see below) and again under the stronger [Assumption A*](#) in place of [B*](#). This observation suggests that centred operators are superior to uncentred ones in terms of generality. So far, the theoretical justification for CME using uncentred operators relies on [Fukumizu et al. \(2013, Theorems 1 and 2\)](#), which require rather strong assumptions. Our improvement can be summarised as follows:

- Since we use ${}^u C_X^\dagger$ instead of ${}^u C_X^{-1}$ our theory can cope with non-injective operators ${}^u C_X$. This is only a minor advance, since ${}^u C_X$ is injective under rather mild conditions on X and k (see [Fukumizu et al. \(2013, Footnote 3\)](#)).
- In contrast to [Fukumizu et al. \(2013, Theorem 2\)](#), we do not require the assumption that $\varphi(x)$ lies in the range of ${}^u C_X$. The reason for this is that the operator $({}^u C_X^\dagger {}^u C_{XY})^*$ in [\(4.3\)](#) is globally defined whereas ${}^u C_{YX} {}^u C_X^{-1}$ is not. This is an important improvement since the assumption that $\varphi(x) \in \text{ran } {}^u C_X$ is typically hard to verify (see [Footnote 4](#)).
- We state a version of the CME formula under [Assumption A*](#), which is a verifiable condition since it follows from the kernel k being L^2 -universal.
- As explained in [Remark 1.1](#), the condition in [Fukumizu et al. \(2013, Theorem 2\)](#) on $\mathbb{E}[g(Y)|X = \cdot]$ to lie in \mathcal{H} for each $g \in \mathcal{G}$ is ill posed, since these functions are uniquely defined only \mathbb{P}_X -a.e. However, in our case, [Assumption 2.1\(f\)](#) ensures that [Assumptions A](#) and [A*](#) are unambiguous.

As mentioned above, using centred operators instead of uncentred ones yields the important advantage of requiring only the weaker [Assumption B](#) in place of [A](#) or [Assumption B*](#) in place of [A*](#), respectively. Further, [Theorem 5.4](#) provides weaker statements than its centred analogue, [Theorem 4.4](#): we show only convergence in $L^2(\mathbb{P}_X; \mathcal{G})$, which does not guarantee convergence for \mathbb{P}_X -a.e. $x \in \mathcal{X}$.

Theorem 5.1. *Under [Assumption 2.1](#), the following statements are equivalent:*

- [Assumption \${}^u C\$](#) holds.*
- For each $g \in \mathcal{G}$ there exists $h_g \in \mathcal{H}$ such that ${}^u C_X h_g = {}^u C_{XY} g$.*
- $\text{ran } {}^u C_{XY} \subseteq \text{ran } {}^u C_X$.*

Proof. The proof is identical to that of [Theorem 4.1](#) (apart from using uncentred covariance operators in place of centred ones). ■

Similar to [Proposition 4.2](#), the element $h_g \in \mathcal{H}$ in [Assumption \${}^u\mathbf{C}\$](#) can always be chosen as $h_g = {}^u C_X^\dagger {}^u C_{XY} g$.

Proposition 5.2. *Let [Assumption 2.1](#) hold. Under [Assumption \${}^u\mathbf{C}\$](#) , h_g may be chosen as*

$$h_g = {}^u C_X^\dagger {}^u C_{XY} g. \quad (5.1)$$

More precisely, ${}^u \text{Cov}[({}^u C_X^\dagger {}^u C_{XY} g)(X) - f_g(X), h(X)] = 0$ for all $h \in \mathcal{H}$ and $g \in \mathcal{G}$. If $f_g \in \mathcal{H}$ for some $g \in \mathcal{G}$, then the identity $f_g = {}^u C_X^\dagger {}^u C_{XY} g$ holds \mathbb{P}_X -a.e.

Proof. By [Theorem 5.1](#), (5.1) is well defined. If [Assumption \${}^u\mathbf{C}\$](#) holds, then, by [Theorem 5.1](#) and [Lemma A.6](#), for all $g \in \mathcal{G}$ and $h \in \mathcal{H}$,

$$\begin{aligned} {}^u \text{Cov}[h(X), ({}^u C_X^\dagger {}^u C_{XY} g)(X)] &= \langle h, {}^u C_X {}^u C_X^\dagger {}^u C_{XY} g \rangle_{\mathcal{H}} \\ &= \langle h, {}^u C_{XY} g \rangle_{\mathcal{H}} = {}^u \text{Cov}[h(X), f_g(X)]. \end{aligned}$$

If $f_g \in \mathcal{H}$ holds for some $g \in \mathcal{G}$, then [Lemma A.6](#) implies that ${}^u C_X f_g = {}^u C_{XY} g$ for all $g \in \mathcal{G}$ and the claim follows from [Lemma A.5](#). \blacksquare

Let us now formulate and prove the analogues of [Theorems 4.3](#) and [4.4](#) for uncentred operators.

Theorem 5.3 (Uncentred CME). *Under [Assumptions 2.1](#) and [Assumption \${}^u\mathbf{C}\$](#) , the linear operator ${}^u C_X^\dagger {}^u C_{XY} : \mathcal{G} \rightarrow \mathcal{H}$ is bounded¹⁰ and, for all $y \in \mathcal{Y}$ and $h \in \mathcal{H}$,*

$$\langle h, \mu_{Y|X=\cdot}(y) \rangle_{L^2(\mathbb{P}_X)} = \left\langle h, ({}^u C_X^\dagger {}^u C_{XY})^* \varphi(\cdot)(y) \right\rangle_{L^2(\mathbb{P}_X)}. \quad (5.2)$$

Suppose in addition that any of the following four conditions holds:

- (i) the kernel k is L^2 -universal;
- (ii) \mathcal{H} is dense in $L^2(\mathbb{P}_X)$;
- (iii) [Assumption A](#) holds;
- (iv) $f_{\psi(y)} \in \mathcal{H}$ for each $y \in \mathcal{Y}$.

Then, for \mathbb{P}_X -a.e. $x \in \mathcal{X}$,

$$\mu_{Y|X=x} = ({}^u C_X^\dagger {}^u C_{XY})^* \varphi(x). \quad (5.3)$$

Proof. First note that, by [Theorems 5.1](#) and [A.1](#), ${}^u C_X^\dagger {}^u C_{XY}$ is well defined and bounded¹⁰ and that for each $g \in \mathcal{G}$ we may choose the function $h_g \in \mathcal{H}$ in [Assumption \${}^u\mathbf{C}\$](#) as $h_g = {}^u C_X^\dagger {}^u C_{XY} g$ by [Proposition 5.2](#). By [Lemma A.7](#) we obtain, for all $x \in \mathcal{X}$ and $y \in \mathcal{Y}$,

$$h_{\psi(y)}(x) = (({}^u C_X^\dagger {}^u C_{XY})^* \varphi(x))(y).$$

This yields (5.2) via

$$\begin{aligned} \left\langle h, (\mu_{Y|X=\cdot} - ({}^u C_X^\dagger {}^u C_{XY})^* \varphi(\cdot))(y) \right\rangle_{L^2(\mathbb{P}_X)} &= \langle h, f_{\psi(y)} - h_{\psi(y)} \rangle_{L^2(\mathbb{P}_X)} \\ &= {}^u \text{Cov}[h(X), (f_{\psi(y)} - h_{\psi(y)})(X)] = 0, \end{aligned}$$

which implies (5.3) under any of the four conditions stated in the theorem (possibly using [Proposition 5.2](#)). \blacksquare

Theorem 5.4 (Uncentred CME under finite-rank approximation). *Let [Assumption 2.1](#) hold. Further, let $(h_n)_{n \in \mathbb{N}}$ be a complete orthonormal system of \mathcal{H} that is an eigenbasis of C_X , let $\mathcal{H}^{(n)} := \text{span}\{h_1, \dots, h_n\}$, let $\mathcal{F} := \mathcal{G} \oplus \mathcal{H}$, let $P^{(n)}: \mathcal{F} \rightarrow \mathcal{F}$ be the orthogonal projection onto $\mathcal{G} \oplus \mathcal{H}^{(n)}$, and let*

$${}^u C := \begin{pmatrix} {}^u C_Y & {}^u C_{YX} \\ {}^u C_{XY} & {}^u C_X \end{pmatrix}, \quad {}^u C^{(n)} := P^{(n)} {}^u C P^{(n)} = \begin{pmatrix} {}^u C_Y & {}^u C_{YX}^{(n)} \\ {}^u C_{XY}^{(n)} & {}^u C_X^{(n)} \end{pmatrix}.$$

Then $\text{ran } {}^u C_{XY}^{(n)} \subseteq \text{ran } {}^u C_X^{(n)}$ and therefore ${}^u h_g^{(n)} := {}^u C_X^{(n)\dagger} {}^u C_{XY}^{(n)} g \in \mathcal{H}$ is well defined for each $g \in \mathcal{G}$. For each $y \in \mathcal{Y}$ and $h \in \mathcal{H}$,

$$\langle h, \mu_{Y|X=\cdot}(y) \rangle_{L^2(\mathbb{P}_X)} = \lim_{n \rightarrow \infty} \langle h, \mu^{(n)}(\cdot, y) \rangle_{L^2(\mathbb{P}_X)}, \quad (5.4)$$

where, for $x \in \mathcal{X}$ and $y \in \mathcal{Y}$,

$${}^u \mu^{(n)}(x, y) := (({}^u C_X^{(n)\dagger} {}^u C_{XY}^{(n)})^* \varphi(x))(y). \quad (5.5)$$

Suppose in addition that any of the following four conditions holds:

- (i) the kernel k is L^2 -universal;*
- (ii) \mathcal{H} is dense in $L^2(\mathbb{P}_X)$;*
- (iii) [Assumption A*](#) holds;*
- (iv) $f_{\psi(y)} \in \overline{\mathcal{H}}^{L^2}$ for each $y \in \mathcal{Y}$.*

Then

$$\|{}^u \mu^{(n)}(X, \cdot) - \mu_{Y|X}\|_{L^2(\mathbb{P}; \mathcal{G})} \xrightarrow{n \rightarrow \infty} 0. \quad (5.6)$$

Proof. The proof goes analogously to the one of [Theorem 4.4](#) up to equation (4.9), using uncentred operators instead of centred ones and the statements (c), (d) instead of (a), (b) of [Lemmas A.9](#) and [A.10](#). However, we cannot draw the final conclusion of convergence almost everywhere since we do not have the martingale property, which is provided by [Lemma A.11](#) for the centred case. Note that our proof relies on [Baker \(1973, Theorem 1\)](#) which, strictly speaking, only treats the centred case, but its uncentred version can be proven similarly. ■

Corollary 5.5. *Under the assumptions of [Theorem 5.3](#) (including either of the additional ones),*

$$\mu_Y = ({}^u C_X^\dagger {}^u C_{XY})^* \mu_X.$$

Under the assumptions of [Theorem 5.4](#) (including either of the additional ones),

$$\|\mu_Y - ({}^u C_X^{(n)\dagger} {}^u C_{XY}^{(n)})^* \mu_X\|_{\mathcal{G}} \xrightarrow{n \rightarrow \infty} 0.$$

Proof. As stated in [Theorem 5.3](#), ${}^u C_X^\dagger {}^u C_{XY}$ is a well-defined and bounded¹⁰ linear operator. Hence, by the law of total expectation and [Theorem 5.3](#),

$$\mu_Y = \mathbb{E}[\mu_{Y|X}] = \mathbb{E}[({}^u C_X^\dagger {}^u C_{XY})^* \varphi(X)] = ({}^u C_X^\dagger {}^u C_{XY})^* \mathbb{E}[\varphi(X)] = ({}^u C_X^\dagger {}^u C_{XY})^* \mu_X,$$

proving the first claim. The second one follows from Jensen's inequality and [Theorem 5.4](#) via

$$\begin{aligned} \|\mu_Y - ({}^u C_X^{(n)\dagger} {}^u C_{XY}^{(n)})^* \mu_X\|_{\mathcal{G}}^2 &= \|\mathbb{E}[\mu_{Y|X}] - ({}^u C_X^{(n)\dagger} {}^u C_{XY}^{(n)})^* \mathbb{E}[\varphi(X)]\|_{\mathcal{G}}^2 \\ &= \|\mathbb{E}[\mu_{Y|X} - ({}^u C_X^{(n)\dagger} {}^u C_{XY}^{(n)})^* \varphi(X)]\|_{\mathcal{G}}^2 \\ &\leq \|\mu_{Y|X} - ({}^u C_X^{(n)\dagger} {}^u C_{XY}^{(n)})^* \varphi(X)\|_{L^2(\mathbb{P}_X; \mathcal{G})}^2 \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

■

6. Gaussian Conditioning in Hilbert spaces

This section gives a review of conditioning theory for Gaussian random variables in separable Hilbert spaces, summarising the work of [Owhadi and Scovel \(2018\)](#). Our only somewhat novel contribution here is the explicit characterisation of the essential operator $\widehat{Q}_{C,\mathcal{H}}$ in terms of the Moore–Penrose pseudo-inverse, which appears as an exercise for the reader in [Arias et al. \(2008, Remark 2.3\)](#).

In the following let $\mathcal{F} = \mathcal{G} \oplus \mathcal{H}$ be the sum of two separable Hilbert spaces \mathcal{G} and \mathcal{H} and let (U, V) be an \mathcal{F} -valued jointly Gaussian random variable with mean $\mu \in \mathcal{F}$ and covariance operator $C: \mathcal{F} \rightarrow \mathcal{F}$ given by the following block structures:

$$\begin{pmatrix} U \\ V \end{pmatrix} \sim \mathcal{N}(\mu, C), \quad \mu = \begin{pmatrix} \mu_U \\ \mu_V \end{pmatrix}, \quad C = \begin{pmatrix} C_U & C_{UV} \\ C_{VU} & C_V \end{pmatrix} \geq 0$$

with $\mu_U \in \mathcal{G}$, etc. We denote by $L(\mathcal{F})$ the Banach algebra of bounded linear operators on \mathcal{F} and by $L_+(\mathcal{F}) = \{A \in L(\mathcal{F}) \mid A \geq 0\}$ the set of positive operators, i.e. those self-adjoint operators A for which $\langle x, Ax \rangle \geq 0$ for all $x \in \mathcal{F}$. The theory of Gaussian conditioning relies on the concept of so-called *oblique projections*:

Definition 6.1. Let $\mathcal{F} = \mathcal{G} \oplus \mathcal{H}$ be a direct sum of two Hilbert spaces \mathcal{G} and \mathcal{H} and let $C \in L_+(\mathcal{F})$ be a positive operator. The set of (C -symmetric) *oblique projections* onto \mathcal{H} is given by

$$\mathcal{P}(C, \mathcal{H}) = \{Q \in L(\mathcal{F}) \mid Q^2 = Q, \text{ ran } Q = \mathcal{H}, CQ = Q^*C\}.$$

The pair (C, \mathcal{H}) is said to be *compatible* if $\mathcal{P}(C, \mathcal{H})$ is non-empty.

The first two conditions $Q^2 = Q$ and $\text{ran } Q = \mathcal{H}$ imply that Q has the block structure

$$Q = \begin{pmatrix} 0 & 0 \\ \widehat{Q} & \text{Id}_{\mathcal{H}} \end{pmatrix}, \quad \widehat{Q}: \mathcal{G} \rightarrow \mathcal{H}. \quad (6.1)$$

Then, the condition $CQ = Q^*C$ is equivalent to $C_V \widehat{Q} = C_{VU}$ (which follows from a straightforward blockwise multiplication, see [Lemma 6.3](#)) and implies in particular $\text{ran } C_{VU} \subseteq \text{ran } C_V$. The other way round, as we will see later on, the condition $\text{ran } C_{VU} \subseteq \text{ran } C_V$ guarantees the existence of an oblique projection $Q \in \mathcal{P}(C, \mathcal{H})$ and will provide a crucial link between the theory of Gaussian conditioning and conditional mean embeddings in [Section 7](#).

The results on conditioning Gaussian measures can then be summarised as follows:

Theorem 6.2 ([Owhadi and Scovel, 2018](#), Theorem 3.3, Corollary 3.4). *If (C, \mathcal{H}) is compatible, then conditioning U on $V = v \in \mathcal{H}$ results in a Gaussian random variable on \mathcal{G} with mean $\mu_{U|V=v}$ and covariance operator $C_{U|V=v}$ given by*

$$\begin{cases} \mu_{U|V=v} = \mu_U + \widehat{Q}^*(v - \mu_V), \\ C_{U|V=v} = C_U - C_{UV} \widehat{Q} \end{cases} \quad (6.2)$$

for any oblique projection $Q \in \mathcal{P}(C, \mathcal{H})$ given in the form (6.1). Also, in this case, $\mathcal{P}(C, \mathcal{H})$ contains a unique element

$$Q_{C,\mathcal{H}} = \begin{pmatrix} 0 & 0 \\ \widehat{Q}_{C,\mathcal{H}} & \text{Id}_{\mathcal{H}} \end{pmatrix}$$

that fulfils the properties (6.4) defined below.

If (C, \mathcal{H}) is incompatible, then conditioning U on $V = v \in \mathcal{H}$ still yields a Gaussian random variable on \mathcal{G} , but the corresponding formulae for the conditional mean $\mu_{U|V=v}$ and covariance

operator $C_{U|V=v}$ are given by a limiting process using finite-rank approximations of C in the following way. Let $(h_n)_{n \in \mathbb{N}}$ be a complete orthonormal system of \mathcal{H} , $P^{(n)}: \mathcal{F} \rightarrow \mathcal{F}$ denote the orthogonal projection onto $\mathcal{G} \oplus \text{span}\{h_1, \dots, h_n\}$ and $C^{(n)} = P^{(n)}CP^{(n)}$. Then $(C^{(n)}, \mathcal{H})$ is compatible for each $n \in \mathbb{N}$ and, for \mathbb{P}_V -a.e. $v \in \mathcal{H}$ (with \mathbb{P}_V denoting the distribution of V),

$$\begin{cases} \mu_{U|V=v} = \mu_U + \lim_{n \rightarrow \infty} \hat{Q}_{C^{(n)}, \mathcal{H}}^*(v - \mu_V), \\ C_{U|V=v} = C_U - \lim_{n \rightarrow \infty} C_{UV} \hat{Q}_{C^{(n)}, \mathcal{H}}, \end{cases} \quad (6.3)$$

where the second limit is in the trace norm.

In the following we will revisit some theory on oblique projections which will be necessary to establish the connection between Gaussian conditioning and conditional mean embeddings. We will also characterise the special oblique projection $Q_{C, \mathcal{H}} \in \mathcal{P}(C, \mathcal{H})$ by means of the Moore–Penrose pseudo-inverse.

Lemma 6.3. *If $\hat{Q}: \mathcal{G} \rightarrow \mathcal{H}$ is a bounded linear operator such that $C_V \hat{Q} = C_{VU}$, then*

$$Q = \begin{pmatrix} 0 & 0 \\ \hat{Q} & \text{Id}_{\mathcal{H}} \end{pmatrix} \in \mathcal{P}(C, \mathcal{H}).$$

In particular, the pair (C, \mathcal{H}) is compatible.

Proof. The properties $Q^2 = Q$ and $\text{ran } Q = \mathcal{H}$ are clear from the definition of Q and a straightforward blockwise multiplication shows that $CQ = Q^*C$. \blacksquare

Proposition 6.4. *In the setup of Definition 6.1, if (C, \mathcal{H}) is compatible, then there exists a unique bounded operator $\hat{Q}_{C, \mathcal{H}}: \mathcal{G} \rightarrow \mathcal{H}$ such that*

$$C_V \hat{Q}_{C, \mathcal{H}} = C_{VU}, \quad \ker \hat{Q}_{C, \mathcal{H}} = \ker C_{VU}, \quad \text{ran } \hat{Q}_{C, \mathcal{H}} \subseteq \overline{\text{ran } C_V}. \quad (6.4)$$

By Lemma 6.3 the first property implies that

$$Q_{C, \mathcal{H}} = \begin{pmatrix} 0 & 0 \\ \hat{Q}_{C, \mathcal{H}} & \text{Id}_{\mathcal{H}} \end{pmatrix} \in \mathcal{P}(C, \mathcal{H}).$$

Proof. See Douglas (1966, Theorem 1) or Fillmore and Williams (1971, Theorem 2.1) for the existence and uniqueness of $\hat{Q}_{C, \mathcal{H}}$ and Corach et al. (2001) or Owhadi and Scovel (2018) for its connection to oblique projections. \blacksquare

If one follows the original construction of Douglas (1966, Theorem 1) or Fillmore and Williams (1971, Theorem 2.1), it is easy to see how this unique element can be characterised in terms of the Moore–Penrose pseudo-inverse C_V^\dagger of C_V :

Theorem 6.5. *If $\text{ran } C_{VU} \subseteq \text{ran } C_V$, then $\hat{Q} = C_V^\dagger C_{VU}: \mathcal{G} \rightarrow \mathcal{H}$ is a well-defined bounded operator which uniquely fulfils the conditions (6.4).*

Proof. This is a direct application of Theorem A.1. \blacksquare

Theorem 6.6. *In the setup of Definition 6.1, the following statements are equivalent:*

- (i) (C, \mathcal{H}) is compatible.
- (ii) $\text{ran } C_{VU} \subseteq \text{ran } C_V$.

If either of these conditions holds, then the unique element $Q_{C,\mathcal{H}} \in \mathcal{P}(C, \mathcal{H})$ in [Proposition 6.4](#) is given by

$$\widehat{Q}_{C,\mathcal{H}} = C_V^\dagger C_{VU}. \quad (6.5)$$

Proof. If (C, \mathcal{H}) is compatible, there exists an element $\widehat{Q}_{C,\mathcal{H}}: \mathcal{G} \rightarrow \mathcal{H}$ with $C_V \widehat{Q}_{C,\mathcal{H}} = C_{VU}$ by [Proposition 6.4](#), which implies (ii). If $\text{ran } C_{VU} \subseteq \text{ran } C_V$, then [Theorem 6.5](#) and [Lemma 6.3](#) imply (i). [Theorem 6.5](#) and the uniqueness of $\widehat{Q}_{C,\mathcal{H}}$ in [Proposition 6.4](#) imply (6.5). ■

Remark 6.7. [Lemma 6.3](#) and the equivalence part of [Theorem 6.6](#) were already proved by [Corach et al. \(2001\)](#); we state them for the sake of readability. The second part of [Theorem 6.6\(ii\)](#) characterises the operator $\widehat{Q}_{C,\mathcal{H}}$ in terms of the Moore–Penrose pseudo-inverse, without an assumption of closed range, as anticipated by [Arias et al. \(2008, Remark 2.3\)](#).

There are covariance operators C for which the above conditions do not hold:

Example 6.8. Let $\mathcal{H} = \mathcal{G}$ be any (separable) infinite-dimensional Hilbert space with complete orthonormal basis $(e_j)_{j \in \mathbb{N}}$. Let

$$\begin{aligned} C_U &:= \sum_{j \in \mathbb{N}} j^{-2} e_j \otimes e_j, \\ C_V &:= \sum_{j \in \mathbb{N}} j^{-4} e_j \otimes e_j, \\ C_{VU} = C_{UV} &:= C_U^{1/2} \text{Id}_{\mathcal{H}} C_V^{1/2} = \sum_{j \in \mathbb{N}} j^{-3} e_j \otimes e_j. \end{aligned}$$

By [Baker \(1973, Theorem 2\)](#),

$$C := \begin{pmatrix} C_U & C_{UV} \\ C_{VU} & C_V \end{pmatrix}$$

is a legitimate positive definite covariance operator on $\mathcal{F} = \mathcal{G} \oplus \mathcal{H}$. However,

$$\text{ran } C_{VU} = \left\{ \sum_{j \in \mathbb{N}} \alpha_j e_j \mid (j^3 \alpha_j)_{j \in \mathbb{N}} \in \ell^2 \right\} \not\subseteq \left\{ \sum_{j \in \mathbb{N}} \alpha_j e_j \mid (j^4 \alpha_j)_{j \in \mathbb{N}} \in \ell^2 \right\} = \text{ran } C_V.$$

7. Connection between CME and Gaussian Conditioning

If we compare the theories of CMEs and Gaussian conditioning in Hilbert spaces, we make the following observations:

- Formula [\(4.3\)](#) for CME and formula [\(6.2\)](#) for Gaussian conditioning look very similar (in view of [Theorem 6.6](#)).
- The assumptions under which the conditioning process is “easy” — namely [Assumption C](#) (as long as [Assumption B*](#) holds as well) and the compatibility of (C, \mathcal{H}) — are equivalent to the conditions that $\text{ran } C_{XY} \subseteq \text{ran } C_X$ and $\text{ran } C_{VU} \subseteq \text{ran } C_V$ respectively ([Theorems 4.1](#) and [6.6](#)).

This motivates us to connect these two theories by working in the setup of [Section 2](#) and introducing new jointly Gaussian random variables U and V that take values in the RKHSs \mathcal{G} and \mathcal{H} respectively, where the means μ_U and μ_V and (cross-)covariance operators C_U, C_{UV}, C_{VU} , and C_V are chosen to coincide with the kernel mean embeddings μ_Y and μ_X and the kernel (cross-)covariance operators C_Y, C_{YX}, C_{XY} , and C_X respectively:

$$\begin{pmatrix} U \\ V \end{pmatrix} \sim \mathcal{N}(\mu, C), \quad \mu = \begin{pmatrix} \mu_U \\ \mu_V \end{pmatrix} := \begin{pmatrix} \mu_Y \\ \mu_X \end{pmatrix}, \quad C = \begin{pmatrix} C_U & C_{UV} \\ C_{VU} & C_V \end{pmatrix} := \begin{pmatrix} C_Y & C_{YX} \\ C_{XY} & C_X \end{pmatrix}. \quad (7.1)$$




Figure 7.1: A normally-distributed $\mathcal{G} \oplus \mathcal{H}$ -valued normal random variable (U, V) can be defined with the same mean and covariance structure as $(\psi(Y), \varphi(X))$. While the latter will typically fail to be normally distributed, surprisingly, the conditional means of the two random variables happen to agree! Since $C_{U|V=v}$ does not depend on the realisation v , a specific property of Gaussian random variables that cannot be expected from $C_{Y|X=x}$, a similar agreement for the conditional covariance operators cannot be obtained. Instead, the identity provided by [Theorem 7.1](#) holds, which is open to interpretation.

By [Baker \(1973, Theorem 1\)](#) and since [Assumption 2.1\(e\)](#) implies that C is a trace-class covariance operator, the Gaussian random variable (U, V) is well defined in $\mathcal{G} \oplus \mathcal{H}$. Note that the random variables $W = (U, V)$ and $Z = (\psi(Y), \varphi(X))$ do not coincide even though they have the same mean and covariance operator, since the latter will not generally be Gaussian. Surprisingly, their conditional means agree, as long as we condition on $V = v = \varphi(x)$ and $X = x$ respectively. This is obvious when one compares [\(4.3\)](#) with [\(6.2\)](#) (and [\(4.6\)](#) with [\(6.3\)](#) using [Theorem 6.5](#)). A natural question is whether a similar equality holds for the conditional covariance operator $C_{Y|X=x}$. However, the covariance operator $C_{U|V=v}$ obtained from Gaussian conditioning is independent of v , a special property of Gaussian measures that cannot be expected of the conditional kernel covariance operator $C_{Y|X=x}$. Instead, $C_{U|V=v}$ equals the mean of $C_{Y|X=x}$ when averaged over all possible outcomes $x \in \mathcal{X}$.¹¹ These insights are summarised in the following proposition and illustrated in [Figure 7.1](#).

Note that the distributions of $\varphi(X)$ and V might have different (and even disjoint!) supports, and so one must be particularly careful with “almost every” statements in this context.

Theorem 7.1. *Let [Assumption 2.1](#) and [Assumption B*](#) hold, (U, V) be the random variable defined by [\(7.1\)](#) and let $\mathbb{P}_{\varphi(X)}$ and \mathbb{P}_V denote the probability distributions of $\varphi(X)$ and V , respectively. Then, for \mathbb{P}_V -a.e. $v \in \mathcal{H}$,*

$$C_{U|V=v} = \mathbb{E}[C_{Y|X}] = \int_{\mathcal{X}} C_{Y|X=x} d\mathbb{P}_X(x).$$

Further, there exist $N_1, N_2 \subseteq \Omega$ with $\mathbb{P}_{\varphi(X)}(N_1) = 0$ and $\mathbb{P}_V(N_2) = 0$, such that, for every $v = \varphi(x) \notin N_1 \cup N_2$, $\mu_{U|V=v} = \mu_{Y|X=x}$.

¹¹This observation has already been made by [Fukumizu et al. \(2004a, Proposition 5\)](#) under stronger assumptions and by [Fukumizu et al. \(2009, Proposition 3\)](#) in a weaker form.

Proof. By Lemma A.12, $\mathbb{E}[C_{Y|X}]$ is well defined. The identity $\mu_{U|V=v} = \mu_{Y|X=x}$ for the means follows directly from Theorems 4.4, 6.2, and 6.5. For the covariance identity, using the notation of Theorem 4.4, note that $\|[h_{\psi(y)}^{(n)}] - [f_{\psi(y)}]\|_{L_C^2} \xrightarrow{n \rightarrow \infty} 0$ by (4.8). Therefore, for $y, y' \in \mathcal{Y}$, $g = \psi(y)$, and $g' = \psi(y')$,

$$\begin{aligned} \mathbb{Cov}[f_g(X), f_{g'}(X)] &= \lim_{n \rightarrow \infty} \mathbb{Cov}[h_g^{(n)}(X), h_{g'}^{(n)}(X)] \\ &= \lim_{n \rightarrow \infty} \langle C_X h_g^{(n)}, h_{g'}^{(n)} \rangle_{\mathcal{H}} \\ &= \lim_{n \rightarrow \infty} \langle C_{XY}^{(n)} g, C_X^{(n)\dagger} C_{XY}^{(n)} g' \rangle_{\mathcal{H}} \\ &= \lim_{n \rightarrow \infty} \langle g, C_{YX}^{(n)} C_X^{(n)\dagger} C_{XY}^{(n)} g' \rangle_{\mathcal{G}} \\ &= \lim_{n \rightarrow \infty} \langle g, C_{UV} C_V^{(n)\dagger} C_{VU}^{(n)} g' \rangle_{\mathcal{G}}. \end{aligned}$$

By the law of total covariance and (6.3), (6.5) this implies that, for $g = \psi(y)$ and $g' = \psi(y')$,

$$\begin{aligned} \langle g, \mathbb{E}[C_{Y|X}] g' \rangle_{\mathcal{G}} &= \mathbb{E}[\mathbb{Cov}[g(Y), g'(Y)|X]] \\ &= \mathbb{Cov}[g(Y), g'(Y)] - \mathbb{Cov}[f_g(X), f_{g'}(X)] \\ &= \langle g, C_U g' \rangle_{\mathcal{G}} - \lim_{n \rightarrow \infty} \langle g, C_{UV} C_V^{(n)\dagger} C_{VU}^{(n)} g' \rangle_{\mathcal{G}} \\ &= \langle g, C_{U|V=v} g' \rangle_{\mathcal{G}} \end{aligned}$$

for \mathbb{P}_V -a.e. $v \in \mathcal{H}$. Since $\text{span}\{\psi(y) \mid y \in \mathcal{Y}\}$ is dense in \mathcal{G} , this finishes the proof. \blacksquare

Remark 7.2. Theorem 7.1 implies in particular that the posterior mean $\mu_{U|V=v}$ of the U -component of a jointly Gaussian random variable (U, V) in an RKHS $\mathcal{G} \oplus \mathcal{H}$ is not just some element in \mathcal{G} , but in fact the KME of some probability distribution on \mathcal{Y} , as long as we condition on an event of the form $V = v = \varphi(x)$ outside the null events N_1 and N_2 . Note, though, that these null sets could be geometrically quite large.

As mentioned above, there is another analogy between CMEs and Gaussian conditioning, namely the assumption under which the formula for the conditional mean is particularly nice, i.e. does not require finite-rank approximations of the (cross-)covariance operators:

Theorem 7.3. Under Assumption 2.1 and with the random variable (U, V) defined by (7.1), Assumption C is equivalent to the compatibility of (C, \mathcal{H}) .

Proof. By Theorems 4.1 and 6.6, both conditions are equivalent to $\text{ran } C_{XY} \subseteq \text{ran } C_X$. \blacksquare

8. Closing Remarks

This article has demonstrated rigorous foundations for the method of conditional mean embedding in reproducing kernel Hilbert spaces. Mild and verifiable sufficient conditions have been provided for the centred and uncentred variants of the CME formula to yield an element $\mu_{Y|X=x}$ that is indeed the kernel mean embedding of the conditional distribution $\mathbb{P}_{Y|X=x}$ on \mathcal{Y} . The CME formula required a correction in the centred case but, modulo this correction, it is more generally applicable than its uncentred counterpart and provides stronger statements: Theorem 4.4 proves convergence in $L^2(\mathbb{P}; \mathcal{G})$ as well as \mathbb{P}_X -almost everywhere convergence, while its analogue Theorem 5.4 yields only convergence in $L^2(\mathbb{P}; \mathcal{G})$. The reason is that

$({}^u\mu^{(n)}(X, \cdot))_{n \in \mathbb{N}}$ defined by (5.5), in contrast to $(\mu^{(n)}(X, \cdot))_{n \in \mathbb{N}}$, may fail to be a martingale (cf. Lemma A.11) and we cannot apply Diestel and Uhl (1977, Theorem V.2.8). Therefore, we advocate for the centred version of the CME formula as the preferred formulation in practice. We have also demonstrated the precise relationship between CMEs and well-established formulae for the conditioning of Gaussian random variables in Hilbert spaces.

Some natural directions for further research suggest themselves:

First, in practice, the KMEs and kernel (cross-)covariance operators will often be estimated using sampled data, and so empirical versions of the CME, along with convergence guarantees, are of great practical importance. Various empirical CMEs have already been considered and applied in the literature (Fukumizu, 2015; Fukumizu et al., 2013; Grünewälder et al., 2012; Park and Muandet, 2020), but their approximation accuracy is not at all trivial to analyse, conditions for validity along the lines of our Assumptions A–^uC are not yet known, and a detailed treatment would be too long to consider in this work, which has deliberately focused on the population CME. Appendix B gives an overview of the technical obstacles that must be overcome in the empirical setting, existing results in the area, and work yet to do.

Second, when using CMEs for inference, a remaining step might be to undo the kernel mean embedding, i.e. to recover the conditional distribution $\mathbb{P}_{Y|X=x}$ on \mathcal{Y} from its embedding $\mu_{Y|X=x} \in \mathcal{G}$, or its density with respect to a reference measure on \mathcal{Y} . This is a particular instance of a non-parametric inverse problem and a principled solution, based upon Tikhonov regularisation, has been proposed in the context of the *kernel conditional density operator* (KCDO) by Schuster et al. (2020). The relationship between this KCDO approach and the sufficient conditions for CME that have been considered in this article remains to be precisely formulated; given the intimate relationship between Tikhonov regularisation and the Moore–Penrose pseudo-inverse, this should be a fruitful avenue of research.

Acknowledgements

The research presented here was supported in part by the German Research Foundation (Deutsche Forschungsgemeinschaft) through project TrU-2 “Demand modelling and control for e-commerce using RKHS transfer operator approaches” of the Excellence Cluster “MATH+ The Berlin Mathematics Research Centre” (EXC-2046/1, project ID: 390685689). The authors also wish to thank S. Klus, H. C. Lie, M. Mollenhauer, and B. Sprungk for helpful and collegial discussions.

A. Technical Results

This section contains several technical results used in the proofs of the theorems given in the article. The following well-known result due to Douglas (1966, Theorem 1) (see also Fillmore and Williams (1971, Theorem 2.1)) is used several times:

Theorem A.1. *Let \mathcal{H} , \mathcal{H}_1 and \mathcal{H}_2 be Hilbert spaces and let $A: \mathcal{H}_1 \rightarrow \mathcal{H}$ and $B: \mathcal{H}_2 \rightarrow \mathcal{H}$ be bounded linear operators with $\text{ran } A \subseteq \text{ran } B$. Then $Q := B^\dagger A: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is a well-defined bounded linear operator, where B^\dagger denotes the Moore–Penrose pseudo-inverse of B . It is the unique operator that satisfies the conditions*

$$A = BQ, \quad \ker Q = \ker A, \quad \text{ran } Q \subseteq \overline{\text{ran } B^*}. \quad (\text{A.1})$$

Remark A.2. In the original work of Douglas (1966) only the existence of a bounded operator Q such that $A = BQ$ was shown. However, the construction of Q in the proof is identical to

that of B^\dagger (multiplied by A). This connection has been observed before by [Arias et al. \(2008, Corollary 2.2 and Remark 2.3\)](#), where it was proven in the case of closed range operators, leaving the proof of the general case to the reader.

The following result partially generalises ([De Vito et al., 2006](#), Proposition 4.1):

Lemma A.3. *Let \mathcal{H} be a separable Hilbert space, let \mathcal{G} be an RKHS over \mathcal{Y} with canonical feature map ψ , and suppose that \mathcal{G} is a subset of $\mathcal{L}^2(\nu)$, where ν is a σ -finite measure on \mathcal{Y} . Then any bounded linear operator $A: \mathcal{H} \rightarrow \mathcal{G}$ is Hilbert–Schmidt as an operator $A: \mathcal{H} \rightarrow L^2(\nu)$.*

Proof. Let $h \in \mathcal{H}$ and $y \in \mathcal{Y}$. Then $(Ah)(y) = \langle \psi(y), Ah \rangle_{\mathcal{G}} = \langle A^* \psi(y), h \rangle_{\mathcal{H}}$. Thus A is a Carleman operator and the claim follows from [Weidmann \(1980, Theorem 6.15\)](#). ■

The following results are used in the proofs of [Sections 3, 4, and 7](#). Note that [Lemma A.4](#) is essentially one direction of Proposition 5 in [Fukumizu et al. \(2009\)](#), but does not require k to be bounded, which makes a separate proof necessary.

Lemma A.4. *Under [Assumption 2.1](#), if k is a characteristic kernel, then $\mathcal{H}_{\mathcal{C}}$ is dense in $L^2_{\mathcal{C}}(\mathbb{P}_X)$.*

Proof. Suppose that $\mathcal{H}_{\mathcal{C}}$ is not dense in $L^2_{\mathcal{C}}(\mathbb{P}_X)$. Then there exists $f \in L^2(\mathbb{P}_X)$ that is not \mathbb{P}_X -a.e. constant such that $[f] \perp_{L^2_{\mathcal{C}}(\mathbb{P}_X)} \mathcal{H}_{\mathcal{C}}$. Choose $\tilde{f} := f - \mathbb{E}[f(X)]$ and set

$$Q_1(E) := \int_E |\tilde{f}| d\mathbb{P}_X, \quad Q_2(E) := \int_E (|\tilde{f}| - \tilde{f}) d\mathbb{P}_X$$

for every Borel-measurable subset $E \subseteq \mathcal{X}$. Since $\|\tilde{f}\|_{L^1(\mathbb{P}_X)} \neq 0$, we may assume without loss of generality that $\|\tilde{f}\|_{L^1(\mathbb{P}_X)} = 1$, making Q_1 and Q_2 two *distinct* probability distributions. Since, for every $h \in \mathcal{H}$,

$$\langle \tilde{f}, h \rangle_{L^2(\mathbb{P}_X)} = \langle f - \mathbb{E}[f(X)], h \rangle_{L^2(\mathbb{P}_X)} \stackrel{[f] \perp \mathcal{H}_{\mathcal{C}}}{=} \langle f - \mathbb{E}[f(X)], \mathbb{E}[h(X)] \rangle_{L^2(\mathbb{P}_X)} = 0,$$

it follows that $\tilde{f} \perp_{L^2(\mathbb{P}_X)} \mathcal{H}$. Let $Z_1 \sim Q_1$ and $Z_2 \sim Q_2$ and $x \in \mathcal{X}$. Since $\varphi(x) \in \mathcal{H}$,

$$(\mathbb{E}[\varphi(Z_1)] - \mathbb{E}[\varphi(Z_2)])(x) = \langle \tilde{f}, \varphi(x) \rangle_{L^2(\mathbb{P}_X)} = 0,$$

which contradicts the assumption that k is characteristic. Note that, by [Assumption 2.1](#), $\mathbb{E}[\varphi(Z_1)]$ and $\mathbb{E}[\varphi(Z_2)]$ are well defined. In fact, by the Cauchy–Schwarz inequality,

$$\mathbb{E}[\|\varphi(Z_1)\|_{\mathcal{H}}] = \int_{\mathcal{X}} \|\varphi(x)\|_{\mathcal{H}} |\tilde{f}(x)| d\mathbb{P}_X(x) \leq \mathbb{E}[\|\varphi(X)\|_{\mathcal{H}}^2]^{1/2} \mathbb{E}[\tilde{f}(X)^2]^{1/2} < \infty$$

and similarly for Z_2 . ■

Lemma A.5. *Under [Assumption 2.1](#), $\ker C_X = \{h \in \mathcal{H} \mid h \text{ is } \mathbb{P}_X\text{-a.e. constant in } \mathcal{X}\}$ and $\ker {}^u C_X = \{h \in \mathcal{H} \mid h = 0 \text{ } \mathbb{P}_X\text{-a.e. in } \mathcal{X}\}$.*

Proof. This is a direct consequence of the facts that $\langle h, C_X h \rangle_{\mathcal{H}} = \mathbb{V}[h(X)]$ and that $\langle h, {}^u C_X h \rangle_{\mathcal{H}} = \|h\|_{L^2(\mathbb{P}_X)}^2$. ■

Lemma A.6. *Under [Assumption 2.1](#), for all $h \in \mathcal{H}$ and $g \in \mathcal{G}$,*

$$\text{Cov}[h(X), f_g(X)] = \langle h, C_{XY} g \rangle_{\mathcal{H}}, \quad {}^u \text{Cov}[h(X), f_g(X)] = \langle h, {}^u C_{XY} g \rangle_{\mathcal{H}}.$$

Proof. Let $h \in \mathcal{H}$ and $g \in \mathcal{G}$ be arbitrary. Then

$$\begin{aligned} \text{Cov}[h(X), f_g(X)] &= \mathbb{E}[h(X)\mathbb{E}[g(Y)|X]] - \mathbb{E}[h(X)]\mathbb{E}[\mathbb{E}[g(Y)|X]] \\ &= \mathbb{E}[h(X)g(Y)] - \mathbb{E}[h(X)]\mathbb{E}[g(Y)] \\ &= \text{Cov}[h(X), g(Y)] \\ &= \langle h, C_{XY}g \rangle_{\mathcal{H}}, \end{aligned}$$

as required. The second statement is proved analogously using uncentred covariance operators and without subtracting the (products of) expected values. \blacksquare

Lemma A.7. Under [Assumption 2.1](#), let $A: \mathcal{G} \rightarrow \mathcal{H}$ be a bounded linear operator. Then, for all $x \in \mathcal{X}$ and $y \in \mathcal{Y}$,

$$(A\psi(y))(x) = (A^*\varphi(x))(y), \quad \mathbb{E}[(A\psi(y))(X)] = (A^*\mu_X)(y).$$

Proof. By the reproducing properties of ψ , φ , and μ_X ,

$$\begin{aligned} (A\psi(y))(x) &= \langle A\psi(y), \varphi(x) \rangle_{\mathcal{H}} = \langle \psi(y), A^*\varphi(x) \rangle_{\mathcal{G}} = (A^*\varphi(x))(y), \\ \mathbb{E}[(A\psi(y))(X)] &= \langle A\psi(y), \mu_X \rangle_{\mathcal{H}} = \langle \psi(y), A^*\mu_X \rangle_{\mathcal{G}} = (A^*\mu_X)(y), \end{aligned}$$

as claimed. \blacksquare

Lemma A.8. Let V be a Hilbert space, let $U_1 \subseteq U_2 \subseteq \dots$ be an increasing sequence of closed subspaces $U_n \subseteq V$, $n \in \mathbb{N}$, and let $U := \bigcup_{n \in \mathbb{N}} U_n$. Further, let $P_{U_n}: V \rightarrow U_n$ denote the orthogonal projection onto U_n . Then, for all $v \in \overline{U}$,

$$P_{U_n}v \xrightarrow{n \rightarrow \infty} v.$$

Proof. Let $v \in \overline{U}$ and $\varepsilon > 0$. Then there exists $u \in U$ such that $\|u - v\| < \varepsilon$. Since the sequence $(U_n)_{n \in \mathbb{N}}$ is increasing and U is its union, there exists an $n_0 \in \mathbb{N}$ such that $u \in U_{n_0}$ and thereby $P_{U_n}u = u$ for all $n \geq n_0$. We therefore obtain, for $n \geq n_0$,

$$\|P_{U_n}v - v\| \leq \|P_{U_n}v - P_{U_n}u\| + \|P_{U_n}u - u\| + \|u - v\| \leq \|P_{U_n}\|\|v - u\| + \|u - v\| < 2\varepsilon,$$

by the triangle inequality and non-expansivity of orthogonal projection. \blacksquare

Lemma A.9. Under [Assumption 2.1](#), with $\mathcal{H}^{(n)}$, $C^{(n)}$, $h_g^{(n)}$ as in [Theorem 4.4](#) and ${}^u C^{(n)}$, ${}^u h_g^{(n)}$ as in [Theorem 5.4](#),

- (a) $\text{Cov}[h(X), f_g(X)] = \lim_{n \rightarrow \infty} \text{Cov}[h(X), h_g^{(n)}(X)]$ for all $h \in \mathcal{H}$;
- (b) $[h_g^{(n)}]$ is the $L^2_{\mathcal{C}}$ -orthogonal projection of $[f_g]$ onto $\mathcal{H}_{\mathcal{C}}^{(n)}$ for all $g \in \mathcal{G}$;
- (c) ${}^u \text{Cov}[h(X), f_g(X)] = \lim_{n \rightarrow \infty} {}^u \text{Cov}[h(X), {}^u h_g^{(n)}(X)]$ for all $h \in \mathcal{H}$;
- (d) ${}^u h_g^{(n)}$ is the L^2 -orthogonal projection of f_g onto $\mathcal{H}^{(n)}$ for all $g \in \mathcal{G}$.

Proof. We only give the proofs of (a) and (b); (c) and (d) can be proven similarly. It is clear that that $C^{(n)} \rightarrow C$ (in the strong and thereby in the weak sense) as $n \rightarrow \infty$ and that C_X and $C_X^{(n)}$ agree on $\mathcal{H}^{(n)} \ni h_g^{(n)}$. Using [Lemma A.6](#) we obtain, for all $h \in \mathcal{H}$,

$$\begin{aligned} \text{Cov}[h(X), f_g(X)] &= \langle h, C_{XY}g \rangle_{\mathcal{H}} \\ &= \lim_{n \rightarrow \infty} \langle h, C_{XY}^{(n)}g \rangle_{\mathcal{H}} \\ &= \lim_{n \rightarrow \infty} \langle h, C_X^{(n)}h_g^{(n)} \rangle_{\mathcal{H}} \\ &= \lim_{n \rightarrow \infty} \langle h, C_X h_g^{(n)} \rangle_{\mathcal{H}} \\ &= \lim_{n \rightarrow \infty} \text{Cov}[h(X), h_g^{(n)}(X)], \end{aligned}$$

which yields (a). Also, for arbitrary $h^{(n)} \in \mathcal{H}^{(n)}$, Lemma A.6 yields

$$\begin{aligned}
\langle [h^{(n)}], [f_g] \rangle_{L^2_{\mathcal{C}}} &= \text{Cov}[h^{(n)}(X), f_g(X)] \\
&= \langle h^{(n)}, C_{XY}g \rangle_{\mathcal{H}} \\
&= \langle C_{YX}h^{(n)}, g \rangle_{\mathcal{G}} \\
&= \langle C_{YX}^{(n)}h^{(n)}, g \rangle_{\mathcal{G}} \\
&= \langle h^{(n)}, C_{XY}^{(n)}g \rangle_{\mathcal{H}} \\
&= \langle h^{(n)}, C_X^{(n)}h_g^{(n)} \rangle_{\mathcal{H}} \\
&= \langle h^{(n)}, C_X h_g^{(n)} \rangle_{\mathcal{H}} \\
&= \text{Cov}[h^{(n)}(X), h_g^{(n)}(X)] \\
&= \langle [h^{(n)}], [h_g^{(n)}] \rangle_{L^2_{\mathcal{C}}},
\end{aligned}$$

which yields (b). ■

Lemma A.10. Let Assumption 2.1 hold and $\mathcal{H}^{(1)} \subseteq \mathcal{H}^{(2)} \subseteq \dots$ be an increasing sequence of closed subspaces $\mathcal{H}^{(n)}$ of $L^2(\mathbb{P}_X)$, $n \in \mathbb{N}$. Further, let $\mathbf{m}, \mathbf{m}^{(n)} \in L^2(\mathbb{P}_X; \mathcal{G}) \simeq L^2(\mathbb{P}_X) \otimes \mathcal{G}$ and denote $\bar{f} := f - \mathbb{E}[f(X)]$ for $f \in L^2(\mathbb{P}_X)$ and $\bar{\mathbf{f}} := \mathbf{f} - \mathbb{E}[\mathbf{f}(X)]$ for $\mathbf{f} \in L^2(\mathbb{P}_X; \mathcal{G})$.

- (a) If $([\mathbf{m}^{(n)}](\cdot))(y)$ is the orthogonal projection in $L^2_{\mathcal{C}}(\mathbb{P}_X)$ of $([\mathbf{m}](\cdot))(y)$ onto $\mathcal{H}^{(n)}_{\mathcal{C}}$ for each $y \in \mathcal{Y}$, then $[\mathbf{m}^{(n)}]$ is the orthogonal projection in $L^2_{\mathcal{C}}(\mathbb{P}_X; \mathcal{G})$ of $[\mathbf{m}]$ onto $(\mathcal{H}^{(n)} \otimes \mathcal{G})_{\mathcal{C}}$.
- (b) If, in addition to the assumption in (a), $([\mathbf{m}^{(n)}](\cdot))(y) \rightarrow ([\mathbf{m}](\cdot))(y)$ in $L^2_{\mathcal{C}}(\mathbb{P}_X)$ as $n \rightarrow \infty$ for each $y \in \mathcal{Y}$, then $[\mathbf{m}^{(n)}] \rightarrow [\mathbf{m}]$ in $L^2_{\mathcal{C}}(\mathbb{P}_X; \mathcal{G})$, or, in other words, $\bar{\mathbf{m}}^{(n)}(X) \rightarrow \bar{\mathbf{m}}(X)$ in $L^2(\mathbb{P}; \mathcal{G})$.
- (c) If $(\mathbf{m}^{(n)}(\cdot))(y)$ is the orthogonal projection in $L^2(\mathbb{P}_X; \mathbb{R})$ of $(\mathbf{m}(\cdot))(y)$ onto $\mathcal{H}^{(n)}$ for each $y \in \mathcal{Y}$, then $\mathbf{m}^{(n)}$ is the orthogonal projection in $L^2(\mathbb{P}_X; \mathcal{G})$ of \mathbf{m} onto $\mathcal{H}^{(n)} \otimes \mathcal{G}$.
- (d) If, in addition to the assumption in (c), $(\mathbf{m}^{(n)}(\cdot))(y) \rightarrow (\mathbf{m}(\cdot))(y)$ in $L^2(\mathbb{P}_X; \mathbb{R})$ as $n \rightarrow \infty$ for each $y \in \mathcal{Y}$, then $\mathbf{m}^{(n)} \rightarrow \mathbf{m}$ in $L^2(\mathbb{P}_X; \mathcal{G})$, or, in other words, $\mathbf{m}^{(n)}(X) \rightarrow \mathbf{m}(X)$ in $L^2(\mathbb{P}; \mathcal{G})$.

Proof. We only give the proofs of (a) and (b); (c) and (d) can be proven similarly with fewer technicalities. Let $h \in \mathcal{H}^{(n)}$ and $y \in \mathcal{Y}$. Then

$$\begin{aligned}
\langle [\mathbf{m}^{(n)}] - [\mathbf{m}], [h \otimes \psi(y)] \rangle_{L^2_{\mathcal{C}}(\mathbb{P}_X; \mathcal{G})} &= \mathbb{E}[\langle \bar{\mathbf{m}}^{(n)}(X) - \bar{\mathbf{m}}(X), \psi(y) \rangle_{\mathcal{G}} \bar{h}(X)] \\
&= \mathbb{E}\left[\left(\bar{\mathbf{m}}^{(n)}(X)(y) - \bar{\mathbf{m}}(X)(y)\right) \bar{h}(X)\right] \\
&= \langle ([\mathbf{m}^{(n)}](\cdot))(y) - ([\mathbf{m}](\cdot))(y), [h] \rangle_{L^2_{\mathcal{C}}(\mathbb{P}_X; \mathbb{R})} \\
&= 0,
\end{aligned}$$

which proves (a). Hence, by Lemma A.8, $[\mathbf{m}^{(n)}]$ converges in $L^2_{\mathcal{C}}(\mathbb{P}_X; \mathcal{G})$ to some limit $[\mathbf{m}']$. This implies pointwise convergence for each $y \in \mathcal{Y}$ in the following sense:

$$\begin{aligned}
\|([\mathbf{m}^{(n)}](\cdot))(y) - ([\mathbf{m}'](\cdot))(y)\|_{L^2_{\mathcal{C}}(\mathbb{P}_X; \mathbb{R})}^2 &= \mathbb{E}[\langle \psi(y), \bar{\mathbf{m}}^{(n)}(X) - \bar{\mathbf{m}}'(X) \rangle_{\mathcal{G}}^2] \\
&\leq \|\psi(y)\|_{\mathcal{G}}^2 \mathbb{E}[\|\bar{\mathbf{m}}^{(n)}(X) - \bar{\mathbf{m}}'(X)\|_{\mathcal{G}}^2] \\
&= \|\psi(y)\|_{\mathcal{G}}^2 \|\mathbf{m}^{(n)} - \mathbf{m}'\|_{L^2(\mathbb{P}_X; \mathcal{G})}^2 \\
&\xrightarrow{n \rightarrow \infty} 0.
\end{aligned}$$

Therefore, by assumption, $[\mathbf{m}']$ agrees with $[\mathbf{m}]$ \mathbb{P}_X -a.e., proving (b). ■

Lemma A.11. *Under the assumptions and notation of [Theorem 4.4](#), $(\mu^{(n)}(X, \cdot))_{n \in \mathbb{N}}$ is a martingale in $L^2(\Omega, \Sigma, \mathbb{P}; \mathcal{G})$ with respect to the filtration $(\sigma(V^{(n)}))_{n \in \mathbb{N}}$ of Σ , where $V^{(n)} := P_{\mathcal{H}^{(n)}}(\varphi(X))$ and $P_{\mathcal{H}^{(n)}}: \mathcal{H} \rightarrow \mathcal{H}^{(n)}$ denotes the orthogonal projection in \mathcal{H} onto $\mathcal{H}^{(n)}$.*

Proof. Consider the Karhunen–Loève expansion of $\varphi(X)$,

$$\varphi(X) = \mu_X + \sum_{i \in \mathbb{N}} Z_i h_i,$$

where $Z_i: (\Omega, \Sigma, \mathbb{P}) \rightarrow \mathbb{R}$ are uncorrelated real-valued random variables with $\mathbb{E}[Z_i] = 0$ and $\mathbb{V}[Z_i] = \sigma_i$ for all $i \in \mathbb{N}$, $\sigma_i \geq 0$ denoting the eigenvalue of C_X corresponding to the eigenvector h_i . We observe that $\sigma(V^{(n)}) = \sigma(Z^{(n)})$, where $Z^{(n)} := (Z_1, \dots, Z_n)$. Now let $A^{(n)} = (C_X^{(n)\dagger} C_{XY}^{(n)})^*$, $n \in \mathbb{N}$, and observe that $A^{(n)}v = A^{(n)}P_{\mathcal{H}^{(n)}}v$ and that $A^{(n+1)}$ and $A^{(n)}$ agree on $\mathcal{H}^{(n)}$. Hence, for $n \in \mathbb{N}$,

$$\begin{aligned} \mathbb{E}[\mu^{(n+1)}(X, \cdot) | V^{(n)}] &= \mu_Y + A^{(n+1)} \mathbb{E}[V - \mu_X | Z^{(n)}] \\ &= \mu_Y + A^{(n+1)} \mathbb{E}\left[\sum_{i \in \mathbb{N}} Z_i h_i \mid Z^{(n)}\right] \\ &= \mu_Y + A^{(n)} \sum_{i=1}^n Z_i h_i \\ &= \mu_Y + A^{(n)} (\varphi(X) - \mu_X) \\ &= \mu^{(n)}(X, \cdot), \end{aligned}$$

proving the martingale property. ■

Lemma A.12. *Let Assumptions [2.1](#) and [B*](#) hold. Then $\mathbb{E}[C_{Y|X}] = \int_{\mathcal{X}} C_{Y|X=x} d\mathbb{P}_X(x)$ is well defined as a strong (Bochner) integral, i.e. $\int_{\mathcal{X}} \|C_{Y|X=x}\| d\mathbb{P}_X(x) < \infty$.*

Proof. The Cauchy–Schwarz inequality and [\(2.2\)](#) imply that, for $x \in \mathcal{X}_Y$,

$$\begin{aligned} \|C_{Y|X=x}\| &= \sup_{\|g\|_{\mathcal{G}} \leq 1, \|\tilde{g}\|_{\mathcal{G}} \leq 1} \langle g, C_{Y|X=x} \tilde{g} \rangle_{\mathcal{G}} \\ &= \sup_{\|g\|_{\mathcal{G}} \leq 1, \|\tilde{g}\|_{\mathcal{G}} \leq 1} \langle g, \tilde{g} \rangle_{\mathcal{L}_{\mathcal{G}}^2(\mathbb{P}_{Y|X=x})} \\ &\leq \sup_{\|g\|_{\mathcal{G}} \leq 1, \|\tilde{g}\|_{\mathcal{G}} \leq 1} \|g\|_{\mathcal{L}^2(\mathbb{P}_{Y|X=x})} \|\tilde{g}\|_{\mathcal{L}^2(\mathbb{P}_{Y|X=x})} \\ &\leq \mathbb{E}[\|\psi(Y)\|_{\mathcal{G}}^2 | X = x], \end{aligned}$$

which, by the law of total expectation and [\(2.1\)](#), yields that

$$\mathbb{E}[\|C_{Y|X}\|] \leq \mathbb{E}[\mathbb{E}[\|\psi(Y)\|_{\mathcal{G}}^2 | X]] = \mathbb{E}[\|\psi(Y)\|_{\mathcal{G}}^2] < \infty,$$

as claimed. ■

B. Empirical Estimates for CMEs

In practice, the kernel mean embeddings and kernel (cross-)covariance operators will often be estimated empirically from observed data, and so empirical versions of the CME, along with convergence guarantees, are of great importance. As mentioned in [Remark 1.2](#), this topic is

beyond the scope of this paper. However, we wish to point out why this is a complex problem and briefly address the main difficulties.

In the simplest setting, given $J \in \mathbb{N}$ independent samples $(X_1, Y_1), \dots, (X_J, Y_J) \sim \mathbb{P}_{XY}$, we have the empirical estimators

$$\mu_X \approx \hat{\mu}_X := \frac{1}{J} \sum_{j=1}^J \varphi(X_j), \quad C_{XY} \approx \hat{C}_{XY} := \frac{1}{J} \sum_{j=1}^J (\varphi(X_j) - \hat{\mu}_X) \otimes (\psi(Y_j) - \hat{\mu}_Y),$$

and so on. (To simplify the notation, we suppress the obvious J -dependence of these estimators.) Laws of large numbers for these empirical estimators have already been established — see e.g. [Smola et al. \(2007, Theorem 2\)](#) and [Mollenhauer \(2018, Lemma 5.8\)](#) — but the impact of this approximation error upon conditioning is, to the best of our knowledge, not yet fully quantified. One natural approach to approximate the CME $\mu_{Y|X=x}$ is the regularisation of \hat{C}_X or \hat{C}_X^\dagger ,

$$\mu_{Y|X=x} \approx \left((\hat{C}_X + \varepsilon \text{Id}_{\mathcal{H}})^\dagger \hat{C}_{XY} \right)^* \varphi(x) = \hat{C}_{YX} (\hat{C}_X + \varepsilon \text{Id}_{\mathcal{H}})^{-1} \varphi(x),$$

where $\varepsilon > 0$ is a regularisation parameter which may depend on J . Note that such a regularisation can be viewed as an approximation both to the new CME formula derived in [Theorem 5.3](#), $\mu_{Y|X=x} = (\hat{C}_X^\dagger \hat{C}_{XY})^* \varphi(x)$, as well as to the original (uncentred) one, $\mu_{Y|X=x} = C_{YX} C_X^{-1} \varphi(x)$. Therefore, this approach is rather well studied and convergence rates for this strategy have been established under certain conditions ([Fukumizu, 2015](#); [Grünwälder et al., 2012](#); [Park and Muandet, 2020](#)).

However, the new formulae (4.3), (4.6), and (5.3) relying on the Moore–Penrose pseudo-inverse suggest another type of approximation, where we will focus on the centred case from now on. The naïve estimate would be

$$\mu_{Y|X=x} \approx \hat{\mu}_Y + (\hat{C}_X^\dagger \hat{C}_{XY})^* (\varphi(x) - \hat{\mu}_X). \quad (\text{B.1})$$

Note that $\text{ran } \hat{C}_{XY} \subseteq \text{ran } \hat{C}_X$ and so (B.1) is well defined. However, the convergence of \hat{C}_X to C_X (e.g. in the Hilbert–Schmidt norm, as $J \rightarrow \infty$) translates badly to the convergence of \hat{C}_X^\dagger to the pseudo-inverse C_X^\dagger . One problem is that small eigenvalues of C_X might be approximated by eigenvalues of \hat{C}_X that are orders of magnitude smaller, causing \hat{C}_X^\dagger to “blow up”. So, in addition to the convergence of \hat{C}_X in the classical norms (such as the Hilbert–Schmidt norm or operator norm), we need to control the the smallest eigenvalue of \hat{C}_X .

A natural workaround, inspired by the finite-rank approximation in [Theorem 4.4](#), is to truncate¹² the (cross-)covariance operators to a subspace $\mathcal{H}^{(n)} = \text{span}\{h_1, \dots, h_n\}$ of \mathcal{H} with $\dim \mathcal{H}^{(n)} = n = n(J) \ll J$. One might thus hope to approximate the dominant n eigenvalues of C_X well while artificially setting the others to zero and preventing the blow-up of \hat{C}_X^\dagger . There are several results from random matrix theory that control the behaviour of the n^{th} eigenvalue of (truncated) empirical covariance matrices for growing J and $n = n(J)$ ([Bai and Silverstein, 2010](#); [Bai, 1999](#); [Bai and Yin, 1993](#); [Heiny and Mikosch, 2018](#)). Most of these results are formulated for the case where the true mean is known to be zero and the true covariance matrix is the identity matrix and are typically of the following form, where $\lambda_{\max}(M)$ and $\lambda_{\min}(M)$ denote the largest and smallest eigenvalues of a matrix M , respectively:

¹²Naturally, truncation can be viewed as another form of regularisation. For further regularised estimates of large covariance and precision matrices by tapering, banding, sparsifying or similar see e.g. [Bickel and Levina \(2008a,b\)](#); [Cai et al. \(2010\)](#); [Yuan \(2010\)](#) and references therein.

Theorem B.1 (Bai and Yin (1993, Theorem 2)). *Let $(\xi_{ij})_{i,j \in \mathbb{N}}$ be a double array of independent and identically distributed random variables with zero mean and unit variance. For $J \in \mathbb{N}$, let $n = n(J)$ be such that $n(J) \rightarrow \infty$ and $n(J)/J \rightarrow \gamma \in (0, 1)$ for $J \rightarrow \infty$ and let*

$$A_J = (\xi_{ij})_{i=1, \dots, n(J), j=1, \dots, J}, \quad S_J = \frac{1}{J} A_J A_J^\top. \quad (\text{B.2})$$

Then, if $\mathbb{E}[\xi_{11}^4] < \infty$,

$$\lambda_{\max}(S_J) \xrightarrow[J \rightarrow \infty]{a.e.} (1 + \sqrt{\gamma})^2, \quad \lambda_{\min}(S_J) \xrightarrow[J \rightarrow \infty]{a.e.} (1 - \sqrt{\gamma})^2.$$

In our case the covariance operator C_X is not the identity; however, our case follows partially from Theorem B.1 using the eigendecomposition of C_X ,

$$C_X = \sum_{i \in \mathbb{N}} \sigma_i h_i \otimes h_i, \quad \sigma_1 \geq \sigma_2 \geq \dots \geq 0,$$

and the Karhunen–Loève expansion of $V = \varphi(X) - \mu_X$,

$$V = \sum_{i \in \mathbb{N}} \sqrt{\sigma_i} \xi_i h_i,$$

where $\xi_n: \Omega \rightarrow \mathbb{R}$ are uncorrelated random variables with $\mathbb{E}[\xi_n] = 0$ and $\mathbb{V}[\xi_i] = 1$ for each $i \in \mathbb{N}$ and $(h_i)_{i \in \mathbb{N}}$ is an orthonormal eigenbasis of \mathcal{H} . To this end, let $V_j := \varphi(X_j) - \mu_X$ be i.i.d. copies of V and let $V_j^{(n)}$ be their respective orthogonal projections onto $\mathcal{H}^{(n)} := \text{span}\{h_1, \dots, h_n\}$, i.e.

$$V_j = \sum_{i \in \mathbb{N}} \sqrt{\sigma_i} \xi_{ij} h_i \text{ with } \xi_{ij} \stackrel{\text{i.i.d.}}{\sim} \xi_i \text{ for all } i, j, \quad V_j^{(n)} = \sum_{i=1}^n \sqrt{\sigma_i} \xi_{ij} h_i.$$

To simplify notation, let us work with $(n \times n)$ -matrices instead of operators, expressed in the basis (h_1, \dots, h_n) of $\mathcal{H}^{(n)}$. Then

$$\widehat{C}_X^{(n)} = \frac{1}{J} \sum_{j=1}^J V_j \otimes V_j = (C_X^{(n)})^{1/2} S_J (C_X^{(n)})^{1/2}, \quad (\text{B.3})$$

where S_J is defined by (B.2). So, we are *nearly* in the setup of Theorem B.1 and ready to conclude

$$\lim_{J \rightarrow \infty} \lambda_{\min}(\widehat{C}_X^{(n)}) \geq \lim_{J \rightarrow \infty} \sigma_n^{1/2} \lambda_{\min}(S_J) \sigma_n^{1/2} \geq \sigma_n (1 - \sqrt{\gamma})^2,$$

with $n = n(J)$ and γ as in Theorem B.1. (Here we make use of the fact that $\|Ax\| \geq \lambda_{\min}(A)\|x\|$ for all x when A is positive semi-definite.) However, some obstacles remain:

- Since the random variables V_j are independent copies of V , the distributions of ξ_{ij} and $\xi_{ij'}$ agree for all j, j' , but we cannot expect the distributions of ξ_{ij} and $\xi_{i'j}$ to agree for all i, i' . Hence, ξ_{ij} do not fulfil the requirement of being identically distributed. However, there are generalisations of Theorem B.1 to this case under technical assumptions; see e.g. Bai (1999, Theorem 2.8).
- It is unclear when the condition $\mathbb{E}[\xi_{11}^4] < \infty$ is fulfilled. There are, however, some results in the case of infinite fourth moments; see Heiny and Mikosch (2018) and references therein.
- The random variables ξ_i are uncorrelated but, in general, not independent. We are not aware of a result similar to Theorem B.1 for the uncorrelated case.¹³

¹³Chen et al. (2013) discuss the estimation of covariance and precision matrices for time series data where the random variables V_j are not assumed to be independent. However, we need to drop independence in the rows of A_J , not in its columns.

Even if we manage to formulate a version of [Theorem B.1](#) which suits our needs, note that the considerations so far, if executed rigorously, only guarantee that C_X^\dagger does not blow up. This is still some way short of establishing the convergence of the corresponding CME estimator

$$\mu_{Y|X=x} \approx \hat{\mu}_Y + (\hat{C}_X^{(n)\dagger} \hat{C}_{XY}^{(n)})^* (\varphi(x) - \hat{\mu}_X), \quad n = n(J). \quad (\text{B.4})$$

We conclude here by summarising some of the necessary steps:

- In the above considerations we assumed μ_X to be known. In practice, however, we may only employ its empirical estimate $\hat{\mu}_X$. Since $\mu_X \otimes \mu_X$ is a rank-one estimator, this issue might be partially resolved by [Gohberg and Kreĭn \(1969, Corollary 2.1\)](#), which implies that the eigenvalues of ${}^u C_X$ and $C_X = {}^u C_X - \mu_X \otimes \mu_X$ have a similar decay rate.
- In order to project onto $\mathcal{H}^{(n)} = \text{span}\{h_1, \dots, h_n\}$, we require the eigenvectors h_i of C_X . While the n dominant eigenvectors of \hat{C}_X can be used as estimates of h_i , it is unclear how the approximation error affects the theoretical results presented above.
- Controlling the eigenvalues alone is insufficient. For a reasonable approximation of $C_X^{(n)}$ in [\(B.3\)](#), we would need a result which tells us that S_J becomes close to the identity matrix for large J , which requires us to understand the behaviour of its eigenvectors as well. The investigation of the eigenvectors of S_J turns out to be extremely challenging; see [Bai and Silverstein \(2010, Chapter 10\)](#) for a survey of existing results.

References

- M. L. Arias, G. Corach, and M. C. Gonzalez. Generalized inverses and Douglas equations. *Proc. Amer. Math. Soc.*, 136(9):3177–3183, 2008. URL <https://doi.org/10.1090/S0002-9939-08-09298-8>.
- Z. Bai and J. W. Silverstein. *Spectral Analysis of Large Dimensional Random Matrices*. Springer Series in Statistics. Springer, New York, second edition, 2010. URL <https://doi.org/10.1007/978-1-4419-0661-8>.
- Z. D. Bai. Methodologies in spectral analysis of large-dimensional random matrices, a review. *Statist. Sinica*, 9(3):611–677, 1999. doi: 10.1142/9789812793096_0015. With comments by G. J. Rodgers and Jack W. Silverstein; and a rejoinder by the author.
- Z. D. Bai and Y. Q. Yin. Limit of the smallest eigenvalue of a large-dimensional sample covariance matrix. *Ann. Probab.*, 21(3):1275–1294, 1993. URL <https://doi.org/10.1214/aop/1176989118>.
- C. R. Baker. Joint measures and cross-covariance operators. *Trans. Amer. Math. Soc.*, 186:273–289, 1973. URL <https://doi.org/10.2307/1996566>.
- A. Berliet and C. Thomas-Agnan. *Reproducing Kernel Hilbert Spaces in Probability and Statistics*. Springer, Boston, 2004. URL <https://doi.org/10.1007/978-1-4419-9096-9>.
- P. J. Bickel and E. Levina. Covariance regularization by thresholding. *Ann. Statist.*, 36(6):2577–2604, 2008a. URL <https://doi.org/10.1214/08-AOS600>.
- P. J. Bickel and E. Levina. Regularized estimation of large covariance matrices. *Ann. Statist.*, 36(1):199–227, 2008b. URL <https://doi.org/10.1214/009053607000000758>.
- T. T. Cai, C.-H. Zhang, and H. H. Zhou. Optimal rates of convergence for covariance matrix estimation. *Ann. Statist.*, 38(4):2118–2144, 2010. URL <https://doi.org/10.1214/09-AOS752>.
- X. Chen, M. Xu, and W. B. Wu. Covariance and precision matrix estimation for high-dimensional time series. *Ann. Statist.*, 41(6):2994–3021, 2013. URL <https://doi.org/10.1214/13-AOS1182>.
- G. Corach, A. Maestripieri, and D. Stojanoff. Oblique projections and Schur complements. *Acta Sci. Math. (Szeged)*, 67(1-2):337–356, 2001.

- E. De Vito, L. Rosasco, and A. Caponnetto. Discretization error analysis for Tikhonov regularization. *Analysis and Applications*, 04(01):81–99, 2006. URL <https://doi.org/10.1142/S0219530506000711>.
- J. Diestel and J. J. Uhl. *Vector Measures*, volume 15 of *Mathematical Surveys*. American Mathematical Society, Providence, RI, 1977. URL <https://doi.org/10.1090/surv/015>.
- R. G. Douglas. On majorization, factorization, and range inclusion of operators on Hilbert space. *Proc. Amer. Math. Soc.*, 17:413–415, 1966. URL <https://doi.org/10.2307/2035178>.
- H. W. Engl, M. Hanke, and A. Neubauer. *Regularization of Inverse Problems*, volume 375 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1996.
- P. A. Fillmore and J. P. Williams. On operator ranges. *Adv. Math.*, 7(3):254–281, 1971. URL [https://doi.org/10.1016/S0001-8708\(71\)80006-3](https://doi.org/10.1016/S0001-8708(71)80006-3).
- K. Fukumizu. Nonparametric Bayesian inference with kernel mean embedding. In *Modern Methodology and Applications in Spatial-Temporal Modeling*, pages 1–24. Springer Japan, Tokyo, 2015. URL https://doi.org/10.1007/978-4-431-55339-7_1.
- K. Fukumizu, F. R. Bach, and M. I. Jordan. Dimensionality reduction for supervised learning with reproducing kernel Hilbert spaces. *J. Mach. Learn. Res.*, 5(Jan):73–99, 2004a. URL <http://www.jmlr.org/papers/volume5/fukumizu04a/fukumizu04a.pdf>.
- K. Fukumizu, F. R. Bach, and M. I. Jordan. Erratum: Dimensionality reduction for supervised learning with reproducing kernel Hilbert spaces. *J. Mach. Learn. Res.*, 2004b. URL <http://www.jmlr.org/papers/volume5/fukumizu04a/fukumizu04a-erratum.pdf>.
- K. Fukumizu, A. Gretton, X. Sun, and B. Schölkopf. Kernel measures of conditional dependence. In *Advances in Neural Information Processing Systems 20*, pages 489–496. Curran Associates, Inc., 2008. URL <https://papers.nips.cc/paper/3340-kernel-measures-of-conditional-dependence.pdf>.
- K. Fukumizu, F. R. Bach, and M. I. Jordan. Kernel dimension reduction in regression. *Ann. Statist.*, 37(4):1871–1905, 2009. URL <https://doi.org/10.1214/08-AOS637>.
- K. Fukumizu, L. Song, and A. Gretton. Kernel Bayes’ rule: Bayesian inference with positive definite kernels. *J. Mach. Learn. Res.*, 14(1):3753–3783, 2013. URL <http://jmlr.org/papers/volume14/fukumizu13a/fukumizu13a.pdf>.
- I. C. Gohberg and M. G. Kreĭn. *Introduction to the Theory of Linear Nonselfadjoint Operators*, volume 18 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, R.I., 1969. Transl. A. Feinstein.
- S. Grünewälder, G. Lever, L. Baldassarre, S. Patterson, A. Gretton, and M. Pontil. Conditional mean embeddings as regressors. In *Proceedings of the 29th International Conference on Machine Learning*, pages 1823–1830, 2012. URL <https://icml.cc/2012/papers/898.pdf>.
- J. Heiny and T. Mikosch. Almost sure convergence of the largest and smallest eigenvalues of high-dimensional sample correlation matrices. *Stochastic Process. Appl.*, 128(8):2779–2815, 2018. URL <https://doi.org/10.1016/j.spa.2017.10.002>.
- O. Kallenberg. *Foundations of Modern Probability*. Springer, New York, 2006. URL <https://doi.org/10.1007/978-1-4757-4015-8>.
- M. Mollenhauer. Singular value decomposition of operators on reproducing kernel Hilbert spaces. Master’s thesis, Freie Universität Berlin, 2018.
- H. Owhadi and C. Scovel. Separability of reproducing kernel spaces. *Proc. Amer. Math. Soc.*, 145(5):2131–2138, 2017. URL <https://doi.org/10.1090/proc/13354>.
- H. Owhadi and C. Scovel. Conditioning Gaussian measure on Hilbert space. *J. Math. Stat. Anal.*, 2018. URL <https://arxiv.org/abs/1506.04208>.
- J. Park and K. Muandet. A measure-theoretic approach to kernel conditional mean embeddings, 2020. URL <https://arxiv.org/abs/2002.03689>.

- S. Saitoh and Y. Sawano. *Theory of Reproducing Kernels and Applications*, volume 44 of *Developments in Mathematics*. Springer, Singapore, 2016. URL <https://doi.org/10.1007/978-981-10-0530-5>.
- V. V. Sazonov. On characteristic functionals. *Teor. Veroyatnost. i Primenen.*, 3:201–205, 1958.
- I. Schuster, M. Mollenhauer, S. Klus, and K. Muandet. Kernel conditional density operators. In *Proceedings of the 23rd International Conference on Artificial Intelligence and Statistics, AISTATS 2020, 3–5 June 2020, Palermo, Sicily, Italy*, Proceedings of Machine Learning Research, 2020. URL <https://arxiv.org/abs/1905.11255>.
- A. J. Smola, A. Gretton, L. Song, and B. Schölkopf. A Hilbert space embedding for distributions. In *Proceedings of the 18th International Conference on Algorithmic Learning Theory*, pages 13–31, Berlin, Heidelberg, 2007. Springer. URL https://doi.org/10.1007/978-3-540-75225-7_5.
- L. Song, J. Huang, A. Smola, and K. Fukumizu. Hilbert space embeddings of conditional distributions with applications to dynamical systems. In *Proceedings of the 26th Annual International Conference on Machine Learning*, pages 961–968, 2009. URL <https://doi.org/10.1145/1553374.1553497>.
- L. Song, B. Boots, S. M. Siddiqi, G. Gordon, and A. Smola. Hilbert space embeddings of hidden Markov models. In *Proceedings of the 27th International Conference on Machine Learning, ICML2010*, pages 991–998, 2010a. URL <https://dl.acm.org/citation.cfm?id=3104322.3104448>.
- L. Song, A. Gretton, and C. Guestrin. Nonparametric tree graphical models. In Y. W. Teh and M. Titterton, editors, *Proceedings of the Thirteenth International Conference on Artificial Intelligence and Statistics*, volume 9 of *Proceedings of Machine Learning Research*, pages 765–772, 2010b. URL <http://proceedings.mlr.press/v9/song10a/song10a.pdf>.
- B. K. Sriperumbudur, A. Gretton, K. Fukumizu, B. Schölkopf, and G. R. G. Lanckriet. Hilbert space embeddings and metrics on probability measures. *J. Mach. Learn. Res.*, 11:1517–1561, 2010. URL <http://www.jmlr.org/papers/volume11/sriperumbudur10a/sriperumbudur10a.pdf>.
- B. K. Sriperumbudur, K. Fukumizu, and G. R. G. Lanckriet. Universality, characteristic kernels and RKHS embedding of measures. *J. Mach. Learn. Res.*, 12:2389–2410, 2011. URL <http://www.jmlr.org/papers/volume12/sriperumbudur11a/sriperumbudur11a.pdf>.
- I. Steinwart and A. Christmann. *Support Vector Machines*. Information Science and Statistics. Springer, New York, 2008. URL <https://doi.org/10.1007/978-0-387-77242-4>.
- I. Steinwart, D. Hush, and C. Scovel. An explicit description of the reproducing kernel Hilbert spaces of Gaussian RBF kernels. *IEEE Trans. Inf. Thy.*, 52(10):4635–4643, 2006. URL <https://doi.org/10.1109/tit.2006.881713>.
- J. Weidmann. *Linear Operators in Hilbert Spaces*, volume 68 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1980. URL <https://doi.org/10.1007/978-1-4612-6027-1>. Transl. J. Szücs.
- M. Yuan. High dimensional inverse covariance matrix estimation via linear programming. *J. Mach. Learn. Res.*, 11:2261–2286, 2010. URL <http://www.jmlr.org/papers/volume11/yuan10b/yuan10b.pdf>.