

# Individual-based models under various time-scales

Aurelien Velleret \*

December 3, 2019

**Abstract :** Through the recent results that we present, we wish to highlight the various time-scales on which the demographic dynamics and the natural selection can be retrieved from individual-based models. Although these results are by no means exhaustive, both on the mathematical and the biological level, they are very complementary. Indeed, they provide a viewpoint for the most classical time-scales of interest. They namely encompass the time-scale of the life of a single individual, the longer one on which a population can be characterized, the ecological one and at least four nested ones on which selection occurs. The limiting behavior is generally deterministic. Yet, since the selection acts on the alea in the history of individuals, the probabilities are shown to be a key element of understanding. Besides, the scarcity of mutations fixing in the population also induces some stochasticity.

**Résumé :** Les résultats récents présentés dans cet article décrivent des limites d'échelles de modèles individus-centrés. Grâce à eux, nous allons mettre en lumière les différentes échelles de temps caractéristiques des variations démographiques et de la sélection naturelle. L'objectif n'est pas d'être exhaustif, autant au niveau mathématique que biologique. Néanmoins, ces résultats sont très complémentaires car ils fournissent un aperçu des principales échelles de temps d'intérêt. Ils englobent notamment l'échelle de temps de la vie d'un individu, celle plus longue où la population peut être décrite comme une entité avec ses caractéristiques propres, celle démographique bien sûr, et au moins quatre autres échelles sur lesquelles la sélection joue un rôle. La dynamique limite est généralement déterministe. Pour autant, puisque l'action de la sélection se base sur l'aléa présent dans l'histoire des individus, les probabilités apparaissent comme un élément clé de compréhension. Par ailleurs, l'exceptionnalité des événements de fixation de mutations induit aussi une stochasticité.

## Introduction

For more than fifteen years, there has been a significant activity around a new probabilistic framework for the mathematical modeling of ecology, population genetics and character evolution. The Individual-Based Models (IBM) are a priori the best representation for the validation of a macroscopic model of population dynamics. These processes detail at the individual level the births, the deaths and the interactions among the population. The description may as well precise the

---

\*Aix-Marseille Université, CNRS, Centrale Marseille, I2M, UMR 7373 13453 Marseille, France, email : aurelien.velleret@univ-amu.fr

migration patterns, the ageing or the competition for resources between the individuals. This setting is clearly the closest to the actual simulations designed by computational biologists to validate theoretical models, or simply some predictions. In this view, such setting is also the one with the least number of simplifying assumptions. It also certainly the one where calculations of event probabilities are the most complicated. One of the first aim of such probabilistic modeling has actually been to connect these individual-based models and simulations to much simpler systems of Ordinary Differential Equations (ODE) or Partial Differential Equation (PDE) models.

To describe the natural selection on heritable traits, systems of ODE usually provide the most classic models to simply express their effects. They have been refined by introducing PDE to deal with continuum of the traits among the population. Thanks to these models, one can cover most of the actual mathematical modeling of ecology, population genetics and character evolution. Yet, this deterministic approach does not take any account of the large amount of variability observed when reproducing similar experiments. Parts of this noise is due to external perturbations. Many Stochastic Differential Equations (SDE) have been introduced to include these effects. They usually assume that the parameters expressed in the systems of ODE / PDE are themselves subject to stochasticity. The other main component of this noise is internal and inherent to the discreteness of the system : the selection between traits emerges through the competition between a finite number of individuals. Taking this component into account is the purpose of these IBM.

To justify the simpler models from the IBM, the usual and main assumption is to assume the population so large that the law of large number makes the probabilistic fluctuations negligible. Yet, to be completely accurate, such assumption would mean that every single individual pattern should be present among the population, with quite a precise accuracy and more importantly with a large number of occurrences. Given the finite population sizes the ecologists are interested in, an elementary description may well be accurately designed while a finer structuring of the population is much more demanding.

It is thus only natural that, as mathematicians, we try to describe simplified behavior through an asymptotic in the number of generations. It is at the core of Darwin's theory of natural selection to assume a separation of time-scale between the apparently incidental variability among contemporary individuals and the evolution of the traits, and even of the species. The goal is surely to obtain exact results of convergence to a non-trivial process for the new dynamics in a new time-scale. It is however very challenging, and it might happen that an approximation cannot be justified this way, while still being helpful. There is also no reason to restrict ourselves purely to deterministic limiting processes, all the more if one can get a SDE governing the first order of fluctuations. Such a next order helps notably to precise the conditions of validity of the deterministic approximations.

An archetypal example is the stochastic approximation of the population size in a continuous setting (i.e. with Brownian-type fluctuations). Beyond such result is the idea that individual events (of births or deaths) have quite a negligible impact on the population on a whole, yet the randomness still occurs in the varying accumulation of such effects. Thus, to obtain the exact convergence, one needs –besides the population size going to infinity– to also amplify the frequency of birth and death events. It means notably that the selective effects appear in a much longer time-scale than the life-expectancy of the individuals (cf. notably chapter 3 in [4]). The population size scale provides then the essential relation between the parameters at the individual time-scale (associated for instance to a birth rate of 1) and at the population level (cf. section 1). In case there is more

than one parameter to adjust, the analysis may be even less intuitive and may even lead to different behaviors associated to different scalings.

Our purpose with the following illustrations is to highlight the variety of time-scales involved in population models. We will also particularly focus on these aspects of having sufficiently large a population size to relate the individual-based random models to the population dynamics.

In Section 1, we discuss this notion of individual-based models and how to relate the individual histories to the law that governs these behaviors. In this view, we shall refer to [16], which gives some insight on the number of generations involved before one can actually distinguish distinct population models. On a shorter time-scale, natural selection may favor some trajectories. But those should be observed with close frequencies between distinct populations and the heritability of such a selection is a priori very small. As presented for instance in [16], natural selection seen as some statistical estimation of the traits requires a sufficiently large population history. For the trait to be efficiently selected, one needs to wait for the whole range of history to be expressed. A very favorable combination of trait and environment may for instance not be so beneficial in the long term if most of the descendants shall be exposed to much less favorable environment. This "sample size" shall thus be larger when there are strong correlations between the successive generations. In the cases where such correlations are sufficiently weak, Hoffmann and Marguet (cf. [16]) study how to take them into account in the statistical analysis. They could indicate the precision of some estimators in terms of the size of the genealogical tree on which the observations are indexed. The focus in [16] is on the random process governing birth events, for which they propose the first statistical analysis in a structured population.

Although this individual-based dispersion is thus usually neglected (only some averaged effect is considered), we shall see in Section 2 that it may have a significant effect when two components of selection are competing. Both the strength and the time-scale of selection between two sub-populations may in fact involve the amount of a priori non-selective variability. Assuming however that this variability has only a small effect on the 'local' reproductive value of the individuals and rescaling time properly, Champagnat and Henry could describe in [9] the evolution history as an almost deterministic behavior. This will be our concern in Section 3. Finally, it may happen that the natural selection rely on the emergence of very rare mutations. The associated time-scale is then given by the mean effect of successful mutations (that invade the population). Such models will be taken from [7], [8], [5] and discussed in Section 4. One may either include the randomness of the mutation emergence or consider it in an averaged deterministic manner like in Section 3.

## 1 Recognition of the population from the individuals

At an individual-based level, each individual is characterized by some value  $x \in \mathcal{X}$ . Typically  $\mathcal{X} \subset \mathbb{R}^k$  or  $\mathcal{X} \subset \mathbb{R}^k \times F$ , where we may consider  $k$  characteristics such as the size, the spatial position, the age, the amount of certain proteins, of certain resources, of parasites and so on, while  $F$  denotes a finite (or numerable) set of classes, such as the sex, the eye color etc. Before the death of the individual, this trait  $x$  may be constant, but it can also evolve according to some stochastic differential equation (SDE). The interaction with the other individuals can then appear in the parameter of this equation, possibly depending on the trait  $x'$  of the other individuals around. In order to have a Markov process, the death of the individuals happens usually at a rate  $d$ , that may

depend first on the trait  $x$  of the individual, but also on the population on a whole. We recall that we could include the age of the individual in the information carried by  $x$ , so that there is almost no restriction. Death at fixed age is also not difficult to include in the model. The same kind of dependencies may be considered for the births, and apply to the number of offspring and their states at birth. We then need an additional kernel to model the birth events conditionally on the number of offspring. Independent exponential variables or Poisson Point Processes are thus usually a very convenient way to encode all of these events. In a nutshell, it gathers quite any population model we could compute (or at least approximate when dealing with solutions to SDEs), as soon as one can follow the individuals one by one. We refer notably to Subsection 4.2 for a more concrete description of such measure-valued process.

The model in [16] is concerned with a trait supposedly governed by a stochastic flow of the form :

$dX_t = b(X_t) dt + \sigma(X_t) dB_t$ , where  $B$  refers to a standard Brownian Motion, and  $(X_t)$  evolves on  $\mathcal{X} \subset \mathbb{R}^k$ . It can for instance represent the size of the cell or the propagation of a parasite (that we take here as an archetype). The main issue is here to estimate the effect of the amount of parasites in a cell on its reproduction. Note that the individual level is the one of a given cell, and not of a given parasite, as it would be in a completely discretized model. Yet, the number of parasites is assumed to be so large that a continuous description by a random process is a well-justified simplification. The estimation is focused on the effect of these parasites on the birth rates of the hosts and on the law of the offspring. We restrict ourselves to the continuous trait-space in order to exploit the regularity of the estimated functions.

## 1.1 Discussion on the assumptions in [16]

### 1.1.1 A process on an incomplete tree

The main specificity of this statistical approach is to deal with the dependency given by the genealogy of the individuals. Indeed, the more parasites the mother-cell has, the more are transmitted to its daughter-cells. With these experiments where one follows each cell individually, one can also obtain easily the entire associated lineages. Yet, many of these cells are not observable anymore because of the design of the experimental process. Although the authors do not mention it, the death of some cells is likely to be included. The biological data is thus indexed by some incomplete tree, with binary divisions. In order to relate the precision of the estimators to the number of birth events, the authors assumed that the generation size evolves either asymptotically with a given growth rate, or stays bounded :

**Definition 1.1.1.** *Define the potential generation  $n$  as  $G_n$ . Note that  $\text{Card}(G_n) = 2^n$ .*

*A regular incomplete tree is a family of subsets  $U_n$  of the complete tree at generation  $n$  (for  $n \geq 1$ ) such that :*

*(i) the parent of any individual of  $U_n$  is also in  $U_n$*

*and (ii) for some  $0 \leq \rho \leq 1$  :*

$$0 < \liminf_{n \rightarrow \infty} 2^{-\rho n} \text{Card}(U_n \cap G_n) \leq \limsup_{n \rightarrow \infty} 2^{-\rho n} \text{Card}(U_n \cap G_n) < \infty.$$

In the case  $\rho \in (0, 1]$ , the data characterize a growing population of infected cells. It corresponds for instance to an experiment where one lets these cells freely duplicate in a rich medium. In the case  $\rho = 0$ , one rather expect some population size at equilibrium, although it is not required here.

It also corresponds to the experiments where only one cell is followed after each division (which is the case in some experiments in microfluidic environment).

In the following results, this indexation tree is considered as a given. The expectations are taken on the random dynamics of parasitic growth inside the cells and of splitting between the two daughter-cells. The purpose of such estimations is to precise the law of birth events but certainly not conditionally on the specific realization of the tree. It means that there is an implicit assumption that the shape of the indexation tree is independent of the parasitic dynamics.

With a single lineage experiment, this is not really an issue. But if one has to consider natural death events, it seems much more questionable. It appears a bit surprising to have a birth rate depending on the number of parasites while the probability of giving birth in one life-span is independent of it. A constant death rate would be a more natural assumption than a prescribed tree. Yet, it would be much more difficult to analyze and would possibly not reflect very well the design of the experiment.

### 1.1.2 Ergodicity of the trait

The other main assumption in [16] is the very rapid convergence to the ergodic measure  $\nu$  in the transmission of the trait at birth from the parent to its offspring, driven by the generator  $Q$ . The convergence of the estimators is stated uniformly as long as there exist  $C > 0$  and a function  $V$  such that for any  $m \geq 1$ , and  $\varphi$  satisfying  $\|\varphi\|_V := \sup_x |\varphi(x)|/[1 + V(x)] < \infty$  :

$$\|Q^m(\varphi) - \nu(\varphi)\|_V \leq C 2^{-m} \|\varphi - \nu(\varphi)\|_V. \quad (1)$$

The set of such transmission process  $Q$  is denoted  $\mathbb{Q}_{1/2}$ . Note in this view that the authors choose to focus exclusively on the trait at birth for their statistical estimations.

### 1.1.3 Regularity and confinement of the stochastic process

By Assumption 2 on the drift and diffusion coefficient, the authors ensure that the trait  $\mathcal{X}$  stays somewhat confined around 0, with a uniformly elliptic and non-singular diffusion. By Assumption 3 on the birth rate, they ensure some regularity in  $x$ , a boundedness condition on the potential explosion of birth events and they prevent the vanishing of these events. Finally, by Assumption 4 on the splitting of  $x$  at birth between the two newborns, they ensure a lower-bounded density on the fragmentation parameter, and prevent too asymmetric partitioning. The authors mention that this assumption could probably be relaxed.

## 1.2 Main results for the estimations of the Generation kernel and the birth rate

For any function  $\psi : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ , and  $y \in \mathcal{X}$ , consider  $\psi_\star(y) := \sup_{x \in \mathcal{X}} |\psi(x, y)|$ ,  $\psi^\star(x) := \sup_{y \in \mathcal{X}} |\psi(x, y)|$  and, given any positive measure  $\rho$  on  $\mathcal{X}$ , and noting  $\wedge$  for the minimum :

$$|\psi|_\rho := \int_{\mathcal{X}^2} |\psi(x, y)| \rho(dx) dy + \left( \int_{\mathcal{X}^2} |\psi(x, y)| dx dy \wedge \int_{\mathcal{X}} |\psi_\star(y)| dy \right)$$

$$\mathcal{M}_{U_n}(\psi) := \frac{1}{\text{Card}(U_n^\star)} \sum_{u \in U_n^\star} \psi(X_{u-}, X_u),$$

where  $U_n \setminus U_n^*$  is only the root of the tree, and  $X_{u-}, X_u$  denotes respectively the trait at birth of the parent of  $u$  and the one of individual  $u$  itself. Note also that the transition from  $X_{u-}$  to  $X_u$  is given by the kernel  $Q(X_{u-}, dx)$ .

**Proposition 1.2.1.** *Work under Assumptions 2, 3 and 4. Let  $\mu$  be a probability measure on  $\mathcal{X}$  such that  $\mu(V^2) < \infty$ . Let  $\psi : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  a bounded function such that  $\psi_\star$  is compactly supported. If  $U_n$  is a regular incomplete tree, the following estimate holds true:*

$$\mathbb{E}_\mu[(\mathcal{M}_{U_n}(\psi) - \nu(Q\psi))^2] \leq \text{Card}(U_n)^{-1}(|\psi^2|_{\mu+\nu} + |\psi^\star\psi|_\mu + (1 + \mu(V^2))|\psi_\star|_1|\psi|_\nu), \quad (2)$$

where the symbol  $\leq$  means up to an explicitly computable constant that depends only on  $\text{supp}(\psi_\star)$  as long as  $Q \in \mathbb{Q}_{1/2}$ . More generally, it would also depend on  $Q$ .

For the following theorem, we assume that the operator  $Q$  has a density  $(q(x, y))_{(x, y) \in \mathcal{X}^2}$  w.r.t. Lebesgue measure on  $\mathcal{X}$ , with some Hölder regularity. We refer to [16] for the precise definition of the sets  $\mathbb{Q}_{1/2}^{\alpha, \beta}(R)$ , where the  $R > 0$  defines a bound on the Hölder regularity of respectively order  $\alpha$  in  $x$  and  $\beta$  in  $y$ . Recall that the "1/2" refers to the property (1). Given such a regularity, the authors propose to adjust the order of the estimation kernel and explain how to choose the associated window sizes given  $|U_n|$  (for some large  $n \geq 1$ ). The estimation of  $q$  also depends on a threshold  $\varpi_n$  that is also to adjust.

**Theorem 1.1.** *Work under Assumptions 2, 3 and 4. Assume that the initial distribution  $\mu$  is absolutely continuous w.r.t. the Lebesgue measure with a locally bounded density function and satisfies  $\mu(V^2) < \infty$ . Let  $\alpha, \beta > 0$ . Specify  $\hat{\nu}_n(y_0)$  and  $\hat{q}_n(x_0, y_0)$  as previously evoked. Then, for any  $\varrho$ -regular incomplete tree  $U_n$  and any  $R > 0$ ,*

$$\sup_{Q \in \mathbb{Q}_{1/2}^{\alpha, \beta}(R)} (\mathbb{E}_\mu[(\hat{\nu}_n(y_0) - \nu(y_0))^2])^{1/2} \leq \text{Card}(U_n)^{-\beta/(2\beta+1)}.$$

$$\text{and} \quad \sup_{Q \in \mathbb{Q}_{1/2}^{\alpha, \beta}(R)} (\mathbb{E}_\mu[(\hat{q}_n(x_0, y_0) - q(x_0, y_0))^2])^{1/2} \leq \varpi_n^{-1} \text{Card}(U_n)^{-s(\alpha, \beta)/(2s(\alpha, \beta)+1)}$$

hold true, where  $s(\alpha, \beta)^{-1} := (\alpha \wedge \beta)^{-1} + \beta^{-1}$  is the effective anisotropic smoothness associated with  $(\alpha, \beta)$ .

Note that the rate of convergence depends on the regularity of  $q$  because the higher it is, the more useful information we can gather from observed transitions from vicinities of  $x$  to vicinities of  $y$ .

The estimation of the division rate is much more complicated. The study is thus restricted to diffusion of the trait on a compact space, so that (2) can be strengthened with a right-hand side only depending on the  $L^\infty$ -norm of  $\psi$ . Again, we restrict the study to  $Q \in \mathbb{Q}_{1/2}$ , according to assumption 14. A parametric approach is also considered, coded with the parameter  $\vartheta$ . By Assumption 17, the authors prevent any degeneracy and singularities of the birth rate. Moreover, this birth rate  $B$  is assumed globally monotone (i.e. uniformly in  $\mathcal{X}$ ) with  $\vartheta$ . By Assumption 19, they assume some upper-bounds on the derivative of  $B$  along  $\vartheta$  up to the third order. Finally, by Assumption 20, the Fisher matrix  $\Psi(\vartheta)$  is non-singular. Although  $\vartheta$  is possibly considered multi-dimensional, the monotonicity conditions seem in practice to reduce the estimation to a family of laws for  $B$  that is essentially one-dimensional.

**Theorem 1.2.** *Work under Assumptions 2, 4, 14, 17, 19 and 20. For every  $\vartheta$  in the interior of  $\Theta$ , if  $U_n$  is a  $\varrho$ -regular incomplete tree :*

$$\sqrt{\text{Card}(U_n)} \times (\hat{\vartheta}_n - \vartheta) \rightarrow \text{N}(0, \Psi(\vartheta)^{-1})$$

*in distribution as  $n \rightarrow \infty$ , where  $\text{N}(0, \Psi(\vartheta)^{-1})$  denote the  $d$ -dimensional Gaussian distribution with mean 0 and covariance the inverse of  $\Psi(\vartheta)$ .*

Note that we find in this case the classical rate of convergence with order the square root of the number of observations, as for classical Markov-Chains. It hints at the fact that every observed transition from parents to offspring contribute to the estimation of  $\vartheta$ .

**Concluding remarks :**

As mentioned after the assumption on the genealogical tree, considering this tree as prescribed is much more convenient but questionable. The fact that the convergence results of Theorems 1.1 and 1.2 depend on  $U_n$  only through its cardinal indicates that these results shall be quite robust concerning the specific realization of the tree. It does not exclude however that the estimators are possibly biased by the fact that certain transmission patterns might lead to a greater probability of survival (of the lineage). For instance, it may be beneficial to divide early, in order not to let the parasites divide for long before the next division. This bias depends on how strongly these effects may have shaped the genealogical tree. Besides, the estimation of the biased birth rate may be the main interest. It only means that one infers the dynamics of a "successful" lineage, with luck playing a role in this success, as well as the environmental conditions.

This issue of estimating individual parameters by looking at the population as a whole leads to similar issues of independence. Notably, the population around the focal individual can be considered as a component of its environment. One may wish that the parameters of its dynamics depend on the interaction it has with this extended environment. The models become much simpler if one can neglect the precise interactions between the individuals, are rather consider averaged interactions. In a limit of a large population size, for a system at equilibrium, such assumption may be justified by the Law of Large Number. Indeed, the effects of the interactions are averaged, as long as they are not too local, and thus almost globally constant. In results of propagation of chaos, such property is generalized in cases where the population is not at equilibrium. We refer to Sznitman's lecture at Saint-Flour [23] for an overview on this topic. In such a limiting case, known as McKean-Vlasov equations, the law of the process itself acts on the individual dynamics, together with fluctuations specific to this individual. At equilibrium, this law is all the more constant that the population size is large. Thus, we can deal with the effects of these interactions through some hidden parameters (like an effective death or birth rate). And as for the analysis of this section, estimating them by simply looking at the death and reproduction times would produce a bias if they depend on heritable factors. But this bias is presumably small if the heritability is weak.

To study the evolution of population traits, that is for instance  $\vartheta$  if mutations could alter this value, it is very classical to assume that the population size is large. Already even demographic changes of the whole population is associated to an amount of birth (and death) events of the order of the number of individuals (also depending on the observed stochasticity). Since even those are generally considered in a way shorter time-scale than the evolutionary process, there is clearly time

for the individual fluctuations to be averaged in terms of the heritable traits to be selected upon. Of course, for a variable environment, some periodicity effects may alter such an homogenization. When the environment itself is affected by the traits of the individuals, like in the next model, it may also not be so clear what an averaged effect would be.

## 2 Selection with two levels

The model presented in this section is an illustration that selective effects may be strongly dependent upon the precise local ecological dynamics (cf. [24]). For simplicity, we focus here on the competition between two types in a population of fixed and large size. We generally consider neutral the fluctuations in the proportion due to the inexact compensation between events of deaths and births occurring with the same large rate. Biologists usually refer to it as the genetic drift, and often neglect it in first estimation for large populations. One shall see however that this component may not be neutral at all in a model of selection at two levels : individuals and the groups they constitute. Having little of the above mentioned fluctuations inside each group strongly hinders the response to selection at the group level.

Our main interest is when these two components of selection are competing. In a nutshell, the issue is to know if and how the inefficient or cheater individuals can be regulated through natural selection at this group level. Another crucial requirement for such a confrontation is a strong heritability for the groups when they split. Indeed, even small transmission between these communities is known to greatly disrupt the stability of cooperative strategies (cf. for instance [25]).

The individual-based model is taken from [18], where a formalism for group selection is introduced. Each group is made up of  $n \in \mathbb{N}$  individuals. There are two types of individuals: the type I individuals are selectively advantageous at the individual (I) level and the type G individuals are selectively advantageous at the group (G) level. The replication and the selection occur concurrently at the individual and group level according to coupled Moran processes. Type I individuals replicate at rate  $w_I(1+s)$ ,  $s \geq 0$  and type G individuals at rate  $w_I$ . When an individual gives birth, another individual in the same group is selected uniformly at random to die. To reflect the antagonism at the higher level of selection, groups replicate at a rate that depends on the number of type G individuals they contain. As a simple case, we take this rate to be  $w_G \times [1 + r(k/n)]$ , where  $k/n$  is the fraction of individuals in the group that are type G,  $r(x)$ ,  $x \in [0, 1]$  is the selection coefficient at the group level. Like at the individual level, the population of groups is maintained at  $m$  by selecting a group uniformly at random to die whenever a group replicates. The offspring of groups are assumed to be identical to their parent.

Two limits of large population are justified from the individual-based model in [18] : both include a term for each selective force, but only the second includes effects of neutral fluctuations of reproduction, both within the hosts and between hosts. By choosing the right scaling of the parameters, one could actually choose to keep any combination of these 4 terms. In [24], another limit is considered, where the neutral fluctuations are kept only inside the hosts, together with the selective forces. Although such limit can be mathematically justified, it is undoubtedly more realistic to keep also neutral fluctuations in the reproduction of groups. Yet, the analysis is notably much more technical with much less clear interpretation. In this more complete model, the ulti-



mate fixation of one type is expected to happen almost surely, with always a positive probability for each. Yet, one hopes that for large enough group populations, the limit obtained by neglecting these fluctuations provides an interesting view on the main features of the dynamics.

Let  $X_t^i$  be the number of type G individuals in group  $i$  at time  $t$ . Then :

$$\mu_t^{m;n} := \frac{1}{m} \sum_{i \leq m} \delta_{X_t^i/n}$$

is the empirical measure at time  $t$  of the proportion of type G by group, for a given number of groups  $m$  and individuals per group  $n$ . Here,  $\delta_x$  is the Dirac at  $x$ . The  $X_t^i$  are divided by  $n$  so that  $\mu_t^{m;n}$  is a probability measure on  $E_n := [0; 1/n; \dots; 1]$ . For fixed  $T > 0$ ,  $(\mu_t^{m;n})_{t \leq T} \in D([0; T]; \mathcal{M}_1(E_n))$ , the set of càdlàg processes on  $[0; T]$  taking values in  $\mathcal{M}_1(E_n)$ , where  $\mathcal{M}_1(S)$  is the set of probability measures on a set  $S$ . With the particle process described above,  $\mu_t^{m;n}$  has generator

$$(\mathcal{L}^{m;n} \psi)(v) = \sum_{i,j} (w_I R_I^{i,j} + w_G R_G^{i,j})(v) \times (\psi[v + 1/m (\delta_{j/n} - \delta_{i/n})] - \psi[v])$$

where  $\psi \in C_b(\mathcal{M}_1([0; 1]))$  is a bounded continuous function, and  $v \in \mathcal{M}_1(E_n) \subset \mathcal{M}_1([0; 1])$ . The transition rates  $(w_I R_I^{i,j} + w_G R_G^{i,j})$  are given by

$$R_I^{i,j}(v) := \begin{cases} m v(i/n) i (1 - i/n) (1 + s) & \text{if } j = i - 1; i < n, \\ m v(i/n) i (1 - i/n) & \text{if } j = i + 1; i > 0 \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad R_G^{i,j}(v) := m v(i/n) v(j/n) (1 + r[j/n]). \quad (3)$$

$R_I^{i,j}$  and  $R_G^{i,j}$  represent respectively individual-level events and group-level events.

**Theorem 2.1.** *Suppose  $n, m \rightarrow \infty$  and the rates  $w_I/n \rightarrow \omega_I$ ,  $n s \rightarrow \sigma$ ,  $w_G$  and  $\{r(x)\}_{x \in [0,1]}$  constant. Suppose the particles in the process  $\mu_t^{m;n}$  are initially independently and identically distributed according to the measure  $\mu_0^{m;n}$ , where  $\mu_0^{m;n} \rightarrow \mu_0$  as  $m, n \rightarrow \infty$ . Then,  $\mu_t^{m;n}$  converges weakly to  $\mu_t \in D([0; T]; \mathcal{M}_1([0; 1]))$ .  $\mu_t$  is the unique solution with initial condition  $\mu_0$  of the differential equation :*

$$\partial_t \langle \mu_t | f \rangle = \langle \mu_t | \mathcal{L}_{WF} f \rangle + \langle \mu_t | r f \rangle - \langle \mu_t | f \rangle \times \langle \mu_t | r \rangle, \quad (4)$$

$$\text{where } \langle \mu | f \rangle := \int_0^1 \mu(dx) f(x) \text{ and } \mathcal{L}_{WF} f(x) = -s x (1 - x) \partial_x f(x) + (\sigma^2/2) \cdot x (1 - x) \partial_{xx}^2 f(x). \quad (5)$$

## 2.1 Definition as a conditional law

We describe the solution of such equation through a Feynman-Kac penalization of the stochastic process with generator  $\mathcal{L}_{WF}$ . Consider thus  $X_t$ , with initial condition  $X_0 \sim \mu_0$  and the solution to the problem :

$$dX_t := -s X_t (1 - X_t) dt + \sigma \sqrt{X_t (1 - X_t)} dB_t. \quad (6)$$

The existence and uniqueness of such process can be found e.g. in chapter 5.3.1 of [11]. It is linked there with an individual-based model, namely the neutral 2-allele Wright-Fischer Markov Chains. We also consider the Feynman-Kac penalization :

$$Z_t := \exp \int_0^t r(X_s) ds, \quad \langle \mu_t | f \rangle := \mathbb{E} [f(X_t) Z_t] / \mathbb{E} [Z_t],$$

for any  $f \in \mathcal{C}_b^2$ . Note that  $r$  is bounded so that for any  $t > 0$ ,  $\mathbb{E}(Z_t) \in (0, \infty)$  and these definitions make sense.

By the Ito formula, it is not difficult to see that  $(\mu_t)$  indeed defines a solution to (4).

The following presentation give another view on this law  $\mu_t$ , that will simplify our notations and clarify our point. Since subtracting a constant to  $r$  does not change the value of  $\langle \mu_t | r f \rangle - \langle \mu_t | f \rangle \langle \mu_t | r \rangle$ , we assume in the following that  $r \leq 0$ , so that we can represent it as a death rate.  $Z_t$  can then be interpreted as the probability that the process has survived until time  $t$ , while confronted to a death rate of  $r$ , conditionally on  $(X_u)_{u \geq 0}$ . More formally, with  $T_\partial \sim \text{Exp}(1)$  (exponential r.v. with rate 1), we can define the extinction time as :

$$\tau_\partial := \inf \{t \geq 0; -\ln(Z_t) \geq T_\partial\}, \quad \text{so that } \mathbb{P}(t < \tau_\partial | X_u, u > 0) = \mathbb{P}(-\ln(Z_t) < T_\partial | X_u, u \leq t) = Z_t$$

$$\langle \mu_t | f \rangle = \mathbb{E} [f(X_t); t < \tau_\partial] / \mathbb{P}[t < \tau_\partial] = \mathbb{E}(f(X_t) | t < \tau_\partial).$$

Any equilibrium of  $\mu_t$  is thus a quasi-stationary distribution (QSD) of  $X$  under the death rate  $r$ . Applying the last techniques on QSD thus enables us to obtain the limiting behavior and the speed of convergence of  $\mu_t$ .

Similarly, the hitting times of 0 and 1 are denoted  $\tau_0$  and  $\tau_1$ . Since they are absorbing, it is natural to be interested in the law of the marginal with these extended extinction times :

$$\tau_{0,\partial} := \tau_\partial \wedge \tau_0, \quad \tau_{1,\partial} := \tau_\partial \wedge \tau_1, \quad \tau_{0,1} := \tau_0 \wedge \tau_1, \quad \tau_{0,1,\partial} := \tau_\partial \wedge \tau_0 \wedge \tau_1.$$

Then :

$$\mu_t = x_t^0 \delta_0 + x_t^1 \delta_1 + x_t^\xi \xi_t, \quad \text{where } x_t^\xi := \mathbb{E} [Z_t; t < \tau_{0,1}] / \mathbb{E} [Z_t],$$

$$x_t^0 := \mathbb{E} [Z_{\tau_0} \exp[-r_0(t - \tau_0)]; \tau_0 < t] / \mathbb{E} [Z_t], \quad x_t^1 := \mathbb{E} [Z_{\tau_1} \exp[-r_1(t - \tau_1)]; \tau_1 < t] / \mathbb{E} [Z_t],$$

$$\langle \xi_t | f \rangle := \mathbb{E} [f(X_t) Z_t; t < \tau_{0,1}] / \mathbb{E} [Z_t; t < \tau_{0,1}] = \mathbb{E} [f(X_t) | t < \tau_{0,1,\partial}], \quad \text{for any } f \in \mathcal{C}_b^2.$$

Remarks : • It is not difficult to generalize the model to include frequency dependent effects of selection at the individual level. We would then replace  $s > 0$  by some smooth function  $(s(x))_{x \in [0,1]}$ . Generalizations of  $\sigma$  as a function are also not a mathematical issue. Yet a priori, it does not seem biologically justified.

• This Feynman-Kac penalization is the analogous of the many-to-one formula described in the case of Markov processes associated to branching Galton-Watson Trees (cf. [3]). Behind is the idea that there is a bias towards larger offspring in the genealogy of a typical individual in the population, favoring the individuals with a larger birth rate. Contrary to [3], where no interaction occurs between lineages, one cannot expect to represent the law of  $\mu_t$  directly through another Markov Process, i.e. without penalization (note in (4) the quadratic term in  $\mu_t$ ).

## 2.2 QSDs and exponential convergence

In any case,  $\delta_0$  and  $\delta_1$  are QSDs for the extinction  $\tau_\partial$ , i.e. stable distributions for the dynamics given by (4).

We define the semi-groups associated to our different extinctions :

$$\mu A_t(dx) := \mathbb{P}_\mu(X_t \in dx \mid t < \tau_\partial), \quad \mu A_t^{01}(dx) := \mathbb{P}_\mu(X_t \in dx \mid t < \tau_{0,1,\partial}), \quad \mu A_t^1(dx) := \mathbb{P}_\mu(X_t \in dx \mid t < \tau_{1,\partial})$$

**Proposition 2.2.1.** *There exists a unique QSD  $\alpha \in \mathcal{M}_1[(0,1)]$  and a survival capacity of  $\eta$  associated to the extinction  $\tau_{0,1,\partial}$ . With the associated extinction rate  $\rho_\alpha$ , it means :*

$$\forall t > 0, \quad \mathbb{P}_\alpha(X_t \in dx ; t < \tau_{0,1,\partial}) = \exp[-\rho_\alpha t] \alpha(dx), \quad \forall x, \quad \eta(x) = \exp[\rho_\alpha t] \mathbb{E}_x(\eta(X_t) ; t < \tau_\partial)$$

Moreover, we have the following exponential convergences at rate  $\zeta > 0$  :

$$\exists C > 0, \quad \forall \mu \in \mathcal{M}_1[(0,1)], \quad \|\mu A_t^{01} - \alpha\|_{TV} \leq C \exp[-\zeta t]. \quad (7)$$

$$\exists C' > 0, \quad \forall \mu \in \mathcal{M}_1[(0,1)], \quad |\exp[\rho_\alpha t] \mathbb{P}_\mu(t < \tau_{0,1,\partial}) - \langle \mu \mid \eta \rangle| \leq C' \exp[-\zeta t] \quad (8)$$

$$\text{a fortiori } \eta(x) := \lim_{t \rightarrow \infty} \exp[\rho_\alpha t] \mathbb{P}_x(t < \tau_{0,1,\partial}) \quad (9)$$

$$\text{and } \|\eta_\bullet\| := \sup_{\{x \in (0,1), t > 0\}} \exp[\rho_\alpha t] \mathbb{P}_x(t < \tau_{0,1,\partial}) < \infty \quad (10)$$

Let  $\rho_0 = -r_0$  ( $\rho_1 = -r_1$ ) the extinction rate of  $\delta_0$  (resp.  $\delta_1$ ). We show in the following that the long-time behavior of the process with only the local extinction rate depends mainly on  $\rho_\alpha$ ,  $\rho_0$  and  $\rho_1$ .

In the following convergences, we will often have uniform bounds for probability measures belonging for some  $n \geq 1$  and  $\xi > 0$  to :

$$\begin{aligned} \mathcal{M}_{n,\xi} &:= \{\mu \in \mathcal{M}_1([0,1]) \mid \mu[1/n, 1] \geq \xi\}, \quad \bigcup_{n,\xi} \mathcal{M}_{n,\xi} = \mathcal{M}_1([0,1]) \setminus \{\delta_0\}. \\ \text{or in } \mathcal{M}_{n,\xi}^{0,1} &:= \{\mu \in \mathcal{M}_1([0,1]) \mid \mu[1/n, 1 - 1/n] \geq \xi\}, \quad (n \geq 3, \xi > 0) \\ \bigcup_{n,\xi} \mathcal{M}_{n,\xi}^{0,1} &= \mathcal{M}_1([0,1]) \setminus \{u \delta_0 + (1-u) \delta_1 \mid u \in [0,1]\}. \end{aligned}$$

### 2.2.1 Despite the fixation events, polymorphic groups maintain themselves in the population

**Proposition 2.2.2.** *Assume that  $\rho_\alpha < \rho_0 \wedge \rho_1 := \rho$ . Then, there is only one stable QSD  $\alpha_{0,1}$ , with convergence rate  $\rho - \rho_\alpha$ , i.e. :*

$$\forall n \geq 1, \quad \forall \xi > 0, \quad \exists C_{n,\xi} > 0, \quad \forall \mu \in \mathcal{M}_{n,\xi}^{0,1}, \quad \|\mu A_t - \alpha_{0,1}\|_{TV} \leq C_{n,\xi} \exp[-(\rho_\alpha - \rho) t],$$

where  $\alpha_{0,1}$  has extinction rate  $\rho_\alpha$  and is given as  $\alpha_{0,1} = y_0 \delta_0 + y_1 \delta_1 + y_\alpha \alpha$  with :

$$\frac{y_0}{y_\alpha} = \frac{\rho_\alpha \times \mathbb{P}_\alpha(\tau_0 = \tau_{0,1,\partial})}{(\rho_0 - \rho_\alpha)}, \quad \frac{y_1}{y_\alpha} = \frac{\rho_\alpha \times \mathbb{P}_\alpha(\tau_1 = \tau_{0,1,\partial})}{(\rho_1 - \rho_\alpha)},$$

and of course  $y_0 + y_1 + y_\alpha = 1$ .

If  $\rho_1 < \rho_0$ , for any initial condition  $\mu_0 = u \delta_0 + (1 - u) \delta_1$  with  $u \in (0, 1)$ ,  $\mu_t$  converges at rate  $\rho_0 - \rho_1$  to  $\delta_1$ .

If  $\rho_1 = \rho_0$ , then any such distribution is a QSD with the extinction rate  $\rho_0$ .

Pure groups are continuously generated from polymorphic groups without any reversed transition. Yet, in this case, these polymorphic groups are sufficiently selected upon through their better survival to persist in the population. Pure groups are like the remnants of these polymorphic groups : their proportion reaches a steady state where their faster decay compensate for the fixation rate from polymorphic groups. The stabilization of the polymorphic profile induces the convergence of both this fixation rate and of the maintenance rate of the polymorphic groups to  $\rho_\alpha$ .

In any case, the polymorphism is maintained by any sufficiently large group selection favoring it, since :

**Proposition 2.2.3.** *Given any  $\sigma > 0$ ,  $s \geq 0$ , and a bounded continuous function  $r^0$  with its maximum only in the interior of  $(0, 1)$ , there exists a critical value  $R_\vee > 0$  such that for any  $R > R_\vee$  and considering the system with  $r = Rr^0$ , we indeed have  $\rho_\alpha < \rho_0 \wedge \rho_1$ .*

Conversely, when the group selection is too small, the polymorphic groups cannot be maintained :

**Proposition 2.2.4.** *Conversely, given any  $\sigma > 0$ ,  $s \geq 0$ , and a bounded measurable function  $r^0$ , there exists a critical value  $R_\wedge > 0$  such that for any  $R < R_\wedge$  and considering the system with  $r = Rr^0$ , we indeed have  $\rho_0 \wedge \rho_1 < \rho_\alpha$ .*

In a similar way :

**Proposition 2.2.5.** *Given any  $s > 0$  and any bounded function  $r$ ,  $\lim_{\sigma \rightarrow \infty} \rho_\alpha(\sigma) = +\infty$ .*

### **Conditions for the polymorphism to be maintained :**

For this maintenance rate  $\rho_\alpha$  to be higher than the decay of pure groups, it is clearly necessary that  $r$  is maximal in  $(0, 1)$ . Of course, it is not sufficient, because first of the "neutral fluctuations" ( $\sigma$ ) in the proportion of both types and second because of the selection effects inside each group. As stated in Proposition 2.6, too large  $\sigma$  would namely induce too large a rate of fixation (to 0 or 1). Even strong effects of selection through  $r$  would then be unable to make their reproduction large enough to compensate for this loss.

But on the other hand, when there are internal selective effects pushing towards type  $I$  individuals, having small neutral fluctuations limits the effectiveness of the selection at the group level. Indeed, all groups with similar initial condition evolve too closely for such effects of selection to really distinguish between them : they are essentially driven by the flow of the ODE :  $\partial_t x_t := -s x_t (1 - x_t)$ . Even if the perturbations are amplified by this selection towards the opposite direction, strong deviations would be much too costly.

## **2.2.2 Fixation on either side is the most stable case and the type G is favored by group selection**

**Proposition 2.2.6.** *Assume that  $\rho_1 < \rho_0 < \rho_\alpha$ . Then,  $\delta_1$  is the only stable QSD, with convergence rate  $\rho_0 - \rho_1$  :*

$$\forall n \geq 1, \forall \xi > 0, \exists C_{n,\xi} > 0, \quad \forall \mu \in \mathcal{M}_{n,\xi}, \quad \|\mu A_t - \delta_1\|_{TV} \leq C_{n,\xi} \exp[-(\rho_0 - \rho_1) t].$$

We also have an additional level of convergence :

**Proposition 2.2.7.** *Assume that  $\rho_0 < \rho_\alpha$ . Then, there exists  $C > 0$  s.t. :*

$$\forall \mu \in \mathcal{M}_1([0, 1]) \setminus \{\delta_1\}, \quad \|\mu A_t^1 - \delta_0\|_{TV} \leq C \exp[-(\rho_\alpha - \rho_0) t].$$

Both results are asymptotic and might not reflect exactly the dynamics on a short time-scale. Yet, their justification gives us some insight on what can happen.

***When the polymorphism gets quickly negligible :***

If  $\sigma$  is large, except for initial conditions very close to 0, a non-negligible proportion of pure type G groups quickly emerges and dominates the distribution. For initial conditions very close to 0, the emergence time of these pure G groups is nearly given through the proportion of groups able to quickly escape such vicinity of 0. The distribution of these survivor groups then rapidly stabilizes to  $\alpha$ , with the first pure G groups also rapidly fixed. The growth of these groups depends soon rather on their own growth rate, so that the role of the still polymorphic groups gets negligible. Finally one observes the competition between the two types of pure groups, with the at first low-frequency G outnumbering the first dominant I.

***When the trajectories are drifted with little fluctuations :***

As explained in the previous subsection, the flow of the equation  $\partial_t x_t := -s x_t (1 - x_t)$  dominates the dynamics as long as  $\mu_t$  stays localized. If  $\sigma$  is rather small, the profile  $\mu_t$  is essentially given by the integration of the growth rate along the trajectories of the flow. Notably, the case where the initial condition has a light tail towards 1 is simpler to get. There is a time that mainly depends on this tail after which the distribution is mainly absorbed at 0. If the proportion of pure G groups is still negligible at this time, we shall observe a decay of the absorbed distribution larger than an exponential rate of  $\rho_\alpha - \rho_0$ . It is a priori unclear that the rate  $\rho_\alpha - \rho_0$  is actually observed, since the QSD  $\alpha$  might be localized around 1 and very difficult to reach for initial conditions rather far away. Yet, we expect some stabilization to occur, where  $\mu_t$  restricted to  $(0, 1)$  gets close to some  $\alpha'$ . This distribution  $\alpha'$  is presumably supported mainly close to 0, with a much larger extinction rate  $\rho'_\alpha$  than  $\rho_0$ . One shall have the right intuition by considering  $\alpha'$  (resp.  $\rho'_\alpha$ ) instead of  $\alpha$  (resp.  $\rho_\alpha$ ) in the reasoning of Propositions 2.2.6 and 2.2.7. Our simulations seem to comfort us in this idea.

As stated in Proposition 2.2.6, the pure G groups shall prevail even in that case. A rate  $\rho_1 - \rho_0$  of decay of the remaining groups is very likely to be seen, but possibly after a domination by type I groups on a much longer time-scale. This time-scale depends actually much on the initial distribution, and particularly on its tail towards 1. Indeed, the transitions to 1 is especially costly in term of its probability of occurrence.

***Compensate the flow of invasion by I individuals :***

By the way, the next section provides a precise description of the strength of this group selection needed to compensate the flow in the limit of vanishing  $\sigma$ . In this large deviation regime, the

process would seem to evolve for most of the time according to a modification of the initial flow. Yet, it is not as simple : one might observe for instance the abrupt emergence of a type negligible until then, whose growth rate is much higher than the previously dominant type.

This is exactly what shall presumably happen in this model in the case  $r_1 < r_0$ . Undoubtedly, the initial proportion of  $G$  groups or the neutral fixation for initially almost pure  $G$  groups concerns a very tiny proportion of ancestors. Yet, the ancestors initially drifted towards larger proportion of  $I$  individuals soon loose any  $G$  individual (with very few exceptions). So there is a point in time at which we cannot neglect anymore the more prolific descendants the former exceptional ancestor groups have. This shall certainly correspond to the time at which  $G$  individuals eventually dominate. Are pure groups the ones that finally dominate ? Due to this potential long-term persistence of the process in the vicinity of 1 when the neutral fluctuations are very small, the behavior of  $\rho_\alpha$  as  $\sigma$  tends to 0 is quite unclear. So it might happen that some almost pure  $G$  groups actually dominate.

### ***Selection upon the initial condition :***

Still, the selection between the initial conditions may be effective if they are sufficiently apart. It may postpone for some significant time the trend towards 0. Yet, only a very specific behavior of the initial condition law could make the polymorphism persist for long. This has been precised in [18] in the limit where the neutral fluctuations are neglected. Such long-term persistence is effectively possible because the flow is vanishing in the vicinity of 1. The authors consider only function  $r$  that are linear with the proportion of type  $G$  individuals. But it should generalize to a much more general case, provided  $r$  is larger around 1 than around 0.

Note however that in such regime, the groups dominating at a given time are essentially not the ancestors of those dominating at a much larger time. The descendants of the former have namely be transported by the flow towards 0, where this sub-population has decayed faster. On the other hand, the ancestors of the latter must have stayed for long in a very tiny and precise region very close to 1, and are thus very few (while the former dominate). Indeed, going backward through the ancestry lines means essentially following the flow backwards. This backward flow goes quickly towards 1, which it approaches at exponential speed. With initial conditions irregular in the vicinity of 1, we thus might observe a surprising sequence of vanishing and reemerging polymorphism between periods of domination by  $I$  individuals.

### ***Is this observable in the individual-based model ?***

Any transition involving a reasonable amount of groups is expected to be indeed observed. This can be estimated through the number of ancestors from the initial population on which such transitions shall rely. The fluctuations shall vanish with the number of them, so that the approximation should be qualitatively good with around 20 ancestries. It means that one expects something very specific, and rather unclear, concerning exceptional behavior when the group neutral birth and death rates are non-negligible anymore.

The main issue is that exceptional transitions are very unlikely to be observed. The escape from a too close vicinity of 0 happens for instance with too small a probability, with a threshold depending on the population size. The most likely is to observe the complete fixation. For the fixation of pure  $G$  groups to be observed, the most probable is then to see one group escaping the

vicinity of 0, reaching the other boundary and generating a sufficient family there for the extinction to become negligible. Only after such exceptional realization becomes the fixation of pure G groups likely to occur. Similarly, when  $\sigma$  is so small that transitions towards 1 become negligible, and for an initial condition with a light tail (towards 1), the fixation of pure G groups happens after a very exceptional behavior.

This mathematical complexity is presumably not so relevant in terms of the biology. As soon as the initial condition has a sufficiently slight tail in this vicinity of 1, one mainly observe a massive proportion of the groups fixing as pure  $I$  types and becoming dominant for a very long time.

### 2.2.3 Polymorphic groups are more stable than type I groups but less than pure type G groups

This case is also treated in [24]. The result is a combination of the ones in the two previous subsections. The dynamics for the domination by pure  $G$  groups relies on similar principles as in Proposition 2.2.6. The main difference is that the intermediate convergence is stated towards a polymorphic QSD rather than pure  $I$  groups. This polymorphic QSD  $\alpha_1$  is described as in Proposition 2.2.2 when one subtracts pure G groups before the renormalization. The asymptotic rate of convergence towards the Dirac at pure  $G$  groups is deduced from this intermediate convergence result :  $\rho_1 - \rho_\alpha$  (smaller than  $\rho_1 - \rho_0$ ).

Again, in the comparison to IBM, one may be faced to the same limitations regarding the origin of the firsts pure  $G$  groups as for Proposition 2.2.6. In practice, the convergence towards the polymorphic QSD might also not actually reflect the main dynamics of convergence. It might happen that the "emergence" of pure type 1 groups can actually be almost concomitant to the emergence of  $\alpha^1$ . Looking at simulations for low values of  $\sigma$ , an alternative metastable regime around 0 may visibly dominate for a significant time the marginal law restricted to  $(0, 1)$ . Comparing with the case of initial condition close to 1, this distribution  $\alpha'$  is very different of the actual QSD  $\alpha$ , with very separate supports. Even if  $\rho_\alpha < \rho_0$ , it would not be surprising that for the alternative distribution  $\rho'_\alpha > \rho_0$ . For a large range of initial condition, even if  $\rho_\alpha < \rho_0$ , the initially observed dynamics is rather the one described in Propositions 2.2.6 and 2.2.7 by the interplay between 1, 0 and  $\alpha'$  where  $\rho_1 < \rho_0 < \rho'_\alpha$ . The first step of the dynamics is thus manifestly a convergence to 0 with a rate expected to stabilize towards  $\rho'_\alpha - \rho_0$ .

### 2.2.4 Validation by some simulations ?

In order to evaluate the exceptionality of transitions towards 1, one can propose the following simulation experiment, which is a work in progress. Let us consider some parameters for which most of the marginals tend towards 0, with  $\sigma$  sufficiently low. To observe the upheaval of the distribution by type 1 individuals, the marginals should be encoded rather not by the masses at the different grid points, but by the logarithms of these quantities. The case of the absorbing states is treated separately from the marginal restricted to  $(0, 1)$ . To be precise in the vicinities of 0 and 1, it may be useful to refine the grid there.

We shall succeed in observing the mass towards 1 increasing up to the point of exceeding the mass towards 0. Yet, it is then unclear at which concentration the regeneration of these cooperative groups exceeds the effect of having more and more exceptional transitions leading there. Since the quickest transitions are expected to bring the less mass towards 0, our idea is to truncate densities

to prevent the most exceptional transitions. So at each simulation step, we suppress from the marginal the mass on states that contain less than the threshold. By varying the threshold, we would have a better view on the number of groups required to observe the transition from 0 to 1 in individual-based models.

If the dynamics is almost unchanged after the truncation, we could conclude that the cost of the transitions that mainly contribute to this upheaval is lower than the threshold. If the upheaval arises later on, it would mean that less costly transitions could have been sufficient to make type  $G$  emerge. Yet, their contribution becomes negligible when compared to quicker transitions. Finally, if the marginal becomes supported on some interval that does not approach 1, it means that any transition towards 1 would be at least as costly as the threshold.

### 2.3 Conclusion of the section

Generally, the interplay between different types happens in the time-scale at which they differentiate their carriers. Yet, we have seen in this example that the trade-off between different kinds of advantages can be particularly tricky. The a priori neutral genetic drift might happen to be strongly coupled to the efficiency of some components of selection. We clearly do not claim however that such effect is prevalent, because the assumption of a strong heritability of the local dynamics is not so common. Also, the selection at the group level might rely on possibly exceptional dynamics. Although this exceptionality could be compensated by a larger asymptotic growth rate, two aspects should be remembered : first, it might be much too unlikely for actual populations, so that the mathematical model could be misleading; second, it will take a possibly very long time. It is thus questionable for real life that no other event happens to disturb any of the sub-populations before the emergence of cooperative groups. Finally, once any type has been dominant, it may be much more difficult than neutral for the other type to invade : for the invasion of cooperatives, the genetic drift inside the groups must lead the process out from the very stable cheater quasi-equilibrium; while, for the invasion of cheaters, the genetic drift between groups must disrupt the also very stable cooperative quasi-equilibrium. Such large deviations are known to be generally especially costly.

## 3 Large deviation estimate for the adaptation by mutations of weak effect

In this section, the focus is on the trajectories of evolution, with the selection of favorable mutations. Interestingly, the authors of [9] show a mathematical similarity between two limiting behaviors : a first one where the trait variations are very small; and the second where the mutation rate is very small (as compared to the selective effects). This first case can notably be associated to a limit where mutations of very small effects accumulate. For the results we present above to hold, the main requirement is that natural selection is very strong in the timescale where one observes the trait dispersion under neutrality.

Again, the population size is assumed to be sufficiently large to include the whole range of the stochastic variations of the trait. The resulting purely deterministic model shall provide a good approximation to the dynamics of trait proportions among this population. This relation to the trait proportions can be retrieved from the convergence result established by the time-scale



separation.

### 3.1 The continuous-space limiting behavior

One considers the family  $(u^\epsilon)_\epsilon$  of deterministic solutions to respectively :

$$\begin{aligned} \partial_t u^\epsilon(t, x) &= \frac{\epsilon}{2} \Delta u^\epsilon(t, x) + \frac{R(x, \psi_t^\epsilon)}{\epsilon} u^\epsilon(t, x), \quad x \in \mathbb{R}^d, t \geq 0, \\ -\epsilon \log(u^\epsilon(0, x)) &= h_\epsilon(x) \end{aligned} \quad (11)$$

where the competition effect  $\psi_t^\epsilon = (\psi_t^{i,\epsilon})_{i \leq r}$  is defined, for  $r$  resources, with a competition kernel  $\Psi_i : \mathbb{R}^d \rightarrow \mathbb{R}_+$  for resource  $i$  by :

$$\psi_t^{i,\epsilon} := \int_{\mathbb{R}^d} \Psi^i(x) u^\epsilon(t, x) dx. \quad (12)$$

Considering mutation effects through some heat kernel corresponds to the case where lots of mutations with very small effects occur and are dispersed throughout the population. In asexual populations, we rather expect selective mutations to invade and fix one after the other. Such a model is thus rather justified for sexual populations, where many recombination events occur and many different alleles with small selective effects may coexist. It is known for instance that many of the Human traits are the consequence of many different loci in the genome, with lots of variability in the loci for a given phenotype. The heat kernel seems then quite relevant to characterize the variability of response for such a trait. It would be of interest to prove that the limiting behavior for  $u^\epsilon$  can also be retrieved starting from these individual-based models approximated by  $u^\epsilon$ .

The following assumptions are required in [9] in order to justify a limit to  $\varphi_\epsilon$  through a variational representation.

1. *Assumptions on  $\Psi_i$*  : For any  $1 \leq i \leq r$ ,  $\Psi_i \in W^{2,\infty}(\mathbb{R}^d)$ .

Moreover, there exist  $0 < \Psi_{\min} < \Psi_{\max}$  such that  $\forall 1 \leq i \leq r, \forall x \in \mathbb{R}^d, \Psi_{\min} \leq \Psi_i(x) \leq \Psi_{\max}$ .

2. *Assumptions on  $R$*  : (a)  $R$  is continuous on  $\mathbb{R}^d \times \mathbb{R}^r$ .

(b) There exists  $A > 0$  such that

$$\forall 1 \leq i \leq r, \forall x \in \mathbb{R}^d, \forall (v_\ell) \in \mathbb{R}^r, \quad -A \leq \partial_{v_i} R(x, (v_\ell)_{\ell \leq r}) \leq -A^{-1}$$

(c) There exist two positive constants  $0 < v_{\min} < v_{\max}$  such that  $\min_{x \in \mathbb{R}^d} R(x, v) > 0$  as soon as  $\|v\|_1 < v_{\min}$ ,

and  $\max_{x \in \mathbb{R}^d} R(x, v) < 0$  as soon as  $\|v\|_1 > v_{\max}$ , where  $\|v\|_1 = \sum_{\ell \leq r} |v_\ell|$ .

(d) Let  $H$  denotes the annulus  $B(x, 2v_{\max}) \setminus B(x, v_{\min}/2)$  (for the  $\|\cdot\|_1$  norm). Then  $\sup_{v \in H} \|R(\cdot, v)\|_{W^{2,\infty}} < \infty$ .

3. *Assumptions on  $h_\epsilon$*  : (a)  $h_\epsilon$  is Lipschitz-continuous on  $\mathbb{R}^d$ , uniformly with respect to  $\epsilon > 0$ .

(b)  $h_\epsilon$  converges in  $L^\infty(\mathbb{R}^d)$  as  $\epsilon$  tends to 0 to a function  $h$ .

(c) For all  $\epsilon > 0$  and all  $1 \leq i \leq d$ ,  $v_{\min} \leq \int_{\mathbb{R}^d} \Psi_i(x) \exp(-h_\epsilon(x)/\epsilon) dx \leq v_{\max}$ .

In particular,  $u_\epsilon(0, x)$  is bounded in  $L^1(\mathbb{R}^d)$ .

**Remark :** Such competition may at first appear as a global competition where every trait is competing against each other for the same resources. The first assumption done in [9] requires indeed for the  $\Psi_i$  to be lower-bounded by a strictly positive constant. However, in a limit where  $r$  is large and  $\Psi_i$  is very concentrated, the model may become a good approximation of a local competition (in the trait space). Thus, one may hope to extend the model naturally to such a framework.

### 3.2 Results for the continuous case

We gather here the main results presented in [9], where we have changed some notations for clarity, notably  $\psi_t$  and  $\varphi$  :

**Theorem 3.1.** (Feynman-Kac representation of the solution of (11))

Let  $u^\varepsilon$  be the unique weak solution of (11), then, under the assumptions given in [9] :

$$\forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d, \quad u^\varepsilon(t, x) = \mathbb{E}_x \left[ \exp \left( \frac{-h_\varepsilon(X_t^\varepsilon)}{\varepsilon} + \frac{1}{\varepsilon} \int_0^t R(X_s^\varepsilon, \psi_{t-s}^\varepsilon) ds \right) \right], \quad (13)$$

where for all  $x \in \mathbb{R}^d$ ,  $\mathbb{E}_x$  is the expectation associated to the probability measure  $\mathbb{P}_x$ , under which  $X_0^\varepsilon = x$  almost surely and the process  $B_t = (X_t^\varepsilon - x)/\sqrt{\varepsilon}$  is a standard Brownian motion in  $\mathbb{R}^d$ .

**Remark :** This representation of the solution through a penalization of a stochastic process is in practice of the same kind as the one we have presented for the empirical law  $\mu_t$  in Section 2. There is only a little difference in the fact that  $X_t$  describes now the process when we look at the lineage backward in time. A similar convergence result should thus justify  $u^\varepsilon(t, x) dx$  as the limit of some individual-based measure-valued processes  $(\nu_t^{\varepsilon, K})$ . In particular,  $(1/\varepsilon) \cdot R(x, \psi_t^{\varepsilon, K})$  should be the additional growth rate of individuals with trait  $x$ , where for  $i \leq r$ ,  $\psi_t^{i, \varepsilon, K} := \langle \nu_t^{\varepsilon, K} | \Psi^i \rangle$ . With fixed  $\varepsilon$ , this is a priori a specific case of the McKean-Vlasov equations mentioned in the conclusion of section 2, where the law of the process itself acts on the individual dynamics (again, we refer to [23]).

**Lemma 3.2.1.** The function  $I_t^{R, \varepsilon} : C([0, t]) \rightarrow \mathbb{R}$  defined by  $I_t^{R, \varepsilon}(y) = \int_0^t R(y_s, \psi_s^\varepsilon) ds$  is Lipschitz continuous on  $C([0, t])$  endowed with the  $L^\infty$ -norm. The Lipschitz constant is uniform with respect to  $\varepsilon$  for  $\varepsilon$  small enough. Moreover, there exists a kernel  $\mathcal{M}$  on  $\mathbb{R}_+ \times \mathcal{B}(\mathbb{R}^k)$  such that, along a subsequence  $(\varepsilon_k)_{k \geq 1}$  converging to 0 :

$$\forall y \in C([0, t]) \quad I_t^R(y) := \lim_{k \rightarrow \infty} I_t^{R, \varepsilon_k}(y) = \int_0^t \int_{\mathbb{R}^k} R(y_s, \psi) \mathcal{M}_s(d\psi) ds.$$

We recall that a kernel  $\mathcal{M}$  is a function from  $\mathbb{R}_+ \times \mathcal{B}(\mathbb{R}^d)$  into  $\mathbb{R}_+$  such that, for all  $t \in \mathbb{R}_+$ ,  $\mathcal{M}_s$  is a measure on  $\mathcal{B}(\mathbb{R}^d)$  and, for all  $A \in \mathcal{B}(\mathbb{R}^d)$ , the function  $s \rightarrow \mathcal{M}_s(A)$  is measurable.

**Remarks :** • Since the level of competition actually depends on the global composition in traits of the population, and that some information on the density is lost in the limit  $\varepsilon \rightarrow 0$ , the implicit dependency of  $v$  on  $\varphi$  is much less clear in the limiting model. Therefore, the uniqueness of solutions to such equations is quite demanding.

• Given a lineage described by  $y$ ,  $I^R(y)$  encodes an averaged effect of selection due to the perceived growth rate. As one can infer from (13) when  $\varepsilon$  is very small, the density around such path

is approximately amplified by the exponential of this quantity divided by  $\epsilon$ . Such large amplification is the core of the Large Deviation theory, where the stochastic behavior is concentrated close to specific paths.

**Theorem 3.2.** *For all  $(t, x)$  in  $\mathbb{R}_+ \times \mathbb{R}^d$ ,*

$$\varphi(t, x) := \lim_{k \rightarrow \infty} \epsilon_k \log u^{\epsilon_k}(t, x) = \sup_{y \in \mathcal{G}_{t,x}} \{-h(y_0) + I_t^R(y) - I_t^L(y)\}$$

where the convergence holds uniformly on compact sets and the limit  $\varphi(t, x)$  is Lipschitz w.r.t.  $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$  while  $I_t^L(y) = \begin{cases} \int_0^t \|y'(s)\|^2 ds & \text{if } y \text{ is absolutely continuous,} \\ +\infty & \text{otherwise.} \end{cases}$

$\mathcal{G}_{t,x}$  denotes the set of continuous functions from  $[0, t]$  to  $\mathbb{R}^d$  such that  $y_t = x$ , and  $I_t^R$  and  $(\epsilon_k)_{k \geq 1}$  are associated as in Lemma 3.2.1.

Note that in this optimization,  $y$  defines the main ancestral line of the individuals dominant with trait  $x$  at time  $t$ . The Large Deviation theory namely ensures that as  $\epsilon \rightarrow 0$ , their ancestral lines are very concentrated on such specific and deterministic histories.

Finally, also in [9] the authors are indeed able to relate this variational limit to the solution of the Hamilton-Jacobi equation :

**Theorem 3.3.** *Under many assumptions on the behavior of  $R$  and  $h_\epsilon$ , one can ensure that  $\psi^{\epsilon_k}$  converges in  $L_{loc}^1(\mathbb{R}_+)$  along the subsequence  $\epsilon_k$  of Lemma 3.2.1 to a nondecreasing limit  $\bar{\psi}$ , the kernel  $\mathcal{M}$  satisfies*

$$\forall s \geq 0, \quad \mathcal{M}_s(d\psi) = \delta_{\bar{\psi}_s}(d\psi),$$

and the limit  $\varphi$  of Theorem 3.2 solves in the viscosity sense :

$$\begin{cases} \partial_t \varphi(t, x) = R(x, \bar{\psi}_t) + \frac{1}{2} |\nabla \varphi(t, x)|, & \forall t \geq 0, x \in \mathbb{R}^d, \\ \max_{x \in \mathbb{R}^d} \varphi(t, x) = 0, & \forall t \geq 0. \end{cases}$$

### **Behavior of the solution to Hamilton-Jacobi equation**

From Theorem 3.2, one deduces the following approximation for small  $\epsilon$  :  $u^\epsilon(t, x) \approx C(t, x) \exp[\varphi(t, x)/\epsilon]$ . We thus expect to observe a concentration of the individual traits on typically a unique value or a few well-separated values. In the limiting model, these values may be driven by a continuous displacement in the direction of natural selection. They may also split, which is especially interesting for a model of speciation. It is in fact the original purpose of such Hamilton-Jacobi analysis in [13] to extend the model of Adaptive Dynamics (cf. Subsection 4.5). We could also imagine fusions, for the recovery of a single population from two populations spatially separated for a long time. Finally, there could also be some jumps, leading to a complete renewal of the eco-system. A wide variety of events are thus observable with this simplified model.

### **Discussion on the involved time-scales :**

At a given  $\epsilon > 0$ , looking at equation (11), we can infer from the growth rate that this ecological and competition fluctuations occurs in a time-scale of order  $\epsilon$ , as compared to the time-scale at

which the traits evolve. The variance of the trait evolves on the contrary at a much larger time scale of order  $\epsilon^{-1}$ , like in the case of neutral traits. Of course, the precise description of the demographics as presented in [16] should occur in the short time-scale of order  $\epsilon$ . In such a large population, the life-expectancy of any particular individual is on an even shorter time-scale, depending on the ratio between the birth and the death rates. The closer it is to one, the shorter this time-scale. A separation of time-scale with the evolutionary trajectory can still be justified for the competition between different effects of selection. But as we have seen in Section 2, this competition might last for very long times.

In practice, the description given by equation (11) is not so easily related to macroscopic observation of the response to selection, because the neutral variability is difficult to scale (the factor  $\epsilon$  before the Laplacian). We can reasonably assume that we know the density  $u^\epsilon(t, x) = u^\epsilon(t, x)$  (as long as it is not too negligible) and the fitness effect  $R_t^\epsilon(x) = (1/\epsilon) \cdot R(x, \psi_t)$ . For at least positive values, it can be inferred in some laboratory experiment by artificially introducing individuals with trait  $x$  in a much larger sample of the population at time  $t$  and see how they grow. One could then estimate the value  $\epsilon$  for which  $\varphi^\epsilon(t, x) = \epsilon \log u^\epsilon(t, x)$  seems reasonably close to the solution  $\varphi$  of the Hamilton-Jacobi equation :

$$\partial_t \varphi(t, x) = \epsilon R_t^\epsilon(x) + \frac{1}{2} |\nabla \varphi(t, x)|^2, \quad (14)$$

or the optimization problem. Note that a Gaussian distribution for  $u^\epsilon$  corresponds to  $\varphi(t, x) \sim -(x - x_t)^2 / \sigma_t^2$ , that is the Second order Taylor expansion around  $x_t$ , with  $x_t$  the optimal trait and  $\epsilon \sigma_t^2$  the variance of the distribution. Moreover, if  $\varphi$  is a solution to equation (14), then for  $\lambda > 0$ ,  $\hat{\varphi}(x, t) := \varphi(\lambda t, \lambda x)$  is a solution to (14) with  $\epsilon$  replaced by  $\lambda \epsilon$ . Since we expect the dependency in  $\sigma_0$  on  $\sigma_t$  to vanish very quickly, the identification of  $\epsilon$  from  $\sigma_t$  would not be a too difficult option. Yet, as mentioned in the next paragraph, the speed of response to selection might be driven by an exceptional proportion of the population at time  $t$ . An estimate such as  $\sigma_t$  that summarizes rather the core of the distribution  $u^\epsilon(t, x) dx$  might thus not be so relevant.

### ***Connection with individual-based models***

To justify the connection with individual-based models, the concentration effects may appear favorable. Yet, as we can see from Theorem 3.2, the adaptation of the population is driven by exceptional profiles of stochastic variations. The stochastic techniques of Large Deviations is then crucial to obtain the limiting behavior. It means notably that the selection is mainly driven by individuals at the front. This optimization provides a very interesting insight on the history of the genealogies that led to the traits observed generally in the population at time  $t$ . We can notably observe cases where the traits at time  $s \leq t$  for the ancestors of the dominants at time  $t$  may not be typical at all as compared to the dominant traits at time  $s$ . This of course raises the issue of biological relevance of such transitions. We face similar limitations as in previous Section 2 for the relevance of events involving very low population headcount. Still, it is much easier to relate the above description to some individual-based model than from the more classical description only relying on Hamilton-Jacobi equations (cf. the conclusion of the section).

### 3.3 Results for the discrete case

This analysis of Feynman-Kac operators and Large Deviation principles can be adapted to a discrete space. Simplifying assumptions are then more easily obtained and makes it possible to have a clearer view on the implications of this model.

In [9] is introduced the following system of ordinary differential equations, modeling the mass dynamics on a finite set  $E$  :

$$\begin{cases} \partial_t u^\varepsilon(t, k) = \sum_{j \in E \setminus \{k\}} \exp\left(\frac{-\mathfrak{I}(k, j)}{\varepsilon}\right) (u^\varepsilon(t, j) - u^\varepsilon(t, k)) + \frac{1}{\varepsilon} u^\varepsilon(t, k) R(k, \psi_t^\varepsilon) , & \forall t \in [0, T] , \forall k \in E \\ u^\varepsilon(0, k) = \exp\left(-\frac{h^{\varepsilon j}(k)}{\varepsilon}\right) , & \psi_t^{i, \varepsilon} = \sum_{k \in E} \Psi^i(k) u^\varepsilon(t, k), \quad \forall i \leq r \end{cases}$$

First, the solution  $u^\varepsilon$  of the system is described in [9] by using an integral representation similar to (13). Let  $(X_s^\varepsilon, s \in [0, T])$  be the Markov process in  $E$  with infinitesimal generator

$$L^\varepsilon f(i) = \sum_{j \in E} (f(j) - f(i)) \exp\left[\frac{-\mathfrak{I}(i, j)}{\varepsilon}\right]$$

i.e. the Markov process which jumps from state  $i \in E$  to  $j \neq i$  with rate  $\exp(-\mathfrak{I}(i, j)/\varepsilon)$  .

**Proposition 3.3.1.** (*Integral representation*)

For any positive real number  $t$  and any element  $i$  of  $E$ , we have

$$u^\varepsilon(t, i) = \mathbb{E}_i \left[ \exp \left( \frac{-h^\varepsilon(X_t^\varepsilon)}{\varepsilon} + \frac{1}{\varepsilon} \int_0^t R(X_s^\varepsilon, \psi_{t-s}^\varepsilon) ds \right) \right]$$

The interpretation is very similar to the one of Theorem 3.1.

**Proposition 3.3.2.** (*Weak LDP*)  $(X^\varepsilon)_{\varepsilon \geq 0}$  satisfies a weak LDP with rate function

$$I_T : \begin{cases} \text{D}([0, T], E) \rightarrow \mathbb{R} \\ y \mapsto \sum_{\ell=1}^{N_y} \mathfrak{I}(y_{t_\ell^y-}, y_{t_\ell^y}) , \end{cases}$$

where  $\text{D}([0, T], E)$  is the space of càdlàg functions from  $[0, T]$  to  $E$ ,  $N_y$  is the number of jumps of  $y$  and  $(t_\ell^y)_{1 \leq \ell \leq N_y}$  the increasing sequence of jump times of  $y$ .

We shall also make use of the notation  $I_T(y) := \sum_{0 < s \leq T} \mathfrak{I}(y_{s-}, y_s)$  with the implicit convention that  $\mathfrak{I}(i, i) = 0$  for all  $i \in E$ .

**Theorem 3.4.** Lemma 3.2.1 is also satisfied in the discrete case and for all  $(t, i)$  in  $(0, +\infty) \times E$ , with the associated subsequence  $(\varepsilon_k)$  :

$$\begin{aligned} \varphi(t, i) &:= \lim_{k \rightarrow \infty} \varepsilon_k \log u^{\varepsilon_k}(t, i) \\ &= \sup_{y \in \text{D}([0, t], E) \text{ s.t. } y_0 = i} \left\{ -h(y_t) + \int_0^t \int_{\mathbb{R}^r} R(y_s, \psi) \mathcal{M}_{t-s}(d\psi) ds - \sum_{0 < s \leq t} \mathfrak{I}(y_{s-}, y_s) \right\} . \end{aligned}$$

Like in the expression in Theorem 3.2, one shall remark that the optimal  $y$  in this expression describes the transition of states for the lineage of a typical individual at time  $t$ . This evolution is backward in time, as can be also seen in the term  $R(y_s, \psi) \mathcal{M}_{t-s}(d\psi)$ , where  $\mathcal{M}_{t-s}(d\psi)$  provides the law of  $\psi_{t-s}$ .

**Theorem 3.5.** *For all subsequence  $(\varepsilon_k)_{k \geq 1}$  as in Theorem 3.4, the limit  $\varphi(t, i)$  of  $\varepsilon \log u^\varepsilon(t, i)$  is Lipschitz with respect to the time variable  $t$  on  $(0, +\infty)$ . In addition, if  $h(i) \leq h(j) + \mathfrak{F}(i, j)$  for all  $i \neq j$ , the function  $\varphi$  is Lipschitz on  $\mathbb{R}_+$ .*

We see that all the previous results in continuous space have an equivalent for discrete space. By the way, the continuous problem is very probably well-approximated by sufficiently fine discretization of the state space.

In the following, we focus on a specific case where the uniqueness can be obtained and the whole dynamics precised with much more precision. The result presented in [9] relies on some assumption on the stability of the dynamics restricted to any subset of  $E$ , which is called Assumption (H). We present next the main intuition behind this assumption and refer to [9] for the precise definition.

By this Assumption (H), the authors notably ensure that, for any subset  $A$  of  $k$  different types involved, there exists a unique strongly attractive equilibrium. Since the focus is on the dynamical system, the estimation of the dynamics is valid as soon as the  $k$  types are initially in non-negligible proportion. Yet, some of the components of the steady-state may be 0, meaning an exponential decay of the mass on those. Of course, by restricting the dynamics on some subset  $B$  of  $A$  while keeping the non-vanishing components of such steady state for  $A$ , one shall retrieve as steady state for  $B$  the restriction on  $B$  of the steady state for  $A$ . Assumption (H) also ensures that any other (unstable) equilibrium is quickly escaped. The dominant traits at time  $t$  (for which  $\varphi(t, \cdot) = 0$ ) are then proved to stay piecewise constant, while the emergence of favorable competitors is easily computed. Once such competitor has emerged, it disrupts the equilibrium of traits, and assumption (H) enables us to predict the issue of the equilibrium that follows.

Given any set  $A$  of present types, the equilibrium is given by  $(u_{A,j}^*)_{j \in A}$ . Thus, the competition exerted on the  $i$ -th resource by this eco-system is  $\Psi^i(A) := \sum_{j=1}^r \Psi^i(j) \cdot u_{A,j}^*$ .

**Proposition 3.3.3.** *Assume that (H) is satisfied. Let  $(\varepsilon_k)_{k \geq 1}$  be as in Theorem 3.4. For any  $t \geq 0$ , there exists  $\rho_t > 0$  such that, for all  $s \in (t, t + \rho_t]$ ,  $\psi_s^\varepsilon$  converges to  $\psi_s = \psi_t := \Psi(\{\varphi(t, \cdot) = 0\})$ , where the convergence is uniform in all compact subsets of  $(t, t + \rho_t]$  and where we define for any  $A \subset E$ :  $F(A) = (\sum_{j=1}^r \eta_i(j) u_{A,j}^*)_{1 \leq i \leq r}$ .*

*In particular, the weak limit  $\mathcal{M}_s$  of  $\delta_{\psi_s^\varepsilon}$  obtained in Lemma 3.2.1 satisfies  $\mathcal{M}_s = \delta_{\psi_t}$ , for almost all  $s \in (t, t + \rho_t)$  and the function  $t \mapsto \psi_t$  is right-continuous.*

Like in the previous case, we shall observe a concentration effect. It means here that there are brutal transitions between one demographic equilibrium and the next one generated by the first invasion by a mutant sub-population.

**Proposition 3.3.4.** *Assume that (H) is satisfied. Any limit  $\varphi$  of  $\varepsilon_k \log u^{\varepsilon_k}$  along a subsequence as in Theorem 3.4 satisfies for all  $i \in E$  and for all  $t \geq 0$ :  $\varphi(0, i) = -h(i)$*

$$\varphi(t, i) = \sup_{y(0)=i} \{-h(y(t)) + \int_0^t R(y(u), \psi_{t-u}) du - I_t(y)\},$$

and its dynamic programming version

$$\varphi(t, i) = \sup_{y(s)=i} \{\varphi(s, y(t)) + \int_0^t R(y(u), \psi_{t+s-u}) du - I_{s,t}(y)\}. \quad (15)$$

In addition, the problem (15) admits a unique solution s.t.  $t \mapsto \psi_t := F(\{\varphi(t, \cdot) = 0\})$  is right-continuous. In particular, the full sequence  $(\varepsilon \log u^\varepsilon)_{\varepsilon > 0}$  converges to this unique solution when  $\varepsilon \rightarrow 0$ .

**Theorem 3.6.** *Under Hypothesis (H) and assuming that  $h(i) \leq h(j) + \mathfrak{T}(i, j)$  for all  $i \neq j$ , the problem*

$$\begin{cases} \partial_t \varphi(t, i) = \sup\{R(j, \psi_t) \mid j \in E \text{ s.t. } \varphi(t, j) - \mathfrak{T}(j, i) = \varphi(t, i)\}, \\ \forall i \in E, \quad \varphi(0, i) = h(i) \end{cases} \quad (\text{with the convention } \mathfrak{T}(i, i) = 0)$$

admits a unique solution such that  $t \mapsto \psi_t = F(\{\varphi(t, \cdot) = 0\})$  is right-continuous and it is the unique solution to the variational problem (15).

The system follows a succession of equilibria (with possibly different types that equilibrate), with the population headcount of mal-adapted types vanishing linearly in their logarithm, while this headcount logarithm increases for the adapted ones. It goes on until one adapted type reaches the threshold for a non-negligible frequency. Then occurs a kind of "catastrophe", where the whole equilibrium is immediately renewed (in the time-scale of evolution). Given the similarity between the description in this discrete case and the continuous case presented in Subsection 3.2, one can infer that similar "catastrophes" might happen even in the continuous case. This is less expected, since we see meanwhile a dynamics for the dominant trait, but cannot be excluded a priori.

Without this assumption of a unique stable equilibrium, it would not be clear what happens to the population at such "catastrophe" events. The only information provided by  $\varphi(\cdot, t)$  is namely the knowledge of the non-negligible types, for which  $\varphi(x, t) = 0$ , with a priori no means to infer otherwise the headcounts at equilibrium. Notably, different issues or an unstable behavior could lead to different competition effects, completely changing the following dynamics.

The main restriction for this model is that the tails of the distribution are not too much involved. Indeed, when there is some barrier, where the growth rate is very low, the trajectory that gets selected upon through the optimization procedure in Theorem 3.2 might not be realistic. It might well be deduced from this optimization the ancestors of the population at time  $t$  shall represent at time  $s \leq t$  a proportion less than  $10^{-9}$  of the entire population at time  $s$ . Of course, it would only be reasonable for populations with a size quite larger than  $10^9$  ! And all the more since families with low population size have a high risk of becoming extinct. Considering the actual population size considered, such optimization procedure could be an efficient way to truncate too rare such transitions as we have done for the analysis in Section 2. For a threshold  $-\varphi_V$ , we can namely define some  $\tilde{\varphi}(t, x)$  as the supremum over the same quantity as in Theorem 3.2 under the additional condition on  $y \in \mathcal{G}_{t,x}$  that :

$$\forall s \leq t, \quad -h(y_0) + I_s^R(y) - I_s(y) > -V_V, \quad \text{where } I_s^R(y) := \int_0^s \int_{\mathbb{R}^k} R(y_u, y) \mathcal{M}_u(dy) du.$$

Naturally, we define  $\tilde{\varphi}(t, x) = -\infty$  if no function  $y \in \mathcal{G}_{t,x}$  is able to satisfy this condition. By this, we mean that such transitions rely on too exceptional ancestors to be treated in such a deterministic way. Alternative strategies have been proposed in [22] and [21] with more regularities than the proposed truncation, yet not much more biological justification. To prevent such exceptional

densities, the authors use a singular term for the growth rate, which gets very negative when the density of the state is too small.

To be even more realistic, one should treat such events of crossing barriers as punctual events occurring at a very low rate. In fact, large deviation approaches may well be suited to estimate these rates. One should only remark that the growth rate on the other side of the barrier is not responsible for any increase on the rate at which such transitions occur. It only increases the probability that such exceptional crossing event leads actually to an invasion. And that is why such events are much more exceptional than with the asymptotic model given by  $\varphi$ , where the growth rate immediately and regularly increases the log-density on this other side. The punctual transitions that we are to describe in the next section is thus quite suited to deal also with such rare crossing events.

## 4 Mutations as the limiting factor

This last Section addresses the longest time-scales, where the limiting factor for evolution is the emergence of the mutations. By its emergence, we mean not only that some individuals carry such mutations at time  $t$  but rather that it has been stabilized at a non-negligible frequency in the population. For simplicity, many authors assume that selective traits cannot coexist in the same population, and that the population size is very stable at a value depending on this dominant trait. Namely, the demographic fluctuations that has been crucial in Section 2 are neglected, and a fortiori the individual variability detailed in Section 1 is only seen through an averaged effect. In such a deterministic approximation, the fixation or the vanishing of an invasive trait is generally described, as soon as the proportion of invaders is not anymore negligible, by Lotka-Volterra equations. For elementary models of interactions (notably not frequency-dependent), it is then easy to impose conditions ensuring that the fixation is the only stable equilibrium of the dynamical system.

Given the assumption that such ecological transitions are in a much shorter time-scale than the time between occurrences of new mutations (at first in a unique individual), such conditions, called "invasion implies fixation", ensures that the population stays almost always monomorphic in the evolutionary time-scale. This means in particular that the invasion of new mutations can be treated as punctual events that refreshes the equilibrium. In this view, this behavior is similar to the evolution of traits given for the discrete space in Section 3 (cf. the concluding remarks for more details on this link). Yet, these punctual fixation events happens at unpredictable times, corresponding to the occurrence of a successful mutation.

Note that this shall hold true even if coexistence of traits are allowed in the model. Yet, the rate at which it occurs and the issue of the invasion might depend on any variations of the frequencies (like oscillation of frequencies) and of the population size, which makes the analysis way trickier. Of course, one could also generalize these models to include the competition for resources like in Section 3. Yet, the specificity of these more complicated models can only be seen when several traits can coexist. If one wishes to include events of speciation, this is however a very reasonable way to justify it (specialization of two sub-populations).

For strong selection effects, one can also prove negligible even the time between the arrival of the first mutant and its descendants invading the population (at least reaching a non-negligible frequency). Namely, a favorable mutant has a larger expectancy of offspring than individuals of resident type and shall thus generate a family that either dies out quite early or thrive at an



exponential rate. Contrary to the model of discrete trait evolution described in Subsection 3.3, the time-scale for the occurrence of mutations is chosen here in such a way that this growth period is not a limiting factor. By "successful mutation", we mean the occurrence of the mutation on its first carrier followed by a successful invasion of the resident population by its descendants. By assuming that the mutation events are sufficiently separated (in time), we infer that successful mutations happen nearly independently of the time waiting for it (where the monomorphic population stays the same) and of the unsuccessful invasions. This explains why the new event of invasion is given by an exponential law without memory.

Such a process with piecewise constant population equilibria, including a dominant trait, has been originally introduced as the Trait Substitution Sequence in [20] and more formally related to individual-based models in [7]. A large family of extensions has emerged, notably to include coexistence of traits (cf. [10], [1]), or more complicated interactions, for instance with horizontal transfer [5] or ageing [19]. Since they deal with two additional time-scales, we rather focus on [6], where the events of invasion affects the dynamics of some marker, and [2], which demonstrates that adaptive dynamics can indeed be a precise approximation of some individual-based models.

#### 4.1 A model combining a marker dynamics to the evolution of traits

In [6], the authors consider an asexual population driven by births and deaths where each individual is characterized by hereditary types: a phenotypic trait under selection and a neutral marker. This analysis is motivated by research on the prevalence of selection between various species or clades that could be based on the observed variability of neutral markers.

The trait and marker spaces  $\mathcal{X}$  and  $\mathcal{U}$  are assumed to be compact subsets of  $\mathbb{R}$ . The type of individual  $i$  is thus a pair  $(x_i, u_i)$ ,  $x_i \in \mathcal{X}$  being the trait value and  $u_i \in \mathcal{U}$  its neutral marker. The individual-based microscopic model from which we start is a stochastic birth and death process with density-dependence whose demographic parameters are functions of the trait under selection and are independent of the marker. We assume that the population size scales with an integer parameter  $K$  tending to infinity so that individuals are weighted with  $\frac{1}{K}$  to observe a non-trivial limit of the empirical measure. The state of the population at time  $t \geq 0$ , rescaled by  $K$ , is described by the point measure :

$$\nu_t^K = \frac{1}{K} \sum_{i=1}^{N_t^K} \delta_{(x_i, u_i)} = X_t^K(dx) \pi_t^K(x, du), \quad \text{where } X_t^K = \frac{1}{K} \sum_{i=1}^{N_t^K} \delta_{x_i} \text{ and } \pi_t^K(x, du) = \frac{\sum_{i=1}^{N_t^K} \mathbf{1}_{x_i=x} \delta_{u_i}}{\sum_{i=1}^{N_t^K} \mathbf{1}_{x_i=x}}$$

are respectively the trait marginal and the marker distribution for a given trait value  $x$ . Here,  $\delta_{(x,u)}$ ,  $\delta_x$  are respectively the Dirac measure at  $(x, u)$  and  $x$ .

With the following definitions, the authors could ensure that the mutations happen at different time scales for the trait and for the marker, both longer than the individuals lifetime scale. Thus, the limiting behavior results from the interplay of these three time scales: births and deaths, trait mutations and marker mutations. To precise such a separation of time scales for the mutations, the proof relies strongly on the fact that the population size remains around the equilibrium of some dominant trait(s) when a mutation on the trait occurs. Although the authors could include fluctuations of the parameter  $u$ , it is only possible because  $u$  has no effect on the probability that the mutation succeeds to invade the population.

## 4.2 The individual-based model

**Definition 4.2.1.** • An individual with trait  $x$  and marker  $u$  reproduces with birth rate given by  $0 \leq b(x) \leq \bar{b}$ , the function  $b$  being continuous and  $\bar{b}$  a positive real.

• Reproduction produces a single offspring which usually inherits the trait and marker of its ancestor except when a mutation occurs. Mutations on trait and marker occur independently with probabilities  $p_K$  and  $q_K$  respectively. Mutations are rare and the marker mutates much more often than the trait. We assume that

$$q_K = p_K r_K, \text{ with } p_K = \frac{1}{K^2}, q_K \xrightarrow{K \rightarrow \infty} 0, r_K \xrightarrow{K \rightarrow \infty} +\infty.$$

• When a trait mutation occurs, the new trait of the descendant is  $x+k \in \mathcal{X}$  with  $k$  chosen according to the probability measure  $m(x, k)dk$ .

• When a marker mutation occurs, the new marker of the descendant is  $u+h \in \mathcal{U}$  with  $h$  chosen according to the probability measure  $G_K(u, dh)$ . For any  $u \in \mathcal{U}$ ,  $G_K(u, \cdot)$  is approximated as follows when  $K$  tends to infinity:

$$\lim_{K \rightarrow +\infty} \sup_{u \in \mathcal{U}} \left| \frac{r_K}{K} \int_{\mathcal{U}} (\phi(u+h) - \phi(u)) G_K(u, dh) - A\phi \right| = 0,$$

where  $(A, \mathcal{D}(A))$  is the generator of a Feller semigroup and  $\phi \in \mathcal{D}(A) \subseteq C_b(\mathcal{U}, \mathbb{R})$ , the set of continuous bounded real functions on  $\mathcal{U}$ .

• An individual with trait  $x$  and marker  $u$  dies with intrinsic death rate  $0 \leq d(x) \leq \bar{d}$ , the function  $d$  being continuous and  $\bar{d}$  a positive real. Moreover the individual experiences competition the effect of which is an additional death rate

$$\eta(x) \cdot C * \nu_t^K(x) = \frac{\eta(x)}{K} \sum_{i=1}^{N_t^K} C(x - x_i).$$

The quantity  $C(x - x_i)$  describes the competition pressure exerted by an individual with trait  $x_i$  on an individual with trait  $x$ . We assume that the functions  $C$  and  $\eta$  are continuous and that there exists  $\underline{\eta} > 0$  such that

$$\forall x, y \in \mathcal{X}, \eta(x) C(x - y) \geq \underline{\eta} > 0. \quad (2.6)$$

A classical choice of competition function  $C$  is  $C \equiv 1$  which is called "mean field case" or "logistic case". In that case the competition death rate is  $\eta(x)N_t^K/K$ .

As mentioned in the beginning of the Section, the authors work under the simplifying assumption that follows, ensuring that the population remains monomorphic between two events of invasion by a new trait.

**Assumption "Invasion implies fixation"** For all  $x \in \mathcal{X}$  and for almost every  $y \in \mathcal{X}$  :

either  $\frac{b(y) - d(y)}{\eta(y)C(y - x)} < \frac{b(x) - d(x)}{\eta(x)C(0)}$ , or  $\frac{b(y) - d(y)}{\eta(y)C(y - x)} > \frac{b(x) - d(x)}{\eta(x)C(0)}$  and  $\frac{b(x) - d(x)}{\eta(x)C(x - y)} < \frac{b(y) - d(y)}{\eta(y)C(0)}$ .

**Remark :**  $\hat{n}_x := \frac{b(x)-d(x)}{\eta(x)C(0)}$  is the equilibrium of the dynamical system that a large population size of individuals with trait  $x$  approximate.

Moreover, in the case of logistic populations with a constant  $C$ , this assumption is satisfied as soon as  $x \rightarrow \hat{n}_x$  is strictly monotonous.

### 4.3 Fleming-Viot process and marker evolution

The dynamics of the marker is first defined for a constant trait  $x$  via a Fleming-Viot process as defined below. This process generalizes to a potential infinity of markers for instance the measure-valued process describing two markers :  $F_t(dv) = X_t \delta_0(dv) + (1 - X_t) \delta_1(dv)$  with  $X$  for instance the neutral Wright-Fisher diffusion defined in Section 2, cf. (6) with  $s = 0$  and remarks below.

In the sequel, we denote by  $\mathcal{P}(\mathcal{U})$  and  $\mathcal{P}(\mathcal{X} \times \mathcal{U})$  the probability measure spaces respectively on  $\mathcal{U}$  and on  $\mathcal{X} \times \mathcal{U}$ , while  $\langle \nu | \phi \rangle$  denotes the integral of the measurable function  $\phi$  against the measure  $\nu$ . Also,  $\mathcal{M}_F(\mathcal{X} \times \mathcal{U})$  is the set of finite measures on  $\mathcal{X} \times \mathcal{U}$ .

**Definition 4.3.1.** *Given  $x \in \mathcal{X}$  and  $u \in \mathcal{U}$ , the Fleming-Viot process  $(F_t^u(x, t) \geq 0)$  indexed by  $x$ , started at time 0 with initial condition  $\delta_u$  and associated with the mutation operator  $A$  is the  $\mathcal{P}(\mathcal{U})$ -valued process whose law is characterized as the unique solution of the following martingale problem. For any  $\phi \in \mathcal{D}(A)$ ,*

$$M_t^x(\phi) = \langle F_t^u(x, \cdot) | \phi \rangle - \phi(u) - b(x) \int_0^t \langle F_s^u(x, \cdot) | A\phi \rangle ds \quad (16)$$

is a continuous square integrable martingale with quadratic variation process

$$\langle M^x(\phi) \rangle_t = \frac{2b(x)}{\hat{n}_x} \int_0^t (\langle F_s^u(x, \cdot) | \phi^2 \rangle - \langle F_s^u(x, \cdot) | \phi \rangle^2) ds.$$

**Remark :** The model presented in Section 2 provides simple illustrations for such kind of processes. For the solution  $X$  of (6), define :

$$F_t(dv) := X_t \delta_0(dv) + (1 - X_t) \delta_1(dv).$$

Then, with  $A \equiv 0$  (the only transitions for the traits are between 0 and 1),  $S$  a continuous function such that  $S(0) = s$  (the parameter of selection at the individual level) and  $S(1) = 0$ , for any  $\varphi$  measurable :

$$\begin{aligned} M_t(\phi) &:= \langle F_t | \phi \rangle - \langle F_0 | \phi \rangle - (\langle F_t | S \times \phi \rangle - \langle F_t | S \rangle \times \langle F_t | \phi \rangle) \\ &= [\phi(0) - \phi(1)] (X_t - x - \int_0^t s X_r (1 - X_r) dr) = [\phi(0) - \phi(1)] \sigma \int_0^t \sqrt{X_r(1 - X_r)} dB_r, \end{aligned}$$

where we used Ito's formula, is clearly a square-integrable martingale with quadratic variation :

$$\langle M(\phi) \rangle_t = [\phi(0) - \phi(1)]^2 \sigma^2 \int_0^t X_r (1 - X_r) dr = \sigma^2 \int_0^t (\langle F_r | \phi^2 \rangle - \langle F_r | \phi \rangle^2) dr.$$

Note that  $s$  is here a selective effect on the distribution that is not present in (16) because the marker is neutral.

Moreover, recalling the equation that described the state  $\mu_t$  of the population of groups in Section 2, one can also associate it to equation (16) with  $b(x)A = \mathcal{L}_{WF}$  and the martingale  $M(\phi)$  being identically zero (i.e. with no quadratic variations). In [18], the authors in fact derive another description of the population of groups in the limit of large population sizes (intra-groups and inter-groups). This limit is also described as such Fleming-Viot process, with a non-zero martingale because one does not neglect anymore the non-selective birth and death events of the groups. Again, its quadratic variations satisfies :  $d\langle M(\phi) \rangle_t \propto \int_0^t (\langle F_u | \phi^2 \rangle - \langle F_u | \phi \rangle^2) du$ . Of course, there is still the additional term involving  $r$  in the finite variation part (16). We refer for instance to [14] and [12] for a detailed presentation of Fleming-Viot processes.

#### 4.4 Convergence to the Substitution Fleming-Viot Process

We can now state the main theorem that describes the slow-fast dynamics of adaptive traits and neutral markers at the (trait) evolutionary time scale.

**Theorem 4.1.** *We work under Definition 4.2.1 and the assumption that "invasion implies fixation". Consider the initial condition  $\nu_0^K(dy, dv) = n_0^K \delta_{(x_0, u_0)}(dy, dv)$  with  $\lim_{K \rightarrow \infty} n_0^K = \hat{n}_{x_0}$  and  $\sup_{K \in \mathbb{N}^*} \mathbb{E}((n_0^K)^3) < +\infty$ . Then, the population process  $(\nu_{Kt}^K, t \geq 0)$  converges in law to the  $\mathcal{M}_F(\mathcal{X} \times \mathcal{U})$ -valued process  $(V_t(dy, dv), t \geq 0)$ .*

To define this Markov process, with initial condition  $\hat{n}_{x_0} \delta_{x_0} \delta_{u_0}$ , we only need to describe it until the first jump of the trait, which is given by an exponential law. Namely, the trait jumps from  $x_0$  to  $x_0 + k$  with rate :

$$b(x_0) \hat{n}_{x_0} \frac{[f(x_0 + k; x_0)]_+}{b(x_0 + k)} m(x_0, k) dk.$$

Then, given that this first jump occurs at time  $T$ , the law of the new marker is given by :

$$U \sim F_T^{u_0}(x_0, du), \text{ so that } V_T(dy, dv) = \hat{n}_{x_0+k} \delta_{x_0+k} \delta_U.$$

Then, for any  $t < T$ , we have :

$$V_t(dy, dv) = \hat{n}_{x_0} \delta_{x_0}(dy) F_t^{u_0}(x_0, dv),$$

and the process is defined recursively like this before the next jumps.

The convergence holds in the sense of finite dimensional distributions on  $\mathcal{M}_F(\mathcal{X} \times \mathcal{U})$ . In addition, the convergence also holds in the sense of occupation measures, i.e. the measure  $\nu_{KT}^K(dy; dv) dt$  on  $\mathcal{X} \times \mathcal{U} \times [0; T]$  converges weakly to the measure  $V_t(dy; dv) dt$  for any  $T > 0$ .

This process is called by the authors the Substitution Fleming-Viot Process (SFVP). It generalizes the Trait Substitution Process (TSP) introduced in [20] and also obtained as a limit of individual-based model in [7]. The TSP is in fact the marginal of the SFVP on the trait space. The jump rate of the TSP can be interpreted quite easily. The mutation rate of any resident is given by  $b(x_0) m(x_0, k) dk$ . The number of such residents at equilibrium is nearly  $\hat{n}_{x_0} K$ . While in competition with the resident population, the survival of the lineage of the mutant depends mainly on the period where the associated sub-population is too small to disrupt the resident population. From classical results of Branching process, it survives with probability  $[f(x_0 + k; x_0)]_+ / b(x_0 + k)$ .

Note that only beneficial mutations pass through, so that they shall invade quickly after reaching a non-negligible proportion in the population. The product provides the rate of occurrence of such successful mutation in the whole population, that gets divided by  $K$  in the new time-scale.

Given the recent and impressive progress in sequencing and analyzing these data between species, one has partly access to the marker dynamics. The selective dynamics is however much more difficult to follow, since one shall be able to trace very specifically the mutation effects and their benefice. It would thus be of high interest to be able to infer strong selective effects from the dynamics of the marker.

At each sweep, a very specific marker is selected. This effect is referred to as a genetic hitchhiking. It shall be chosen according to the law  $F_T^{u_0}(x_0, du)$ . We may expect that numerous selective sweeps should increase the variability of the markers. Yet, without assuming any effect on the marker dynamics from the selective component, it is not very clear that doing these steps more frequently shall increase the variance in the selected marker (after a comparable time span).

Nonetheless, such hitchhiking events usually leave a mark, at least for sexual reproduction species. Because of the frequent recombination events, the hitchhiking effect is mostly effective for markers linked to the selected allele. The diversity of variants is all the smaller that one gets closer in the genome to the selected allele. A well-known example is given by the selection for genes that favor the digestion of milk. As one can imagine, the analysis is much more demanding than the convergence to the SFVP. But the latter is a significant next step in this direction.

#### 4.5 The last time-scale of adaptive-dynamics

As the last time-scale presented in this work, Adaptive Dynamics is probably the one that gives to natural selection the most predictable effect. For the discussion above, we refer to [2], where the connection of Adaptive dynamics with the individual-based model is demonstrated through a single step of convergence, in the absence of singularity. In this context, the canonical equation of adaptive dynamics (CEAD) states that the population of interest can be considered monomorphic, and its trait  $x_t$  follows an ordinary differential equation of first order. The speed of the trait namely involves the mutation rate, the population size at equilibrium, some derivative of the fitness of invasion, and the squared effect of mutations in the direction of invasion. Remark that this last term is not the mean effect of mutations in this direction because the more effective is a mutation, the more likely it is to fix. In order to obtain such deterministic behavior, we again need the assumptions for the TSS, i.e. rare mutations as compared to the ecological time-scale, with negligible fluctuations around the size equilibrium and invasion implying fixation (cf. previous Subsection). Moreover, the CEAD relies on the assumption that mutations have infinitesimal effects, so that the trait evolves continuously by the accumulation of such mutations in large number. For the connection of the TSS to the CEAD, it means a coupled rescaling of time and of the fitness effects, which is rather natural. Yet, for the actual connection with the individual-based model, it introduces the major difficulty that any favorable mutation step shall be quite insignificant and yet shall replace effectively the dominant trait. In the same idea, the dominant population shall prevent deleterious yet almost neutral mutations to invade and filter favorable mutation with an invasion success still proportional to the mutation effect.

Nonetheless, the authors of [2] actually manage to demonstrate a regime of convergence to the CEAD, where these issues are rigorously controlled. Their individual-based model is quite close

to the previous one, except that there is no marker anymore and that the possible mutation steps are assumed to be on some discrete and finite grid (whose mesh size goes to 0), preventing large mutations. We thus use the same notations as previously (rather than the one of [2]) and refer to Subsection 4.2. The main difference is also a scaling parameter for the mutation effect :

- Here, when a trait mutation occurs (in a population with trait  $x$ ), the new trait of the descendant is  $x + \sigma_K k \in \mathcal{X}$  with  $k$  chosen according to the probability measure  $\{m(x, k)\}_{k \in [-A, A]}$ . The mutation rate is also allowed to depend on  $x$ , and is thus given by  $q_K M(x)$ .

There is also an additive issue of regularity for the birth rate  $b$ , the death rate  $d$ , the mutation rate  $M$ , the sensibility to competition  $\eta$  and the competition kernel  $C$ . This is stated in Assumption 1 with the other assumptions we have made in Subsection 4.2, plus the fact that  $b(x) > d(x)$  and  $C(x, x)$  is uniformly upper-bounded for any  $x \in \mathcal{X}$ .

There is also a last assumption to ensure the absence of singularity, based on the invasion fitness of a mutant  $y$  in a resident population  $x$  :

$$f(y, x) := b(y) - d(y) - \eta(y) C(y, x) \hat{n}_x.$$

It indicates the mean growth rate of a mutant population  $y$  as long as it is still negligible as compared to the resident population  $x$ . As stated in the convergence to the TSS, the invasion probability of such a mutant population initiated by a single individual tends to  $f(y, x)_+/b(y)$  (for large population size). Here,  $f_+$  is the positive part of  $f$ , meaning that deleterious mutations cannot invade.

**Assumption 3 :** For all  $x \in \mathcal{X}$ ,  $\partial_1 f(x, x) \neq 0$ .

Assumption 3 implies that either  $\forall x \in \mathcal{X} : \partial_1 f(x, x) > 0$  or  $\forall x \in \mathcal{X} : \partial_1 f(x, x) < 0$ . Therefore, coexistence of two traits is not possible. Without loss of generality, we can assume that,  $\forall x \in \mathcal{X}, \partial_1 f(x, x) > 0$ . In fact, a weaker assumption is sufficient, see Remark (i).

**Theorem 4.2.** Assume that Assumptions 1 and 3 hold and that there exists a small  $\alpha > 0$  such that

$$K^{-1/2+\alpha} \ll \sigma_K \ll 1$$

$$\text{and} \quad \exp(-K^\alpha) \ll q_K \ll \frac{\sigma_K^{1+\alpha}}{K \ln K}, \quad \text{as } K \rightarrow \infty.$$

Fix  $x_0 \in \mathcal{X}$  and let  $(N_0^K)_{K \geq 0}$  be a sequence of  $\mathbb{N}$ -valued random variables such that  $N_0^K/K$  converges in law, as  $K \rightarrow \infty$ , to the positive constant  $\hat{n}(x_0)$  and is bounded in  $L^p$ , for some  $p > 1$ .

For each  $K \geq 0$ , let  $\nu_t^K$  be the process generated by  $\mathcal{L}^K$  with monomorphic initial state  $(N_0^K/K) \cdot \delta_{\{x_0\}}$ . Then, for all  $T > 0$ , the sequence of rescaled processes,  $(\nu_{t/(K q_K \sigma_K^2)}^K)_{0 \leq t \leq T}$ , converges in probability, as  $K \rightarrow \infty$ , with respect to the Skorokhod topology on  $\mathbb{D}([0, T], \mathbb{M}(\mathcal{X}))$  to the measure-valued process  $\hat{n}(x_t) \delta_{x_t}$ , where  $(x_t)_{0 \leq t \leq T}$  is given as a solution of the CEAD,

$$\frac{dx_t}{dt} = \sum_{k=-A}^A k [k M(x_t) \hat{n}(x_t) \partial_1 f(x_t, x_t)]_+ m(x_t, k), \quad \text{with initial condition } x_0.$$

**Remarks :**

(i) The main result of the paper actually holds under weaker assumptions. More precisely, Assumption 3 can be replaced by the following :

**Assumption 3'.** *The initial state  $v_0^K$  has a.s. support  $\{x_0\}$  with  $x_0 \in \mathcal{X}$  satisfying  $\partial_1 f(x_0, x_0) \neq 0$ .*

The reason is that, since  $x \mapsto \partial_1 f(x, x)$  is continuous, Assumption 3 is satisfied locally. Since moreover  $x \mapsto \partial_1 f(x, x)$  is Lipschitz-continuous, the CEAD never reaches in finite time an evolutionary singularity (i.e. a value  $y \in \mathcal{X}$  such that  $\partial_1 f(y, y) = 0$ ). In particular, for a fixed  $T > 0$ , the CEAD only visits traits in some interval  $I$  of  $\mathcal{X}$  where  $\partial_1 f(x, x) \neq 0$ . By modifying the parameters of the model out of  $I$  in such a way that  $\partial_1 f(x, x) \neq 0$  everywhere in  $\mathcal{X}$ , we can apply Theorem 4.2 to this modified process  $\tilde{\nu}$ . Then, we deduce that  $\tilde{\nu}_{t/(Ku_K\sigma_K^2)}$  has support included in  $I$  for  $t \in [0, T]$  with high probability, and hence coincides with  $\nu_{t/(Ku_K\sigma_K^2)}$  on this time interval.

(ii) The condition  $q_K \ll \frac{\sigma_K^{1+\alpha}}{K \ln K}$  allows mutation events during an invasion phase of a mutant trait, but ensures that there is no "successful mutation" event during this phase.

(iii) The fluctuations of the resident population are of order  $K^{-1/2}$ , thus  $K^{-1/2+\alpha} \ll \sigma_K$  ensures that the sign of the initial growth rate is not influenced by the fluctuations of the population size. If a mutant trait  $y$  appears in a monomorphic population with trait  $x$ , then its initial growth rate is  $b - d(y) - \eta(y) C(y, x) \langle \nu_t^K | 1 \rangle = f(y, x) + o(\sigma_K) = (y - x) \partial_1 f(x, x) + o(\sigma_K)$  since  $y - x = O(\sigma_K)$ .

(iv)  $\exp(K^\alpha)$  is the time the resident population stays with high probability in an  $O(\varepsilon\sigma_K)$ -neighborhood of an attractive domain. This is a moderate deviation result. Thus, the condition  $\exp(-K^\alpha) \ll u_K$  ensures that the resident population is still in this neighborhood when a mutant occurs.

(vi) The time scale is  $(K q_K \sigma_K^2)^{-1}$  since the expected time for a mutation event is  $(K q_K)^{-1}$ , the probability that a mutant invades is of order  $\sigma_K$  and one needs  $O(\sigma_K^{-1})$  mutant invasions to see an  $O(1)$  change of the resident trait value.

Still, such a strong filtering of mutations is probably the most questionable issue of realism in term of the modeling of evolution. The fluctuations around the deterministic system shall be extremely small and the slightly deleterious mutations well-separated for such conclusions to be satisfied. It seems unlikely that selective effects are so dominant even for rather large populations (a million of individuals or so). Especially since there is usually a structuring of the population in term of non-heritable or loosely heritable characteristics, that are not neutral for survival. Think for instance of the individual positions or their level of infection by parasites. As we have seen in Section 1, it may induce much more variability compared to the case where all the individuals are identical. In this time-scale of infinitesimal mutations, we may expect to see, in addition to these selective effect mainly driven by favorable mutations, also some noise due to the fixation of almost neutral mutations. The trait of the population is still quite likely to follow the direction given by the fitness of invasion, yet its displacement might be quite different from the one given by the CEAD and not as regular.

## 5 Conclusion

As we have seen, there is a large class of processes that can be rigorously obtained as limits of individual-based models appropriately rescaled. This is to be expected since this representation fits the closest to simulations of populations, with the minimal set of assumptions to include any interaction of interest. Nonetheless, the precise convergence results are still quite challenging and impose to be very precise on the way time-scales are well-separated. By the coupled observation both of the proofs and the simulations, the main weaknesses of the model usually appear much more salient. Also, the next step for modeling the random fluctuations due to finite population sizes can be inferred as some second order expansion.

Notably, we could evaluate the difficulty in estimating the birth rate in Section 1 from the unique knowledge of the trait at birth (because there is a lot of fluctuations until the birth event). In Sections 2 and 3, the main issue appears to be that the predicted selection effects might be driven by too exceptional realizations of the stochastic process describing the dynamics of a typical individual. In simulations and actual life, such transitions from a very stable equilibrium to another one quite separated shall not happen exactly the way these models predict, notably for the time at which they occur. Although corrections can be made by some truncation, or by adding another term to the equation governing the density, the most realistic first step would possibly be to consider the Fleming-Viot representation introduced in Section 4. Yet, such measured-value stochastic process is quite more challenging to describe, with the white noise acting a bit everywhere. Also in Section 4, we have to discuss the assumption that mutations have only an infinitesimal effect, although sufficient to keep the population monomorphic and filter the mutations according to their effects. Given the difficulty of this demonstration, there is still much to do for the next steps, but we shall certainly continue to progress in this direction, like these processes of evolution.

This is the core of science to start with the most elementary models, like the ODE defining the growth rate of a population, and then to progressively incorporate more realistic features. With the current probabilistic tools at our disposal, it is clearly time to relate most of these models describing the dynamics of densities to the individual-based measure-valued processes. The main requirement is clearly that one averages over a large number of individuals, but this can be obtained in very various ways depending on the interactions of interest. Notably, such convergence results can be stated for very diverse time-scales, from the rapid adaptation of cells to the propagation of parasites to the evolution of species over millions of years. They provide an elementary way to unify the models of micro-biology, ecology and evolution. Thus, they enable to justify more rigorously the separation of the related time-scales or on the contrary motivate interesting coupling. Notably, the demographics and the selection effects are closely linked in Section 2, while in Section 4, the evolution of the marker as a measure-valued process is coupled in a very specific way to the punctual events of fixation for the traits under selection.

## References

- [1] Baar, M., Bovier, A.; The polymorphic evolution sequence for populations with phenotypic plasticity, *Electron. J. Probab.*, Volume 23, no. 72, pp. 1-27 (2018)
- [2] Baar, M., Bovier, A., Champagnat, N.; From stochastic, individual-based models to the canonical equation of adaptive dynamics in one step. *Ann. Appl. Probab.* 27, no. 2, 1093–1170. (2017)



- [3] Bansaye, V., Delmas, J-F., Marsalle, L., Tran, V.; Limit theorems for Markov processes indexed by continuous time Galton–Watson trees *Ann. Appl. Probab.*, V 21, Nbr 6, 2263-2314 (2011)
- [4] Bansaye, V., Méléard, S.; *Stochastic Models for Structured Populations, Scaling Limits and Long Time Behavior*, *Stochastics in Biological Systems*, 1.4 (2015)
- [5] Billiard, S., Collet, P., Fèrrière, R., Méléard, S., Tran, V.; The effect of competition and horizontal trait inheritance on invasion, fixation and polymorphism. *Journal of Theoretical Biology*, Elsevier, 411, pp.48-58 (2016)
- [6] Billiard, S., Ferrière, R., Méléard, S., Tran, V.; Stochastic dynamics of adaptive trait and neutral marker driven by eco-evolutionary feedbacks *J. Math. Biol.* 71: 1211-1242 (2015)
- [7] Champagnat, N.; A microscopic interpretation for adaptive trait substitution sequence model. *Stoch Process Appl* 116:1127–1160 (2006)
- [8] Champagnat, N., Ferrière, R., Méléard, S.; Unifying evolutionary dynamics: from individual stochastic processes to macroscopic models. *Theoretical Population Biology*, Elsevier, 69 (3), pp.297-321 (2006)
- [9] Champagnat, N., Henry, B.; A probabilistic approach to Dirac concentration in non-local models of adaptation with several resources, *Ann. Appl. Probab.*, V.29, N.4 (2019)
- [10] Champagnat, N., Méléard, S.; Polymorphic evolution sequence and evolutionary branching. *Probab Theory Relat Fields* 151:45–94 (2011)
- [11] Dawson, D.A.; *Introductory Lectures on Stochastic Population Systems*, Technical Report Series No. 451, Laboratory for Research in Stat. and Probab. (2010)
- [12] Dawson, D. A.; *Mesure-valued markov processes*. v. 1541 of *Lectures Notes in Math.*, Ecole d’Eté de probabilités de Saint-Flour XXI, Springer, pp 1–260, New York (1993)
- [13] Diekmann, O., Jabin, P.E., Mischler, S., Perthame, B.; The dynamics of adaptation: An illuminating example and a Hamilton-Jacobi approach. *Theor. Pop. Biol.*, 67:257–271 (2005)
- [14] Etheridge, A. *An introduction to superprocesses*, University Lecture Series, v. 20. American Mathematical Society, Providence (2000)
- [15] Marc Hoffmann. Adaptive estimation in diffusion processes. *Stochastic Process. Appl.*, 79(1):135-163 (1999)
- [16] Hoffmann, M., Marguet, A.; Statistical estimation in a randomly structured branching population, *Stochastic Process. Appl.*, to appear <https://doi.org/10.1016/j.spa.2019.02.015> (2019)
- [17] Kutoyants, Y. A.; *Statistical inference for ergodic diffusion processes*. Springer Science and Business Media (2013)
- [18] Luo, S., Mattingly, J.; Scaling limits of a model for selection at two scales. *Nonlinearity*. 30. 10.1088/1361-6544/aa5499. (2015)

- [19] Méléard, S., Tran, V.; Trait Substitution Sequence process and Canonical Equation for age-structured populations. *J. Math. Biology*, Springer (Germany), 58 (6), pp.881-921 (2009)
- [20] J.A.J. Metz and al. Adaptive dynamics, a geometrical study of the consequences of nearly faithful reproduction. S.J. Van Strien and S.M. Verduyn Lunel (ed.), *Stochastic and Spatial Structures of Dynamical Systems*, 45:183–231 (1996)
- [21] Mirrahimi S., Barles G., Perthame B., Souganidis P. E.: Singular hamilton-jacobi equation for the tail problem. *SIAM J. Math. Anal.* 44(6), 4297–4319 (2012)
- [22] Perthame, B., Gauduchon, M.; Survival thresholds and mortality rates in adaptive dynamics : conciliating deterministic and stochastic simulations. *Math. Med. Biol.* 27, 195–210 (2010)
- [23] Sznitman, A.S.; Topics in propagation of chaos. In *École d’Été de Probabilités de Saint-Flour XIX—1989*, Lecture Notes in Math.1464, pp 165–251. Springer, Berlin (1991).
- [24] Velleret, A.; Two level natural selection under the light of quasi-stationary distributions, on ArXiv : <https://arxiv.org/abs/1903.10161> (2019).
- [25] West, SA., Griffin, AS., Gardner A., Diggle SP. Social evolution theory for microorganisms. *Nat. Rev. Microbiol.* 4:597–607 (2006)