

Global existence, smooth and stabilization in a three-dimensional Keller-Segel-Navier-Stokes system with rotational flux

Jiashan Zheng*

School of Mathematics and Statistics Science,
Ludong University, Yantai 264025, P.R.China

Abstract

We consider the spatially 3-D version of the following Keller-Segel-Navier-Stokes system with rotational flux

$$\begin{cases} n_t + u \cdot \nabla n = \Delta n - \nabla \cdot (nS(x, n, c)\nabla c), & x \in \Omega, t > 0, \\ c_t + u \cdot \nabla c = \Delta c - c + n, & x \in \Omega, t > 0, \\ u_t + \kappa(u \cdot \nabla)u + \nabla P = \Delta u + n\nabla\phi, & x \in \Omega, t > 0, \\ \nabla \cdot u = 0, & x \in \Omega, t > 0 \end{cases} \quad (*)$$

under no-flux boundary conditions in a bounded domain $\Omega \subseteq \mathbb{R}^3$ with smooth boundary, where $\phi \in W^{2,\infty}(\Omega)$ and $\kappa \in \mathbb{R}$ represent the prescribed gravitational potential and the strength of nonlinear fluid convection, respectively. Here the matrix-valued function $S(x, n, c) \in C^2(\bar{\Omega} \times [0, \infty)^2; \mathbb{R}^{3 \times 3})$ denotes the rotational effect which satisfies $|S(x, n, c)| \leq C_S(1+n)^{-\alpha}$ with some $C_S > 0$ and $\alpha \geq 0$. Compared with the signal consumption case as in chemotaxis-(Navier-)Stokes system, the quantity c of system (*)

*Corresponding author. E-mail address: zhengjiashan2008@163.com (J.Zheng)

is no longer a priori bounded by its initial norm in L^∞ , which means that we have less regularity information on c . Moreover, the tensor-valued sensitivity functions result in new mathematical difficulties, mainly linked to the fact that a chemotaxis system with such rotational fluxes thereby loses an energy-like structure. In this paper, by seeking some new functionals and using the bootstrap arguments on system (*), we establish the existence of global weak solutions to system (*) for arbitrarily large initial data under the assumption $\alpha \geq 1$. Moreover, under an explicit condition on the size of C_S relative to C_N , we can secondly prove that in fact any such **weak** solution (n, c, u) becomes smooth ultimately, and that it approaches the unique spatially homogeneous steady state $(\bar{n}_0, \bar{n}_0, 0)$, where $\bar{n}_0 = \frac{1}{|\Omega|} \int_\Omega n_0$ and C_N is the best Poincaré constant. To the best of our knowledge, there are the first results on asymptotic behavior of the system.

Key words: Navier-Stokes system; Keller-Segel model; Stabilization; Global existence; Tensor-valued sensitivity

2010 Mathematics Subject Classification: 35K55, 35Q92, 35Q35, 92C17

1 Introduction

In this paper, we investigate the global existence and stabilization of the solutions to the following Keller-Segel-Navier-Stokes system with rotational flux

$$\begin{cases} n_t + u \cdot \nabla n = \Delta n - \nabla \cdot (nS(x, n, c)\nabla c), & x \in \Omega, t > 0, \\ c_t + u \cdot \nabla c = \Delta c - c + n, & x \in \Omega, t > 0, \\ u_t + \kappa(u \cdot \nabla)u + \nabla P = \Delta u + n\nabla\phi, & x \in \Omega, t > 0, \\ \nabla \cdot u = 0, & x \in \Omega, t > 0 \end{cases} \quad (1.1)$$

in a physical smoothly bounded domain $\Omega \subseteq \mathbb{R}^3$, under zero-flux boundary conditions

$$(\nabla n - nS(x, n, c)\nabla c) \cdot \nu = \nabla c \cdot \nu = 0, u = 0, \quad x \in \partial\Omega, t > 0 \quad (1.2)$$

and with prescribed initial data

$$n(x, 0) = n_0(x), c(x, 0) = c_0(x), u(x, 0) = u_0(x), \quad x \in \Omega, \quad (1.3)$$

where n, c, u, P, ϕ , respectively denote the bacterial density, chemical concentration, fluid velocity, the associated pressure and the prescribed gravitational potential. This system was initially proposed by Wang-Winkler-Xiang ([39, 40]) to describe chemotaxis-fluid interaction in cases when the evolution of the chemoattractant is essentially dominated by production through cells.

Before going into our mathematical analysis, we recall some important progresses on system (1.1) and its variants. When $u \equiv 0$, the PDE system (1.1) is reduced to the Keller-Segel system

$$\begin{cases} n_t = \Delta n - \nabla \cdot (nS(x, n, c)\nabla c), & x \in \Omega, t > 0, \\ c_t = \Delta c - c + n, & x \in \Omega, t > 0, \end{cases} \quad (1.4)$$

where the tensor-valued function (or the scalar function) S measures the chemotactic sensitivity and satisfies

$$S \in C^2(\bar{\Omega} \times [0, \infty)^2; \mathbb{R}^{N \times N}) \quad (1.5)$$

and

$$|S(x, n, c)| \leq C_S(1+n)^{-\alpha} \quad \text{for all } (x, n, c) \in \Omega \times [0, \infty)^2 \quad (1.6)$$

with some $C_S > 0$ and $\alpha \geq 0$. Here N denotes the space dimension. The global existence and finite time blow-up of system (1.4) have been widely investigated in the past decades. In fact, in the case $S(x, n, c) = 1$, solutions of (1.4) may blow up in finite time when $N \geq 2$ ([?, 46]). On the other hand, such explosion phenomena can be ruled out when $S(x, n, c) := S(n)$ is related to the prototypical assumption of volume-filling effect ([17, 42]). More precisely, $S(x, n, c) := (1 + n)^{-\alpha}$ is a scalar function, when $\alpha > 1 - \frac{2}{N}$, all solutions are global and uniformly bounded ([17]); while if $\Omega \subseteq \mathbb{R}^N (N \geq 2)$ is a ball and $\alpha < 1 - \frac{2}{N}$, the solution may blow up under some technical assumptions ([17, 42]). Quite a number of results have been investigated, including the system with the logistic terms (see [67, 38, 43, 36], for instance), the system with nonlinear diffusion (see [6, 20, 60, 59]) and so on. We refer to [2, 16] for the further reading.

When $n = c = 0$, the PDE system (1.1) becomes the Navier-Stokes system

$$\begin{cases} u_t + \kappa(u \cdot \nabla)u + \nabla P = \Delta u, & x \in \Omega, t > 0, \\ \nabla \cdot u = 0, & x \in \Omega, t > 0. \end{cases} \quad (1.7)$$

As we all know that the corresponding Navier-Stokes system (1.7) does not admit a satisfactory solution theory up to now. In fact, global weak solvability was studied in [22, 31], some local existence of smooth solutions are proved by [12] and [41].

In various situations, however, the migration of bacteria may be more complex because it can be effected by the changes in their living environment (see [7] and [28]). Considering that in nature some bacteria, like *Bacillus subtilis* and *Escherichia coli*, often live in a viscous fluid, Tuval and his cooperators ([37]) described the above biological phenomena and proposed the mathematical model consisting of oxygen diffusion and **consumption**, chemotaxis, and fluid dynamics. In fact, Tuval et al. ([37]) proposed the following chemotaxis-(Navier-)Stokes system in the context of signal **consumption** by cells

$$\begin{cases} n_t + u \cdot \nabla n = \Delta n - \nabla \cdot (nS(n)\nabla c), & x \in \Omega, t > 0, \\ c_t + u \cdot \nabla c = \Delta c - nf(c), & x \in \Omega, t > 0, \\ u_t + \kappa(u \cdot \nabla)u + \nabla P = \Delta u + n\nabla\phi, & x \in \Omega, t > 0, \\ \nabla \cdot u = 0, & x \in \Omega, t > 0, \end{cases} \quad (1.8)$$

where $f(c)$ and κ represent the consumption rate of the bacteria and the strength of nonlinear fluid convection, respectively. In particular, when the fluid flow is slow, we can use the Stokes equation instead of the Navier-Stokes one, i.e., $\kappa = 0$ (see [28, 10]). The chemotaxis fluid system has been studied in the last few years. In fact, by making use of the energy-type functionals, system (1.8) and its variants have attracted extensive attention (see, e.g., Chae et al. [5], Duan et al. [8], Liu and Lorz [27, 28], Tao and Winkler [35, 45, 47, 50], Zhang and Zheng [58], and references therein). For example, the well-posedness and stabilization in a two-dimensional bounded domain have been investigated in [45] and [47], respectively. In [47], Winkler showed that in bounded **convex** domains $\Omega \subseteq R^2$, the global classical solutions obtained in [45] stabilize to the spatially uniform equilibrium $(\frac{1}{|\Omega|} \int_{\Omega} n_0, 0, 0)$ as $t \rightarrow \infty$. For $\Omega \subseteq R^3$, the global existence of weak solution for system (1.8) has been proved in [50]. Recently, the eventual smoothness of weak solutions in the three-dimensional case has also been investigated by Winkler in [51]. For more literature related to this model, we can refer to [33, 34, 48, 52, 24] and the references therein.

Compared with the chemotaxis system and the chemotaxis-(Navier-)Stokes system, the Keller-Segel-Navier-Stokes system (1.1) is much less understood. In fact, compared with the **signal consumption case** as in chemotaxis-(Navier-)Stokes system (1.9), the quantity c of system (1.1) is no longer a priori bounded by its initial norm in L^∞ , which means that we have less regularity information on c . Now, let us briefly recapitulate the recent developments for system (1.1). To this end, in 2018, Wang-Winkler-Xiang ([39]) showed that when the tensor-valued sensitivity S satisfying (1.5) and (1.6) with $\alpha > 0$ and $\Omega \subseteq \mathbb{R}^2$ is a bounded **convex** domain with smooth boundary, system (1.1) admits a global-in-time classical and bounded solution. By deriving a new type of entropy-energy estimate, Zheng ([63]) extends the result of [39] to general bounded domains. In comparison to the result for the corresponding fluid-free system, it is easy to see that the restriction on α here is optimal (see [63, 39]). In the case of $N = 3$ and tensor-valued $S(x, n, c)$ satisfying (1.5) and (1.6), global weak solutions were shown to exist for $\alpha \geq \frac{3}{7}$ ([26]). Recently, Zheng and Ke ([19]) presented the existence of global and weak solutions for Keller-Segel-Navier-Stokes system (1.1) under the assumption that S satisfies (1.5) and (1.6) with $\alpha > \frac{1}{3}$, which, in light of the

known results for the fluid-free system, is an optimal restriction on α . For more works about the chemotaxis models with tensor sensitivity (and nonlinear diffusion), we refer readers to see [62, 53, 66, 3, 29, 25, 23, 62].

Experiments (see [56, 57]) find that the oriented migration of bacteria or cells may not be parallel to the gradient of the chemical substance, but may rather involve rotational flux components. This suggests that the sensitivity function $S(n)$ should be replaced by a matrix $S(x, n, c)$. As a consequence, chemotaxis systems with such rotational flux lose their energy structure, which plays a key role in the mathematical analysis. Thus, a generalization of the chemotaxis-fluid system (1.8) should be of the form

$$\begin{cases} n_t + u \cdot \nabla n = \Delta n - \nabla \cdot (nS(x, n, c)\nabla c), & x \in \Omega, t > 0, \\ c_t + u \cdot \nabla c = \Delta c - nf(c), & x \in \Omega, t > 0, \\ u_t + \kappa(u \cdot \nabla)u + \nabla P = \Delta u + n\nabla\phi, & x \in \Omega, t > 0, \\ \nabla \cdot u = 0, & x \in \Omega, t > 0, \end{cases} \quad (1.9)$$

where $S(x, n, c)$ is a tensor-valued function, which indicates the rotational effect. This generalization results in considerable mathematical difficulties due to the fact that chemotaxis systems (1.9) with such rotational fluxes lose some energy structure, which has served as a key to the analysis for scalar-valued S (see Winkler et al. [39, 49, 54] and Cao-Lankeit [4], Zheng [64]). Consequently, some new approaches are necessary when studying the global existence or even boundedness (see [49]). Quite a number of results on global existence and boundedness properties have also been obtained for the variant of (1.9) obtained on replacing Δn by nonlinear diffusion operators generalizing the porous medium-type choice Δn^m for several ranges of $m > 1$ (see e.g. [52, 48, 34, 33]).

Motivated by the above works, the first purpose of this paper is to show the global solvability of weak solutions to the problem (1.1).

Throughout this paper, we assume ϕ and the initial data satisfy

$$\phi \in W^{2,\infty}(\Omega) \quad (1.10)$$

and

$$\begin{cases} n_0 \in C(\bar{\Omega}) & \text{with } n_0 \geq 0, \\ c_0 \in W^{1,\infty}(\Omega) & \text{with } c_0 \geq 0 \text{ in } \bar{\Omega}, \\ u_0 \in D(A^\gamma) & \text{for some } \gamma \in (\frac{3}{4}, 1), \end{cases} \quad (1.11)$$

where A denotes the Stokes operator with domain $D(A) := W^{2,2}(\Omega) \cap W_0^{1,2}(\Omega) \cap L_\sigma^2(\Omega)$, and $L_\sigma^2(\Omega) := \{\varphi \in L^2(\Omega) \mid \nabla \cdot \varphi = 0\}$ ([31]).

In the context of these assumptions, the first of our main results can be read as follows.

Theorem 1.1. *Let $\Omega \subseteq \mathbb{R}^3$ be a bounded domain with smooth boundary. Suppose that the assumptions (1.5)–(1.6) and (1.10)–(1.11) hold. If*

$$\alpha \geq 1, \quad (1.12)$$

then problem (1.1)–(1.3) possesses at least one global weak solution (n, c, u) in terms of Definition 2.1 below. Furthermore, this solution satisfies

$$\begin{cases} n \in L_{loc}^\infty([0, \infty), L^2(\Omega)) \cap L_{loc}^2([0, \infty), W^{1,2}(\Omega)), \\ c \in L_{loc}^\infty([0, \infty), L^p(\Omega)) \cap L_{loc}^2([0, \infty), W^{1,2}(\Omega)) \quad \text{for any } p > 1, \\ u \in L_{loc}^2([0, \infty), W_{0,\sigma}^{1,2}(\Omega)) \cap L_{loc}^\infty([0, \infty), L_\sigma^2(\Omega)) \cap L_{loc}^{\frac{10}{3}}(\bar{\Omega} \times [0, \infty)) \end{cases} \quad (1.13)$$

and there exists $C > 0$ such that

$$\|n(\cdot, t)\|_{L^{2\alpha}(\Omega)} + \|c(\cdot, t)\|_{L^p(\Omega)} + \|u(\cdot, t)\|_{L^2(\Omega)} \leq C \quad \text{for all } t > 0. \quad (1.14)$$

Moreover, (n, c, u) can be obtained as the limit of solutions $(n_\varepsilon, c_\varepsilon, u_\varepsilon, P_\varepsilon)$ to the regularized problems (2.1) below in the sense that there exists $(\varepsilon_j)_{j \in \mathbb{N}} \subset (0, 1)$ such that $\varepsilon_j \searrow 0$ as $j \rightarrow \infty$ and

$$n_\varepsilon \rightarrow n \text{ as well as } c_\varepsilon \rightarrow c \text{ and } u_\varepsilon \rightarrow u \text{ a.e. in } \Omega \times (0, \infty)$$

as $\varepsilon = \varepsilon_j \searrow 0$.

Going beyond these existence statements, the study of global dynamic is a natural continuation. In fact, in [55], Winkler proved that in the three-dimensional bounded **convex** domains, Keller-Segel-Navier-Stokes system with logistic source $\rho n - \mu n^2$ possesses at least one globally generalized solution, moreover, if $\mu > \frac{\chi\sqrt{\rho+}}{4}$, then this solution converge to

the homogeneous steady state with respect to the topology of $L^1(\Omega) \times L^p(\Omega) \times L^2(\Omega)$ for $p \in [1, 6)$. However, the regularity properties and the eventual smoothness of arbitrary weak solutions are still open. The present paper shows that if $C_S < 2\sqrt{C_N}$, after some relaxation time, this weak solution enjoys further regularity properties and thereby complies with the concept of weak solutions. In fact, under an explicit condition on the size of C_S relative to C_N , we can secondly prove that in fact any such **weak** solution (n, c, u) becomes smooth ultimately, and that it approaches the unique spatially homogeneous steady state $(\bar{n}_0, \bar{n}_0, 0)$, where $\bar{n}_0 = \frac{1}{|\Omega|} \int_{\Omega} n_0$ and C_N is the best Poincaré constant and C_S is given by (1.6).

Theorem 1.2. *Under the assumptions of Theorem 1.1, and assume that in addition*

$$C_S < 2\sqrt{C_N}, \quad (1.15)$$

where C_N is the best Poincaré constant and C_S is given by (1.6). Then whenever (n_0, c_0, u_0) satisfies (1.11) with $n_0 \not\equiv 0$. Then there exist $T > 0$ and $P \in C^{1,0}(\bar{\Omega} \times [T, \infty))$ such that

$$\begin{cases} n \in C^{2,1}(\bar{\Omega} \times (T, \infty)), \\ c \in C^{2,1}(\bar{\Omega} \times (T, \infty)), \\ u \in C^{2,1}(\bar{\Omega} \times (T, \infty); \mathbb{R}^3), \end{cases} \quad (1.16)$$

and such that (n, c, u) solves the boundary value problem in (1.1)–(1.3) classically in $\bar{\Omega} \times [T, \infty)$. Moreover,

$$n(\cdot, t) \rightarrow \bar{n}_0, \quad c(\cdot, t) \rightarrow \bar{n}_0 \quad \text{and} \quad u(\cdot, t) \rightarrow 0 \quad \text{in} \quad L^\infty(\Omega)$$

as $t \rightarrow \infty$, where $\bar{n}_0 = \frac{1}{|\Omega|} \int_{\Omega} n_0$.

1.1 Mathematical challenges and Organization of the paper

The global existence in three dimensions requires a more delicate proof and only results in a weak solvability. The main reason for these difficulties is the dimension dependency of the Gagliardo-Nirenberg inequalities. System (1.1)–(1.3) incorporates structures of **3-dimensional** fluid dynamics and rotational flux, which involves more complex chemotactic cross-diffusion reinforced by **signal production** and brings about many considerable mathematical difficulties. In fact, according to [56, 57] (see also [39, 48, 49]), the tensor-valued

sensitivity functions result in new mathematical difficulties, mainly linked to the fact that a chemotaxis system with such rotational fluxes thereby loses an energy-like structure, which is totally different from [53]. Moreover, unlike the **signal consumption** system (1.9), we cannot gain the $L^\infty(\Omega)$ estimates of c via the maximum principle directly. To overcome these difficulties, some new approaches are necessary when studying the global existence and stabilization for system (1.1)–(1.3). The strategy for the proof of the Theorem 1.2 lies in identifying the functional

$$\|n_\varepsilon - \bar{n}_0\|_{L^2(\Omega)}^2 + \frac{B}{2} \|c_\varepsilon - \bar{n}_0\|_{L^2(\Omega)}^2 \quad (\text{for some } B > 0)$$

as eventual Lyapunov functional. Apparent estimates for its dissipation rate show L^2 -convergence, which together with the L^p - L^q estimates for the Stokes semigroup and a contraction mapping implies the decay of u_ε in $L^p(\Omega)$ with some $p \geq 6$. Next, applying the variation-of-constants formula for $c_\varepsilon - \bar{n}_0$ and n_ε , we can derive the exponential decay for $\int_\Omega |\nabla c_\varepsilon(x, t)|^2$ and $\int_\Omega |\nabla c_\varepsilon(x, t)|^4$ and boundeness of $\|n_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)}$ and $\|c_\varepsilon(\cdot, t)\|_{W^{1,\infty}(\Omega)}$ for some large enough $t > 2$. Then based on maximal Sobolev regularity in the Stokes evolution system and inhomogeneous linear heat equations, one can successively obtain further ultimate regularity properties of $u_\varepsilon, c_\varepsilon$ and n_ε which by standard Schauder theory imply eventual smoothness (see Lemmas 6.11–6.17). Finally, collecting the regularity for $n_\varepsilon, c_\varepsilon$ and u_ε allows for turning the weak decay information previously gathered into the desired uniform convergence statements and thereby complete the proof of Theorem 1.2 (see Lemmas 6.18–6.20).

2 Preliminaries

Our intention is to construct a global weak solution as the limit of smooth solutions of appropriately regularized problem. Since, the nonlinear boundary condition on n ($(\nabla n - nS(x, n, c)) \cdot \nu = 0 \quad x \in \partial\Omega, t > 0$) and the nonlinear convective term ($\kappa \neq 0$ in the Navie-Stokes subsystem of (1.1)–(1.3)) brings about a great challenge to the study of system (1.1)–(1.3), we shall first follow an idea of [39] (see also [19, 48]) to deal with some regularized

approximate problems. To this end, we consider the following approximate system of (1.1)–(1.3):

$$\left\{ \begin{array}{l} n_{\varepsilon t} + u_{\varepsilon} \cdot \nabla n_{\varepsilon} = \Delta n_{\varepsilon} - \nabla \cdot (n_{\varepsilon} F_{\varepsilon}(n_{\varepsilon}) S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon}) \nabla c_{\varepsilon}), \quad x \in \Omega, t > 0, \\ c_{\varepsilon t} + u_{\varepsilon} \cdot \nabla c_{\varepsilon} = \Delta c_{\varepsilon} - c_{\varepsilon} + n_{\varepsilon}, \quad x \in \Omega, t > 0, \\ u_{\varepsilon t} + \nabla P_{\varepsilon} = \Delta u_{\varepsilon} - \kappa(Y_{\varepsilon} u_{\varepsilon} \cdot \nabla) u_{\varepsilon} + n_{\varepsilon} \nabla \phi, \quad x \in \Omega, t > 0, \\ \nabla \cdot u_{\varepsilon} = 0, \quad x \in \Omega, t > 0, \\ \nabla n_{\varepsilon} \cdot \nu = \nabla c_{\varepsilon} \cdot \nu = 0, u_{\varepsilon} = 0, \quad x \in \partial\Omega, t > 0, \\ n_{\varepsilon}(x, 0) = n_0(x), c_{\varepsilon}(x, 0) = c_0(x), u_{\varepsilon}(x, 0) = u_0(x), \quad x \in \Omega, \end{array} \right. \quad (2.1)$$

where

$$F_{\varepsilon}(s) := \frac{1}{(1 + \varepsilon s)^3} \quad \text{for all } s \geq 0 \text{ and } \varepsilon > 0, \quad (2.2)$$

as well as

$$S_{\varepsilon}(x, n, c) := \rho_{\varepsilon}(x) S(x, n, c), \quad x \in \bar{\Omega}, \quad n \geq 0, \quad c \geq 0 \quad (2.3)$$

and

$$Y_{\varepsilon} w := (1 + \varepsilon A)^{-1} w \quad \text{for all } w \in L^2_{\sigma}(\Omega)$$

is a standard Yosida approximation and A is the realization of the Stokes operator (see [31]). Here, $(\rho_{\varepsilon})_{\varepsilon \in (0,1)} \in C^{\infty}_0(\Omega)$ is a family of standard cutoff functions satisfying $0 \leq \rho_{\varepsilon} \leq 1$ in Ω and $\rho_{\varepsilon} \nearrow 1$ in Ω as $\varepsilon \searrow 0$.

Due to the nonlinear convective term $(u \cdot \nabla)u$ in third equation of (1.1)–(1.3), it is hard to find a classical solution. We intend to obtain a weak solution by approximation. To begin with, the definition of weak solutions to (1.1)–(1.3) is given as follows:

Definition 2.1. By a global weak solution of (1.1)–(1.3) we mean a triple (n, c, u) of functions

$$\left\{ \begin{array}{l} n \in L^1_{loc}([0, \infty); \bar{\Omega}), \\ c \in L^1_{loc}([0, \infty); W^{1,1}(\Omega)), \\ u \in L^1_{loc}([0, \infty); W^{1,1}_0(\Omega); \mathbb{R}^3), \end{array} \right. \quad (2.4)$$

such that $n \geq 0$ and $c \geq 0$ a.e. in $\Omega \times (0, \infty)$,

$$\begin{aligned} u \otimes u &\in L^1_{loc}(\bar{\Omega} \times [0, \infty); \mathbb{R}^{3 \times 3}) \text{ and } n \text{ belongs to } L^1_{loc}(\bar{\Omega} \times [0, \infty)), \\ cu, nu, \text{ and } nS(x, n, c)\nabla c &\text{ belong to } L^1_{loc}(\bar{\Omega} \times [0, \infty); \mathbb{R}^3) \end{aligned} \quad (2.5)$$

that $\nabla \cdot u = 0$ a.e. in $\Omega \times (0, \infty)$, and that

$$\begin{aligned} & - \int_0^\infty \int_\Omega n \varphi_t - \int_\Omega n_0 \varphi(\cdot, 0) \\ = & - \int_0^\infty \int_\Omega \nabla n \cdot \nabla \varphi + \int_0^\infty \int_\Omega n S(x, n, c) \nabla c \cdot \nabla \varphi + \int_0^\infty \int_\Omega n u \cdot \nabla \varphi \end{aligned} \quad (2.6)$$

for any $\varphi \in C_0^\infty(\bar{\Omega} \times [0, \infty))$,

$$\begin{aligned} & - \int_0^\infty \int_\Omega c \varphi_t - \int_\Omega c_0 \varphi(\cdot, 0) \\ = & - \int_0^\infty \int_\Omega \nabla c \cdot \nabla \varphi - \int_0^T \int_\Omega c \varphi + \int_0^T \int_\Omega n \varphi + \int_0^\infty \int_\Omega c u \cdot \nabla \varphi \end{aligned} \quad (2.7)$$

for any $\varphi \in C_0^\infty(\bar{\Omega} \times [0, \infty))$ as well as

$$\begin{aligned} & - \int_0^\infty \int_\Omega u \varphi_t - \int_\Omega u_0 \varphi(\cdot, 0) - \kappa \int_0^\infty \int_\Omega u \otimes u \cdot \nabla \varphi \\ = & - \int_0^\infty \int_\Omega \nabla u \cdot \nabla \varphi - \int_0^\infty \int_\Omega n \nabla \phi \cdot \varphi \end{aligned} \quad (2.8)$$

for any $\varphi \in C_0^\infty(\bar{\Omega} \times [0, \infty); \mathbb{R}^3)$ fulfilling $\nabla \varphi \equiv 0$.

The following notion of weak solutions to (1.1)–(1.3) is taken from [39]. Here and in the sequel, for vectors $v \in \mathbb{R}^3$ and $w \in \mathbb{R}^3$ we let $v \otimes w$ denote the matrix $(a_{ij})_{i,j \in \{1,2,3\}} \in \mathbb{R}^{3 \times 3}$ defined on setting $a_{ij} := v_i w_j$ for $i, j \in \{1, 2, 3\}$.

The first lemma concerns the local solvability of system (2.1) in the classical sense. Applying the Schauder fixed point theorem as well as the standard regularity theory of parabolic equations and the bootstrap arguments, the local existence of solutions to system (1.1)–(1.3) can be established by the similar arguments as in Lemma 2.1 of [45] (see also [49]), therefore, we give the following lemma without proof.

Lemma 2.1. *Let $\Omega \subseteq \mathbb{R}^3$ be a bounded domain with smooth boundary. Suppose that (1.5) and (1.6) hold. Then there exist $T_{max,\varepsilon} \in (0, \infty]$ and a classical solution $(n_\varepsilon, c_\varepsilon, u_\varepsilon, P_\varepsilon)$ of (1.1)–(1.3) in $\Omega \times (0, T_{max,\varepsilon})$ such that*

$$\begin{cases} n_\varepsilon \in C^0(\bar{\Omega} \times [0, T_{max,\varepsilon})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max,\varepsilon})), \\ c_\varepsilon \in C^0(\bar{\Omega} \times [0, T_{max,\varepsilon})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max,\varepsilon})), \\ u_\varepsilon \in C^0(\bar{\Omega} \times [0, T_{max,\varepsilon})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max,\varepsilon})), \\ P_\varepsilon \in C^{1,0}(\bar{\Omega} \times (0, T_{max,\varepsilon})) \end{cases} \quad (2.9)$$

classically solving (1.1)–(1.3) in $\Omega \times [0, T_{max,\varepsilon})$. Moreover, n_ε and c_ε are nonnegative in $\Omega \times (0, T_{max,\varepsilon})$, and

$$\|n_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} + \|c_\varepsilon(\cdot, t)\|_{W^{1,\infty}(\Omega)} + \|A^\gamma u_\varepsilon(\cdot, t)\|_{L^2(\Omega)} \rightarrow \infty \text{ as } t \nearrow T_{max,\varepsilon}, \quad (2.10)$$

where γ is given by (1.11).

Now, we let $(n_\varepsilon, c_\varepsilon, u_\varepsilon)$ denote the local solution to (2.1) with the maximal time given by $T_{max,\varepsilon}$. In what follows, let $C, C_i, C_{i,*}, \tilde{C}_i, \tilde{C}_{i,*}, \rho_i, \lambda_{i,*}, \tilde{\rho}_{i,*}, M, M_i, \gamma_i, \alpha_i$ and μ_i denote some different constants, which are independent of ε and $T_{max,\varepsilon}$, and if no special explanation, they depend at most on $\Omega, \alpha, \nabla\phi, n_0, c_0$ and u_0 .

Let us start with recalling some properties which have been established in previous studies and are fundamental when discussing results concerning the global existence of classical solutions in the setting of (1.1)–(1.3). The following estimates of n_ε and c_ε are basic but important in the proof of our result.

Lemma 2.2. *The solution of (1.1)–(1.3) satisfies*

$$\int_{\Omega} n_\varepsilon = \int_{\Omega} n_0 \text{ for all } t \in (0, T_{max,\varepsilon}) \quad (2.11)$$

as well as

$$\int_{\Omega} c_\varepsilon \leq \max\left\{\int_{\Omega} n_0, \int_{\Omega} c_0\right\} \text{ for all } t \in (0, T_{max,\varepsilon}). \quad (2.12)$$

In the following, we proceed to derive ε -independent estimates. In fact, using the idea comes from [19] and [63], one could propose some regularity estimates for n_ε and c_ε by tracking the time evolution of a certain combinational functional of them. Here we only state it as a lemma, while for the detailed proof readers can refer to Lemma 3.3 of [19].

Lemma 2.3. *Suppose that (1.10)–(1.11) and (1.5)–(1.6) hold with $\alpha > \frac{1}{3}$. Then one can find $C > 0$ independent of ε such that for all $t \in (0, T_{max,\varepsilon})$, the solution of (1.1)–(1.3) satisfies*

$$\int_{\Omega} n_\varepsilon^{2\alpha}(x, t) + \int_{\Omega} c_\varepsilon^2(x, t) + \int_{\Omega} |u_\varepsilon(x, t)|^2 \leq C \text{ for all } t \in (0, T_{max,\varepsilon}). \quad (2.13)$$

Moreover, for all $t \in (0, T_{max,\varepsilon} - \tau)$, it holds that one can find a constant $C > 0$ such that

$$\int_t^{t+\tau} \int_{\Omega} [n_{\varepsilon}^{2\alpha-2} |\nabla n_{\varepsilon}|^2 + |\nabla c_{\varepsilon}|^2 + |\nabla u_{\varepsilon}|^2] \leq C, \quad (2.14)$$

where

$$\tau = \min\{1, \frac{1}{6}T_{max,\varepsilon}\}. \quad (2.15)$$

With Lemma 2.3 at hand, we are now in the position to show the solution of the approximate problem (2.1) is actually global in time, that is, $T_{max,\varepsilon} = \infty$.

Lemma 2.4. *Let $\alpha \geq 1$. Then for all $\varepsilon \in (0, 1)$, the solution of (2.1) is global in time.*

Proof. Firstly, we will prove that for any $\tau \in (0, T_{max,\varepsilon})$,

$$\|A^{\gamma}u_{\varepsilon}(\cdot, t)\|_{L^2(\Omega)} + \|c_{\varepsilon}(\cdot, t)\|_{W^{1,\infty}(\Omega)} + \|n_{\varepsilon}(\cdot, t)\|_{L^{\infty}(\Omega)} \leq C(\tau) \quad \text{for all } t \in (\tau, T_{max,\varepsilon}) \quad (2.16)$$

holds with some $C(\tau) > 0$. To do so, we fix $\tau \in (0, T_{max,\varepsilon})$ such that $\tau < 1$. In view of (2.2), we derive

$$F_{\varepsilon}(n_{\varepsilon}) \leq \frac{1}{\varepsilon^3 n_{\varepsilon}^3},$$

so that, by multiplying the first equation in (2.1) by $n_{\varepsilon}^{5+2\alpha}$ and using $\nabla \cdot u_{\varepsilon} = 0$,

$$\begin{aligned} & \frac{1}{6+2\alpha} \frac{d}{dt} \|n_{\varepsilon}\|_{L^{6+2\alpha}(\Omega)}^{6+2\alpha} + (5+2\alpha) \int_{\Omega} n_{\varepsilon}^{4+2\alpha} |\nabla n_{\varepsilon}|^2 \\ &= - \int_{\Omega} n_{\varepsilon}^{5+2\alpha} \nabla \cdot \left(n_{\varepsilon} \frac{1}{(1+\varepsilon n_{\varepsilon})^3} S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon}) \cdot \nabla c_{\varepsilon} \right) \\ &\leq (5+2\alpha) \int_{\Omega} n_{\varepsilon}^{5+2\alpha} \frac{1}{(1+\varepsilon n_{\varepsilon})^3} |S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon})| |\nabla n_{\varepsilon}| |\nabla c_{\varepsilon}| \quad \text{for all } t \in (0, T_{max,\varepsilon}). \end{aligned} \quad (2.17)$$

Recalling (1.6), by Young inequality, one can see that

$$\begin{aligned} & \int_{\Omega} n_{\varepsilon}^{5+2\alpha} \frac{1}{(1+\varepsilon n_{\varepsilon})^3} |S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon})| |\nabla n_{\varepsilon}| |\nabla c_{\varepsilon}| \\ &\leq C_S \int_{\Omega} n_{\varepsilon}^{5+2\alpha} \frac{1}{(1+\varepsilon n_{\varepsilon})^3} (1+n_{\varepsilon})^{-\alpha} |\nabla n_{\varepsilon}| |\nabla c_{\varepsilon}| \\ &\leq \frac{1}{\varepsilon^3} C_S \int_{\Omega} n_{\varepsilon}^{2+\alpha} |\nabla n_{\varepsilon}| |\nabla c_{\varepsilon}| \\ &\leq \frac{(5+2\alpha)}{2} \int_{\Omega} n_{\varepsilon}^{4+2\alpha} |\nabla n_{\varepsilon}|^2 + C_1 \int_{\Omega} |\nabla c_{\varepsilon}|^2 \quad \text{for all } t \in (0, T_{max,\varepsilon}), \end{aligned} \quad (2.18)$$

where C_1 is a positive constant, as all subsequently appearing constants C_2, C_3, \dots possibly depend on ε . This combined with (2.14) implies that

$$\int_{\Omega} n_{\varepsilon}^{6+2\alpha}(x, t) \leq C_2 \quad \text{for all } t \in (0, T_{max,\varepsilon}) \quad (2.19)$$

holds.

Relying on properties of the Yosida approximation Y_ε , we can also derive that

$$\|Y_\varepsilon u_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} = \|(I + \varepsilon A)^{-1} u_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq C_3 \|u_\varepsilon(\cdot, t)\|_{L^2(\Omega)} \leq C_4 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \quad (2.20)$$

and some positive constants C_3 and C_4 depend on ε .

Next, testing the projected Stokes equation $u_{\varepsilon t} + Au_\varepsilon = \mathcal{P}[-\kappa(Y_\varepsilon u_\varepsilon \cdot \nabla)u_\varepsilon + n_\varepsilon \nabla \phi]$ by Au_ε , we derive from the Young inequality as well as (2.19) and (2.20) that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|A^{\frac{1}{2}} u_\varepsilon\|_{L^2(\Omega)}^2 + \int_{\Omega} |Au_\varepsilon|^2 \\ &= \int_{\Omega} Au_\varepsilon \mathcal{P}(-\kappa(Y_\varepsilon u_\varepsilon \cdot \nabla)u_\varepsilon) + \int_{\Omega} \mathcal{P}[n_\varepsilon \nabla \phi] Au_\varepsilon \\ &\leq \frac{1}{2} \int_{\Omega} |Au_\varepsilon|^2 + \kappa^2 \int_{\Omega} |(Y_\varepsilon u_\varepsilon \cdot \nabla)u_\varepsilon|^2 + \|\nabla \phi\|_{L^\infty(\Omega)}^2 \int_{\Omega} n_\varepsilon^2 \\ &\leq \frac{1}{2} \int_{\Omega} |Au_\varepsilon|^2 + C_5 \int_{\Omega} |\nabla u_\varepsilon|^2 + C_6 \quad \text{for all } t \in (0, T_{max, \varepsilon}), \end{aligned} \quad (2.21)$$

which together with (2.14) implies

$$\int_{\Omega} |\nabla u_\varepsilon(x, t)|^2 \leq C_7 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \quad (2.22)$$

by some basic calculation. Now, let $h_\varepsilon(x, t) = \mathcal{P}[n_\varepsilon \nabla \phi - \kappa(Y_\varepsilon u_\varepsilon \cdot \nabla)u_\varepsilon]$. Then combined with (2.22) yields to

$$\|h_\varepsilon(\cdot, t)\|_{L^2(\Omega)} \leq C_8 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \quad (2.23)$$

by using (2.19) and (2.22). Now, we express $A^\gamma u_\varepsilon$ by its variation-of-constants representation and make use of well-known smoothing properties of the Stokes semigroup ([15]) to obtain $C_9 > 0$ such that for any $\tau \in (0, T_{max, \varepsilon})$,

$$\|A^\gamma u_\varepsilon(\cdot, t)\|_{L^2(\Omega)} \leq C_9 \quad \text{for all } t \in (\tau, T_{max, \varepsilon}), \quad (2.24)$$

where $\gamma \in (\frac{3}{4}, 1)$. Since, $D(A^\gamma)$ is continuously embedded into $L^\infty(\Omega)$ by $\gamma > \frac{3}{4}$, so that, (2.24) yields to for some positive constant C_{10} such that

$$\|u_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq C_{10} \quad \text{for all } t \in (\tau, T_{max, \varepsilon}). \quad (2.25)$$

Now, test the second equation of (2.1) by $-\Delta c_\varepsilon$ and obtain, upon two applications of Young's inequality, that

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\nabla c_\varepsilon\|_{L^2(\Omega)}^2 + \int_{\Omega} |\Delta c_\varepsilon|^2 + \int_{\Omega} |\nabla c_\varepsilon|^2 \\
&= - \int_{\Omega} n_\varepsilon \Delta c_\varepsilon + \int_{\Omega} \Delta c_\varepsilon u_\varepsilon \cdot \nabla c_\varepsilon \\
&\leq \frac{1}{2} \int_{\Omega} |\Delta c_\varepsilon|^2 + \int_{\Omega} n_\varepsilon^2 + \sup_{t \in (\tau, T_{max, \varepsilon})} \|u_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)}^2 \int_{\Omega} |\nabla c_\varepsilon|^2 \quad \text{for all } t \in (\tau, T_{max, \varepsilon}).
\end{aligned} \tag{2.26}$$

Recalling the bounds provided by (2.25) as well as (2.19) and (2.13), this immediately implies

$$\|c_\varepsilon(\cdot, t)\|_{W^{1,2}(\Omega)} \leq C_{11} \quad \text{for all } t \in (\tau, T_{max, \varepsilon}) \tag{2.27}$$

by some calculation. In light of Lemma 2.1 of [18] and the Young inequality, we have

$$\begin{aligned}
& \|\nabla c_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \\
&\leq C_{12}(1 + \|c_\varepsilon(\cdot, t) - u_\varepsilon(\cdot, t) \cdot c_\varepsilon(\cdot, t)\|_{L^4(\Omega)}) \\
&\leq C_{12}(1 + \|c_\varepsilon(\cdot, t)\|_{L^4(\Omega)} + \|u_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \|\nabla c_\varepsilon(\cdot, t)\|_{L^4(\Omega)}) \\
&\leq C_{12}(1 + \|c_\varepsilon(\cdot, t)\|_{L^4(\Omega)} + \|u_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \|\nabla c_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)}^{\frac{1}{2}} \|\nabla c_\varepsilon(\cdot, t)\|_{L^2(\Omega)}^{\frac{1}{2}}) \\
&\leq C_{13}(1 + \|\nabla c_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)}^{\frac{1}{2}}) \quad \text{for all } t \in (\tau, T_{max, \varepsilon}),
\end{aligned} \tag{2.28}$$

which combined with (2.27) implies that

$$\|c_\varepsilon(\cdot, t)\|_{W^{1,\infty}(\Omega)} \leq C_{14} \quad \text{for all } t \in (\tau, T_{max, \varepsilon}) \tag{2.29}$$

by using the Gagliardo–Nirenberg inequality.

Furthermore, applying the variation-of-constants formula to the n_ε -equation in (2.1), we derive that for any $T \in (\tau, T_{max, \varepsilon})$,

$$n_\varepsilon(\cdot, t) = e^{(t-\tau)\Delta} n_\varepsilon(\cdot, \tau) - \int_{\tau}^t e^{(t-s)\Delta} \nabla \cdot (n_\varepsilon(\cdot, s) \tilde{h}_\varepsilon(\cdot, s)) ds, \quad t \in (\tau, T), \tag{2.30}$$

where $\tilde{h}_\varepsilon := \frac{1}{(1+\varepsilon n_\varepsilon)^3} S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon + u_\varepsilon$. Next, by (1.6), (2.29) and (2.25), we also have

$$\|\tilde{h}_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq C_{15} \quad \text{for all } t \in (\tau, T_{max, \varepsilon}),$$

so that, we can thus estimate

$$\begin{aligned}
\|n_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} &\leq \|e^{(t-\tau)\Delta}n_\varepsilon(\cdot, \tau)\|_{L^\infty(\Omega)} + \int_\tau^t \|e^{(t-s)\Delta}\nabla \cdot (n_\varepsilon(\cdot, s)\tilde{h}_\varepsilon(\cdot, s))\|_{L^\infty(\Omega)} ds \\
&\leq C_{16}(t-\tau)^{-\frac{3}{2}}\|n_0\|_{L^1(\Omega)} + C_{16} \int_\tau^t (t-s)^{-\frac{7}{8}}e^{-\lambda_1(t-s)}\|n_\varepsilon(\cdot, s)\tilde{h}_\varepsilon(\cdot, s)\|_{L^4(\Omega)} ds \\
&\leq C_{18} + C_{17} \int_\tau^t (t-s)^{-\frac{7}{8}}e^{-\lambda_1(t-s)}\|n_\varepsilon(\cdot, s)\|_{L^4(\Omega)} ds \\
&\leq C_{18} + C_{19} \int_\tau^t (t-s)^{-\frac{7}{8}}e^{-\lambda_1(t-s)}\|n_\varepsilon(\cdot, s)\|_{L^\infty(\Omega)}^{\frac{1}{2}}\|n_\varepsilon(\cdot, s)\|_{L^2(\Omega)}^{\frac{1}{2}} ds \\
&\leq C_{18} + C_{20} \sup_{s \in (\tau, T_{max, \varepsilon})} \|n_\varepsilon(\cdot, s)\|_{L^\infty(\Omega)}^{\frac{1}{2}} \quad \text{for all } t \in (\tau, T),
\end{aligned} \tag{2.31}$$

where

$$C_{20} = \int_0^\infty s^{-\frac{7}{8}}e^{-\lambda_1 s}\|n_\varepsilon(\cdot, s)\|_{L^2(\Omega)}^{\frac{1}{2}} < +\infty,$$

λ_1 is the first nonzero eigenvalue of $-\Delta$ on Ω under the Neumann boundary condition. And thereby

$$\|n_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq C_{21} \quad \text{for all } t \in (\tau, T_{max, \varepsilon}) \tag{2.32}$$

by some basic calculation and $T \in (\tau, T_{max, \varepsilon})$ was arbitrary.

Together with (2.24), (2.29) and (2.32), yields to (2.16). Assume that $T_{max, \varepsilon} < \infty$. Due to (2.16), we apply Lemma 2.1 to reach a contradiction and hence the proof of this Lemma is completed. \square

3 Global existence for problem (1.1)–(1.3)

In order to prove Theorem 1.1, we must state some further regularity properties for the approximate solutions obtained in the last section. To this end, by Lemma 3.1, we immediately obtain the boundedness of c_ε with respect to the norm in $L^p(\Omega)$ for any $p > 1$.

Lemma 3.1. *Let $(n_\varepsilon, c_\varepsilon, u_\varepsilon)$ be the solution of (2.9). Then there exists a positive constant C independent of ε such that*

$$\|c_\varepsilon(\cdot, t)\|_{L^p(\Omega)} + \|n_\varepsilon(\cdot, t)\|_{L^2(\Omega)} \leq C \quad \text{for all } t > 0 \tag{3.1}$$

and any $p > 2$. Moreover, there exists $C > 0$ independent of ε such that, for each $T > 0$,

$$\int_0^T \int_\Omega c_\varepsilon^{p-2} |\nabla c_\varepsilon|^2 + \int_0^T \int_\Omega |\nabla n_\varepsilon|^2 \leq C(T+1). \tag{3.2}$$

Proof. Firstly, multiplying the first equation in (2.1) by n_ε , in view of (2.2) and using $\nabla \cdot u_\varepsilon = 0$, we derive

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|n_\varepsilon\|_{L^2(\Omega)}^2 + \int_{\Omega} |\nabla n_\varepsilon|^2 \\
&= - \int_{\Omega} n_\varepsilon \nabla \cdot (n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \cdot \nabla c_\varepsilon) \\
&\leq \int_{\Omega} n_\varepsilon F_\varepsilon(n_\varepsilon) |S_\varepsilon(x, n_\varepsilon, c_\varepsilon)| |\nabla n_\varepsilon| |\nabla c_\varepsilon| \text{ for all } t > 0.
\end{aligned} \tag{3.3}$$

Recalling (1.6) and (2.2) and using $\alpha \geq 1$, via the Young inequality, we derive

$$\begin{aligned}
& \int_{\Omega} n_\varepsilon F_\varepsilon(n_\varepsilon) |S_\varepsilon(x, n_\varepsilon, c_\varepsilon)| |\nabla n_\varepsilon| |\nabla c_\varepsilon| \\
&\leq C_S \int_{\Omega} |\nabla n_\varepsilon| |\nabla c_\varepsilon| \\
&\leq \frac{1}{2} \int_{\Omega} |\nabla n_\varepsilon|^2 + \frac{C_S^2}{2} \int_{\Omega} |\nabla c_\varepsilon|^2 \text{ for all } t > 0.
\end{aligned} \tag{3.4}$$

Here we have used the fact that

$$n_\varepsilon F_\varepsilon(n_\varepsilon) |S_\varepsilon(x, n_\varepsilon, c_\varepsilon)| \leq C_S n_\varepsilon (1 + n_\varepsilon)^{-\alpha} \leq C_S$$

by using (1.6) and $\alpha \geq 1$. Therefore, collecting (3.3) and (3.4) and using (2.14), we conclude that there exist positive constants C_1 and C_2 such that

$$\int_{\Omega} n_\varepsilon^2(x, t) \leq C_1 \text{ for all } t > 0 \tag{3.5}$$

and

$$\int_0^T \int_{\Omega} |\nabla n_\varepsilon|^2 \leq C_2(T + 1) \tag{3.6}$$

for all $T > 0$. Let $p > 2$. Testing the second equation in (2.1) by c_ε^{p-1} , using the fact $\nabla \cdot u_\varepsilon = 0$, and applying the Hölder inequality, we have

$$\begin{aligned}
& \frac{1}{p} \frac{d}{dt} \int_{\Omega} c_\varepsilon^p + (p-1) \int_{\Omega} c_\varepsilon^{p-2} |\nabla c_\varepsilon|^2 + \int_{\Omega} c_\varepsilon^p \\
&= \int_{\Omega} c_\varepsilon^{p-1} n_\varepsilon \\
&\leq \|n_\varepsilon(\cdot, t)\|_{L^2(\Omega)} \left(\int_{\Omega} c_\varepsilon^{2(p-1)} \right)^{\frac{1}{2}} \\
&\leq C_1^{\frac{1}{2}} \left(\int_{\Omega} c_\varepsilon^{2(p-1)} \right)^{\frac{1}{2}} \text{ for all } t > 0
\end{aligned} \tag{3.7}$$

by using (3.5). Then the Gagliardo–Nirenberg inequality and (2.12) enables us to see that

$$\begin{aligned}
& \left(\int_{\Omega} c_{\varepsilon}^{2(p-1)} \right)^{\frac{1}{2}} \\
&= \|c_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{4(p-1)}{p}}(\Omega)}^{\frac{2(p-1)}{p}} \\
&\leq C_2 \left(\|\nabla c_{\varepsilon}^{\frac{p}{2}}\|_{L^2(\Omega)}^{\frac{6(p-2)}{3p-2}} \|c_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{2}{p}}(\Omega)}^{\frac{2(p-1)}{p} - \frac{6(p-2)}{(3p-2)}} + \|c_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{2}{p}}(\Omega)}^{\frac{2(p-1)}{p}} \right) \\
&\leq C_3 \left(\|\nabla c_{\varepsilon}^{\frac{p}{2}}\|_{L^2(\Omega)}^{\frac{6(p-2)}{3p-2}} + 1 \right) \text{ for all } t > 0
\end{aligned}$$

and some positive constants C_2 and C_3 . Here we use the Young inequality to obtain that there is $C_4 > 0$ such that

$$\begin{aligned}
& \frac{1}{p} \frac{d}{dt} \int_{\Omega} c_{\varepsilon}^p + (p-1) \int_{\Omega} c_{\varepsilon}^{p-2} |\nabla c_{\varepsilon}|^2 + \int_{\Omega} c_{\varepsilon}^p \\
&\leq \frac{(p-1)}{2} \int_{\Omega} c_{\varepsilon}^{p-2} |\nabla c_{\varepsilon}|^2 + C_4 \text{ for all } t > 0.
\end{aligned} \tag{3.8}$$

Whereupon, our conclusion comes from (3.5) as well as (3.6) and (6.68) by using some basic calculation. \square

By interpolation and the embedding theorem, the estimates from Lemma 3.1 and Lemma 2.3 imply bounds for further spatio-temporal integrals.

Lemma 3.2. *Let $\alpha \geq 1$. Then there exists $C > 0$ independent of ε such that, for each $T > 0$, the solution of (2.1) satisfies*

$$\int_0^T \int_{\Omega} \left[|u_{\varepsilon}|^{\frac{10}{3}} + n_{\varepsilon}^{\frac{6+4\alpha}{3}} \right] dt + \int_0^T \|u_{\varepsilon}(\cdot, t)\|_{L^6(\Omega)}^2 dt \leq C(T+1) \tag{3.9}$$

as well as

$$\int_0^T \int_{\Omega} \left[|n_{\varepsilon} F_{\varepsilon}(n_{\varepsilon}) S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon}) \nabla c_{\varepsilon}|^2 + |n_{\varepsilon} u_{\varepsilon}|^{\frac{5(3+2\alpha)}{3(4+\alpha)}} \right] dt \leq C(T+1) \tag{3.10}$$

and

$$\int_0^T \int_{\Omega} \left[|u_{\varepsilon} \cdot \nabla c_{\varepsilon}|^{\frac{5}{4}} + |c_{\varepsilon} u_{\varepsilon}|^q \right] dt \leq C(T+1), \tag{3.11}$$

where $q < \frac{10}{3}$.

Proof. From Lemma 2.3 and (1.6), in light of the Gagliardo–Nirenberg inequality, we derive that there exist positive constants C_1, C_2, C_3, C_4, C_5 and C_6 such that

$$\begin{aligned}
\int_0^T \int_{\Omega} n_{\varepsilon}^{\frac{6+4\alpha}{3}} dt &\leq C_1 \int_0^T \left(\|\nabla n_{\varepsilon}\|_{L^2(\Omega)}^2 \|n_{\varepsilon}\|_{L^{2\alpha}(\Omega)}^{\frac{4\alpha}{3}} + \|n_{\varepsilon}\|_{L^{2\alpha}(\Omega)}^{\frac{6+4\alpha}{3}} \right) dt \\
&\leq C_2(T+1) \text{ for all } T > 0
\end{aligned} \tag{3.12}$$

as well as

$$\begin{aligned} \int_0^T \int_{\Omega} |u_{\varepsilon}|^{\frac{10}{3}} dt &\leq C_3 \int_0^T \left(\|\nabla u_{\varepsilon}\|_{L^2(\Omega)}^2 \|u_{\varepsilon}\|_{L^2(\Omega)}^{\frac{4}{3}} + \|u_{\varepsilon}\|_{L^2(\Omega)}^{\frac{10}{3}} \right) dt \\ &\leq C_4(T+1) \text{ for all } T > 0 \end{aligned} \quad (3.13)$$

and

$$\begin{aligned} \int_0^T \|u_{\varepsilon}(\cdot, t)\|_{L^6(\Omega)}^2 dt &\leq C_5 \int_0^T \|\nabla u_{\varepsilon}(\cdot, t)\|_{L^2(\Omega)}^2 dt \\ &\leq C_6(T+1) \text{ for all } T > 0, \end{aligned} \quad (3.14)$$

where the last inequality we have used the embedding $W_{0,\sigma}^{1,2}(\Omega) \hookrightarrow L^6(\Omega)$ and the Poincaré inequality. Now, in light of (1.6), applying (2.2), (3.12) and Lemma 2.3, one can derive from the Hölder inequality that for some positive constant C_7 such that

$$\begin{aligned} &\int_0^T \int_{\Omega} \left[|n_{\varepsilon} F_{\varepsilon}(n_{\varepsilon}) S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon}) \nabla c_{\varepsilon}|^2 + |u_{\varepsilon} \cdot \nabla c_{\varepsilon}|^{\frac{5}{4}} + |n_{\varepsilon} u_{\varepsilon}|^{\frac{5(3+2\alpha)}{3(4+\alpha)}} \right] \\ &\leq C_S \int_0^T \int_{\Omega} n_{\varepsilon}^2 (1+n_{\varepsilon})^{-2\alpha} |\nabla c_{\varepsilon}|^2 \\ &\quad + \left(\int_0^T \int_{\Omega} |\nabla c_{\varepsilon}|^2 \right)^{\frac{5}{8}} \left(\int_0^T \int_{\Omega} |u_{\varepsilon}|^{\frac{10}{3}} \right)^{\frac{3}{8}} + \left(\int_0^T \int_{\Omega} n_{\varepsilon}^{\frac{6+4\alpha}{3}} \right)^{\frac{5}{2(4+\alpha)}} \left(\int_0^T \int_{\Omega} |u_{\varepsilon}|^{\frac{10}{3}} \right)^{\frac{3+2\alpha}{2(4+\alpha)}} \\ &\leq C_S \int_0^T \int_{\Omega} |\nabla c_{\varepsilon}|^2 + \left(\int_0^T \int_{\Omega} |\nabla c_{\varepsilon}|^2 \right)^{\frac{5}{8}} \left(\int_0^T \int_{\Omega} |u_{\varepsilon}|^{\frac{10}{3}} \right)^{\frac{3}{8}} \\ &\quad + \left(\int_0^T \int_{\Omega} n_{\varepsilon}^{\frac{6+4\alpha}{3}} \right)^{\frac{5}{2(4+\alpha)}} \left(\int_0^T \int_{\Omega} |u_{\varepsilon}|^{\frac{10}{3}} \right)^{\frac{3+2\alpha}{2(4+\alpha)}} \\ &\leq C_7(T+1) \text{ for all } T > 0 \end{aligned} \quad (3.15)$$

by using $\alpha \geq 1$, where C_S is the same as (1.6).

Next, for any $q < \frac{10}{3}$, by using (3.1), one can find $C_8 > 0$ independent of ε such that

$$\|c_{\varepsilon}(\cdot, t)\|_{L^{\frac{10q}{10-3q}}(\Omega)} \leq C_8 \text{ for all } t > 0, \quad (3.16)$$

so that, combined with (3.13) yields to

$$\begin{aligned} &\int_0^T \int_{\Omega} [|c_{\varepsilon} u_{\varepsilon}|^q] \\ &\leq \left(\int_0^T \int_{\Omega} c_{\varepsilon}^{\frac{10}{3}} \right)^{\frac{10-3q}{10}} \left(\int_0^T \int_{\Omega} |u_{\varepsilon}|^{\frac{10}{3}} \right)^{\frac{3q}{10}} \\ &\leq C_9(T+1) \text{ for all } T > 0 \end{aligned} \quad (3.17)$$

with some positive constant C_9 independent of ε . Therefore a combination of (3.12)–(3.17) directly implies this lemma. \square

4 Regularity properties of time derivatives

To construct a weak solution of problem (1.1)–(1.3) by a limit process from the regularized problems (2.1), we shall rely on the following regularity property with respect to the time variable.

Lemma 4.1. *Let $\alpha \geq 1$, and assume that (1.10) and (1.11) hold. Then there exists $C > 0$ independent of ε such that*

$$\int_0^T \|\partial_t n_\varepsilon(\cdot, t)\|_{(W^{1, \frac{5}{2}}(\Omega))^*}^{\frac{5}{3}} dt \leq C(T+1) \text{ for all } T \in (0, \infty) \quad (4.1)$$

as well as

$$\int_0^T \|\partial_t c_\varepsilon(\cdot, t)\|_{(W^{1,2}(\Omega))^*}^2 dt \leq C(T+1) \text{ for all } T \in (0, \infty) \quad (4.2)$$

and

$$\int_0^T \|\partial_t u_\varepsilon(\cdot, t)\|_{(W_{0,\sigma}^{1,5}(\Omega))^*}^{\frac{5}{4}} dt \leq C(T+1) \text{ for all } T \in (0, \infty). \quad (4.3)$$

Proof. Firstly, testing the first equation of (2.1) by certain $\varphi \in C^\infty(\bar{\Omega})$, we have

$$\begin{aligned} & \left| \int_{\Omega} (n_{\varepsilon,t}) \varphi \right| \\ &= \left| \int_{\Omega} [\Delta n_\varepsilon - \nabla \cdot (n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon) - u_\varepsilon \cdot \nabla n_\varepsilon] \varphi \right| \\ &= \left| \int_{\Omega} [-\nabla n_\varepsilon \cdot \nabla \varphi + n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon \cdot \nabla \varphi + n_\varepsilon u_\varepsilon \cdot \nabla \varphi] \right| \\ &\leq \left| \int_{\Omega} \left[\|\nabla n_\varepsilon\|_{L^{\frac{5}{3}}(\Omega)} + \|n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon\|_{L^{\frac{5}{3}}(\Omega)} + \|n_\varepsilon u_\varepsilon\|_{L^{\frac{5}{3}}(\Omega)} \right] \|\varphi\|_{W^{1, \frac{5}{2}}(\Omega)} \right| \end{aligned}$$

for all $t > 0$.

Therefore, in view of $\alpha \geq 1$, Lemma 2.3 and Lemma 3.2, we deduce from the Young inequality that for some $C_1 > 0$ and $C_2 > 0$ such that

$$\begin{aligned} & \int_0^T \|\partial_t n_\varepsilon(\cdot, t)\|_{(W^{1, \frac{5}{2}}(\Omega))^*}^{\frac{5}{3}} dt \\ &\leq C_1 \left\{ \int_0^T \int_{\Omega} |\nabla n_\varepsilon|^2 + \int_0^T \int_{\Omega} |n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon|^2 + \int_0^T \int_{\Omega} |n_\varepsilon u_\varepsilon|^{\frac{5}{3}} + T \right\} \quad (4.4) \\ &\leq C_2(T+1) \text{ for all } T > 0. \end{aligned}$$

Likewise, given any $\varphi \in C^\infty(\bar{\Omega})$, we may test the second equation in (2.1) against φ to conclude that

$$\begin{aligned} \left| \int_{\Omega} \partial_t c_\varepsilon(\cdot, t) \varphi \right| &= \left| \int_{\Omega} [\Delta c_\varepsilon - c_\varepsilon + n_\varepsilon - u_\varepsilon \cdot \nabla c_\varepsilon] \cdot \varphi \right| \\ &= \left| - \int_{\Omega} \nabla c_\varepsilon \cdot \nabla \varphi - \int_{\Omega} c_\varepsilon \varphi + \int_{\Omega} n_\varepsilon \varphi + \int_{\Omega} c_\varepsilon u_\varepsilon \cdot \nabla \varphi \right| \\ &\leq \left\{ \|\nabla c_\varepsilon\|_{L^2(\Omega)} + \|c_\varepsilon\|_{L^2(\Omega)} + \|n_\varepsilon\|_{L^2(\Omega)} + \|c_\varepsilon u_\varepsilon\|_{L^2(\Omega)} \right\} \|\varphi\|_{W^{1,2}(\Omega)} \end{aligned}$$

for all $t > 0$. Thus, from Lemma 3.2 and Lemma 2.3 again, we invoke the Young inequality again and obtain that there exist positive constants C_3 and C_4 such that

$$\begin{aligned} &\int_0^T \|\partial_t c_\varepsilon(\cdot, t)\|_{(W^{1,2}(\Omega))^*}^2 dt \\ &\leq C_3 \left(\int_0^T \int_{\Omega} |\nabla c_\varepsilon|^2 + \int_0^T \int_{\Omega} n_\varepsilon^2 + \int_0^T \int_{\Omega} c_\varepsilon^5 + \int_0^T \int_{\Omega} |u_\varepsilon|^{\frac{10}{3}} + T \right) \\ &\leq C_4(T+1) \text{ for all } T > 0. \end{aligned}$$

Finally, for any given $\varphi \in C_{0,\sigma}^\infty(\Omega; \mathbb{R}^3)$, we infer from the third equation in (2.1) that

$$\left| \int_{\Omega} \partial_t u_\varepsilon(\cdot, t) \varphi \right| = \left| - \int_{\Omega} \nabla u_\varepsilon \cdot \nabla \varphi - \kappa \int_{\Omega} (Y_\varepsilon u_\varepsilon \otimes u_\varepsilon) \cdot \nabla \varphi + \int_{\Omega} n_\varepsilon \nabla \phi \cdot \varphi \right| \text{ for all } t > 0.$$

Now, Lemma 2.3, Lemma 3.2 and (1.10), we also derive that there exist positive constants C_5 as well as C_6 and C_7 such that

$$\begin{aligned} &\int_0^T \|\partial_t u_\varepsilon(\cdot, t)\|_{(W_{0,\sigma}^{1,2}(\Omega))^*}^2 dt \\ &\leq C_5 \left(\int_0^T \int_{\Omega} |\nabla u_\varepsilon|^2 + \int_0^T \int_{\Omega} |Y_\varepsilon u_\varepsilon \otimes u_\varepsilon|^2 + \int_0^T \int_{\Omega} n_\varepsilon^2 \right) \\ &\leq C_6 \left(\int_0^T \int_{\Omega} |\nabla u_\varepsilon|^2 + \int_0^T \int_{\Omega} |Y_\varepsilon u_\varepsilon|^2 + \int_0^T \int_{\Omega} |u_\varepsilon|^{\frac{10}{3}} + \int_0^T \int_{\Omega} n_\varepsilon^2 + T \right) \\ &\leq C_7(T+1) \text{ for all } T > 0, \end{aligned}$$

which implies (4.3) and whereby the proof is completed. \square

5 Passing to the Limit: Proof of Theorem 1.1

In this section we discuss the convergence of approximate solutions and complete the proof of Theorem 1.1. To this end, based on the above lemmas (see Lemma 2.3, Lemma 3.1 as well as Lemma 3.2 and Lemma 4.1), we can take the limit of the approximated solution

$(n_\varepsilon, c_\varepsilon, u_\varepsilon)$ as $\varepsilon \searrow 0$ and show that the limit is a global weak solution in the sense of the above definition (see Definition 2.1).

Lemma 5.1. *Let (1.5), (1.6), (1.10), and (1.11) hold, and suppose that $\alpha \geq 1$. There exists $(\varepsilon_j)_{j \in \mathbb{N}} \subset (0, 1)$ such that $\varepsilon_j \searrow 0$ as $j \rightarrow \infty$ and such that as $\varepsilon = \varepsilon_j \searrow 0$ we have*

$$n_\varepsilon \rightarrow n \text{ a.e. in } \Omega \times (0, \infty) \text{ and strongly in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \quad (5.1)$$

$$n_\varepsilon \rightharpoonup n \text{ weakly in } L_{loc}^{\frac{6+4\alpha}{3}}(\bar{\Omega} \times [0, \infty)) \quad (5.2)$$

$$\nabla n_\varepsilon \rightharpoonup \nabla n \text{ weakly in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \quad (5.3)$$

$$c_\varepsilon \rightarrow c \text{ strongly in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \text{ and a.e. in } \Omega \times (0, \infty), \quad (5.4)$$

$$\nabla c_\varepsilon^{\frac{p}{2}} \rightharpoonup \nabla c^{\frac{p}{2}} \text{ weakly in } L_{loc}^2([0, \infty); L^2(\Omega)) \text{ for all } p > 1, \quad (5.5)$$

$$c_\varepsilon \rightharpoonup c \text{ weakly }^* \text{ in } L_{loc}^\infty([0, \infty); L^p(\Omega)) \text{ for all } p > 1, \quad (5.6)$$

$$\nabla c_\varepsilon \rightarrow \nabla c \text{ a.e. in } \Omega \times (0, \infty), \quad (5.7)$$

$$u_\varepsilon \rightarrow u \text{ strongly in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \text{ and a.e. in } \Omega \times (0, \infty), \quad (5.8)$$

$$\nabla c_\varepsilon \rightharpoonup \nabla c \text{ weakly in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \quad (5.9)$$

as well as

$$\nabla u_\varepsilon \rightharpoonup \nabla u \text{ weakly in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \quad (5.10)$$

and

$$u_\varepsilon \rightharpoonup u \text{ weakly in } L_{loc}^{\frac{10}{3}}(\bar{\Omega} \times [0, \infty)) \quad (5.11)$$

with some triple (n, c, u) that is a global weak solution of (1.1)–(1.3) in the sense of Definition 2.1.

Proof. Based on Lemma 3.1, Lemma 2.3, Lemma 4.1, Lemma 3.2 and the Aubin–Lions lemma (see e.g. [30]), we can pick a sequence $(\varepsilon_j)_{j \in \mathbb{N}} \subset (0, 1)$ with $\varepsilon = \varepsilon_j \searrow 0$ as $j \rightarrow \infty$ such that (5.1)–(5.6) and (5.8)–(5.11) are valid with certain limit functions n, c and u belonging to the indicated spaces. Next, let $g_\varepsilon(x, t) := -c_\varepsilon + n_\varepsilon - u_\varepsilon \cdot \nabla c_\varepsilon$. With this notation, the second equation of (2.1) can be rewritten in component form as

$$c_{\varepsilon t} - \Delta c_\varepsilon = g_\varepsilon. \quad (5.12)$$

Thus, recalling Lemmas 3.1–3.2 and applying the Hölder inequality, we conclude that, for any $\varepsilon \in (0, 1)$, g_ε is bounded in $L^{\frac{5}{4}}(\Omega \times (0, T))$, and then we may invoke the standard parabolic regularity theory to (5.12) and infer that $(c_\varepsilon)_{\varepsilon \in (0, 1)}$ is bounded in $L^{\frac{5}{4}}((0, T); W^{2, \frac{5}{4}}(\Omega))$. Here, noting from (4.2) and the Aubin–Lions lemma that $(c_\varepsilon)_{\varepsilon \in (0, 1)}$ is relative compact in $L^{\frac{5}{4}}((0, T); W^{1, \frac{5}{4}}(\Omega))$. We can pick an appropriate subsequence that is still written as $(\varepsilon_j)_{j \in \mathbb{N}}$ such that $\nabla c_{\varepsilon_j} \rightarrow z_1$ in $L^{\frac{5}{4}}(\Omega \times (0, T))$ for all $T \in (0, \infty)$ and some $z_1 \in L^{\frac{5}{4}}(\Omega \times (0, T))$ as $j \rightarrow \infty$. Therefore, by (4.2), we can also derive that $\nabla c_{\varepsilon_j} \rightarrow z_1$ a.e. in $\Omega \times (0, \infty)$ as $j \rightarrow \infty$. In view of (5.9) and Egorov’s theorem, we conclude that $z_1 = \nabla c$ and hence (5.7) holds.

In the following proof, we shall prove that (n, c, u) is a weak solution of problem (1.1)–(1.3) in the sense of Definition 2.1, n, c and u satisfy (1.7), (3.9) and (3.10), (3.11), (3.12). Therefore, with the help of (5.1)–(5.4) and (5.8)–(5.10), we can derive (2.4). Now, by the nonnegativity of n_ε and c_ε , we obtain $n \geq 0$ and $c \geq 0$. Next, from (5.10) and $\nabla \cdot u_\varepsilon = 0$, we conclude that $\nabla \cdot u = 0$ a.e. in $\Omega \times (0, \infty)$. However, in view of (2.2), (1.5), as well as (3.10) and Lemma 3.2, we conclude that

$$n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon \rightharpoonup z_2 \text{ in } L^2(\Omega \times (0, T)) \text{ as } \varepsilon = \varepsilon_j \searrow 0 \text{ for each } T \in (0, \infty). \quad (5.13)$$

However, it follows from (1.5), (2.3), (3.10), (5.1), (5.4) and (5.7) that

$$n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon \rightarrow n S(x, n, c) \nabla c \text{ a.e. in } \Omega \times (0, \infty) \text{ as } \varepsilon = \varepsilon_j \searrow 0. \quad (5.14)$$

Again by Egorov’s theorem, we gain $z_2 = n S(x, n, c) \nabla c$, and therefore (5.13) can be rewritten as

$$n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon \rightharpoonup n S(x, n, c) \nabla c \text{ weakly in } L^2(\Omega \times (0, T)) \text{ as } \varepsilon = \varepsilon_j \searrow 0 \quad (5.15)$$

for each $T \in (0, \infty)$, which implies that the integrability of $n S(x, n, c) \nabla c$ in (2.5) as well. Thereupon, recalling (5.8), (5.1) and (3.10), we infer that, for each $T \in (0, \infty)$,

$$n_\varepsilon u_\varepsilon \rightharpoonup z_3 \text{ weakly in } L^{\frac{5(3+2\alpha)}{3(4+\alpha)}}(\Omega \times (0, T)), \text{ as } \varepsilon = \varepsilon_j \searrow 0, \quad (5.16)$$

which together with (5.1) and (5.8) implies

$$n_\varepsilon u_\varepsilon \rightarrow nu \text{ a.e. in } \Omega \times (0, \infty) \text{ as } \varepsilon = \varepsilon_j \searrow 0. \quad (5.17)$$

(5.16) along with (5.17) and Egorov's theorem guarantees that $z_3 = nu$, whereupon we derive from (5.16) that

$$n_\varepsilon u_\varepsilon \rightharpoonup nu \text{ weakly in } L^{\frac{5(3+2\alpha)}{3(4+\alpha)}}(\Omega \times (0, T)) \text{ as } \varepsilon = \varepsilon_j \searrow 0, \quad (5.18)$$

for each $T \in (0, \infty)$.

As a straightforward consequence of (5.4) and (5.8), it holds that

$$c_\varepsilon u_\varepsilon \rightarrow cu \text{ in } L^1_{loc}(\bar{\Omega} \times (0, \infty)) \text{ as } \varepsilon = \varepsilon_j \searrow 0. \quad (5.19)$$

Thus, the integrability of nu and cu in (2.5) is verified by (5.4) and (5.8).

Moreover, a combination of (5.8) and arguments in the proof of (6.26) of [19] yields that

$$Y_\varepsilon u_\varepsilon \otimes u_\varepsilon \rightarrow u \otimes u \text{ in } L^1_{loc}(\bar{\Omega} \times [0, \infty)) \text{ as } \varepsilon = \varepsilon_j \searrow 0. \quad (5.20)$$

Therefore, the integrability of $nS(x, n, c)\nabla c$, nu , cu , and $u \otimes u$ in (2.5) is verified by (5.15), (5.18), (5.19) and (5.20). Finally, according to (5.1)–(5.4), (5.8)–(5.10), (5.15), (5.18), (5.19) and (5.20), we may pass to the limit in the respective weak formulations associated with the regularized system (2.1) and obtain the integral identities (2.6)–(2.8). \square

Thereby our main result on global existence has actually been established already:

Proof of Theorem 1.1. Theorem 1.1 is a direct result of Lemma 5.1.

6 Asymptotic behavior

As the existence part of the main theorem is covered by the previous section, we will now turn our attention to verifying the desired convergence result. In the following, we will state some auxiliary lemmas, which will be frequently used in the sequel. To this end, we first briefly collect some known facts concerning the Stokes operator from [11, 48].

Lemma 6.1. *(Theorem 1 and Theorem 2 of [13]) The Helmholtz projection \mathcal{P} defines a bounded linear operator $\mathcal{P} : L^p(\Omega, \mathbb{R}^3) \rightarrow L^p_\sigma(\Omega)$; in particular, for any $p \in (1, \infty)$, there is $k_0(p) > 0$ such that*

$$\|\mathcal{P}w\|_{L^p(\Omega)} \leq k_0(p)\|w\|_{L^p(\Omega)} \quad (6.1)$$

for every $w \in L^p(\Omega)$.

Lemma 6.2. ([17, 44, 67]) *The Stokes operator A generates the analytic semigroup $(e^{-tA})_{t>0}$ in $L^2_\sigma(\Omega)$. Its spectrum satisfies $\lambda_1 := \inf \operatorname{Re}\sigma(A) > 0$ and we fix $\mu \in (0, \lambda_1)$. For any such μ , the following holds:*

(i) *For any $p \in (1, \infty)$ and $\gamma \geq 0$, there is $\kappa_1(p, \gamma) > 0$ such that*

$$\|A^\gamma e^{-tA} \varphi\|_{L^p(\Omega)} \leq \kappa_1(p, \gamma) t^{-\gamma} e^{-\mu t} \|\varphi\|_{L^p(\Omega)} \quad \text{for all } t > 0 \quad \text{and any } \varphi \in L^p_\sigma(\Omega). \quad (6.2)$$

(ii) *For p, q satisfying $1 < p \leq q < \infty$, there exists $\kappa_2(p, q) > 0$ such that*

$$\|e^{-tA} \varphi\|_{L^q(\Omega)} \leq \kappa_2(p, q) t^{-\frac{3}{2}(\frac{1}{p} - \frac{1}{q})} e^{-\mu t} \|\varphi\|_{L^p(\Omega)} \quad \text{for all } t > 0 \quad \text{and any } \varphi \in L^p_\sigma(\Omega). \quad (6.3)$$

(iii) *For p, q satisfying $1 < p \leq q < \infty$, there exists a positive constant $\kappa_3(p, q)$ such that*

$$\|\nabla e^{-tA} \varphi\|_{L^q(\Omega)} \leq \kappa_3(p, q) t^{-\frac{1}{2} - \frac{3}{2}(\frac{1}{p} - \frac{1}{q})} e^{-\mu t} \|\varphi\|_{L^p(\Omega)} \quad \text{for all } t > 0 \quad \text{and any } \varphi \in L^p_\sigma(\Omega). \quad (6.4)$$

(iv) *If $\gamma \geq 0$ and $1 < q < p < \infty$ satisfy $2\gamma - \frac{3}{q} \geq 1 - \frac{3}{p}$, then there is $\kappa_4(\gamma, p, q) > 0$ such that for all $\varphi \in D(A^\gamma)$*

$$\|\varphi\|_{W^{1,p}(\Omega)} \leq \kappa_4(\gamma, p, q) \|A^\gamma \varphi\|_{L^q(\Omega)}. \quad (6.5)$$

Lemma 6.3. (Lemma 4.6 of [9]) *Let $\lambda > 0, C > 0$, and suppose that $y \in C^1([0, \infty))$ and $h \in C^0([0, \infty))$ are nonnegative functions satisfying $y'(t) + \lambda y(t) \leq h(t)$ for some $\lambda > 0$ and all $t > 0$. Then if $\int_0^\infty h(s) ds \leq C$, we have $\lim_{t \rightarrow \infty} y(t) = 0$.*

Lemma 6.4. *Suppose that*

$$C_S < 2\sqrt{C_N}, \quad (6.6)$$

where C_N is the best Poincaré constant and C_S is given by (1.6). Then there exists $B > 0$ such that for each $\varepsilon \in (0, 1)$ we have

$$\begin{aligned} & \frac{B}{2} \frac{d}{dt} \|n_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2 + \frac{1}{2} \frac{d}{dt} \|c_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2 \\ & + \left(\frac{BC_N}{2} - \frac{1}{4}\right) \int_\Omega |n_\varepsilon - \bar{n}|^2 + \left(1 - \frac{BC_S^2}{2}\right) \int_\Omega |\nabla c_\varepsilon|^2 \\ & \leq 0 \quad \text{for all } t > 0, \end{aligned} \quad (6.7)$$

where

$$\bar{n}_0 = \frac{1}{|\Omega|} \int_\Omega n_0. \quad (6.8)$$

Proof. Firstly, we use the first equation in (1.1)–(1.3) and the fact that $\nabla \cdot u_\varepsilon = 0$ to compute

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|n_\varepsilon(\cdot, t) - \bar{n}_\varepsilon\|_{L^2(\Omega)}^2 \\
&= \int_{\Omega} (n_\varepsilon - \bar{n}_\varepsilon) [\Delta n_\varepsilon - u_\varepsilon \cdot \nabla n_\varepsilon - \nabla \cdot (n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon)] \\
&\leq - \int_{\Omega} |\nabla n_\varepsilon|^2 + \int_{\Omega} |n_\varepsilon| |S_\varepsilon(x, n_\varepsilon, c_\varepsilon)| |\nabla n_\varepsilon| |\nabla c_\varepsilon| \\
&\leq - \frac{1}{2} \int_{\Omega} |\nabla n_\varepsilon|^2 + \frac{C_S^2}{2} \int_{\Omega} \frac{n_\varepsilon^2}{(1+n_\varepsilon)^{2\alpha}} |\nabla c_\varepsilon|^2 \\
&\leq - \frac{1}{2} \int_{\Omega} |\nabla n_\varepsilon|^2 + \frac{C_S^2}{2} \int_{\Omega} |\nabla c_\varepsilon|^2 \quad \text{for all } t > 0
\end{aligned} \tag{6.9}$$

by using (2.2) as well as (1.6) and $\alpha \geq 1$, where

$$\bar{n}_\varepsilon = \frac{1}{|\Omega|} \int_{\Omega} n_\varepsilon \tag{6.10}$$

and C_S is the same as (1.6). We note from the Poincaré inequality that there is $C_N > 0$ such that

$$\|\varphi - \frac{1}{|\Omega|} \int_{\Omega} \varphi\|_{L^2(\Omega)}^2 \leq C_N \int_{\Omega} |\nabla \varphi|^2 \quad \text{for all } \varphi \in W^{1,2}(\Omega). \tag{6.11}$$

This combined with (6.9) yields to

$$\frac{1}{2} \frac{d}{dt} \|n_\varepsilon(\cdot, t) - \bar{n}_\varepsilon\|_{L^2(\Omega)}^2 \leq - \frac{C_N}{2} \int_{\Omega} |n_\varepsilon - \bar{n}_\varepsilon|^2 + \frac{C_S^2}{2} \int_{\Omega} |\nabla c_\varepsilon|^2 \quad \text{for all } t > 0, \tag{6.12}$$

which together with (2.11) implies

$$\frac{1}{2} \frac{d}{dt} \|n_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2 \leq - \frac{C_N}{2} \int_{\Omega} |n_\varepsilon - \bar{n}_0|^2 + \frac{C_S^2}{2} \int_{\Omega} |\nabla c_\varepsilon|^2 \quad \text{for all } t > 0. \tag{6.13}$$

Next, by means of the testing procedure, we may derive from the Young inequality that

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|c_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2 \\
&= \int_{\Omega} (c_\varepsilon - \bar{n}_0) [\Delta c_\varepsilon - u_\varepsilon \cdot \nabla c_\varepsilon - (c_\varepsilon - \bar{n}_0) + (n_\varepsilon - \bar{n}_0)] \\
&= \int_{\Omega} (c_\varepsilon - \bar{n}_0) (\Delta c_\varepsilon - u_\varepsilon \cdot \nabla c_\varepsilon) - \int_{\Omega} (c_\varepsilon - \bar{n}_0)^2 + \int_{\Omega} (c_\varepsilon - \bar{n}_0) (n_\varepsilon - \bar{n}_0) \\
&\leq - \int_{\Omega} |\nabla c_\varepsilon|^2 + \frac{1}{4} \int_{\Omega} (n_\varepsilon - \bar{n}_0)^2 \quad \text{for all } t > 0,
\end{aligned} \tag{6.14}$$

where we have used the fact that $\nabla \cdot u_\varepsilon = 0$ and $u_\varepsilon|_{\partial\Omega} = 0$.

In view of (6.6), we can choose $B > 0$ such that

$$1 - \frac{B}{2} C_S^2 > 0 \tag{6.15}$$

and

$$\frac{B}{2}C_N - \frac{1}{4} > 0. \quad (6.16)$$

Collecting (6.13)–(6.16), we thus infer that

$$\begin{aligned} & \frac{B}{2} \frac{d}{dt} \|n_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2 + \frac{1}{2} \frac{d}{dt} \|c_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2 \\ & + \left(\frac{BC_N}{2} - \frac{1}{4}\right) \int_\Omega |n_\varepsilon - \bar{n}_0|^2 + \left(1 - \frac{BC_S^2}{2}\right) \int_\Omega |\nabla c_\varepsilon|^2 \\ & \leq 0 \quad \text{for all } t > 0, \end{aligned} \quad (6.17)$$

which completes the proof. \square

Lemma 6.5. *Under the assumptions of Lemma 6.4, then for any $t > 0$, there exists $\rho_{*,1} > 0$ such that for all $\varepsilon \in (0, 1)$,*

$$\begin{aligned} & \|n_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2 + \|c_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2 \\ & \leq e^{-\rho_{*,1}t} [\|n_0(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2 + \|c_0(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2], \end{aligned} \quad (6.18)$$

and there exists $C_{*,1} > 0$ such that

$$\int_0^\infty \int_\Omega |\nabla c_\varepsilon|^2 + \int_0^\infty \int_\Omega |n_\varepsilon - \bar{n}_0|^2 + \int_0^\infty \int_\Omega |\nabla n_\varepsilon|^2 \leq C_{*,1} \quad \text{for all } \varepsilon \in (0, 1), \quad (6.19)$$

where \bar{n}_0 is given by (6.8).

Proof. Let $y(t) = \frac{B}{2} \frac{d}{dt} \|n_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2 + \frac{1}{2} \frac{d}{dt} \|c_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)}^2$. Then by (6.7), one can conclude that there exists a positive constant $\kappa_{1,*}$ (e.g. $\kappa_{1,*} \leq \min\{\frac{2(\frac{BC_N}{2} - \frac{1}{4})}{B}, 2C_N(1 - \frac{BC_S^2}{2})\}$) such that

$$y(t) + \kappa_{1,*}y(t) \leq 0,$$

so that, integrating the above inequality and (6.7) in time, we can get (6.18) and (6.20) by using (6.9). \square

In view of the interpolation, the estimates from Lemma 6.5 yield the following Lemma.

Proposition 6.1. *Under the assumptions of Lemma 6.4, there exists $C_{*,2} > 0$ such that for all $\varepsilon \in (0, 1)$,*

$$\int_0^\infty \int_\Omega n_\varepsilon^{\frac{10}{3}} \leq C_{*,2} \quad \text{for all } \varepsilon \in (0, 1). \quad (6.20)$$

Proof. Firstly, in view of (6.18), there exists a positive constant $\kappa_{1,**}$ such that

$$\|n_\varepsilon(\cdot, t)\|_{L^2(\Omega)} \leq \kappa_{1,**} \quad \text{for all } t > 0.$$

Therefore, in light of the Gagliardo–Nirenberg inequality, we derive from Lemma 6.5 that there exist positive constants $\kappa_{2,**}$ and $\kappa_{3,**}$ such that

$$\begin{aligned} \int_0^\infty \int_\Omega n_\varepsilon^{\frac{10}{3}} dt &\leq \kappa_{2,**} \int_0^\infty \left(\|\nabla n_\varepsilon\|_{L^2(\Omega)}^2 \|n_\varepsilon\|_{L^2(\Omega)}^{\frac{4}{3}} + \|n_\varepsilon\|_{L^2(\Omega)}^{\frac{10}{3}} \right) dt \\ &\leq \kappa_{3,**}. \end{aligned} \quad (6.21)$$

□

We are now prepared to prove the claimed asymptotic behavior of u_ε . In fact, exploiting the basic identity (6.18), this time in a more elaborate manner, yields the decay of u_ε with respect to the norm in $L^2(\Omega)$ in the first instance.

Lemma 6.6. *Under the assumptions of Lemma 6.4, for any $\eta > 0$, there are $T_{*,3} > 0, \rho_{*,3}$ and $C_{*,3} > 0$ such that for any $t > T_{*,3}$ such that whenever $\varepsilon \in (0, 1)$, we have*

$$\|u_\varepsilon(\cdot, t)\|_{L^2(\Omega)} \leq C_{*,3} e^{-\rho_{*,3}t}, \quad (6.22)$$

$$\int_t^{t+1} \|\nabla u_\varepsilon(\cdot, s)\|_{L^2(\Omega)}^2 ds < \eta \quad (6.23)$$

as well as

$$\int_t^{t+1} \|u_\varepsilon(\cdot, s)\|_{L^q(\Omega)}^2 ds < \eta \quad (6.24)$$

and

$$\int_t^{t+1} \|u_\varepsilon(\cdot, s)\|_{L^q(\Omega)} ds < \eta \quad (6.25)$$

for any $q \in [1, 6)$.

Proof. From straightforward calculations, while relying on (1.10), we derive the third equation in (2.1) that

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \|u_\varepsilon(\cdot, t)\|_{L^2(\Omega)}^2 \\ &= - \int_\Omega |\nabla u_\varepsilon|^2 + \int_\Omega n_\varepsilon \nabla \phi \cdot u_\varepsilon - \int_\Omega \nabla P_\varepsilon \cdot u_\varepsilon \\ &= - \int_\Omega |\nabla u_\varepsilon|^2 + \int_\Omega (n_\varepsilon - \bar{n}_0) \nabla \phi \cdot u_\varepsilon \\ &\leq - \int_\Omega |\nabla u_\varepsilon|^2 + \|\nabla \phi\|_{L^\infty(\Omega)} \left(\int_\Omega |n_\varepsilon - \bar{n}_0|^2 \right)^{\frac{1}{2}} \left(\int_\Omega |u_\varepsilon|^2 \right)^{\frac{1}{2}} \quad \text{for all } t > 0, \end{aligned} \quad (6.26)$$

where we have used the fact that $\nabla \cdot u_\varepsilon = 0$ and $u_\varepsilon|_{\partial\Omega} = 0$. Due to the Poincaré inequality again, there exists a positive constant η_N such that

$$\eta_N \int_{\Omega} |u_\varepsilon|^2 \leq \int_{\Omega} |\nabla u_\varepsilon|^2,$$

therefore, collecting (6.18) and (6.26), we derive that for some positive constants $\kappa_{1,***}$ and $\varrho_{1,***}$

$$\int_{\Omega} |u_\varepsilon(x, t)|^2 dx \leq \kappa_{1,***} e^{-\varrho_{1,***} t} \quad \text{for all } t > 0 \quad (6.27)$$

and

$$\int_t^{t+1} \|\nabla u_\varepsilon(\cdot, s)\|_{L^2(\Omega)}^2 ds \rightarrow 0 \quad \text{as } t \rightarrow \infty \quad (6.28)$$

and thereby proves (6.22)–(6.23). Finally, we make use of the embedding $W^{1,2}(\Omega) \hookrightarrow L^q(\Omega)$ (for any $q \in [1, 6)$) and the Young inequality to find that (6.24) and (6.25) hold. \square

With the decay property of $n_\varepsilon(\cdot, t) - \bar{n}_0$ (see Lemma 6.5) at hand, applying Lemma 6.6, we can now make sure that (6.18) implies the following a certain eventual regularity and decay of u_ε in $L^p(\Omega)$ with some $p \geq 6$.

Lemma 6.7. *Under the assumptions of Lemma 6.4, then for any $p \in [6, \infty)$, and $\varepsilon \in (0, 1)$, the solution of (2.1) satisfies*

$$\|u_\varepsilon\|_{L^\infty((t, \infty); L^p(\Omega))} \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (6.29)$$

Proof. We let $p \geq 6$ and choose $q \in (3, 6)$ which is close to 6 (e.g. $q = \frac{6p}{p+1}$) such that

$$\frac{1}{2} - \frac{3}{2p} - 3\left(\frac{1}{q} - \frac{1}{p}\right) > 0. \quad (6.30)$$

Now, we can fix $\gamma_0 = \frac{3}{2}(\frac{1}{q} - \frac{1}{p})$ and r_0 such that $r_0 \in (\frac{3}{2}(\frac{1}{2} - \frac{1}{p}), 1)$. Then, in view of Lemmas 6.1 and 6.2, for any $\mu \in (0, \min\{\frac{\rho_*-1}{2}, \lambda_1\})$, we recall the L^p - L^q estimates for the Stokes semigroup (see also Lemma 2.3 of [4]) to choose positive constants $\tilde{k}_1, \tilde{k}_2, \tilde{k}_3$ and \tilde{k}_4 such that

$$\|e^{-tA} \mathcal{P}\varphi\|_{L^p(\Omega)} \leq \tilde{k}_1 t^{-\gamma_0} \|\varphi\|_{L^q(\Omega)} \quad \text{for all } t \in (0, 2) \quad \text{and any } \varphi \in L^q(\Omega) \quad (6.31)$$

as well as

$$\|e^{-tA} \mathcal{P}\nabla \cdot \varphi\|_{L^p(\Omega)} \leq \tilde{k}_2 t^{-\frac{1}{2} - \frac{3}{2}(\frac{2}{p} - \frac{1}{p})} e^{-\mu t} \|\varphi\|_{L^{\frac{p}{2}}(\Omega)} \quad \text{for all } t \in (0, 2) \quad \text{and any } \varphi \in L^{\frac{p}{2}}(\Omega) \quad (6.32)$$

and

$$\begin{aligned}
\|e^{-tA}\mathcal{P}\varphi\|_{L^p(\Omega)} &\leq \|A^{r_0}e^{-tA}A^{-r_0}\mathcal{P}\varphi\|_{L^p(\Omega)} \\
&\leq \tilde{\kappa}_3 t^{-r_0} e^{-\mu t} \|A^{-r_0}\mathcal{P}\varphi\|_{L^p(\Omega)} \\
&\leq \tilde{\kappa}_4 t^{-r_0} e^{-\mu t} \|\varphi\|_{L^2(\Omega)} \quad \text{for all } t \in (0, 2) \quad \text{and any } \varphi \in L^2(\Omega),
\end{aligned} \tag{6.33}$$

where A is the Stokes operator as well as $\lambda_1 = \inf \operatorname{Re}\sigma(A) > 0$ and $\rho_{*,1}$ is the same as Lemma 6.5. Next, in view of (6.30) and using $p \geq 6$, $q \in (3, 6)$, we have

$$-2\gamma_0 - \frac{1}{2} - \frac{3}{2p} > -1 \quad \text{and} \quad -\frac{1}{2} - \frac{3}{2p} > -1.$$

Therefore,

$$2\tilde{\kappa}_2 \int_0^1 (1-\tau)^{-\frac{1}{2}-\frac{3}{2p}} \tau^{-2\gamma_0} d\tau < +\infty, \tag{6.34}$$

where $\tilde{\kappa}_2$ is the same as (6.32).

We finally invoke the Cauchy-Schwarz inequality to fix $\tilde{\kappa}_5 > 0$ such that

$$\|v \oplus w\|_{L^{\frac{p}{2}}(\Omega)} \leq \tilde{\kappa}_5 \|v\|_{L^p(\Omega)} \|w\|_{L^p(\Omega)} \quad \text{for all } v, w \in L^p(\Omega).$$

We now take any positive $\eta \leq \eta_0 := \min\left\{\frac{1}{3\tilde{\kappa}_2\tilde{\kappa}_5|\kappa|2^{\frac{1}{2}-\frac{3}{2p}-\gamma_0} \int_0^1 (1-\tau)^{-\frac{1}{2}-\frac{3}{2p}} \tau^{-2\gamma_0} d\tau}, 1\right\}$ and let

$$\eta_1 := \frac{\eta}{3\tilde{\kappa}_2} \quad \text{and} \quad \eta_2 := \frac{\eta}{3 \cdot 2^{\gamma_0} \tilde{\kappa}_4 \|\nabla\phi\|_{L^\infty(\Omega)} \int_0^{+\infty} s^{-r_0} e^{-\mu s} ds},$$

and note that as a consequence of Lemma 6.5 and Lemma 6.6, for any such η we can pick $T(\eta) > 2$ such that for all $t > T(\eta) - 2$

$$\int_t^{t+1} \|u_\varepsilon(\cdot, s)\|_{L^q(\Omega)}^2 ds \leq \eta_1^2 \tag{6.35}$$

and

$$\|n_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)} \leq \eta_2. \tag{6.36}$$

To see that these choices ensure that

$$\|u_\varepsilon(\cdot, t_0)\|_{L^p(\Omega)} \leq \eta, \quad \text{for all } t_0 > T(\eta), \tag{6.37}$$

we fix any such t_0 and then infer from (6.35) that there exists $t_* \in (t_0 - 2, t_0 - 1)$ such that

$$\|u_\varepsilon(\cdot, t_*)\|_{L^q(\Omega)} \leq \eta_1. \tag{6.38}$$

We now follow a standard reasoning to construct, independently of u_ε , another weak solution \tilde{u}_ε of the initial value problem associated with the Navier-Stokes system $\tilde{u}_{\varepsilon t} + A\tilde{u}_\varepsilon = -\mathcal{P}[\kappa(Y_\varepsilon u_\varepsilon \cdot \nabla)u_\varepsilon] + \mathcal{P}[n_\varepsilon \nabla \phi]$ in $\Omega \times (t_*, t_* + 2)$ with $\tilde{u}_\varepsilon(\cdot, t_*) = u_\varepsilon(\cdot, t_*)$ and some favorable additional properties, finally implying by a uniqueness argument that actually $u_\varepsilon = \tilde{u}_\varepsilon$ and that hence u_ε itself has these properties. To this end, for the above t_*, p, γ_0 and η , we let

$$X := \{v \in C^0((t_*, t_* + 2]; L_\sigma^p(\Omega)) \mid \|v\|_X := \sup_{t \in (t_*, t_* + 2)} (t - t_*)^{\gamma_0} \|v(\cdot, t)\|_{L^p(\Omega)} < \infty\}. \quad (6.39)$$

Then we consider the mapping Ψ defined by

$$\varphi(v)(\cdot, t) := e^{-(t-t_*)A}u_\varepsilon(\cdot, t_*) + \int_{t_*}^t e^{-(t-s)A}\mathcal{P}[-\kappa \nabla \cdot (Y_\varepsilon v(\cdot, \tau) \otimes v(\cdot, \tau) + n_\varepsilon(\cdot, \tau) \nabla \phi)] d\tau \quad t \in (t_*, t_* + 2],$$

for v belonging to the closed subset

$$S := \{v \in X \mid \|v\| \leq \eta\}$$

of X . Then for $v \in S$ we can use (6.31)–(6.33) as well as (6.36) and (6.38) to estimate

$$\begin{aligned} & \|\varphi(v)(\cdot, t)\|_{L^p(\Omega)} \\ & \leq \tilde{\kappa}_1(t - t_*)^{-\gamma_0} \|u_\varepsilon(\cdot, t_*)\|_{L^q(\Omega)} + \tilde{\kappa}_2|\kappa| \int_{t_*}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2}(\frac{2}{p} - \frac{1}{p})} e^{-\mu(t-\tau)} \|v(\cdot, \tau) \oplus v(\cdot, \tau)\|_{L^{\frac{p}{2}}(\Omega)} d\tau \\ & \quad + \int_{t_*}^t \|\mathcal{P}\{[n_\varepsilon(\cdot, \tau) - \bar{n}_0] \nabla \phi\}\|_{L^p(\Omega)} d\tau \\ & \leq \tilde{\kappa}_1(t - t_*)^{-\gamma_0} \|u_\varepsilon(\cdot, t_*)\|_{L^q(\Omega)} + \tilde{\kappa}_2|\kappa| \int_{t_*}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2}(\frac{2}{p} - \frac{1}{p})} e^{-\mu(t-\tau)} \|v(\cdot, \tau) \oplus v(\cdot, \tau)\|_{L^{\frac{p}{2}}(\Omega)} d\tau \\ & \quad + \tilde{\kappa}_4 \|\nabla \phi\|_{L^\infty(\Omega)} \int_{t_*}^t (t - \tau)^{-r_0} e^{-\mu(t-\tau)} \|n_\varepsilon(\cdot, \tau) - \bar{n}_0\|_{L^2(\Omega)} d\tau \\ & \leq \tilde{\kappa}_1(t - t_*)^{-\gamma_0} \|u_\varepsilon(\cdot, t_*)\|_{L^q(\Omega)} + \tilde{\kappa}_2\tilde{\kappa}_5|\kappa| \int_{t_*}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2p}(\frac{2}{p} - \frac{1}{p})} e^{-\mu(t-\tau)} \|v(\cdot, \tau)\|_{L^p(\Omega)}^2 d\tau \\ & \quad + \tilde{\kappa}_4 \|\nabla \phi\|_{L^\infty(\Omega)} \eta_2 \int_{t_*}^{t_*+2} (t - \tau)^{-r_0} e^{-\mu(t-\tau)} d\tau \\ & \leq \tilde{\kappa}_1(t - t_*)^{-\gamma_0} \eta_1 + \tilde{\kappa}_2\tilde{\kappa}_5|\kappa|\eta^2 \int_{t_*}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2p}\tau} \tau^{-2\gamma_0} d\tau \\ & \quad + \tilde{\kappa}_4 \|\nabla \phi\|_{L^\infty(\Omega)} \eta_2 \int_0^{+\infty} s^{-r_0} e^{-\mu s} ds \quad \text{for all } t \in (t_*, t_* + 2), \end{aligned} \quad (6.40)$$

where μ and r_0 are the same as (6.33). Therefore, in light of the definition of γ_0 , (6.34),

(6.38) as well as (6.39) and (6.40), we see that for every $t \in (t_*, t_* + 2)$ and every $v \in S$

$$\begin{aligned}
& (t - t_*)^{\gamma_0} \|\varphi(v)(\cdot, t)\|_{L^p(\Omega)} \\
& \leq \tilde{\kappa}_1 \eta_1 + \tilde{\kappa}_2 \tilde{\kappa}_5 |\kappa| \eta^2 (t - t_*)^{\gamma_0 - \frac{1}{2} - \frac{3}{2p} - 2\gamma_0 + 1} \int_0^1 (1 - \tau)^{-\frac{1}{2} - \frac{3}{2p}} \tau^{-2\gamma_0} d\tau \\
& \quad + \tilde{\kappa}_4 \|\nabla \phi\|_{L^\infty(\Omega)} \eta_2 (t - t_*)^{\gamma_0} \int_0^{+\infty} s^{-r_0} e^{-\mu s} ds \\
& \leq \tilde{\kappa}_1 \eta_1 + \tilde{\kappa}_2 \tilde{\kappa}_5 |\kappa| \eta^2 2^{\frac{1}{2} - \frac{3}{2p} - \gamma_0} \int_0^1 (1 - \tau)^{-\frac{1}{2} - \frac{3}{2p}} \tau^{-2\gamma_0} d\tau \\
& \quad + \tilde{\kappa}_4 \|\nabla \phi\|_{L^\infty(\Omega)} \eta_2 2^{\gamma_0} \int_0^{+\infty} s^{-r_0} e^{-\mu s} ds \\
& \leq \frac{\eta}{3} + \frac{\eta}{3} + \frac{\eta}{3} \\
& < \eta,
\end{aligned} \tag{6.41}$$

from which it readily follows that $\varphi(S) \subset S$. Likewise, for $v \in S$ and $w \in S$ we can use Lemma 6.2 to find that

$$\begin{aligned}
& \|(\varphi(v) - \varphi(w))(\cdot, t)\|_{L^p(\Omega)} \\
& \leq \tilde{\kappa}_2 |\kappa| \int_{t_*}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2p}} \|v(\cdot, \tau) \oplus v(\cdot, \tau) - w(\cdot, \tau) \oplus w(\cdot, \tau)\|_{L^{\frac{p}{2}}(\Omega)} d\tau \\
& = \tilde{\kappa}_2 |\kappa| \int_{t_*}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2p}} \|v(\cdot, \tau) \oplus (v(\cdot, \tau) - w(\cdot, \tau)) + (v(\cdot, \tau) - w(\cdot, \tau)) \oplus w(\cdot, \tau)\|_{L^{\frac{p}{2}}(\Omega)} d\tau \\
& \leq \tilde{\kappa}_2 \tilde{\kappa}_5 |\kappa| \int_{t_*}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2p}} (\|v(\cdot, \tau)\|_{L^p} + \|w(\cdot, \tau)\|_{L^p}) \|v(\cdot, \tau) - w(\cdot, \tau)\|_{L^p(\Omega)} d\tau \\
& \leq \tilde{\kappa}_2 \tilde{\kappa}_5 |\kappa| \int_{t_*}^t \tau^{-\frac{1}{2} - \frac{3}{2p}} 2\eta \tau^{-\gamma_0} \cdot \tau^{-\gamma_0} \|v - w\|_X d\tau \quad \text{for all } t \in (t_*, t_* + 2),
\end{aligned} \tag{6.42}$$

which implies that

$$\begin{aligned}
& (t - t_*)^{\gamma_0} \|(\varphi(v) - \varphi(w))(\cdot, t)\|_{L^p(\Omega)} \\
& \leq 2^{\frac{3}{2} - \frac{3}{2p} - \gamma_0} \tilde{\kappa}_2 \tilde{\kappa}_5 |\kappa| \int_0^1 (1 - \tau)^{-\frac{1}{2} - \frac{3}{2p}} \tau^{-2\gamma_0} d\tau \eta \|v - w\|_X \quad \text{for all } t \in (t_*, t_* + 2).
\end{aligned} \tag{6.43}$$

Recalling

$$\eta \leq \frac{1}{3\tilde{\kappa}_2 \tilde{\kappa}_5 |\kappa| 2^{\frac{1}{2} - \frac{3}{2p} - \gamma_0} \int_0^1 (1 - \tau)^{-\frac{1}{2} - \frac{3}{2p}} \tau^{-2\gamma_0} d\tau},$$

thus,

$$2^{\frac{3}{2} - \frac{3}{2p} - \gamma_0} \tilde{\kappa}_2 \tilde{\kappa}_5 |\kappa| \int_0^1 (1 - \tau)^{-\frac{1}{2} - \frac{3}{2p}} \tau^{-2\gamma_0} d\tau \eta \leq \frac{2}{3} < 1,$$

so that, (6.43) implies that φ acts as a contraction on S and hence possesses a unique fixed point \tilde{u}_ε . In view of a standard arguments (see e.g. [31]), it follows that \tilde{u}_ε in fact is a

weak solution of the Navier-Stokes subsystem of (2.1) in $\Omega \times (t_*, t_* + 2)$ subject to the initial condition $\tilde{u}_\varepsilon(\cdot, t_*) = u_\varepsilon(\cdot, t_*)$. Next, using the definition of S and X , we also derive that

$$\int_{t_*}^{t_*+2} \|\tilde{u}_\varepsilon(\cdot, t)\|_{L^p(\Omega)}^{q_0} dt \leq \eta, \quad (6.44)$$

where

$$q_0 = \frac{2p}{p-3}$$

satisfy

$$\frac{2}{q_0} + \frac{3}{p} = 1 \quad \text{and} \quad q_0 \gamma_0 = \frac{2p}{p-3} \times \frac{3}{2} \left(\frac{1}{q} - \frac{1}{p} \right) < 1 \quad \text{by using } q > 3. \quad (6.45)$$

Collecting (6.44) and (6.45), a well-known uniqueness property of the Navier-Stokes equations (see e.g. Theorem V.2.5.1 of [31]) entails that \tilde{u}_ε must coincide with u_ε in $\Omega \times (t_*, t_* + 2)$.

In particular, since $t_0 \in (t_* + 1, t_* + 2)$, (6.44) thus implies that

$$\|u_\varepsilon(\cdot, t_0)\|_{L^p(\Omega)} \leq \|\tilde{u}_\varepsilon(\cdot, t_0)\|_{L^p(\Omega)} \leq \eta(t_0 - t_*)^{-\gamma_0} \leq \eta,$$

and thereby establishes (6.37), which together with the fact that $\eta \in (0, \eta_0]$ was arbitrary yields to (6.29). \square

With the help of Lemma 6.5 as well as Lemma 2.3 and Lemma 6.7, we can achieve the boundedness of ∇c_ε in $L^\infty((T, \infty); L^4(\Omega))$ and n_ε in $L^\infty(\Omega \times (T, \infty))$ for some large $T > 2$.

Lemma 6.8. *Under the assumptions of Theorem 1.2, then one can find that $T_{*,4} > 2$, $\rho_{*,4} \in (0, 1)$ and $C_{*,4} > 0$ such that for each $\varepsilon \in (0, 1)$,*

$$\int_{\Omega} |\nabla c_\varepsilon(x, t)|^2 < C_{*,4} e^{-\rho_{*,4} t} \quad \text{for all } t > T_{*,4} \quad (6.46)$$

and

$$\int_{\Omega} |\nabla c_\varepsilon(x, t)|^4 < C_{*,4} e^{-\rho_{*,4} t} \quad \text{for all } t > T_{*,4}. \quad (6.47)$$

Proof. To show this Lemma, it will be convenient to introduce the following notation:

$$w_\varepsilon = c_\varepsilon - \bar{n}_0.$$

Accordingly, in light of the second equation of (2.1), by a straightforward computation, we see w_ε has the following property:

$$w_{\varepsilon t} - \Delta w_\varepsilon + u_\varepsilon \cdot \nabla w_\varepsilon + w_\varepsilon = n_\varepsilon - \bar{n}_0, \quad x \in \Omega, \quad t > 0.$$

For any $\eta \in (0, 1)$, let us apply Corollary 6.5, Lemma 6.7 to fix $T_{1,*} > 2$ and constants $\rho_{1,*} \in (0, 1)$, $C_{1,*} > 0$ and $C_{2,*} > 0$ such that for any $t > T_{1,*}$

$$\|u_\varepsilon\|_{L^\infty((t,+\infty);L^p(\Omega))} < \eta \quad \text{for all } p > 1 \quad (6.48)$$

as well as

$$\|n_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^\infty((t,+\infty);L^2(\Omega))} < C_{1,*}e^{-\rho_{1,*}t} \quad (6.49)$$

and

$$\|w_\varepsilon\|_{L^\infty((t,+\infty);L^2(\Omega))} < C_{2,*}e^{-\rho_{1,*}t}. \quad (6.50)$$

On the other hand, by Lemma 6.5, we also have

$$\int_{T_{1,*}}^{\infty} \int_{\Omega} |\nabla w_\varepsilon|^2 \leq C_{3,*} \quad \text{for all } \varepsilon \in (0, 1) \quad (6.51)$$

and some positive constant $C_{3,*}$. Then (6.51) in particular implies that we can find $\tilde{t}_* \in (T_{1,*}, \infty)$ such that

$$\|\nabla w_\varepsilon(\cdot, \tilde{t}_*)\|_{L^2(\Omega)} \leq C_{3,*}. \quad (6.52)$$

Next, testing the second equation in (2.1) against $-\Delta w_\varepsilon$ and employing the Young inequality yields

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\nabla w_\varepsilon(\cdot, t)\|_{L^2(\Omega)}^2 &= \int_{\Omega} -\Delta w_\varepsilon (\Delta w_\varepsilon - w_\varepsilon + n_\varepsilon - \bar{n}_0 - u_\varepsilon \cdot \nabla w_\varepsilon) \\ &\leq -\frac{1}{4} \int_{\Omega} |\Delta w_\varepsilon|^2 - \int_{\Omega} |\nabla w_\varepsilon|^2 + \int_{\Omega} (n_\varepsilon - \bar{n}_0)^2 + \int_{\Omega} |u_\varepsilon \cdot \nabla w_\varepsilon| |\Delta w_\varepsilon| \end{aligned} \quad (6.53)$$

for all $t > t_*$. Next, one needs to estimate the last term on the right-hand side of (6.53).

Indeed, in view of (6.48) and (6.50), we derive from the Hölder inequality, the Gagliardo–Nirenberg inequality, and the Young inequality that there exist positive constants $C_{4,*}$ as well as $C_{5,*}$ and $C_{6,*}$ such that

$$\begin{aligned} \int_{\Omega} |u_\varepsilon \cdot \nabla w_\varepsilon| |\Delta w_\varepsilon| &\leq \|u_\varepsilon\|_{L^6(\Omega)} \|\nabla w_\varepsilon\|_{L^3(\Omega)} \|\Delta w_\varepsilon\|_{L^2(\Omega)} \\ &\leq C_{4,*} \|\nabla w_\varepsilon\|_{L^3(\Omega)} \|\Delta w_\varepsilon\|_{L^2(\Omega)} \\ &\leq C_{5,*} \left(\|\Delta w_\varepsilon\|_{L^2(\Omega)}^{\frac{3}{4}} \|w_\varepsilon\|_{L^2(\Omega)}^{\frac{1}{4}} + \|w_\varepsilon\|_{L^2(\Omega)}^2 \right) \|\Delta w_\varepsilon\|_{L^2(\Omega)} \\ &\leq C_{5,*} \max\{C_{2,*}^2, C_{2,*}^{\frac{1}{4}}\} e^{-\frac{\rho_{1,*}}{4}t} (\|\Delta w_\varepsilon\|_{L^2(\Omega)}^{\frac{7}{4}} + \|\Delta w_\varepsilon\|_{L^2(\Omega)}) \\ &\leq \frac{1}{4} \|\Delta w_\varepsilon\|_{L^2(\Omega)}^2 + C_{6,*} e^{-\frac{\rho_{1,*}}{2}t} \quad \text{for all } t > \tilde{t}_*, \end{aligned} \quad (6.54)$$

where we have used the fact that $t > \tilde{t}_* \geq T_{1,*} > 2$. Inserting (6.54) into (6.53) and using (6.76), we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\nabla w_\varepsilon(\cdot, t)\|_{L^2(\Omega)}^2 &\leq -\frac{1}{2} \int_{\Omega} |\Delta w_\varepsilon|^2 - \int_{\Omega} |\nabla w_\varepsilon|^2 + C_{6,*} e^{-\frac{\rho_{1,*}}{2}t} + \int_{\Omega} (n_\varepsilon - \bar{n}_0)^2 \\ &\leq -\frac{1}{2} \int_{\Omega} |\Delta w_\varepsilon|^2 - \int_{\Omega} |\nabla w_\varepsilon|^2 + C_{6,*} e^{-\frac{\rho_{1,*}}{2}t} + C_{1,*} e^{-\rho_{1,*}t} \\ &\leq -\frac{1}{2} \int_{\Omega} |\Delta w_\varepsilon|^2 - \int_{\Omega} |\nabla w_\varepsilon|^2 + \max\{C_{6,*}, C_{1,*}\} e^{-\frac{\rho_{1,*}}{2}t} \quad \text{for all } t > \tilde{t}_*. \end{aligned}$$

Integrating the above inequality in time and using (6.66), we derive that there exist positive constants $\rho_{2,*}$ and $C_{7,*}$ such that

$$\int_{\Omega} |\nabla w_\varepsilon(x, t)|^2 dx \leq C_{7,*} e^{-\rho_{2,*}t} \quad \text{for all } t \geq \tilde{t}_*. \quad (6.55)$$

However, from (6.55) and (6.50), with the help of the Sobolev imbedding theorem, it follows that there exist positive constants $C_{8,*}$ and $\rho_{3,*}$ such that

$$\|w_\varepsilon(\cdot, t)\|_{L^5(\Omega)} \leq C_{8,*} e^{-\rho_{3,*}t} \quad \text{for all } t \geq \tilde{t}_*. \quad (6.56)$$

Now, applying the variation-of-constants formula for w_ε and applying $\nabla \cdot u_\varepsilon = 0$ in $x \in \Omega, t > 0$, we obtain that for any $t > 2\tilde{t}_*$,

$$w_\varepsilon(\cdot, t) = e^{(t-\tilde{t}_*)(\Delta-1)} w_\varepsilon(\cdot, \tilde{t}_*) + \int_{\tilde{t}_*}^t e^{(t-s)(\Delta-1)} (n_\varepsilon(\cdot, s) - \bar{n}_0 + \nabla \cdot (u_\varepsilon(\cdot, s) w_\varepsilon(\cdot, s))) ds, \quad (6.57)$$

so that, we have

$$\begin{aligned} &\|\nabla w_\varepsilon(\cdot, t)\|_{L^4(\Omega)} \\ &\leq \|\nabla e^{(t-\tilde{t}_*)(\Delta-1)} w_\varepsilon(\cdot, \tilde{t}_*)\|_{L^4(\Omega)} + \int_{\tilde{t}_*}^t \|\nabla e^{(t-s)(\Delta-1)} (n_\varepsilon(\cdot, s) - \bar{n}_0)\|_{L^4(\Omega)} ds \\ &\quad + \int_{\tilde{t}_*}^t \|\nabla e^{(t-s)(\Delta-1)} \nabla \cdot (u_\varepsilon(\cdot, s) w_\varepsilon(\cdot, s))\|_{L^4(\Omega)} ds \quad \text{for all } t > 2\tilde{t}_*. \end{aligned} \quad (6.58)$$

To address the right-hand side of (6.58), in view of (6.56), we first use Lemma 6.2 to get for some positive constants $\mu_{1,*} > 0, \rho_{4,*} > 0$ and $C_{9,*} > 0$ such that

$$\begin{aligned} \|\nabla e^{(t-\tilde{t}_*)(\Delta-1)} w_\varepsilon(\cdot, \tilde{t}_*)\|_{L^4(\Omega)} &\leq \kappa_3 (t - \tilde{t}_*)^{-\frac{1}{2} - \frac{3}{2}(\frac{1}{5} - \frac{1}{4})} e^{-\mu_{1,*}t} \|w_\varepsilon(\cdot, t)\|_{L^5(\Omega)} \\ &\leq \kappa_3 \|w_\varepsilon(\cdot, t)\|_{L^5(\Omega)} \\ &\leq C_{9,*} e^{-\rho_{4,*}t} \quad \text{for all } t > 2\tilde{t}_* \end{aligned} \quad (6.59)$$

by using $\tilde{t}_* > 1$, where κ_3 is the same as Lemma 6.2. Since

$$-\frac{1}{2} - \frac{3}{2} \left(\frac{1}{2} - \frac{1}{4} \right) > -1,$$

together with this, by using Lemma 6.2 and (6.76), the second term on the right-hand side of (6.58) is estimated as

$$\begin{aligned} & \int_{\tilde{t}_*}^t \|\nabla e^{(t-s)(\Delta-1)}(n_\varepsilon(\cdot, s) - \bar{n}_0)\|_{L^4(\Omega)} ds \\ & \leq C_{10,*} \int_{\tilde{t}_*}^t [1 + (t-s)^{-\frac{1}{2} - \frac{3}{2}(\frac{1}{2} - \frac{1}{4})}] e^{-\lambda_{1,*}(t-s)} \|n_\varepsilon(\cdot, s) - \bar{n}_0\|_{L^2(\Omega)} ds \\ & \leq C_{10,*} \int_{\tilde{t}_*}^t [1 + (t-s)^{-\frac{1}{2} - \frac{3}{2}(\frac{1}{2} - \frac{1}{4})}] e^{-\lambda_{1,*}(t-s)} C_{1,*} e^{-\rho_{1,*} s} ds \\ & \leq C_{11,*} e^{-\rho_{5,*} t} \quad \text{for all } t > 2\tilde{t}_* \end{aligned} \tag{6.60}$$

and some positive constants $C_{10,*}$, $\rho_{5,*}$, $\lambda_{1,*}$ and $C_{11,*}$. Here $C_{1,*}$ and $\rho_{1,*}$ are the same as (6.76). Next, we will address the third term on the right-hand side of (6.58). To this end, we choose $\iota = \frac{23}{48}$ and $\tilde{\kappa} = \frac{1}{96}$, then $\frac{1}{2} + \iota + \tilde{\kappa} < 1$. Therefore, in view of the Hölder inequality, we derive from Lemma 6.2, (6.48) and (6.56) that there exist positive constants $\lambda_{2,*}$, $\rho_{6,*}$, $C_{12,*}$, $C_{13,*}$, $C_{14,*}$, $C_{15,*}$ and $C_{16,*}$ such that

$$\begin{aligned} & \int_{\tilde{t}_*}^t \|\nabla e^{(t-s)(\Delta-1)} \nabla \cdot (u_\varepsilon(\cdot, s) w_\varepsilon(\cdot, s))\|_{L^4(\Omega)} ds \\ & \leq C_{12,*} \int_{\tilde{t}_*}^t \|e^{(t-s)(\Delta-1)} \nabla \cdot (u_\varepsilon(\cdot, s) w_\varepsilon(\cdot, s))\|_{W^{1,4}(\Omega)} ds \\ & \leq C_{13,*} \int_{\tilde{t}_*}^t \|(-\Delta + 1)^t e^{(t-s)(\Delta-1)} \nabla \cdot (u_\varepsilon(\cdot, s) w_\varepsilon(\cdot, s))\|_{L^{\frac{9}{2}}(\Omega)} ds \\ & \leq C_{14,*} \int_{\tilde{t}_*}^t (t-s)^{-\iota - \frac{1}{2} - \tilde{\kappa}} e^{-\lambda_{2,*}(t-s)} \|u_\varepsilon(\cdot, s) w_\varepsilon(\cdot, s)\|_{L^{\frac{9}{2}}(\Omega)} ds \\ & \leq C_{15,*} \int_{\tilde{t}_*}^t (t-s)^{-\iota - \frac{1}{2} - \tilde{\kappa}} e^{-\lambda_{2,*}(t-s)} \|u_\varepsilon(\cdot, s)\|_{L^{45}(\Omega)} \|w_\varepsilon(\cdot, s)\|_{L^5(\Omega)} ds \\ & \leq C_{16,*} e^{-\rho_{6,*} t} \quad \text{for all } t > 2\tilde{t}_*. \end{aligned} \tag{6.61}$$

Here we have used the fact that

$$\int_{\tilde{t}_*}^t (t-s)^{-\iota - \frac{1}{2} - \tilde{\kappa}} e^{-\lambda_{2,*}(t-s)} ds \leq \int_0^\infty \sigma^{-\iota - \frac{1}{2} - \tilde{\kappa}} e^{-\lambda_{2,*} \sigma} d\sigma < +\infty.$$

Finally, collecting (6.58)–(6.61), we can obtain that there exists a positive constant $C_{17,*}$ and $\rho_{7,*}$ such that

$$\int_{\Omega} |\nabla c_\varepsilon(x, t)|^4 dx \leq C_{17,*} e^{-\rho_{7,*} t} \quad \text{for all } t > 2\tilde{t}_*. \tag{6.62}$$

This completes the proof of Lemma 6.8. \square

Lemma 6.9. *Under the assumptions of Theorem 1.2, one can find that $T_{*,5} > 2$ and $C_{*,5} > 0$ with the property that for all $\varepsilon \in (0, 1)$ we have*

$$\|n_\varepsilon(\cdot, t)\|_{L^{\frac{10}{3}}(\Omega)} \leq C_{*,5} \quad \text{for all } t \geq T_{*,5}. \quad (6.63)$$

Proof. Firstly, let us apply Lemma 6.8 to fix $T_{1,**} > 2$ as well as constants $\rho_{1,**} \in (0, 1)$ and $C_{1,**} > 0$ such that for any $t \geq T_{1,**}$

$$\|\nabla c_\varepsilon\|_{L^\infty((t, +\infty); L^4(\Omega))} < C_{1,**} e^{-\rho_{1,**} t}. \quad (6.64)$$

On the other hand, by Proposition 6.1, we also derive that

$$\int_{T_{1,**}}^{\infty} \int_{\Omega} n_\varepsilon^{\frac{10}{3}} \leq C_{2,**} \quad \text{for all } \varepsilon \in (0, 1) \quad (6.65)$$

and some positive constant $C_{2,**}$. Then (6.65) in particular implies that we can find $t_{**} \in (T_{1,**}, \infty)$ such that

$$\|n_\varepsilon(\cdot, t_{**})\|_{L^{\frac{10}{3}}(\Omega)} \leq C_{2,**}. \quad (6.66)$$

Let $p = \frac{10}{3}$. Testing the first equation in (2.1) by n_ε^{p-1} , using the fact $\nabla \cdot u_\varepsilon = 0$, and applying the Young inequality and the Hölder inequality, we have

$$\begin{aligned} & \frac{1}{p} \frac{d}{dt} \int_{\Omega} n_\varepsilon^p + (p-1) \int_{\Omega} n_\varepsilon^{p-2} |\nabla c_\varepsilon|^2 \\ &= - \int_{\Omega} n_\varepsilon^{p-1} \nabla \cdot (n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon) \\ &= (p-1) \int_{\Omega} n_\varepsilon^{p-2} F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon \cdot \nabla n_\varepsilon \\ &\leq \frac{p-1}{2} \int_{\Omega} n_\varepsilon^{p-2} |\nabla n_\varepsilon|^2 + \frac{p-1}{2} \int_{\Omega} n_\varepsilon^p (1+n_\varepsilon)^{-2\alpha} |\nabla c_\varepsilon|^2 \\ &\leq \frac{p-1}{2} \int_{\Omega} n_\varepsilon^{p-2} |\nabla n_\varepsilon|^2 + \frac{p-1}{2} \left(\int_{\Omega} n_\varepsilon^{2p} (1+n_\varepsilon)^{-4\alpha} \right)^{\frac{1}{2}} \left(\int_{\Omega} |\nabla c_\varepsilon|^4 \right)^{\frac{1}{2}} \quad \text{for all } t > t_{**} \end{aligned} \quad (6.67)$$

by using (1.6) and (2.2). Without loss of generality, we may assume that $\alpha < \frac{17}{12}$, since $\alpha \geq \frac{17}{12}$ can be proved similarly and easily. Then the Gagliardo–Nirenberg inequality as well

as (6.64) and (2.11) enables us to see that

$$\begin{aligned}
& \frac{p-1}{2} \left(\int_{\Omega} n_{\varepsilon}^{2p} (1+n_{\varepsilon})^{-4\alpha} \right)^{\frac{1}{2}} \left(\int_{\Omega} |\nabla C_{\varepsilon}|^4 \right)^{\frac{1}{2}} \\
& \leq \frac{p-1}{2} \left(\int_{\Omega} n_{\varepsilon}^{2p-4\alpha} \right)^{\frac{1}{2}} C_{1,**}^2 e^{-2\rho_{1,**}t} \\
& \leq C_{1,**}^2 \frac{p-1}{2} \|n_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{4p-8\alpha}{p}}(\Omega)}^{\frac{2p-4\alpha}{p}} \\
& \leq C_{3,**} (\|\nabla n_{\varepsilon}^{\frac{p}{2}}\|_{L^2(\Omega)}^{\frac{6p-12\alpha-3}{3p-1}} \|n_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{2}{p}}(\Omega)}^{\frac{2(p-1)}{p} - \frac{6p-12\alpha-3}{3p-1}} + \|n_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{2}{p}}(\Omega)}^{\frac{2(p-1)}{p}}) \\
& \leq C_{4,**} (\|\nabla n_{\varepsilon}^{\frac{p}{2}}\|_{L^2(\Omega)}^{\frac{6p-12\alpha-3}{3p-1}} + 1) \quad \text{for all } t > t_{**}
\end{aligned}$$

and some positive constants $C_{3,**}$ and $C_{4,**}$, where $C_{1,**}$ and $\rho_{1,**}$ are the same as (6.64)

Here we use the Young inequality to obtain that there is $C_{5,**} > 0$ such that

$$\frac{1}{p} \frac{d}{dt} \int_{\Omega} n_{\varepsilon}^p + \frac{(p-1)}{2} \int_{\Omega} n_{\varepsilon}^{p-2} |\nabla n_{\varepsilon}|^2 \leq C_{5,**} \quad \text{for all } t > t_{**}. \quad (6.68)$$

Next, in view of (2.11), we derive from the Gagliardo–Nirenberg inequality that for some positive constants $C_{6,**}$ and $C_{7,**}$ such that

$$\begin{aligned}
& \|n_{\varepsilon}\|_{L^p(\Omega)}^p \\
& = \|n_{\varepsilon}^{\frac{p}{2}}\|_{L^2(\Omega)}^2 \\
& \leq C_{6,**} (\|\nabla n_{\varepsilon}^{\frac{p}{2}}\|_{L^2(\Omega)}^{\frac{6(p-1)}{3p-1}} \|n_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{2}{p}}(\Omega)}^{2 - \frac{6(p-1)}{3p-1}} + \|n_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{2}{p}}(\Omega)}^2) \\
& \leq C_{7,**} (\|\nabla n_{\varepsilon}^{\frac{p}{2}}\|_{L^2(\Omega)}^{\frac{6(p-1)}{3p-1}} + 1) \quad \text{for all } t > t_{**},
\end{aligned}$$

which together with the Young inequality implies that

$$\begin{aligned}
\|n_{\varepsilon}\|_{L^p(\Omega)}^p & \leq \|\nabla n_{\varepsilon}^{\frac{p}{2}}\|_{L^2(\Omega)}^2 + C_{8,**} \\
& = \frac{p^2}{4} \int_{\Omega} n_{\varepsilon}^{p-2} |\nabla n_{\varepsilon}|^2 + C_{8,**} \quad \text{for all } t > t_{**}.
\end{aligned} \quad (6.69)$$

with some positive constant $C_{8,**}$. Inserting (6.69) into (6.68), there exist positive constants $C_{9,**}$ and $C_{10,**}$ such that

$$\frac{d}{dt} \int_{\Omega} n_{\varepsilon}^p + C_{9,**} \int_{\Omega} n_{\varepsilon}^p \leq C_{10,**} \quad \text{for all } t > t_{**} \quad (6.70)$$

by some basic calculation. Finally, employing the Gronwall lemma to (6.70), we can obtain (6.74) by using (6.66). \square

Lemma 6.10. *Under the assumptions of Theorem 1.2, one can find that $T_{*,6} > 2$ and $C_{*,6} > 0$ such that for all $t \geq T_{*,6}$ and $\varepsilon \in (0, 1)$,*

$$\|c_\varepsilon(\cdot, s)\|_{W^{1,\infty}(\Omega)} \leq C_{*,6}. \quad (6.71)$$

Proof. Firstly, applying Lemma 6.5 as well as Lemmas 6.7–6.9, one can derive that for some positive constants $T_{1,***} > 2$ and $C_{1,***}$ such that

$$\|c_\varepsilon(\cdot, s)\|_{W^{1,4}(\Omega)} + \|c_\varepsilon(\cdot, s)\|_{L^\infty(\Omega)} + \|u_\varepsilon(\cdot, s)\|_{L^q(\Omega)} + \|n_\varepsilon(\cdot, s)\|_{L^{\frac{10}{3}}(\Omega)} \leq C_{1,***} \quad \text{for all } s \geq T_{1,***} \quad (6.72)$$

and $q > 2$. Here we have used the Sobolev embedding $W^{1,4}(\Omega) \hookrightarrow L^\infty(\Omega)$. To prove the boundedness of $\|\nabla c_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)}$, we rewrite the variation-of-constants formula for c_ε in the form

$$c_\varepsilon(\cdot, t) = e^{(t-T_{1,***})(\Delta-1)}c_\varepsilon(\cdot, T_{1,***}) + \int_{T_{1,***}}^t e^{(t-s)(\Delta-1)}[n_\varepsilon(\cdot, s) - u_\varepsilon(\cdot, s) \cdot \nabla c_\varepsilon(\cdot, s)]ds \quad \text{for all } t > 2T_{1,***}.$$

Now, we choose $\theta \in (\frac{1}{2} + \frac{3}{2} \times \frac{3}{10}, 1)$, then the domain of the fractional power $D((-\Delta + 1)^\theta) \hookrightarrow W^{1,\infty}(\Omega)$ (see [17]). Hence, in view of L^p – L^q estimates associated with the heat semigroup, (6.72), we derive that there exist positive constants $\lambda_{1,***}$, $C_{2,***}$, $C_{3,***}$, $C_{4,***}$, and $C_{5,***}$ such that

$$\begin{aligned} & \|c_\varepsilon(\cdot, t)\|_{W^{1,\infty}(\Omega)} \\ & \leq C_{2,***} \|(-\Delta + 1)^\theta c_\varepsilon(\cdot, t)\|_{L^{\frac{10}{3}}(\Omega)} \\ & \leq C_{3,***} (t - T_{1,***})^{-\theta} e^{-\lambda_{1,***}(t-T_{1,***})} \|c_\varepsilon(\cdot, T_{1,***})\|_{L^{\frac{10}{3}}(\Omega)} \\ & \quad + C_{3,***} \int_{T_{1,***}}^t (t-s)^{-\theta} e^{-\lambda_{1,***}(t-s)} \|n_\varepsilon(\cdot, s) - u_\varepsilon(\cdot, s) \cdot \nabla c_\varepsilon(\cdot, s)\|_{L^{\frac{10}{3}}(\Omega)} ds \\ & \leq C_{4,***} + C_{4,***} \int_{T_{1,***}}^t (t-s)^{-\theta} e^{-\lambda_{1,***}(t-s)} [\|n_\varepsilon(\cdot, s)\|_{L^{\frac{10}{3}}(\Omega)} + \|u_\varepsilon(\cdot, s)\|_{L^{20}(\Omega)} \|\nabla c_\varepsilon(\cdot, s)\|_{L^4(\Omega)}] ds \\ & \leq C_{5,***} \quad \text{for all } t > 2T_{1,***}. \end{aligned} \quad (6.73)$$

Here we have used Hölder's inequality as well as

$$\int_{T_{1,***}}^t (t-s)^{-\theta} e^{-\lambda_{1,***}(t-s)} \leq \int_0^\infty \sigma^{-\theta} e^{-\lambda_{1,***}\sigma} d\sigma < +\infty$$

and $t - T_{1,***} > T_{1,***} > 1$. Finally, collecting (6.72) and (6.73) imply (6.71) and thereby completes the proof of Lemma 6.10. \square

Lemma 6.11. *Under the assumptions of Theorem 1.2, one can find that $T_{*,7} > 2$ and $C_{*,7} > 0$ with the property that for all $\varepsilon \in (0, 1)$ we have*

$$\|n_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq C_{*,7} \quad \text{for all } t \geq T_{*,7}. \quad (6.74)$$

Proof. In view of Lemma 6.5, Lemma 6.8 and Lemma 6.10, we can pick $T_{1,****} > 2$ as well as $\rho_{1,****} \in (0, 1)$ and $C_{1,****} > 0$ such that for any $t \geq T_{1,****}$

$$\|u_\varepsilon\|_{L^\infty((t, +\infty); L^p(\Omega))} < C_{1,****} \quad \text{for all } p > 1 \quad (6.75)$$

as well as

$$\|n_\varepsilon(\cdot, t) - \bar{n}_0\|_{L^\infty((t, +\infty); L^2(\Omega))} < C_{1,****} \quad (6.76)$$

and

$$\|\nabla c_\varepsilon\|_{L^\infty((t, +\infty); L^\infty(\Omega))} < C_{1,****}. \quad (6.77)$$

Next, let $\tilde{h}_\varepsilon := F_\varepsilon(n_\varepsilon)S_\varepsilon(x, n_\varepsilon, c_\varepsilon)\nabla c_\varepsilon + u_\varepsilon$. In view of (1.6), (2.2) as well as (6.75) and (6.77), there exists $C_{2,****} > 0$ such that

$$\|\tilde{h}_\varepsilon(\cdot, t)\|_{L^{\frac{190}{3}}(\Omega)} \leq C_{2,****} \quad \text{for all } t \geq T_{1,****}. \quad (6.78)$$

Hence, due to the fact that $\nabla \cdot u_\varepsilon = 0$, again, by means of an associate variation-of-constants formula for n_ε , we can derive

$$n_\varepsilon(\cdot, t) = e^{(t-T_{1,****})\Delta} n_\varepsilon(\cdot, T_{1,****}) - \int_{T_{1,****}}^t e^{(t-s)\Delta} \nabla \cdot (n_\varepsilon(\cdot, s) \tilde{h}_\varepsilon(\cdot, s)) ds, \quad t > 2T_{1,****}. \quad (6.79)$$

Since, $T_{1,****} > 1$, then in light of the L^p - L^q estimates for the Neumann heat semigroup and Lemma 6.5, we conclude that for some positive constants $C_{3,****}$ as well as $C_{4,****}$ and $\tilde{\rho}_{2,****}$ such that for all $t > 2T_{1,****}$

$$\|e^{(t-T_{1,****})\Delta} n_\varepsilon(\cdot, T_{1,****})\|_{L^\infty(\Omega)} \leq C_{3,****} (t - T_{1,****})^{-\frac{3}{4}} e^{-\tilde{\rho}_{2,****} t} \|n_\varepsilon(\cdot, T_{1,****})\|_{L^2(\Omega)} \leq C_{4,****}. \quad (6.80)$$

In the following, we will estimate the last term on the right-hand side of (6.79). In fact, applying the smoothing properties of the Stokes semigroup as well as (3.1) and the Hölder

inequality to find $C_{5,****} > 0$, $C_{6,****} > 0$ and $\tilde{\lambda}_{1,****}$ such that

$$\begin{aligned}
& \int_{T_{1,****}}^t \|e^{(t-s)\Delta} \nabla \cdot (n_\varepsilon(\cdot, s) \tilde{h}_\varepsilon(\cdot, s))\|_{L^\infty(\Omega)} ds \\
& \leq C_{5,****} \int_{T_{1,****}}^t (t-s)^{-\frac{1}{2}-\frac{9}{19}} e^{-\tilde{\lambda}_{1,****}(t-s)} \|n_\varepsilon(\cdot, s) \tilde{h}_\varepsilon(\cdot, s)\|_{L^{\frac{19}{8}}(\Omega)} ds \\
& \leq C_{5,****} \int_{T_{1,****}}^t (t-s)^{-\frac{1}{2}-\frac{9}{19}} e^{-\tilde{\lambda}_{1,****}(t-s)} \|n_\varepsilon(\cdot, s)\|_{L^{\frac{19}{3}}(\Omega)} \|\tilde{h}_\varepsilon(\cdot, s)\|_{L^{\frac{190}{3}}(\Omega)} ds \\
& \leq C_{6,****} \quad \text{for all } t > 2T_{1,****},
\end{aligned} \tag{6.81}$$

where we have used the fact that

$$\int_{T_{1,****}}^t (t-s)^{-\frac{1}{2}-\frac{9}{19}} e^{-\tilde{\lambda}_{1,****}(t-s)} ds \leq \int_1^\infty \sigma^{-\frac{1}{2}-\frac{9}{19}} e^{-\tilde{\lambda}_{1,****}\sigma} d\sigma < +\infty.$$

In combination with (6.79)–(6.81), we may finally derive

$$\|n_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq C_{7,****} \quad \text{for all } t \geq 2T_{1,****} \tag{6.82}$$

by some basic calculation. This completes the proof of Lemma 6.11. \square

Along the idea of the proof of Section 7.3 in [51], we could derive the Hölder regularity of the components of a solution on intervals of the form $(T_0, T_0 + 1)$ for any large $T_0 > 2$. For this purpose, we firstly introduce the following cut-off functions, which will play a key role in deriving higher order regularity for solution of problem (2.1).

Definition 6.1. Let $\xi : \mathbb{R} \rightarrow [0, 1]$ be a smooth, monotone function, satisfying $\xi \equiv 0$ on $(-\infty, 0]$ and $\xi \equiv 1$ on $(1, \infty)$ and for any $t_0 \in \mathbb{R}$ we let $\xi_{t_0} := \xi(t - t_0)$.

Due to the above cut-off function, it follows from maximal Sobolev regularity that the solution $(n_\varepsilon, c_\varepsilon, u_\varepsilon)$ even satisfies estimates in appropriate Hölder spaces:

Lemma 6.12. *Let $(n_\varepsilon, c_\varepsilon, u_\varepsilon)$ be a solution of (2.1) as well as (1.12) and (1.15) hold. Then for all $p \geq 2$, there exist $T_{*,8} > 0$ and $C_{*,8} > 0$ such that for all $\varepsilon \in (0, 1)$,*

$$\|u_\varepsilon\|_{L^p((t, t+1); W^{2,p}(\Omega))} + \|u_{\varepsilon t}\|_{L^p(\Omega \times (t, t+1))} \leq C_{*,8} \quad \text{for all } t \geq T_{*,8}. \tag{6.83}$$

Proof. Firstly, for any $p \geq 2$, we invoke Lemmas 6.7 and 6.11 to fix $\tilde{T}_{1,*}$ as well as $\tilde{C}_{1,*} > 0$ and $\tilde{C}_{2,*} > 0$ such that with $q := \max\{6p, \frac{6p}{2p-3}\}$ we have

$$\|n_\varepsilon\|_{L^\infty((t-1, t+1); L^{\frac{3p}{2}}(\Omega))} \leq \tilde{C}_{1,*} \quad \text{for all } t > \tilde{T}_{1,*} \tag{6.84}$$

and

$$\|u_\varepsilon\|_{L^\infty((t-1; t+1); L^q(\Omega))} \leq \tilde{C}_{2,*} \quad \text{for all } t > \tilde{T}_{1,*}. \quad (6.85)$$

For given $t_0 > \tilde{T}_{1,*}$, we let ξ_{t_0} be as defined in Definition 6.1. Then the function $v_\varepsilon : \Omega \times (t_0 - 1, \infty) \rightarrow \mathbb{R}^3$ defined by $v_\varepsilon(x, t) := \xi_{t_0}(t)u_\varepsilon(x, t)$, $(x, t) \in \Omega \times (t_0 - 1, \infty)$, is a weak solution in $L^\infty_{loc}([t_0 - 1, \infty); L^2_\sigma(\Omega)) \cap L^2_{loc}([t_0 - 1, \infty); W^{1,2}_{0,\sigma}(\Omega) \cap W^{2,2}(\Omega))$ of

$$v_{\varepsilon t} + Av_\varepsilon = h_\varepsilon \quad \text{in } \Omega \times (t_0 - 1, \infty), \quad (6.86)$$

where

$$v_\varepsilon(\cdot, t_0 - 1) \equiv 0 \quad \text{and} \quad h_\varepsilon = \xi_{t_0}(t)\mathcal{P}[-\kappa(Y_\varepsilon u_\varepsilon \cdot \nabla)u_\varepsilon + n_\varepsilon \nabla \phi] + \xi'_{t_0} u_\varepsilon.$$

To estimate the inhomogeneity h_ε herein, we first note that the known maximal Sobolev regularity estimate for the Stokes semigroup ([14]) yields that there exist positive constants $\tilde{k}_{1,*} > 0, \tilde{k}_{2,*} > 0$ and $\tilde{k}_{3,*} > 0$ such that

$$\begin{aligned} & \int_{t_0-1}^{t_0+1} \|\xi_{t_0}(t)\mathcal{P}[n_\varepsilon(\cdot, t)\nabla\phi]\|_{L^{\frac{3p}{2}}(\Omega)}^{\frac{3p}{2}} dt \\ & \leq \tilde{k}_{1,*} \int_{t_0-1}^{t_0+1} \|n_\varepsilon(\cdot, t)\nabla\phi\|_{L^{\frac{3p}{2}}(\Omega)}^{\frac{3p}{2}} dt \\ & \leq \tilde{k}_{1,*} \|\nabla\phi\|_{L^\infty(\Omega)} \int_{t_0-1}^{t_0+1} \|n_\varepsilon(\cdot, t)\|_{L^{\frac{3p}{2}}(\Omega)}^{\frac{3p}{2}} dt \\ & \leq \tilde{k}_{2,*} \end{aligned} \quad (6.87)$$

and

$$\int_{t_0-1}^{t_0+1} \|\xi'_{t_0} u_\varepsilon(\cdot, t)\|_{L^{\frac{3p}{2}}(\Omega)}^{\frac{3p}{2}} dt \leq \tilde{k}_{3,*}. \quad (6.88)$$

From the boundedness of the Helmholtz projection in L^p -spaces and the Hölder inequality we derive from Lemma 6.7 that there exist positive constant $\tilde{k}_{i,*}, i = 4, \dots, 8$ such that for any $t_0 > \tilde{T}_{1,*}$

$$\begin{aligned} & \int_{t_0-1}^{t_0+1} \|\xi_{t_0}\mathcal{P}[Y_\varepsilon u_\varepsilon \cdot \nabla u_\varepsilon](\cdot, t)\|_{L^{\frac{3p}{2}}(\Omega)}^{\frac{3p}{2}} dt \\ & \leq \tilde{k}_{4,*} \int_{t_0-1}^{t_0+1} \|\nabla u_\varepsilon(\cdot, t)\|_{L^{2p}(\Omega)}^{\frac{3p}{2}} \|u_\varepsilon(\cdot, t)\|_{L^{6p}(\Omega)}^{\frac{3p}{2}} dt \\ & \leq \tilde{k}_{5,*} \int_{t_0-1}^{t_0+1} \|\nabla u_\varepsilon(\cdot, t)\|_{L^{2p}(\Omega)}^{\frac{3p}{2}} dt \\ & \leq \tilde{k}_{6,*} \int_{t_0-1}^{t_0+1} \|u_\varepsilon(\cdot, t)\|_{W^{2,p}(\Omega)}^p \|u_\varepsilon(\cdot, t)\|_{L^{\frac{6p}{2p-3}}(\Omega)}^{\frac{p}{2}} dt \\ & \leq \tilde{k}_{7,*} \int_{t_0-1}^{t_0+1} \|u_\varepsilon(\cdot, t)\|_{W^{2,p}(\Omega)}^p dt \\ & \leq \tilde{k}_{8,*}. \end{aligned} \quad (6.89)$$

As a consequence of (6.87)–(6.89), once more by maximal Sobolev regularity estimates, now applied to (6.86), we obtain $k_{9,****} > 0$ and $k_{10,****} > 0$ satisfying

$$\begin{aligned}
& \int_{t_0}^{t_0+1} \|u_\varepsilon(\cdot, t)\|_{W^{2, \frac{3p}{2}}(\Omega)}^{\frac{3p}{2}} dt + \int_{t_0}^{t_0+1} \|v_{\varepsilon t}(\cdot, t)\|_{W^{2, \frac{3p}{2}}(\Omega)}^{\frac{3p}{2}} dt \\
\leq & \int_{t_0-1}^{t_0+1} \|u_\varepsilon(\cdot, t)\|_{W^{2, \frac{3p}{2}}(\Omega)}^{\frac{3p}{2}} dt + \int_{t_0-1}^{t_0+1} \|v_{\varepsilon t}(\cdot, t)\|_{W^{2, \frac{3p}{2}}(\Omega)}^{\frac{3p}{2}} dt \\
\leq & \tilde{k}_{9,*} \int_{t_0-1}^{t_0+1} \|h_\varepsilon(\cdot, t)\|_{L^{\frac{3p}{2}}(\Omega)}^{\frac{3p}{2}} dt \\
\leq & \tilde{k}_{10,*}.
\end{aligned} \tag{6.90}$$

Therefore, by the definition of ξ , for any $p > 1$, there exist positive constants $T_{*,8} > 0$ and $C_{*,8} > 0$ such that for any $t > T_{*,8}$ and for all $\varepsilon \in (0, 1)$,

$$\|u_\varepsilon(\cdot, t)\|_{L^p((t,t+1);W^{2,p}(\Omega))} + \|u_{\varepsilon t}(\cdot, t)\|_{L^p(\Omega \times (t,t+1))} \leq C_{*,8}. \tag{6.91}$$

This establishes (6.83) and thereby completes the proof. \square

Lemma 6.13. *Let $\alpha \geq 1$ and (1.15) hold. Then one can find $\mu_{*,9} \in (0, 1)$ and $T_{*,9} > 2$ and $C_{*,9} > 0$ such that for each $\varepsilon \in (0, 1)$ and $t > T_{*,9}$ we have*

$$\|u_\varepsilon(\cdot, t)\|_{C^{1+\mu_{*,9}, \frac{\mu_{*,9}}{2}}(\bar{\Omega} \times [t,t+1]); \mathbb{R}^3} \leq C_{*,9}. \tag{6.92}$$

Proof. Using sufficiently high values of p , in view of Lemma 6.14, an application of the embedding result in [1] refines this into the assertion on Hölder continuity. \square

Applying a similar reasoning (see the proof of Lemma 6.12), concerning c_ε we obtain bounds of the same kind by using Lemmas 6.11 and 6.13–6.10.

Lemma 6.14. *Let $(n_\varepsilon, c_\varepsilon, u_\varepsilon)$ be a solution of (2.1). If (1.12) and (1.15) holds, then for all $p > 1$, there exist $T_{*,10} > 2$ and $C_{*,10} > 0$ such that for any $\varepsilon \in (0, 1)$ and $t \geq T_{*,10}$ we have*

$$\|c_\varepsilon(\cdot, t)\|_{L^p((t,t+1);W^{2,p}(\Omega))} + \|c_{\varepsilon t}(\cdot, t)\|_{L^p(\Omega \times (t,t+1))} \leq C_{*,10}. \tag{6.93}$$

Proof. Firstly, applying Lemmas 6.11, 6.10 and 6.13, there exist $\tilde{T}_{1,**} > 2$ and positive constants $\tilde{C}_{1,**}$ as well as $\tilde{C}_{2,**}$ and $\tilde{C}_{3,**}$ such that

$$\|n_\varepsilon\|_{L^{2p}(\Omega \times (t_0-1, t_0+1))} \leq \tilde{C}_{1,**} \quad \text{for all } t_0 \geq \tilde{T}_{1,**} \tag{6.94}$$

as well as

$$\|u_\varepsilon\|_{L^\infty((t_0-1, t_0+1); L^\infty(\Omega))} \leq \tilde{C}_{2,**} \quad \text{for all } t_0 \geq \tilde{T}_{1,**} \quad (6.95)$$

and

$$\|\nabla c_\varepsilon\|_{W^{1,2p}(\Omega \times (t_0-1, t_0+1))} \leq \tilde{C}_{3,**} \quad \text{for all } t_0 \geq \tilde{T}_{1,**} \quad (6.96)$$

Now for fixed $t_0 > \tilde{T}_{1,**}$ we let ξ_{t_0} be as given by Definition 6.1 and consider the following problem

$$\begin{cases} \tilde{v}_{\varepsilon t} - \Delta \tilde{v}_\varepsilon = \tilde{h}_\varepsilon & x \in \Omega, t > t_0 - 1, \\ \frac{\partial \tilde{v}_\varepsilon}{\partial \nu} = 0, & x \in \partial\Omega, t > t_0 + 1, \\ \tilde{v}_\varepsilon(x, t_0 - 1) = 0, & x \in \Omega, \end{cases} \quad (6.97)$$

where

$$\tilde{h}_\varepsilon(\cdot, t) := \xi_{t_0}(t)(-c_\varepsilon + n_\varepsilon - u_\varepsilon \cdot \nabla c_\varepsilon) + \xi'_{t_0}(t)c_\varepsilon.$$

Here using (6.94)–(6.96) we see that for some $\tilde{C}_{4,**} > 0$ and $\tilde{C}_{5,**} > 0$ such that

$$\begin{aligned} & \int_{t_0-1}^{t_0+1} \|\tilde{h}_\varepsilon(\cdot, t)\|_{L^{2p}(\Omega)}^{2p} dt \\ & \leq \int_{t_0-1}^{t_0+1} \|\xi_{t_0}(t)(-c_\varepsilon + n_\varepsilon - u_\varepsilon \cdot \nabla c_\varepsilon) + \xi'_{t_0}(t)c_\varepsilon\|_{L^{2p}(\Omega)}^{2p} dt \\ & \leq \tilde{C}_{4,**} \int_{t_0-1}^{t_0+1} \left[\|c_\varepsilon(\cdot, t)\|_{L^{2p}(\Omega)}^{2p} + \|n_\varepsilon(\cdot, t)\|_{L^{2p}(\Omega)}^{2p} \right] dt \\ & \quad + \tilde{C}_{4,**} \int_{t_0-1}^{t_0+1} \left[\|u_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)}^{2p} \|\nabla c_\varepsilon(\cdot, t)\|_{L^{2p}(\Omega)}^{2p} + 2\|\xi'_{t_0}\|_{L^\infty(\mathbb{R})}^{2p} \|c_\varepsilon(\cdot, t)\|_{L^{2p}(\Omega)}^{2p} \right] dt \\ & \leq \tilde{C}_{5,**}. \end{aligned} \quad (6.98)$$

Therefore, we may argue as above to conclude from maximal Sobolev regularity estimates that there exist $\tilde{C}_{6,**} > 0$ and $\tilde{C}_{7,**} > 0$ such that

$$\begin{aligned} & \int_{t_0-1}^{t_0+1} \|\tilde{v}_\varepsilon(\cdot, t)\|_{W^{2,p}(\Omega)}^{2p} dt + \|\tilde{v}_{\varepsilon t}(\cdot, t)\|_{L^{2p}(\Omega)}^{2p} dt \\ & \leq \tilde{C}_{6,**} \int_{t_0-1}^{t_0+1} \|\tilde{h}_\varepsilon(\cdot, t)\|_{L^{2p}(\Omega)}^{2p} dt \\ & \leq \tilde{C}_{7,**}. \end{aligned} \quad (6.99)$$

Again since $\tilde{v}_\varepsilon = c_\varepsilon$ a.e. in $\Omega \times (t_0, t_0 + 1)$ by definition of ξ_{t_0} , this shows (6.93) and thereby completes the proof of Lemma 6.14. \square

Again, this implies the Hölder estimates as follows.

Lemma 6.15. *Suppose that α and C_S satisfy (1.12) and (1.15), respectively, where C_S is the same as (1.6). Then one can find $\mu_{*,11} \in (0, 1)$ as well as $T_{*,11} > 2$ and $C_{*,11} > 0$ such that for all $\varepsilon \in (0, 1)$ and $t > T_{*,11}$,*

$$\|c_\varepsilon(\cdot, t)\|_{C^{1+\mu_{*,11}, \frac{\mu_{*,11}}{2}}(\bar{\Omega} \times [t, t+1])} \leq C_{*,11}. \quad (6.100)$$

Proof. In precisely the same manner as Lemma 6.13 was derived from Lemma 6.12, this follows from Lemma 6.14 by application of a standard embedding result. \square

Using once more Lemma 6.11 and Lemma 2.3, we can improve our knowledge on the regularity of n_ε .

Lemma 6.16. *Under the assumptions of Theorem 1.2, then one can find $\mu_{*,12} \in (0, 1)$ as well as $T_{*,12} > 0$ and $C_{*,12} > 0$ such that whenever $\varepsilon \in (0, 1)$, we have*

$$\|n_\varepsilon(\cdot, t)\|_{C^{\mu_{*,12}, \frac{\mu_{*,12}}{2}}(\bar{\Omega} \times [t, t+1])} \leq C_{*,12} \quad \text{for all } t > T_{*,12}. \quad (6.101)$$

Proof. Firstly, in light of Lemma 6.11 as well as Lemma 6.15 and Lemma 6.13, there exist $\tilde{\mu}_{1,***} \in (0, 1)$ as well as $\tilde{T}_{1,***} > 2$ and $\tilde{C}_{1,***} > 0$ such that for all $t > \tilde{T}_{1,***}$,

$$\|\nabla c_\varepsilon\|_{C^{\tilde{\mu}_{1,***}, \frac{\tilde{\mu}_{1,***}}{2}}(\bar{\Omega} \times [t, t+1])} + \|n_\varepsilon\|_{L^\infty(\bar{\Omega} \times [t, t+1])} + \|u_\varepsilon\|_{C^{\tilde{\mu}_{1,***}, \frac{\tilde{\mu}_{1,***}}{2}}(\bar{\Omega} \times [t, t+1])} \leq \tilde{C}_{1,***}. \quad (6.102)$$

In order to derive (6.101), we may now use that $\nabla \cdot (n_\varepsilon u_\varepsilon) = u_\varepsilon \cdot \nabla n_\varepsilon$ thanks to the fact that $\nabla u_\varepsilon \equiv 0$, we have

$$\begin{aligned} & n_\varepsilon(\cdot, t) \\ = & e^{(t-\tilde{T}_{1,***})\Delta} n_\varepsilon(\cdot, \tilde{T}_{1,***}) \\ & + \int_{\tilde{T}_{1,***}}^t e^{(t-s)\Delta} \nabla \cdot \{n_\varepsilon(\cdot, s) F_\varepsilon(n_\varepsilon(\cdot, s)) S_\varepsilon(x, n_\varepsilon(\cdot, s), c_\varepsilon(\cdot, s)) \nabla c_\varepsilon(\cdot, s) + n_\varepsilon(\cdot, s) u_\varepsilon(\cdot, s)\} ds \end{aligned} \quad (6.103)$$

for all $t > \tilde{T}_{1,***}$. Therefore, in view of (6.102) and the standard regularity arguments involving well-known smoothing properties of the Neumann heat semigroup $(e^{\tau\Delta})_{\tau \geq 0}$ (see e.g. [43]), one can readily derive the existence of $\mu_{*,12} \in (0, 1)$ as well as $T_{*,12}$ and $C_{*,12} > 0$ such that for any $\varepsilon > 0$, (6.101) holds. \square

Straightforward applications of standard Schauder estimates for the Stokes evolution equation and the heat equation, respectively, finally yield eventual smoothness of the solution $(n_\varepsilon, c_\varepsilon, u_\varepsilon)$.

Lemma 6.17. *Under the assumptions of Theorem 1.2, then one can find $\mu_{*,13} \in (0, 1)$ and $T_{*,13} > 2$ such that for some $C_{*,13} > 0$ with the property that for each $\varepsilon \in (0, 1)$,*

$$\|u_\varepsilon(\cdot, t)\|_{C^{2+\mu_{*,13}, 1+\frac{\mu_{*,13}}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3)} \leq C_{*,13} \quad \text{for all } t > T_{*,13} \quad (6.104)$$

as well as

$$\|c_\varepsilon(\cdot, t)\|_{C^{2+\mu_{*,13}, 1+\frac{\mu_{*,13}}{2}}(\bar{\Omega} \times [t, t+1])} \leq C_{*,13} \quad \text{for all } t > T_{*,13} \quad (6.105)$$

and

$$\|n_\varepsilon(\cdot, t)\|_{C^{2+\mu_{*,13}, 1+\frac{\mu_{*,13}}{2}}(\bar{\Omega} \times [t, t+1])} \leq C_{*,13} \quad \text{for all } t > T_{*,13}. \quad (6.106)$$

Proof. We first combine Lemma 6.13 as well as Lemma 6.15 and Lemma 6.16 to infer the existence of $\tilde{\mu}_{1,****} \in (0, 1)$ as well as $\tilde{T}_{1,****} > 0$ and $\tilde{C}_{1,****} > 0$ such that for all $t > \tilde{T}_{1,****}$,

$$\begin{aligned} & \|u_\varepsilon \cdot \nabla c_\varepsilon\|_{C^{\tilde{\mu}_{1,****}, \frac{\tilde{\mu}_{1,****}}{2}}(\bar{\Omega} \times [t, t+1])} + \|u_\varepsilon \cdot \nabla n_\varepsilon\|_{C^{\tilde{\mu}_{1,****}, \frac{\tilde{\mu}_{1,****}}{2}}(\bar{\Omega} \times [t, t+1])} \\ & + \|n_\varepsilon\|_{C^{\tilde{\mu}_{1,****}, \frac{\tilde{\mu}_{1,****}}{2}}(\bar{\Omega} \times [t, t+1])} + \|c_\varepsilon\|_{C^{\tilde{\mu}_{1,****}, \frac{\tilde{\mu}_{1,****}}{2}}(\bar{\Omega} \times [t, t+1])} \\ & \leq \tilde{C}_{1,****}. \end{aligned} \quad (6.107)$$

The standard parabolic Schauder estimates applied to the second equation in (2.1) (see [21]) thus provide $\tilde{C}_{2,****} > 0$ fulfilling

$$\|c_\varepsilon\|_{C^{2+\tilde{\mu}_{1,****}, 1+\frac{\tilde{\mu}_{1,****}}{2}}(\bar{\Omega} \times [t, t+1])} \leq \tilde{C}_{2,****} \quad \text{for all } t > \tilde{T}_{1,****}. \quad (6.108)$$

According to Lemma 6.16, it is possible to fix $\tilde{\mu}_{2,****} \in (0, 1)$ as well as $\tilde{T}_{2,****} > 2$ and $\tilde{C}_{3,****} > 0$ such that

$$\|n_\varepsilon\|_{C^{1+\tilde{\mu}_{2,****}, \frac{\tilde{\mu}_{2,****}}{2}}(\bar{\Omega} \times [t, t+1])} + \|u_\varepsilon\|_{C^{1+\tilde{\mu}_{2,****}, \frac{\tilde{\mu}_{2,****}}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3)} \leq \tilde{C}_{3,****} \quad \text{for all } t > \tilde{T}_{2,****}. \quad (6.109)$$

We next set $T := \tilde{T}_{2,****} + 1$ and let $t_0 > T$ be given. Then with ξ_{t_0} taken from Definition 6.1, we again use that $v_\varepsilon(\cdot, t) := \xi_{t_0} u_\varepsilon(\cdot, t)$, $(x \in \Omega, t > t_0 - 1)$, is a solution of

$$\begin{cases} v_{\varepsilon t} - \Delta v_\varepsilon = h_\varepsilon & x \in \Omega, t > t_0 - 1, \\ v_\varepsilon(x, t_0 - 1) = 0, & x \in \Omega, \end{cases} \quad (6.110)$$

where

$$h_\varepsilon = -\kappa(Y_\varepsilon u_\varepsilon \cdot \nabla)v + \nabla(\xi P_\varepsilon) + \xi n_\varepsilon \nabla \phi + \xi' u_\varepsilon.$$

Now from (6.109) and the smoothness of ξ we readily obtain $\tilde{\mu}_{3,****} \in (0, 1)$ and $\tilde{C}_{4,****} > 0$ fulfilling

$$\|h_\varepsilon\|_{C^{\tilde{\mu}_{3,****}, \frac{\tilde{\mu}_{3,****}}{2}}(\bar{\Omega} \times [t_0-1, t_0+1]; \mathbb{R}^3)} \leq \tilde{C}_{4,****}, \quad (6.111)$$

so that, the regularity estimates from Schauder theory for the Stokes evolution equation ([32]) ensure that (6.110) possesses a classical solution $\bar{v} \in C^{2+\tilde{\mu}_{3,****}, 1+\frac{\tilde{\mu}_{3,****}}{2}}(\bar{\Omega} \times [t_0-1, t_0+1])$ satisfying

$$\|\bar{v}\|_{C^{2+\tilde{\mu}_{3,****}, 1+\frac{\tilde{\mu}_{3,****}}{2}}(\bar{\Omega} \times [t_0-1, t_0+1]; \mathbb{R}^3)} \leq \tilde{C}_{5,****} \quad (6.112)$$

with some $\tilde{C}_{5,****} > 0$ which is independent of t_0 . This combined with the uniqueness property of (6.110), one can prove

$$\|u_\varepsilon\|_{C^{2+\tilde{\mu}_{3,****}, 1+\frac{\tilde{\mu}_{3,****}}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3)} \leq \tilde{C}_{6,****}. \quad (6.113)$$

Again relying on Lemma 6.16, this in turn warrants that for some $\tilde{\mu}_{4,****} \in (0, 1)$, $\tilde{T}_{4,****} > 0$ and $\tilde{C}_{7,****} > 0$ such that for all $t > \tilde{T}_{4,****}$

$$\|\nabla \cdot (n_\varepsilon F_\varepsilon(n_\varepsilon) S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon)\|_{C^{\tilde{\mu}_{4,****}, \frac{\tilde{\mu}_{4,****}}{2}}(\bar{\Omega} \times [t, t+1])} + \|u_\varepsilon \cdot \nabla n_\varepsilon\|_{C^{\tilde{\mu}_{4,****}, \frac{\tilde{\mu}_{4,****}}{2}}(\bar{\Omega} \times [t, t+1])} \leq \tilde{C}_{7,****}, \quad (6.114)$$

which along with the Schauder theory establishes

$$\|n_\varepsilon\|_{C^{2+\tilde{\mu}_{4,****}, 1+\frac{\tilde{\mu}_{4,****}}{2}}(\bar{\Omega} \times [t, t+1])} \leq \tilde{C}_{8,****}. \quad (6.115)$$

Finally, choose

$$T_{*,13} = \max\{\tilde{T}_{1,****}, \tilde{T}_{2,****}, \tilde{T}_{3,****}, \tilde{T}_{4,****}\}$$

and

$$\mu_{*,13} = \min\{\tilde{\mu}_{1,****}, \tilde{\mu}_{2,****}, \tilde{\mu}_{3,****}, \tilde{\mu}_{4,****}\},$$

then (6.108), (6.113), (6.115) imply (6.104)–(6.106). \square

Having found uniform Hölder bounds on $n_\varepsilon, c_\varepsilon$ and u_ε for $\varepsilon > 0$ in the previous three lemmas (see Lemmas 6.16 and 6.17), also n, c and u share this regularity and these bounds.

Lemma 6.18. *Assume that $\alpha \geq 1$ and $C_S < 2\sqrt{C_N}$, where C_N is the best Poincaré constant and C_S is given by (1.6). There exist $\theta \in (0, 1)$ as well as $T_0 > 0$ and $(\varepsilon_j)_{j \in \mathbb{N}} \subset (0, 1)$ of the sequence from Lemma 6.4 such that for any $t > T_0$*

$$\begin{cases} n \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]), \\ c \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]), \\ u \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3), \end{cases} \quad (6.116)$$

that $\varepsilon_j \searrow 0$ as $j \rightarrow \infty$ and

$$\begin{cases} n_\varepsilon \rightarrow n \in C^{1+\theta, \frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]), \\ c_\varepsilon \rightarrow c \in C^{1+\theta, \frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]), \\ u_\varepsilon \rightarrow u \in C^{1+\theta, \frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3) \end{cases} \quad \text{and} \quad (6.117)$$

as $\varepsilon = \varepsilon_j \searrow 0$. Moreover, there is $C > 0$ such that

$$\|c(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1])} \leq C \quad \text{for all } t > T_0 \quad (6.118)$$

as well as

$$\|n(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1])} + \|u(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3)} \leq C \quad \text{for all } t > T_0. \quad (6.119)$$

Proof. According to Lemma 6.17 and Lemma 6.4, an application of the Aubin-Lions lemma (see [30]) provides a sequence $(\varepsilon_j)_{j \in \mathbb{N}} \subset (0, 1)$ such that $\varepsilon_j \searrow 0$ as $j \rightarrow \infty$ and such that (6.116)–(6.119) hold. And thereby completes the proof. \square

Lemma 6.19. *Let $\alpha \geq 1$. Then one can find $\theta \in (0, 1)$ and $T > 0$ such that*

$$\|c(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T, \infty))} \leq C \quad (6.120)$$

as well as

$$\|u(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T, \infty); \mathbb{R}^3)} \leq C \quad (6.121)$$

and

$$\|n(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T, \infty))} \leq C. \quad (6.122)$$

Proof. Let $g := -\xi c + n\xi - \xi u \cdot \nabla c + c\xi'$, where $\xi := \xi_{T_0}$ is given by Definition 6.1 and T_0 is same as the previous lemmas. Then we consider the following problem

$$\begin{cases} \tilde{c}_t - \Delta \tilde{c} = g & x \in \Omega, t > T_0, \\ \tilde{c}(T_0) = 0, & x \in \Omega, \\ \frac{\partial \tilde{c}}{\partial \nu} = 0, & x \in \partial\Omega. \end{cases} \quad (6.123)$$

According to Lemma 6.18 and Definition 6.1, we can find $\theta \in (0, 1)$ such that

$$g \text{ is bounded in } C^\theta(\bar{\Omega} \times (T_0, \infty)).$$

Therefore, the regularity estimates from Schauder theory for the parabolic equation (see e.g. III.5.1 of [21]) ensure that problem (6.123) admits a unique solution $\tilde{c} \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T_0 + 1, \infty))$. This combined with the property of ξ implies that

$$c \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T_0 + 1, \infty)). \quad (6.124)$$

Finally, employing almost exactly the same arguments as in the proof of Lemma 6.17 (the minor necessary changes are left as an easy exercise to the reader), and taking advantage of (6.119), we conclude that

$$u \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T_0 + 1, \infty); \mathbb{R}^3) \quad (6.125)$$

and

$$n \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T_0 + 1, \infty)), \quad (6.126)$$

whence combining the result of (6.124) completes the proof. \square

Our main result on eventual smoothness thereby becomes immediate.

Lemma 6.20. *Under the assumptions of Theorem 1.2, the solution (n, c, u) of (2.1) constructed in Lemma 6.4 satisfies*

$$n(\cdot, t) \rightarrow \bar{n}_0 \text{ as well as } c(\cdot, t) \rightarrow \bar{n}_0 \text{ and } u(\cdot, t) \rightarrow 0 \text{ in } L^\infty(\Omega), \quad (6.127)$$

where $\bar{n}_0 = \frac{1}{|\Omega|} \int_{\Omega} n_0$.

Proof. Firstly, due to Lemmas 6.4, we derive from Lemma 6.19 that

$$n(t) \rightarrow \bar{n}_0 \quad \text{as well as} \quad c(t) \rightarrow \bar{n}_0 \quad \text{and} \quad u(t) \rightarrow 0 \quad \text{in} \quad L^2(\Omega) \quad \text{as} \quad t \rightarrow \infty, \quad (6.128)$$

where \bar{n}_0 is given by (6.8). Next, due to Lemma 6.19, one can obtain there exist positive constants M_1 and T such that for all $t > T$

$$\|n(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} + \|c(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} + \|u(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} \leq M_1. \quad (6.129)$$

Therefore, a straightforward reasoning based on interpolation and the compactness of the first among the continuous embeddings $C^{2+\theta}(\bar{\Omega}) \hookrightarrow L^\infty(\Omega) \hookrightarrow L^2(\Omega)$ shows that (6.128) and (6.129) entail (6.127). In fact, for any $\eta > 0$, we may employ an Ehrling-type lemma to pick $M_2 > 0$ fulfilling

$$\|n(\cdot, t) - \bar{n}_0\|_{L^\infty(\Omega)} \leq \frac{\eta}{2M_1} \|n(\cdot, t) - \bar{n}_0\|_{C^{2+\theta}(\bar{\Omega})} + M_2 \|n(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)} \quad (6.130)$$

as well as

$$\|c(\cdot, t) - \bar{n}_0\|_{L^\infty(\Omega)} \leq \frac{\eta}{2M_1} \|c(\cdot, t) - \bar{n}_0\|_{C^{2+\theta}(\bar{\Omega})} + M_2 \|c(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)} \quad (6.131)$$

and

$$\|u(\cdot, t)\|_{L^\infty(\Omega)} \leq \frac{\eta}{2M_1} \|u(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} + M_2 \|u(\cdot, t)\|_{L^2(\Omega)}. \quad (6.132)$$

Now due to (6.128), we may choose $t_0 > \max\{1, T\}$ large enough such that for all $t > t_0$,

$$\|n(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)} + \|c(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)} + \|u(\cdot, t)\|_{L^2(\Omega)} < \frac{\eta}{2M_2}, \quad (6.133)$$

which in conjunction with (6.130)–(6.132) entails that

$$\begin{aligned} \|n(\cdot, t) - \bar{n}_0\|_{L^\infty(\Omega)} &\leq \frac{\eta}{2M_1} \|n(\cdot, t) - \bar{n}_0\|_{C^{2+\theta}(\bar{\Omega})} + M_2 \|n(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)} \\ &< \frac{\eta}{2M_1} M_1 + M_2 \frac{\eta}{2M_2} \\ &= \eta \quad \text{for all } t > t_0, \end{aligned}$$

$$\begin{aligned} \|c(\cdot, t) - \bar{n}_0\|_{L^\infty(\Omega)} &\leq \frac{\eta}{2M_1} \|c(\cdot, t) - \bar{n}_0\|_{C^{2+\theta}(\bar{\Omega})} + M_2 \|c(\cdot, t) - \bar{n}_0\|_{L^2(\Omega)} \\ &< \frac{\eta}{2M_1} M_1 + M_2 \frac{\eta}{2M_2} \\ &= \eta \quad \text{for all } t > t_0 \end{aligned}$$

as well as and

$$\begin{aligned}\|u(\cdot, t)\|_{L^\infty(\Omega)} &\leq \frac{\eta}{2M_1} \|u(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} + M_2 \|u(\cdot, t)\|_{L^2(\Omega)} \\ &< \frac{\eta}{2M_1} M_1 + M_2 \frac{\eta}{2M_2} \\ &= \eta \quad \text{for all } t > t_0,\end{aligned}$$

which combined with the fact that $\eta > 0$ is arbitrary yields to (6.127), and thereby completes the proof. □

In order to prove Theorem 1.2, we now only have to collect the results prepared during this section:

Proof of Theorem 1.2.

Proof. The statement is evidently implied by Lemmas 6.18–6.20. □

Acknowledgement: This work is partially supported by Shandong Provincial Science Foundation for Outstanding Youth (No. ZR2018JL005).

References

- [1] H. Amann, *Compact embeddings of vector-valued Sobolev and Besov spaces*, Glasnik Mat., 55(2000), 161–177.
- [2] N. Bellomo, A. Belloquid, Y. Tao, M. Winkler, *Toward a mathematical theory of Keller–Segel models of pattern formation in biological tissues*, Math. Models Methods Appl. Sci., 25(2015), 1663–1763.
- [3] T. Black, *Global solvability of chemotaxis-fluid systems with nonlinear diffusion and matrix-valued sensitivities in three dimensions*, Nonlinear Analysis, 180(2019), 129–153.
- [4] X. Cao, J. Lankeit, *Global classical small-data solutions for a three-dimensional Chemotaxis Navier-Stokes System involving Matrix-Valued Sensitivities*, Cal. Var. Partial Differential Equations, 55(2016), 107–123.

- [5] M. Chae, K. Kang, J. Lee, *Global Existence and temporal decay in Keller–Segel models coupled to fluid equations*, Comm. Part. Diff. Eqns., 39(2014), 1205–1235.
- [6] T. Cieřlak, M. Winkler, *Finite-time blow-up in a quasilinear system of chemotaxis*, Nonlinearity, 21(2008), 1057–1076.
- [7] C. Dombrowski, et al., *Self-concentration and large-scale coherence in bacterial dynamics*, Physical Review Letters, 93(2004), 098103.
- [8] R. Duan, A. Lorz, P. A. Markowich, *Global solutions to the coupled chemotaxis–fluid equations*, Comm. Part. Diff. Eqns., 35(2010), 1635–1673.
- [9] E. E. Espejo, M. Winkler, *Global classical solvability and stabilization in a two-dimensional chemotaxis–Navier–Stokes system modeling coral fertilization*, Nonlinearity, 31(2018), 1227–1259.
- [10] M. Di Francesco, A. Lorz, P. Markowich, *Chemotaxis–fluid coupled model for swimming bacteria with nonlinear diffusion: global existence and asymptotic behavior*, Discrete Contin. Dyn. Syst., 28(2010), 1437–1453.
- [11] A. Friedman, *Partial Differential Equations*, Holt, Rinehart Winston, New York, 1969.
- [12] H. Fujita, T. Kato, *On the Navier–Stokes initial value problem I*, Arch. Ration. Mech. Anal., 16(1964), 269–315.
- [13] D. Fujiwara, H. Morimoto, *An L^r -theorem of the Helmholtz decomposition of vector fields*, J. Fac. Sci. Univ. Tokyo, 24(1977), 685–700.
- [14] Y. Giga, H. Sohr, *Abstract L^p estimate for the Cauchy problem with applications to the Navier–Stokes equations in exterior domains*, J. Funct. Anal., 102(1991), 72–94.
- [15] Y. Giga, *Solutions for semilinear parabolic equations in L^p and regularity of weak solutions of the Navier–Stokes system*, J. Diff. Eqns., 61(1986), 186–212.
- [16] T. Hillen, K. Painter, *A user’s guide to PDE models for chemotaxis*, J. Math. Biol., 58(2009), 183–217.

- [17] D. Horstmann, M. Winkler, *Boundedness vs. blow-up in a chemotaxis system*, J. Diff. Eqns, 215(2005), 52–107.
- [18] S. Ishida, K. Seki and T, Yokota, *Boundedness in quasilinear Keller–Segel systems of parabolic–parabolic type on non-convex bounded domains*, J. Diff. Eqns., 256(2014), 2993–3010.
- [19] Y. Ke, J. Zheng, *An optimal result for global existence in a three-dimensional Keller–Segel–Navier–Stokes system involving tensor-valued sensitivity with saturation*, Calculus of Variations and Partial Diff. Eqns., 58(3)(2019), 58–109.
- [20] R. Kowalczyk, *Preventing blow-up in a chemotaxis model*, J. Math. Anal. Appl., 305(2005), 566–585.
- [21] O. A. Ladyzenskaja, V. A. Solonnikov and N. N. Ural’ceva, *Linear and quasi-Linear Equations of Parabolic Type*, Amer. Math. Soc. Transl., Providence, RI, 1968.
- [22] J. Leray, *Sur le mouvement d’un liquide visqueux emplissant l’espace*, Acta Math., 63(1934), 193–248.
- [23] D. Li, et al. *Boundedness in a three-dimensional Keller-Segel-Stokes system involving tensor-valued sensitivity with saturation*, Discrete and Continuous Dynamical Systems-B, 24(2019), 831–849.
- [24] Y. Li, Y. Li, *global boundedness of solutions for the chemotaxis-Navier-Stokes system in R^2* , J. Diff. Eqns., 261(2016), 6570–6613.
- [25] J. Liu, Y. Wang, *Boundedness and decay property in a three-dimensional Keller-Segel-Stokes system involving tensor-valued sensitivity with saturation*, J. Diff. Eqns., 261(2016), 967–999.

- [26] J. Liu, Y. Wang, *Global weak solutions in a three-dimensional Keller-Segel-Navier-Stokes system involving a tensor-valued sensitivity with saturation*, J. Diff. Eqns., 262(2017), 5271–5305.
- [27] J.-G. Liu, A. Lorz, *A coupled chemotaxis–fluid model: global existence*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 28(2011), 643–652.
- [28] A. Lorz, *Coupled chemotaxis fluid equations*, Math. Models Methods Appl. Sci., 20(2010), 987–1004.
- [29] Y. Peng, Z. Xiang, *Global existence and boundedness in a 3D Keller–Segel–Stokes system with nonlinear diffusion and rotational flux*, Z. Angew. Math. Phys., (2017), 68:68.
- [30] J. Simon, *Compact sets in the space $L^p(O, T; B)$* , Annali di Matematica Pura ed Applicata, 146(1)(1986), 65–96.
- [31] H. Sohr, *The Navier–Stokes equations, An elementary functional analytic approach*, Birkhäuser Verlag, Basel (2001).
- [32] V. A. Solonnikov, *Schauder estimates for the evolutionary generalized Stokes problem, Non-linear equations and spectral theory*, Amer. Math. Soc. Transl. Ser. 2, vol. 220, Amer. Math. Soc., Providence, RI, 2007, pp. 165–200.
- [33] Y. Tao, M. Winkler, *Global existence and boundedness in a Keller–Segel–Stokes model with arbitrary porous medium diffusion*, Discrete Contin. Dyn. Syst. Ser. A, 32(2012), 1901–1914.
- [34] Y. Tao, M. Winkler, *Locally bounded global solutions in a three-dimensional chemotaxis–Stokes system with nonlinear diffusion*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 30(2013), 157–178.
- [35] Y. Tao, M. Winkler, *Boundedness and decay enforced by quadratic degradation in a three-dimensional chemotaxis–fluid system*, Z. Angew. Math. Phys., 66(2015), 2555–2573.

- [36] J. I. Tello, M. Winkler, *A chemotaxis system with logistic source*, Comm. Partial Diff. Eqns., 32(2007), 849–877.
- [37] I. Tuval, L. Cisneros, C. Dombrowski, et al., *Bacterial swimming and oxygen transport near contact lines*, Proc. Natl. Acad. Sci. USA, 102(2005), 2277–2282.
- [38] L. Wang, C. Mu, P. Zheng, *On a quasilinear parabolic–elliptic chemotaxis system with logistic source*, J. Diff. Eqns., 256(2014), 1847–1872.
- [39] Y. Wang, M. Winkler, Z. Xiang, *Global classical solutions in a two-dimensional chemotaxis–Navier–Stokes system with subcritical sensitivity*, Annali della Scuola Normale Superiore di Pisa–Classe di Scienze. XVIII, (2018), 2036–2145.
- [40] Y. Wang, Z. Xiang, *Global existence and boundedness in a Keller–Segel–Stokes system involving a tensor-valued sensitivity with saturation: the 3D case*, J. Differ. Eqns. 261(2016), 4944–4973.
- [41] M. Wiegner, *The Navier–Stokes equations a neverending challenge?* Jahresber. Deutsch. Math.-Verein., 101(1999), 1–25.
- [42] M. Winkler, *Does a volume-filling effect always prevent chemotactic collapse*, Math. Methods Appl. Sci., 33(2010), 12–24.
- [43] M. Winkler, *Boundedness in the higher-dimensional parabolic–parabolic chemotaxis system with logistic source*, Comm. Partial Diff. Eqns., 35(2010), 1516–1537.
- [44] M. Winkler, *Aggregation vs. global diffusive behavior in the higher-dimensional Keller–Segel model*, J. Diff. Eqns., 248(2010), 2889–2905.
- [45] M. Winkler, *Global large-data solutions in a chemotaxis–(Navier–)Stokes system modeling cellular swimming in fluid drops*, Comm. Partial Diff. Eqns., 37(2012), 319–351.
- [46] M. Winkler, *Finite-time blow-up in the higher-dimensional parabolic–parabolic Keller–Segel system*, J. Math. Pures Appl., 100(2013), 748–767.

- [47] M. Winkler, *Stabilization in a two-dimensional chemotaxis–Navier–Stokes system*, Arch. Ration. Mech. Anal., 211(2014), 455–487.
- [48] M. Winkler, *Boundedness and large time behavior in a three-dimensional chemotaxis–Stokes system with nonlinear diffusion and general sensitivity*, Calculus of Variations and Partial Diff. Eqns., 54(2015), 3789–3828.
- [49] M. Winkler, *Large-data global generalized solutions in a chemotaxis system with tensor-valued sensitivities*, SIAM J. Math. Anal., 47(2015), 3092–3115.
- [50] M. Winkler, *Global weak solutions in a three-dimensional chemotaxis–Navier–Stokes system*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 33(2016), 1329–1352.
- [51] M. Winkler, *How far do oxytaxis-driven forces influence regularity in the Navier-Stokes system?* Trans. Am. Math. Soc., 369(2017), 3067–3125.
- [52] M. Winkler, *Global existence and stabilization in a degenerate chemotaxis–Stokes system with mildly strong diffusion enhancement*, J. Diff. Eqns., 264(2018), 6109–6151.
- [53] M. Winkler, *Does fluid interaction affect regularity in the three-dimensional Keller-Segel System with saturated sensitivity?*, J. Math. Fluid Mechanics, 20(2018), 1889–1909.
- [54] M. Winkler, *Can rotational fluxes impede the tendency toward spatial homogeneity in nutrient taxis (-Stokes) systems?*, International Mathematics Research Notices, 2019.
- [55] M. Winkler, *A three-dimensional Keller-Segel-Navier-Stokes system with logistic source: Global weak solutions and asymptotic stabilization*, J. Func. Anal., 276(2019), 1339–1401.
- [56] C. Xue, *Macroscopic equations for bacterial chemotaxis: integration of detailed biochemistry of cell signaling*, J. Math. Biol., 70(2015), 1–44.
- [57] C. Xue, H. G. Othmer, *Multiscale models of taxis-driven patterning in bacterial populations*, SIAM J. Appl. Math., 70(2009), 133–167.

- [58] Q. Zhang, X. Zheng, *Global well-posedness for the two-dimensional incompressible chemotaxis–Navier–Stokes equations*, SIAM J. Math. Anal., 46(2014), 3078–3105.
- [59] J. Zheng, *Boundedness of solutions to a quasilinear parabolic–elliptic Keller–Segel system with logistic source*, J. Diff. Eqns., 259(2015), 120–140.
- [60] J. Zheng, *A note on boundedness of solutions to a higher-dimensional quasi-linear chemotaxis system with logistic source*, Zeitsc. Angew. Mathe. Mech., 97(2017), 414–421.
- [61] J. Zheng, *Global weak solutions in a three-dimensional Keller–Segel–Navier–Stokes system with nonlinear diffusion*, J. Diff. Eqns., 263(2017), 2606–2629.
- [62] J. Zheng, *Global weak solutions in a three-dimensional Keller–Segel–Navier–Stokes system with nonlinear diffusion*, J. Diff. Eqns., 263(2017), 2606–2629.
- [63] J. Zheng, *An optimal result for global classical and bounded solutions in a two-dimensional Keller–Segel–Navier–Stokes system with sensitivity*, arXiv:1903.01033, 2019.
- [64] J. Zheng, *Global solvability and stabilization in a two-dimensional chemotaxis–Navier–Stokes system with chemotactic sensitivity modeling coral fertilization*, Preprint.
- [65] J. Zheng, *An optimal result for global very weak solutions to a chemotaxis–fluid system with nonlinear diffusion and rotational flux*, Preprint.
- [66] J. Zheng, *An optimal result for global existence and boundedness in a three-dimensional Keller–Segel–Stokes system with nonlinear diffusion*, J. Diff. Eqns., 267(2019), 2385–2415.
- [67] J. Zheng, Y. Ke, *Large time behavior of solutions to a fully parabolic chemotaxis–haptotaxis model in N dimensions*, J. Diff. Eqns., 266(2019), 1969–2018.