

# ON SOME CONNECTED GROUPS OF AUTOMORPHISMS OF WEIL ALGEBRAS

MIROSLAV KUREŠ AND JAN ŠÚTORA

**ABSTRACT.** The method of direct calculation of the group of  $\mathbb{R}$ -algebra automorphisms of a Weil algebra is presented in detail. The paper is focused on the case of a one-componental group and presents two cases of values of the determinant of its linear part.

## INTRODUCTION

The paper is devoted to groups of  $\mathbb{R}$ -algebra automorphisms of Weil algebras, which are important examples of algebraic groups. We consider the usual Euclidean topology. The essential fact ([1]) is that every algebraic group has as irreducible components the cosets modulo a closed connected normal subgroup of finite index, called the *identity component*. (It contains the neutral element which is the identity automorphism.) Those irreducible components turn out to be also the connected components. We denote the group of  $\mathbb{R}$ -algebra automorphisms of a Weil algebra  $A$  by  $\text{Aut } A$  and its identity component by  $G_A$ .

In differential geometry, Weil algebras play a fundamental role in the theory of Weil bundles  $T^A M$  over a smooth manifold  $M$  which generalize well-known higher order velocities bundles  $T_n^r M$ . Contact elements are defined as orbits of the group action. In the classical case, they form the fiber bundle denoted by  $K_n^1 M = \text{reg } T_n^1 M / G_n^1$  where  $G_n^1$  denotes the first order jet group. It is the bundle of contact elements; moreover the bundle of oriented contact elements is obtained if we replace the whole group  $G_n^1$  by its identity component which is the subgroup of matrices with positive determinant, see [2]. We refer to paper [4] for the case of higher order jet groups  $G_n^r$ , bundles  $K_n^r M = \text{reg } T_n^r M / G_n^r$  of higher order contact elements and the generalizations: groups of Weil algebra  $\mathbb{R}$ -algebra automorphisms  $\text{Aut } A$  and bundles  $K^A M = \text{reg } T^A M / \text{Aut } A$  of Weil contact elements.

We investigate the case  $\text{Aut } A = G_A$ , it means Weil algebras possessing only one connected component. They represent an important case where orientation reversing reparametrizations for corresponding velocities are impossible.

## 1. BASIC CONCEPTS, DENOTATIONS AND THE METHOD

This section will, among other things, show how to effectively calculate a group of  $\mathbb{R}$ -algebra automorphisms of a Weil algebra.

The Weil algebra  $\mathbb{D}_2^2 / \langle X^2, Y^2 \rangle$  represents the Weil algebra of the iterated tangent functor. Let the residue classes of  $X$  and  $Y$  will be denoted by the same symbol  $X$  and  $Y$ , respectively. Then we choose the basis  $\{1, X, Y, XY\}$  and write elements  $a + bX + cY + dXY$  of this Weil algebra as vectors  $(a, b, c, d)$ . The endomorphisms

---

2010 *Mathematics Subject Classification.* 13H99, 16W20, 58A32.

*Key words and phrases.* Weil algebra,  $\mathbb{R}$ -algebra automorphism, connected component of a group, contact elements.

have a form

$$(a, b, c, d) \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & A & B & C \\ 0 & D & E & F \\ 0 & G & H & I \end{pmatrix} = (a, Ab + Dc + Gd, Bb + Ec + Hd, Cb + Fc + Id)$$

which follows from the fact  $1 \mapsto 1$ . It is therefore sufficient to investigate the third order submatrix  $\mathcal{M} = \begin{pmatrix} A & B & C \\ D & E & F \\ G & H & I \end{pmatrix}$ . In fact, we restrict to subspace formed by nilpotent elements (which is an ideal in the algebra and which is usually denoted by  $\mathfrak{n}$ ). For automorphisms, the matrix  $\mathcal{M}$  must be non-singular. However, we have to describe *algebra* endomorphisms and automorphisms. As

$$X \mapsto AX + BY + CXY \quad \text{and} \quad Y \mapsto DX + EY + FXY,$$

we compute

$$XY \mapsto ADX^2 + (AE + BD)XY + BEY^2$$

and it follows  $AD = 0$  and  $BE = 0$ . So, we have four variants

- (i)  $B = 0, D = 0$
- (ii)  $A = 0, E = 0$
- (iii)  $D = 0, E = 0$
- (iv)  $A = 0, B = 0$

and, as  $G = AD = 0, H = BE = 0, I = AE + BD$ , they correspond with matrices

$$\begin{pmatrix} A & 0 & C \\ 0 & E & F \\ 0 & 0 & AE \end{pmatrix}, \begin{pmatrix} 0 & B & C \\ D & 0 & F \\ 0 & 0 & BD \end{pmatrix}, \begin{pmatrix} A & B & C \\ 0 & 0 & F \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & C \\ D & E & F \\ 0 & 0 & 0 \end{pmatrix}.$$

It is clear that the third and fourth matrices will be killed in the case of automorphisms (they have zero row). Thus,  $\mathbb{R}$ -algebra automorphisms are represented by matrices

$$\begin{pmatrix} A & 0 & C \\ 0 & E & F \\ 0 & 0 & AE \end{pmatrix}, \begin{pmatrix} 0 & B & C \\ D & 0 & F \\ 0 & 0 & BD \end{pmatrix},$$

in which  $A \neq 0, E \neq 0$  and  $B \neq 0, D \neq 0$ , respectively. As non-zero coefficients can be either positive or negative, we find that the group of  $\mathbb{R}$ -algebra automorphisms has eight connected components. For the determinant  $\mathfrak{D} = \det \mathcal{M}$ , we have

$$\mathfrak{D} = A^2 E^2 \quad \text{or} \quad \mathfrak{D} = -B^2 D^2.$$

If we make a deeper restriction, only for the supspace formed by elements of  $\mathfrak{n}/\mathfrak{n}^2$ , we consider the matrix  $\mathcal{M}_1$  having two possible forms

$$\begin{pmatrix} A & 0 \\ 0 & E \end{pmatrix}, \begin{pmatrix} 0 & B \\ D & 0 \end{pmatrix},$$

( $A \neq 0, E \neq 0, B \neq 0, D \neq 0$ ), and, for the determinant  $\mathfrak{D}_1 = \det \mathcal{M}_1$ , we have

$$\mathfrak{D}_1 = AE \quad \text{or} \quad \mathfrak{D}_1 = -BD.$$

## 2. THEOREM ON A CONNECTED GROUP OF $\mathbb{R}$ -ALGEBRA AUTOMORPHISMS OF A WEIL ALGEBRA

In the introductory example, we saw the group of automorphisms non-connected,  $G_A$  was a proper subgroup of  $\text{Aut } A$ . We now want to focus only on algebras where the group is connected ( $G_A = \text{Aut } A$ ). We recall that in [3] is proved that the Weil

algebra  $\mathbb{D}_2^6 / \langle X^3 + Y^4, X^4 + Y^5 \rangle$  possesses a connected (one-componental) group of  $\mathbb{R}$ -algebra automorphisms and

$$\mathcal{M}_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

(see Theorem 1 and its proof in [3] for more details). and

$$\mathfrak{D}_1 = 1.$$

The following assertion holds.

**THEOREM.** *Let  $A$  is a Weil algebra with connected group of  $\mathbb{R}$ -algebra automorphisms. Then there may occur one of the following two cases for the determinant  $\mathfrak{D}_1$ :*

- (i)  $\mathfrak{D}_1$  is constant with the value 1
- (ii)  $\mathfrak{D}_1$  takes all real values from the interval  $(0, \infty)$ .

*Proof.* It is obvious that the map assigning the determinant  $\mathfrak{D}_1$  to an  $\mathbb{R}$ -algebra automorphism is a continuous function and, moreover, a group homomorphism. It is also clear that its image is a subset of the interval  $(0, \infty)$ . So it is clear that only possibilities (i) and (ii) can occur because we have no other connected subgroup of the group  $(0, \infty)$  (with the usual multiplication of reals) than these two. The case (i) occurs for the Weil algebra

$$\mathbb{D}_2^6 / \langle X^3 + Y^4, X^4 + Y^5 \rangle.$$

It remains to show that the case (ii) may occur, too.

Let us consider  $A = \mathbb{D}_2^4 / \langle X^3Y, X^2Y^2, Y^4, X^3 - Y^3 \rangle$ . We choose the basis

$$\{1, X, Y, X^2, XY, Y^2, X^3, X^2Y, XY^2, X^4\}.$$

Then the endomorphism  $\varphi: A \rightarrow A$  maps

$$\phi(X) = AX + BY + CX^2 + DXY + EY^2 + FX^3 + GX^2Y + HXY^2 + IX^4$$

and

$$\phi(Y) = JX + KY + LX^2 + MXY + NY^2 + PX^3 + QX^2Y + RXY^2 + SX^4.$$

For  $\varphi$  to be an  $\mathbb{R}$ -algebra automorphism, it is necessary

- (i)  $AK - BJ \neq 0$
- (ii)  $\varphi(X^3Y) = \varphi(0) = 0$ , but it means that in the expression of  $\varphi(X^3Y)$  the coefficient standing at  $X^4 - XY^3$  is zero, which gives the equation

$$A^3J - B^3J - 3AB^2K = 0 \quad (1)$$

- (iii)  $\varphi(X^2Y^2) = \varphi(0) = 0$ , but it means that in the expression of  $\varphi(X^2Y^2)$  the coefficient standing at  $X^4 - XY^3$  is zero, which gives the equation

$$A^2J^2 - 2B^2JK - 2ABK^2 = 0 \quad (2)$$

- (iv)  $\varphi(Y^4) = \varphi(0) = 0$ , but it means that in the expression of  $\varphi(Y^4)$  the coefficient standing at  $X^4 - XY^3$  is zero, which gives the equation

$$J^4 - 4JK^3 = 0 \quad (3)$$

We start with the equation (3).

- ( $\alpha$ ) If  $J = 0$ , then  $A \neq 0$ ,  $K \neq 0$  and then  $B = 0$  due to (1) and (2).

- ( $\beta$ ) If  $J \neq 0$ , then  $J = \sqrt[3]{4}K$ , (1) transforms to

$$\left( \sqrt[3]{4}A^3 - \sqrt[3]{4}B^3 - 3AB^2 \right) K = 0 \quad (4)$$

and (2) transforms to

$$\left( \sqrt[3]{8}A^2 - 2\sqrt[3]{4}B^2 - 2AB \right) K^2 = 0, \quad (5)$$

but  $K \neq 0$  as  $J \neq 0$ . So the expressions in brackets have to be zero but the obtained system in unknowns  $A$  and  $B$  has only the solution  $A = B = 0$  which is impossible.

So we continue with  $B = 0$ ,  $J = 0$ ,  $A \neq 0$ ,  $K \neq 0$ . Moreover, it is necessary

- (v)  $\varphi(X^3 - Y^3) = \varphi(0) = 0$ , but it means that in the expression of  $\varphi(X^3 - Y^3)$  the coefficient standing at  $X^3 - Y^3$  is zero, which gives the equation

$$A^3 - K^3 = 0 \quad (6)$$

with the solution  $K = A$  and also in the expression of  $\varphi(X^3 - Y^3)$  the coefficient standing at  $X^4 - XY^3$  is zero, which gives the equation

$$3A^2CD - 3A^2M = 0 \quad (7)$$

with the solution  $M = C$ .

So, we have

$$\begin{aligned} \phi(X) &= AX + CX^2 + DXY + EY^2 + FX^3 + GX^2Y + HXY^2 + IX^4 \\ \phi(Y) &= AY + LX^2 + CXY + NY^2 + PX^3 + QX^2Y + RXY^2 + SX^4 \quad \text{where } A \neq 0. \end{aligned}$$

That is why we obtain

$$\mathcal{M} = \begin{pmatrix} A & 0 & C & D & E & F & G & H & I \\ 0 & A & L & C & N & P & Q & R & S \\ 0 & 0 & A^2 & 0 & 0 & 2AC & 2AD & 2AE & 2AF + C^2 - 2DE \\ 0 & 0 & 0 & A^2 & 0 & AL - AE & 2AC & AD + AN & AP + CL - AH - CE - DN \\ 0 & 0 & 0 & 0 & A^2 & -2AN & 2AL & 2AC & L^2 - 2AR - 2CN \\ 0 & 0 & 0 & 0 & 0 & 2A^3 & 0 & 0 & 6A^2C \\ 0 & 0 & 0 & 0 & 0 & 0 & A^3 & 0 & A^2L - 2A^2E \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & A^3 & -A^2D - 2A^2N \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2A^4 \end{pmatrix},$$

$$\mathcal{M}_1 = \begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix}$$

and

$$\mathfrak{D} = 4A^{21} \quad \text{and} \quad \mathfrak{D}_1 = A^2.$$

□

REMARK. We note that the result remains in force for local algebras over an arbitrary field of the characteristics 0.

## REFERENCES

- [1] Guil-Asensio, F. and Saorín, M., *The group of automorphisms of a commutative algebra*, Mathematische Zeitschrift 219, 31–48, 1995.
- [2] Guggenheimer, H., *Contact elements, contact correspondences and contact invariants*, Annali di Matematica Pura ed Applicata, IV. Ser. 120, 229–261, 1979.
- [3] Ivičič, V. and Kureš, M., *Two general examples of Weil algebras having the group of automorphisms connected*, Int. J. Geom. Meth. in Mod. Phys. 13, No.3, 1650035-1–1650035-19, 2016.
- [4] Kureš, M., *Local approach to higher-order contact elements*, Reports on Mathematical Physics 58, No. 3, 395–410, 2006.

BRNO UNIVERSITY OF TECHNOLOGY  
DEPARTMENT OF MATHEMATICS  
TECHNICKÁ 2, 61669, BRNO, CZECHIA  
E-mail address: kures@fme.vutbr.cz

BRNO UNIVERSITY OF TECHNOLOGY  
DEPARTMENT OF MATHEMATICS  
TECHNICKÁ 2, 61669, BRNO, CZECHIA  
*E-mail address:* `sutora@post.cz`