

# ADJUSTED SUBADDITIVITY OF RELATIVE ENTROPY FOR NON-COMMUTING CONDITIONAL EXPECTATIONS

NICHOLAS LARACUENTE

ABSTRACT. If a set of von Neumann subalgebras has a trivial intersection in finite dimension, then the sum of relative entropies of a given density to its projection in each such algebra is larger than a multiple of its relative entropy to its projection in the trivial intersection. This results in a subadditivity of relative entropy with a dimension and algebra-dependent, multiplicative constant. As a primary application, this inequality lets us derive relative entropy decay estimates in the form of modified logarithmic-Sobolev inequalities for complicated quantum Markov semigroups from those of simpler constituents.

## 1. INTRODUCTION

The strong subadditivity (SSA) of von Neumann entropy states that for a tripartite density  $\rho^{ABC}$  on systems  $ABC$ ,

$$H(AC)_\rho + H(BC)_\rho \geq H(ABC)_\rho + H(C)_\rho, \quad (1)$$

where  $H(\rho) = -\text{tr}(\rho \log \rho)$ , and  $H(A)_\rho = H(\text{tr}_{BC}(\rho))$ . Lieb and Ruskai proved SSA in 1973 [1]. SSA's impacts range from quantum Shannon theory [2] to holographic spacetime [3].

A later form by Petz in 1991 generalizes from subsystems to subalgebras. Let  $\mathcal{M}$  be a von Neumann algebra and  $\mathcal{N}$  be a subalgebra. Associated with  $\mathcal{N}$  is a unique conditional expectation  $\mathcal{E}_\mathcal{N}$  that projects an operator in  $\mathcal{M}$  onto  $\mathcal{N}$ . If  $\mathcal{S}, \mathcal{T} \subseteq \mathcal{M}$  as subalgebras, we call them a commuting square if  $\mathcal{E}_\mathcal{S}\mathcal{E}_\mathcal{T} = \mathcal{E}_\mathcal{T}\mathcal{E}_\mathcal{S} = \mathcal{E}_{\mathcal{S} \cap \mathcal{T}}$ , the conditional expectation onto their intersection. Petz's theorem [4] implies:

**Theorem 1.1** (Petz's Conditional Expectation SSA). *Let  $\mathcal{E}_\mathcal{S}, \mathcal{E}_\mathcal{T}$  be conditional expectations to subalgebras  $\mathcal{S}, \mathcal{T} \subseteq \mathcal{M}$ . If  $\mathcal{S}$  and  $\mathcal{T}$  form a commuting square, then for all densities  $\rho$ ,*

$$H(\mathcal{E}_\mathcal{S}(\rho)) + H(\mathcal{E}_\mathcal{T}(\rho)) \geq H(\mathcal{E}_{\mathcal{S} \cap \mathcal{T}}(\rho)) + H(\rho).$$

In terms of relative entropy,

$$D(\rho \| \mathcal{E}_\mathcal{S}(\rho)) + D(\rho \| \mathcal{E}_\mathcal{T}(\rho)) \geq D(\rho \| \mathcal{E}_{\mathcal{S} \cap \mathcal{T}}(\rho)) \quad (2)$$

for  $\mathcal{S}$  and  $\mathcal{T}$  forming a commuting square. As we note with Gao and Junge in [5], theorem 1.1 implies SSA and an uncertainty relation for mutually unbiased bases, joining these concepts.

In [5], we also show that the condition  $[\mathcal{E}_\mathcal{S}, \mathcal{E}_\mathcal{T}] = 0$  is necessary as well as sufficient in finite dimensions. Hence inequalities of the form

$$D(\rho \| \mathcal{E}_\mathcal{S}(\rho)) + D(\rho \| \mathcal{E}_\mathcal{T}(\rho)) \geq D(\rho \| \mathcal{E}_{\mathcal{S} \cap \mathcal{T}}(\rho)) + c \quad (3)$$

can only hold for all  $\rho$  when  $[\mathcal{E}_S, \mathcal{E}_T] \neq 0$  if  $c < 0$ . Nonetheless, in this work we consider:

**Definition 1.2** (Adjusted Subadditivity (ASA)). *Let  $\{\mathcal{E}_j : j \in 1 \dots J \in \mathbb{N}\}$  be a set of conditional expectations, and  $\mathcal{E}$  the conditional expectation onto their intersection algebra. We call the set  $\alpha$ -subadditive if*

$$\sum_j D(\rho \| \mathcal{E}_j(\rho)) \geq \alpha D(\rho \| \mathcal{E}(\rho))$$

for some  $\alpha > 0$ . We call it completely  $\alpha$ -subadditive if  $\{\hat{1}^C \otimes \mathcal{E}_j\}$  is  $\alpha$ -subadditive for any finite-dimensional extension  $C$ .

We prove  $\alpha$ -subadditivity for sets of conditional expectations  $\{\mathcal{E}_j\}$  onto algebras with trivial intersection, where  $\alpha$  depends on  $\{\mathcal{E}_j\}$  and the dimension  $d$ . When no base is given for the logarithm, we denote the natural logarithm. In example 4.5, we derive such an inequality for a pair of incompatible bases that are not mutually unbiased, avoiding the negative constant appearing in conventional uncertainty relations.

A form of ASA appears in Proposition 2.1 of [6] as quasi-factorization of relative entropy\*. As a conditional expectation may take the form of a Haar-random average over a unitary subgroup, we find conceptual analogies with unitary  $t$ -designs [8], scrambling [9] and rapid mixing [10]. Given a set of mutually commuting mixing or averaging operations, their composition will often show similar mixing in a straightforward way. One can interpret subadditivity of conditional expectation entropies in equation (2) this way. Non-commuting mixing operations should still intuitively imply some sort of overall mixing on the whole, but quantifying how is a much richer and harder problem. Many works focus on how local but overlapping mixing operations compose to mix an extensive system. ASA is particularly strong in that it bounds relative entropy without including additive corrections, as would a conventional uncertainty relation or perturbed subadditivity. Furthermore, ASA does not rely on nearness to independence, distinguishing it from usual quasi-factorization.

Decay and decoherence are some of the most vexing challenges to quantum technology. As a primary application, ASA allows us to derive exponential relative entropy decay rates of continuous-time quantum processes having subprocesses with strong decay. Because ASA bounds relative entropy with a multiplicative constant, it implies merging of a particularly strong form of decay estimate known as a modified logarithmic Sobolev inequality (MLSI). This does not follow from perturbations of subadditivity that include additive constants as in equation (3). Finally, we note that the stronger form in Conjecture 5.1 would have major consequences for decay inequalities that motivate future work. Were Conjecture 5.1 definitively refuted, it would suggest separation between types of decay.

**1.1. Primary Contributions.** First, we prove a basic bound on relative entropy:

---

\*We thank the authors of an in-preparation manuscript [7] for showing us an early draft that also considers and motivates an inequality of this form.

**Theorem 1.3.** *Given  $a \in [0, 1]$  and two densities  $\rho, \sigma$  in dimension  $d$  such that  $\rho \succ \sigma$  ( $\rho$  majorizes  $\sigma$ ),*

$$D(\rho \|(1 - \zeta)\hat{1}/d + \zeta\sigma) - (1 - a)D(\rho \|\hat{1}/d) \geq 0$$

for any  $\beta \in (0, 1)$  and

$$\zeta \leq a \min \left\{ \frac{1 - \beta}{n + a(1 - \beta) + 1}, \frac{\beta}{(1 - a\beta)n + a\beta + 1} \right\}.$$

We may optimize  $\beta$  in Theorem 1.3 for a given  $d$  and  $a$ . If we wish to avoid optimization,  $\beta = 1/2$  is a reasonable value. Theorem 1.3 compares forms of telescopic relative entropy [11].

**Theorem 1.4.** *Let  $\{(\mathcal{N}_j, \mathcal{E}_j) : j = 1 \dots J \in \mathbb{N}\}$  be a set of  $J$  von Neumann algebras of dimension  $d$  and associated conditional expectations. Assume that  $\bigcap_j \mathcal{N}_j = \mathbb{C}1$ , the physically trivial algebra of phase and normalization. Then  $\{\mathcal{E}_j\}$  is  $\alpha$ -subadditive for some  $\alpha > 0$ .*

*In particular, let  $S = \bigcup_{m \in \mathbb{N}} \{1 \dots J\}^{\otimes m}$  be the set of sequences of indices  $1 \dots J$ . For any  $s \in S$ , let  $\mathcal{E}^s$  denote the composition of conditional expectations  $\mathcal{E}_{j_1} \dots \mathcal{E}_{j_{|s|}}$  for  $s = (j_1, \dots, j_{|s|})$ , where  $|s|$  denotes the sequence length. Let  $\mu : S \rightarrow [0, 1]$  be a probability measure on  $S$  that is non-zero only on finite sequences,  $k_s$  be the maximum number of repeats of any index in  $s$ , and  $k = \sum_s \mu(s)k_s$ . If for all  $d$ -dimensional densities  $\rho$ ,*

$$\sum_{s \in S} \mu(s) \mathcal{E}^s(\rho) = (1 - \zeta)\hat{1}/d + \zeta\Phi(\rho)$$

for some unital channel  $\Phi$  such that  $\mathcal{E} \circ \Phi = \mathcal{E}$ , then  $\{\mathcal{E}_j\}$  is  $1/(2k(\lceil \log_\zeta(2/(3d+5)) \rceil))$ -subadditive, where  $\lceil \cdot \rceil$  is the ceiling function.

As a primary application, ASA allows us to combine a particular class of decay estimates known as modified logarithmic-Sobolev inequalities (MLSI). Let  $\Phi^t$  be a family of quantum channels in dimension  $d$  parameterized by  $t \in \mathbb{R}^+$ , such that  $\Phi^s \circ \Phi^t = \Phi^{s+t}$  for all  $s, t \in \mathbb{R}^+$ . This family of channels is thereby a semigroup under composition, and there exists a Lindbladian generator given by  $\mathcal{L} = \lim_{t \rightarrow 0} (\hat{1} - \Phi^t)$  such that  $\Phi^t = e^{-t\mathcal{L}}$ . As studied in [12, 13, 14], we say that  $\mathcal{L}$  has  $\lambda$ -MLSI if for any input density  $\rho$ ,

$$D(\Phi^t(\rho) \|\Phi^\infty(\rho)) \leq e^{-\lambda t} D(\rho \|\Phi^\infty(\rho)). \quad (4)$$

As defined in [14], we say that  $\mathcal{L}$  has  $\lambda$ -CLSI (complete logarithmic-Sobolev inequality) if for any bipartite density  $\rho^{AB}$ ,

$$D((\Phi^t \otimes \hat{1}^B)(\rho) \|\Phi^\infty \otimes \hat{1}^B(\rho)) \leq e^{-\lambda t} D(\rho \|\Phi^\infty \otimes \hat{1}^B(\rho)). \quad (5)$$

We obtain the following result:

**Theorem 1.5** (MLSI Merge). *Let  $\{\Phi_j^t : j \in 1 \dots J \in \mathbb{N}\}$  be selfadjoint quantum Markov semigroups such that  $\Phi_j^t = \text{ad}_{\exp(-\mathcal{L}_j t)}$  with fixpoint conditional expectation  $\Phi_j^\infty = \mathcal{E}_j$ . Let  $\Phi^t$  be the semigroup generated by  $\mathcal{L} = \sum_j \mathcal{L}_j + \mathcal{L}_0$ , where  $\mathcal{L}_0$  generates  $\Phi_0^t$  with fixpoint algebra containing the fixpoint algebra of  $\{\Phi_j\}$ . Let  $\{\mathcal{E}_j\}$  be  $\alpha$ -subadditive, and  $\Phi_j^t$  have  $\lambda_j$ -MLSI for each  $j$ . Then  $\Phi^t$  has  $\alpha \times \min_j \{\lambda_j\}$ -MLSI.*

We obtain a corresponding result for CLSI (Corollary 4.1) given complete adjusted subadditivity.

Section 2 proves Theorem 1.3. Section 3 proves Theorem 1.4. Section 4 describes applications that use ASA to combine modified logarithmic Sobolev inequalities and proves Theorem 1.5. Section 5 introduces the strong adjusted subadditivity conjecture 5.1 and concludes.

## 2. RELATIVE ENTROPY COMPARISONS

Since relative entropy is biconvex,

$$D(\rho\|(1-\zeta)\omega + \zeta\sigma) \leq (1-\zeta)D(\rho\|\omega) + \zeta D(\rho\|\sigma)$$

for any densities  $\rho, \omega, \sigma$  and  $\zeta \in [0, 1]$ . There are however examples of densities  $\rho, \omega, \sigma$  for which  $D(\rho\|(1-\zeta)\omega + \zeta\sigma) = 0$  while

$$D(\rho\|\omega), D(\rho\|\sigma) > 0,$$

or where  $D(\rho\|(1-\zeta)\omega + \zeta\sigma)$  is finite while either term in the sum is infinite. Hence in general, there is no way to multiplicatively restrict the extent of non-concavity of relative entropy.

If we restrict to  $\omega = \hat{1}/d$ , then we see that  $D(\rho\|(1-\zeta)\hat{1}/d + \zeta\sigma)$  and  $D(\rho\|\hat{1}/d)$  are necessarily both finite. These conditions are still insufficient to lower bound  $D(\rho\|(1-\zeta)\hat{1}/d + \zeta\sigma)$  in terms of  $D(\rho\|\hat{1}/d)$ . We may for instance take  $\sigma$  pure, and  $\rho = (1-\zeta)\hat{1}/d + \zeta\sigma$ . The final condition we add is that  $\rho$  majorizes  $\sigma$  ( $\rho \succ \sigma$ ), in which case we obtain Theorem 1.3. The rest of this section contains Lemmas leading up to proof of this Theorem.

**Lemma 2.1.** *Let  $\rho, \omega$  be simultaneously diagonal densities of dimension  $d$ . Let  $\delta > 0$ . Let  $i \neq j \in 1\dots d$  such that  $\rho_i \geq \rho_j$ , and  $\rho_i\omega_j \geq \rho_j\omega_i$ . Let  $\omega \rightarrow \tilde{\omega}$  under the replacement  $\omega_i \rightarrow \tilde{\omega}_i = \omega_i - \delta, \omega_j \rightarrow \tilde{\omega}_j = \omega_j + \delta$ . Then  $D(\rho\|\omega) \leq D(\rho\|\tilde{\omega})$ .*

*Proof.* For any  $a > b \in \mathbb{R}^+$ ,  $(a-b)/a \leq \ln a - \ln b \leq (a-b)/b$ , as one can verify from  $(d/dx)(\ln x) = 1/x$ . Hence

$$\begin{aligned} D(\rho\|\tilde{\omega}) - D(\rho\|\omega) &= \rho_i(\ln \omega_i - \ln(\omega_i - \delta)) + \rho_j(\ln \omega_j - \ln(\omega_j + \delta)) \\ &\geq (\rho_i/\omega_i - \rho_j/\omega_j)\delta \geq 0 \end{aligned}$$

■

**Lemma 2.2.** *Let  $\rho$  and  $\sigma$  be two densities such that  $\rho \succ \sigma$ , and  $\zeta \in [0, 1]$ . Then*

$$D(\rho\|(1-\zeta)\hat{1}/d + \zeta\sigma) \geq D(\rho\|(1-\zeta)\hat{1}/d + \zeta\rho).$$

*Proof.* The main idea of this proof is that if  $\rho \succ \sigma$ , then flattening  $\rho$  until it becomes  $\sigma$  only increases the value of  $D(\rho\|(1-\zeta)\hat{1}/d + \zeta\rho)$ . First,

$$D(\rho\|(1-\zeta)\hat{1}/d + \zeta\sigma) \geq D(\rho\|(1-\zeta)\hat{1}/d + \zeta\mathcal{E}_{\rho'}(\sigma)),$$

by data processing under  $\mathcal{E}_{\rho'}$ , the conditional expectation onto the subalgebra that commutes with  $\rho$ . We hence assume that  $[\rho, \sigma] = 0$ , and they are simultaneously diagonal. Let  $\rho^\zeta = (1-\zeta)\hat{1}/d + \zeta\rho$ , and define  $\sigma^\zeta$  accordingly. Let  $\vec{\rho}^\zeta$  and  $\vec{\sigma}^\zeta$  be  $d$ -dimensional vectors of the eigenvalues of  $\rho^\zeta$  and  $\sigma^\zeta$  respectively, each in non-increasing order. Let  $\omega = \rho^\zeta$ . We alter  $\vec{\omega}$  via a cascading probability redistribution procedure consisting of the following steps, which transform it into a copy of  $\vec{\sigma}^\zeta$ :




FIGURE 1. Visualization of the cascading redistribution algorithm, which converts the distribution on the top-left to that on the bottom-right.

- (1) Start with the index  $i$  set to 1.
- (2) Let  $\Delta = \bar{\omega}_i - \bar{\sigma}_i^\zeta$ . If  $\Delta_i > 0$ , then
  - (a) Subtract  $\Delta$  from  $\bar{\omega}_i$ . Let  $j = i + 1$ .
  - (b) If  $\bar{\omega}_j < \bar{\sigma}_j^\zeta$ , then let  $\delta = \min\{\Delta, \bar{\sigma}_j^\zeta - \bar{\omega}_j\}$ . Add  $\delta$  to  $\bar{\omega}_j$  and subtract it from  $\Delta$ .
  - (c) If  $\Delta = 0$  or  $j = d$ , go on to step (3). Otherwise, increment  $j \rightarrow j + 1$ , and return to the previous substep (2b).
- (3) If  $i < d - 1$ , increment  $i \rightarrow i + 1$  and return to step (2). Otherwise, the procedure is done.

See figure 1. Since this procedure only subtracts from larger eigenvalues and adds to smaller ones, we apply Lemma 2.1 at each step that transfers probability mass from one index to another. If  $\bar{\rho}_i \geq \bar{\rho}_j$ , then  $\bar{\rho}_i/\bar{\rho}_i^\zeta \geq \bar{\rho}_j/\bar{\rho}_j^\zeta$ . Furthermore, it is always the case that  $\bar{\omega}_i \leq \bar{\rho}_i^\zeta$  if we are moving probability mass out of  $\bar{\omega}_i$ , and always that  $\bar{\omega}_j \geq \bar{\rho}_j^\zeta$  if we are moving probability into  $j$ . Hence  $\bar{\rho}_i/\bar{\rho}_j \geq \bar{\omega}_i/\bar{\omega}_j$ . This shows that  $D(\rho\|\rho^\zeta) = D(\bar{\rho}\|\bar{\rho}^\zeta) \leq D(\bar{\rho}\|\bar{\sigma}^\zeta)$ . Finally, using the simultaneous diagonality of  $\rho$  and  $\sigma$ , it is easy to see that  $D(\bar{\rho}\|\bar{\sigma}^\zeta) \leq D(\rho\|\sigma^\zeta)$ . ■

**Lemma 2.3.** *Let  $b \geq 0$ , and  $a \in [0, 1]$ . Then for  $\zeta \leq a/(1 + b)$ ,*

$$a \log(1 + b) \geq \log(1 + \zeta b) .$$

*Proof.* We solve

$$(1 + b)^a \geq 1 + \zeta b$$

yielding

$$\zeta \leq \frac{(1 + b)^a - 1}{b} .$$

We then estimate

$$\frac{(1+b)^a - 1}{b} = \frac{(1+b)^{1+a} - 1 - b}{b(1+b)} \geq \frac{1 + (1+a)b - 1 - b}{b(1+b)} = \frac{a}{1+b}$$

by Bernoulli's inequality. ■

**Lemma 2.4.** *Let  $\rho$  be given in its diagonal basis by  $(\rho_i)_{i=1}^n$ , where  $n$  is the dimension of the system. Let  $a, \beta \in (0, 1)$ ,  $\rho_i \geq 1/n \geq \rho_j$ , and*

$$\zeta \leq a \min \left\{ \frac{1 - \beta}{n + a(1 - \beta) + 1}, \frac{\beta}{(1 - a\beta)n + a\beta + 1} \right\}.$$

Then

$$\left( \frac{\partial}{\partial \rho_i} - \frac{\partial}{\partial \rho_j} \right) \text{tr}(\rho(a \log(n\rho) - \log((1 - \zeta)\hat{1} + \zeta n\rho))) \geq 0.$$

Note: in this Lemma and its proof, and thereafter within this section, we use the letter  $n$  rather than  $d$  for dimension to avoid possible confusion with the derivative.

*Proof.* Define  $\delta \equiv \rho_i - \rho_j \geq 0$ . For the purposes of the derivative,  $\text{tr}(\log(n\rho))$  is equivalent to  $\text{tr}(\log \rho)$ , since they differ only by a constant  $\log n$ . We directly calculate,

$$\frac{\partial}{\partial \rho_k} (\rho_k \log((1 - \zeta) + \zeta n\rho_k)) = \log((1 - \zeta) + \zeta n\rho_k) + \frac{\zeta n\rho_k}{(1 - \zeta) + \zeta n\rho_k}. \quad (6)$$

for any  $k \in 1 \dots n$ . By setting  $\zeta = 1$ ,

$$\left( \frac{\partial}{\partial \rho_i} - \frac{\partial}{\partial \rho_j} \right) \text{tr}(\rho \log \rho) = \log(\rho_i/\rho_j) = \log(1 + \delta/\rho_j).$$

Because

$$\frac{x}{1+x} \leq \log(1+x) \leq x$$

for all  $x \geq 0$ , we have for any  $\beta \in [0, 1]$  that

$$a \log(1 + \delta/\rho_j) \geq a(1 - \beta) \log(1 + \delta/\rho_j) + a\beta \frac{\delta/\rho_j}{1 + \delta/\rho_j}.$$

This will allow us to deal with the two terms in equation (6) individually.

First, we handle the logarithm term,  $\log((1 - \zeta) + \zeta n\rho_k)$ , by finding  $\zeta$  such that

$$a(1 - \beta) \log(1 + \delta/\rho_j) \geq \log \left( \frac{(1 - \zeta) + \zeta n\rho_j + \zeta n\delta}{(1 - \zeta) + \zeta n\rho_j} \right).$$

We rewrite the right hand side as

$$\log \left( 1 + \frac{\zeta n\delta}{(1 - \zeta) + \zeta n\rho_j} \right).$$

On the left hand side, since  $\rho_j \leq 1/n$ ,  $\delta/\rho_j \geq n\delta$ . We aim to show that

$$a(1 - \beta) \log(1 + n\delta) \geq \log \left( 1 + \frac{\zeta}{(1 - \zeta) + \zeta n\rho_j} n\delta \right),$$

which by Lemma 2.3 is achieved when

$$\frac{\zeta}{(1-\zeta) + \zeta n \rho_j} \leq \frac{a(1-\beta)}{1+n\delta}.$$

We know  $\zeta/((1-\zeta) + \zeta n \rho_j) \leq \zeta/(1-\zeta)$  for any  $n \rho_j \geq 0$ . Hence any

$$\zeta \leq \frac{a(1-\beta)}{1+n\delta + a(1-\beta)} \quad (7)$$

is sufficiently small.

Next, we handle the fraction terms by finding  $\zeta$  such that

$$a\beta \frac{\delta/\rho_j}{1+\delta/\rho_j} \geq \frac{\zeta n(\rho_j + \delta)}{(1-\zeta) + \zeta n(\rho_j + \delta)} - \frac{\zeta n \rho_j}{(1-\zeta) + \zeta n \rho_j}.$$

By Taylor expansion,

$$\begin{aligned} & \frac{\zeta n(\rho_j + \delta)}{(1-\zeta) + \zeta n(\rho_j + \delta)} - \frac{\zeta n \rho_j}{(1-\zeta) + \zeta n \rho_j} \\ &= \frac{\zeta n(\rho_j + \delta)}{(1-\zeta) + \zeta n \rho_j} \sum_{k=0}^{\infty} \left( \frac{-\zeta n \delta}{(1-\zeta) + \zeta n \rho_j} \right)^k - \frac{\zeta n \rho_j}{(1-\zeta) + \zeta n \rho_j}. \end{aligned}$$

Canceling the 0th order term,

$$\begin{aligned} \dots &= \frac{\zeta n \rho_j}{(1-\zeta) + \zeta n \rho_j} \sum_{k=1}^{\infty} \left( \frac{-\zeta n \delta}{(1-\zeta) + \zeta n \rho_j} \right)^k + \frac{\zeta n \delta}{(1-\zeta) + \zeta n \delta} \sum_{k=0}^{\infty} \left( \frac{-\zeta n \delta}{(1-\zeta) + \zeta n \rho_j} \right)^k \\ &= \frac{\zeta n \rho_j}{(1-\zeta) + \zeta n \rho_j} \sum_{k=1}^{\infty} \left( \frac{-\zeta n \delta}{(1-\zeta) + \zeta n \rho_j} \right)^k - \sum_{k=1}^{\infty} \left( \frac{-\zeta n \delta}{(1-\zeta) + \zeta n \rho_j} \right)^k \\ &= \left( \frac{\zeta n \rho_j}{(1-\zeta) + \zeta n \rho_j} - 1 \right) \sum_{k=1}^{\infty} \left( \frac{-\zeta n \delta}{(1-\zeta) + \zeta n \rho_j} \right)^k. \end{aligned}$$

Now to turn this back into the form of a fraction,

$$\begin{aligned} \dots &= \frac{-(1-\zeta)}{(1-\zeta) + \zeta n \rho_j} \sum_{k=1}^{\infty} \left( \frac{-\zeta n \delta}{(1-\zeta) + \zeta n \rho_j} \right)^k \\ &= \frac{(1-\zeta)\zeta n \delta}{((1-\zeta) + \zeta n \rho_j)^2} \sum_{k=0}^{\infty} \left( \frac{-\zeta n \delta}{(1-\zeta) + \zeta n \rho_j} \right)^k \\ &= \frac{(1-\zeta)\zeta n \delta}{((1-\zeta) + \zeta n \rho_j)^2} \frac{1}{1 + \frac{\zeta n \delta}{(1-\zeta) + \zeta n \rho_j}} \\ &= \frac{1}{(1-\zeta) + \zeta n \rho_j} \frac{\zeta n \delta}{1 + \frac{\zeta}{1-\zeta}(\rho_j + \delta)n}. \end{aligned}$$

Since  $0 \leq \rho_j \leq 1/n$ ,

$$\dots \leq \frac{\frac{\zeta}{1-\zeta}n\delta}{1 + \frac{\zeta}{1-\zeta}n\delta}.$$

We must find a  $\zeta$  for which

$$a\beta \log(1 + \delta/\rho_j) \geq a\beta \frac{\delta/\rho_j}{1 + \delta/\rho_j} \geq a\beta \frac{n\delta}{1 + n\delta} \geq \frac{\frac{\zeta}{1-\zeta}n\delta}{1 + \frac{\zeta}{1-\zeta}n\delta}.$$

One can easily check that this is satisfied by

$$\frac{\zeta}{1 + \zeta} \leq \frac{a\beta}{(1 - a\beta)n\delta + 1},$$

which follows from

$$\zeta \leq \frac{a\beta}{(1 - a\beta)n\delta + a\beta + 1}. \quad (8)$$

Finally, any

$$\zeta \leq \min \left\{ \frac{a(1 - \beta)}{1 + n + a(1 - \beta)}, \frac{a\beta}{(1 - a\beta)n + a\beta + 1} \right\}$$

satisfies both inequalities 7 and 8. ■

*Proof of Theorem 1.3.* Let  $n$  be the dimension of  $\rho$ , since  $d$  may be confused with a derivative. The goal is to show that given some  $a \in [0, 1]$  and densities  $\rho, \sigma$  such that  $\rho \succ \sigma$ ,

$$D(\rho \| (1 - \zeta)\hat{1}/n + \zeta\sigma) - (1 - a)D(\rho \| \hat{1}/n) \geq 0,$$

for an appropriate value of  $\zeta \in [0, 1]$ . We apply Lemma 2.2 to replace  $\sigma$  by  $\rho$ . For any states  $\rho$  and  $\omega$ ,

$$D(n\rho \| n\omega) = \text{tr}((n\rho)(\log \rho + \log n - \log \omega - \log n)) = nD(\rho \| \omega).$$

Hence it is sufficient to prove that

$$D(n\rho \| (1 - \zeta)\hat{1} + \zeta n\rho) - (1 - a)D(n\rho \| \hat{1}) \geq 0,$$

which expands as

$$\dots = n \text{tr}(\rho(a \log(n\rho) - \log((1 - \zeta)\hat{1} + \zeta n\rho))) \geq 0.$$

The main insight behind this proof is Lemma 2.4. If  $\rho = \hat{1}/n$ , then both terms are 0, and the proof is trivially complete. If  $\rho \neq \hat{1}/n$ , then the total probability mass above  $\hat{1}/n$  must equal that below  $\hat{1}/n$  to maintain normalization. Hence we may apply the variational argument in Lemma 2.4 to continuously redistribute probability from larger to smaller until  $\rho \rightarrow \hat{1}/n$ . ■

### 3. COMBINING CONDITIONAL EXPECTATIONS

**Lemma 3.1** (Chain Expansion). *Let  $\Phi$  be a quantum channel and  $\mathcal{E}$  be a conditional expectation such that  $\mathcal{E}(\Phi(\rho)) = \mathcal{E}(\rho)$ . Then for any state  $\rho$ ,*

$$D(\rho \| \Phi(\rho)) = D(\rho \| \mathcal{E}(\rho)) + D(\mathcal{E}(\rho) \| \Phi(\rho))$$

*Proof.*

$$\begin{aligned}
& D(\rho\|\mathcal{E}(\rho)) + D(\mathcal{E}(\rho)\|\Phi(\rho)) \\
&= \text{tr}(\rho \log \rho - \rho \log \mathcal{E}(\rho) + \mathcal{E}(\rho) \log \mathcal{E}(\rho) - \mathcal{E}(\rho) \log \Phi(\rho)) \\
&= \text{tr}(\rho \log \rho - \rho \log \mathcal{E}(\rho) + \rho \log \mathcal{E}(\rho) - \rho \log \Phi(\rho)) \\
&= \text{tr}(\rho \log \rho - \rho \log \Phi(\rho)) ,
\end{aligned}$$

where the 1st equality follows from expanding the relative entropy, and the 2nd from the defining property of conditional expectations.  $\blacksquare$

**Lemma 3.2.** *Let  $\mathcal{E}$  be a conditional expectation and  $\Phi$  be a quantum channel. Then*

$$D(\rho\|\mathcal{E}(\rho)) + D(\rho\|\Phi(\rho)) \geq D(\rho\|\mathcal{E}(\Phi(\rho)))$$

*Proof.* By data processing on the 2nd term,

$$D(\rho\|\mathcal{E}(\rho)) + D(\rho\|\Phi(\rho)) \geq D(\rho\|\mathcal{E}(\rho)) + D(\mathcal{E}(\rho)\|\mathcal{E}(\Phi(\rho))) .$$

By Lemma 3.1 and the idempotence of conditional expectations, we obtain the desired result.  $\blacksquare$

**Lemma 3.3.** *Let  $\{(\mathcal{N}_j, \mathcal{E}_j) : j = 1 \dots J \in \mathbb{N}\}$  be a set of  $J$  von Neumann algebras and associated conditional expectations. Let  $\mathcal{E}$  be the conditional expectation to the intersection algebra,  $\mathcal{N} = \bigcap_j \mathcal{N}_j$ . Let  $S = \bigcup_{m \in \mathbb{N}} \{1 \dots J\}^{\otimes m}$  be the set of sequences of indices  $1 \dots k$ . For any  $s \in S$ , let  $\mathcal{E}^s$  denote the composition of conditional expectations  $\mathcal{E}_{j_1} \dots \mathcal{E}_{j_m}$  for  $s = (j_1, \dots, j_m)$ . Let  $\mu : S \rightarrow [0, 1]$  be a probability measure on  $S$  that is non-zero only if no index appears more than  $k$  times. If for all input densities  $\rho$ ,*

$$\sum_{s \in S} \mu(s) \mathcal{E}^s(\rho) = (1 - \zeta) \mathcal{E}(\rho) + \zeta \Phi(\rho)$$

for some channel  $\Phi$ , and

$$D(\rho\|(1 - \zeta) \mathcal{E}(\rho) + \zeta \Phi(\rho)) \geq f(\{\mathcal{N}_j\}) D(\rho\|\mathcal{E}(\rho))$$

for some  $f(\{\mathcal{N}_j\}) > 0$ , then  $\{\mathcal{E}_j\}$  is  $f(\{\mathcal{N}_j\})/k$ -subadditive.

*Proof.* By rewriting the identity as a sum weighted by  $\mu$ ,

$$\sum_j D(\rho\|\mathcal{E}_j(\rho)) = \frac{1}{k} \left( \sum_{s \in S} \mu(s) k_s \right) \sum_j D(\rho\|\mathcal{E}_j(\rho)) ,$$

where  $k = \sum_{s \in S} \mu(s) k_s$ . For each  $s$  such that  $\mu(s) > 0$ , we apply Lemma 3.2 iteratively until we obtain the relative entropy of the sequence  $s$ . Since we have  $k$  copies of each conditional expectation involved and know that no more than  $k$  appear in any sequence, we know that this succeeds. Hence

$$\sum_j D(\rho\|\mathcal{E}_j(\rho)) \geq \frac{1}{k} \sum_{s \in S} \mu(s) D(\rho\|\mathcal{E}^s(\rho)) .$$

By convexity of relative entropy and the assumptions of the Lemma,

$$\sum_j D(\rho\|\mathcal{E}_j(\rho)) \geq \frac{1}{k} D(\rho\|(1 - \zeta) \mathcal{E}(\rho) + \zeta \Phi(\rho)) \geq \frac{f(\{\mathcal{N}_j\})}{k} D(\rho\|\mathcal{E}(\rho)) .$$

■

*Proof of Theorem 1.4.* The gist of this proof is Lemma 3.3 with  $f(\{\mathcal{N}_j\})$  given by Theorem 1.3. First, let us assume that there exists such a  $\mu$  and  $\zeta$  as described in the second part of the Theorem.

First we will consider  $D(\rho\|(1-\zeta)\mathcal{E}(\rho) + \zeta\Phi(\rho))$ . By a result of Uhlmann [15],  $\rho \succ \Phi(\rho)$  for any unital channel  $\Phi$ . If  $\zeta$  is not sufficiently small, we use the fact that for any  $m \in \mathbb{N}$ ,

$$((1-\zeta)\mathcal{E} + \zeta\Phi)^m(\rho) = (1-\zeta^m)\mathcal{E}(\rho) + \zeta^m\Phi^m(\rho)$$

by the idempotence of conditional expectations, and by the assumption that  $\mathcal{E} \circ \Phi = \mathcal{E}$ . To construct  $m$  powers of a convex combination of sequences of conditional expectations, we assume that we have  $((1-\zeta)\mathcal{E} + \zeta\Phi)^{m-1}(\rho)$ , then recursively apply Lemma 3.2 to append subsequences of conditional expectations, and then convexify to reach  $((1-\zeta)\mathcal{E} + \zeta\Phi)^m(\rho)$ . For a sequence of conditional expectations  $\mathcal{E}^s = \mathcal{E}_{j_1} \dots \mathcal{E}_{j_{|s|}}$ , we need at most  $k_s$  copies of each  $D(\rho\|\mathcal{E}_j(\rho))$ , where  $k_s$  is the maximum count of any constituent conditional expectation. Hence

$$\mu(s)k_s \sum_j D(\rho\|\mathcal{E}_j(\rho)) \geq \mu(s)D(\rho\|\mathcal{E}^s(\rho)).$$

We may then repeat this process inductively.

In full generality, we would optimize  $a$ ,  $\beta$ ,  $m$ , and  $\epsilon$ . Here however, we will simplify by choosing  $a = \beta = 1/2$ . Hence we may choose

$$\zeta^m \leq \frac{2}{3d^2 + 5} \leq \min \left\{ \frac{2}{4d + 5}, \frac{2}{3d^2 + 5} \right\},$$

which is satisfied by  $m \geq \lceil \log_\zeta(2/(3d^2 + 5)) \rceil$ . Then

$$D(\rho\|(1-\zeta^m)\hat{1}/d + \zeta^m\Phi^m(\rho)) \geq \frac{1}{2}D(\rho\|\hat{1}/d).$$

We apply Lemma 3.3 with  $\zeta^m$  replacing  $\zeta$ ,  $mk$  replacing  $k$ , and with the above bound.

Finally, we must show that some sufficiently long chain of conditional expectations eventually converges to the intersection algebra, so that such a  $\mu$  and  $\zeta$  as in the Theorem always exist. Let  $s = (1, \dots, J)$ , so that  $\mathcal{E}^s$  is a sequence on all constituent conditional expectations. As a convex combination of unitaries,  $s$  is a contraction. One can easily see that it satisfies properties  $(0.I - 0.IV)$  in [16]. As a result, the  $\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{k=0}^m (\mathcal{E}^s)^k \rightarrow \mathcal{E}$  in 2-norm distance. In finite dimension, this eventually implies a bound on the smallest element, so for some  $m$ , it must yield a convex combination of the complete mixture with some unital channel. ■

**Remark 3.4.** For a bipartite classical-quantum state  $\rho^{XB}$  with classical system  $X$  for which  $\rho = \sum_x p_x \vec{e}_x \otimes \rho_x$ , one can expand  $D(\rho\|\mathcal{E}(\rho)) = \sum_{x \in X} p_x D(\rho_x\|\mathcal{E}(\rho_x))$ . Hence we may extend Theorem 1.4 to include classical extensions.

#### 4. APPLICATION: MERGING MODIFIED LOG-SOBOLEV INEQUALITIES

The main part of this section is to derive MLSI merging using a Suzuki-Trotter decomposition, followed by some examples.

*Proof of Theorem 1.5.* We start by writing  $\Phi^t(\rho) = \Phi^\tau(\rho')$ , where  $\tau < t$ , and  $\rho' = \Phi^{t-\tau}(\rho)$ . Let

$$\text{wsb}(\epsilon, \mathcal{N} \subseteq \mathcal{M}) = 2\epsilon D(\mathcal{M} \parallel \mathcal{N}) + (1 + 2\epsilon)h\left(\frac{\epsilon}{1 + 2\epsilon}\right), \quad (9)$$

as in proposition 3.7 of [17], where  $h$  is the binary entropy,  $\mathcal{N} \subset \mathcal{M}$  is a subalgebra, and  $D(\mathcal{M} \parallel \mathcal{N}) = \sup_\rho D(\mathcal{E}_\mathcal{M}(\rho) \parallel \mathcal{E}_\mathcal{N}(\rho))$ . Rather than depend on dimension as would the Fannes-Audenart bound, this bounds entropy in terms of trace distance and subalgebra index. Let  $\mathcal{N}$  be the fixpoint algebra of  $\Phi^t$  with conditional expectation  $\mathcal{E}$ , and  $\mathcal{N}_j$  the fixpoint algebra of  $\Phi_j^t$ . Let  $\mathcal{M}$  be the original algebra containing  $\rho$ . By the Suzuki-Trotter expansion, the Taylor expansion of an exponential, or by direct commutation,  $\Phi^\tau(\rho) = \Phi_1^\tau \dots \Phi_J^\tau \Phi_0^\tau(\rho) + O((J+1)\tau^2)$ . Let  $\hat{\rho} = \Phi_0^\tau(\rho')$ . Hence

$$D(\Phi^\tau(\hat{\rho}) \parallel \mathcal{E}(\hat{\rho})) \leq D(\Phi_1^\tau \dots \Phi_J^\tau(\hat{\rho}) \parallel \mathcal{E}(\hat{\rho})) + \text{wsb}(O(J\tau^2), \mathcal{N} \subseteq \mathcal{M}). \quad (10)$$

Define  $\gamma_j$  by

$$D(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}) \parallel \mathcal{E}_j(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}))) = \gamma_j D(\Phi^\tau(\hat{\rho}) \parallel \mathcal{E}(\hat{\rho})). \quad (11)$$

Then

$$\begin{aligned} & D(\Phi_1^\tau \dots \Phi_J^\tau(\hat{\rho}) \parallel \mathcal{E}(\hat{\rho})) \\ &= D(\Phi_1^\tau \dots \Phi_J^\tau(\hat{\rho}) \parallel \mathcal{E}_j(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}))) + D(\mathcal{E}_j(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho})) \parallel \mathcal{E}(\hat{\rho})) \\ &\leq (1 - \lambda_j \tau + O(\lambda_j^2 \tau^2)) D(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}) \parallel \mathcal{E}_j(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}))) + D(\mathcal{E}_j(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho})) \parallel \mathcal{E}(\hat{\rho})) \\ &\leq (1 - \lambda_j \gamma_j \tau + O(\lambda_j^2 \tau^2)) (D(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}) \parallel \mathcal{E}_j(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}))) + D(\mathcal{E}_j(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho})) \parallel \mathcal{E}(\hat{\rho}))) \\ &= (1 - \lambda_j \gamma_j \tau + O(\lambda_j^2 \tau^2)) D(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}) \parallel \mathcal{E}(\hat{\rho})), \end{aligned}$$

where the first equality follows from Lemma 3.1 and the first inequality from CLSI of  $\Phi_i^\tau$ . The second inequality follows from equation (11), where we replace the  $-\lambda_j \tau D(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}) \parallel \mathcal{E}_j(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho})))$  subtracted term by  $-\lambda_j \gamma_j \tau D(\Phi^\tau(\hat{\rho}) \parallel \mathcal{E}(\hat{\rho}))$  and then note that this is at least as large as the entire original expression. Iterating, we obtain

$$D(\Phi_1^\tau \dots \Phi_J^\tau(\hat{\rho}) \parallel \mathcal{E}(\hat{\rho})) \leq \left(1 - \sum_{j=1}^J \lambda_j \gamma_j \tau + O\left(\tau^2 \sum_j \lambda_j^2\right) D(\mathcal{M} \parallel \mathcal{N}_j)\right) D(\hat{\rho} \parallel \mathcal{E}(\rho)). \quad (12)$$

Now we must show that the subtracted sum  $\sum_j \lambda_j \gamma_j$  is bounded below. Here we invoke  $\alpha$ -subadditivity. First, note that for any  $j$ ,

$$D(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}) \parallel \mathcal{E}_j(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho}))) \geq D(\hat{\rho} \parallel \mathcal{E}_j(\hat{\rho})) + \text{wsb}(O((J-j)\tau^2), \mathcal{N}_j \subseteq \mathcal{M}).$$

Hence

$$\sum_j D^{\mathcal{N}_j}(\Phi_{j+1}^\tau \dots \Phi_J^\tau(\hat{\rho})) = \sum_j \gamma_j D(\hat{\rho} \parallel \mathcal{E}(\hat{\rho})) \geq \alpha D(\hat{\rho} \parallel \mathcal{E}(\hat{\rho})) + \sum_{j=1}^J \text{wsb}(O((J-j)\tau^2), \mathcal{N}_j \subseteq \mathcal{M}).$$

This implies that  $\sum_j \gamma_j \geq \alpha$ .  $D(\hat{\rho} \parallel \mathcal{E}(\rho)) \geq D(\Phi^t(\rho) \parallel \mathcal{E}(\rho))$  by data processing, so either the latter is zero, or the former is at least some  $\epsilon$  that is independent of  $\tau$ . If  $D(\Phi^t(\rho) \parallel \mathcal{E}(\rho)) = 0$ ,

then the Theorem is trivially complete. Otherwise, returning to equations (12) and (10),

$$D(\Phi^\tau(\hat{\rho})\|\mathcal{E}(\hat{\rho})) \leq \left(1 - \alpha\tau \min_j\{\lambda_j\} + \text{err}\right)D(\hat{\rho}\|\mathcal{E}(\hat{\rho})). \quad (13)$$

if we let

$$\begin{aligned} \text{err} &= \frac{\text{wsb}(O(J\tau^2), \mathcal{N} \subseteq \mathcal{M})}{\epsilon} \\ &+ \sum_{j=1}^J \left( \frac{\text{wsb}(O((J-j)\tau^2), \mathcal{N}_j \subseteq \mathcal{M})}{\epsilon} + O(\lambda_j^2\tau^2)D(\mathcal{M}\|\mathcal{N}_j) \right). \end{aligned} \quad (14)$$

Furthermore,

$$D(\Phi_0^\tau(\rho')\|\mathcal{E}(\rho')) \leq D(\rho'\|\mathcal{E}(\rho')), \quad (15)$$

as the left hand side follows from data processing on the right hand side. Since  $\Phi^t = (\Phi^\tau)^{t/\tau}$ ,

$$D(\Phi^t(\rho)\|\mathcal{E}(\rho)) \leq (1 - \alpha \min_j\{\lambda_j\}\tau + \text{err})^{t/\tau} D(\rho\|\mathcal{E}(\rho)). \quad (16)$$

Since  $\text{err} \leq O(J \log J \tau^2 \log \tau (D(\mathcal{M}\|\mathcal{N}) + \sum_j \lambda_j^2 \log \lambda_j D(\mathcal{M}\|\mathcal{N}_j)))$ ,

$$\lim_{\tau \rightarrow 0} (1 - \alpha \min_j\{\lambda_j\}\tau + \text{err})^{t/\tau} = e^{-\alpha \min_j\{\lambda_j\}t}.$$

This completes the Theorem. ■

**Corollary 4.1** (CLSI Merge). *Let  $\{\Phi_j^t\}$  and  $\Phi^t$  be as in Theorem 1.5, with  $\{\mathcal{E}_j\}$  being completely  $\alpha$ -subadditive, and each  $\Phi_j^t$  having  $\lambda_j$ -CLSI. Then  $\Phi^t$  has  $\alpha \times \min_j\{\lambda_j\}$ -CLSI.*

*Proof.* Apply Theorem 1.5 for any extension  $\{\hat{1}^C \otimes \Phi_j^t\}$  and  $\{\hat{1}^C \otimes \mathcal{E}_j\}$ . ■

**Corollary 4.2** (Commuting Hypercube M/CLSI Merge). *Let  $\{\Phi_j^t\}$  be a set of QMSs as in Theorem 1.5 that form a commuting hypercube, in that  $[\mathcal{E}_j, \mathcal{E}_k] = 0$  for any  $j, k \in 1 \dots J$ , and  $\mathcal{E}_{\sigma(J) \dots \mathcal{E}_{\sigma(1)}} = \mathcal{E}$  for any permutation  $\sigma \in S_J$ . Then the set is 1-subadditive with M/CLSI merge.*

*Proof.* Iteratively applying Lemma 3.2, we obtain

$$\sum_{j=1}^J D(\rho\|\mathcal{E}_j(\rho)) \geq D(\rho\|\mathcal{E}_J \dots \mathcal{E}_1(\rho)) = D(\rho\|\mathcal{E}(\rho)). \quad (17)$$

Hence we apply Theorem 1.5 or Corollary 4.1. Note that  $\{1^C \otimes \mathcal{E}_j\}$  automatically form a commuting hypercube for all extensions  $C$ . ■

**Corollary 4.3** (M/CLSI Addition). *Let  $\{\Phi_j^t\}$  be a set of QMSs as in Theorem 1.5 (resp Corollary 4.1), but such that  $\mathcal{E}_j = \mathcal{E}$  for each  $j$ . Then  $\Phi^t$  has  $J \min_j\{\lambda_j\}$ -MLSI (resp CLSI).*

*Proof.* This follows almost immediately from  $J$ -subadditivity, in that

$$\sum_{j=1}^J D(\rho\|\mathcal{E}(\rho)) = JD(\rho\|\mathcal{E}(\rho)). \quad (18)$$

Hence we apply Theorem 1.5 (resp Corollary 4.1). ■

**Example 4.4** (Mutually Unbiased Basis Decay). *Let  $X, Z$  be the Pauli matrices, and  $\langle X \rangle, \langle Z \rangle$  their corresponding generated subalgebras. It is well-known that the subgroup  $\Phi_{\langle X \rangle}^t$  generated by  $\mathcal{L}_{\langle X \rangle}(\rho) = \rho - X\rho X$  has 1-CLSI, and so does  $\Phi_{\langle Z \rangle}^t$ . Here we have*

$$\Phi_{\langle X \rangle}^t(\rho) = e^{-t\mathcal{L}_{\langle X \rangle}}(\rho) = e^{-t}\rho + (1 - e^{-t})X\rho X.$$

Hence

$$\begin{aligned} \Phi_{\langle X \rangle}^t(\mathcal{E}_{\langle Z \rangle}(\rho)) &= \frac{1}{2}(e^{-t}\rho + (1 - e^{-t})X\rho X + e^{-t}Z\rho Z + (1 - e^{-t})XZ\rho ZX) \\ &= \frac{1}{2}(e^{-t}\rho + (1 - e^{-t})X\rho X + e^{-t}Z\rho Z + (1 - e^{-t})ZX\rho XZ) \\ &= \mathcal{E}_{\langle Z \rangle}(\Phi_{\langle X \rangle}^t(\rho)), \end{aligned}$$

and similarly,  $[\Phi_{\langle Z \rangle}^t, \mathcal{E}_{\langle X \rangle}] = 0$ . As these form a commuting square, (Theorem/Corollary) ?? implies that the semigroup generated by  $\mathcal{L} = \mathcal{L}_{\langle X \rangle} + \mathcal{L}_{\langle Z \rangle}$  has 1-CLSI.

This example generalizes to any pair of mutually unbiased bases  $\mathcal{X}, \mathcal{Z}$  in finite dimension.

**Example 4.5** (Tilted Incompatible Basis Decay). *Let  $\mathcal{X}$  and  $\mathcal{Z}$  now denote a pair of generalized bases in dimension  $d$ , not necessarily mutually unbiased. Let  $\{|i_X\rangle : i = 1\dots d\}$  and  $\{|i_Z\rangle : i = 1\dots d\}$  index these bases, and  $\inf_{i,j} |\langle i_X | j_Z \rangle|^2 = \epsilon$  for some  $0 \leq \epsilon \leq 1/d$ . Because a collapse to either basis is completely dephasing in that basis, a subsequent collapse in the other leaves any input state in a classical mixture with minimal probability mass no less than  $\epsilon$ . We may write each collapse as a convex combination of unitaries of the form  $|i\rangle \rightarrow \exp(2\pi i k/d)|i\rangle$  for each  $k \in 1\dots d$ . Hence it depolarizes with probability  $\epsilon$ , so  $\mathcal{E}_{\mathcal{X}} \circ \mathcal{E}_{\mathcal{Z}}(\rho) = \epsilon \hat{1}/d + (1 - \epsilon)\Phi(\rho)$  for some unital  $\Phi$ . By Theorem 1.4,*

$$D(\rho \| \mathcal{E}_{\mathcal{X}}(\rho)) + D(\rho \| \mathcal{E}_{\mathcal{Z}}(\rho)) \geq \frac{D(\rho \| \hat{1}/d)}{4(\log_{1-\epsilon}(2/(3d+5)) + 1)}.$$

Expanding the relative entropies, we find this is equivalent to

$$H(\mathcal{X})_\rho + H(\mathcal{Z})_\rho \geq \left(2 - \frac{1}{4(\lceil \log_{1-\epsilon}(2/(3d+5)) \rceil)}\right) H(\rho) + \frac{\log d}{4(\lceil \log_{1-\epsilon}(2/(3d+5)) \rceil)}.$$

An uncertainty relation in the conventional form would be

$$H(\mathcal{X})_\rho + H(\mathcal{Z})_\rho \geq H(\rho) + \log d - c$$

for some constant  $c > 0$ , as required for non-commuting conditional expectations as shown in [5]. When  $\rho \approx \hat{1}/d$ , the right hand side of the conventional uncertainty relation becomes negative, and the bound becomes trivial. In contrast, adjusted subadditivity still gives a positive, non-trivial bound on the sum of basis entropies.

Furthermore, let  $\Phi_{\mathcal{X}}^t$  and  $\Phi_{\mathcal{Z}}^t$  be quantum Markov semigroups with  $\beta$ -MLSI for some  $\beta > 0$  and respective Lindbladian generators  $\mathcal{L}_{\mathcal{X}}, \mathcal{L}_{\mathcal{Z}}$ . Then the channel given by  $\Phi = e^{-(\mathcal{L}_{\mathcal{X}} + \mathcal{L}_{\mathcal{Z}})t}$  has  $1/(4(\lceil \log_{1-\epsilon}(2/(3d+5)) \rceil))$ -MLSI.

## 5. CONCLUSIONS AND OUTLOOK

The primary limitations of these results are dimension-dependence and restriction to algebras with trivial intersection. In infinite dimensions, we might expect the bound to be trivial in cases for which the intersection of all constituent conditional expectations is completely depolarizing, and indeed, it's not clear that such a case would even exist in type III von Neumann algebras. It is however plausible that the dimension appearing in Theorem 1.4 actually arises from  $D(\mathbb{B}(\mathcal{H})\|\mathbb{C}1)$ , the maximal asymmetry/index as derived in [17]. If so, we might generalize:

**Conjecture 5.1** (Strong Adjusted Subadditivity (SASA)). *Let  $\{(\mathcal{N}_j, \mathcal{E}_j) : j = 1 \dots J \in \mathbb{N}\}$  be a set of  $J$  von Neumann algebras within a larger algebra  $\mathcal{M}$  and associated conditional expectations. Then  $\{\mathcal{E}_j\}$  is  $\alpha$ -subadditive with an  $\alpha$  that depends on the index  $D(\mathcal{M}\|\mathcal{N})$ , but not explicitly on the dimension of  $\mathcal{M}$ .*

The main challenge, as seen in 3.3, is to prove an analog of Theorem 1.3 that replaces  $\hat{1}/d$  by an arbitrary conditional expectation. We might also generalize to a measure on a continuous set of conditional expectations, but that is probably more complicated to formulate than would be appropriate for these concluding remarks.

Between Theorem 1.4 and Conjecture 5.1, we might also consider when the intersection algebra is given by  $\mathcal{N} = \mathbb{C}1 \otimes \mathbb{B}(B)$ , where  $B$  is an auxiliary system that is untouched by any of the conditional expectations in question. This would be an adjusted subadditivity with memory, and with Corollary 4.1, it would create a new method to build up CLSI on complicated semigroups formed from simpler constituents. The main challenge to proving Theorem 1.3 in this case is that since  $\hat{1}/A \otimes \rho^B$  and  $\rho^{AB}$  need not commute, we can't obviously reduce the problem to one of estimating logarithms of commuting vectors. It would not be surprising if all the main components of Theorem 1.3 have matrix analogs, but the proof strategy might differ.

Strong Adjusted Subadditivity would be sufficiently powerful to justify continued investigation. Even its refutation could suggest major differences between where MLSI and CLSI would probably hold, and resultingly, where auxiliary systems might be effective in slowing decoherence.

Though we do not show SASA here, we have shown that such a decay inequality generally holds for algebras with trivial intersection. What is most surprising is that ASA does not rely on any sort of nearness to a commuting square. It is not an approximate tensorization or factorization result. Rather, ASA appears to be in a class of entropy inequalities that do not follow directly from strong subadditivity. Its connection to decoherence and decay estimates motivate inequalities of this type.

## 6. ACKNOWLEDGEMENTS

I would like to thank Marius Junge for early feedback on these results. I would also like to acknowledge interactions with Li Gao as helping to motivate this project.

## REFERENCES

- [1] Elliott H. Lieb and Mary Beth Ruskai. Proof of the strong subadditivity of quantummechanical entropy. *Journal of Mathematical Physics*, 14(12):1938–1941, December 1973.

- [2] Mark M. Wilde. *Quantum Information Theory*. Cambridge University Press, Cambridge, UK. ; New York, 1 edition edition, June 2013.
- [3] Elena Caceres, Arnab Kundu, Juan F. Pedraza, and Walter Tangarife. Strong subadditivity, null energy condition and charged black holes. *Journal of High Energy Physics*, 2014(1):84, January 2014.
- [4] Dénes Petz. On certain properties of the relative entropy of states of operator algebras. *Mathematische Zeitschrift*, 206(1):351–361, January 1991.
- [5] Li Gao, Marius Junge, and Nicholas LaRacuente. Unifying Entanglement with Uncertainty via Symmetries of Observable Algebras. *arXiv:1710.10038 [quant-ph]*, October 2017. arXiv: 1710.10038.
- [6] Filippo Cesi. Quasi-factorization of the entropy and logarithmic Sobolev inequalities for Gibbs random fields. *Probability Theory and Related Fields*, 120(4):569–584, August 2001.
- [7] Ivan Bardet, Angela Capel, and Cambyse Rouzé. Private communications. 2019.
- [8] Fernando G. S. L. Brando, Aram W. Harrow, and Michał Horodecki. Local Random Quantum Circuits are Approximate Polynomial-Designs. *Communications in Mathematical Physics*, 346(2):397–434, September 2016.
- [9] Xiao Chen and Tianci Zhou. Operator scrambling and quantum chaos. *arXiv:1804.08655 [cond-mat, physics:hep-th, physics:quant-ph]*, April 2018. arXiv: 1804.08655.
- [10] Elizabeth Crosson and Aram W. Harrow. Rapid mixing of path integral Monte Carlo for 1d stoquastic Hamiltonians. *arXiv:1812.02144 [cond-mat, physics:quant-ph]*, December 2018. arXiv: 1812.02144.
- [11] Koenraad M. R. Audenaert. Telescopic Relative Entropy. In Dave Bacon, Miguel Martin-Delgado, and Martin Roetteler, editors, *Theory of Quantum Computation, Communication, and Cryptography*, Lecture Notes in Computer Science, pages 39–52, Berlin, Heidelberg, 2014. Springer.
- [12] Michael J. Kastoryano and Kristan Temme. Quantum logarithmic Sobolev inequalities and rapid mixing. *Journal of Mathematical Physics*, 54(5):052202, May 2013.
- [13] Ivan Bardet. Estimating the decoherence time using non-commutative Functional Inequalities. *arXiv:1710.01039 [math-ph, physics:quant-ph]*, October 2017. arXiv: 1710.01039.
- [14] Li Gao, Marius Junge, and Nicolas LaRacuente. Fisher Information and Logarithmic Sobolev Inequality for Matrix Valued Functions. *arXiv:1807.08838 [quant-ph]*, July 2018. arXiv: 1807.08838.
- [15] A. Uhlmann. Relative entropy and the Wigner-Yanase-Dyson-Lieb concavity in an interpolation theory. *Communications in Mathematical Physics*, 54(1):21–32, 1977.
- [16] Marius Junge and Quanhua Xu. Noncommutative Maximal Ergodic Theorems. *Journal of the American Mathematical Society*, 20(2):385–439, 2007.
- [17] Li Gao, Marius Junge, and Nicholas LaRacuente. Relative entropy for von Neumann subalgebras. *arXiv:1909.01906 [math]*, September 2019. arXiv: 1909.01906.