

THE STOCHASTIC KLAUSMEIER SYSTEM AND A STOCHASTIC SCHAUDER-TYCHONOFF TYPE THEOREM

ERIKA HAUSENBLAS AND JONAS M. TÖLLE

ABSTRACT. We investigate the existence of a pair of nonnegative solutions to the stochastic system of advection-diffusion equations proposed by Klausmeier with Gaussian multiplicative noise. The proof of existence is based upon a stochastic version of the Schauder-Tychonoff fixed point theorem, which is also proved here.

1. INTRODUCTION

Pattern formation at the ecosystem level is a rapidly growing area of spatial ecology. Theoretical models are a widely used tool for studying e.g. banded vegetation patterns. One important model is the system of advection-diffusion equations proposed by Klausmeier [16]. This model for vegetation dynamics in semi-deserted areas is based on the water-redistribution hypothesis, using the idea that rainwater in dry regions is eventually infiltrated into the ground. Instead, water mostly runs off downhill towards the next patch of vegetation. The soil in such regions of the world as Australia, Africa, and Southwestern North America is prone to nonlocality of water uptake due to the semiarid environment. Studies of the properties of the system and further developments can be found in e.g. [15, 26–28, 35].

The Klausmeier system is a generalization of the so-called Gray-Scott system [13] (see also [25, 29] for earlier accounts employing similar models) which already exhibits effects similar to Turing patterns [9, 34], see for instance the discussion in [36]. We refer the reader to [18, 22, 23] for further reading on pattern formation in biology.

The underlying mathematics of this model is given by a pair of solutions (u, v) to a partial differential equation system coupled by a nonlinearity. The function u represents the surface water content and v represents the biomass density of the plants. In order to model the spread of water on a terrain without a specific preference for the direction in which the water flows, the original models were extended by replacing the diffusion operator by a nonlinear porous media operator, which represents the situation that the ground is partially filled by interconnected pores conveying fluid under an applied pressure gradient.

To this end, let $\mathcal{O} \subset \mathbb{R}^d$ be a bounded domain, $d = 1, 2, 3$, having C^∞ boundary or $\mathcal{O} = [0, 1]^d$ with periodic or Neumann boundary conditions.

Date: December 3, 2019.

2010 Mathematics Subject Classification. Primary 60H15; secondary 35K57, 37N25, 60G57, 76S05, 92C15.

Key words and phrases. stochastic Klausmeier system; stochastic Schauder-Tychonoff type theorem; pattern formation in ecology; nonlinear stochastic partial differential equation; flows in porous media; nonnegative solution.

The research was partially funded by FWF (Austrian Science Foundation) grant P 28010 “Mathematische Analyse von Flüssigkeitskristallen mit stochastischer Störung”.

We shall look for solutions to the following problem with $\Phi(x) := x^{[\gamma]} := |x|^{\gamma-1}x$, $\gamma > 1$,

$$(1.1) \quad \begin{cases} \dot{u}(t) &= r_u \Delta \Phi(u(t)) - \kappa u(t) v^2(t) + k - f u(t), & t > 0, \\ u(0) &= u_0, \\ \dot{v}(t) &= r_v \Delta v(t) + u(t) v^2(t) - g v(t), & t > 0, \\ v(0) &= v_0, \end{cases}$$

with Neumann (or periodic, if \mathcal{O} is a torus) boundary conditions and initial conditions $u(0) = u_0$ and $v(0) = v_0$. Moreover, r_u , r_v , f , κ , k , and g are positive constants.

The deterministic or macroscopic model is derived from the limiting behavior of interacting diffusions — the so called microscopic model. Applying the strong law of large numbers and passing from the microscopic to the macroscopic equation one is neglecting the random fluctuations. In order to get a more realistic model, it is necessary to add noise, which represents the randomness of the natural environment or the fluctuation of parameters in the model. Our aim is to investigate the system (1.1) driven by a multiplicative infinite dimensional Wiener process. Under suitable regularity assumptions on the initial data (u_0, v_0) , on γ and on the perturbation by noise which we specify below, we find that there exists a nonnegative solution to the system (1.1) in dimensions $d = 1, 2, 3$, see our main result¹ Theorem 3.7.

See [8] for a recent work proving existence of martingale solutions to stochastic cross-diffusion systems, however, their approach is using another methods and is not covering the porous media case.

Due to the nonlinearities standard methods for showing existence and uniqueness of solutions to stochastic partial differential equations cannot be applied directly. Hence we are using the stochastic Schauder-Tychonoff type Theorem 2.1 which we present in Section 2. In the subsequent section, i.e. Section 3, we apply the Schauder-Tychonoff type theorem to show the existence of a martingale solution to the stochastic counterpart (2.1) of the system (1.1). In Section 4, we prove several technical propositions that are need for the main result in Section 3.

2. A SCHAUDER-TYCHONOFF TYPE THEOREM

Our main motivation to prove a probabilistic Schauder-Tychonoff type fixed point theorem² was to show existence of a solution to the stochastic counterpart of the system (1.1). Roughly spoken, we are adding multiplicative noise to the reduced system, i.e., we are considering the following system

$$(2.1) \quad \begin{cases} \dot{u}(t) &= r_u \Delta \Phi(u(t)) - \kappa u(t) v^2(t) + u(t) \xi_1(t), & t > 0, \\ \dot{v}(t) &= r_v \Delta v(t) + u(t) v^2(t) + v(t) \xi_2(t), & t > 0. \end{cases}$$

Here, ξ_1 and ξ_2 denotes some random Gaussian noise specified later on.

The challenging problem in the system given by (1.1), respective the noisy system (2.1), is the nonlinearity appearing once with a negative sign and once with a positive sign. The nonlinearity is not of variational structure, such that energy methods are not available for

¹In fact, we prove existence of nonnegative solutions to the noisy systems (3.4)–(3.5) (Itô noise), (3.1)–(3.2) (Stratonovich noise) respectively.

²See [12] for the classical Schauder and Schauder-Tychonoff fixed point theorem(s).

the analysis, and, in particular, the maximum principle does not work. To overcome this obstacle, we use a Schauder-Tychonoff type fixed point argument, which is the main result of this section. See [21] for a previous work using the classical Schauder theorem for stochastic evolution equations with fractional Gaussian noise.

Let us fix some notation. Let H be a separable Hilbert space, let $Q \in L(H)$ be a nonnegative definite trace class (covariance) operator. Let U_1 be a Banach space. Let $\mathcal{O} \subset \mathbb{R}^d$ be open, $d \geq 1$. Let $\mathbb{X} \subset \{\eta : [0, T] \rightarrow \mathcal{D}'(\mathcal{O})\}$ be a Banach function space³, let $\mathbb{X}' \subset \{\eta : [0, T] \rightarrow \mathcal{D}'(\mathcal{O})\}$ be a reflexive Banach function space embedded compactly into \mathbb{X} . Let $\mathfrak{A} = (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ be a filtered probability space with filtration $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$ satisfying the usual conditions. For $m \geq 1$, define the collection of processes

$$(2.2) \quad \mathcal{M}_{\mathfrak{A}}^m(\mathbb{X}) := \left\{ \xi : \Omega \times [0, T] \rightarrow \mathcal{D}'(\mathcal{O}) : \right. \\ \left. \xi \text{ is } \mathbb{F}\text{-progressively measurable and } \mathbb{E}|\xi|_{\mathbb{X}}^m < \infty \right\}$$

equipped with the semi-norm

$$|\xi|_{\mathcal{M}_{\mathfrak{A}}^m(\mathbb{X})} := (\mathbb{E}|\xi|_{\mathbb{X}}^m)^{1/m}, \quad \xi \in \mathcal{M}_{\mathfrak{A}}^m(\mathbb{X}).$$

We shall assume that $\mathcal{M}_{\mathfrak{A}}^m(\mathbb{X})$ is a reflexive Banach space with respect to the above semi-norm for $m > 1$, which is the case in all of the applications in this work.

Theorem 2.1. *Let H be a Hilbert space, $Q \in L(H)$ such that Q is nonnegative definite and of trace class, let U_1 be a Banach space, and let us assume that we have a compact embedding $\mathbb{X}' \hookrightarrow \mathbb{X}$ as above. Let $m > 1$. Suppose that for any filtered probability space $\mathfrak{A} = (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ and for any Q -Wiener process⁴ W with values in H that is modeled on \mathfrak{A} , we are given a convex closed nonempty subset⁵ $\mathcal{X}(\mathfrak{A})$ of $\mathcal{M}_{\mathfrak{A}}^m := \mathcal{M}_{\mathfrak{A}}^m(\mathbb{X})$ and an operator*

$$\mathcal{V}_{\mathfrak{A}, W} : \mathcal{M}_{\mathfrak{A}}^m \rightarrow \mathcal{M}_{\mathfrak{A}}^m$$

with the following properties:

- (i) $\mathcal{V}_{\mathfrak{A}, W}(\mathcal{X}(\mathfrak{A})) \subset \mathcal{X}(\mathfrak{A})$,
- (ii) the restriction $\mathcal{V}_{\mathfrak{A}, W} \upharpoonright_{\mathcal{X}(\mathfrak{A})}$ is continuous,
- (iii) there exist $R > 0$, $m_0 \geq m$ such that

$$\mathbb{E}|\mathcal{V}_{\mathfrak{A}, W}(v)|_{\mathbb{X}'}^{m_0} \leq R \quad \forall v \in \mathcal{X}(\mathfrak{A}),$$

- (iv) $\mathcal{V}_{\mathfrak{A}, W}(\mathcal{X}(\mathfrak{A})) \subset \mathbb{D}([0, T]; U_1)$ \mathbb{P} -a.s.⁶

Then there exists a filtered probability space $\tilde{\mathfrak{A}} = (\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}})$ (that satisfies the usual conditions) together with a Q -Wiener process \tilde{W} modeled on $\tilde{\mathfrak{A}}$ and an element $\tilde{v} \in \mathcal{M}_{\tilde{\mathfrak{A}}}^m$ such that for all $t \in [0, T]$, $\tilde{\mathbb{P}}$ -a.s.

$$\mathcal{V}_{\tilde{\mathfrak{A}}, \tilde{W}}(\tilde{v})(t) = \tilde{v}(t).$$

Proof. We shall partially follow the proof in [6]. Fix \mathfrak{A} and W and fix $v^{(0)} \in \mathcal{X}(\mathfrak{A})$.

³Here, $\mathcal{D}'(\mathcal{O})$ denotes the space of Schwartz distributions on \mathcal{O} , that is, the topological dual space of smooth functions with compact support $\mathcal{D}(\mathcal{O}) = C_0^\infty(\mathcal{O})$.

⁴See e.g. [7] for this notion.

⁵Here, the notation $\mathcal{X}(\mathfrak{A})$ means that $\text{Law}(u) = \text{Law}(\tilde{u})$ on \mathbb{X} for $u \in \mathcal{X}(\mathfrak{A})$ and $\tilde{u} \in \mathcal{M}_{\mathfrak{A}}^m$ implies $\tilde{u} \in \mathcal{X}(\tilde{\mathfrak{A}})$.

⁶Here, $\mathbb{D}([0, T]; U_1)$ denotes the Skorokhod space of càdlàg paths in U_1 .

Step I: For $n \in \mathbb{N}$, define recursively $v^{(n)} := \mathcal{V}_{\mathfrak{A}, W}(v^{(n-1)})$. Let $W^{(n)} := W$ for every $n \in \mathbb{N}$. We claim that the laws of the sequence $(v^{(n)}, W^{(n)})$, $n \in \mathbb{N}$ are tight on $\mathbb{X} \times \mathbb{D}([0, T]; U_1)$. By compactness of the embedding $\mathbb{X}' \hookrightarrow \mathbb{X}$, and by reflexivity of \mathbb{X}' , it follows by standard arguments that

$$K_R := \{\xi \in \mathbb{X} : |\xi|_{\mathbb{X}'}^{m_0} \leq R\}$$

is compact in \mathbb{X} . By (i), $v^{(n)} \in \mathcal{X}(\mathfrak{A})$ for every $n \in \mathbb{N}$, and thus it follows by (iii) that $v^{(n)} \in K_R$ for any $n \in \mathbb{N}$. Therefore, the laws of the sequence $\{v^{(n)}\}_{n \in \mathbb{N}}$ are tight on \mathbb{X} . By Ulam's theorem, the single measure $\text{Law}(W) = \text{Law}(W^{(n)})$ is tight on the Polish space $C([0, T]; H)$ and thus is the "constant sequence" $\{\text{Law}(W^{(n)})\}_{n \in \mathbb{N}}$. Since the marginal distributions of $\rho_n := \text{Law}(v^{(n)}, W^{(n)}) = \text{Law}(v^{(n)}, W)$ are tight, so is $\{\rho_n\}_{n \in \mathbb{N}}$.

Step II: By Step I and Prokhorov's theorem, we that there exists a (non-reabeled) subsequence $\{\rho_n\}_{n \in \mathbb{N}}$ and a Borel probability law ρ^* on $\mathbb{X} \times C([0, T]; H)$ such that

$$\rho_n \rightharpoonup \rho^*$$

as $n \rightarrow \infty$ weakly in the sense of probability measures. Next, by the Skorokhod lemma, there exists a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, a sequence of $\mathbb{X} \times C([0, T]; H)$ -valued random variables $\{(\tilde{v}^{(n)}, \tilde{W}^{(n)})\}_{n \in \mathbb{N}}$ and (\tilde{v}, \tilde{W}) such that

$$\text{Law}(\tilde{v}^{(n)}, \tilde{W}^{(n)}) = \rho_n, \quad n \in \mathbb{N}, \quad \text{Law}(\tilde{v}, \tilde{W}) = \rho^*,$$

and such that

$$(\tilde{v}^{(n)}, \tilde{W}^{(n)}) \rightarrow (\tilde{v}, \tilde{W}) \quad \tilde{\mathbb{P}}\text{-a.s.}$$

on \mathbb{X} as $n \rightarrow \infty$. We note that the structure of the random variables is preserved by [6, Theorem C.1], see also Step IV below.

Step III: Clearly, by (iii), we can conclude from the application of the Skorokhod lemma in Step II that

$$\mathbb{E}|v^{(n)}|_{\mathbb{X}}^r = \tilde{\mathbb{E}}|\tilde{v}^{(n)}|_{\mathbb{X}}^r,$$

for any $r \in [1, m_0]$, so that we get by (iii) that

$$\sup_n \tilde{\mathbb{E}}|\tilde{v}^{(n)}|_{\mathbb{X}}^{m_0} \leq CR,$$

where $C > 0$ is a constant such that $|\cdot|_{\mathbb{X}} \leq C|\cdot|_{\mathbb{X}'}$.

Hence, we know that $\{|\tilde{v}^{(n)}|_{\mathbb{X}}^r\}$ is uniformly integrable for any $r \in (1, m_0]$ w.r.t. the probability measure $\tilde{\mathbb{P}}$. By Step II, $\tilde{v}^{(n)} \rightarrow \tilde{v}$ $\tilde{\mathbb{P}}$ -a.s., so we get by the Vitali convergence theorem that

$$(2.3) \quad \lim_{n \rightarrow \infty} \tilde{\mathbb{E}} \left| \tilde{v}^{(n)} - \tilde{v} \right|_{\mathbb{X}}^r = 0$$

for any $r \in (1, m_0]$.

Step IV: Let $\tilde{\mathbb{F}} = (\tilde{\mathcal{F}}_t)_{t \geq 0}$ be the filtration defined for $t \in [0, T]$ by

$$(2.4) \quad \tilde{\mathcal{F}}_t := \sigma \left(\sigma \left\{ \tilde{W}(s), \tilde{v}^{(n)}(s), \tilde{v}(s) : n \in \mathbb{N}, 0 \leq s \leq t \right\} \cup \mathcal{N} \right),$$

where \mathcal{N} denotes the σ -ideal of null sets in $\tilde{\mathcal{F}}$. Since $W^{(n)} = W$, it is easy to show that the filtration obtained by replacing \tilde{W} in (2.4) by $\tilde{W}^{(n)}$ is equal to $\tilde{\mathbb{F}}$. Set

$\tilde{\mathfrak{A}} := (\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}})$. Now, by arguments as in the proof of [6, Theorem C.1], we can show that

- For every $n \in \mathbb{N}$, $\tilde{W}^{(n)}$ is an H -valued Wiener process over $\tilde{\mathfrak{A}}$ with covariance Q ;
- \tilde{W} is an H -valued Wiener process over $\tilde{\mathfrak{A}}$ with covariance Q ;
- for every $n \in \mathbb{N}$, $\tilde{v}^{(n)}$ is an $\tilde{\mathbb{F}}$ -progressively measurable process;
- \tilde{v} is an $\tilde{\mathbb{F}}$ -progressively measurable process.

Step V: Fix $\varepsilon > 0$. Set, for simplicity, $\tilde{\mathcal{V}} := \mathcal{V}_{\tilde{\mathfrak{A}}, \tilde{W}}$. We claim that

$$\tilde{\mathbb{E}} \left| \tilde{\mathcal{V}}(\tilde{v}) - \tilde{v} \right|_{\mathbb{X}} \leq \varepsilon.$$

Let $n \in \mathbb{N}$, $n \geq 2$. By Step III, we get that $\tilde{v}^{(n)} \rightharpoonup \tilde{v}$ weakly in the reflexive Banach space $\mathcal{M}_{\tilde{\mathfrak{A}}}^m$. Hence, as $\mathcal{X}(\tilde{\mathfrak{A}})$ is closed and convex, we get by Mazur's lemma that $\mathcal{X}(\tilde{\mathfrak{A}})$ is weakly closed and thus $\tilde{v} \in \mathcal{X}(\tilde{\mathfrak{A}})$. By (ii), $\tilde{\mathcal{V}}$ is continuous on $\mathcal{X}(\tilde{\mathfrak{A}})$ and hence there exists $\delta = \delta(\varepsilon) > 0$ such that

$$\tilde{\mathbb{E}} \left| \tilde{\mathcal{V}}(\tilde{v}) - \tilde{\mathcal{V}}(\tilde{v}^{(n-1)}) \right|_{\mathbb{X}}^m \leq \frac{\varepsilon}{2},$$

whenever $\tilde{\mathbb{E}} |\tilde{v} - \tilde{v}^{(n-1)}|_{\mathbb{X}}^m < \delta$, which is the case by Step III (and the assumption $m \leq m_0$) for some large n . Also, by construction,

$$\tilde{\mathbb{E}} \left| \tilde{\mathcal{V}}(\tilde{v}^{(n-1)}) - \tilde{v}^{(n)} \right|_{\mathbb{X}} = \mathbb{E} \left| \mathcal{V}(v^{(n-1)}) - v^{(n)} \right|_{\mathbb{X}} = 0.$$

Furthermore, by (2.3),

$$\tilde{\mathbb{E}} \left| \tilde{v}^{(n)} - \tilde{v} \right|_{\mathbb{X}} \leq \frac{\varepsilon}{2}$$

for some large n . Altogether, if we write

$$\tilde{\mathcal{V}}(\tilde{v}) - \tilde{v} = \tilde{\mathcal{V}}(\tilde{v}) - \tilde{\mathcal{V}}(\tilde{v}^{(n-1)}) + \tilde{\mathcal{V}}(\tilde{v}^{(n-1)}) - \tilde{v}^{(n)} + \tilde{v}^{(n)} - \tilde{v}.$$

we can easily complete the proof of the claim. As a consequence,

$$\tilde{\mathcal{V}}(\tilde{v}) = \tilde{v}, \quad \tilde{\mathbb{P}}\text{-a.s.}$$

As seen above, $\tilde{v} \in \mathcal{X}(\tilde{\mathfrak{A}})$, so that by (iv), $\tilde{\mathcal{V}}(\tilde{v}) \in \mathbb{D}([0, T]; U_1)$, and therefore $\tilde{v} \in \mathbb{D}([0, T]; U_1)$ $\tilde{\mathbb{P}}$ -a.s. Hence for all $t \in [0, T]$, $\tilde{\mathbb{P}}$ -a.s.

$$\tilde{\mathcal{V}}(\tilde{v})(t) = \tilde{v}(t)$$

and the proof is complete. □

3. EXISTENCE OF A SOLUTION TO THE STOCHASTIC KLAUSMEIER SYSTEM

In this section, we show the existence of a nonnegative solution to the Klausmeier system. First, we will introduce the notation and the definition of the solution and its martingale solution. As pointed out before we will use compactness arguments to show the existence, which implies the loss of the original probability space. Next, we present our main Theorem, which says under which conditions a martingale solution exists. Then, we give the proof of the main result.

Let H_1 and H_2 be two Hilbert spaces, let $\mathfrak{A} = (\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$ be a filtered probability space and W_j , $j = 1, 2$, are two cylindrical Wiener process defined on H_j , $j = 1, 2$, respectively. We are now interested in the solution to the following reduced Klausmeier system for $x \in \mathcal{O}$ and $t > 0$

$$(3.1) \quad du(t, x) = (r_u \Delta u^{[\gamma]}(t, x) - u(t, x) v^2(t, x)) dt + \sigma_1(u(t, x)) \circ dW_1(x),$$

and

$$(3.2) \quad dv(t, x) = (r_v \Delta v(t, x) + u(t, x) v^2(t, x)) dt + \sigma_2(v(t, x)) \circ dW_2(x),$$

with Neumann (or if \mathcal{O} is a torus, periodic) boundary conditions and initial conditions $u(0) = u_0$ and $v(0) = v_0$. Let $r_u, r_v > 0$ be positive constants. Here, we use the abbreviation $x^{[\gamma]} := |x|^{\gamma-1}x$. The mappings σ_j , $j = 1, 2$, will be specified later on. Since the white noise is an approximation of a continuously fluctuating noise with finite memory being much shorter than the dynamical timescales, the representation of the stochastic integral as a Stratonovich stochastic integral is appropriate. In order to show the existence of a solution to the original system, the linear parts have to be incorporated, which can be done by modifying the proof given here. The structure of the proof will remain if one adds a linear gradient term to the equation.

Due to the porous media term, we do not use solutions in the classical sense. Firstly, let us define what is a solution.

Definition 3.1. *We call a couple (u, v) solution to the system (3.1) and (3.2), for initial data (u_0, v_0) if there exists $\rho \in \mathbb{R}$ such that*

$$u \in L^2(\Omega; C([0, T]; H_2^{-1}(\mathcal{O}))) \cap L^{\gamma+1}(\Omega \times (0, T) \times \mathcal{O})$$

and

$$v \in L^2(\Omega; C([0, T]; H_2^\rho(\mathcal{O}))) \cap L^2(\Omega \times (0, T); H_2^{\rho+1}(\mathcal{O}))$$

such that u and v are $(\mathcal{F}_t)_{t \geq 0}$ -adapted, and satisfies for all $t \in [0, T]$, \mathbb{P} -a.s.

$$u(t) = u_0 + \int_0^t (r_u \Delta u^{[\gamma]}(s) - u(s)v^2(s)) ds + \int_0^t \sigma_1(u(s)) \circ dW_1(s),$$

and

$$v(t) = v_0 + \int_0^t (r_v \Delta v(s) + u(s)v^2(s)) ds + \int_0^t \sigma_2(v(s)) \circ dW_2(s).$$

As mentioned before, in the proof of the main result, we are using compactness arguments, which causes the loss of the original probability space, which means the solution will only be a weak solution in the probabilistic sense.

Definition 3.2. *A martingale solution to the problem (3.1) and (3.2) is a system*

$$(3.3) \quad (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P}, (W_1, W_2), (u, v))$$

such that

- $\mathfrak{A} := (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ is a complete filtered probability space with a filtration $\mathbb{F} = \{\mathcal{F}_t : t \in [0, T]\}$ satisfying the usual conditions,
- W_1 and W_2 are H_1 -valued, respectively H_2 -valued Wiener processes over the probability space \mathfrak{A} with covariance Q_1 and Q_2 ;

- $u : [0, T] \times \Omega \rightarrow H_2^{-1}(\mathcal{O})$ and $v : [0, T] \times \Omega \rightarrow H^\rho(\mathcal{O})$ are two \mathbb{F} -progressively measurable processes such that the couple (u, v) is a solution to the system (3.1) and (3.2) over the probability space \mathfrak{A} .

The Stratonovich integral is not a martingale, and, therefore, often, not easy to handle. Hence, for our purposes, instead of the Stratonovich formulation, it is convenient to consider the equation in Itô form for $x \in \mathcal{O}$, $t > 0$.

$$(3.4) \quad \begin{cases} du(t, x) &= (r_u \Delta u^{[\gamma]}(t, x) - u(t, x) v^2(t, x)) dt + \sigma_1(u(t, x)) dW_1(x), \\ u(0) &= u_0, \end{cases}$$

and

$$(3.5) \quad \begin{cases} dv(t, x) &= (r_v \Delta v(t, x) + u(t, x) v^2(t, x)) dt + \sigma_2(v(t, x)) dW_2(x), \\ v(0) &= v_0. \end{cases}$$

In the setting above, we interpret the stochastic integral as an Itô integral. For simplicity, we omit the correction term (cf. [10]). If one is interested in the exact form of the correction term, we refer to [14]. Before presenting our main result, we will first introduce the hypotheses on d , γ , the initial conditions, and on σ_j , $j = 1, 2$.

Hypothesis 3.3. *Let*

$$d < \frac{4(1 + 3\gamma)}{3(1 + \gamma)}$$

Hypothesis 3.4. *Let $d = 1, 2, 3$, $p \in (2\mathbb{N} + 1)$, $p \geq 7$, $\rho \in \mathbb{R}$, and $\delta_0 \in \mathbb{R}$ be some numbers satisfying*

$$\begin{aligned} 1 - \frac{d}{2} &> \rho > -\frac{p+1}{2p}, \\ \frac{2\gamma+1}{\gamma+1} &> \frac{3}{8}d + \frac{2}{p+1}, \\ \gamma &> \frac{1+d-2\rho}{3+2\rho}, \quad \text{and,} \\ \delta_0 &< \frac{\gamma}{\gamma+1}. \end{aligned}$$

Let us assume that

$$\mathbb{E}|u_0|_{L^{p+1}}^{p+1} < \infty$$

and

$$\mathbb{E}|v_0|_{H_2^{-\delta_0}}^{\frac{2\gamma+1}{\gamma-1}} < \infty \quad \text{and} \quad \mathbb{E}|v_0|_{H_2^\rho}^2 < \infty.$$

and that u_0 and v_0 are a.e. nonnegative functions (nonnegative Borel measures that are finite on compact subsets, respectively).

Hypothesis 3.5. *Assume that $H_1 = H_2 = H_2^{-1}(\mathcal{O})$. Assume that there exist constants $\tilde{\sigma}_1, \tilde{\sigma}_2 > 0$ such that*

$$\sigma_1(\xi)h = \tilde{\sigma}_1 \xi (-\Delta)^{-\beta/2} h, \quad h \in H_1, \quad \xi \in L^2(\mathcal{O}),$$

and

$$\sigma_2(\xi)h = \tilde{\sigma}_2 \xi (-\Delta)^{-\beta/2} h, \quad h \in H_2, \quad \xi \in L^2(\mathcal{O}),$$

where $\beta > \frac{3}{2}d$.

Remark 3.6. If $\mathcal{O} = [0, 1]^d$ is a rectangular domain, then the condition $\beta > \frac{3}{2}d$ can be relaxed to $\beta > \frac{d}{2}$ (see [2, p. 7 and p. 22]).

Under these hypotheses the existence of a martingale solution can be shown.

Theorem 3.7. Fix $m = 2\frac{\gamma+1}{\gamma}$ and $m_0 = 2\frac{\gamma+1}{\gamma-1}$. Under the Hypotheses 3.3, 3.4, and 3.5, there exists a martingale solution to system (3.4)–(3.5) satisfying the following properties

- (i) $\mathbb{P} \otimes \text{Leb}$ a.s. $u(x, t) \geq 0$ and $v(x, t) \geq 0$;
- (ii) $\mathbb{E} \left[\sup_{0 \leq s \leq T} |u(s)|_{L^p}^p \right] + p(p-1)C(r_u)\mathbb{E} \int_0^T \int_{\mathcal{O}} u(s, x)^{p-2} |\nabla u(s, x)|^2 dx ds$
 $+ \mathbb{E} \int_0^T \int_{\mathcal{O}} u^{p+1}(s, x) v^2(s, x) dx ds \leq C(T) \mathbb{E} |u_0|_{L^{p+1}}^{p+1},$
- (iii) $\mathbb{E} \int_0^T |v(s)|_{L^m}^{m_0} ds \leq C_2(T) \mathbb{E} |v_0|_{H_2^{-\delta_0}}^{m_0},$
- (iv) and $\mathbb{E} \left[\sup_{0 \leq s \leq T} |v(s)|_{H_2^p}^2 \right] + \mathbb{E} \int_0^T |v(s)|_{H_2^p}^2 ds \leq C_3(T) \mathbb{E} |v_0|_{H_2^p}^2.$

Let H and \tilde{H} be two Hilbert spaces, then the set of Hilbert Schmidt mappings from H to \tilde{H} is denoted by $\mathcal{L}_2(H, \tilde{H})$.

The proof is an application of Theorem 2.1 and consists of five steps. However, to keep the proof itself simple, the proof will use a couple of technical results which are collected in Section 4.

Proof of Theorem 3.7. In the first step, we are specifying the underlying Banach spaces and, in the second step, we construct the operator \mathcal{V} . Then, in the third step, we show that the operator \mathcal{V} satisfies the assumption of Theorem 2.1. In the fifth step, we apply the Theorem 2.1 and show that the fixed point is indeed a martingale solution satisfying the above properties.

Step I: Here we define the spaces on which the operator \mathcal{V} will act. Let the probability space $\mathfrak{A} = (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ be given and let W_1 and W_2 be two independent H_1 and H_2 -valued Wiener processes defined over \mathfrak{A} with covariances Q_1 and Q_2 . Let $W = (W_1, W_2)$, $H = H_1 \times H_2$, with covariance operator

$$Q = \begin{pmatrix} Q_1 & 0 \\ 0 & Q_2 \end{pmatrix}.$$

Fix the reflexive Banach space $\mathbb{X} := L^{m_0}(0, T; L^m(\mathcal{O}))$, $m = 2\frac{\gamma+1}{\gamma}$, $m_0 = 2\frac{\gamma+1}{\gamma-1}$, and fix the compact and reflexive Banach subspace $\mathbb{X}' = L^{m_0}(0, T; H_m^\sigma(\mathcal{O})) \cap \mathbb{W}^{\alpha, m_0}(0, T; H_m^{-\delta_2}(\mathcal{O}))$ equipped with the norm

$$\|x\|_{\mathbb{X}'} := \left(\|x\|_{L^{m_0}(0, T; H_m^\sigma)}^{m_0} + \|x\|_{\mathbb{W}^{\alpha, m_0}(0, T; H_m^{-\delta_2})}^{m_0} \right)^{1/m_0}.$$

Put

$$(3.6) \quad \tilde{R} = \tilde{C}_0(T) \mathbb{E} |u_0|_{L^{p+1}}^{p+1}$$

where $\tilde{C}_0(T)$ is the constant appearing in Proposition 4.5. Let $R_1 > 0$ so large that

$$(3.7) \quad \tilde{C}_1(T) \left(\mathbb{E} |v_0|_{H_2^p}^2 + \mathbb{E} |u_0|_{L^{p+1}}^{p+1} + R_1^{7/8} \right) \leq R_1,$$

where $\tilde{C}_1(T)$ is the constant appearing in Proposition 4.9 and let $R_2 > 0$ so large that

$$(3.8) \quad \tilde{C}_2(T) \left(\mathbb{E}|v_0|_{H_m^{-\delta_0}}^2 + C_0 T^{\beta_0} R_1^{\delta_1} \tilde{R}^{\delta_2} \right) \leq R_2,$$

where $\tilde{C}_2(T)$, C_0 , δ_1 , δ_2 , and β_0 are the constants appearing in Proposition 4.12-(i). Fix the set

$$\begin{aligned} & \mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho) \\ := & \left\{ \xi \in \mathbb{X} : \xi(t, x) \geq 0 \text{ Leb} \otimes \mathbb{P}\text{-a.s. for all } (t, x) \in [0, T] \times \mathcal{O}, \right. \\ & \mathbb{E}\|\xi\|_{\mathbb{X}}^2 \leq R_2 \quad \text{and} \quad \mathbb{E}\|\xi\|_{\mathbb{Z}}^2 \leq R_1, \quad \xi \text{ progressively measurable on } \mathfrak{A}, \\ & \left. \xi(0, x) = v_0(x) \text{ for a.e. } x \in \mathcal{O} \right\}, \end{aligned}$$

where $\mathbb{Z} = L^\infty(0, T; H_2^\rho(\mathcal{O})) \cap L^2(0, T; H_2^{\rho+1}(\mathcal{O}))$, and note that $\mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho)$ is a (strongly) closed convex subset of the reflexive Banach space $\mathcal{M}_{\mathfrak{A}}^m(\mathbb{X})$, which defined as in (2.2), as we have that $m > 2$.

To define the operator \mathcal{V} let us introduce a second auxiliary Banach space $\mathbb{Y} = L^{\gamma+1}(0, T; L^{\gamma+1}(\mathcal{O})) \cap L^\infty(0, T; H_2^{-1}(\mathcal{O}))$ and the space of progressively measurable processes $\mathcal{M}_{\mathfrak{A}}^2(\mathbb{Y})$ by

$$\begin{aligned} \mathcal{M}_{\mathfrak{A}}^2(\mathbb{Y}) := & \left\{ \xi : [0, T] \times \Omega \rightarrow \mathcal{D}'(\mathcal{O}) : \right. \\ & \left. \xi \text{ is progressively measurable on } \mathfrak{A} \text{ and } \mathbb{E}|\xi|_{\mathbb{Y}}^2 < \infty \right\} \end{aligned}$$

equipped with norm

$$|\xi|_{\mathbb{Y}} := \left(\mathbb{E}|\xi|_{\mathbb{Y}}^2 \right)^{\frac{1}{2}}, \quad \xi \in \mathbb{Y}.$$

In addition, let us fix the convex subset

$$\begin{aligned} & \mathcal{Y}_{\mathfrak{A}}(\tilde{R}, p) \\ := & \left\{ \xi \in \mathbb{Y} : \xi(t, x) \geq 0 \text{ Leb} \otimes \mathbb{P}\text{-a.s. for all } (t, x) \in [0, T] \times \mathcal{O}, \right. \\ & \xi \text{ progressively measurable on } \mathfrak{A}, \quad \xi(0, x) = u_0(x) \text{ for a.e. } x \in \mathcal{O}, \\ & \left. \mathbb{E} \left[\sup_{0 \leq s < T} |\xi(s)|_{L^{p+1}}^{p+1} \right] + p(p-1)C(r_u) \mathbb{E} \int_0^T \int_{\mathcal{O}} \xi^{p-2}(s, x) |\nabla \xi(s, x)|^2 dx ds \leq \tilde{R} \right\}, \end{aligned}$$

where $C(r_u)$ is as in Proposition 4.5.

Step II: In this step we define the integral operator \mathcal{V} . First, we introduce the operator

$$\mathcal{U} := \mathcal{U}_{\mathfrak{A}} : \mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho) \rightarrow \mathcal{Y}_{\mathfrak{A}}(\tilde{R}, p)$$

by $\mathcal{U}[v] = u$, where u is a solution to

$$(3.9) \quad du(t) = r_u \Delta u^{[\gamma]}(t) dt - u(t) v^2(t) dt + \sigma_1(u(t)) dW_1(t), \quad u(0) = u_0.$$

The operator \mathcal{U} is well defined. In fact, given $v \in \mathcal{X}(R_1, R_2, \rho)$ it follows by Theorem 4.3 below the existence of a unique solution u to (3.9) such that

$$\mathbb{E} \left[\sup_{0 \leq s \leq T} |u(s)|_{H_2^{-1}}^2 \right] + \mathbb{E} \int_0^T |u(s)|_{L^{\gamma+1}}^{\gamma+1} ds < \infty.$$

The positivity follows by Proposition 4.4 below. By Proposition 4.5 and the choice of \tilde{R} it follows that $u = \mathcal{U}[v] \in \mathcal{Y}_{\mathfrak{A}}(\tilde{R}, p)$. In particular, if u_0 satisfies Hypothesis 3.4 and $v \in \mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho)$, then we have by (3.6)

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq s < T} |u(s)|_{L^{p+1}}^{p+1} \right] + p(p-1)C(r_u) \mathbb{E} \int_0^T \int_{\mathcal{O}} u^{p-2}(s, x) |\nabla u(s, x)|^2 dx ds \\ & + p \mathbb{E} \int_0^T \int_{\mathcal{O}} u^{p+1}(s, x) v^2(s, x) dx ds \leq \tilde{R}. \end{aligned}$$

Next, let us define the operator

$$\mathcal{V} := \mathcal{V}_{\mathfrak{A}} : \mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho) \rightarrow \mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho),$$

where $\mathcal{V}[v] = w$ and w is a solution to

$$(3.10) \quad dw(t) = r_v \Delta w(t) + u(t)v^2(t) + \sigma_2(u(t))dW_2(t), \quad w(0) = v_0,$$

with $u = \mathcal{U}[v]$. It follows by Proposition 4.9 the existence of a unique solution to (3.10). In addition, setting $p = 7$ and $\kappa = 4$ as in Proposition 4.9 we have

$$\mathbb{E} \left[\sup_{0 \leq s \leq T} |w(s)|_{H_2^\rho}^2 \right] + \mathbb{E} \int_0^T |w(s)|_{H_2^{\rho+1}}^2 ds \leq \tilde{C}_1(T) \left\{ \mathbb{E}|v_0|_{H_2^\rho}^2 + \mathbb{E}|u_0|_{L^{p+1}}^{p+1} + R_1^{\frac{7}{8}} \right\}.$$

Since R_1 is satisfying (3.7) we know that given $\mathbb{E}|v_0|_{H_2^\rho}^2$ and $\mathbb{E}|u_0|_{L^{p+1}}^{p+1}$ that we have

$$\left(\mathbb{E} \left[\sup_{0 \leq s \leq T} |w(s)|_{H_2^\rho}^2 \right] + \mathbb{E} \int_0^T |w(s)|_{H_2^{\rho+1}}^2 ds \right) \leq R_1.$$

The nonnegativity follows by Proposition 4.11. Next, by Proposition 4.12-(i) we know

$$\mathbb{E} \|w\|_{L^{m_0}(0, T; L^m)}^m \leq \tilde{C}_2(T) \left(\mathbb{E}|v_0|_{H_m^{-\delta_0}}^2 + C_0 T^{\beta_0} R_1^{\delta_1} \tilde{R}^{\delta_2} \right),$$

where $\tilde{C}_2(T)$, C_0 , δ_1 , δ_2 , and β_0 are the constant appearing in Proposition 4.12-(i). Given R_1 and \tilde{R} , due to (3.8) we have

$$\mathbb{E} \|w\|_{L^{m_0}(0, T; L^m)}^{m_0} \leq R_2.$$

It follows that the operator \mathcal{V} maps $\mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho)$ into $\mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho)$. In particular, assumption (i) of Theorem 2.1 is satisfied.

Step III: Next we show that assumption (ii) of Theorem 2.1 is satisfied. In particular, that the operator $\mathcal{V} : \mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho) \rightarrow \mathcal{M}_{\mathfrak{A}}^m(\mathbb{X})$ is continuous.

Due to Theorem 4.3 and Proposition 4.15 below the operator $\mathcal{U} : \mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho) \rightarrow \mathcal{M}_{\mathfrak{A}}^2(\mathbb{Y})$ is continuous. It follows now from Theorem 4.14, that the operator \mathcal{V} is continuous. In particular, let $v_n \rightarrow v$ in $\mathcal{M}_{\mathfrak{A}}^m(\mathbb{X})$, $v_n, v \in \mathcal{X}_{\mathfrak{A}}(R_1, R_2, \rho)$. Then, $u_n := \mathcal{V}[v_n] \rightarrow u = \mathcal{V}[v]$ in $\mathcal{M}_{\mathfrak{A}}^2(\mathbb{Y})$. By Theorem 4.14 follows that $\mathcal{V}(v_n) \rightarrow \mathcal{V}[v]$

in $\mathcal{M}_{\mathfrak{X}}^m(\mathbb{X})$, and, in particular, that the operator $\mathcal{V} : \mathcal{X}_{\mathfrak{X}}(R_1, R_2, \rho) \rightarrow \mathcal{M}_{\mathfrak{X}}^m(\mathbb{X})$ is continuous.

Step IV: Next we show that assumption (iii) of Theorem 2.1 is satisfied. By Proposition 4.12 it follows that there exists a constant $C > 0$ such that

$$\mathbb{E} |\mathcal{V}[v]|_{\mathbb{X}'}^{m_0} \leq C,$$

for all $v \in \mathcal{X}_{\mathfrak{X}}(R_1, R_2, \rho)$, where \mathbb{X}' is defined in Step I.

To show assumption (d) of Theorem 2.1, we notice first that due to the choice of $U_1 = H_2^\rho(\mathcal{O})$ it follows from Proposition 4.9 and Kolmogorov's continuity criterion that $\mathcal{V}(v) \in C([0, T]; H_2^\rho(\mathcal{O}))$ \mathbb{P} -a.s. for all $v \in \mathcal{X}_{\mathfrak{X}}(R_1, R_2, \rho)$.

Step V: By Theorem 2.1, there exists a probability space $\tilde{\mathfrak{A}} = (\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}})$, a Wiener process $\tilde{W} = (\tilde{W}_1, \tilde{W}_2)$ defined on $\tilde{\mathfrak{A}}$ and an element $\tilde{v} \in \mathcal{X}_{\tilde{\mathfrak{A}}}(R_1, R_2, \rho) \cap \mathbb{D}([0, T]; H_2^\rho(\mathcal{O}))$ such that $\tilde{\mathbb{P}}$ -a.s.

$$\mathcal{V}_{\tilde{\mathfrak{A}}}(v)(t) = v(t)$$

for all $t \in [0, T]$. Let $\tilde{u} = \mathcal{U}_{\tilde{\mathfrak{A}}}(\tilde{v})$. Due to the construction of $\mathcal{V}_{\tilde{\mathfrak{A}}}$ and $\mathcal{U}_{\tilde{\mathfrak{A}}}$ the pair (\tilde{u}, \tilde{v}) solves the system (3.4)–(3.5) over the stochastic basis $\tilde{\mathfrak{A}}$ with the Wiener noise \tilde{W} .

The asserted properties of the solutions follow by the fact that $\tilde{v} \in \mathcal{X}_{\tilde{\mathfrak{A}}}(R_1, R_2, \rho)$ and $\tilde{u} = \mathcal{U}_{\tilde{\mathfrak{A}}}(\tilde{v}) \in \mathcal{Y}_{\tilde{\mathfrak{A}}}(\tilde{R}, p)$ and by Proposition 4.5.

The proof of the main result Theorem 3.7 is complete. \square

4. RESULTS ON REGULARITY AND TECHNICAL PROPOSITIONS

We begin with a remark on the noise coefficients.

Remark 4.1. *From Hypotheses 3.5, one can infer that there exist a constant $c > 0$ such that*

$$\sum_{k=1}^{\infty} [(\sigma_j(\xi) f_k)(x)]^2 \leq c |\xi(x)|^2, \quad \forall \xi \in L^2(\mathcal{O}), x \in \mathcal{O}, \text{ and } j = 1, 2.$$

Here $\{f_k\}$ is an orthonormal basis in $H_2^{-1}(\mathcal{O})$ (compare [2, Hypothesis 3, p. 42]). In addition, note, that

- $\sigma_j : H_2^{-1}(\mathcal{O}) \rightarrow \mathcal{L}_2(H_j, H_2^{-1}(\mathcal{O}))$ is of linear growth and Lipschitz continuous. In particular, there exists constants $C_1, L_1 > 0$ such that

$$|\sigma_j(\xi)|_{\mathcal{L}_2(H_j, H_2^{-1})} \leq C_1(1 + |\xi|_{H_2^{-1}}), \quad \xi \in H_2^{-1}(\mathcal{O});$$

$$|\sigma_j(\xi) - \sigma_j(\eta)|_{\mathcal{L}_2(H_j, H_2^{-1})} \leq L_1 |\xi - \eta|_{H_2^{-1}}, \quad \xi, \eta \in H_2^{-1}(\mathcal{O});$$

- and $\sigma_j : L^2(\mathcal{O}) \rightarrow \mathcal{L}_2(H_j, L^2(\mathcal{O}))$ is of linear growth and Lipschitz continuous. In particular, there exists constants $C_2, L_2 > 0$ such that

$$|\sigma_j(\xi)|_{\mathcal{L}_2(H_j, L^2)} \leq C_2(1 + |\eta|_{L^2}), \quad \eta \in L^2(\mathcal{O});$$

$$|\sigma_j(\xi) - \sigma_j(\eta)|_{\mathcal{L}_2(H_j, L^2)} \leq L_2 |\xi - \eta|_{L^2}, \quad \xi, \eta \in L^2(\mathcal{O}).$$

Similarly, straightforward computations and using the fact that $|f_k|_{L^\infty} \leq \alpha_k^{\frac{d-1}{2}}$ (see [2, p. 46]) we get (see e.g. [4, 37])

$$|\sigma_2(\xi)|_{\gamma(H_2, L^m)} \leq C_1(1 + |\xi|_{L^m}), \quad \xi \in L^m(\mathcal{O});$$

$$|\sigma_2(\xi) - \sigma_2(\eta)|_{\gamma(H_2, L^m)} \leq L_1 |\xi - \eta|_{L^m}, \quad \xi, \eta \in L^m(\mathcal{O}),$$

where $\gamma(H_2, L^m)$ denotes the space of γ -radonifying operators.

4.1. Properties of the System (3.4). In the first part, we analyze the equation (3.4), where we impose following assumption on the process v . Let us remind that we fixed $m = 2\frac{\gamma+1}{\gamma}$ and $m_0 = 2\frac{\gamma+1}{\gamma-1}$.

Assumption 4.2. *Given $K_1 > 0$ let v be a progressively measurable function-valued process over \mathfrak{A} such that*

- (i) $v \in L^{m_0}(0, T; L^m(\mathcal{O}))$ with $\mathbb{E}\|v\|_{L^{m_0}(0, T; L^m)}^{m_0} \leq K_1$;
- (ii) is \mathbb{P} -a.s. nonnegative.

We are interested in the solution to the following system

$$(4.1) \quad \begin{cases} du(t, x) = [r_u \Delta u^{[\gamma]}(t, x) - u(t, x)v^2(t, x)] dt + \sigma_1(u(t, x)) dW_1(t), & x \in \mathcal{O}, t > 0, \\ u(0, x) = u_0(x), & x \in \mathcal{O}. \end{cases}$$

First, we will show that a unique solution to the system (4.1) exists and is nonnegative. In addition, we will show in Proposition 4.15 that the solution depends continuously on v , and, secondly, we will show by variational methods that this solution satisfies some integrability properties, given $u_0 \in L^{p+1}$.

Theorem 4.3. *Assume that v is fixed satisfying (4.2). Then for any $u_0 \in L^2(\Omega; \mathcal{F}_0, \mathbb{P}; H_2^{-1}(\mathcal{O}))$*

$$(4.2) \quad du(t) = [r_u \Delta u^{[\gamma]}(t) dt - u(t)v^2(t) dt + \sigma_1(u(t)) dW_1, \quad t \in [0, T], \quad u(0) = u_0,$$

has a unique solution $u = \{u(t) : t \in [0, T]\}$ and

$$\mathbb{E} \left[\sup_{0 \leq s \leq T} |u(s)|_{H_2^{-1}}^2 \right] + \mathbb{E} \int_0^T |u(s)|_{L^{\gamma+1}}^{\gamma+1} ds < \infty.$$

Proof. Before starting with the proof, we introduce the setting used by the book of Barbu, Da Prato and Röckner [2], the book of Liu and Röckner [20] respectively.

Let $\mathcal{H} := H_2^{-1}(\mathcal{O})$, the dual space of $\mathcal{H}^* = H_2^1(\mathcal{O})$ (corresponding to Neumann boundary conditions). Fix some $p \geq \gamma + 1$ and set $V := L^p(\mathcal{O})$. By the Sobolev embedding theorem, $\mathcal{H}^* \hookrightarrow V^* = L^{p/(p-1)}(\mathcal{O})$ densely and compactly. Thus, upon identifying \mathcal{H} with its dual space via the Riesz-map $(-\Delta)^{-1}$ of \mathcal{H} , we have a Gelfand triple

$$V \subset \mathcal{H} \cong \mathcal{H}^* \subset V^*.$$

We set

$$A(t, u, \omega) := A_v(u) := r_u \Delta u^{[\gamma]} - v^2(t, \omega)u.$$

Note, due to assumption 4.2, v is adapted and there exists a constant $K > 0$ such that

$$(4.3) \quad \mathbb{E} \int_0^T |v(t)|_{L^m}^{m_0} dt \leq K_1.$$

Then, under the assumption (4.3), we shall verify the conditions of [20, Theorem 5.1.3] for $\gamma > 1$. First note that for fixed $t \in [0, T]$ and fixed $\omega \in \Omega$, A maps from V to V^* . In

particular, by Hölder inequality and the Sobolev embedding $L^1(\mathcal{O}) \hookrightarrow H_{\frac{\gamma+1}{\gamma}}^{-2}(\mathcal{O})$, we have for v as above, $d = 1, 2, 3$, and $\gamma > 1$

$$\begin{aligned}
(4.4) \quad |_{V^*} \langle A_v(u), w \rangle_V &= \left| \int_{\mathcal{O}} \left[r_u u^{[\gamma]} w + (-\Delta)^{-1}(v^2(t, \omega)u)w \right] dx \right| \\
&\leq \left[r_u |u|_{L^{\gamma+1}}^\gamma |w|_{L^{\gamma+1}} + |(-\Delta)^{-1}(v^2 u)|_{L^{(\gamma+1)/\gamma}} |w|_{L^{\gamma+1}} \right] \\
&\leq \left[r_u |u|_{L^{\gamma+1}}^\gamma + C |v^2 u|_{L^1} \right] |w|_{L^{\gamma+1}} \\
&\leq \left[r_u |u|_{L^{\gamma+1}}^\gamma + C |v|_{L^{2\frac{\gamma+1}{\gamma}}}^2 |u|_{L^{\gamma+1}} \right] |w|_{L^{\gamma+1}} \\
&\leq \left[r_u |u|_{L^{\gamma+1}}^\gamma + C |v|_{L^{2\frac{\gamma+1}{\gamma}}}^2 |u|_{L^{\gamma+1}} \right] |w|_{L^{\gamma+1}} \\
&\leq \left[r_u |u|_{L^{\gamma+1}}^\gamma + C |v|_{L^{2\frac{\gamma+1}{\gamma}}}^{2\gamma/(\gamma-1)} + C |u|_{L^{\gamma+1}}^\gamma \right] |w|_{L^{\gamma+1}},
\end{aligned}$$

where only v depends on t and ω and C may change from line to line. Next, we verify Hypothesis (H1), (H2'), (H3), and (H4') of [20, Theorem 5.1.3].

(H1): For $\lambda \in \mathbb{R}$, $u_1, u_2, w \in L^{\gamma+1}(\mathcal{O})$, $t \in [0, T]$, $\omega \in \Omega$ consider the map

$$\lambda \mapsto \langle A_v(u_1 + \lambda u_2), w \rangle$$

and show its hemicontinuity. Note, that we have

$$\langle A_v(u_1 + \lambda u_2), w \rangle = -r_u \int_{\mathcal{O}} (u_1 + \lambda u_2)^{[\gamma]} w dx - \int_{\mathcal{O}} (-\Delta)^{-1}[v^2(t)(u_1 + \lambda u_2)]w dx,$$

where only v depends on t and ω . For the first integral in the above identity, we prove hemicontinuity with the same arguments as in [20, Example 4.1.11, p. 87]. For the second integral, we note that $u \mapsto (-\Delta)^{-1}(v^2(t, \omega)u)$ is a linear operator for fixed t and ω which is bounded as a map from $L^{\gamma+1}$ to $L^{(\gamma+1)/\gamma}$ by (4.4) and thus continuous. Due to remark 4.1 the assumption (H1) of Theorem 5.1.3 of [20] is satisfied.

(H2'): Let $u, w \in L^{\gamma+1}(\mathcal{O})$, $t \in [0, T]$, $\omega \in \Omega$. Take (A.1) into account and consider, for some $\delta \in (0, 2^{-2\frac{\gamma-1}{\gamma+1}} r_u^{\frac{2}{\gamma+1}}]$,

$$\begin{aligned}
&|_{V^*} \langle A_v(u) - A_v(w), u - w \rangle_V + |\sigma_1(u) - \sigma_1(w)|_{\mathcal{L}_2(H_1, \mathcal{H})}^2 \\
&\leq - \int_{\mathcal{O}} \left[r_u (u^{[\gamma]} - w^{[\gamma]})(u - w) + (-\Delta)^{-1}(v^2(t, \omega)(u - w))(u - w) \right] d\xi + C |u - w|_{\mathcal{H}}^2 \\
&\leq -2^{1-\gamma} r_u |u - w|_{L^{\gamma+1}}^{\gamma+1} + C |v|_{L^{2\frac{\gamma+1}{\gamma}}}^2 |u - w|_{L^{\gamma+1}} |u - w|_{L^{\gamma+1}} + C |u - w|_{\mathcal{H}}^2 \\
&\leq -2^{1-\gamma} r_u |u - w|_{L^{\gamma+1}}^{\gamma+1} + C |v|_{L^{2\frac{\gamma+1}{\gamma}}}^2 |u - w|_{L^{\gamma+1}}^2 + C |u - w|_{\mathcal{H}}^2 \\
&\leq -2^{1-\gamma} r_u |u - w|_{L^{\gamma+1}}^{\gamma+1} + C(\gamma) \delta^{-(\gamma+1)/(\gamma-1)} |v|_{L^{2\frac{\gamma+1}{\gamma}}}^{2(\gamma+1)/(\gamma-1)} + \delta^{\frac{\gamma+1}{2}} |u - w|_{L^{\gamma+1}}^{\gamma+1} + C |u - w|_{\mathcal{H}}^2 \\
&\leq C(\gamma) \delta^{-(\gamma+1)/(\gamma-1)} |v|_{L^{2\frac{\gamma+1}{\gamma}}}^{2(\gamma+1)/(\gamma-1)} + C |u - w|_{\mathcal{H}}^2.
\end{aligned}$$

Hence, (H2') holds with

$$f := C(\delta, \gamma) |v|_{L^2 \frac{\gamma+1}{\gamma}}^{2(\gamma+1)/(\gamma-1)} \in L^1_+([0, T] \times \Omega).$$

(H3): Let $u \in L^{\gamma+1}(\mathcal{O})$, $t \in [0, T]$, $\omega \in \Omega$. Consider, for some $\delta_0 \in (0, r_u^{\frac{2}{\gamma+1}})$,

$$\begin{aligned} & v^* \langle A_v(u), u \rangle_V + |\sigma_1(u)|_{\mathcal{L}_2(H_1, \mathcal{H})}^2 \\ & \leq - \int_{\mathcal{O}} \left[r_u u^{[\gamma]} u + (-\Delta)^{-1}(v^2(t, \omega)u)u \right] d\xi + C|u|_{\mathcal{H}}^2 \\ & \leq -r_u |u|_{L^{\gamma+1}}^{\gamma+1} + C|v|_{L^2 \frac{\gamma+1}{\gamma}}^2 |u|_{L^{\gamma+1}} |u|_{L^{\gamma+1}} + C|u|_{\mathcal{H}}^2 \\ & \leq -r_u |u|_{L^{\gamma+1}}^{\gamma+1} + C|v|_{L^2 \frac{\gamma+1}{\gamma}}^2 |u|_{L^{\gamma+1}}^2 + C|u|_{\mathcal{H}}^2 \\ & \leq C(\gamma) \delta_0^{-(\gamma+1)/(\gamma-1)} |v|_{L^2 \frac{\gamma+1}{\gamma}}^{2(\gamma+1)/(\gamma-1)} + \left(\delta_0^{\frac{\gamma+1}{2}} - r_u \right) |u|_{L^{\gamma+1}}^{\gamma+1} + C|u|_{\mathcal{H}}^2 \end{aligned}$$

As a consequence, (H3) holds with $f := C(\gamma, \delta_0) |v|_{L^2 \frac{\gamma+1}{\gamma}}^{2(\gamma+1)/(\gamma-1)}$ and $\theta := \left(r_u - \delta_0^{\frac{\gamma+1}{2}} \right)$.

(H4'): Let $u \in L^{\gamma+1}(\mathcal{O})$, $t \in [0, T]$, $\omega \in \Omega$. Then, (H4') holds by (4.4) with $\alpha := \gamma + 1$, $\beta := 0$ and $f := C(1 + |v|_{L^2 \frac{\gamma+1}{\gamma}}^{2(\gamma+1)/\gamma})$.

The rest of the proof follows by an application of [20, Theorem 5.1.3]. \square

Proposition 4.4. *Under the conditions of Theorem 4.3 and the additional condition $u_0(x) \geq 0$ for all $x \in \mathcal{O}$ and $v(t, x) \geq 0$, $(t, x) \in [0, T] \times \mathcal{O}$, the solution u to (4.1) is nonnegative.*

Proof. For the nonnegativity of the solution to (4.1) we refer to the proof of positivity of the stochastic porous medium equation (PME), see Section 2.6 in [2] and see also [1]. Mimicking the proof of nonnegativity in [2] the nonnegativity of (4.1) can be shown. \square

In the next proposition we are using variational methods to verify uniform bounds of u .

Proposition 4.5. *Let us assume that v satisfies the assumption 4.2 and u is a solution to (4.1). Fix $p_0 \geq 2$ and suppose $u_0 \in L^{p_0+1}(\mathcal{O})$. Then, there exists a constant $C_0(p_0, T) > 0$ such that we have for all $2 \leq p \leq p_0$*

$$(4.5) \quad \begin{aligned} & \mathbb{E} \left[\sup_{0 \leq s \leq T} |u(s)|_{L^{p+1}}^{p+1} \right] + p(p-1)C(r_u) \mathbb{E} \int_0^T \int_{\mathcal{O}} u(s, x)^{p-2} |\nabla u(s, x)|^2 dx ds \\ & + p \mathbb{E} \int_0^T \int_{\mathcal{O}} u^{p+1}(s, x) v^2(s, x) dx ds \leq C_0(p, T) \mathbb{E} |u_0|_{L^{p+1}}^{p+1}. \end{aligned}$$

Proof. Proposition 4.5 can be shown by the Itô-formula and straightforward calculations using the fact that u and v are nonnegative. \square

4.2. Properties of the system (3.5). Throughout this section we will assume that v satisfies the assumption 4.2. In addition, we will introduce the following assumptions.

Assumption 4.6. Given a number $K_2 > 0$ and a number $\rho \in \mathbb{R}$ we assume that

$$(4.6) \quad \mathbb{E} \left[\sup_{0 \leq s \leq T} |v(s)|_{H_2^\rho}^2 \right] + C_2 \mathbb{E} \int_0^T |v(s)|_{H_2^{\rho+1}}^2 ds \leq K_2.$$

Remark 4.7. Let (l, α) be a pair of numbers, $1 \leq l < \infty$, $\alpha \in \mathbb{R}$, such that

$$(4.7) \quad \alpha \leq \rho + \frac{2}{l}.$$

Then, it follows by interpolation, see e.g. [3, Theorem 5.1.2, p. 107], that there exist constants $C_1 > 0$ and $C_2 > 0$ such that

$$(4.8) \quad \mathbb{E} \left(\int_0^T |v(s)|_{H_2^\alpha}^l ds \right)^{\frac{2}{l}} \leq C_1 \mathbb{E} \left[\sup_{0 \leq s \leq T} |v(s)|_{H_2^\rho}^2 \right] + C_2 \mathbb{E} \int_0^T |v(s)|_{H_2^{\rho+1}}^2 ds.$$

Assumption 4.8. Given $K_3 > 0$, $p \geq 2$ and let u being a progressively measurable process over \mathfrak{A} with

$$\mathbb{E} \left[\sup_{0 \leq s \leq T} |u(s)|_{L^{p+1}}^{p+1} \right] < \infty$$

and the couple (u, v) satisfies

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq s \leq T} |u(s)|_{L^{p+1}}^{p+1} \right] + p(p-1)C(r_u) \mathbb{E} \int_0^T \int_{\mathcal{O}} u^{p-2}(s, x) |\nabla u(s, x)|^2 dx ds \\ & + p \mathbb{E} \int_0^T \int_{\mathcal{O}} u^{p+1}(s, x) v^2(s, x) dx ds \leq K_3. \end{aligned}$$

Given the couple (u, v) where v satisfies Assumption 4.2 and Assumption 4.6, and the couple (u, v) satisfies Assumption 4.8, we consider the solution w to the auxiliary system

$$(4.9) \quad dw(t) = [r_v \Delta w(t) + u(t)v^2(t)] dt + \sigma_2(w(t, x)) dW_2(t), \quad w(0) = v_0.$$

First, we will In the next proposition we investigate existence and uniqueness and the regularity of w .

Proposition 4.9. Fix $p \geq 2$ and $1 - \frac{d}{2} > \rho > -\frac{p+1}{2p}$. Let v satisfy Assumptions 4.2 and 4.6 with $\rho \in \mathbb{R}$. Then, for all couples (u, v) satisfying Assumption 4.8 with the given p there exists a solution w to the following system

$$dw(t) = [r_v \Delta w(t) + u(t)v^2(t)] dt + \sigma_2(w(t)) dW_2(t), \quad w(0) = w_0,$$

such that

$$w \in L^\infty(0, T; H_2^\rho(\mathcal{O})) \cap L^2(0, T; H_2^{\rho+1}(\mathcal{O})) \quad \mathbb{P} - a.s.$$

In particular, there exists a generic constant $\tilde{C}_2(T) > 0$ such that

$$\mathbb{E} \left[\sup_{0 \leq s \leq T} |w(s)|_{H_2^\rho}^2 \right] + \mathbb{E} \int_0^T |w(s)|_{H_2^{\rho+1}}^2 ds \leq \tilde{C}_2(T) \left(\mathbb{E}|v_0|_{H_2^\rho}^2 + \mathbb{E}|u_0|_{L^{p+1}}^{p+1} + R_1^{\frac{2}{\kappa} \frac{p}{p+1-\kappa}} \right).$$

Before proving Proposition 4.9, we consider the following lemma, which will be essential.

Lemma 4.10. *Let v satisfy assumption 4.2 and 4.6 with $\rho \in \mathbb{R}$. Then, for all $p \geq 2$ and $\kappa < p + 1$ such that $\rho + \frac{p+1-\kappa}{2p} \geq 0$ and for all couples (u, v) satisfying Assumption 4.8, there exists $R > 0$ such that*

$$\mathbb{E}|uv^2|_{L^\kappa(0,T;L^1)} \leq \mathbb{E}|u_0|_{L^{p+1}}^{p+1} + R^{\frac{2}{\kappa} \frac{p}{p+1-\kappa}}.$$

Proof of Lemma 4.10: Applying the Cauchy Schwarz inequality gives for $r = p + 1$, $r' = \frac{p+1}{p}$, and $\gamma = \frac{1}{r}$

$$|uv^2|_{L^1} \leq |uv^{2\gamma}|_{L^r} |v^{2(1-\gamma)}|_{L^{r'}} = |u^{p+1}v^2|_{L^1}^{\frac{1}{r}} |v^2|_{L^1}^{\frac{1}{r'}}.$$

By the Young inequality we have with $q = \frac{r}{\kappa}$ and $q' = \frac{r}{r-\kappa}$ ($r > \kappa$)

$$|u^{p+1}v^2|_{L^1}^{\frac{\kappa}{r}} |v^2|_{L^1}^{\frac{\kappa}{r'}} \leq |u^{p+1}v^2|_{L^1} + |v^2|_{L^1}^{\frac{2}{r-\kappa}} \leq |u^{p+1}v^2|_{L^1} + |v|_{L^2}^{\frac{4}{r-\kappa}}.$$

By Assumption 4.8 we know that $\mathbb{E} \int_0^T |u^{p+1}(s)v^2(s)|_{L^1} ds \leq \mathbb{E}|u_0|_{L^{p+1}}^{p+1}$. By Remark 4.7 and the fact that $\rho \geq -\frac{1}{2} \frac{r-\kappa}{r-1} = -\frac{1}{2} \frac{p+1-\kappa}{p}$, we have that

$$\mathbb{E} \left(\int_0^T |v(s)|_{L^2}^{\frac{4}{r-\kappa}} ds \right)^{\frac{1}{2} \frac{r-\kappa}{r-1}} \leq C_1 \mathbb{E} \left[\sup_{0 \leq s \leq T} |v(s)|_{H_2^\rho}^2 \right] + C_2 \mathbb{E} \int_0^T |v(s)|_{H_2^{\rho+1}}^2 ds \leq R.$$

and therefore, by Assumption 4.6, we get that

$$\mathbb{E} \int_0^T |u(s)v^2(s)|_{L^1}^2 ds \leq \mathbb{E}|u_0|_{L^{p+1}}^{p+1} + R^{\frac{2}{\kappa} \frac{p}{p+1-\kappa}}.$$

□

Proof of Proposition 4.9: Let us consider the following system

$$(4.10) \quad dw(t) = [r_v \Delta w(t) + F(t)] dt + \sigma_2(w(t, x)) dW_2(t), \quad w(0) = w_0.$$

We will prove the existence and uniqueness by applying [20, Theorem 4.2.4]. To be more precise, if $F \in L^2(0, T; L^1(\mathcal{O}))$, then for all $\rho < 1 - \frac{d}{2}$, the system (4.10) has a solution belonging \mathbb{P} -a.s.

$$C([0, T]; H_2^\rho(\mathcal{O})) \cap L^2(0, T; H_2^{\rho+1}(\mathcal{O})).$$

In fact, let us define the Gelfand triplet $V' = H_2^{\rho-1}(\mathcal{O})$, $\mathcal{H} = H_2^\rho(\mathcal{O})$ and $V = H_2^{\rho+1}(\mathcal{O})$. Let $\mathcal{A} : [0, T] \times \Omega \times V \rightarrow V'$ be given by $\mathcal{A}(t, \omega)u := r_v \Delta u + F(\omega, t)$, where

$$\text{dom}(\Delta) = \{u \in V' \mid \text{Neumann or periodic boundary conditions are satisfied}\}$$

and $F \in L^2(\Omega; L^2(0, T; V'))$ with $F(\omega, t) := u(\omega, t)v^2(\omega, t)$. It is straightforward to verify that this operator satisfies the assumptions of [20, Theorem 4.2.4], that is, [20, Conditions (H1)–(H4), p. 70]. In fact, hemicontinuity and weak monotonicity are given by the Laplace term. Observe, $\rho < 1 - \frac{d}{2}$, we know there exists a number $\sigma \in (0, 1)$ with $\frac{d}{2} = \sigma\rho + (1 - \sigma)(\rho + 1)$.

Using the Sobolev embedding $L^1(\mathcal{O}) \hookrightarrow H_2^{-\frac{d}{2}}(\mathcal{O})$ and interpolation, it follows that for all $\varepsilon > 0$

$$\langle F, w \rangle \leq |F|_{H_2^{-\frac{d}{2}}} |w|_{H_2^{\frac{d}{2}}} \leq |F|_{L^1} |w|_{\mathcal{H}}^\sigma |w|_V^{1-\sigma} \leq C_\varepsilon |F|_{L^1}^2 + c_\varepsilon |w|_{\mathcal{H}}^2 + \varepsilon |w|_V^2.$$

Choosing $\varepsilon = \frac{\lambda_1 r_v}{2}$, where $\lambda_1, \lambda_2 > 0$ are such that $\langle r_v \Delta w, w \rangle \leq -\lambda_1 r_v |w|_V^2 + \lambda_2 |w|_{\mathcal{H}}^2$, we get for all $t \in [0, T]$ with $f(t) = |F(t)|_{L^1}^2$

$$\begin{aligned} & \langle \mathcal{A}w, w \rangle + |\sigma_2(w)|_{\mathcal{L}_2(H_2, \mathcal{H})}^2 \\ & \leq -\lambda_1 r_v |w|_V^2 + \langle F(t), w \rangle + (C + \lambda_2) |w|_{\mathcal{H}}^2 \\ & \leq (C + \lambda_2 + c_\varepsilon) |w|_{\mathcal{H}}^2 - \frac{\lambda_1 r_v}{2} |w|_V^2 + (C_\varepsilon + 1) f(t). \end{aligned}$$

By Lemma 4.10 we know that under the assumption on v and u we have $uv^2 \in L^2((0, T) \times \Omega; L^1(\mathcal{O}))$. \square

Proposition 4.11. *Under the conditions of Theorem 4.3 and the additional condition $v_0(x) \geq 0$ for all $x \in \mathcal{O}$ and $u(t, x) \geq 0$, $(t, x) \in [0, T] \times \mathcal{O}$, the solution u to (4.1) is nonnegative.*

Proof. The semigroup generated by the Laplace maps a nonnegative function into a nonnegative function. In this way we refer to the proof of nonnegativity by Tessitore and Zabczyk [32]. The perturbation can be incorporated by comparison results (see Kotelenetz [17]). \square

Proposition 4.12. *Let $m = 2\frac{\gamma+1}{\gamma}$ and $m_0 = 2\frac{\gamma+1}{\gamma-1}$. Then, for*

$$d < \frac{4(1+3\gamma)}{3(1+\gamma)}$$

and $\kappa \in \mathbb{N}$ such that $\frac{1}{m} + 1 > \frac{3}{8}d + \frac{1}{\kappa}$ and put $p \geq \kappa + 1$ such that $1 - \frac{d}{2} > -\frac{p+1-\kappa}{2p}$.

Let the process v satisfy Assumption 4.2 and Assumption 4.6. In addition let us assume that the couple (u, v) satisfy Assumption 4.8 with p . Let w be a solution to

$$dw(t) = [r_v \Delta w(t) + u(t)v^2(t)] dt + \sigma_2(w(t)) dW_2(t), \quad w(0) = v_0, \quad v_0 \in H_m^{-\delta_1}(\mathcal{O}).$$

Then

(i) *there exists a number $r_0 > 0$ such that for any $r \leq r_0$ there exists a $\beta_0 > 0$, $C_0 > 0$ and $\delta_1, \delta_2, \delta_3 \geq 0$ and $\delta_2 \in (0, 1)$ with*

$$\mathbb{E} \|w\|_{L^{m_0}(0, T; H_m^r)}^{m_0} \leq \tilde{C}_2(T) \left\{ \mathbb{E} |v_0|_{H_m^{-\delta_0}}^{m_0} + \mathbb{E} |u_0|_{L^{p+1}}^{p+1} + C_0 T^{\beta_0} K_1^{\delta_1} K_2^{\delta_2} K_3^{\delta_3} \right\};$$

(ii) *there exists a number $\alpha_0 > 0$ such that for any $\alpha \leq \alpha_0$ there exists a $\beta_0 > 0$, $C_0 > 0$ and $\delta_1, \delta_2, \delta_3 \geq 0$ with*

$$\mathbb{E} \|w\|_{\mathbb{W}_\alpha^{m_0}(0, T; H_m^{-\sigma})}^{m_0} \leq \tilde{C}_2(T) \left\{ \mathbb{E} |v_0|_{H_m^{-\delta_0}}^{m_0} + C_0 T^{\beta_0} K_1^{\delta_1} K_2^{\delta_2} K_3^{\delta_3} \right\}.$$

Proof. First, we get by the analyticity of the semigroup for $\delta_0, \delta_1 \geq 0$

$$\begin{aligned} \mathbb{E} \|w\|_{L^{m_0}(0, T; H_m^r)}^{m_0} & \leq \int_0^T \left\{ t^{-2m_0\delta_0} \mathbb{E} |v_0|_{H_m^{-\delta_0}}^{m_0} + \mathbb{E} \left| \int_0^t e^{-r_v \Delta(t-s)} u(s)v^2(s) ds \right|_{H_m^r}^{m_0} \right. \\ & \quad \left. + \mathbb{E} \left| \int_0^t e^{-r_v \Delta(t-s)} \sigma_2(w(s)) dW_2(s) \right|_{H_m^r}^{m_0} \right\} dt \\ & =: I + II + III. \end{aligned}$$

Since

$$2m_0\delta_0 < 1,$$

the first term, i.e. I is bounded. In particular, we have

$$(4.11) \quad I \leq T^{1-2m_0\delta_0} \mathbb{E} |v_0|_{H_m^{-\delta_0}}^{m_0}.$$

Let us continue with the second term. The smoothing property of the semigroup gives for any $\delta_1 \geq 0$

$$II \leq \mathbb{E} \int_0^T \left(\int_0^t (t-s)^{-\frac{1}{2}(r+\delta_1)} |u(s)v^2(s)|_{H_m^{-\delta_1}} ds \right)^{m_0} dt.$$

Using the Sobolev embedding $L^1(\mathcal{O}) \hookrightarrow H_m^{-\delta_1}(\mathcal{O})$, where $\delta_1 = \frac{3}{4}d$, we get

$$II \leq \mathbb{E} \int_0^T \left(\int_0^t (t-s)^{-\frac{1}{2}(r+\delta_1)} |u(s)v^2(s)|_{L^1} ds \right)^{m_0} dt.$$

Supposing $l\frac{\delta_1+r_0}{2} < 1$ the Young inequality for convolutions gives for

$$(4.12) \quad \frac{1}{m_0} + 1 = \frac{1}{l} + \frac{1}{\kappa} \quad \text{and} \quad \beta_0 = \frac{1}{l} - \frac{1}{2}(r + \delta_1)$$

$$(4.13) \quad II \leq C_0 T^{\beta_0} \mathbb{E} \left(\int_0^T |u(s)v^2(s)|_{L^1}^\kappa ds \right)^{\frac{1}{\kappa}}.$$

In case $\kappa < p+1$ and $\rho \geq -\frac{p+1-\kappa}{2p}$, we have

$$II \leq C_0 T^{\beta_0} \left(\mathbb{E} |u_0|_{L^{p+1}}^{p+1} + K_2^{\frac{1}{2} \frac{p+1-\kappa}{p}} \right).$$

In case

$$d < \frac{4(1+3\gamma)}{3(1+\gamma)}$$

the entity $\frac{1}{m_0} + 1 - \frac{3}{8}d > 0$ and one can find a number $\kappa < p+1$ such that (4.12) is satisfied. In addition setting $p = 2\kappa + 1$, $\frac{1}{2} \frac{p+1-\kappa}{p} \leq \frac{1}{4}$.

Next, let us investigate III . We treat the stochastic term by applying Corollary 3.5-(ii) of [5], from which it follows for $\tilde{\sigma} + r < 1$

$$\mathbb{E} \int_0^T \left| \int_0^t e^{-r_v \Delta(t-s)} \sigma_2(w(s)) dW_2(s) \right|_{H_m^r}^{m_0} dt \leq T^\gamma \mathbb{E} \int_0^T |w(t)|_{H_m^{-\tilde{\sigma}}}^{m_0} dt.$$

Due to the Sobolev embedding and interpolation if $\tilde{\sigma} > \frac{d}{2} - \frac{d}{m} - \tilde{\rho}$ there exists some $\theta \in (0, 1)$ such that

$$|w(t)|_{H_m^{-\tilde{\sigma}}} \leq |w(t)|_{L^m}^\theta |w(t)|_{H_m^{\tilde{\rho}}}^{1-\theta} \leq C |w(t)|_{L^m}^\theta |w(t)|_{H_2^{\tilde{\rho}}}^{1-\theta}$$

Thus, if $\tilde{\rho}$ satisfies (4.7), we get for $\gamma > \frac{d-2\rho}{2(2+\rho)}$

$$\begin{aligned} \mathbb{E} \int_0^T |w(t)|_{H_m^{-\tilde{\sigma}}}^{m_0} dt &\leq C \mathbb{E} \left\{ \left[\int_0^T |w(t)|_{H_2^{\tilde{\rho}}}^{m_0} dt \right]^{1-\theta} \left[\int_0^T |u(t)|_{L^m}^{m_0} dt \right]^\theta \right\} \\ &\leq C \left[\mathbb{E} \int_0^T |w(t)|_{H_2^{\tilde{\rho}}}^{m_0} dt \right]^{1-\theta} \left[\mathbb{E} \int_0^T |w(t)|_{L^m}^{m_0} dt \right]^\theta \\ &\leq C(\varepsilon) K_2 + \varepsilon \mathbb{E} \|w\|_{L^{m_0}(0,T;L^m)}^{m_0}. \end{aligned}$$

Choosing $\varepsilon > 0$ sufficiently small and subtracting $\varepsilon \mathbb{E} \|w\|_{L^{m_0}(0,T;L^m)}^{m_0}$ on both sides, (i) follows.

In the next lines we will show item (ii). Note, that for $s < t$

$$w(t) - w(s) = (e^{-r_v \Delta(t-s)} - \text{Id})w(s) + \int_s^t e^{-r_v \Delta(t-\tilde{s})} u(\tilde{s}) v^2(\tilde{s}) d\tilde{s} + \int_0^s e^{-r_v \Delta(t-\tilde{s})} w(\tilde{s}) dW_2(\tilde{s}).$$

Substituting it in the definition of $\mathbb{W}^{\alpha, m_0}([0, T]; H_m^{-\sigma}(\mathcal{O}))$ we can write

$$\begin{aligned} &\int_0^T \int_0^T \frac{|w(t) - w(s)|_{H_m^{-\sigma}}^{m_0}}{|t-s|^{1+\alpha m_0}} ds dt \\ &\leq 2 \int_0^T \int_0^t \frac{|w(t) - w(s)|_{H_m^{-\sigma}}^{m_0}}{|t-s|^{1+\alpha m_0}} ds dt \\ &\leq 2 \int_0^T \int_0^t \frac{|[e^{-r_v \Delta(t-s)} - \text{Id}]w(s)|_{H_m^{-\sigma}}^{m_0}}{|t-s|^{1+\alpha m_0}} ds dt + 2 \int_0^T \int_0^t \frac{|\int_s^t e^{-r_v \Delta(t-\tilde{s})} u(\tilde{s}) v^2(\tilde{s}) d\tilde{s}|_{H_m^{-\sigma}}^{m_0}}{|t-s|^{1+\alpha m_0}} ds dt \\ &\quad + 2 \int_0^T \int_0^t \frac{|\int_s^t e^{-r_v \Delta(t-\tilde{s})} w(\tilde{s}) dW_2(\tilde{s})|_{H_m^{-\sigma}}^{m_0}}{|t-s|^{1+\alpha m_0}} ds dt =: IV + V + VI. \end{aligned}$$

First we consider the first term IV . First, let us note that by Proposition 4.12-(i) we know that there exists a number $r > 0$ such that

$$\mathbb{E} \int_0^T |w(s)|_{H_m^r}^{m_0} ds \leq C(T) \left(\mathbb{E}|v_0| + \mathbb{E}|u_0|_{L^{p+1}}^{p+1} + R_1 \right).$$

Since $|(e^{-r_v \Delta(t-s)} - \text{Id})x|_{H_m^{-\sigma}} = C \int_0^{t-s} \tilde{s}^{2(r+\sigma)-1} d\tilde{s} |x|_{H_m^r} \leq C(t-s)^{2(r+\sigma)} |x|_{H_m^r}$, we can infer

$$\dots \leq 2 \int_0^T \int_0^t \frac{(t-s)^{2(r+\sigma)m_0} |w(s)|_{H_m^r}^{m_0}}{|t-s|^{1+\alpha m_0}} ds dt.$$

The Young inequality for convolutions (with all exponents equal to 1) gives

$$\dots \leq \left(\int_0^T s^{m_0 2(r+\delta)-1-\alpha m_0} ds \right) \cdot \left(\int_0^T |w(s)|_{H_m^r}^{m_0} ds \right).$$

In particular, since $2(r+\delta) > \alpha$, the right hand side is bounded.

Now, we consider the second term V . In particular, we consider

$$\int_0^T \int_0^t \frac{|\int_s^t e^{-r_v \Delta(t-\tilde{s})} u(\tilde{s}) v^2(\tilde{s}) d\tilde{s}|_{H_m^{-\sigma}}^{m_0}}{|t-s|^{1+\alpha m_0}} ds dt.$$

The Cauchy Schwarz inequality gives for any $m_0 > 1$

$$\begin{aligned} \dots &\leq \left| \int_s^t e^{-r_v \Delta(t-r)} u(r) v^2(r) dr \right|_{H_m^{-\sigma}} \\ &\leq (t-s)^{\frac{1}{m'}} \left(\int_s^t |u(\tilde{s}) v^2(\tilde{s})|_{H_m^{-\sigma}}^{m_0} d\tilde{s} \right). \end{aligned}$$

Hence,

$$\begin{aligned} &\int_0^T \int_0^t \frac{\left| \int_s^t e^{-r_v \Delta(t-\tilde{s})} u(\tilde{s}) v^2(\tilde{s}) d\tilde{s} \right|_{H_m^{-\sigma}}^{m_0}}{|t-s|^{1+\alpha m_0}} ds dt \\ &\leq \int_0^T \int_0^t \frac{(t-s)^{\frac{m_0}{m'}} \int_s^t |u(\tilde{s}) v^2(\tilde{s})|_{H_m^{-\sigma}}^{m_0} d\tilde{s}}{|t-s|^{1+\alpha m_0}} ds dt \\ &\leq \left[\int_0^T \int_0^t (t-s)^{\frac{m_0}{m'}-1-\alpha m_0} ds dt \right] \left[\int_0^T |u(\tilde{s}) v^2(\tilde{s})|_{H_m^{-\sigma}}^{m_0} d\tilde{s} \right]. \end{aligned}$$

Taking into account that $\alpha < \frac{1}{m_0'}$, Integration and the Sobolev embedding gives

$$\begin{aligned} \dots &\leq \left[\int_0^T t^{\frac{m_0}{m'}-\alpha m_0} dt \right] \int_0^T |u(r) v^2(r)|_{H_m^{-\tilde{\sigma}}}^{m_0} dr \\ &\leq T^{\frac{m_0}{m'}+1-\alpha m_0} \int_0^T |u(r) v^2(r)|_{L^1}^{m_0} dr. \end{aligned}$$

It remains to give an estimate to the second term. The same computations as in Proposition 4.12 part (i), i.e. starting at (4.13), we can show the assertion.

Next, let us investigate VI . Here, the Burkholder-Davis-Gundy inequality gives

$$\mathbb{E} \left| \int_s^t e^{-r_v \Delta(t-\tilde{s})} \sigma_2(w(\tilde{s})) dW_2(\tilde{s}) \right|_{H_m^{-\sigma}}^{m_0} \leq (t-s)^{m_0-2} \mathbb{E} \int_s^t |w(\tilde{s})|_{H_m^{-\tilde{\sigma}}}^{m_0} d\tilde{s}.$$

In this way we get

$$\begin{aligned} \mathbb{E} VI &\leq 2 \int_0^T \int_0^t \frac{|t-s|^{m_0-2}}{|t-s|^{1+\alpha m_0}} \mathbb{E} \int_s^t |w(\tilde{s})|_{H_m^{-\tilde{\sigma}}}^{m_0} d\tilde{s} ds dt \\ &\leq 2 \int_0^T \int_0^t |t-s|^{m_0(1-\alpha)-3} \left(\mathbb{E} \int_s^t |w(\tilde{s})|_{H_m^{-\tilde{\sigma}}}^{m_0} d\tilde{s} \right) ds dt \end{aligned}$$

Applying the Hölder inequality, we show that

$$\mathbb{E} VI \leq C(T) \mathbb{E} \int_0^T |w(s)|_{H_m^{-\tilde{\sigma}}}^{m_0} ds.$$

Collecting all together gives (ii). □

Proposition 4.13 (Technical proposition). *Let us assume that (u, v) satisfies Assumptions 4.2, 4.6 and 4.8. Then, for any $q \geq 1$, there exists some $\delta_1, \delta_2 > 0$ such that*

$$\begin{aligned} & \mathbb{E} \int_0^T |u_1(s)v_1^2(s) - u_2(s)v_2^2(s)|_{L^1}^q ds \\ & \leq C(K_1, K_2, K_3, T) \left(\left(\mathbb{E} \|v_1 - v_2\|_{L^{m_0}(0, T; L^m)}^{m_0} \right)^{\delta_1} + \left(\mathbb{E} \|u_1 - u_2\|_{L^{\gamma+1}(0, T; L^{\gamma+1})}^{\gamma+1} \right)^{\delta_2} \right). \end{aligned}$$

Proof. Observe, for any $n \in \mathbb{N}$ and any $a, b \geq 0$, we have

$$\begin{aligned} |a - b| & \leq \left| \sum_{k=1}^n a^{\frac{k-1}{n}} (a^{\frac{1}{n}} - b^{\frac{1}{n}}) b^{\frac{n-k}{n}} \right| \\ & \leq |a - b|^{\frac{1}{n}} \sum_{k=1}^n a^{\frac{k-1}{n}} b^{\frac{n-k}{n}} \leq C |a - b|^{\frac{1}{n}} (n-1) \left(a^{\frac{n-1}{n}} + b^{\frac{n-1}{n}} \right). \end{aligned}$$

where the last inequality follows from an application of Young's inequality.

Note, that

$$\int_0^T |u_1(s)v_1^2(s) - u_2(s)v_2^2(s)|_{L^1}^q ds \leq \int_0^T \left| |v_1^2(s)u_1(s) - v_2^2(s)u_2(s)|^{\frac{1}{n}} \Psi(s) \right|_{L^1}^q ds$$

where

$$(4.14) \quad \Psi(s) = (n-1) \left((u_1 v_1^2)^{\frac{n-1}{n}}(s) + (u_2 v_2^2)^{\frac{n-1}{n}}(s) \right)$$

First, note that

$$\begin{aligned} & |v_1^2(s, x)u_1(s, x) - v_2^2(s, x)u_2(s, x)| \\ & \leq |v_1(s, x) - v_2(s, x)|v_1(s, x) + v_2(s, x)|u_1(s, x)| + |v_1^2(s, x) + v_2^2(s, x)||u_1(s, x) - u_2(s, x)|. \end{aligned}$$

Let $n \in \mathbb{N}$ be that large such that $\frac{2q}{m_0 n} + \frac{q}{n(\gamma+1)} + \frac{q}{\kappa} \leq 1$. The Hölder inequality gives

$$\begin{aligned} & \left| |v_1^2(s)u_1(s) - v_2^2(s)u_2(s)|^{\frac{1}{n}} \Psi(s) \right|_{L^1} \leq \left| |v_1^2(s)u_1(s) - v_2^2(s)u_2(s)|^{\frac{1}{n}} \right|_{L^n} |\Psi(s)|_{L^{\frac{n}{n-1}}} \\ & \leq \| |v_1(s) - v_2(s)|(v_1(s) + v_2(s)) \|_{L^m} \| |u_1(s)| + |u_1(s) - u_2(s)|v_2^2(s) \|_{L^1}^{\frac{1}{n}} |\Psi(s)|_{L^{\frac{n}{n-1}}} \\ & \leq \left(\| |v_1(s) - v_2(s)|^{\frac{1}{n}} \|_{L^m} \| (v_1(s) + v_2(s)) \|_{L^m}^{\frac{1}{n}} \| |u_1(s)|^{\frac{1}{n}} \|_{L^{\gamma+1}} \right. \\ & \quad \left. + \| |u_1(s) - u_2(s)|^{\frac{1}{n}} \|_{L^{\gamma+1}} \| |v_2(s)|^{\frac{2}{n}} \|_{L^m} \right) |\Psi(s)|_{L^{\frac{n}{n-1}}}. \end{aligned}$$

Next, we obtain

$$\begin{aligned}
& \mathbb{E} \int_0^T |u_1(s)v_1^2(s) - u_2(s)v_2^2(s)|_{L^1}^q ds \\
& \leq \mathbb{E} \int_0^T \left(|v_1(s) - v_2(s)|_{L^m}^{\frac{q}{n}} |(v_1(s) + v_2(s))|_{L^m}^{\frac{q}{n}} |u_1(s)|_{L^{\gamma+1}}^{\frac{q}{n}} \right. \\
& \quad \left. + |u_1(s) - u_2(s)|_{L^{\gamma+1}}^{\frac{q}{n}} |v_2(s)|_{L^m}^{\frac{2q}{n}} \right) |\Psi(s)|_{L^{\frac{n}{n-1}}}^q ds \\
& \leq \mathbb{E} \left\{ \left[\int_0^T |v_1(s) - v_2(s)|_{L^m}^{m_0} ds \right]^{\frac{q}{m_0 n}} \left[\int_0^T |v_1(s) + v_2(s)|_{L^m}^{m_0} ds \right]^{\frac{q}{m_0 n}} \right. \\
& \quad \left. \times \left[\sup_{0 \leq s \leq T} |u_1(s)|_{L^{\gamma+1}}^{\gamma+1} \right]^{\frac{q}{(\gamma+1)n}} \left[\int_0^T |\Psi(s)|_{L^{\frac{n}{n-1}}}^{\kappa} ds \right]^{\frac{q}{\kappa}} \right\} \\
& \quad + \mathbb{E} \left\{ \left[\int_0^T |u_1(s) - u_2(s)|_{L^{\gamma+1}}^{\gamma+1} ds \right]^{\frac{q}{2n}} \left[\int_0^T |v_2(s)|_{L^m}^q ds \right]^{\frac{2}{n}} \left[\int_0^T |\Psi(s)|_{L^{\frac{n}{n-1}}}^{\kappa} ds \right]^{\frac{q}{\kappa}} \right\}
\end{aligned}$$

Rewriting and applying the Hölder inequality again gives

$$\begin{aligned}
\cdots & \leq \left\{ \left[\mathbb{E} \left[\sup_{0 \leq s \leq T} |u_1(s)|_{L^{\gamma+1}}^{\gamma+1} \right] \right]^{\frac{q}{(\gamma+1)n}} \right. \\
& \quad \times \left[\mathbb{E} \|v_1 - v_2\|_{L^{m_0}(0,T;L^m)}^{m_0} \right]^{\frac{q}{m_0 n}} \left[\mathbb{E} \|v_1 + v_2\|_{L^{m_0}(0,T;L^m)}^{m_0} \right]^{\frac{q}{m_0 n}} \mathbb{E} \left[\|\Psi\|_{L^{\kappa}(0,T;L^{\frac{n}{n-1}})}^{\kappa} \right]^{\frac{q}{\kappa}} \\
& \quad \left. + \left[\mathbb{E} \|u_1 - u_2\|_{L^{\gamma+1}(0,T;L^{\gamma+1})}^{\gamma+1} \right]^{\frac{q}{n}} \left[\mathbb{E} \|v_2\|_{L^{m_0}(0,T;L^m)}^{m_0} \right]^{\frac{2q}{n}} \left[\mathbb{E} \|\Psi\|_{L^{\kappa}(0,T;L^{\frac{n}{n-1}})}^{\kappa} \right]^{\frac{q}{\kappa}} \right\}.
\end{aligned}$$

Reminding definition (4.14) we know

$$\|\Psi\|_{L^{\kappa}(0,T;L^{\frac{n}{n-1}})} \leq C(n-1) \left\{ \|u_1 v_1^2\|_{L^{\kappa}(0,T;L^1)} + \|u_2 v_2^2\|_{L^{\kappa}(0,T;L^1)} \right\}.$$

Now, by Proposition 4.10 and taking into account that $\rho \geq -\frac{1}{\kappa} \frac{r-\kappa}{r-1} = -\frac{1}{\kappa} \frac{p+1-\kappa}{p}$ we can infer that

$$\mathbb{E} \int_0^t |\Psi(s)|_{L^1}^{\kappa} ds \leq K_3 + K_1^{\kappa} \frac{1}{p} \frac{p+1-\kappa}{\kappa}.$$

Due to the Assumptions 4.2, 4.6 and 4.8, we know that there exists some $\delta_1, \delta_2 > 0$ such that

$$\begin{aligned}
& \mathbb{E} \|u_1 v_1^2 - u_2 v_2^2\|_{L^q(0,T;L^1)}^q \\
& \leq C(K_1, K_2, K_3, T) \left(\left(\mathbb{E} \|v_1 - v_2\|_{L^{m_0}(0,T;L^m)}^{m_0} \right)^{\delta_1} + \left(\mathbb{E} \|u_1 - u_2\|_{L^{\gamma+1}(0,T;L^{\gamma+1})}^{\gamma+1} \right)^{\delta_2} \right).
\end{aligned}$$

□

The next proposition gives the continuity property.

Proposition 4.14. *Let the couples (u_1, v_1) and (u_2, v_2) satisfy the assumptions 4.2, 4.6, and 4.8 with constants K_1, K_2 , and K_3 , respectively. Let w_1 and w_2 be the solutions to*

$$dw_j(t) = [r_v \Delta w_j(t) + u_j(t)v_j^2(t)] dt + \sigma_2(u_j(t))dW_2(t), \quad t \in (0, T], \quad w_j(0) = v_0, j = 1, 2.$$

Then, under the assumption of Theorem 3.7 there exists a constant $C = C(K_1, K_2, K_3, T) > 0$ and numbers $\delta_1, \delta_2 > 0$ such that

$$\begin{aligned} & \mathbb{E} \|w_1 - w_2\|_{L^{m_0}(0, T; L^m)} \\ & \leq C(K_1, K_2, K_3, T) \left\{ \left[\mathbb{E} \|v_1 - v_2\|_{L^{m_0}(0, T; L^m)}^{m_0} \right]^{\delta_1} + \left[\mathbb{E} \|u_1 - u_2\|_{L^{\gamma+1}(0, T; L^{\gamma+1})}^{\gamma+1} \right]^{\delta_2} \right\}. \end{aligned}$$

Proof. Now, let us start with the proof. First, we get by the analyticity of the semigroup for $\delta > 0$

$$(4.15) \quad \begin{aligned} \mathbb{E} \int_0^T |w_1(t) - w_2(t)|_{L^m}^{m_0} dt & \leq \mathbb{E} \int_0^T \int_0^t (t-s)^{-\frac{\delta}{2}} |u_1(s)v_1^2(s) - u_2(s)v_2^2(s)|_{H_m^{-\delta}} ds dt \\ & \quad + \left| \int_0^T \mathbb{E} \int_0^t (\sigma_2(w_1(s)) - \sigma_2(w_2(s))) dW_2(s) \right|_{L^m} dt \\ & =: I + II. \end{aligned}$$

Next, we use the Sobolev embedding $L^1(\mathcal{O}) \hookrightarrow H_m^{-\delta}(\mathcal{O})$, where $\delta \geq d(1 - \frac{1}{m}) = d\frac{2+\gamma}{1+\gamma}$

$$I \leq \mathbb{E} \int_0^t (t-s)^{-\frac{\delta}{2}} |u_1(s)v_1^2(s) - u_2(s)v_2^2(s)|_{L^1} ds.$$

Next, the Young inequality for convolution gives for $\frac{1}{m_0} + 1 = \frac{1}{l} + \frac{1}{q}$, $\frac{\delta}{2} < \frac{1}{l}$, and $q > \frac{4(1+\gamma)}{2+6\gamma-d(2+\gamma)}$

$$I \leq T^{\beta_0} \mathbb{E} \left(\int_0^T |u_1(s)v_1^2(s) - u_2(s)v_2^2(s)|_{L^1}^q ds \right)^{\frac{m_0}{q}}.$$

Now, by Proposition 4.13 and taking into account that $\rho \geq -\frac{1}{\kappa} \frac{r-\kappa}{r-1} = -\frac{1}{\kappa} \frac{p+1-\kappa}{p}$ we can infer that there exists a constant $C(K_1, K_2, K_3, T) > 0$ and $\delta_1, \delta_2 > 0$ such that

$$I \leq C(K_1, K_2, K_3, T) \left(\left(\mathbb{E} \|v_1 - v_2\|_{L^{m_0}(0, T; L^m)}^{m_0} \right)^{\delta_1} + \left(\mathbb{E} \|u_1 - u_2\|_{L^{\gamma+1}(0, T; L^{\gamma+1})}^{\gamma+1} \right)^{\delta_2} \right).$$

It remains to tackle the second term II . This can be done by standard arguments using the Burkholder-Davis-Gundy inequality. \square

Finally we have to show the continuity of u with respect to v .

Proposition 4.15. *Let the couples (u_1, v_1) and (u_2, v_2) satisfy the assumptions 4.2, 4.6, and 4.8 with constants K_1, K_2 , and K_3 , respectively. Let w_1 and w_2 be the solutions to*

$$du_j(t) = [r_u \Delta u_j^{[\gamma]}(t) - u_j(t)v_j^2(t)] dt + \sigma_2(w_j(t)) dW_2(t), \quad t \in (0, T], \quad u_j(0) = u_0, \quad j = 1, 2.$$

Then, under the assumption of Theorem 3.7 there exists a constant $C = C(R_1, R_2, R_3, T) > 0$ and numbers $\delta_1, \delta_2 > 0$ such that

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq s \leq T} |u_1(s) - u_2(s)|_{H_2^{-1}}^2 \right] \\ & \leq C(K_1, K_2, K_3, T) \left\{ \left[\mathbb{E} \|v_1 - v_2\|_{L^{m_0}(0, T; L^m)}^{m_0} \right]^{\delta_1} + \left[\mathbb{E} \|u_1 - u_2\|_{L^{\gamma+1}(0, T; L^{\gamma+1})}^{\gamma+1} \right]^{\delta_2} \right\}. \end{aligned}$$

Proof. Applying the Itô formula and integration by parts gives

$$\begin{aligned} & |u_1(t) - u_2(t)|_{H_2^{-1}}^2 + 2r_u \int_0^t \langle u_1(s) - u_2(s), u_1^{[\gamma]}(s) - u_2^{[\gamma]}(s) \rangle ds \\ &= -2 \int_0^t \langle \nabla^{-1}(u_1(s) - u_2(s)), \nabla^{-1}(u_1(s)v_1^2(s) - u_2(s)v_2^2(s)) \rangle ds \\ &\quad + 2\sigma_1 \int_0^t \langle u_1(s) - u_2(s), (u_1(s) - u_2(s))dW_1(s) \rangle + \sigma_1^2 \int_0^t |u_1(s) - u_2(s)|_{H_2^{-1}}^2 ds, \end{aligned}$$

where $\nabla^{-1} := -(-\Delta)^{1/2}$. The Young inequality, Lemma A.1, and the embedding $L^1(\mathcal{O}) \hookrightarrow H_{\frac{\gamma+1}{\gamma}}^{-2}(\mathcal{O})$ give for $\varepsilon \in \left(0, 2^{\frac{2-\gamma}{\gamma+1}} r_u^{\frac{1}{\gamma+1}}\right]$,

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq s \leq t} |u_1(s) - u_2(s)|_{H_2^{-1}}^2 \right] + 2^{2-\gamma} r_u \mathbb{E} \int_0^t |u_1(s) - u_2(s)|_{L^{\gamma+1}}^{\gamma+1} ds \\ & \leq \varepsilon^{\gamma+1} \mathbb{E} \int_0^t |u_1(s) - u_2(s)|_{L^{\gamma+1}}^{\gamma+1} ds + C(\gamma) \varepsilon^{-\frac{\gamma+1}{\gamma}} \mathbb{E} \int_0^t |(-\Delta)^{-1}(u_1(s)v_1^2(s) - u_2(s)v_2^2(s))|_{L^{\frac{\gamma+1}{\gamma}}}^{\frac{\gamma+1}{\gamma}} ds \\ & \quad + 2\sigma_1 \mathbb{E} \int_0^t \langle \nabla^{-1}(u_1(s) - u_2(s)), \nabla^{-1}dW_1(s) \rangle + \sigma_1^2 \mathbb{E} \int_0^t |u_1(s) - u_2(s)|_{H_2^{-1}}^2 ds \\ & \leq \varepsilon^{\gamma+1} \mathbb{E} \int_0^t |u_1(s) - u_2(s)|_{L^{\gamma+1}}^{\gamma+1} ds + C(\gamma) \varepsilon^{-\frac{\gamma+1}{\gamma}} \mathbb{E} \int_0^t |u_1(s)v_1^2(s) - u_2(s)v_2^2(s)|_{L^1}^{\frac{\gamma+1}{\gamma}} ds \\ & \quad + C\sigma_1 \mathbb{E} \int_0^t |u_1(s) - u_2(s)|_{H_2^{-1}}^2 ds + \sigma_1^2 \mathbb{E} \int_0^t |u(s)_1 - u_2(s)|_{H_2^{-1}}^2 ds \\ & =: I + II + III + IV, \end{aligned}$$

where we have used the Burkholder-Davis-Gundy inequality. Term I cancels with the corresponding term on the left hand side. To tackle the second term, we will apply the technical Proposition 4.13. We apply Gronwall's lemma for terms III and IV and obtain the assertion. \square

APPENDIX A. AN ELEMENTARY INEQUALITY

Lemma A.1. *Let $\gamma > 1$, $x, y \in \mathbb{R}$. Then it holds that*

$$(A.1) \quad (x^{[\gamma]} - y^{[\gamma]})(x - y) \geq 2^{1-\gamma} |x - y|^{\gamma+1},$$

where $z^{[\gamma]} := |z|^{\gamma-1}z$ for $z \in \mathbb{R}$.

Proof. See e.g. [19, Lemma 3.1]. \square

APPENDIX B. POWERS AND MULTIPLICATION

Let $\sigma_p := d \max(0, \frac{1}{p} - 1)$ and $\sigma_{p,q} := d \max(0, \frac{1}{p} - 1, \frac{1}{q} - 1)$, where d is the dimension. Assume $s_1 < 0 < s_2$.

For the definition of the spaces $B_{p,q}^s$ and $F_{p,q}^s$, we refer to [24, 30, 33]. They translate to classical function spaces as in e.g. [30, Remark 2.1.1], in particular, $F_{p,2}^0 = L_p$, $1 < p < \infty$ (Lebesgue spaces), $F_{p,2}^m = W_p^m$, $m \in \mathbb{N}$ (Sobolev spaces), $1 < p < \infty$ and $F_{p,2}^s = H_p^s$, $s \in \mathbb{R}$, $1 < p < \infty$ (fractional Sobolev spaces).

Theorem B.1. *Assume $s = s_1 \leq s_2$, $s_1 + s_2 > d \cdot \max(0, \frac{1}{p} - 1)$, and $q \geq \max(q_1, q_2)$. Then*

- *if $s_2 > s_1$, then $F_{p,q_1}^{s_1}(\mathbb{R}^d) \cdot B_{\infty,q_2}^{s_2}(\mathbb{R}^d) \hookrightarrow F_{p,q_1}^{s_1}(\mathbb{R}^d)$;*
- *and, if $s_1 = s_2$ then $F_{p,q_1}^{s_1}(\mathbb{R}^d) \cdot B_{\infty,q_2}^{s_1}(\mathbb{R}^d) \hookrightarrow F_{p,q}^{s_1}(\mathbb{R}^d)$.*

Proof. See [24, p. 229]. □

Theorem B.2. *Let $s_1 \leq s_2$ and $s_1 + s_2 > d \cdot \max(0, \frac{1}{p} - 1)$.*

- *Assume $s_2 > \frac{d}{p}$ and $q \geq \max(q_1, q_2)$. Then, if $s_2 > s_1$, $F_{p,q_1}^{s_1}(\mathbb{R}^d) \cdot F_{p,q_2}^{s_2}(\mathbb{R}^d) \hookrightarrow F_{p,q_1}^{s_1}(\mathbb{R}^d)$; if $s_2 = s_1$, $F_{p,q_1}^{s_1}(\mathbb{R}^d) \cdot F_{p,q_2}^{s_2}(\mathbb{R}^d) \hookrightarrow F_{p,q}^{s_1}(\mathbb{R}^d)$;*
- *Let $s_1 = s_2 = \frac{d}{p}$ and $q \geq \max(q_1, q_2)$. If $0 < p \leq 1$, then $F_{p,q_1}^{s_1}(\mathbb{R}^d) \cdot F_{p,q_2}^{s_2}(\mathbb{R}^d) \hookrightarrow F_{p,q}^{s_1}(\mathbb{R}^d)$;*
- *If $s_2 < \frac{d}{p}$, then $F_{p,q_1}^{s_1}(\mathbb{R}^d) \cdot F_{p,q_2}^{s_2}(\mathbb{R}^d) \hookrightarrow F_{p,q}^{s_1+s_2-\frac{d}{p}}(\mathbb{R}^d)$.*

Proof. See [24, p. 190]. □

APPENDIX C. A COMPACTNESS RESULT

Let B be a separable Banach space, $I \subset \mathbb{R}_+$ be an interval. Recall that

$$\mathbb{L}^p(I; B) = \left\{ u : I \rightarrow B : u \text{ measurable and } \int_I |u(t)|_B^p dt < \infty \right\}.$$

Given $p > 1$, $\alpha \in (0, 1)$, let $\mathbb{W}^{\alpha,p}(I; B)$ be the Sobolev space of all $u \in \mathbb{L}^p(I; B)$ such that

$$\int_I \int_{I \cap [t, t+1]} \frac{|u(t) - u(s)|_B^p}{|t - s|^{1+\alpha p}} ds dt < \infty.$$

equipped with the norm

$$\|u\|_{\mathbb{W}^{\alpha,p}(I; B)} := \left(\int_I \int_{I \cap [t, t+1]} \frac{|u(t) - u(s)|_B^p}{|t - s|^{1+\alpha p}} ds dt \right)^{\frac{1}{p}}.$$

Theorem C.1. *Let $B_0 \subset B \subset B_1$ be Banach spaces, B_0 and B_1 reflexive, with compact embedding of B_0 in B . Let $p \in (1, \infty)$ and $\alpha \in (0, 1)$ be given. Let X be the space*

$$X = \mathbb{L}^p([0, T]; B_0) \cap \mathbb{W}^{\alpha,p}([0, T]; B_1).$$

Then the embedding of X in $\mathbb{L}^p([0, T]; B)$ is compact.

Proof. See [31, p. 86, Corollary 5] or [11, Theorem 2.1]. □

REFERENCES

- [1] V. Barbu, G. Da Prato, and M. Röckner. Existence and uniqueness of nonnegative solutions to the stochastic porous media equation. *Indiana Univ. Math. J.*, 57(1):187–211, 2008.
- [2] V. Barbu, G. Da Prato, and M. Röckner. *Stochastic porous media equations*. Lecture Notes in Mathematics. Springer International Publishing, 2016.
- [3] J. Bergh and J. Löfström. *Interpolation Spaces. An Introduction*, volume 223 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin-New York, 1976.

- [4] Z. Brzeźniak. Stochastic partial differential equations in M-type 2 Banach spaces. *Potential Anal.*, 4(1):1–45, 1995.
- [5] Z. Brzeźniak. On stochastic convolution in Banach spaces and applications. *Stochastics Stochastics Rep.*, 61(3-4):245–295, 1997.
- [6] Z. Brzeźniak, E. Hausenblas, and P. A. Razafimandimby. Stochastic reaction-diffusion equations driven by jump processes. *Potential Anal.*, 49:131–201, 2018.
- [7] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 152 of *Encyclopedia of mathematics and its applications*. Cambridge University Press, 2nd edition, 2014.
- [8] G. Dhariwal, F. Huber, A. Jüngel, C. Kuehn, and A. Neamțu. Global martingale solutions for quasilinear SPDEs via the boundedness-by-entropy method. *Preprint*, pages 1–33, 2019. <https://arxiv.org/abs/1909.08892>.
- [9] E. Dulos, J. Boissonade, J. J. Perraud, B. Rudovics, and P. De Kepper. Chemical morphogenesis: turing patterns in an experimental chemical system. *Acta Biotheor.*, 44(3–4):249–261, 1996.
- [10] L. C. Evans. *An Introduction to Stochastic Differential Equations*. American Mathematical Society, Providence, 2012.
- [11] F. Flandoli and D. Gałtarek. Martingale and stationary solutions for stochastic Navier-Stokes equations. *Probab. Theory Related Fields*, 102(3):367–391, 1995.
- [12] A. Granas and J. Dugundji. *Fixed point theory*. Springer, 2003.
- [13] P. Gray and S. K. Scott. Autocatalytic reactions in the isothermal, continuous stirred tank reactor: isolas and other forms of multistability. *Chemical Engineering Science*, 38(1):29–43, 1983.
- [14] E. Hausenblas, T. A. Randrianasolo, and M. Thalhaammer. Theoretical study and numerical simulation of pattern formation in the deterministic and stochastic Gray-Scott equations. *J. Comput. Appl. Math.*, 364:112335, 27, 2020.
- [15] B. J. Kealy and D. J. Wollkind. A nonlinear stability analysis of vegetative turing pattern formation for an interaction–diffusion plant–surface water model system in an arid flat environment. *Bulletin of Mathematical Biology*, 74(4):803–833, Apr 2012.
- [16] C. A. Klausmeier. Regular and irregular patterns in semi-arid vegetation. *Science*, 284:1826–1828, 1999.
- [17] P. Kotelenez. Comparison methods for a class of function valued stochastic partial differential equations. *Probab. Theory Related Fields*, 93(1):1–19, 1992.
- [18] S. A. Levin and L. A. Segel. Hypothesis for origin of planktonic patchiness. *Nature*, 259:659, 1976.
- [19] W. Liu. Harnack inequality and applications for stochastic evolution equations with monotone drifts. *Journal of Evolution Equations*, 9(4):747–770, 2009.
- [20] W. Liu and M. Röckner. *Stochastic Partial Differential Equations: An Introduction*. Universitext. Springer, 2015.
- [21] B. Maslowski and D. Nualart. Evolution equations driven by a fractional Brownian motion. *Journal of Functional Analysis*, 202(1):277–305, 2003.
- [22] J. Murray. How the leopard gets its spots. *Scientific American*, 258:80–87, 1988.
- [23] J. Murray. *Mathematical biology: Spatial models and biomedical applications*. Springer Science & Business Media, 2003.

- [24] T. Runst and W. Sickel. *Sobolev spaces of fractional order, Nemytskij operators, and nonlinear partial differential equations*. Walter de Gruyter, 1996.
- [25] L. A. Segel and J. L. Jackson. Dissipative structure: An explanation and an ecological example. *J. Theor. Biol.*, 37:545–592, 1972.
- [26] J. A. Sherratt. An analysis of vegetation stripe formation in semi-arid landscapes. *J. Math. Biol.*, 51:183–197, 2005.
- [27] J. A. Sherratt. Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments I. *Nonlinearity*, 23:2657–2675, 2010.
- [28] J. A. Sherratt and G. J. Lord. Nonlinear dynamics and pattern bifurcations in a model for vegetation stripes in semi-arid environments. *Theor. Popul. Biol.*, 71:1–11, 2007.
- [29] N. Shigesada, K. Kawasaki, and E. Teramoto. Spatial segregation of interacting species. *J. Theor. Biol.*, 79(1):83–99, 1979.
- [30] W. Sickel and H. Triebel. Hölder inequalities and sharp embeddings in function spaces of B_{pq}^s and F_{pq}^s type. *Zeitschrift für Analysis und ihre Anwendungen*, 14(1):105–140, 1995.
- [31] J. Simon. Compact sets in the space $L^p(0, T; B)$. *Annali di Matematica Pura ed Applicata*, 146(1):65–96, 1986.
- [32] G. Tessitore and J. Zabczyk. Strict positivity for stochastic heat equations. *Stochastic Processes Appl.*, 77:83–98, 1998.
- [33] H. Triebel. *Interpolation theory, function spaces, differential operators*, volume 18 of *North-Holland Mathematical Library*. North-Holland, Amsterdam–New York–Oxford, 1978.
- [34] A. M. Turing. The chemical basis of morphogenesis. *Philosophical Transactions of the Royal Society of London B*, 237(641):37–72, 1952.
- [35] N. Ursino. The influence of soil properties on the formation of unstable vegetation patterns on hillsides of semiarid catchments. *Advances in Water Resources*, 28(9):956–963, 2005.
- [36] S. van der Stelt, A. Doelman, G. Hek, and J. D. M. Rademacher. Rise and fall of periodic patterns for a generalized Klausmeier–Gray–Scott model. *Journal of Nonlinear Science*, 23(1):39–95, 2012.
- [37] J. van Neerven, M. Veraar, and L. Weis. Stochastic integration in Banach spaces—a survey. In *Stochastic analysis: a series of lectures*, volume 68 of *Progr. Probab.*, pages 297–332. Birkhäuser/Springer, Basel, 2015.

MONTANUNIVERSITÄT LEOBEN, DEPARTMENT MATHEMATIK UND INFORMATIONSTECHNOLOGIE, FRANZ JOSEF STRASSE 18, 8700 LEOBEN, AUSTRIA

E-mail address: erika.hausenblas@unileoben.ac.at

UNIVERSITY OF HELSINKI, DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX 68 (GUSTAF HÄLLSTRÖMIN KATU 2), FI-00014 UNIVERSITY OF HELSINKI, FINLAND

UNIVERSITÄT AUGSBURG, INSTITUT FÜR MATHEMATIK, UNIVERSITÄTSSTRASSE 14, 86135 AUGSBURG, GERMANY

E-mail address: jonas.tolle@helsinki.fi