

# Gravity and Matters on a pure geometric thick polynomial $f(R)$ brane

Heng Guo <sup>a†</sup>

*School of Physics, Xidian University, Xi'an 710071, China*

Lang-Lang Wang<sup>‡</sup>

*School of Physics, Xidian University, Xi'an 710071, China and*

*School of General Education, Shanxi Institute of*

*Science and Technology, Jincheng 048000, China*

Chun-E Fu<sup>§</sup>

*School of Science, Xi'an Jiaotong University, Xi'an 710049, China*

Qun-Ying Xie<sup>¶</sup>

*School of Information Science and Engineering,*

*Lanzhou University, Lanzhou 730000, China*

(Dated: May 10, 2023)

## Abstract

In this paper, writing the most general form  $f(R) = \sum_{i=1}^n a_i R^i + \Lambda$  the solutions of the pure geometric thick  $f(R)$  brane are investigated. For the certain value of  $n$ , the analytical thick brane solution can be calculated, and when  $n = 3$ ,  $n = 4$ , and  $n = 10$ , the thick brane solutions are presented. The solutions are stable against linear tensor perturbations. The zero mode of gravity and scalar field can be localized on thick  $f(R)$  branes naturally. The zero mode of vector field and left-chiral fermion can be localized on thick  $f(R)$  branes by introducing the coupling with scalar curvature  $R$  of spacetime, and the massive resonant modes can be quasilocalized on the brane with the large coupling coefficients.

---

<sup>a</sup> Corresponding author

<sup>†</sup> hguo@xidian.edu.cn

<sup>‡</sup> wllsrr@163.com

<sup>§</sup> fuche13@mail.xjtu.edu.cn

<sup>¶</sup> xieqy@lzu.edu.cn

## I. INTRODUCTION

The idea that our four-dimensional (4D) Universe can be considered as a brane embedded in a higher-dimensional spacetime, can supply new insights for solving the gauge hierarchy problem [1–8] and the cosmological constant problem [9–21]. In the Randall-Sundrum (RS) braneworld model [22], the effective 4D gravity could be recovered even in the case of non-compact extra dimensions, however, singularities are present at the position of the branes. The smooth thick braneworld solutions are generally based on gravity coupled to bulk scalar fields [23–62]. For some comprehensive reviews about thick branes, please see Refs. [63–67]. There are also thick branes arising from pure geometry without the inclusion of bulk scalar fields at all [68–77].

In braneworld scenarios, the important issue is the localization of gravity and various bulk matter fields for the purpose of recovering the effective 4D gravity and building up the standard model. Generally the gravity and the scalar field zero mode can be localized on the brane naturally. For five-dimensional (5D) free spin-1 Abelian vector field, the zero mode can not be trapped on the Minkowski ( $\mathcal{M}_4$ ) brane, however, it can be localized on the thick de Sitter brane and Weyl thick brane [8, 80–83]. In some ways, by introducing the coupling with background geometry, the zero mode of the vector field can also be trapped on the Minkowski brane [84, 85]. For the spin-1/2 fermion, zero mode cannot be localized on the branes without introducing the coupling. By introducing the coupling between the fermion and background scalar fields or fermion and background geometry, the left-chiral fermion zero mode can be localized on the brane [8, 16, 48, 80–82, 86–131].

However, due to the fact that Einstein’s general relativity is not renormalizable, the effects of higher order curvature terms are suggested to consider.  $f(R)$  gravity theories, which the Lagrangians are proportional to some functions of the scalar curvature  $R$ , were created in the investigation of cosmology, and offer pure geometric explanations for cosmological inflation and dark energy in physics [132–139]. Nevertheless, there are studies devoted to embedding branes into various types of  $f(R)$  gravities [6, 41, 50, 57–60, 73, 74, 77–79, 140–150]. In Ref. [77], two cases of the thick brane solutions have been investigated in pure geometry  $f(R)$  gravities: one with a triangular  $f(R)$  and the other a simple polynomial  $f(R)$ .

In this paper, we intend to investigate the thick branes generated by only geometry under

pure  $f(R)$  gravity. The general form of  $f(R)$  would be considered, which is chosen as a  $n$ -th order polynomial of the scalar curvature  $R$ . When the degree of the polynomial  $f(R)$  is  $n = 3$ , the solution is equivalent to the one of Ref. [77], When  $n = 4$  and  $n = 10$ , the solutions are new solutions. These solutions will be presented in next section. The stability of tensor perturbations and the localization of gravity are discussed in Sec.III. The zero mode of gravity can be localized on the brane naturally. In Sec.IV, the zero mode of the scalar field can be trapped on the branes naturally. In Sec.V, by introducing the coupling between the vector field and the background geometry, the zero mode vector can be localized on the brane and the massive resonant Kaluza-Klein (KK) modes can also be quasilocalized on the brane. In Sec.VI, for a pure geometry brane, the coupling between the fermion and the background scalars can not be introduced, however, we introduce the coupling between the fermion and the background geometry, and the zero mode of the left-chiral fermion and the massive resonant KK modes can be localized and quasilocalized on the brane. Finally, the conclusion and discussion are given in Sec.VII.

## II. THE PURE GEOMETRIC THICK $f(R)$ BRANES

We start with the following 5D action for a pure geometric thick  $f(R)$  braneworld

$$S = \frac{1}{2\kappa_5^2} \int d^5x \sqrt{-g} f(R), \quad (1)$$

where  $\kappa_5^2 = 8\pi G_5$  with  $G_5$  being the 5D Newton constant,  $R$  is the 5D scalar curvature, and  $g = \det(g_{MN})$  is the determinant of the metric. Throughout this paper, capital Latin letters  $M, N, \dots = 0, 1, 2, 3, 5$  and Greek letters  $\mu, \nu, \dots = 0, 1, 2, 3$  are used to represent the bulk and brane indices, respectively. From this action, the 5D Einstein equation is given by

$$R_{MN} f_R - \frac{1}{2} g_{MN} f(R) + (g_{MN} \square - \nabla_M \nabla_N) f_R = 0, \quad (2)$$

where  $f_R \equiv \frac{df(R)}{dR}$ ,  $\square = g^{MN} \nabla_M \nabla_N$  is the 5D d'Alembert operator, and  $R_{MN}$  is the 5D Ricci tensor, defined in terms of the Riemann tensor  $R_{MN} = R_{MQN}^Q$ .

The line element of the  $\mathcal{M}_4$  brane is assumed as

$$ds^2 = g_{MN} dx^M dx^N = e^{2A(y)} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2, \quad (3)$$

where  $e^{2A(y)}$  is the warp factor,

$$\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1) \quad (4)$$

is the metric of the 4D Minkowski spacetime, and  $y$  denotes the extra dimensional coordinate. From the metric (3), the Ricci tensor and scalar curvature can be computed:

$$R_{\mu\nu} = -e^{2A(y)}(A'' + 4A'^2)\eta_{\mu\nu}, \quad R_{55} = -4A'' - 4A'^2, \quad (5)$$

$$R = -20A'^2 - 8A'', \quad (6)$$

where the prime denotes the derivatives with respect to extra dimensional coordinate  $y$ . The Einstein equations (2) can be rewritten as:

$$f(R) + 2f_R(4A'^2 + A'') - 6f'_R A' - 2f''_R = 0, \quad (7a)$$

$$-f(R) - 8f_R(A'' + A'^2) + 8f_R A' = 0. \quad (7b)$$

Above two equations are not independent, because the left side of Eq. (2) is divergence free [136]. Hence, we choose to solve Eq. (7a) in the following discussion.

In this paper, we are interested in investigating the general form of  $f(R)$ , chosen as a  $n$ -th order polynomial of the scalar curvature  $R$ :

$$f(R) = \sum_{i=1}^n a_i R^i + \Lambda, \quad (8)$$

where the  $a_i$  coefficients have the appropriate dimensions,  $\Lambda$  is 5D cosmological constants, and the parameter  $n$  should be an integer.

According to Eqs. (6) and (8), one can easily prove that  $f(R)$  includes  $(A'')^n$  and  $(A')^{2n}$ ,  $f'_R$  includes  $A'''$  and  $f''_R$  includes  $A''''$ . By taking  $f(R)$ ,  $f'_R$  and  $f''_R$  back into Eqs. (7), obviously, the Einstein equations are high-order differential equations, and very hard to be solve. In order to find a solution, we begin with a simple  $A(y)$ , which consider as [77, 150]

$$A(y) = -\delta \ln(\cosh(ky)) \quad (9)$$

with  $\delta$  a dimensionless positive constant, and  $k$  another positive constant with the dimension of length inverse. At the infinity, the behavior of the  $A(y \rightarrow \infty)$  is

$$A(y \rightarrow \infty) \rightarrow -\delta k |y|. \quad (10)$$

Defining a new parameter by  $\tilde{k} = \delta k$ , it is easy to proof that the warp factor

$$e^{2A(y \rightarrow \infty)} \rightarrow e^{-2\tilde{k}|y|}, \quad (11)$$

which is the same as the warp factor of the RS II braneworld and implies that the metric (3) reduces an Anti-de Sitter (AdS) one, which is essential for the localization of gravitation. The following expressions can also be taken

$$A'(y) = -\delta k \tanh(ky), \quad (12)$$

$$A'^2(y) = \delta^2 k^2 (1 - \text{sech}^2(ky)), \quad (13)$$

$$A''(y) = -\delta k^2 \text{sech}^2(ky), \quad (14)$$

and the scalar curvature takes a simple form:

$$R(y) = 20\delta^2 k^2 (\varepsilon \text{sech}^2(ky) - 1), \quad (15)$$

where  $\varepsilon = \frac{5\delta+2}{5\delta}$ . Furthermore, we can obtain the following expressions

$$f(R) = \sum_{i=1}^n a_i (20\delta^2 k^2)^i (\varepsilon \text{sech}^2(ky) - 1)^i + \Lambda, \quad (16)$$

$$f_R = \sum_{i=1}^n i a_i (20\delta^2 k^2)^{i-1} (\varepsilon \text{sech}^2(ky) - 1)^{i-1}, \quad (17)$$

$$\begin{aligned} f_R' &= \sum_{i=1}^n 2i(1-i)a_i k \varepsilon (20\delta^2 k^2)^{i-1} \text{sech}^2(ky) \tanh(ky) \\ &\quad \times (\varepsilon \text{sech}^2(ky) - 1)^{i-2}, \end{aligned} \quad (18)$$

$$\begin{aligned} f_R' A' &= \sum_{i=1}^n 2i(i-1)a_i \delta k^{2i} \varepsilon (20\delta^2)^{i-1} \text{sech}^2(ky) \\ &\quad \times (1 - \text{sech}^2(ky)) (\varepsilon \text{sech}^2(ky) - 1)^{i-2}, \end{aligned} \quad (19)$$

$$\begin{aligned} f_R'' &= \sum_{i=1}^n 2i(i-1)a_i \varepsilon k^{2i} (20\delta^2)^{i-1} (\varepsilon \text{sech}^2(ky) - 1)^{i-3} \\ &\quad \times \text{sech}^2(ky) [2(i-2)\varepsilon \text{sech}^2(ky)(1 - \text{sech}^2(ky)) \\ &\quad + (2 - 3\text{sech}^2(ky))(\varepsilon \text{sech}^2(ky) - 1)] \end{aligned} \quad (20)$$

By taking above expressions back into Eq. (7a), it is clear that all the terms of the Einstein equation are the function of  $\text{sech}^2(ky)$ , and the order of the highest is  $2n$ . The Einstein equation (7a) can be expressed as the following algebraic equation:

$$\sum_{i=0}^n B_i \text{sech}^{2i}(ky) = 0, \quad (21)$$

where the coefficient  $B_i$  can be written as follows

$$\begin{aligned}
B_i = & \sum_{j=i}^n (-1)^{j-i} 2a_j (20\delta^2)^{j-1} k^{2j} \varepsilon^{i-1} \\
& \times \left[ 10\delta^2 C_j^i \varepsilon - j \delta \left( 4\delta C_{j-1}^i \varepsilon + (1+4\delta) C_{j-1}^{i-1} \right) \right. \\
& + 2j(j-1) \left( (2+3\delta) C_{j-2}^{i-1} \varepsilon + (3+3\delta) C_{j-2}^{i-2} \right) \\
& \left. + 4j(j-1)(j-2) (C_{j-3}^{i-2} \varepsilon + C_{j-3}^{i-3}) \right], \tag{22}
\end{aligned}$$

with  $a_0 \equiv \Lambda$  and  $C_j^i$  the binomial coefficient. Here we define  $C_j^i \equiv 0$  when  $i > j$ . The  $B_i$  can also be expressed as a column matrix:

$$\begin{pmatrix} B_0 \\ B_1 \\ \vdots \\ B_{n-1} \\ B_n \end{pmatrix} = \begin{pmatrix} b_{0,0} & b_{0,1} & \cdots & b_{0,n-1} & b_{0,n} \\ b_{1,0} & b_{1,1} & \cdots & b_{1,n-1} & b_{1,n} \\ \vdots & \vdots & b_{i,j} & \vdots & \vdots \\ b_{n-1,0} & b_{n-1,1} & \cdots & b_{n-1,n-1} & b_{n-1,n} \\ b_{n,0} & b_{n,1} & \cdots & b_{n,n-1} & b_{n,n} \end{pmatrix}_{((n+1) \times (n+1))} \times \begin{pmatrix} a_0 \\ a_1 \\ \vdots \\ a_{n-1} \\ a_n \end{pmatrix}, \tag{23}$$

where the element of the matrix  $b_{i,j}$  is

$$\begin{aligned}
b_{i,j} = & (-1)^{j-i} 2(20\delta^2)^{j-1} k^{2j} \varepsilon^{i-1} \\
& \times \left[ 10\delta^2 C_j^i \varepsilon - j \delta \left( 4\delta C_{j-1}^i \varepsilon + (1+4\delta) C_{j-1}^{i-1} \right) \right. \\
& + 2j(j-1) \left( (2+3\delta) C_{j-2}^{i-1} \varepsilon + (3+3\delta) C_{j-2}^{i-2} \right) \\
& \left. + 4j(j-1)(j-2) (C_{j-3}^{i-2} \varepsilon + C_{j-3}^{i-3}) \right]. \tag{24}
\end{aligned}$$

By setting all the coefficients to zero  $B_i = 0$ , the solution to the Eq (21) can be calculated. Analyzing the expression (24), when  $i > j$ , it is easy to find that  $b_{i,j} = 0$ , since all the binomial coefficients are zero. Specially, except for  $b_{n,n}$ , the elements of the last row of the matrix are zero, so the coefficient  $B_n$  satisfies the following formulation

$$\begin{aligned}
B_n = & 2k^2 a_n (20\delta^2 k^2)^{n-1} \varepsilon^{n-1} \left[ (10-4n)\delta^2 \right. \\
& \left. + (6n^2 - 7n + 4)\delta + (4n^3 - 6n^2 + 2n) \right] = 0. \tag{25}
\end{aligned}$$

For non-zero solution of  $a_n$ , the relationship between  $\delta$  and  $n$  can be calculated as follows

$$\delta = \frac{4n^2 - 6n + 2}{2n - 5}. \tag{26}$$

For each certain value of  $n$ , the parameter  $\delta$  can be obtained as it is depicted in Fig. 1. In order to ensure that the behavior of warp factor at infinity tends to zero,  $\delta$  must be positive




FIG. 1. The relation between the parameter  $\delta$  and  $n$ .

definite, so the minimum value of the positive integer  $n$  is  $n_{\min} = 3$ . It is easy to see that when  $n = 4$ , the minimum  $\delta$  can be achieved  $\delta_{\min} = 14$ , and when  $n$  tends to infinity,  $\delta$  is proportional to  $n$  and tends to divergence.

Furthermore, Eq. (23) can be reexpressed as follows

$$\begin{pmatrix} 1 & b_{0,1} & \cdots & b_{0,n-1} & b_{0,n} \\ 0 & b_{1,1} & \cdots & b_{1,n-1} & b_{1,n} \\ \vdots & \vdots & b_{i,j} & \vdots & \vdots \\ 0 & 0 & \cdots & b_{n-1,n-1} & b_{n-1,n} \\ 0 & 0 & \cdots & 0 & 0 \end{pmatrix}_{(n+1) \times (n+1)} \times \begin{pmatrix} \Lambda \\ a_1 \\ \vdots \\ a_{n-1} \\ a_n \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}. \quad (27)$$

By considering the Eqs. (24) and (26), all the elements of the matrix  $b_{i,j}$  can be calculated for the given value of  $n$ , so the matrix equation (27) contains  $n$  linear equations and  $n + 1$  unknown parameters  $\Lambda, a_1, a_2, a_3, \cdots, a_n$ , and the relationships of these parameters could be concluded by solving Eq. (27).

When  $n = 1$  and  $f(R) = a_1 R + \Lambda$ , the case of general relativity is recovered. Because  $\delta = 0$  and  $A(y) = 0$ , there is no pure geometric thick  $\mathcal{M}_4$  brane solution. However, by using a different metric ansatz, a de Sitter  $dS_4$  brane solution can be obtained, which has been studied in Refs. [8, 151], or by introducing the tension of the brane, the RS braneworld model [3, 22] can also be concluded.

When  $n = 2$  and  $f(R) = a_1 R + a_2 R^2 + \Lambda$ , the parameter  $\delta = -6$  and  $A(y) = 6 \ln(\cosh(ky))$ , thus the warp factor  $e^{2A(y)} = \cosh^{12}(ky)$  is divergent at infinity of the extra dimensional coordinate. By introducing a background scalar, the divergence of warp factor is removed and the thick branes can be constructed in Ref. [50, 57, 147].

When  $n = 3$ , the parameter  $\delta = 20$  and warp factor  $A(y) = -20 \ln(\cosh(ky))$ , and a

$n$	$n = 3$	$n = 4$	$n = 10$
$\delta$	$\delta = 20$	$\delta = 14$	$\delta = \frac{114}{5}$
$a_1$	$-\frac{3147}{94400k^2}\Lambda \approx -0.0333\frac{\Lambda}{k^2}$	$-0.0478\frac{\Lambda}{k^2}$	$-0.0292\frac{\Lambda}{k^2}$
$a_2$	$\frac{249}{6041600k^4}\Lambda \approx 4.121 \times 10^{-5}\frac{\Lambda}{k^4}$	$2.488 \times 10^{-5}\frac{\Lambda}{k^4}$	$3.166 \times 10^{-5}\frac{\Lambda}{k^4}$
$a_3$	$-\frac{13}{1933312000k^6}\Lambda \approx -6.724 \times 10^{-9}\frac{\Lambda}{k^6}$	$-2.166 \times 10^{-8}\frac{\Lambda}{k^6}$	$-4.403 \times 10^{-9}\frac{\Lambda}{k^6}$
$a_4$		$7.927 \times 10^{-13}\frac{\Lambda}{k^8}$	$-2.834 \times 10^{-14}\frac{\Lambda}{k^8}$
$a_5$			$-1.002 \times 10^{-18}\frac{\Lambda}{k^{10}}$
$a_6$			$-3.689 \times 10^{-23}\frac{\Lambda}{k^{12}}$
$a_7$			$-1.130 \times 10^{-27}\frac{\Lambda}{k^{14}}$
$a_8$			$-2.550 \times 10^{-32}\frac{\Lambda}{k^{16}}$
$a_9$			$-3.684 \times 10^{-37}\frac{\Lambda}{k^{18}}$
$a_{10}$			$-2.530 \times 10^{-42}\frac{\Lambda}{k^{20}}$

TABLE I. The solutions of the pure geometric  $f(R)$  thick brane for  $n = 3$ ,  $n = 4$ , and  $n = 10$ .

pure geometric thick  $\mathcal{M}_4$  brane can be received. When  $n = 4$ , the parameter  $\delta$  can be achieved the minimum value  $\delta = \delta_{\min} = 14$ , and a pure geometric thick  $\mathcal{M}_4$  brane can also be constructed. When  $n = 5, 6, \dots$ , the solutions of a thick  $\mathcal{M}_4$  brane can be concluded, however, as the value of the parameter  $n$  increases, the calculation also increases. In this paper, the solutions will be discussed when  $n = 3$  (for the minimum  $n$ ),  $n = 4$  (for the minimum  $\delta$ ), and  $n = 10$  (for a large value  $n$ ), and these cases are very typical. The parameters of the solutions are shown in Table I. Here, the coefficient  $a_1$  is set to positive, so the 5D cosmological constant must be negative  $\Lambda < 0$ , and the 5D spacetime is AdS one. Moreover, the coefficients  $a_i$  for the solutions are shown in Fig. 2, and it can be found that the coefficients  $a_i$  significantly decreases with increasing the order  $i$ . For the case of  $n = 3$ , if we set  $\Lambda = -\frac{377600k^2}{7803}$ , the result is the same as example 2 in Ref.[77].

The pure geometric thick branes can be constructed, and the behaviors of the scalar curvature (15) at zero and infinity can be analyzed as follows:

$$\begin{cases} R(y=0) = 8\delta k^2, \\ R(y \rightarrow \infty) \rightarrow -20\delta^2 k^2. \end{cases} \quad (28)$$

The shapes of  $A(y)$  and  $R(y)$  are shown in Figs.3(a) and 3(b). We can find that both of  $A(y)$  and  $R(y)$  are smooth of the extra dimensional and  $R(y)$  tends to a constant  $-20\delta^2 k^2$






FIG. 2. The coefficients  $|\frac{a_i}{\Lambda}|$  in logarithmic coordinate.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .




FIG. 3. The shapes of  $A(y)$  in (a) and  $R(y)$  in (b) for different thick brane solutions.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .

at infinity.

### III. TENSOR PERTURBATION AND THE LOCALIZATION OF GRAVITY ON THE THICK $f(R)$ BRANES

The stability of the tensor perturbations of the gravity will be studied in this section. The small perturbations  $h_{\mu\nu}$  are introduced into the metric [144]:

$$ds^2 = e^{2A(y)}(\eta_{\mu\nu} + h_{\mu\nu})dx^\mu dx^\nu + dy^2 \quad (29)$$

where  $h_{\mu\nu} = h_{\mu\nu}(x^\rho, y)$  depend on all the spacetime coordinates. The following relations can be obtained immediately:

$$\begin{aligned}\delta R_{\mu\nu} = & -\frac{1}{2}(\hat{\square}h_{\mu\nu} + \partial_\mu\partial_\nu h - \partial_\nu\partial_\sigma h_\mu^\sigma - \partial_\mu\partial_\sigma h_\nu^\sigma) \\ & - 2e^{2A}A'h'_{\mu\nu} - 3h_{\mu\nu}e^{2A}A'^2 - \frac{1}{2}e^{2A}h''_{\mu\nu} \\ & - \frac{1}{2}e^{2A}A'\eta_{\mu\nu}h',\end{aligned}\tag{30}$$

$$\delta R_{\mu 5} = \frac{1}{2}\partial_y(\partial_\lambda h_\mu^\lambda - \partial_\mu h),\tag{31}$$

$$\delta R_{55} = -\frac{1}{2}(2A'h' + h''),\tag{32}$$

$$\begin{aligned}\delta R = & \delta(g^{\mu\nu}R_{\mu\nu}) \\ = & -e^{-2A}\hat{\square}h + e^{-2A}\partial_\mu\partial_\nu h^{\mu\nu} - 5A'h' - h'',\end{aligned}\tag{33}$$

Where  $\hat{\square} = \eta^{\mu\nu}\partial_\mu\partial_\nu$  is the 4D d'Alembert operator, and  $h = \eta^{\mu\nu}h_{\mu\nu}$ . The tensor perturbations satisfy the transverse-traceless (TT) condition  $\partial_\mu h_\nu^\mu = h = 0$ .

The perturbed Einstein equations can be expressed:

$$\begin{aligned}\delta R_{MN}f_R + R_{MN}f_{RR}\delta R - \frac{1}{2}\delta g_{MN}f(R) \\ - \frac{1}{2}g_{MN}f_R\delta R + \delta(g_{MN}\square f_R) - \delta(\nabla_M\nabla_N f_R) = 0,\end{aligned}\tag{34}$$

where

$$f_{RR} \equiv \frac{df_R}{dR} = \frac{d^2 f(R)}{dR^2},$$

$$\delta(\nabla_M\nabla_N f_R) = (\partial_M\partial_N - \Gamma_{MN}^P\partial_P)(f_{RR}\delta R) - \delta\Gamma_{MN}^P\partial_P f_R,\tag{35}$$

$$\delta(g_{MN}\square f_R) = \delta g_{MN}\square f_R + g_{MN}\delta g^{AB}(\nabla_A\nabla_B f_R) + g_{MN}g^{AB}\delta(\nabla_A\nabla_B f_R).\tag{36}$$

By substituting the Eqs. (30),(31),(32), and (33) into the Eq.(34), the perturbed Einstein equations can be simplified:

$$\left(e^{-2A}\hat{\square}h_{\mu\nu} + 4A'h'_{\mu\nu} + h''_{\mu\nu}\right)f_R + h'_{\mu\nu}f'_R = 0,\tag{37}$$

or, we can have a more simpler form

$$\square h_{\mu\nu} - \frac{f'_R}{f_R}\partial_y h_{\mu\nu} = 0.\tag{38}$$

By following the method given in Refs [3, 22], a coordinate transformation can be introduced

$$dz = e^{-A(y)}dy,\tag{39}$$

and the conformal flat metric can be taken:

$$ds^2 = e^{2A(z)}(\eta_{\mu\nu}dx^\mu dx^\nu + dz^2). \quad (40)$$

By using the coordinate transformation Eq.(39), the perturbed Einstein equations Eq.(38) are rewritten as:

$$\left[ \partial_z^2 + (3\partial_z A + \frac{\partial_z f_R}{f_R})\partial_z + \hat{\square} \right] h_{\mu\nu} = 0. \quad (41)$$

By doing the KK decomposition

$$h_{\mu\nu}(x^\rho, z) = e^{-\frac{3}{2}A} f_R^{-\frac{1}{2}} \varepsilon_{\mu\nu}(x^\rho) \psi(z), \quad (42)$$

a Schrödinger-like equation for the KK modes  $\psi(z)$  can be reexpressed

$$[-\partial_z^2 + W(z)]\psi(z) = m^2\psi(z) \quad (43)$$

where the effective potential  $W(z)$  is

$$\begin{aligned} W(z) = & \frac{9}{4}(\partial_z A)^2 + \frac{3}{2}\partial_z^2 A \\ & + \frac{3}{2} \frac{\partial_z A \partial_z f_R}{f_R} - \frac{1}{4} \frac{(\partial_z f_R)^2}{f_R^2} + \frac{\partial_z^2 f_R}{f_R}. \end{aligned} \quad (44)$$

Above equation (44) can be factorized as

$$QQ^\dagger \psi(z) = m^2 \psi(z) \quad (45)$$

with

$$Q = +\partial_z + \left( \frac{3}{2}\partial_z A + \frac{1}{2} \frac{\partial_z f_R}{f_R} \right), \quad (46)$$

$$Q^\dagger = -\partial_z + \left( \frac{3}{2}\partial_z A + \frac{1}{2} \frac{\partial_z f_R}{f_R} \right), \quad (47)$$

which ensures that there is no gravitational mode with  $m^2 < 0$  and the braneworld solutions are stable against the tensor perturbations. However, the scalar perturbations are still not clear because of the higher derivatives in the perturbation equations. It is well-known that a pure  $f(R)$  gravity is conformally equivalent to a theory with a minimally coupled scalar in Einstein's gravity [133]. By following the method given by in Ref. [77], our solutions are stable against the small linear metric perturbations, including tensor, vector, and scalar perturbations, in the Einstein frame.

For Eq.(43), the zero mode with  $m = 0$  is

$$\psi_{(0)}(z) = N_G e^{\frac{3}{2}A(z)} f_R^{\frac{1}{2}}(z), \quad (48)$$

where  $N_G$  is the normalization constant, and the  $\psi_{(0)}(z)$  satisfies the normalizable condition

$$\int_{-\infty}^{+\infty} |\psi_{(0)}(z)|^2 dz = 1. \quad (49)$$

The coordinate transformation Eq.(39) can be rewritten as follows:

$$z = \int e^{-A(y)} dy = \int \cosh^\delta(ky) dy. \quad (50)$$

However, the above equation can not be analytically calculated for an arbitrary  $\delta$ , and it is also difficult to directly get the analytical expressions for the effective potential  $W(z)$  and the zero mode  $\psi_{(0)}(z)$  under the conformal coordinate  $z$ . So the solutions under the coordinate  $y$  can be calculated firstly, and the numerical solutions under the conformal coordinate  $z$  can also be obtained.

By using the relation

$$e^{-A(y)} \xrightarrow{y \rightarrow +\infty} 2^{-\delta} e^{\delta ky}, \quad (51)$$

the asymptotic behavior of the relation between  $z$  and  $y$  can be analyzed

$$z \xrightarrow{y \rightarrow +\infty} \frac{1}{2^\delta \delta k} e^{\delta ky}. \quad (52)$$

We can solve the asymptotic behavior of the relation between  $y$  and  $z$  for  $z$  tends to infinity

$$y \xrightarrow{z \rightarrow +\infty} \frac{1}{\delta k} \ln(2^\delta \delta k z). \quad (53)$$

And the relations between  $z$  and  $y$  can be solved by numerical method, and shown in Fig 4, when  $\delta$  are the determined values given by  $n = 3, 4, 10$  respectively.

Then the behavior of  $A(z)$  and the scalar curvature  $R(z)$  at  $z = 0$  and  $z \rightarrow \pm\infty$  can be analyzed respectively,

$$\begin{cases} A(z = 0) = 0 \\ A(z \rightarrow \pm\infty) \rightarrow -\ln(\delta k |z|) \end{cases}, \quad (54)$$

$$\begin{cases} R(z = 0) = 8\delta k^2 \\ R(z \rightarrow \pm\infty) \rightarrow -20\delta^2 k^2 + C_1 |z|^{-\frac{2}{\delta}} \end{cases} \quad (55)$$




FIG. 4. The relation between  $z$  and  $y$ .  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .




FIG. 5. The shapes of  $A(z)$  for different solutions.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .

with  $C_1 = (8\delta k^2 + 20\delta^2 k^2)(\delta k)^{-\frac{2}{\delta}}$ , and we can have conclusions that the warp factor  $e^{2A}$  tends to 0 and  $R$  tends to a constant  $-20\delta^2 k^2$  for  $z \rightarrow \pm\infty$ . The numerical solutions of  $A(z)$  and  $R(z)$  are also solved and shown in Figs.5 and 6.

Since there is no analytical expression for the effective potential  $W(z)$  with respect to the coordinate  $z$ , the expression for it in the coordinate  $y$  can be expressed as follows:

$$\begin{aligned}
 W(z(y)) = & \frac{15}{4}e^{2A(y)}A'^2(y) + \frac{3}{2}e^{2A(y)}A''(y) \\
 & + \frac{1}{2}e^{2A(y)}\frac{4A'(y)f'_R + f''_R}{f_R} \\
 & - \frac{1}{4}e^{2A(y)}\left(\frac{f'_R}{f_R}\right)^2.
 \end{aligned} \tag{56}$$

And the shapes of  $W(z(y))$  are shown in Fig.7(a). The numerically solution of  $W(z)$  are shown in Fig.7(b), which are similar to Fig.7(a). The asymptotic behavior of these poten-




FIG. 6. The shapes of the scalar curvature  $R(z)$  for different solutions.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The range of the coordinate is around zero for (a), and the range of the coordinate is far away from zero for (b). The parameter is set to  $k = 1$ .




FIG. 7. The shapes of the potential  $W(z(y))$  in 7(a) and  $W(z)$  in 7(b) for different solutions.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .

tials tends zero at infinity. The height of the potential is related to the parameter  $\delta$  decided by  $n$ . When  $n = 10$  and  $\delta = \frac{114}{5}$ , the height of the potential is the tallest of all, and when  $n = 4$  and  $\delta = 14$ , it is the lowest.

This potential supports a normalizable zero mode, and the normalization constant  $N_G$

satisfies the following condition:

$$\begin{aligned}
1 &= \int_{-\infty}^{+\infty} |\psi_{(0)}(z)|^2 dz = N_G^2 \int_{-\infty}^{+\infty} e^{3A(z)} f_R(z) dz \\
&= \frac{N_G^2}{k} \int_{-\infty}^{+\infty} e^{2A(\omega)} f_R(\omega) d\omega,
\end{aligned} \tag{57}$$

where  $\omega = ky$  and  $dz = e^{-A} \frac{d\omega}{k}$ . For our solutions, the above integration can be done analytically. For  $n = 3$ , the integration gives

$$-\frac{N_G^2}{k} \frac{3723736645632}{245251477631875k^2} \Lambda = 1, \tag{58}$$

and  $N_G \approx 8.115k^{\frac{3}{2}}/\sqrt{-\Lambda}$ . For  $n = 4$  and  $n = 10$ , similarly, we can get  $N_G \approx 6.181k^{\frac{3}{2}}/\sqrt{-\Lambda}$ , and  $N_G \approx 8.965k^{\frac{3}{2}}/\sqrt{-\Lambda}$  respectively. Thus, the gravitational zero mode is normalizable and can be localized on the brane, which results in the familiar Newton's law on the brane. The shape of zero mode  $|\psi_{(0)}(z)|^2$  is shown in Fig. 8. There is a platform for the gravitational zero mode around zero, so the zero mode is localized near zero.

Starting from  $m^2 > 0$ , the continuum of the massive KK modes might lead to a correction to the Newtonian potential on the brane. As have been addressed in ref.[27], if the potential  $\lim_{z \rightarrow \infty} W(z) = \frac{\beta(\beta+1)}{z^2}$ , the massive modes will contribute a correction  $\Delta U \propto 1/r^{2\beta}$  to the Newton's law at large distance (see also [46]). For our solutions, the asymptotic behavior of  $z^2 W(z)$  with  $z \rightarrow \infty$  can be analytically obtained. Instead of writing down the explicit expressions, it is shown in Fig. 9 that for our solutions  $\lim_{z \rightarrow \infty} z^2 W(z) = \frac{15}{4}$ , namely  $\beta = 3/2$ . Thus, the corrections to the Newtonian potential  $\Delta U \propto 1/r^3$  are suppressed at large  $r$  for the braneworld.

Note that for KK modes with  $0 < m^2 < W_{\max}(z)$ , there might exist some resonant KK modes which tend to plane waves when  $z \rightarrow \infty$  and cannot be normalized for this type potential  $W(z)$  [52]. Here  $W_{\max}(z)$  is the maximum of the potential. However, according to the numerical calculation, there exists no resonant state.

#### IV. LOCALIZATION OF SPIN-0 SCALAR FIELDS ON THE THICK $f(R)$ BRANES

In this section, we will investigate the localization of scalar field on the thick  $f(R)$  branes. We start with the action of a 5D massless scalar field:

$$S_0 = \int d^5x \sqrt{-g} \left[ -\frac{1}{2} g^{MN} \partial_M \Phi \partial_N \Phi \right], \tag{59}$$




FIG. 8. The shapes of the zero mode for gravity  $|\psi_{(0)}(z)|^2$  for different solutions.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .




FIG. 9. The shapes of  $z^2W(z)$  for different solutions.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .

and the equation of motion from the above action(59) is read as:

$$\frac{1}{\sqrt{-g}}\partial_M(\sqrt{-g}g^{MN}\partial_N\Phi) = 0. \quad (60)$$

Using the conformal metric(40), and introducing the KK decomposition

$$\Phi(x^\mu, z) = \sum_{\theta} \phi_{\theta}(x^\mu) \chi_{\theta}(z) e^{-\frac{3}{2}A}, \quad (61)$$

we can get the 4D Klein-Gordon equation

$$\frac{1}{\sqrt{-\hat{g}}}\partial_{\mu}(\sqrt{-\hat{g}}\eta^{\mu\nu}\partial_{\nu})\phi_{\theta} = m_{\theta}^2\phi_{\theta}, \quad (62)$$



where  $\hat{g} = \det(\eta_{\mu\nu})$  is the determinant of the 4D effective metric, and the Schrödinger-like equation of the scalar KK modes respect to the extra dimensional coordinate

$$[-\partial_z^2 + V_0(z)]\chi_\theta(z) = m_\theta^2 \chi_\theta(z), \quad (63)$$

where  $m_\theta$  is the mass of the  $\theta$ -th KK mode of the scalar field, and the potential is given by:

$$V_0(z) = \frac{3}{2}\partial_z^2 A + \frac{9}{4}(\partial_z A)^2. \quad (64)$$

The scalar KK modes should satisfy the following orthogonal normalization conditions:

$$\int_{-\infty}^{\infty} dz \chi_\theta(z) \chi_\sigma(z) = \delta_{\theta\sigma}, \quad (65)$$

and the action (59) of the 5D free massless scalar field turns to the 4D effective action of a massless ( $m_0 = 0$ ) and a series of massive ( $m_\theta > 0$ ) scalar fields:

$$S_0 = \sum_{\theta} \int d^4x \sqrt{-g} \left[ -\frac{1}{2} g^{\mu\nu} \partial_\mu \phi_\theta \partial_\nu \phi_\theta - \frac{1}{2} m_\theta^2 \phi_\theta^2 \right] \quad (66)$$

By setting  $m = 0$ , the scalar zero mode can be solved from Eq.(63)

$$\chi_0(z) = N_S e^{\frac{3}{2}A(z)}, \quad (67)$$

where  $N_S$  is the normalization constant given by

$$\begin{aligned} 1 &= \int_{-\infty}^{+\infty} |\chi_0(z)|^2 dz = N_S^2 \int_{-\infty}^{+\infty} e^{3A(z)} dz \\ &= \frac{N_S^2}{k} \int_{-\infty}^{+\infty} e^{2A(\omega)}(\omega) d\omega. \end{aligned} \quad (68)$$

When  $n = 3$ ,  $n = 4$ , and  $n = 10$ ,  $N_S \approx 1.583\sqrt{k}$ ,  $N_S \approx 1.446\sqrt{k}$ , and  $N_S \approx 1.637\sqrt{k}$  respectively. The zero mode of scalar can be localized on the brane.

Considering Eq.(54), the asymptotic behavior of  $V_0(z)$  (64) and  $\chi_0(z)$  (67) can be analyzed as follows:

$$\begin{cases} V_0(z=0) = -\frac{3}{2}\delta k^2 \\ V_0(z \rightarrow \pm\infty) \rightarrow \frac{15}{4}|z|^{-2} \rightarrow 0, \end{cases} \quad (69)$$

$$\begin{cases} \chi_0(z=0) = N_S \\ \chi_0(z \rightarrow \pm\infty) \rightarrow N_S(\delta k)^{-\frac{3}{2}}|z|^{-\frac{3}{2}}. \end{cases} \quad (70)$$

The numerical results of  $V_0(z)$  and  $|\chi_0(z)|^2$  are also shown in Fig. 10 and Fig. 11 respectively.

We can have a conclusion that the zero mode of the scalar is localized at  $z = 0$ .




FIG. 10. The shapes of the effective potential  $V_0(z)$  for different solutions.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .




FIG. 11. The shapes of scalar field zero mode  $|\chi_0(z)|^2$  for different solutions.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .

## V. LOCALIZATION OF SPIN-1 VECTOR FIELDS ON THE THICK $f(R)$ BRANES

In this section, we will investigate the localization of vector fields on the thick  $f(R)$  brane. We start with a 5D gauge invariant action for a vector field coupled to the scalar curvature [85]

$$S_1 = -\frac{1}{4} \int d^5x \sqrt{-g} g_V(R) g^{MN} g^{RS} F_{MR} F_{NS}, \quad (71)$$

where  $F_{MN} = \partial_M A_N - \partial_N A_M$  is the field strength tensor, and  $g_V(R)$  is the function of scalar curvature  $R$ . Setting  $g_V(R) = 1$ , the action of the 5D free vector field can be recovered. The equation of motion is read as follows:

$$\frac{1}{\sqrt{-g}} \partial_M (\sqrt{-g} g_V(R) g^{MN} g^{RS} F_{NS}) = 0. \quad (72)$$

Using the conformal metric (40), introducing the following general KK decomposition:

$$A_\mu(x^\lambda, z) = \sum_\theta a_\mu^{(\theta)}(x^\lambda) \rho_\theta(z) e^{-\frac{A}{2}} (g_V(R))^{-\frac{1}{2}}, \quad (73)$$

and choosing the gauge freedom  $A_5 = 0$ , the vector KK modes  $\rho_\theta(z)$  should satisfy the following Schrödinger-like equation:

$$[-\partial_z^2 + V_1(z)]\rho_\theta(z) = m_\theta^2 \rho_\theta(z), \quad (74)$$

where the effective potential  $V_1(z)$  is

$$\begin{aligned} V_1(z) = & \frac{1}{2}\partial_z^2 A + \frac{1}{4}(\partial_z A)^2 \\ & + \frac{\partial_z A \partial_z g_V(R)}{2g_V(R)} + \frac{\partial_z^2 g_V(R)}{2g_V(R)} - \frac{(\partial_z g_V(R))^2}{4(g_V(R))^2}. \end{aligned} \quad (75)$$

The full 5D action (71) can be reduced to the 4D effective action for a massless and series of massive vectors

$$\begin{aligned} S_1 = & \sum_\theta \int d^4x \sqrt{-\hat{g}} \left( -\frac{1}{4} \eta^{\mu\alpha} \eta^{\nu\beta} f_{\mu\nu}^{(\theta)} f_{\alpha\beta}^{(\theta)} \right. \\ & \left. - \frac{1}{2} m_\theta^2 \eta^{\mu\nu} a_\mu^{(\theta)} a_\nu^{(\theta)} \right), \end{aligned} \quad (76)$$

when integrated over the extra dimension, with the requirement that Eq. (74) is satisfied and the following orthonormalization conditions are obeyed:

$$\int_{-\infty}^{\infty} \rho_\theta(z) \rho_\sigma(z) dz = \delta_{\theta\sigma}. \quad (77)$$

By setting  $m = 0$ , the solution of zero mode of the KK modes can be calculated:

$$\rho_0(z) = N_V e^{\frac{1}{2}A(z)} (g_V(R(z)))^{\frac{1}{2}}, \quad (78)$$

where  $N_V$  is the normalization constant given by

$$\begin{aligned} 1 = & \int_{-\infty}^{+\infty} |\rho_0(z)|^2 dz = N_V^2 \int_{-\infty}^{+\infty} e^{A(z)} g_V(R(z)) dz \\ = & \frac{N_V^2}{k} \int_{-\infty}^{+\infty} g_V(R(\omega)) d\omega. \end{aligned} \quad (79)$$

Next we will investigate the relation between the coupling with the gravity and the localization mechanism of the KK modes for the vector field.

#### A. Without the coupling: $g_V(R) = 1$

Firstly, we do not introduce the coupling between the vector field and the background spacetime, i.e.,  $g_V(R) = 1$ . From Eq.(75), the potential of the vector can be reduced to

$$V_1(z) = \frac{1}{2}A'' + \frac{1}{4}A'^2, \quad (80)$$




FIG. 12. The shapes of vector field effective potential  $V_1(z)$  without coupling term for different solutions.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .

and the asymptotic behavior of  $V_1(z)$  at  $z = 0$  and  $z \rightarrow \infty$  can be analyzed as follows

$$\begin{cases} V_1(z = 0) = \frac{1}{2}\delta k^2 \\ V_1(z \rightarrow \pm\infty) \rightarrow \frac{3}{4}z^{-2} \rightarrow 0. \end{cases} \quad (81)$$

The shapes of  $V_1(z)$  for the thick brane solutions are shown in Fig.12 by numerical method.

The zero mode (78) can also be reduced to

$$\rho_0(z) = N_V e^{\frac{1}{2}A}, \quad (82)$$

and the asymptotic behavior of  $\rho_0(z)$  has be calculated

$$\begin{cases} \rho_0(z = 0) = N_V \\ \rho_0(z \rightarrow \pm\infty) \rightarrow \frac{1}{\sqrt{\delta k}} z^{-\frac{1}{2}} \rightarrow 0. \end{cases} \quad (83)$$

From Eqs. (79) and (83), it is clear that the zero mode does not satisfy the normalization constant, i.e.,

$$\int_1^\infty |\rho_0(z)|^2 dz \approx \int_1^\infty \frac{1}{z} dz \rightarrow \infty. \quad (84)$$

Therefore the zero mode  $\rho_0(z)$  Eq.(82) can not be localized on the brane without the coupling between the vector and the gravity. The situation is the same as the RS model.

### B. With the coupling: $g_V(R) = (1 + \frac{R}{20\delta^2 k^2})^q$

In order to localization of the zero mode of the vector field, the coupling between the vector and the gravity must be introduced, and the following simple form is choose in this

paper:

$$g_V(R) = (1 + \frac{R}{20\delta^2 k^2})^q, \quad (85)$$

where  $q$  is coupling coefficient.

By using Eq.(75), the effective potential can be given by

$$V_1(z) = \frac{A''}{2} + \frac{A'^2}{4} + \frac{q(A'R' + R'')}{40k^2\delta^2 + 2R} + \frac{q(q-2)R'^2}{4(20k^2\delta^2 + R)^2}, \quad (86)$$

and the asymptotic behaviors of  $V_1(z)$  at zero and infinity are

$$\begin{cases} V_1(z=0) = -\frac{1}{2}(\delta + 2q)k^2 \\ V_1(z \rightarrow \infty) \rightarrow (\frac{3}{4} + \frac{2q}{\delta} + \frac{q^2}{\delta^2})z^{-2}. \end{cases} \quad (87)$$

The zero mass mode can also be given by

$$\rho_0 = N_V e^{\frac{1}{2}A(z)} (1 + \frac{R}{20k^2\delta^2})^{\frac{q}{2}}, \quad (88)$$

and the asymptotic behaviors of the zero mode  $\rho_0(z)$  at  $z=0$  and  $z \rightarrow \infty$  are

$$\begin{cases} \rho_0(z=0) = N_V (\frac{5\delta+2}{5\delta})^{\frac{q}{2}} \\ \rho_0(z \rightarrow \infty) \rightarrow N_V (\frac{5\delta+2}{5\delta})^{\frac{q}{2}} (\delta k)^{-\frac{\delta+2q}{2\delta}} z^{-\frac{\delta+2q}{2\delta}}. \end{cases} \quad (89)$$

For the purpose of trapping the zero mode of the vector field on the brane, the normalization condition (79) should be satisfied, i.e.,

$$\int_1^\infty \rho_0^2(z \rightarrow +\infty) dz \approx \int_1^\infty z^{-1-2\frac{q}{\delta}} dz < \infty, \quad (90)$$

which require that the coupling coefficient must be positive  $q > 0$ . In this paper, we will investigate the effects of the coupling coefficient  $q$  on the localization of the vector KK modes, so here set  $q = \frac{1}{2}\delta, 5\delta, 10\delta, 20\delta$ . The numerical solutions of the potential  $V_1(z)$  for the  $n=3$  thick brane are shown Fig. 13. It is clear that the potentials have a negative well around  $z=0$  and have two symmetrical barriers at both sides of the origin of the extra dimension, which increased with the coupling coefficient  $q$ . For the case of  $n=4$  and  $n=10$ , the potentials are similar to  $n=3$ . When  $q = \frac{\delta}{2}$ , the numerical results of the vector zero mode for  $n=3$ ,  $n=4$ , and  $n=10$  have been also shown in Fig. 14. Generally, this type potential implies that there may exist resonant states, which tend to plane waves when  $z \rightarrow \infty$  and




FIG. 13. The shapes of the effective potential for the vector fields  $V_1(z)$  with coupling term for different values of the coupling coefficient  $q$ . The parameters are set to  $n = 3$  and  $k = 1$ .




FIG. 14. The shapes of vector field zero modes  $|\rho_0(z)|^2$  with the coupling coefficient  $q = \frac{\delta}{2}$  for different solutions.  $n = 3$  for the dashed blue line,  $n = 4$  for the green line, and  $n = 10$  for the thin red line. The parameter is set to  $k = 1$ .

cannot be normalized. Following the method presented in Refs. [107, 111, 112, 152], the relative probability function of a resonance on the brane is defined as follows:

$$P_V(m^2) = \frac{\int_{-z_b}^{z_b} |\rho(z)|^2 dz}{\int_{-z_{\max}}^{z_{\max}} |\rho(z)|^2 dz}, \quad (91)$$

where  $2z_b$  is approximately the width of the thick brane, and  $z_{\max} = 10z_b$ . It is clear that the KK modes are approximately taken as plane waves and the corresponding probability  $P_V(m^2)$  tend to  $1/10$ , when  $m^2 \gg V_1^{\max}$  ( $V_1^{\max}$  is the maximum value of the corresponding potential  $V_1$ ). The lifetime  $\tau$  of a resonant state is  $\tau \sim \Gamma^{-1}$  with  $\Gamma = \delta m$  being the full width at half maximum of the peak. Equation (74) can be solved by the numerical method, and we will set the coupling coefficient  $q$  as different values  $q = \frac{1}{2}\delta, 5\delta, 10\delta, 20\delta$ , for each thick brane solution ( $n = 3, n = 4$ , and  $n = 10$ ) respectively.

When  $q = \frac{1}{2}\delta$ , there is no resonant state vector KK mode for each thick brane solution. However, when  $q = 5\delta$ , there is one resonant KK mode for each thick brane solution, and

the total number of resonant KK modes increases with the coupling coefficient  $q$ . The mass, width, and lifetime of the vector resonant KK modes with different values of  $q$  for each thick brane solution are listed in Tables II.

For the case of  $n = 3$  thick brane solution, the profiles of the relative probability  $P_V$  corresponding to different coupling coefficient  $q$  are shown in Fig. 15. In these figures, each peak corresponds to a resonant state, and the corresponding mass spectra with the effective potentials are also shown in Fig. 15. For the mass spectra of the vector KK modes, it can be seen that the ground state is zero mode (bound state), and all the massive KK modes are resonant KK modes. From Tables II and Fig. 15(f), it is clear that there are six resonant KK modes when  $q = 20\delta$ , and all the resonant KK modes are shown in Fig. 16. So we can summarize that the vector KK zero mode can be localized on the pure geometric thick  $f(R)$  brane, and the massive KK modes can be quasilocalized on the brane. For the other cases  $n = 4(\delta = 14)$  and  $n = 10(\delta = \frac{114}{5})$ , the situation is similar to the case of  $n = 3(\delta = 20)$ . Furthermore, the coupling function  $g_V(R)$  can also be set as other formulations and the property of the localization will be different. And the discussion in more detail has been investigated in Ref. [85].

## VI. LOCALIZATION OF SPIN-1/2 FERMION FIELDS ON THE THICK $f(R)$ BRANES

In this section, the localization of Spin-1/2 fermion field on the thick  $f(R)$  branes will be investigated, and the 5D Dirac action of fermion can be expressed as

$$S_{\frac{1}{2}} = \int d^5x \sqrt{-g} [\bar{\Psi} \Gamma^M (\partial_M + \omega_M) \Psi - \eta \bar{\Psi} \Gamma^M \partial_M g_f(R) \Psi], \quad (92)$$

where  $\omega_M$  is the spin connection defined as

$$\omega_M = \frac{1}{4} \omega_M^{\bar{M}\bar{N}} \Gamma_{\bar{M}} \Gamma_{\bar{N}} \quad (93)$$

with

$$\begin{aligned} \omega_M^{\bar{M}\bar{N}} = & \frac{1}{2} E^{N\bar{M}} (\partial_M E_N^{\bar{N}} - \partial_N E_M^{\bar{N}}) \\ & - \frac{1}{2} E^{N\bar{N}} (\partial_M E_N^{\bar{M}} - \partial_N E_M^{\bar{M}}) \\ & - \frac{1}{2} E^{P\bar{M}} E^{Q\bar{N}} E_M^{\bar{R}} (\partial_P E_{Q\bar{R}} - \partial_Q E_{P\bar{R}}). \end{aligned} \quad (94)$$

$n$	$\delta$	$q$	$V_1^{\max}$	$\theta$	$m^2$	$m$	$\Gamma$	$\tau$
3	20	$5\delta$	220.1707	1	173.1216	13.1575	0.2574	3.8840
				1	377.0040	19.4165	$6.092 \times 10^{-4}$	$1.641 \times 10^3$
				2	659.3324	25.6774	0.1214	8.2327
		$10\delta$	794.7859	3	829.4739	28.8005	1.4998	0.6667
				1	778.1259	27.8949	$1.431 \times 10^{-11}$	$6.984 \times 10^{10}$
				2	$1.470 \times 10^3$	38.3434	$6.533 \times 10^{-7}$	$1.530 \times 10^6$
				3	$2.070 \times 10^3$	45.5006	$3.880 \times 10^{-4}$	$2.577 \times 10^3$
		$20\delta$	$3.020 \times 10^3$	4	$2.567 \times 10^3$	50.6670	0.0273	36.5568
				5	$2.931 \times 10^3$	54.1458	0.4101	2.4379
				6	$3.176 \times 10^3$	56.3604	2.4695	0.4049
4	14	$5\delta$	152.5827	1	120.7882	10.9903	0.2455	4.0724
				1	263.5687	16.2348	$6.529 \times 10^{-4}$	$1.531 \times 10^3$
				2	459.9604	21.4466	0.1200	8.3271
		$10\delta$	550.7041	3	581.1552	24.1071	1.5808	0.6325
				1	544.3748	23.3318	$1.988 \times 10^{-11}$	$5.028 \times 10^{10}$
				2	$1.027 \times 10^3$	32.0602	$8.152 \times 10^{-7}$	$1.226 \times 10^6$
		$20\delta$	$2.092 \times 10^3$	3	$1.446 \times 10^3$	38.0281	$4.501 \times 10^{-4}$	$2.221 \times 10^3$
				4	$1.790 \times 10^3$	42.3200	0.0295	33.8816
				5	$2.042 \times 10^3$	45.1897	0.4127	2.4229
				6	$2.230 \times 10^3$	47.2321	3.6694	0.2725
10	$\frac{114}{5}$	$5\delta$	251.7237	1	197.5419	14.0549	0.2653	3.7686
				1	429.9404	20.7350	$6.001 \times 10^{-4}$	$1.666 \times 10^3$
				2	752.3325	27.4286	0.1222	8.1773
		$10\delta$	908.7305	3	945.6910	30.7520	1.5185	0.6585
				1	887.2097	29.7860	$1.301 \times 10^{-11}$	$7.686 \times 10^{10}$
				2	$1.676 \times 10^3$	40.9470	$6.170 \times 10^{-7}$	$1.620 \times 10^6$
		$20\delta$	$3.453 \times 10^3$	3	$2.361 \times 10^3$	48.5960	$3.745 \times 10^{-4}$	$2.670 \times 10^3$
				4	$2.929 \times 10^3$	54.1234	0.0268	37.2132
				5	$3.346 \times 10^3$	57.8474	0.4162	2.4025
				6	$3.615 \times 10^3$	60.1269	2.5566	0.3911

TABLE II. The mass, width, and lifetime of resonant KK modes of the vector with  $k = 1$ .






FIG. 15. The mass spectra, the effective vector potential  $V_1$ , and corresponding relative probability  $P_V$  with different coupling coefficient  $q = 5\delta, 10\delta, 20\delta$ .  $V_1$  for the black line, the even parity KK mode for the blue line, and the odd parity KK mode for the red line. The parameters are set to  $n = 3$  and  $k = 1$ .

The letters with barrier  $\bar{M}, \bar{N}$  are the five dimensional local Lorentz indices and the vielbein  $E_M^{\bar{M}}$  satisfies  $E_M^{\bar{M}} E_N^{\bar{N}} \eta^{\bar{M}\bar{N}} = g^{MN}$ . The relation between the gamma matrices  $\Gamma^M$  and  $\Gamma^{\bar{M}} = (\Gamma^{\bar{\mu}}, \Gamma^{\bar{5}}) = (\gamma^{\bar{\mu}}, \gamma^{\bar{5}})$  is given by  $\Gamma^M = E_{\bar{m}}^M \Gamma^{\bar{M}}$ .




FIG. 16. The shapes of vector resonance KK modes with the coupling coefficient  $q = 20\delta$ . The parameters are set to  $n = 3$  and  $k = 1$ .

Here the coupling between Dirac fermion and background spacetime [131]  $-\eta\bar{\Psi}\Gamma^M\partial_M g_f(R)\Psi$  with  $\eta$  coupling coefficient is introduced. Here we assume that the coupling coefficient  $\eta$  is positive.

Considering the conformally flat metric Eq.(40), the component of the spin connection is given by  $\omega_\mu = \frac{1}{2}(\partial_z A(z))\gamma_\mu\gamma_5$  and  $\omega_5 = 0$ . The equation of motion of the 5D Dirac fermion can be derived as

$$[\gamma^\mu\partial_\mu + \gamma^5(\partial_z + 2\partial_z A(z)) + \eta\partial_z g_f(R)]\Psi = 0. \quad (95)$$

The chiral decomposition for  $\Psi(x, z)$  can be introduced,

$$\Psi(x, z) = e^{-2A(z)} \sum_{\theta} [\psi_{L\theta}(x)L_{\theta}(z) + \psi_{R\theta}(x)R_{\theta}(z)], \quad (96)$$

where  $\Psi_{L\theta} = -\gamma^5\Psi_{L\theta}$  and  $\Psi_{R\theta} = \gamma^5\Psi_{R\theta}$  are the left- and right-chiral components of the 4D Dirac fermion field, respectively.

By taking the following orthonormalization conditions for the KK modes  $L_\theta$  and  $R_\theta$

$$\int_{-\infty}^{\infty} L_\theta L_\sigma dz = \delta_{\theta\sigma}, \quad (97)$$

$$\int_{-\infty}^{\infty} R_\theta R_\sigma dz = \delta_{\theta\sigma}, \quad (98)$$

$$\int_{-\infty}^{\infty} L_\theta R_\theta dz = 0, \quad (99)$$

one can take the effective action of the 4D massless and massive Dirac fermions from the 5D Dirac action (92)

$$S_{\frac{1}{2}} = \sum_{\theta} \int d^4x \sqrt{-\hat{g}} \bar{\psi}_\theta [\gamma^\mu (\partial_\mu + \hat{\omega}_\mu) - m_\theta] \psi_\theta. \quad (100)$$

By introducing the chiral decomposition Eq.(96), the Schrödinger-like equations of motion for the left- and right-chiral fermion KK modes  $L_\theta(z)$  and  $R_\theta(z)$  can be obtained:

$$[-\partial_z^2 + V_L(z)]L_\theta(z) = m_\theta^2 L_\theta(z), \quad (101a)$$

$$[-\partial_z^2 + V_R(z)]R_\theta(z) = m_\theta^2 R_\theta(z), \quad (101b)$$

where the effective potentials  $V_{L,R}(z)$  of the fermion KK modes are read as

$$V_L(z) = (\eta \partial_z g_f(R))^2 + \eta \partial_z^2 g_f(R), \quad (102a)$$

$$V_R(z) = (\eta \partial_z g_f(R))^2 - \eta \partial_z^2 g_f(R). \quad (102b)$$

By setting  $m_\theta = 0$  in Eq. (101), the solution of zero modes  $L_0$  and  $R_0$  can be obtained

$$L_0 \propto e^{\int_0^z dz \eta \partial_z g_f(R)} = e^{\eta g_f(R(z))} \quad (103a)$$

$$R_0 \propto e^{-\int_0^z dz \eta \partial_z g_f(R)} = e^{-\eta g_f(R(z))}. \quad (103b)$$

From the above relations (103), it is impossible to make both massless left- and right-chiral KK modes to be localized on the brane at the same time, since when one is normalizable, the other one is not.

From Eqs. (101) and (102), it is clear that, if we do not introduce the coupling in the action (92), i.e.,  $\eta = 0$ , the effective potentials for left- and right-chiral KK modes  $V_{L,R}(z) = 0$  and both left- and right-chiral fermions can not be localized on the thick brane, so the coupling term must be introduced. Moreover, since  $R$  is even function of  $z$ ,  $V_{L,R}(z)$  are naturally  $Z_2$  even with respect to  $z$ . Here we set a simple formulation:

$$g_f(R) = -\frac{20\delta^2 k^2 + 8\delta k^2}{R + 20\delta^2 k^2}, \quad (104)$$

By using Eqs.(54,55) and (104), the asymptotic behaviors of  $V_{L,R}(z)$  at  $z = 0$  and  $z \rightarrow \infty$  are as follows

$$\begin{cases} V_L(z = 0) = -2\eta k^2, \\ V_L(z \rightarrow \infty) \rightarrow \frac{2\eta}{\delta^2}(1 - \delta)z^{-2}, \end{cases} \quad (105)$$

$$\begin{cases} V_R(z = 0) = 2\eta k^2, \\ V_R(z \rightarrow \infty) \rightarrow \frac{2\eta}{\delta^2}(\delta - 1)z^{-2}. \end{cases} \quad (106)$$

Considering the coupling formulation (104) and Eq.(103), for the positive coupling coefficient  $\eta$  and positive parameter  $\delta$  and  $k$ , only left-chiral fermion zero mode may be localized on the brane. The asymptotic behavior of  $L_0(z)$  is also given by

$$\begin{cases} L_0(z = 0) = N_L e^{-\eta}, \\ L_0(z \rightarrow \infty) \rightarrow N_L e^{-\eta C_L |z|^s}, \end{cases} \quad (107)$$

where  $N_L$  is a constant,  $C_L = (\delta k)^{\frac{\delta}{2}} > 0$  and  $s = \frac{\delta}{2} \geq \frac{\delta_{\min}}{2} = 7$  are positive constants, respectively. It is easy to see that the normalization condition of the left-chiral fermion zero mode is satisfied, and it can be localized on the brane.

Next, the effect of the coupling between fermion and background spacetime for the property of the localization will be investigated, so the coupling coefficient  $\eta$  will be taken different values:  $\eta = 1, 50, 100$ . This type potential implies that resonant left- and right-chiral KK modes may exist. Mimic to the case of vector, the fermion relative probabilities for finding the left- and right-chiral fermion resonant states with mass  $m$  can be defined as:

$$P_{L,R}(m^2) = \frac{\int_{-z_b}^{z_b} |L, R(z)|^2 dz}{\int_{-z_{max}}^{z_{max}} |L, R(z)|^2 dz}. \quad (108)$$

For the case of  $n = 3$ , when the coupling coefficient  $\eta = 1$ , the effective potentials and the mass spectra for the left- and right-chiral fermion KK modes are shown in Figs. 17(a) and 17(b). Only the left-chiral fermion zero mode (bound state) can be localized on the brane. When the coupling coefficient  $\eta = 50$ , the effective potentials and the mass spectra for the left- and right-chiral fermion KK modes are shown in Figs. 17(c) and 17(d). For the left-chiral fermion KK modes, there is only one bound zero mode and one massive resonant KK mode, which is an even-parity. However, for the right-chiral fermion KK mode, there is only one massive resonant KK mode, which is an odd-parity. Both the mass of the left- and

right-chiral fermion KK modes are the same. In fact, these conclusions are originated from the coupled equations of the left- and right-chiral fermions. When  $\eta = 100$ , the effective potentials and the mass spectra for the left- and right-chiral fermion KK modes are shown in Figs. 17(e) and 17(f). The profiles of the relative probability  $P_{L,R}(m^2)$  are shown in Figs. 18(a) and 18(b), respectively. And the left- and right-chiral fermion KK modes are shown in Figs. 19 and 20. So we can summarize that the 4D massless left-chiral fermion can be localized on the brane, and the 4D massive Dirac fermions can also be quasilocalized on the brane, which consist of the pairs of coupled left- and right-chiral KK modes with different parities. The total number of resonant KK modes increases with the coupling coefficient  $\eta$ .

For the cases of  $n = 4$  and  $n = 10$ , the situation is similar to the case of  $n = 3$ , and the mass, width, and lifetime of the left- and right-chiral fermion KK resonant modes are listed in Table III.

Furthermore, the coupling function  $g_f(R)$  can also be set as other formulations and the property of the localization will be different. And the discussion in more detail has been investigated in Ref. [131].

## VII. CONCLUSION AND DISCUSSION

In this paper, we investigate a pure geometric thick  $\mathcal{M}_4$  brane, embedded in a  $\text{AdS}_5$  space-time, in general  $f(R)$  gravity theory. Here, the form of  $f(R)$  is set as  $f(R) = \sum_{i=1}^n a_i R^i + \Lambda$ , and any background scalar has not been introduced. For the pure geometric thick  $\mathcal{M}_4$  brane, the parameter  $n$  must be satisfied  $n \geq 3$ , and for a certainty value of  $n$ , a thick brane solution can be obtained. In this paper, the solutions of the  $\mathcal{M}_4$  brane for  $n = 3$ ,  $n = 4$ , and  $n = 10$ , have been studied, and found that the solutions are stable against tensor perturbations. Moreover, we investigate the localization of gravity and various bulk matter fields on the branes.

For the gravity, it is found that the gravitational zero mode can be localized near zero. All the massive modes are continuous-spectrum wave functions and can not be localized on the brane. For the scalar field, the zero mode is localized on the zero. For the vector field, if the coupling between vector and background spacetime is not introduced, the vector zero mode can not be localized on the brane, with the same case of RS brane. However, by introducing the coupling, the zero mode can be localized on the brane. For a large

$n$	$\eta$	Chrial	Height of $V_{L,R}$	$\theta$	$m^2$	$m$	$\Gamma$	$\tau$
3	50	Left	201.3102	1	157.0089	12.5303	0.1732	5.7749
		Right	202.9641	1	156.9938	12.5297	0.1801	5.5527
	100	Left	798.2219	1	360.5018	18.9869	$9.592 \times 10^{-5}$	$1.042 \times 10^4$
				2	635.9341	25.2177	0.0464	21.5500
				3	803.5692	28.3473	1.2032	0.8311
		Right	798.9972	1	360.5020	18.9868	$9.555 \times 10^{-5}$	$1.046 \times 10^4$
				2	635.9326	25.2177	0.0470	21.2549
				3	803.7101	28.3497	1.3213	0.7568
4	50	Left	296.0509	1	172.5166	13.1345	0.0057	173.6421
				2	281.7977	16.7868	0.4054	2.4664
		Right	296.7473	1	172.5166	13.1345	0.0058	172.3199
				2	281.6335	16.7819	0.4227	2.3652
	100	Left	$1.179 \times 10^3$	1	373.4193	19.3240	$1.844 \times 10^{-9}$	$5.421 \times 10^8$
				2	691.9976	26.3058	$4.681 \times 10^{-5}$	$2.136^4$
				3	950.5699	30.8313	0.0113	88.4840
				4	$1.134 \times 10^3$	33.6808	0.2806	3.5632
		Right	$1.179 \times 10^3$	1	373.4196	19.3240	$1.868 \times 10^{-9}$	$5.353 \times 10^8$
				2	691.9974	26.3058	$4.647 \times 10^{-5}$	$2.151 \times 10^4$
				3	950.5701	30.8313	0.0114	87.4471
				4	$1.134 \times 10^3$	33.6800	0.2929	3.4134
10	50	Left	175.4931	1	149.2316	12.2160	0.3738	2.6751
		Right	177.7526	1	149.0260	12.2076	0.4045	2.4718
	100	Left	694.0240	1	354.2681	18.8220	0.0010	929.1899
				2	606.2331	24.6218	0.1903	5.2540
		Right	695.0366	1	354.2683	18.8220	0.0011	909.9818
				2	606.3846	24.6248	0.1982	5.0438

TABLE III. The mass, width, and lifetime of resonant KK modes of the fermions with the parameter  $k = 1$ .




FIG. 17. The mass spectra and the effective potentials of the left- and right-chiral fermions with different coupling coefficient  $\eta = 1$ ,  $\eta = 50$ , and  $\eta = 100$ . The parameters are set to  $n = 3$  and  $k = 1$ .

coupling coefficient, there exist vector resonant states, and the number of the resonances increases with the coupling coefficient. For the spin-1/2 fermion field, in order to localize the fermion zero mode, the coupling between the fermion and the background spacetime must be introduced. With a small coupling coefficient, only the left-chiral fermion zero mode is localized on the brane. However, with a large coupling coefficient, the left-chiral fermion zero




FIG. 18. The shapes of the relative probability  $P_L(m^2)$  for the left-chiral fermion and  $P_R(m^2)$  for right-chiral fermion resonant. The parameters are set to  $\eta = 100$ ,  $n = 3$  and  $k = 1$ .




FIG. 19. The shapes of the left-chiral fermion KK modes, which is set to  $\eta = 100$ ,  $n = 3$  and  $k = 1$ .

mode is localized on the brane, and a finite number of resonant massive KK modes of the left- and right-chiral fermions are quasilocalized on the brane. And the number of resonances also increases with the coupling coefficient. Hence, the massless fermion is localized on the brane consists of just the left-chiral KK mode, and the massive fermions are quasilocalized




FIG. 20. The shapes of the right-chiral fermion KK modes, which is set to  $\eta = 100$ ,  $n = 3$  and  $k = 1$ .



on the brane consist of the left- and right-chiral fermion KK modes, represented the the 4D Dirac massive fermions. The lifetime of the fermion KK resonant modes decreases with their masses.

## ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (Grants No. 11305119, No. 11705070, and No. 11405121), the Natural Science Basic Research Plan in Shaanxi Province of China (Program No. 2020JM-198), the Fundamental Research Funds for the Central Universities (Grants No. JB170502), and the 111 Project (B17035).

- 
- [1] N. Arkani-Hamed, S. Dimopoulos, G. Dvali, *The hierarchy problem and new dimensions at a millimeter* Physics Letters B **429**(3-4), (1998) 263.
  - [2] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, G. Dvali, *New dimensions at a millimeter to a fermi and superstrings at a TeV* Physics Letters B **436**(3-4), (1998) 257.
  - [3] L. Randall, R. Sundrum, *Large Mass Hierarchy from a Small Extra Dimension* Physical Review Letters **83**, (1999) 3370.
  - [4] M. Gogberashvili, *Hierarchy problem in the shell-universe model* International Journal of Modern Physics D **11**(10), (2002) 1635.
  - [5] J.A. Cabrer, G.V. Gersdorff, M. Quirós, *Soft-wall stabilization* New Journal of Physics **12**(7), (2010) 075012.
  - [6] J.M. Hoff da Silva, M. Dias, *Five-dimensional  $f(R)$  braneworld models* Physical Review D **84**(6), (2011) 066011.
  - [7] K. Yang, Y. X. Liu, Y. Zhong, X. L. Du, S. W. Wei, *Gravity localization and mass hierarchy in scalar-tensor branes* Physical Review D **86**(12), (2012) 127502.
  - [8] H. Guo, A. Herrera-Aguilar, Y. X. Liu, D. Malagón-Morejón, R.R. Mora-Luna, *Localization of bulk matter fields, the hierarchy problem and corrections to Coulomb's law on a pure de Sitter thick braneworld* Physical Review D **87**(9), (2013) 095011.
  - [9] V.A. Rubakov, M.E. Shaposhnikov, *Do we live inside a domain wall?* Physics Letters B **125**(2-3), (1983) 136.

- [10] V.A. Rubakov, M.E. Shaposhnikov, *Extra space-time dimensions: Towards a solution to the cosmological constant problem* Physics Letters B **125**(2-3), (1983) 139.
- [11] S. Randjbar-Daemi, C. Wetterich, *Kaluza-Klein solutions with non-compact internal spaces* Physics Letters B **166**(1), (1986) 65.
- [12] E.J. Squires, *Dimensional reduction caused by a cosmological constant* Physics Letters B **167**(3), (1986) 286.
- [13] C. Csáki, J. Erlich, C. Grojean, T.J. Hollowood, *General properties of the self-tuning domain wall approach to the cosmological constant problem* Nuclear Physics B **584**(1-2), (2000) 359.
- [14] N. Arkani-Hamed, S. Dimopoulos, N. Kaloper, R. Sundrum, *A small cosmological constant from a large extra dimension* Physics Letters B **480**(1-2), (2000) 193.
- [15] S. Kachru, M. Schulz, E. Silverstein, *Self-tuning flat domain walls in 5D gravity and string theory* Physics Letters D **62**(4), (2000) 045021.
- [16] B. Bajc, G. Gabadadze, *Localization of matter and cosmological constant on a brane in anti de Sitter space* Physics Letters B **474**(3-4), (2000) 282.
- [17] J.E. Kim, B. Kyae, H.M. Lee, *Randall-Sundrum Model for Self-Tuning the Cosmological Constant* Physical Review Letters **86**(19), (2001) 4223-4226.
- [18] G.D. Starkman, D. Stojkovic, M. Trodden, *Homogeneity, Flatness, and "Large" Extra Dimensions* Physical Review Letters **87**(23), (2001) 231303.
- [19] A. Kehagias, K. Tamvakis, *A self-tuning solution of the cosmological constant problem* Modern Physics Letters A **17**(27), (2002) 1767.
- [20] A. Kehagias, *A conical tear drop as a vacuum-energy drain for the solution of the cosmological constant problem* Physics Letters B **600**(1-2), (2004) 133.
- [21] I.P. Neupane, *De Sitter brane-world, localization of gravity, and the cosmological constant* Physical Review D **83**(8), (2011) 086004.
- [22] L. Randall, R. Sundrum, *An Alternative to Compactification* Physical Review Letters **83**(23), (1999) 4690.
- [23] F. Bonjour, C. Charmousis, R. Gregory, *Thick domain wall universes* Classical and Quantum Gravity **16**(7), (1999) 2427.
- [24] W.D. Goldberger, M.B. Wise, *Modulus Stabilization with Bulk Fields* Physical Review Letters **83**(24), (1999) 4922.
- [25] M. Gremm, *Four-dimensional gravity on a thick domain wall* Physics Letters B **478**(4),

- (2000) 434.
- [26] O. DeWolfe, D.Z. Freedman, S.S. Gubser, A. Karch, *Modeling the fifth dimension with scalars and gravity* Physical Review D **62**(4), (2000) 046008.
  - [27] C. Csáki, J. Erlich, T.J. Hollowood, Y. Shirman, *Universal aspects of gravity localized on thick branes* Nuclear Physics B **581**(1-2), (2000) 309.
  - [28] S. Ichinose, *Some properties of domain wall solution in the Randall-Sundrum model* Classical and Quantum Gravity **18**(23), (2001) 5239.
  - [29] S. Ichinose, *A solution of the Randall-Sundrum model and the mass hierarchy problem* Classical and Quantum Gravity **18**(3), (2001) 421.
  - [30] A. Campos, *Critical Phenomena of Thick Branes in Warped Spacetimes* Physical Review Letters **88**(14), (2002) 141602.
  - [31] R. Gregory, A. Padilla, *Nested braneworlds and strong brane gravity* Physical Review D **65**(8), (2002) 084013.
  - [32] S. Kobayashi, K. Koyama, J. Soda, *Thick brane worlds and their stability* Physical Review D **65**(6), (2002) 064014.
  - [33] K.A. Bronnikov, B.E. Meierovich, *A general thick brane supported by a scalar field* Grav. Cosmol. **9**, (2003) 313.
  - [34] M. Eto, N. Sakai, *Solvable models of domain walls in  $\mathcal{N} = 1$  supergravity* Physical Review D **68**(12), (2003) 125001.
  - [35] A. Melfo, N. Pantoja, A. Skirzewski, *Thick domain wall spacetimes with and without reflection symmetry* Physical Review D **67**(10), (2003) 105003.
  - [36] D. Bazeia, C. Furtado, A.R. Gomes, *Brane structure from a scalar field in warped spacetime* Journal of Cosmology and Astroparticle Physics **2004**(02), (2004) 002.
  - [37] D. Bazeia, A. Gomes, A.R. Gomes, *Bloch brane* Journal of High Energy Physics **2004**(05), (2004) 012.
  - [38] D. Bazeia, F.A. Brito, L. Losano, *Scalar fields, bent branes, and RG flow* Journal of High Energy Physics **2006**(11), 064 (2006).
  - [39] V.I. Afonso, D. Bazeia, L. Losano, *First-order formalism for bent brane* Physics Letters B **634**, (2006) 526.
  - [40] V. Dzhunushaliev, *Thick brane solution in the presence of two interacting scalar fields* Grav. Cosmol. **13**, (2007) 302.

- [41] V. Afonso, D. Bazeia, R. Menezes, A. Petrov, *f(R)-branes* Physics Letters B **658**(1), (2007) 71.
- [42] D.P. George, R.R. Volkas, *Kink modes and effective four dimensional fermion and Higgs brane models* Physical Review D **75**(10), (2007) 105007.
- [43] V. Dzhunushaliev, V. Folomeev, D. Singleton, S. Aguilar-Rudametkin, *6D thick branes from interacting scalar fields* Physical Review D **77**(4), 044006 (2008).
- [44] V. Dzhunushaliev, V. Folomeev, M. Minamitsuji, *Thick de Sitter brane solutions in higher dimensions* Physical Review D **79**(2), (2009) 024001.
- [45] D. Bazeia, A. Gomes, L. Losano, R. Menezes, *Braneworld models of scalar fields with generalized dynamics* Physical Review B **671**(3), (2009) 402.
- [46] D. Bazeia, A.R. Gomes, L. Losano, *Gravity localization on thick branes: a numerical approach* International Journal of Modern Physics A **24**(06), (2009) 1135.
- [47] A.E.R. Chumbes, M.B. Hott, *Nonpolynomial potentials with deformable topological structures* Physical Review D **81**(4), (2010) 045008.
- [48] Z. H. Zhao, Y. X. Liu, H. T. Li, Y. Q. Wang, *Effects of the variation of mass on fermion localization and resonances on thick branes* Physical Review D **82**(8), (2010) 084030.
- [49] Y. X. Liu, Y. Zhong, K. Yang, *Scalar-kinetic branes* Europhysics Letters **90**(5), (2010) 51001.
- [50] Y. X. Liu, Y. Zhong, Z. H. Zhao, H. T. Li, *Domain wall brane in squared curvature gravity* Journal of High Energy Physics **1106**(6), (2011) 135.
- [51] H. T. Li, Y. X. Liu, Z. H. Zhao, H. Guo, *Fermion resonances on a thick brane with a piecewise warp factor* Physical Review D **83**(4), (2011) 045006.
- [52] H. Guo, Y. X. Liu, Z. H. Zhao, F. W. Chen, *Thick branes with a nonminimally coupled bulk-scalar field* Physical Review D **85**(12), (2012)124033.
- [53] H. Guo, Y. X. Liu, S. W. Wei, C. E. Fu, *Gravity localization and effective Newtonian potential for Bent thick branes* Europhysics Letters **97**(6), (2012) 60003.
- [54] D. Bazeia, F.A. Brito, F.G. Costa, *Braneworld solutions from scalar field in bimetric theory* Physical Review D **87**, (2013) 065007.
- [55] A. de Souza Dutra, G.P. de Brito, J.M.H. da Silva, *Method for obtaining thick brane models* Physical Review D **91**, (2014) 086016.
- [56] G. Germán, A. Herrera-Aguilar, D. Malagón-Morejón, I. Quiros, R. da Rocha, *Study of field*

- fluctuations and their localization in a thick braneworld generated by gravity nonminimally coupled to a scalar field with the Gauss-Bonnet term* Physical Review D **89**(2), (2014) 026004.
- [57] D. Bazeia, A.S. Lobáó. Jr, R. Menezes, A.Y. Petrov, A.J. da Silva, *Braneworld solutions for  $F(R)$  models with non-constant curvature* Physical Review B **729**, (2014) 127.
  - [58] D. Bazeia, A.S. Lobáó. Jr, R. Menezes, *Thick brane models in generalized theories of gravity* Physical Review B **743**, (2015) 98.
  - [59] D. Bazeia, A.S. Lobáó. Jr, L. Losano, R. Menezes, G.J. Olmo, *Braneworld solutions for modified theories of gravity with nonconstant curvature* Physical Review D **91**(12), (2015) 124006.
  - [60] D. Bazeia, L. Losano, R. Menezes, G.J. Olmo, D. Rubiera-Garcia, *Thick brane in  $f(R)$  gravity with Palatini dynamics* European Physical Journal C **75**(12), (2015) 569.
  - [61] B. Guo, Y.X. Liu, K. Yang, *Brane worlds in gravity with auxiliary fields* European Physical Journal C **75**, (2015) 63.
  - [62] L. Visinelli, N. Bolis, and S. Vagnozzi, *Brane-world extra dimensions in light of GW170817* Physical Review D **97**, (2018) 064039.
  - [63] S. SenGupta, *Aspects of warped braneworld models* , arXiv:0812.1092[hep-th].
  - [64] V. Dzhunushaliev, V. Folomeev, and M. Minamitsuji, *Thick brane solutions* Reports on Progress in Physics **73**(6), (2010) 066901.
  - [65] M. Shifman, *LARGE EXTRA DIMENSIONS: Becoming acquainted with an alternative paradigm* International Journal of Modern Physics A **25**(02n03), (2010) 199.
  - [66] T.G. Rizzo, H.A. Morales-Tecotl, L.A. Urena-Lopez, R. Linares-Romero, H.H. Garcia-Compean, *Introduction to Extra Dimensions* AIP Conf. Proc. **1256**, (2010) 27.
  - [67] R. Maartens, K. Koyama, *Brane-World Gravity* Living Reviews in Relativity **13**(1), (2010) 5.
  - [68] O. Arias, R. Cardenas, I. Quiros, *Thick brane worlds arising from pure geometry* Nuclear Physics B **643**(1-3), (2002) 187.
  - [69] N. Barbosa-Cendejas, A. Herrera-Aguilar, *4D gravity localized in non  $\mathbb{Z}_2$  -symmetric thick branes* Journal of High Energy Physics **2005**(10), (2005) 101.
  - [70] N. Barbosa-Cendejas, A. Herrera-Aguilar, *Localization of 4D gravity on pure geometrical thick branes* Physical Review D **73**, (2006) 084022.
  - [71] N. Barbosa-Cendejas, A. Herrera-Aguilar, M.A.R. Santos, C. Schubert, *Mass gap for gravity*

- localized on Weyl thick branes* Physical Review D **77**(12), (2008) 126013.
- [72] Y. X. Liu, K. Yang, Y. Zhong, *de Sitter thick brane solution in Weyl geometry* Journal of High Energy Physics **2010**(10), (2010) 066901.
- [73] V. Dzhunushaliev, V. Folomeev, B. Kleihaus, J. Kunz, *Some thick brane solutions in  $f(R)$ -gravity* Journal of High Energy Physics **2010**(4), (2010) 130.
- [74] H. Liu, H. Lü, Z.L. Wang,  *$f(R)$  gravities, Killing spinor equations, "BPS" domain walls and cosmology* Journal of High Energy Physics **2012**(2), (2012) 083.
- [75] J. Yang, Y. L. Li, Y. Zhong, Y. Li, *Thick brane split caused by spacetime torsion* Physical Review D **85**(8), (2012) 084033.
- [76] N. Barbosa-Cendejas, A. Herrera-Aguilar, K. Kanakoglou, U. Nucamendi, I. Quiros, *Mass hierarchy, mass gap and corrections to Newton's law on thick branes with Poincaré symmetry* General Relativity and Gravitation **46**(1), (2014) 1631.
- [77] Y. Zhong, Y. X. Liu, *Pure geometric thick  $f(R)$ -branes: stability and localization of gravity* European Physical Journal C **76**(6), (2016) 321.
- [78] R. Myrzakulov, L. Sebastiani, S. Vagnozzi, *Inflation in  $f(R, \phi)$ -theories and mimetic gravity scenario* European Physical Journal C **75**(6), (2015) 444.
- [79] V. Dzhunushaliev, V. Folomeev, G. Nurtaeva, S. D. Odintsov, *Thick branes in higher-dimensional  $f(R)$  gravity* International Journal of Geometric Methods in Modern Physics **17**(3), (2020) 2050036.
- [80] Y. X. Liu, X. H. Zhang, L. D. Zhang, Y. S. Duan, *Localization of matters on pure geometrical thick branes* Journal of High Energy Physics **2008**(02), (2008) 067.
- [81] Y. X. Liu, L. D. Zhang, L. J. Zhang, Y. S. Duan, *Fermions on thick branes in the background of sine-Gordon kinks* Physical Review D **78**(6), (2008) 065025.
- [82] Y. X. Liu, L. D. Zhang, S. W. Wei, Y. S. Duan, *Localization and mass spectrum of matters on Weyl thick branes* Journal of High Energy Physics **2008**(08), (2008) 041.
- [83] Y. X. Liu, H. Guo, C. E. Fu, H. T. Li, *Localization of gravity and bulk matters on a thick anti-de Sitter brane* Physical Review D **84**(4), (2011) 044033.
- [84] Z. H. Zhao, Q. Y. Xie, Y. Zhong, *New localization method of  $U(1)$  gauge vector field on flat branes in (asymptotic)  $AdS_5$  spacetime* Classical and Quantum Gravity **32**(3), (2015) 035020.
- [85] Z. H. Zhao, Q. Y. Xie, *Localization of  $U(1)$  gauge vector field on flat branes with five-*

- dimension (asymptotic) AdS<sub>5</sub> spacetime* Journal of High Energy Physics **2018**(5), (2018) 72.
- [86] S. Randjbar-Daemi, M. Shaposhnikov, *Fermion zero-modes on brane-worlds* Physics Letters B **492**(3-4), (2000) 361.
  - [87] S.L. Dubovsky, V.A. Rubakov, P.G. Tinyakov, *Brane world: Disappearing massive matter* Physical Review D **62**(10), (2000) 105011.
  - [88] T. Gherghetta, M. Shaposhnikov, *Localizing Gravity on a Stringlike Defect in Six Dimensions* Physical Review Letters **85**(2), (2000) 240.
  - [89] Y. Grossman, M. Neubert, *Neutrino masses and mixings in non-factorizable geometry* Physics Letters B **474**(3-4), (2000) 361.
  - [90] A. Kehagias, K. Tamvakis, *Localized gravitons, gauge bosons and chiral fermions in smooth spaces generated by a bounce* Physics Letters B **504**(1-2), (2001) 38.
  - [91] I. Oda, *Localization of matters on a string-like defect* Physics Letters B **496**(1-2), (2000) 113.
  - [92] I.P. Neupane, *Consistency of higher derivative gravity in the brane background* Journal of High Energy Physics **2000**(09), (2000) 040.
  - [93] I.P. Neupane, *Completely localized gravity with higher curvature terms* Classical and Quantum Gravity **19**(21), (2002) 5507.
  - [94] S. Ichinose, *Fermions in Kaluza-Klein and Randall-Sundrum theories* Physical Review D **66**(10), (2002) 104015.
  - [95] C. Ringeval, P. Peter, J.P. Uzan, *Localization of massive fermions on the brane* Physical Review D **65**, (2002) 044016.
  - [96] A.A. Andrianov, V.A. Andrianov, P. Giacconi, R. Soldati, *Domain wall generation by fermion self-interaction and light particles* Journal of High Energy Physics **2003**(07), (2003) 063.
  - [97] A.A. Andrianov, V.A. Andrianov, P. Giacconi, R. Soldati, *Brane world generation by matter and gravity* Journal of High Energy Physics **2005**(07), (2005) 003.
  - [98] R. Koley, S. Kar, *A novel braneworld model with a bulk scalar field* Physics Letters B **623**(3), (2005) 244.
  - [99] R. Koley, S. Kar, *Scalar kinks and fermion localization in warped spacetimes* Classical and Quantum Gravity **22**(4), (2005) 753.



- [100] Y. Q. Wang, T. Y. Si, Y. X. Liu, Y. S. Duan, *Fermionic zero modes in self-dual vortex background* Modern Physics Letters A **20**(39), (2005) 3045.
- [101] A. Melfo, N. Pantoja, J.D. Tempo, *Fermion localization on thick branes* Physical Review D **73**(4), (2006) 044033.
- [102] A. Salvio, M. Shaposhnikov, *Chiral asymmetry from a 5D Higgs mechanism* Journal of High Energy Physics **0711**, (2007) 037.
- [103] T.R. Slatyer, R.R. Volkas, *Cosmology and fermion confinement in a scalar-field-generated domain wall brane in five dimensions* Journal of High Energy Physics **2007**(04), (2007) 062.
- [104] Y. X. Liu, L. Zhao, Y. S. Duan, *Localization of fermions on a string-like defect* Journal of High Energy Physics **0704**, (2007) 097.
- [105] Y. X. Liu, L. Zhao, X. H. Zhang, Y. S. Duan, *Fermions in self-dual vortex background on a string-like defect* Nuclear Physics B **785**(3), (2007) 234.
- [106] Y. Brihaye, T. Delsate, *Remarks on bell-shaped lumps: Stability and fermionic modes* Physical Review D **78**(2), (2008) 025014.
- [107] L. Zhao, Y. X. Liu, Y. S. Duan, *Fermions in gravity and gauge backgrounds on a brane world* Modern Physics Letters A **23**(15), (2008) 1129.
- [108] M.O. Tahim, W.T. Cruz, C.A.S. Almeida, *Tensor gauge field localization in branes* Physical Review D **79**, (2009) 085022.
- [109] A. Flachi, M. Minamitsuji, *Field localization on a brane intersection in anti-de Sitter space-time* Physical Review D **79**(10), (2009) 104021.
- [110] Y. Kodama, K. Kokubu, N. Sawado, *Localization of massive fermions on the baby-Skyrmion branes in 6 dimensions* Physical Review D **79**(6), (2009) 065024.
- [111] Y. X. Liu, J. Yang, Z. H. Zhao, C. E. Fu, Y. S. Duan, *Fermion localization and resonances on a de Sitter thick brane* Physical Review D **80**(6), (2009) 065019.
- [112] Y. X. Liu, C. E. Fu, L. Zhao, Y. S. Duan, *Localization and mass spectra of fermions on symmetric and asymmetric thick branes* Physical Review D **80**(6), (2009) 065020.
- [113] Y. X. Liu, Z. H. Zhao, S. W. Wei, Y. S. Duan, *Bulk matters on symmetric and asymmetric de Sitter thick branes* Journal of Cosmology and Astroparticle Physics **2009**(02), (2009) 003.
- [114] C.A.S. Almeida, R. Casana, M.M. Ferreira, A.R. Gomes, *Fermion localization and resonances on two-field thick branes* Physical Review D **79**(12), (2009) 125022.
- [115] R. Koley, J. Mitra, S. SenGupta, *Fermion localization in a generalized Randall-Sundrum*



- model* Physical Review D **79**, (2009) 041902.
- [116] L.B. Castro, *Fermion localization on two-field thick branes* Physical Review D **83**, (2011) 045002.
  - [117] A.E.R. Chumbes, A.E.O. Vasquez, M.B. Hott, *Fermion localization on a split brane* Physical Review D **83**(10), (2011) 105010.
  - [118] W.T. Cruz, A.R. Gomes, C.A.S. Almeida, *Fermions on deformed thick branes* European Physical Journal C **71**(11), (2011) 1790.
  - [119] C. E. Fu, Y. X. Liu, H. Guo, *Bulk matter fields on two-field thick branes* Physical Review D **84**(4), (2011) 044036.
  - [120] O. Castillo-Felisola, I. Schmidt, *Localization of fermions in different domain wall models* Physical Review D **86**, (2012) 024014.
  - [121] T. Delsate, N. Sawado, *Localizing modes of massive fermions and a  $U(1)$  gauge field in the inflating baby-Skyrmion branes* Physical Review D **85**(6), (2012) 065025.
  - [122] A.A. Andrianov, V.A. Andrianov, O.O. Novikov, *CP Violation in the models of fermion localization on a domain wall (brane)* Theor. Math. Phys. **175**, (2013) 735.
  - [123] V.K. Oikonomou, *Localized fermions on domain walls and extended supersymmetric quantum mechanics* Classical and Quantum Gravity **31**(2), (2013) 025018.
  - [124] M. Gogberashvili, P. Midodashvili, *Fermions in the 5D gravity-scalar standing wave braneworld* International Journal of Modern Physics A **29**(24), 1450141 (2014).
  - [125] L.J.S. Sousa, C.A.S. Silva, D.M. Dantas, C.A.S. Almeida, *Vector and fermion fields on a bouncing brane with a decreasing warp factor in a string-like defect* Physics Letters B **731**, (2014) 64.
  - [126] Y. X. Liu, Z. G. Xu, F. W. Chen, S. W. Wei, *New localization mechanism of fermions on braneworlds* Physical Review D **89**(8), (2014) 086001.
  - [127] N. Barbosa-Cendejas, D. Malagón-Morejón, R.R. Mora-Luna, *Universal spin-1/2 fermion field localization on a 5D braneworld* General Relativity and Gravitation **47**(7), (2015).
  - [128] H. Guo, Q. Y. Xie, C. E. Fu, *Localization and quasilocalization of a spin-1/2 fermion field on a two-field thick braneworld* Physical Review D **92**(10), (2015) 106007.
  - [129] I.C. Jardim, G. Alencar, R.R. Landim, R.N.C. Filho, *Solutions to the problem of Elko spinor localization in brane models* Physical Review D **91**, (2015) 085008.
  - [130] R. Cartas-Fuentevilla, A. Escalante, G. Germán, A. Herrera-Aguilar, R.R. Mora-Luna,

- Coulomb's law corrections and fermion field localization in a tachyonic de Sitter thick braneworld* Journal of Cosmology and Astroparticle Physics **1605**, 026 (2016).
- [131] Y. Y. Li, Y. P. Zhang, W. D. Guo, Y. X. Liu, *Fermion localization mechanism with derivative geometrical coupling on branes* Physical Review D **92**, (2017) 115003.
- [132] V.F. Mukhanov, L.A. Kofman, D.Y. Pogosyan, *Cosmological perturbations in the inflationary universe* Physics Letters B **193**(4), (1987) 427.
- [133] J.D. Barrow, S. Cotsakis, *Inflation and the conformal structure of higher-order gravity theories* Physics Letters B **214**(4), (1988) 515.
- [134] G. Cognola, E. Elizalde, S. Nojiri, S.D. Odintsov, L. Sebastiani, S. Zerbini, *Class of viable modified  $f(R)$  gravities describing inflation and the onset of accelerated expansion* Physical Review D **77**(4), (2008) 046009.
- [135] A.D. Felice, S. Tsujikawa,  *$f(R)$  Theories* Living Reviews in Relativity **13**(1), (2010) 3.
- [136] T.P. Sotiriou, V. Faraoni,  *$f(R)$  theories of gravity* Reviews of Modern Physics **82**, (2010) 451.
- [137] S. Nojiri, S.D. Odintsov, *Unified cosmic history in modified gravity: From  $F(R)$  theory to Lorentz non-invariant models* Physics Reports **505**(2-4), (2011) 59.
- [138] N. Dimakis, A. Giacomini, A. Paliathanasis, *Black branes in four-dimensional conformal equivalent theories* European Physical Journal C **78**(9), (2018) 751.
- [139] N. Dimakis, A. Giacomini, A. Paliathanasis, *(Compactified) black branes in four dimensional  $f(R)$ -gravity* Physics Letters B **777**(361-368), (2018) 059.
- [140] M. Parry, S. Pichler, D. Deeg, *Higher-derivative gravity in brane world models* Journal of Cosmology and Astroparticle Physics **2005**(04), (2005) 014.
- [141] N. Deruelle, M. Sasaki, Y. Sendouda, *Junction Conditions in  $f(R)$  Theories of Gravity* Progress of Theoretical Physics **119**(2), (2008) 237.
- [142] A. Balcerzak, M.P. Dabrowski, *Generalized Israel junction conditions for a fourth-order brane world* Physical Review D **77**(2), (2008) 023524.
- [143] A. Balcerzak, M.P. Dabrowski, *Gibbons-Hawking boundary terms and junction conditions for higher-order brane gravity models* Journal of Cosmology and Astroparticle Physics **0901**(01), (2009) 018.
- [144] Y. Zhong, Y. X. Liu, K. Yang, *Tensor perturbations of  $f(R)$ -branes* Physics Letters B **699**(5), (2011) 398.

- [145] T.R.P. Carames, M.E.X. Guimaraes, J.M. H. da Silva, *Effective gravitational equations for  $f(R)$  braneworld models* Physical Review D **87**(10), (2013) 106011.
- [146] D. Bazeia, R. Menezes, A.Y. Petrov, A.J. da Silva, *On the many-field  $f(R)$  brane* Physics Letters B **726**(1-3), (2013) 523.
- [147] Z. G. Xu, Y. Zhong, H. Yu, Y. X. Liu, *The structure of  $f(R)$ -brane model* European Physical Journal C **75**(8), (2015) 368.
- [148] B. M. Gu, B. Guo, H. Yu, Y. X. Liu, *Tensor perturbations of Palatini  $f(\mathbf{R})$  branes* Physical Review D **92**(2), (2015) 02401.
- [149] H. Yu, Y. Zhong, B. M. Gu, Y. X. Liu, *Gravitational resonances on  $f(R)$ -brane* European Physical Journal C **76**(4), (2016) 195.
- [150] Z. Q. Cui, Y. X. Liu, B. M. Gu, L. Zhao, *Linear stability of  $f(R, \varphi, X)$  thick branes: tensor perturbations* Journal of High Energy Physics **2018**(11), (2018) 083.
- [151] A. Herrera-Aguilar, D. Malagón-Morejón, R. Rigel Mora-Luna, *Localization of gravity on a de Sitter thick braneworld without scalar fields* Journal of High Energy Physics **2010**(11), (2010) 015.
- [152] Q. Y. Xie, J. Yang, L. Zhao, *Resonance mass spectra of gravity and fermion on Bloch branes* Physical Review D **88**(10), (2013) 105014 .