

Supernovae Ia in 2019 (review): a rising demand for spherical explosions

NOAM SOKER^{1,2}

¹*Department of Physics, Technion Israel Institute of Technology, Haifa 3200003, Israel*

²*Guangdong Technion Israel Institute of Technology, Guangdong Province, Shantou 515069, China*

ABSTRACT

I review new studies of type Ia supernovae (SNe Ia) from 2019, and use these to improve the comparison between the five binary SN Ia scenarios. New low polarisation measurements solidify the claim that most SN Ia explosions are globally spherically symmetric (clumps are possible). Explosions by dynamical processes, like explosions that take place during a merger process of two white dwarfs (WDs) in the double degenerate (DD) scenario, or during an accretion process in the double detonation (DDet) scenario and in the single degenerate (SD) scenario, lead to non-spherical explosions, in contradiction with observations of normal SNe Ia. I argue that these (DD, DDet, SD) scenarios account mainly for peculiar SNe Ia. The explosion of a Chandrasekhar mass (M_{CH}) WD (deflagration to detonation process) has a global spherical structure that is compatible with observations. To reach spherical explosions, SN Ia scenarios should allow for a time delay between the formation of an M_{CH} -WD and its explosion. As such, I split the DD scenario to a channel without merger to explosion delay (MED) time (that forms mainly peculiar SNe Ia), and a channel with a MED, the DD-MED channel (scenario). I speculate that the main contributors to normal SNe Ia are the core degenerate (CD) scenario, the DD-MED scenario, both have M_{CH} spherical explosions, and the DD scenario that has sub- M_{CH} non-spherical explosions.

Keywords: supernovae: general; binaries: close; white dwarfs; Astrophysics - Solar and Stellar Astrophysics; Astrophysics - High Energy Astrophysical Phenomena

1. INTRODUCTION

The year 2019 saw the continuation of the debate between the supporters of the five binary scenarios for type Ia supernovae (SNe Ia; in Soker 2019 I justify this partition to five scenarios). Despite several scientific meetings and tens of papers (e.g., Liu, & Stancliffe 2018; Ablimit, & Maeda 2019b; Ashall et al. 2019b; Bravo et al. 2019; Brown et al. 2019; Calder et al. 2019; Chakradhari et al. 2019; Chen et al. 2019b; Graur 2019; Graur et al. 2019; Heringer et al. 2019b; Konyves-Toth et al. 2019; Liu et al. 2019; Miles et al. 2019; Moriya et al. 2019; Siebert et al. 2019; Taubenberger et al. 2019; Wu et al. 2019; Meng 2019; Yao et al. 2019; Di Stefano 2020; Han et al. 2020; Kushnir, & Katz 2020a,b; Leung, & Nomoto 2020; Mazzali et al. 2020; Miller et al. 2020; Pan et al. 2020; Pellegrino et al. 2020; Shingles et al. 2020), the community did not come closer to a consensus. A prominent example is

the existence of studies that argue for Chandrasekhar SN Ia explosion alongside studies that argue for sub-Chandrasekhar SNe Ia, and studies that argue for both (e.g., Levanon, & Soker 2019b; Sarbadhicary et al. 2019; Seitzzahl et al. 2019). However, I do find some significant results in studies from 2019 that constrain some scenarios and some channels within scenarios. Since there are several reviews of SNe Ia from the years 2018-2019 (Livio & Mazzali 2018; Wang 2018b; Soker 2018; Jha et al. 2019; Ruiz-Lapuente 2019; older reviews include, e.g., Maoz et al. 2014 and Maeda, & Terada 2016, a summary of several evolutionary routes to SNe Ia Tutukov, & Fedorova 2007), and a review that was written after this review (Ruiter 2020), that together cover all relevant aspects of SNe Ia, in the present study I concentrate on the new results from the year 2019. I discuss only scenarios that involve binary stellar systems, and will not touch single star scenarios (e.g., Clavelli 2019; Antoniadis et al. 2020; I note that the evolution after the formation of a massive core inside a single AGB star in the scenario of Antoniadis et al.

2020 has some similar ingredients to the core degenerate scenario).

In an earlier review (Soker 2018) I summarised the five binary SN Ia scenarios in a table, which I find to be the most convenient presentation for this review as well. I bring an updated version of that table in Table 1, and I will refer to this table throughout the review (note that in the new table here I split one scenario to two channels). The contributions of the scenarios to normal and peculiar SNe Ia are my present estimates, and one should note that other researchers might have largely different estimates, e.g., Ruiter 2020.

One can characterise the scenarios by the following properties. (1) Number of stars at explosion, N_{exp} , 1 or 2. (2) Whether a companion survives the explosion, ($N_{\text{sur}} = 0$ or 1). (3) Whether the mass of the exploding WD(s) at explosion is around the Chandrasekhar mass limit, M_{Ch} , or whether it is sub- M_{Ch} . I list these characteristics in the third row of Table 1. To these we can add the partition to channels that have a merger (or accretion) to explosion delay time (MED; Soker 2018), and those that do not (second row of Table 1). Specifically, in systems that have MED a merger event or an accretion process bring the WD to a mass of close to M_{Ch} , and the explosion occurs a time t_{MED} after the WD has reached the mass $M_{\text{WD}} \simeq M_{\text{Ch}}$, where t_{MED} is much larger than the dynamical time.

Both the core degenerate (CD) and the double degenerate (DD) with MED (DD-MED) scenarios have $[N_{\text{exp}}, N_{\text{sur}}, \mathbf{M}] = [1, 0, M_{\text{Ch}}]$. The difference between the CD scenario and the DD-MED scenario is that in the DD-MED scenario there is a delay time from the formation of the two WDs to merger that is determined by gravitational waves, while in the CD scenario the merger occurs during the common envelope evolution (CEE) phase, and so gravitational waves play no role. A big challenge to the DD-MED scenario is that in many cases the merger of two CO WDs does not lead to a SNe Ia with a MED (e.g., Wu et al. 2019).

The non-detection of the expected companions inside supernova remnants (SNRs) in the single degenerate (SD) scenario (e.g., Li, C. et al. 2019; but I note that Ablimit, & Maeda 2019a argue that it will be hard to detect the companion for a magnetic WD), the general no detection of hydrogen (e.g., Holmbo et al. 2019), and/or the very low mass of hydrogen even when rare observations do show hydrogen (e.g., Kollmeier et al. 2019; Prieto et al. 2020; Dessart et al. 2020), suggest that for normal SNe Ia *all* cases must have a MED, what originally was termed spin-up/spin-down evolution as rotation keeps the WD from exploding until it loses angular momentum (e.g., Piersanti et al. 2003;

Di Stefano et al. 2011; Justham 2011; Boshkayev et al. 2014; Wang et al. 2014; Benvenuto et al. 2015). For example, Kuuttila et al. (2019) and Graur, & Woods (2019) strengthen earlier claims that rule out hot and luminous progenitors as expected in some channels of the SD scenario. Their arguments do not apply to the SD-MED scenario (spin-up/spin-down) that in principle might account for a fraction of normal SNe Ia. Meng, & Li (2019) suggest that the common-envelope wind channel of the SD-MED scenario (Meng, & Podsiadlowski 2017), can leave a surviving sdB companion which is hard to detect. They further estimate that SNe Ia with an sdB companion might contribute 22% of all SNe Ia. In the present study I refer to the SD-MED channel when referring to normal SNe Ia, and to the SD scenario without MED when referring to peculiar SNe Ia, unless stated otherwise. For that, I do not list as severe problems in the table the no detection of a surviving companion and the no detection of hydrogen in the ejecta. However, a companion should still survive even if a faint one, and in some cases we should expect hydrogen.

I notice that recent results by Toonen et al. (2018) solidified earlier claims that the WD-WD collision (WWC) scenario might at most account for less than one per cent of all SNe Ia (later studies have reached the same qualitative conclusion, e.g., Haim, & Katz 2018; Hallakoun & Maoz 2019; Hamers & Thompson 2019). The second severe problem of the WWC scenario is that the collision of the two WDs lead to a highly non-spherical explosion (e.g., Kushnir et al. 2013), contrary to both the morphologies of SNRs Ia and to the new polarisation studies of SNe Ia. I do note that there are some studies in 2019 that claim for the WWC scenario (e.g., Wygoda et al. 2019a,b), and that this scenario might account for rare types of peculiar SNe Ia. In particular, Valley et al. (2020) find that some SNe Ia show bimodal velocity distributions of ^{56}Ni decay products, and Livneh, & Katz (2020) find evidence for asymmetrical Si distribution. These papers claim that their results support the WWC scenario for many SNe Ia. In the present review I attribute such bimodal and asymmetrical distributions to large ^{56}Ni and Si clumps in the explosion mechanism, although the global explosion in normal SNe Ia is spherical (see, e.g., Levanon, & Soker 2019b; Magee et al. 2020, for papers on the ^{56}Ni distribution).

Table 1 has several significant differences from the previous comparison table (Soker 2018).

1. In my previous review (Soker 2018) I used the comparison of the five scenarios with each other and with observations to argue that scenarios must al-

Table 1. Confronting SN Ia scenarios with observations

Scenario ^[1,2]	Core Degenerate (CD)	Double Degenerate (DD)	Double Degenerate (DD-MED)	Double Detonation (DDet)	Single Degenerate (SD-MED)	WD-WD collision (WWC)
Channel by MED	MED built-in.	No MED	MED	No MED (not allowed)	MED (with no MED peculiar SNe Ia)	No MED
$[N_{\text{exp}}, N_{\text{sur}}, M]$	$[1, 0, M_{\text{Ch}}]$	$[2, 0, \text{sub-}M_{\text{Ch}}]$	$[1, 0, M_{\text{Ch}}]$	$[2, 1, \text{sub-}M_{\text{Ch}}]$	$[2, 1, M_{\text{Ch}}]$	$[2, 0, \text{sub-}M_{\text{Ch}}]$
Presence of 2 opposite Ears in some SNR Ia ^[3]	Explained by the SN inside planetary nebula (SNIP) mechanism.	Low mass Ears if jets during merger (Tsebreko & Soker 2013).	Requires a short gravitational waves delay time shortly after CEE; unlikely.	No Ears are expected for He WD companion.	OP; ^[4] Ears by jets from accreting WD (Tsebreko & Soker 2013).	No Ears are expected
Spherical SNRs + low polarisations	Expected in all cases.	Cannot explain.	Expected in all cases.	Cannot explain.	Explained in most cases (with MED).	Cannot explain.
$\approx 1M_{\odot}$ CSM in Keplers SNR	The massive CSM shell might be a PN.	No CSM shell	Requires a short gravitational waves delay time shortly after CEE; unlikely.	Any CSM is of a much lower mass.	OP; ^[4] Can be explained by heavy mass loss from an AGB donor.	No CSM shell
The need to synthesize ^{55}Mn and other elements.	M_{Ch} can do it	Not possible	M_{Ch} can do it	Not possible	M_{Ch} can do it	Unlikely
Main Scenario Predictions	1. Single WD explodes 2. Massive CSM in some cases (SNIP). 3. $M_{\text{WD}} \simeq M_{\text{Ch}}$	1. Sufficient WD-WD close binaries 2. $\text{DTD} \propto t^{-1}$ 3. $M_{\text{WD}} < 1.2M_{\odot}$	1. Single WD explodes 2. $M_{\text{WD}} \simeq M_{\text{Ch}}$	1. A companion survives 2. Asymmetrical explosion 3. $M_{\text{WD}} < 1.2M_{\odot}$	1. A companion survives 2. $M_{\text{WD}} \simeq M_{\text{Ch}}$	Asymmetrical explosion
General Strong Characteristics	1. Explains some SN Ia with H-CSM 2. Spherical explosions 3. Many explosions with $M_{\text{WD}} \simeq M_{\text{Ch}}$ 4. Explains large SNe Ia population shortly after CEE	1. Explains very well the delay time distribution (DTD) 2. Ignition easily achieved	Explains very well the delay time distribution (DTD) 2. Many explosions with $M_{\text{WD}} \simeq M_{\text{Ch}}$ 3. Spherical explosions	Ignition achieved	1. Accreting massive WDs exist 2. Many explosions with $M_{\text{WD}} \simeq M_{\text{Ch}}$ 3. Spherical explosions	Ignition easily achieved
Work for future studies	1. Ignition process 2. Merge during CEE 3. To solidify the claim for M_{Ch} WDs (Bear, & Soker 2018) 4. DTD	1. To derive spherical explosions	1. Ignition process 2. Merge process 3. To solidify the claim for M_{Ch} WDs (Bear, & Soker 2018)	1. To explain the non-detection of helium 2. To find surviving companions	1. Ignition process 2. To explain the DTD and number of SNe Ia 3. To find surviving companions	1. Find a way to account for $> 1\%$ of normal SNe Ia. 2. Examine which peculiar SNe Ia the WWC might account for.
Contribution to normal SNe Ia ^[5]	$\approx 20 - 50\%$	$\approx 20 - 40\%$	$\approx 20 - 40\%$	$\approx 0 - 10\%$	$\approx 0 - 10\%$	$\ll 1\%$
Contribution to peculiar SNe Ia ^[5]	$\approx 0 - 10\%$	$\approx 30 - 70\%$	$\approx 0 - 10\%$	$\approx 10 - 30\%$	$\approx 20 - 50\%$ by the SD scenario without MED	$\approx 1\%$

Notes:

[1] Scenarios for SN Ia by alphabetical order.

[2] The first three rows are the names and explosion characteristics; N_{exp} is the number of stars at explosion (1 or 2); $N_{\text{sur}} = 1$ if a companion survives the explosion and $N_{\text{sur}} = 0$ if no companion survives the explosion; M_{Ch} and $\text{sub-}M_{\text{Ch}}$, indicate whether the mass of the exploding WD(s) at explosion is around the Chandrasekhar mass limit or whether it is sub-Chandrasekhar, respectively.

[3] The observations in rows 4-10 refer to normal SNe Ia.

[4] OP (only peculiar) means that the SD scenario without MED can explain this observation, but not the SD-MED scenario. Therefore, explanation of the particular observation is possible within the SD channels only for peculiar SNe Ia.

[5] Last two rows present my very crude estimates of the contribution of each channel to normal and peculiar SNe Ia, respectively.

low, at least in some cases, for a time delay between the dynamical process of merger or the dynamical process of mass accretion to explosion. This merger to explosion delay (MED) was the main conclusion then. In the present study I incorporate the MED as a property that splits scenarios to different channels (second row in Table 1).

2. The outcomes of the DD scenario when explosion occurs within several dynamical time scales after merger or at much later times (the DD-MED) are very different. For that, I find it necessary to split this scenario to two columns in Table 1. A MED might take place when the merger remnant has a mass of $M_{\text{rem}} \simeq M_{\text{CH}}$.
3. I changed somewhat the rows. For example, to emphasise the new polarisation studies that suggest spherical explosions, I added the fifth row.
4. Following the above changes and new results from 2019, I changed my estimates of the contribution of each channel to normal and peculiar SNe Ia.

I turn to discuss some new studies from 2018/2019 and their implications to the different scenarios. These new studies join older studies (that I do not review here) in challenging one or more scenarios. When comparing theoretical studies to observations these challenges cannot be ignored. There is no scenario that is free from challenges, some that I find hard, or even impossible, to overcome. For example, I do not see how the WWC scenario overcome the two challenges I mentioned above.

I note again that I cite mainly papers from 2019, although in most cases there are earlier relevant papers, because this review is about studies from 2019 (and to some extend from 2018) and their new implications.

★ **Summary of section 1.** It is mandatory to consider all scenarios when comparing observations to theory. There are five (5; not 2) scenarios, with some of them having multiple channels. Most significant splitting to channels is that to the two channels of the DD scenario according to the presence (the DD-MED channel) or not of a MED (merger to explosion delay) time. I find the observation that most normal SNe Ia have globally spherical explosions to be one of the most significant in 2019. This finding, alongside others observations, strongly limit the contribution of the DDet scenario, the SD scenario without MED, and the DD scenario without MED to normal SNe Ia.

2. SPHERICALLY SYMMETRIC EXPLOSIONS

2.1. New polarisation observations

I consider the observations of very low polarisation of SNe Ia (Cikota et al. 2019; Yang et al. 2019) to be the most significant type of observations published in 2019. These observations suggest that most SNe Ia have spherical explosions. The globally spherical morphologies of many SNRs Ia (but not all are spherical, e.g., Alsaberi et al. 2019) also suggest that most SNe Ia have spherical explosions (review by Soker 2018), but the new observations put this conclusion on the forefront (for a summary of polarisation of SNe Ia see, e.g., Meng et al. 2017). I note that Fang et al. (2020) show that they can reproduce the departure of the SNR of SN 1006 from spherical morphology by an interaction of a spherical explosion with an ambient medium with a density discontinuity. As well, Luken et al. (2020) conclude that the young type Ia (Borkowski et al. 2013) SNR G1.9+0.3 that has an asymmetrical structure expands into an inhomogeneous interstellar medium (ISM).

Some features, such as ‘Ears’ and other protrusions can exist in otherwise spherical explosions. Some of these are iron-rich protrusions (‘iron-bullets’; e.g., Tsebreiko, & Soker 2015b; Sato et al. 2020).

Cikota et al. (2019) study the polarisation of 35 SNe Ia, and argue that their results support the possibility of two distinct explosion mechanisms. Their analysis shows the peak polarisation of the Si II line to be consistent with the expectation of the DDet scenario and of the delayed-detonation mechanism of M_{CH} explosions. The violent merger of the DD scenario predicts (e.g., Bulla et al. 2016a) too high polarisation (e.g., for 34 out of 35 SNe Ia).

Yang et al. (2019) note that their detection of low polarisation in the normal SN Ia SN 2018gv implies a high degree of spherically symmetric explosion, and that this in turn is consistent with the expected morphology of the delayed detonations explosion mechanism and is inconsistent with the merger-induced explosion mechanism.

For the benefit of the discussion in section 5, I recall that not all SNe Ia have low polarisation. Cikota et al. (2017) find that the polarisation curves of some SNe Ia sight-lines are similar to those of some proto-planetary nebulae (PNe). They claim that this suggests that some SNe Ia explode inside the wind of a post-asymptotic giant branch star. The case of a SN Ia that explodes inside a proto-PN or a PN is termed SNIP (Tsebreiko & Soker 2015a), for a SN inside a PN (Dickel & Jones 1985).

The findings that most SNe Ia have global spherically symmetric explosions strongly challenges several explosion mechanisms. Ferrand et al. (2019) show in a recent paper that morphological signatures of the explosion can still be detected hundreds of years after explosion, so

that an interaction with the ISM cannot erase global asymmetries of SN Ia explosions.

I here discuss only two recent papers on explosion mechanisms. But I do note that alongside a global spherical structure of the SN Ia ejecta, the ejecta might be clumpy (e.g., Millard et al. 2019; Sato et al. 2019), particularly the iron group elements (e.g., Maguire et al. 2018).

2.2. The DDet scenario

I consider first the DDet scenario. This scenario has some strong points, e.g., Townsley et al. (2019) perform hydrodynamical simulations of the DDet scenario and find that it can account for the brightness and spectra of SNe Ia (see also, e.g., Polin et al. 2019a). Bulla et al. (2020) find the early light curve of many SNe Ia to be compatible with the DDet scenario. However, the observations that suggest spherical SN Ia explosions strongly challenge this scenario. I do note that Bulla et al. (2016b) calculate the polarisation of the DDet scenario and find it to be low enough to be compatible with observations. However, Bulla et al. (2016b) use an explosion model from Fink et al. (2010) that does not include the roles of the companion (helium-donor) star in causing deviation from a spherical explosion. Namely, the companion forms an empty conical region in the ejecta by blocking the SN ejecta along its direction, and the companion causes a departure from symmetry by its orbital motion (see below). Bulla et al. (2016b) perform their simulations in 2D by adopting an azimuthal symmetry about the z-axis, rather than in full 3D. For that, it might be that Bulla et al. (2016b) underestimate the polarisation of the DDet scenario.

The first process that leads to a non-spherical explosion is the collision of the ejecta with the surviving companion. This collision forms a conical region behind the companion where the density is very low (e.g., Tanikawa et al. 2018, 2019; Bauer et al. 2019). But there is another process that causes deviation from spherical explosion.

Consider the “dynamically driven double-degenerate double-detonation” (D^6) scenario that Shen et al. (2018) suggest as an explanation to the three hypervelocity WDs that they identify with Gaia (see simulation by Tanikawa et al. 2019). Each of this three WDs is the companion that survived a DDet SN Ia. Neunteufel et al. (2019) find that the parameter space of the DDet scenario with a non-degenerate helium donor might account for no more than 3% of all SNe Ia. In the D^6 model, though, the helium-donor is a WD.

According to Shen et al. (2018), the respective velocities of the three hypervelocity WDs relative to the Galaxy are $v_2 \simeq 1300 \text{ km s}^{-1}$, 2300 km s^{-1} , and 2400 km s^{-1} , with large uncertainties. In the DDet scenario these velocities are the pre-explosion orbital velocities of the respective helium-donor WD companion in each system. To achieve a high velocity of $v_2 \simeq 2000 \text{ km s}^{-1}$ the mass of the WD companion is $M_2 \simeq 1M_\odot$ (Shen et al. 2018). Since the exploding WD has a similar mass, the orbital velocity of the exploding WD is $v_{1,\text{orb}} \simeq v_2 \simeq 2300 \text{ km s}^{-1}$ in the two cases above with high hypervelocities. What causes departure from spherical explosion is that during the explosion the companion and the ejecta do not move on straight lines, but rather continue to curve in the sense of their original orbital motion. This implies that the direction of ejection of the center of mass of low-velocity ejecta is not as that of higher velocity ejecta. Let us demonstrate this qualitatively.

Consider then the explosion process of the WD in the presence of a helium-donor star of similar mass. For a demonstrative case I take the spherical density profiles from Dwarkadas & Chevalier (1998)

$$\rho_0(v, t) = \frac{M_e}{8\pi (v_e t)^3} e^{-v/v_e}, \quad v_e = \left(\frac{E_k}{6M_e} \right)^{1/2}, \quad (1)$$

where M_e and E_k are the total ejecta mass and kinetic energy, respectively. For $M_e = 1M_\odot$ and $E_k = 10^{51} \text{ erg}$ we find $v_e = 2895 \text{ km s}^{-1}$. Integration over the volume shows that half of the ejecta mass is within a velocity of $v_h = 7750 \text{ km s}^{-1} \simeq 3.4v_{1,\text{orb}}$, and quarter of the mass within a velocity of $v_q = 5000 \text{ km s}^{-1}$. By the time the coordinate of half the mass expands beyond the companion, half of the mass is still under the influence of the gravity of the companion, and it changes its direction as if the companion continues to orbit the center of mass more or less. This time is $t_h \simeq 2R_{1,\text{orb}}/7750 \text{ km s}^{-1}$, where $R_{1,\text{orb}}$ is the orbit of the exploding WD around the center of mass, which is about equal to that of the surviving companion. During that time the mass still within the orbit of the secondary star would crudely orbit an angle of $\alpha_h \approx 360^\circ v_{1,\text{orb}} t_h / 2\pi R_{1,\text{orb}} \simeq 360^\circ / 3.4\pi = 34^\circ$. In the same manner we find that for quarter of the mass $v_q = 5000 \text{ km s}^{-1} \simeq 2.2v_{1,\text{orb}}$ and $\alpha_q = 53^\circ$.

There is a need for more accurate calculations to calculate the polarisation days after explosion, and to reveal the later SNR morphology. But this simple calculation shows that the D^6 scenario with a companion that escapes the system after explosion with $v_2 \gtrsim 2000 \text{ km s}^{-1}$ has highly non-spherical explosion, contrary to observations.

In the case of $v_2 \simeq 1300 \text{ km s}^{-1}$ the mass of the hypervelocity WD is $M_2 \approx 0.2 M_\odot$, and the orbital velocity of the WD that exploded was $v_{1,\text{orb}} \simeq 300 \text{ km s}^{-1}$. In that case the orbital motion has a minor influence on the explosion morphology, which will nonetheless be non-spherical due to the collision of the ejecta with the companion.

The production of a non-spherical SNe Ia is not the only recent problem of the DDet scenario. [Polin et al. \(2019b\)](#) show that although the DDet scenario can qualitatively reproduce sub-luminous SNe Ia spectra in the nebular phase, these explosions produce too much [CaII] emission compared to most normal SNe Ia.

Another general problem is that sub- M_{Ch} explosions cannot yield some isotopes (e.g., [Bravo 2019](#)), such as ^{55}Mn (seventh row in Table 1).

2.3. The hybrid HeCO channel of the DD scenario

The second study I consider deals with the hybrid HeCO channel of the DD scenario ([Zenati et al. 2019](#); [Perets et al. 2020](#)). (The HeCO hybrid channel of the DD scenario is different than the hybrid CONe channel of M_{Ch} deflagration to detonation explosions, e.g., [Augustine et al. 2019](#)). [Perets et al. \(2020\)](#) simulate the destruction of a HeCO WD by a CO WD with 2D hydrodynamical simulations that include nuclear reactions. [Perets et al. \(2020\)](#) set their initial conditions to have a HeCO torus around the CO WD. They obtain a double detonation process and ignite both the CO WD and the HeCO torus. They can account for several important properties of SNe Ia, like the light curves, spectra, and the range of peak-luminosities. They further argue that together with the contribution from the DD scenario of two massive CO WDs they can reproduce the rate and delay-time distribution of SNe Ia.

However, they face several challenges. Their 2D code forces them to simulate axisymmetrical flow. For that, they could not actually follow the destruction of the HeCO WD and establish whether the destructed HeCO WD forms a torus before ignition. The second challenge is the high density along the symmetry axis inside the CO WD that they obtain in their simulations (their figure 1). Their 2D axisymmetrical numerical grid necessarily forces the shock wave that the ignition of the HeCO torus induces in the CO WD to converge on, or near, the axis (their figure 1 at $t = 7.2 \text{ s}$). Namely, their 2D grid ignites a ring around the symmetry axis, rather than a point in reality. In reality, the ignition that starts in one point will not lead to an axisymmetrical shock wave, but rather will converge on an off-center point (e.g., [Pakmor et al. 2013](#); [Townesley et al. 2019](#); [Gronow et al. 2020](#)). It is not clear whether

[Perets et al. \(2020\)](#) can obtain high enough densities inside the CO WD. I also note that even with these high densities (which might be overestimated) they do not synthesise some elements, such as ^{55}Mn .

I consider the third challenge to be the strongest one. [Perets et al. \(2020\)](#) present their numerical grid up to a distance of $6 \times 10^9 \text{ cm}$ from the center of the flow. From what they present (their figure 1), one sees that the explosion is highly non-spherical. There is a fast polar outflow and much slower equatorial ejecta. It is a challenge for the hybrid HeCO channel of the DD scenario to show they can account for the new low polarisation measurements ([Cikota et al. 2019](#); [Yang et al. 2019](#)), and for globally spherical SNRs Ia (section 2.1).

★ **Summary of section 2.** The new polarisation studies from 2019 ([Cikota et al. 2019](#); [Yang et al. 2019](#)) impose firm constraints on explosion mechanisms to yield explosions that are globally spherical, at least in most cases, if not in all normal SNe Ia. (Small features, such as ‘Ears’ are possible; Table 1.) Interestingly, interactions that seem more easily to explode in numerical simulations have highly non-spherical ejecta. These include the violent merger of the DD scenario, the WWC scenario, possibly the D^6 channel of the DDet scenario, and the newly simulated hybrid HeCO channel of the DD scenario, as I discussed above. The deflagration-to-detonation explosion mechanism of M_{Ch} WDs lead to spherical explosions, and so I favour this explosion mechanism. Although there is a need to work out several steps in this explosion mechanism, the new study by [Poludnenko et al. \(2019\)](#) strongly supports the case for the deflagration-to-detonation explosion mechanism.

3. EARLY EXCESS EMISSION

An interesting class of SNe Ia is that of SNe Ia that show early ($\lesssim 5 \text{ days}$) excess emission in their light curve (e.g., [Jiang et al. 2018](#); [Li, W. et al. 2019](#); [Shappee et al. 2019](#); [Dimitriadis et al. 2019a](#)). In this section I use the class of SNe with early excess emission to emphasise that in many cases we cannot attribute a given type of observations (here it is the early excess emission) to only one scenario (like the SD scenario). We must examine the possibility that other SN Ia scenarios might also account for that specific type of observations.

The SD scenario predicts this kind of early emission in most SNe Ia (e.g. [Kasen 2010](#)), unless there is a very long MED during which the mass-donor radius decrease by an order of magnitude or more, i.e., a giant donor that becomes a WD. My point here is that such an emission is possible also in the DD scenario if the explosion takes place within a time of about $10^3 \text{ s} \lesssim t_{\text{MED}} \lesssim 1 \text{ day}$, as

the ejecta collides with disk-originated matter (DOM; Levanon, & Soker 2019a).

The positive side for the DD scenario of the requirement of $t_{\text{MED}} \gtrsim 10^3 \text{ s} \gg t_{\text{dyn}}$, where $t_{\text{dyn}} \simeq 10 \text{ s}$ is the dynamical time of the merger process, is that the merger product has time to acquire spherical structure that will lead to the required spherical explosion (section 2.1). The rarity of the early excess emission might suggest that in most cases the MED of the DD scenario is longer than tens of years to allow the DOM to disperse (Levanon et al. 2015), again, leading to spherical explosion.

The finding that early excess emission is rare (e.g., Fausnaugh et al. 2019), and the non-detection of hydrogen in the ejecta (e.g., Dimitriadis et al. 2019b), or only a very low mass of hydrogen in the ejecta (section 4), imply that if the SD scenario accounts for some SNe Ia, there must be a very long MED. Namely, it must be the SD-MED scenario.

★ **Summary of section 3.** Early excess emission can occur in principle both in the SD and in the DD scenarios. The new findings of early excess emission and their rarity suggest that in both scenarios the common channel is the one where there is a merger (or accretion) to explosion delay (MED) time. Namely, for normal SNe Ia the DD-MED and SD-MED scenarios dominate the DD and the SD scenarios, respectively.

4. HYDROGEN IN SNE IA

Most recent studies find no hydrogen in SNe Ia (e.g., Tucker et al. 2020). This by itself almost rules-out the SD scenario that has no MED. It leaves the SD-MED channel (spin-up/spin-down channel) for normal SNe Ia. But there is still the problem that population synthesis studies cannot attribute all SNe Ia to the SD scenario.

Even the recent detection of hydrogen in two SNe Ia is much below that expected in the SD scenario. Kollmeier et al. (2019) estimate the hydrogen mass in the sub-luminous SN Ia SN2018fhw/ASASSN-18tb to be $M_{\text{H}} \approx 2 \times 10^{-3} M_{\odot}$ (or maximum up to $M_{\text{H}} \approx 0.01 M_{\odot}$). To explain the $\text{H}\alpha$ line they consider stripped gas from the companion in the SD scenario, fluorescent UV pumping in a slowly expanding shell of material, and interaction with CSM (also Valley et al. 2019a). However, Valley et al. (2019a) note that SN2018fhw is very different from other known CSM-interacting SNe Ia.

Prieto et al. (2020) infer a hydrogen mass of $M_{\text{H}} \approx 10^{-3} M_{\odot}$ from their detection of $\text{H}\alpha$ emission line in the low-luminosity fast declining SN Ia SN2018cqj/ATLAS18qtd. Although some properties of the $\text{H}\alpha$ line are consistent with stripped hydrogen, the hydrogen mass is significantly less than predictions of theoretical calculations

of the SD scenario without MED (e.g., Botyánszki et al. 2018).

Another explanation for the presence of little hydrogen, although speculative, is the rare evaporation of a planet (or two) by SNe Ia (Soker 2019). The Jupiter-like planet should reside within $a \simeq 50 R_{\odot}$ from the SN. Interestingly, all SN Ia binary scenarios might in principle account for the presence of planets.

★ **Summary of section 4.** The general non-detection of hydrogen in SNe Ia, and the very low hydrogen mass of $M_{\text{H}} \approx 10^{-3} M_{\odot}$ in rare cases, rules out the SD scenario that has no long MED ($t_{\text{MED}} \gtrsim 10^7 \text{ yr}$; Soker 2018). The very low hydrogen mass in rare cases might come from a planet that the SN Ia evaporates.

5. DELAY TIMES AND THEIR IMPLICATIONS

The delay time distribution (DTD) refers to the distribution of SNe Ia with time from star formation to explosion, $t_{\text{SF-E}}$. A common approximation is a power-law (see Strolger et al. 2020 for an exponential form)

$$\dot{N}_{\text{DTD}} \equiv \left(\frac{dN_{\text{Ia}}}{dt} \right)_{\text{DTD}} = A \left(\frac{t}{1 \text{ Gyr}} \right)^{\alpha}. \quad (2)$$

In the past, many studies have obtained $\alpha \simeq -1$. Two significant recent papers find steeper dependence. Friedmann & Maoz (2018) derive $A = 5 - 8 \times 10^{-13} M_{\odot}^{-1} \text{ yr}^{-1}$ and $\alpha = -1.30^{+0.23}_{-0.16}$, and Heringer et al. (2019a) derive $A = 7 \pm 2 \times 10^{-13} M_{\odot}^{-1} \text{ yr}^{-1}$ and $\alpha = -1.34^{+0.19}_{-0.17}$, although other derivation exist (e.g., Frohmaier et al. 2019).

The time $t_{\text{SF-E}}$ can be composed from several times. For scenarios that involve two WDs there are in principle 3 times. These are the time from star formation to the formation of the two WDs in the post-CEE phase, $t_{\text{SF-CE}}$, the time for gravitational waves to merge the two WDs, t_{GW} , and the time from the merger of the two WDs to explosion, the MED time t_{MED} . Namely,

$$t_{\text{SF-E}}(\text{DD}) = t_{\text{SF-CE}} + t_{\text{CEED}} = t_{\text{SF-CE}} + t_{\text{GW}} + t_{\text{MED}}, \quad (3)$$

where $t_{\text{CEED}} = t_{\text{GW}} + t_{\text{MED}}$ is the time from the end of the CEE to explosion. In the DD and DD-MED scenario the longest time is t_{GW} . In the DD scenario and in the DDet scenario with a WD helium-donor, such as the D⁶ channel, $t_{\text{MED}} = 0$. In the CD scenario the merger takes place during the CEE and so $t_{\text{GW}} = 0$ and $t_{\text{CEED}} = t_{\text{MED}}$.

There are some SNe Ia that occur while there is a CSM around the explosion cite, so called SNe-CSM. Graham et al. (2019) estimate the fraction of SNe Ia-CSM with close CSM to be $f_{\text{CSM}} < 0.06$ of all SNe Ia. In addition, there are cases with CSM further away.

Over all, the presence of a CSM implies that in scenarios that involve a CEE the explosion occurs within a short time after the CE, $t_{\text{CEED}} \lesssim 10^6$ yr. Of course, in most cases explosions occur much later, e.g., as non-detection of radio emission reveals (e.g., Cendes et al. 2020; Lundqvist et al. 2020).

I used the new findings of Graham et al. (2019) and of Friedmann & Maoz (2018) and Heringer et al. (2019a), to study the relation of SNe Ia that occur at short times (within a million years) after the CEE, and those that take place long after the CEE. Based on these studies that show from observations this population of SNe Ia to be young (Graham et al. 2019), I concluded (Soker 2019) that the population of SNe Ia with $t_{\text{CEED}} \lesssim 10^6$ yr amounts to $\approx \text{few} \times 0.1$ of all SNe Ia. I further concluded that the expression for the rate of these SNe Ia is different from that of the DTD billions of years after star formation (equation 2). In a recent study Jerkstrand et al. (2020) suggest that the superluminous SN 2006gy exploded in the frame of the CD scenario shortly after the CEE. Their suggestion, if holds, supports my earlier conclusion on the importance of the SN Ia population with short t_{CEED} .

From these conclusions I suggested that the physical processes that determine the explosion time shortly after the CEE are different (at least to some extend) than the processes that determine the explosion times long after the CEE. I here argue that at least those that occur within a million years after the CEE come from the CD scenario. I do note thought that there are claims that the CD scenario contributes less than my estimate in the Table 1 (e.g., Wang et al. 2017).

★ **Summary of section 5.** New results from 2018-2019 strengthen the case for a population of SNe Ia that occur within a million years after the CEE of their progenitors. The other population is of SNe Ia that occur much later, over a timescale of $\approx 10^7$ yr and longer. The processes that determine the delay time from the CEE to explosion are different (at least somewhat) in these two populations. I suggest that the short-delay population comes mainly from the CD scenario.

6. PECULIAR SNE IA

Peculiar SNe Ia include SNe Iax, Ca-rich transients, SN 2002es-like SNe Ia, SN 1991bg-like SNe Ia, 2009dc-like SNe Ia-pec, and more (e.g., review by Jha et al. 2019). Chen et al. (2019a) study a luminous SNe Ia, and suggest to replace the name “Super-Chandrasekhar SN Ia” by “2009dc-like SN Ia-pec”, that I use here. In Table 1 I list my crude estimate of the contribution of different scenarios to peculiar SNe Ia. (Maeda, & Terada 2016 present a different type of table of some scenarios

and their properties in relation to normal and peculiar SNe Ia but that table does not include the CD scenario nor estimates of the fractional contribution of each scenario.)

In the summary of section 2 I noted that dynamical binary interactions that numerical simulations show to ignite one or both WDs (e.g., the violent merger, the D⁶, the hybrid HeCO channels of the DD scenario, and the WWC scenario), have highly non-spherical explosions. Although some studies show that the ignition in the DDet scenario might not be easy (e.g., Wu, & Wang 2019), I assume that ignition does take place (e.g., Gronow et al. 2020) but lead to highly asymmetrical explosions (section 2.2). For that, I find that these scenarios and channels are unlikely to account for a large fraction of normal SNe Ia. However, they might explain a large fraction of peculiar SNe Ia and other transients (e.g., like accretion-induced collapse, e.g., Wang 2018a), and the easy ignition in these processes suggests that peculiar SNe Ia (and similar transients) are very common, possibly even more common than normal SNe Ia. Not surprisingly, there are relatively many recent papers on peculiar SNe Ia and similar transients (e.g., Lyutikov, & Toonen 2019; Panther et al. 2019; Prentice et al. 2020).

The studies of peculiar SNe Ia from 2019 add to the growing evidence that peculiar SNe Ia come from several channels and scenarios, possibly from all scenarios listed in Table 1. Indeed, at least some peculiar SNe Ia show low polarisation as SNe Ia show (Meng et al. 2017).

The DDet scenario seems to be popular by studies of peculiar SNe Ia (e.g., Magee et al. 2019), and in particular Ca-rich transients (e.g., De et al. 2020). Polin et al. (2019b) study the DDet scenario and argue that even a small amount of calcium can make a Ca-rich transient. Jacobson-Galan et al. (2019b) argue that the Ca-rich SN 2016hmk is consistent with the DDet scenario. Takaro et al. (2020) estimate the delay time from star formation to explosion of SNe Iax, and conclude that they better fit the DDet scenario than massive WR progenitors. However, other scenarios exist for these types of peculiar SNe Ia. For example, the hybrid HeCO channel of the DD scenario for Ca-rich transients (Zenati et al. 2020).

Sand et al. (2019) find no H α emission in fast-declining SNe Ia, suggesting these do not come from the SD scenario. But not all peculiar SNe Ia are sub- M_{Ch} explosions. Galbany et al. (2019) argue that the peculiar SN 2016hmk, a Ca-strong 1991bg-like SN, comes from an M_{Ch} explosion. Jacobson-Galan et al. (2019a) spectroscopically model 44 SNe Iax. They find helium in two SNe Iax that better fit a CSM helium than a

helium in the ejecta. In the majority of SNe Iax they find no helium, but still suggest that the SD scenario with helium donor might be the main contribute of SNe Iax. Meng, & Podsiadlowski (2018) propose that SNe Iax that are SNe Ia-CSM result from the hybrid CONe common-envelope wind channel of the SD scenario.

Raddi et al. (2019) report the observation of three runaway stars, that are chemically peculiar due to enrichment of nuclear ashes of partial oxygen and silicon burning. They deduce their masses to be in the range of $0.2 - 0.28M_{\odot}$. They further speculate that these inflated WDs are the partly burnt remnants of either electron-capture supernovae or peculiar SNe Iax (e.g., Zhang et al. 2019). My view is that *any* thermonuclear explosion that leaves a runaway companion, e.g., simulations by Bauer et al. (2019) of the DDet (D^6) scenario, belongs to peculiar SNe Ia (and other transients), such as SNe Iax.

★ **Summary of section 6.** The relative easy ignition of WDs by dynamical effects in numerical simulations of some scenarios and channels (e.g., accretion at a high rate or collision), as evidence also from studies from 2019, suggests that these explosions are common. However, since they are highly globally non-spherical, they cannot explain many normal SNe Ia (section 2). These non-spherical explosions, I suggest, make the majority of peculiar SNe Ia, as I summarise in Table 1. I expect, therefore, that most peculiar SNe Ia have non-spherical explosions, unlike normal SNe Ia.

7. SUMMARY

The question of the main progenitors of normal SNe Ia is unsolved yet. Not only there is no consensus on the most promising SN Ia scenario(s), but there is no consensus even on the different potential scenarios. The correct counting as of the end of 2019 is of five binary scenarios: the CD, the DD, the DDet, the SD and the WWC scenario (section 1). I list them in Table 1, where I also split the DD scenario to a channel without a merger to explosion delay (MED) time, and channel with a MED, the DD-MED channel (scenario). (It might be that we need to consider the DD channel and the DD-MED channel of the DD scenarios as two separate scenarios.)

According to population synthesis studies there are not enough double WD systems with combined mass of $\gtrsim M_{\text{Ch}}$ (e.g., Cheng et al. 2020; I note that Rebassa-Mansergas et al. 2019 argue that observationally it is not easy to detect WD binary systems). Therefore, the DD-MED scenario cannot account for all SNe Ia.

The strongest point of sub- M_{Ch} scenarios (DD, DDet, WWC scenarios) is that they can achieve easy ignition by dynamical processes. However, they lead to highly non-spherical explosions, in contradiction with observations (Table 1). On the other hand, the main challenge of the M_{Ch} scenarios (CD, DD-Med and SD scenarios) is to achieve ignition. Recent studies (e.g., Fisher et al. 2019; Poludnenko et al. 2019) show that turbulence can drive detonation (deflagration-to-detonation) in WDs, and by that turbulence increases the range of conditions for the onset of carbon detonation. This result helps the M_{Ch} explosions. On the observational side, Kawabata et al. (2019) study the high-velocity SNe Ia SN 2019ein, and find its properties to be compatible with deflagration to detonation M_{Ch} explosion. However, there are still open issues and observations that the M_{Ch} model cannot explain (yet), e.g., Byrohl et al. (2019).

The entire set of observations brings many researchers to the conclusion that SNe Ia come both from M_{Ch} explosions and from sub- M_{Ch} explosions (e.g., Levanon, & Soker 2019b; Polin et al. 2019a,b; Sarbadhicary et al. 2019; Seitenzahl et al. 2019). The new study by Seitenzahl et al. (2019) which presents a new approach to study SNRs is a good example for that. By spatially resolving some nebular emission lines, they effectively perform a "tomography", i.e., they reveal the location of the reverse shock. By using evolutionary model they could constrain the type of explosion in two SNRs. They conclude that SNR 0519-69.0 was a standard M_{Ch} explosion, while SNR 0509-67.5 was an energetic sub- M_{Ch} explosion. As another example, Ashall et al. (2019a) support M_{Ch} explosion for most SNe Ia, but note that the subluminal SN 2015bo might come from the merger of two WDs. They claim that this demonstrates the diversity of explosion scenarios of faint SNe Ia. Kobayashi et al. (2019) study galactic chemical evolution from which they conclude that sub- M_{Ch} SNe Ia contribute up to 25% of SNe Ia in the solar neighbourhood. In dwarf spheroidal galaxies sub- M_{Ch} SNe Ia contribute a higher fraction when star formation took place. Kirby et al. (2019) reach similar conclusions, i.e., that sub- M_{Ch} explosions dominated in dwarf galaxies when they formed stars in the past. At present, M_{Ch} SNe Ia might dominate, as they do in the Milky Way and in dwarf galaxies with extended star formation. These studies of chemical evolution, and the very recent study by de los Reyes et al. (2020), also point to both M_{Ch} and sub- M_{Ch} explosions.

★ **Summary of section 7.** The summary of this section, and in turn of the entire review, are the last two rows of Table 1 where I list my crude estimates of

the contribution of the different scenarios. This table presents my view that normal SNe Ia come both from M_{Ch} explosions, mainly the CD and DD-MED scenarios, and from sub- M_{Ch} explosions, mainly the DD scenario. There is a possible small contribution from the SD and DDet scenarios. Peculiar SNe Ia come mainly from the DD, DDet, and SD scenarios, all without MED.

ACKNOWLEDGEMENTS

I thank an anonymous referee for detailed and very useful comments. I thank Mattia Bulla, Keiichi Maeda, Paolo Mazzali, Michael Tucker, Ivo Seitenzahl, and Xiangcun Meng for useful comments. This research was supported by the Israel Science Foundation, by the Prof. Amnon Pazy Research Foundation, and by the E. and J. Bishop Research Fund at the Technion. I presented an earlier version of this review at the Progenitors of Type Ia Supernovae meeting, Lijiang, China, August 05-09, 2019.

REFERENCES

- Ablimit, I., & Maeda, K. 2019a, *ApJ*, 871, 31
- Ablimit, I., & Maeda, K. 2019b, *ApJ*, 885, 99
- Alsaberi, R. Z. E., Barnes, L. A., Filipovic, M. D., et al. 2019, *Ap&SS*, 364, 204
- Antoniadis, J., Chanlaridis, S., Gräfenr, G., & Langer, N. 2020, *A&A*, 635, A72
- Ashall, C., Hoefflich, P., Hsiao, E. Y., et al. 2019a, *ApJ*, 878, 86
- Ashall, C., Hsiao, E. Y., Hoefflich, P., et al. 2019b, *ApJL*, 875, L14
- Augustine, C. N., Willcox, D. E., Brooks, J., Townsley, D. M., & Calder, A. C. 2019, *ApJ*, 887, 188
- Bauer, E. B., White, C. J., & Bildsten, L. 2019, *ApJ*, 887, 68
- Bear, E., & Soker, N. 2018, *MNRAS*, 480, 3702
- Benvenuto, O. G., Panei, J. A., Nomoto, K., Kitamura, H., & Hachisu, I. 2015, *ApJL*, 809, L6
- Borkowski, K. J., Reynolds, S. P., Hwang, U., Green D. A., Petre R., Krishnamurthy K., & Willett R., 2013, *ApJL*, 771, L9
- Boshkayev, K., Rueda, J. A., Ruffini, R., & Siutsou, I. 2014, *Journal of Korean Physical Society*, 65, 855
- Botyánszki, J., Kasen, D., & Plewa, T. 2018, *ApJL*, 852, L6
- Bravo, E. 2019, *A&A*, 624, A139
- Bravo, E., Badenes, C., & Martínez-Rodríguez, H. 2019, *MNRAS*, 482, 4346
- Brown, P. J., Hosseinzadeh, G., Jha, S. W., et al. 2019, *ApJ*, 877, 152
- Bulla, M., Miller, A. A., Yao, Y., et al. 2020, *arXiv:2001.00587*
- Bulla, M., Sim, S. A., Pakmor, R., et al. 2016a, *MNRAS*, 455, 1060
- Bulla, M., Sim, S. A., Kromer, M., et al. 2016b, *MNRAS*, 462, 1039
- Byrohl, C., Fisher, R., & Townsley, D. 2019, *ApJ*, 878, 67
- Calder, A. C., Willcox, D. E., DeGrendele, C. J., Shangase, D., Zingale, M., & Townsley, D. M. 2019, *Journal of Physics Conference Series*, 012002
- Cendes, Y., Drout, M. R., Chomiuk, L., & Sarbadhicary S. K., 2020, *arXiv:2001.03558*
- Chakradhari, N. K., Sahu, D. K., & Anupama, G. C. 2019, *MNRAS*, 487, 1886
- Chen, P., Dong, S., Katz, B., et al. 2019a, *ApJ*, 880, 35
- Chen, X., Hu, L., & Wang, L. 2019b, *arXiv:1911.05209*
- Cheng, S., Cummings, J. D., Ménard, B., & Toonen, S. 2020, *ApJ*, 891, 160
- Cikota, A., Patat, F., Cikota, S., Spyromilio, J., & Rau, G. 2017, *MNRAS*, 471, 2111
- Cikota, A., Patat, F., Wang, L., et al. 2019, *MNRAS*, 490, 578
- Clavelli, L. 2019, *arXiv:1908.11397*
- De, K., Kasliwal, M. M., Tzanidakis, A., et al. 2020, *arXiv e-prints*, *arXiv:2004.09029*
- de los Reyes, M. A. C., Kirby, E. N., Seitenzahl, I. R., & Shen, K. J. 2020, *ApJ*, 891, 85
- Dessart, L., Leonard, D., & Prieto, J.-L. 2020, *arXiv:2004.03986*
- Dickel, J. R., & Jones, E. M. 1985, *ApJ*, 288, 707
- Dimitriadis, G., Foley, R. J., Rest, A., et al. 2019a, *ApJL*, 870, L1
- Dimitriadis, G., Rojas-Bravo, C., Kilpatrick, C. D., et al. 2019b, *ApJL*, 870, L14
- Di Stefano, R. 2020, *MNRAS*, 491, 495
- Di Stefano, R., Voss, R., & Claeys, J. S. W. 2011, *ApJL*, 738, LL1
- Dwarkadas, V. V., & Chevalier, R. A. 1998, *ApJ*, 497, 807.
- Fang, J., Yan, J., Wen, L., Lu C., & Yu H., 2020, *MNRAS*, 491, 2460
- Fausnaugh, M. M., Vallely, P. J., Kochanek, C. S., et al. 2019, *arXiv:1904.02171*
- Ferrand, G., Warren, D. C., Ono, M., Nagataki, S., Röpke, F. K., & Seitenzahl, I. R., 2019, *ApJ*, 877, 136

- Fink, M., Röpke, F. K., Hillebrandt, W., Seitenzahl, I. R., Sim, S. A., & Kromer, M., 2010, *A&A*, 514, A53
- Fisher, R., Mozumdar, P., & Casabona, G. 2019, *ApJ*, 876, 64
- Friedmann, M., & Maoz, D. 2018, *MNRAS*, 479, 3563
- Frohmaier, C., Sullivan, M., Nugent, P. E., et al. 2019, *MNRAS*, 486, 2308.
- Galbany, L., Ashall, C., Höflich, P., et al. 2019, *A&A*, 630, A76
- Graham, M. L., Harris, C. E., Nugent, P. E., et al. 2019, *ApJ*, 871, 62
- Graur, O. 2019, *ApJ*, 870, 14
- Graur, O., Maguire, K., Ryan, R., et al. 2019, *Nature Astronomy*, 463
- Graur, O., & Woods, T. E. 2019, *MNRAS*, 484, L79
- Gronow, S., Collins, C., Ohlmann, S. T., et al. 2020, *A&A*, 635, A169
- Hallakoun, N., & Maoz, D. 2019, *MNRAS*, 490, 657
- Haim, N., & Katz, B. 2018, *MNRAS*, 479, 3155
- Hamers, A. S. & Thompson, T. A. 2019, *ApJ*, 882, 24
- Han, X., Zheng, W., Stahl, B. E., et al. 2020, *arXiv:1911.07734*
- Heringer, E., Pritchet, C., & van Kerkwijk, M. H. 2019a, *ApJ*, 882, 52
- Heringer, E., van Kerkwijk, M. H., Sim, S. A., Kerzendorf, W. E., & Graham, M. L. 2019b, *ApJ*, 871, 250
- Holmbo, S., Stritzinger, M. D., Shappee, B. J., et al. 2019, *A&A*, 627, A174
- Jacobson-Galán, W. V., Foley, R. J., Schwab, J., et al. 2019a, *MNRAS*, 487, 2538
- Jacobson-Galan, W. V., Polin, A., Foley, R. J., et al. 2019b, *arXiv:1910.05436*
- Jerkstrand, A., Maeda, K., & Kawabata, K. S. 2020, *Science*, 367, 415
- Jha, S. W., Maguire, K., & Sullivan, M. 2019, *Nature Astronomy*, 3, 706
- Jiang, J.-A., Doi, M., Maeda, K., et al. 2018, *ApJ*, 865, 149
- Justham, S. 2011, *ApJL*, 730, LL34
- Kawabata, M., Maeda, K., Yamanaka, M., et al. 2019, *arXiv:1908.03001*
- Kasen, D. 2010, *ApJ*, 708, 1025
- Kirby, E. N., Xie, J. L., Guo, R., et al. 2019, *ApJ*, 881, 45
- Kobayashi, C., Leung, S.-C., & Nomoto, K. 2019, *arXiv:1906.09980*
- Kollmeier, J. A., Chen, P., Dong, S., et al. 2019, *MNRAS*, 486, 3041
- Konyves-Toth, R., Vinko, J., Ordasi, A., et al. 2019, *arXiv:1908.00582*
- Kushnir, D., & Katz, B. 2020a, *MNRAS*, 491, 3760
- Kushnir, D., & Katz, B. 2020b, *MNRAS*, 493, 5413
- Kushnir, D., Katz, B., Dong, S., Livne, E., & Fernandez, R. 2013, *ApJL*, 778, L37
- Kuuttila, J., Gilfanov, M., Seitenzahl, I. R., Woods T. E., & Vogt F. P. A. 2019, *MNRAS*, 484, 1317
- Leung, S.-C., & Nomoto, K. 2020, *ApJ*, 888, 80
- Levanon, N., & Soker, N. 2019a, *ApJ*, 872, L7.
- Levanon, N., & Soker, N. 2019b, *MNRAS*, 486, 5528
- Levanon, N., Soker, N., & García-Berro, E. 2015, *MNRAS*, 447, 2803
- Li, C.-J., Kerzendorf, W. E., Chu, Y.-H., et al. 2019, *ApJ*, 886, 99
- Li, W., Wang, X., Vinkó, J., et al. 2019, *ApJ*, 870, 12
- Liu, Z.-W., & Stancliffe, R. J. 2018, *MNRAS*, 475, 5257
- Liu, D., Wang, B., Ge, H., Chen X., & Han Z., 2019, *A&A*, 622, A35
- Livio, M., & Mazzali, P. 2018, *Physics Reports*, 736, 1
- Livneh, R., & Katz, B. 2020, *arXiv:1912.04313*
- Luken, K. J., Filipović, M. D., Maxted, N. I., et al. 2020, *MNRAS*, 492, 2606
- Lundqvist, P., Kundu, E., Perez-Torres, M. A., et al. 2020, *ApJ*, 890, 159
- Lyutikov, M., & Toonen, S. 2019, *MNRAS*, 487, 5618
- Maeda, K., & Terada, Y. 2016, *International Journal of Modern Physics D*, 25, 1630024
- Magee, M. R., Maguire, K., Kotak, R., Sim, S. A., Gillanders, J. H., Prentice, S. J., & Skillen, K., 2020, *A&A*, 634, A37
- Magee, M. R., Sim, S. A., Kotak, R., Maguire, K., & Boyle, A. 2019, *A&A*, 622, A102
- Maguire, K., Sim, S. A., Shingles, L., et al. 2018, *MNRAS*, 477, 3567
- Maoz, D., Mannucci, F., & Nelemans, G. 2014, *ARA&A*, 52, 107
- Mazzali, P. A., Bikmaev, I., Sunyaev, R., et al. 2020, *MNRAS*, doi:10.1093/mnras/staa839
- Meng, X. 2019, *ApJ*, 886, 58
- Meng, X., & Li, J. 2019, *MNRAS*, 482, 5651
- Meng, X., & Podsiadlowski, P. 2017, *MNRAS*, 469, 4763
- Meng, X., & Podsiadlowski, P. 2018, *ApJ*, 861, 127
- Meng, X., Zhang, J., & Han, Z. 2017, *ApJ*, 841, 62
- Miles, B. J., Townsley, D. M., Shen, K. J., Timmes, F. X., & Moore, K. 2019, *ApJ*, 871, 154
- Millard, M. J., Bhalerao, J., Park, S., et al. 2019, *arXiv:1905.04475*
- Miller, A. A., Yao, Y. Bulla, M., et al. 2020, *arXiv:2001.00598*
- Moriya, T. J., Liu, D., Wang, B., & Liu Z.-W., 2019, *MNRAS*, 488, 3949
- Neunteufel, P., Yoon, S.-C., & Langer, N. 2019, *A&A*, 627, A14

- Pakmor, R., Kromer, M., Taubenberger, S., & Springel, V. 2013, *ApJL*, 770, L8
- Pan, Y.-C., Foley, R. J., Jones, D. O., Filippenko A. V., & Kuin N. P. M. 2020, *MNRAS*, 491, 5897
- Panther, F. H., Seitzzahl, I. R., Ruiter, A. J., et al. 2019, *PASA*, 36, e031
- Pellegrino, C., Howell, D. A., Sarbadhicary, S. K., et al. 2020, *arXiv:2003.05946*
- Perets, H. B., Zenati, Y., Toonen, S., & Bobrick, A. 2020, *arXiv:1910.07532*
- Piersanti, L., Gagliardi, S., Iben, I., Jr., & Tornambé, A. 2003, *ApJ*, 583, 885
- Polin, A., Nugent, P., & Kasen, D. 2019a, *ApJ*, 873, 84
- Polin, A., Nugent, P., & Kasen, D. 2019b, *arXiv:1910.12434*
- Poludnenko, A., Chambers, J., Ahmed, K., Gamezo V., & Taylor B. 2019, *Science*, 2019, 366
- Prentice, S. J., Maguire, K., Flörs, A., et al. 2020, *A&A*, 635, A186
- Prieto, J. L., Chen, P., Dong, S., et al. 2020, *ApJ*, 889, 100
- Raddi, R., Hollands, M. A., Koester, D., et al. 2019, *MNRAS*, 489, 1489
- Rebassa-Mansergas, A., Toonen, S., Korol, V., & Torres, S. 2019, *MNRAS*, 482, 3656
- Ruiter, A. J. 2020, *arXiv:2001.02947*
- Ruiz-Lapuente, P. 2019, *NewAR*, 85, 101523
- Sand, D. J., Amaro, R. C., Moe, M., et al. 2019, *ApJL*, 877, L4
- Sarbadhicary, S. K., Chomiuk, L., Badenes, C., Tremou E., Soderberg A. M., & Sjouwerman L. O. 2019, *ApJ*, 872, 191
- Sato, T., Bravo, E., Badenes, C., Hughes, J. P., Williams, B. J., Yamaguchi, H., 2020, *ApJ*, 890, 104
- Sato, T., Hughes, J. P., Williams, B. J., et al. 2019, *ApJ*, 879, 64
- Seitzzahl, I. R., Ghavamian, P., Laming, J. M., & Vogt F. P. A., 2019, *PhRvL*, 123, 041101
- Shappee, B. J., Holoien, T. W.-S., Drout, M. R., et al. 2019, *ApJ*, 870, 13
- Shen, K. J., Boubert, D., Gänsicke, B. T., et al. 2018, *ApJ*, 865, 15
- Shingles, L. J., Sim, S. A., Kromer, M., et al. 2020, *MNRAS*, 492, 2029
- Siebert, M. R., Foley, R. J., Jones, D. O., et al. 2019, *MNRAS*, 486, 5785
- Soker, N. 2018, *Science China Physics, Mechanics, and Astronomy*, 61, 49502
- Soker, N. 2019, *MNRAS*, 490, 2430
- Soker, N. 2019, *Research Notes of the American Astronomical Society*, 3, 153
- Soker, N., García-Berro, E., & Althaus, L. G. 2014, *MNRAS*, 437, L66
- Strolger, L.-G., Rodney, S. A., Pacifici, C., Narayan, G., & Graur O. 2020, *ApJ*, 890, 140
- Takaro, T., Foley, R. J., McCully, C., et al. 2020, *MNRAS*, 493, 986
- Tanikawa, A., Nomoto, K., & Nakasato, N. 2018, *ApJ*, 868, 90
- Tanikawa A., Nomoto K., Nakasato N., Maeda K., 2019, *ApJ*, 885, 103
- Taubenberger, S., Floers, A., Vogl, C., et al. 2019, *MNRAS*, 488, 5473
- Toonen, S., Perets, H. B., & Hamers, A. S. 2018, *A&A*, 610, A22
- Townsley, D. M., Miles, B. J., Shen, K. J., & Kasen, D. 2019, *ApJL*, 878, L38
- Tsebrenko, D., & Soker, N. 2013, *MNRAS*, 435, 320
- Tsebrenko, D., & Soker, N. 2015a, *MNRAS*, 447, 2568
- Tsebrenko, D., & Soker, N. 2015b, *MNRAS*, 453, 166
- Tucker, M. A., Shappee, B. J., Valley, P. J., et al. 2020, *MNRAS*, 493, 1044
- Tutukov, A. V., & Fedorova, A. V. 2007, *Astronomy Reports*, 51, 291
- Valley, P. J., Fausnaugh, M., Jha, S. W., et al. 2019a, *MNRAS*, 487, 2372
- Valley, P. J., Tucker, M. A., Shappee, B. J., Brown J. S., Stanek K. Z., & Kochanek C. S. 2020, *MNRAS*, 492, 3553
- Wang, B. 2018a, *MNRAS*, 481, 439
- Wang, B. 2018b, *RAA*, 18, 49
- Wang, B., Justham, S., Liu, Z.-W., Zhang, J.-J., Liu, D.-D., & Han, Z. 2014, *MNRAS*, 445, 2340
- Wang, B., Zhou, W.-H., Zuo, Z.-Y., et al. 2017, *MNRAS*, 464, 3965
- Wu, C., & Wang, B. 2019, *MNRAS*, 486, 2977
- Wu, C., Wang, B., & Liu, D. 2019, *MNRAS*, 483, 263
- Wygoda, N., Elbaz, Y., & Katz, B. 2019a, *MNRAS*, 484, 3941
- Wygoda, N., Elbaz, Y., & Katz, B. 2019b, *MNRAS*, 484, 3951
- Yang, Y., Hoefflich, P. A., Baade, D., et al. 2019, *arXiv:1903.10820*
- Yao, Y., Miller, A. A., Kulkarni, S. R., et al. 2019, *ApJ*, 886, 152
- Zenati, Y., Perets, H. B., & Toonen, S. 2020, Talk given at the Progenitors of Type Ia Supernovae meeting, Lijiang, China, August 05-09, 2019.
- Zenati, Y., Toonen, S., & Perets, H. B. 2019, *MNRAS*, 482, 1135
- Zhang, M., Fuller, J., Schwab, J., & Foley R. J., 2019, *ApJ*, 872, 29