

# Birational geometry for the covering of a nilpotent orbit closure II

Yoshinori Namikawa

## Abstract

Let  $O$  be a nilpotent orbit of a complex semisimple Lie algebra  $\mathfrak{g}$  and let  $\pi : X \rightarrow \bar{O}$  be the finite covering associated with the universal covering of  $O$ . In the previous article [Part 1] we have explicitly constructed a  $\mathbf{Q}$ -factorial terminalization  $\tilde{X}$  of  $X$  when  $\mathfrak{g}$  is classical. In this article we count how many non-isomorphic  $\mathbf{Q}$ -factorial terminalizations  $X$  has. We construct the universal Poisson deformation of  $\tilde{X}$  over  $H^2(\tilde{X}, \mathbf{C})$  and look at the action of the Weyl group  $W(X)$  on  $H^2(\tilde{X}, \mathbf{C})$ . The main result is an explicit geometric description of  $W(X)$ .

MSC2020: 14E15, 17B08

## Introduction

Let  $O$  be a nilpotent orbit of a complex semisimple Lie algebra  $\mathfrak{g}$  and let  $\pi_0 : X^0 \rightarrow O$  be an étale covering. Then  $\pi_0$  extends to a finite covering  $\pi : X \rightarrow \bar{O}$ . A birational projective morphism  $f : \tilde{X} \rightarrow X$  is called a  $\mathbf{Q}$ -factorial terminalization if  $\tilde{X}$  has only  $\mathbf{Q}$ -factorial terminal singularities and  $f$  is crepant (i.e.  $K_{\tilde{X}} = f^*K_X$ , where  $K_X$  and  $K_{\tilde{X}}$  are respectively the canonical divisors of  $X$  and  $\tilde{X}$ ). Two  $\mathbf{Q}$ -factorial terminalizations  $f_1 : \tilde{X}_1 \rightarrow X$  and  $f_2 : \tilde{X}_2 \rightarrow X$  are called *isomorphic* if there is an isomorphism  $\tau : \tilde{X}_1 \rightarrow \tilde{X}_2$  such that  $f_1 = f_2 \circ \tau$ . In the present article we count how many non-isomorphic  $\mathbf{Q}$ -factorial terminalizations  $X$  has. The main idea is to construct the universal Poisson deformation  $\tilde{\mathcal{X}} \rightarrow H^2(\tilde{X}, \mathbf{C})$  of  $\tilde{X}$  and look at the action of the Weyl group  $W(X)$  on the base space  $H^2(\tilde{X}, \mathbf{C})$ . For the definition and the properties of the Weyl group  $W(X)$ , see [Na 2], §1 and [Na 4]. In fact,  $H^2(\tilde{X}, \mathbf{R})$  is divided into a finite number of chambers, and the movable cone  $\text{Mov}(\tilde{X})$  is a union of some chambers. The Weyl group  $W(X)$  acts on  $H^2(\tilde{X}, \mathbf{R})$  and  $\text{Mov}(\tilde{X})$  is a fundamental domain for this  $W(X)$ -action. Each chamber inside  $\text{Mov}(\tilde{X})$  corresponds to a  $\mathbf{Q}$ -factorial terminalization of  $X$ . Hence

$$\#\{\mathbf{Q}\text{-factorial terminalizations of } X\} = \#\{\text{chambers}\}/|W(X)|.$$

Let us recall how we constructed a  $\mathbf{Q}$ -factorial terminalization  $\tilde{X}$  of  $X$ . Let  $G$  be a complex semisimple Lie group with  $\text{Lie}(G) = \mathfrak{g}$  and  $\pi_1(G) = \{1\}$ . Let  $Q$  be a parabolic subgroup of  $G$ . Fix a maximal torus  $T$  of  $Q$ . Then we have a Levi decomposition  $Q = U \cdot L$ . Here  $U$  is the unipotent radical and  $L$  is the Levi subgroup containing  $T$ . Let  $\mathfrak{n}$  be the nilradical of  $\mathfrak{q}$ . Then  $\mathfrak{n}$  coincides with the Lie algebra of  $U$ . Let  $\mathfrak{l}$  be the Lie algebra of  $L$ . Then we have a direct sum decomposition  $\mathfrak{q} = \mathfrak{n} \oplus \mathfrak{l}$ . Let  $O'$  be a nilpotent orbit of  $\mathfrak{l}$ . Then there is a unique nilpotent orbit  $O$  of  $\mathfrak{g}$  such that  $O$  meets  $\mathfrak{n} + O'$  in a non-empty Zariski open set of  $\mathfrak{n} + O'$ . In such a case we say that  $O$  is induced from  $O'$  and write  $O = \text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(O')$ . There is a generically finite map

$$\mu : G \times^Q (\mathfrak{n} + \bar{O}') \rightarrow \bar{O} \quad ([g, z] \mapsto \text{Ad}_g(z)),$$

which we call a generalized Springer map. Assume that an  $L$ -equivariant etale cover  $(X')^0 \rightarrow O'$  is given. Let  $X' \rightarrow \bar{O}'$  be the associated finite covering. Then the  $Q$ -action on  $\mathfrak{n} + \bar{O}'$  lifts to a  $Q$ -action on  $\mathfrak{n} \times X'$ . Hence we can make  $G \times^Q (\mathfrak{n} \times X')$  and get a commutative diagram

$$\begin{array}{ccc} G \times^Q (\mathfrak{n} \times X') & \xrightarrow{\mu'} & Z \\ \pi' \downarrow & & \downarrow \\ G \times^Q (\mathfrak{n} + \bar{O}') & \xrightarrow{\mu} & \bar{O}, \end{array} \quad (1)$$

where  $Z$  is the Stein factorization of  $\mu \circ \pi'$ . When  $Z = X$ , we say that the  $G$ -cover  $X^0$  of  $O$  is birationally induced from an  $L$ -cover  $(X')^0$  of  $O'$ . When  $X^0$  is not birationally induced from any other one except itself,  $X^0$  is called a birationally rigid cover. It is proved in [Ma, Corollary 4.3], [LMM, §7] that, for any  $G$ -cover  $X^0$  of  $O$ , there exists a unique  $(L, O', (X')^0)$  such that  $\mu'$  is birational and  $(X')^0$  is a birationally rigid cover. Moreover, in such a case,  $X'$  has only  $\mathbf{Q}$ -factorial terminal singularities and  $\mu'$  gives a  $\mathbf{Q}$ -factorial terminalization of  $X$ . In [Part 1] we have given an explicit algorithm for finding such  $(L, O', (X')^0)$  when  $\mathfrak{g}$  is a classical simple Lie algebra and  $X^0$  is the universal covering of  $O$ .

Therefore, in this article, we start with the situation when a commutative diagram

$$\begin{array}{ccc} G \times^Q (\mathfrak{n} \times X') & \xrightarrow{\mu'} & X \\ \pi' \downarrow & & \downarrow \\ G \times^Q (\mathfrak{n} + \bar{O}') & \xrightarrow{\mu} & \bar{O}, \end{array} \quad (2)$$

is already given and  $\mu'$  is a  $\mathbf{Q}$ -factorial terminalization of  $X$ . Put

$$N_G(L, T) := \{g \in G \mid gTg^{-1} = T, gLg^{-1} = L\}.$$

Then  $N_G(L, T)$  acts on  $\mathfrak{l}$ . We are mainly interested in the case when  $X^0$  is the universal covering of  $O$ . So we assume the following three conditions

(0) If we write  $O'$  as  $L/L^z$  for  $z \in O'$ , then  $(X')^0 = L/(L^z)^0$ , where  $(L^z)^0$  is the identity component of  $L^z$ .

(1)  $X'$  has only  $\mathbf{Q}$ -factorial terminal singularities.

(2)  $N_G(L, T)$  stabilizes  $O'$ .

In fact, (0) is always satisfied when  $X^0$  is the universal covering of  $O$ . Moreover, (2) is satisfied for all  $(L, O', X')$  constructed in [Part 1].

Let  $\mathfrak{k}$  be the center of  $\mathfrak{l}$ . Then the solvable radical  $\mathfrak{r}$  of  $\mathfrak{q}$  is written as

$$\mathfrak{r} = \mathfrak{k} \oplus \mathfrak{n}.$$

The adjoint  $Q$ -action stabilizes  $\mathfrak{r} + \bar{O}' \subset \mathfrak{r} \oplus \mathfrak{l}$  and one can make a space  $G \times^Q (\mathfrak{r} + \bar{O}')$ . Write  $x \in \mathfrak{r} + \bar{O}'$  as  $x = x_1 + x_2 + x_3$ , where  $x_1 \in \mathfrak{k}$ ,  $x_2 \in \mathfrak{n}$  and  $x_3 \in \bar{O}'$ . Define a map

$$\eta : G \times^Q (\mathfrak{r} + \bar{O}') \rightarrow \mathfrak{k}$$

by  $[g, x] \in G \times^Q (\mathfrak{r} + \bar{O}') \mapsto x_1$ . Since  $G \times^Q (\mathfrak{r} \times X')$  is a finite cover of  $G \times^Q (\mathfrak{r} + \bar{O}')$ , we have a map  $G \times^Q (\mathfrak{r} \times X') \rightarrow \mathfrak{k}$ . This map is a flat map with the central fiber  $G \times^Q (\mathfrak{n} \times X')$ .

Moreover, it is actually the universal Poisson deformation of  $G \times^Q (\mathfrak{n} \times X')$  and  $\mathfrak{k}$  is identified with  $H^2(G \times^Q (\mathfrak{n} \times X'), \mathbf{C})$ . We put

$$Z_{\mathfrak{l}, X'} := \text{Spec } \Gamma(G \times^Q (\mathfrak{r} \times X'), \mathcal{O}).$$

As the subscript indicates,  $Z_{\mathfrak{l}, X'}$  only depends on the Levi part  $\mathfrak{l}$  and  $X'$ . By definition, there is a map  $Z_{\mathfrak{l}, X'} \rightarrow \mathfrak{k}$  whose central fiber is  $X$ . By the general theory of the Poisson deformation of a conical symplectic variety, the Weyl group  $W(X)$  for  $X$  acts on both  $\mathfrak{k}$  and  $Z_{\mathfrak{l}, X'}$  so that the induced map  $Z_{\mathfrak{l}, X'}/W(X) \rightarrow \mathfrak{k}/W(X)$  gives the universal Poisson deformation of  $X$ :

$$\begin{array}{ccc} G \times^Q (\mathfrak{r} \times X') & \longrightarrow & Z_{\mathfrak{l}, X'}/W(X) \\ \downarrow & & \downarrow \\ \mathfrak{k} & \longrightarrow & \mathfrak{k}/W(X) \end{array} \quad (3)$$

The main result of this article is an explicit description of the  $W(X)$  (Theorem 9). In [Na 1] we already observed that a certain finite group  $W'$  acts on  $\mathfrak{k}$ . The group  $W'$  is defined as  $W' := N_G(L, T)/N_L(T)$ , where

$$N_L(T) := \{g \in L \mid gTg^{-1} = T\}.$$

Here we put

$$Y_{\mathfrak{l}, O'} := \text{Spec } \Gamma(G \times^Q (\mathfrak{r} + O'), \mathcal{O}).$$

The map  $Z_{\mathfrak{l}, X'} \rightarrow \mathfrak{k}$  factorizes as

$$Z_{\mathfrak{l}, X'} \rightarrow Y_{\mathfrak{l}, O'} \rightarrow \mathfrak{k}.$$

Then  $\text{Aut}(Z_{\mathfrak{l}, X'}/Y_{\mathfrak{l}, O'})$  is isomorphic to  $\text{Aut}((X')^0/O')$ . The group  $W'$  acts on  $Y_{\mathfrak{l}, O'}$ , but does not act on  $Z_{\mathfrak{l}, X'}$  in general. Instead of  $W'$  we consider a new group  $\tilde{W}'$  obtained as an extension of  $W'$  by  $\text{Aut}((X')^0/O')$ :

$$1 \rightarrow \text{Aut}((X')^0/O') \rightarrow \tilde{W}' \rightarrow W' \rightarrow 1$$

Then  $\tilde{W}'$  acts on  $Z_{\mathfrak{l}, X'}$  in such a way that the map  $Z_{\mathfrak{l}, X'} \rightarrow \mathfrak{k}$  is  $\tilde{W}'$ -equivariant. Since  $W'$  fixes the origin  $0 \in \mathfrak{k}$ , an element of  $\tilde{W}'$  acts on the central fiber  $X$  of  $Z_{\mathfrak{l}, X'} \rightarrow \mathfrak{k}$  as an element of  $\text{Aut}(X/\bar{O})$ . In this way we have a homomorphism

$$\rho_X : \tilde{W}' \rightarrow \text{Aut}(X/\bar{O}).$$

We then define

$$W_X := \text{Ker}(\rho_X).$$

This  $W_X$  equals the Weyl group  $W(X)$  (Theorem 9). We remark that Losev [Lo], Proposition 4.6 has given a similar description of  $W(X)$  when  $X'$  is the normalization of  $\bar{O}'$ . In this case we do not need to take  $\tilde{W}'$ .

In §2 we first show that  $\rho_X$  is surjective. This is enough to calculate  $|W_X|$  and the number of non-isomorphic  $\mathbf{Q}$ -factorial terminalizations of  $X$  (cf. Theorem 14). But we need more information on  $\rho_X$  to know the structure of  $W_X$ . When  $\mathfrak{g}$  is a classical simple

Lie algebra, we study  $\rho_X(\tilde{w})$  for an element  $\tilde{w} \in \tilde{W}'$  such that its image  $w \in W'$  is a reflection. In §3, we actually count the numbers of  $\mathbf{Q}$ -factorial terminalizations of  $X$  in two examples as an application of the theory developed in the previous sections.

*Acknowledgement.* The author thanks Ivan Losev for letting him know Proposition 4.7 in [Lo] after the first version of this paper (arXiv:1912.01729) was written.

## § 1. Universal Poisson deformations of $\mathbf{Q}$ -factorial terminalizations and Weyl groups

Let  $G$  be a complex semisimple Lie group and let  $Q$  be a parabolic subgroup of  $G$ . We take a maximal torus  $T$  of  $G$  so that  $T \subset Q$ . Let  $Q = U \cdot L$  be the Levi decomposition with  $T \subset L$ . Let  $\mathfrak{n}$  be the nilradical of  $\mathfrak{q}$ . Then  $\mathfrak{n}$  coincides with the Lie algebra of  $U$ . Let  $\mathfrak{l}$  be the Lie algebra of  $L$ . Then we have a direct sum decomposition  $\mathfrak{q} = \mathfrak{n} \oplus \mathfrak{l}$ . Let  $O' \subset \mathfrak{l}$  be a nilpotent orbit. Then we have a generalized Springer map

$$\mu : G \times^Q (\mathfrak{n} + \bar{O}') \rightarrow \mathfrak{g}, \quad [g, x] \mapsto \text{Ad}_g(x).$$

The image of  $\mu$  is the closure  $\bar{O}$  of a certain nilpotent orbit  $O$  of  $\mathfrak{g}$ . We write  $O'$  as  $L/L^z$  for  $z \in O'$ . Put  $(X')^0 := L/(L^z)^0$  and consider an etale covering  $(X')^0 \rightarrow O'$ . Let  $X' \rightarrow \bar{O}'$  be the associated finite covering. Then we have a finite covering  $\mathfrak{n} \times X' \rightarrow \mathfrak{n} \times \bar{O}' (= \mathfrak{n} + O')$ . The  $Q$ -action on  $\mathfrak{n} + \bar{O}'$  lifts to a  $Q$ -action on  $\mathfrak{n} \times X'$ . We assume

(1)  $X'$  has only  $\mathbf{Q}$ -factorial terminal singularities.

We can make  $G \times^Q (\mathfrak{n} \times X')$  and get a commutative diagram

$$\begin{array}{ccc} G \times^Q (\mathfrak{n} \times X') & \xrightarrow{\mu'} & X \\ \pi' \downarrow & & \pi \downarrow \\ G \times^Q (\mathfrak{n} + \bar{O}') & \xrightarrow{\mu} & \bar{O}, \end{array} \quad (4)$$

where  $X$  is the Stein factorization of  $\mu \circ \pi'$ .  $X$  is a conical symplectic variety and  $\mu'$  gives a  $\mathbf{Q}$ -factorial terminalization of  $X$ .

Let  $\mathfrak{k}$  be the center of  $\mathfrak{l}$ . We put  $\mathfrak{t} := \text{Lie}(T)$ . Then the root system  $\Phi$  is determined from  $(\mathfrak{g}, \mathfrak{t})$ . The quotient  $\mathfrak{l}_{\text{s.s.}} := \mathfrak{l}/\mathfrak{k}$  is a semisimple Lie algebra, and  $\mathfrak{t}/\mathfrak{k}$  is a Cartan subalgebra of  $\mathfrak{l}_{\text{s.s.}}$ . The root system  $\Phi_{\mathfrak{l}_{\text{s.s.}}}$  for  $(\mathfrak{l}_{\text{s.s.}}, \mathfrak{t}/\mathfrak{k})$  is a root subsystem of  $\Phi$ . We have

$$\mathfrak{k} = \{h \in \mathfrak{t} \mid \alpha(h) = 0, \forall \alpha \in \Phi_{\mathfrak{l}}\}.$$

We define

$$\mathfrak{k}^{\text{reg}} := \{h \in \mathfrak{k} \mid \alpha(h) \neq 0 \forall \alpha \in \Phi \setminus \Phi_{\mathfrak{l}_{\text{s.s.}}}\}.$$

The solvable radical  $\mathfrak{r}$  of  $\mathfrak{q}$  is written as

$$\mathfrak{r} = \mathfrak{k} \oplus \mathfrak{n}.$$

The adjoint  $Q$ -action stabilizes  $\mathfrak{r} + \bar{O}' \subset \mathfrak{r} \oplus \mathfrak{l}$  and one can make a space  $G \times^Q (\mathfrak{r} + \bar{O}')$ . Write  $x \in \mathfrak{r} + \bar{O}'$  as  $x = x_1 + x_2 + x_3$ , where  $x_1 \in \mathfrak{k}$ ,  $x_2 \in \mathfrak{n}$  and  $x_3 \in \bar{O}'$ . Define a map

$$\eta : G \times^Q (\mathfrak{r} + \bar{O}') \rightarrow \mathfrak{k}$$

by  $[g, x] \in G \times^Q (\mathfrak{r} + \bar{O}') \mapsto x_1$ . This is a well-defined map; in fact, for  $q \in Q$ , we have  $(Ad_q(x_1))_1 = x_1$ ,  $Ad_q(x_2) \in \mathfrak{n}$  and  $Ad_q(x_3) \in \mathfrak{n} + \bar{O}'$ . By definition  $\eta^{-1}(0) = G \times^Q (\mathfrak{n} + \bar{O}')$  and  $\eta$  gives a flat deformation of  $G \times^Q (\mathfrak{n} + \bar{O}')$  over  $\mathfrak{k}$ . Moreover, the  $Q$ -action on  $\mathfrak{r} + \bar{O}'$  lifts to a  $Q$ -action on  $\mathfrak{r} \times X'$ . Then  $G \times^Q (\mathfrak{r} \times X')$  is a finite covering of  $G \times^Q (\mathfrak{r} + \bar{O}')$ . We have a commutative diagram

$$\begin{array}{ccc}
G \times^Q (\mathfrak{n} \times X') & \longrightarrow & G \times^Q (\mathfrak{r} \times X') \\
\pi' \downarrow & & \Pi' \downarrow \\
G \times^Q (\mathfrak{n} + \bar{O}') & \longrightarrow & G \times^Q (\mathfrak{r} + \bar{O}') \\
\downarrow & & \eta \downarrow \\
\{0\} & \longrightarrow & \mathfrak{k},
\end{array} \tag{5}$$

where  $\eta \circ \Pi'$  gives a flat deformation of  $G \times^Q (\mathfrak{n} \times X')$  over  $\mathfrak{k}$ . Note that  $\mathfrak{k}$  is naturally identified with  $H^2(G \times^Q (\mathfrak{r} \times X'), \mathbf{C})$ . In fact,  $H^2(G \times^Q (\mathfrak{r} \times X'), \mathbf{C}) = \text{Pic}(G \times^Q (\mathfrak{r} \times X')) \otimes_{\mathbf{Z}} \mathbf{C}$  and we have isomorphisms

$$\text{Hom}_{alg.gp}(L, \mathbf{C}^*) \cong \text{Pic}(G/Q) \cong \text{Pic}(G \times^Q (\mathfrak{r} \times X'))$$

The isomorphism  $\mathfrak{t} \rightarrow \mathfrak{t}^*$  determined by the Killing form sends  $\mathfrak{k} \subset \mathfrak{t}$  isomorphically onto the subspace  $\text{Hom}_{alg.gp}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{C} \subset \mathfrak{t}^*$ .

Let us consider a fiber of  $\eta$ . For  $t \in \mathfrak{k}$ ,  $\eta^{-1}(t) = G \times^Q (t + \mathfrak{n} + \bar{O}')$ .

**Lemma 1.** (*[Na 3], Lemma (3.2)*) *For  $t \in \mathfrak{k}^{reg}$ , any orbit of the  $Q$ -variety  $t + \mathfrak{n} + \bar{O}'$  is of the form  $Q(t + y)$  with  $y \in \bar{O}'$ .*

Let  $W$  be the Weyl group of  $\mathfrak{g}$  with respect to  $\mathfrak{t}$ .  $W$  acts on  $\mathfrak{t}$ . Let  $\mathfrak{g}/G$  be the adjoint quotient of  $\mathfrak{g}$ . The map  $\mathfrak{t}/W \rightarrow \mathfrak{g}/G$  is an isomorphism by Chevalley's restriction theorem. We consider the adjoint quotient map  $\chi : \mathfrak{g} \rightarrow \mathfrak{g}/G \cong \mathfrak{t}/W$ . Let  $G \cdot (\mathfrak{r} + \bar{O}') \subset \mathfrak{g}$  be the space of all  $G$ -translates of the elements of  $\mathfrak{r} + \bar{O}'$ . We denote by the same  $\chi$  the restriction of  $\chi$  to  $G \cdot (\mathfrak{r} + \bar{O}')$ . By [Na 3], p. 21, the following diagram commutes

$$\begin{array}{ccc}
G \times^Q (\mathfrak{r} + \bar{O}') & \longrightarrow & G \cdot (\mathfrak{r} + \bar{O}') \\
\eta \downarrow & & \chi \downarrow \\
\mathfrak{k} & \xrightarrow{\iota} & \mathfrak{t}/W,
\end{array} \tag{6}$$

where the top horizontal map is defined by  $[g, x] \mapsto Ad_g(x)$  and  $\iota$  is the composite of the natural inclusion  $\mathfrak{k} \rightarrow \mathfrak{t}$  and the quotient map  $\mathfrak{t} \rightarrow \mathfrak{t}/W$ . By Lemma 1,

$$G \cdot (\mathfrak{r} + \bar{O}') = \overline{G \cdot (\mathfrak{k}^{reg} + \bar{O}')}.$$

Note that  $\mathfrak{k}$  only depends on the Levi part  $\mathfrak{l}$  of  $\mathfrak{q}$ . Hence  $G \cdot (\mathfrak{r} + \bar{O}')$  depends only on  $\mathfrak{l}$  and  $O'$ .

The commutative diagram above induces a commutative diagram

$$\begin{array}{ccc}
G \times^Q (\mathfrak{r} + \bar{O}') & \longrightarrow & \mathfrak{k} \times_{\mathfrak{t}/W} G \cdot (\mathfrak{r} + \bar{O}') \\
\eta \downarrow & & \downarrow \\
\mathfrak{k} & \xrightarrow{id} & \mathfrak{k}
\end{array} \tag{7}$$

We put  $N_G(T) := \{g \in G \mid gTg^{-1} = T\}$ ,  $N_G(L, T) := \{g \in N_G(T) \mid gLg^{-1} = L\}$  and  $N_L(T) := \{g \in L \mid gTg^{-1} = T\}$ . Then  $T \subset N_L(T) \subset N_G(L, T) \subset N_G(T)$ . The Weyl group  $W$  for  $G$  is defined as  $W := N_G(T)/T$  and the Weyl group  $W(L)$  for  $L$  is defined as  $W(L) := N_L(T)/T$ . We have inclusions

$$W(L) \subset N_G(L, T)/T \subset W.$$

Define

$$W' := N_G(L, T)/N_L(T) \tag{8}$$

Obviously  $W'$  is the quotient of  $N_G(L, T)/T$  by  $W(L)$ . Then  $W'$  acts on  $\mathfrak{k}$  and the map  $\iota : \mathfrak{k} \rightarrow \mathfrak{t}/W$  factorizes as  $\mathfrak{k} \rightarrow \mathfrak{k}/W' \rightarrow \mathfrak{t}/W$ . Here  $\mathfrak{k}/W'$  is the normalization of  $\iota(\mathfrak{k})$ . Let  $G \cdot (\mathfrak{r} + \bar{O}')^n$  be the normalization of  $G \cdot (\mathfrak{r} + \bar{O}')$ . Then the map  $G \cdot (\mathfrak{r} + \bar{O}')^n \rightarrow \mathfrak{t}/W$  factors through  $\mathfrak{k}/W'$ . Hence

$$(\mathfrak{k} \times_{\mathfrak{t}/W} G \cdot (\mathfrak{r} + \bar{O}'))^n = (\mathfrak{k} \times_{\mathfrak{k}/W'} G \cdot (\mathfrak{r} + \bar{O}')^n)^n.$$

We denote this variety by  $Y_{\mathfrak{l}, O'}$ . As the notation indicates,  $Y_{\mathfrak{l}, O'}$  depends only on  $\mathfrak{l}$  and  $O'$ . Let  $W'$  act on  $\mathfrak{k}$  in the usual way and act trivially on  $G \cdot (\mathfrak{r} + \bar{O}')^n$ . Then  $W'$  acts on  $\mathfrak{k} \times_{\mathfrak{k}/W'} G \cdot (\mathfrak{r} + \bar{O}')^n$ ; hence acts on its normalization  $Y_{\mathfrak{l}, O'}$ . Let  $\tilde{O}'$  be the normalization of  $\bar{O}'$ . Then the normalization of  $G \times^Q (\mathfrak{r} + \bar{O}')$  is  $G \times^Q (\mathfrak{r} \times \tilde{O}')$ . There is a natural map

$$G \times^Q (\mathfrak{r} \times \tilde{O}') \rightarrow Y_{\mathfrak{l}, O'}.$$

By definition  $N_G(L, T)$  acts on  $\mathfrak{l}$ . We assume that

(2)  $N_G(L, T)$  stabilizes  $O'$ .

Then the map is a birational projective morphism by [Na 3], Proposition 3.6. Here let us consider the composite  $G \times^Q (\mathfrak{r} \times X') \rightarrow G \times^Q (\mathfrak{r} \times \tilde{O}') \rightarrow Y_{\mathfrak{l}, O'}$  and take its Stein factorization  $Z_{\mathfrak{l}, X'}$ . Then we get a commutative diagram

$$\begin{array}{ccc}
G \times^Q (\mathfrak{r} \times X') & \xrightarrow{m'_Q} & Z_{\mathfrak{l}, X'} \\
\downarrow & & \downarrow \\
G \times^Q (\mathfrak{r} \times \tilde{O}') & \xrightarrow{m_Q} & Y_{\mathfrak{l}, O'}
\end{array} \tag{9}$$

We claim that  $Z_{\mathfrak{l}, X'}$  depends only on  $\mathfrak{l}$  and  $X'$  as the subscript indicates. More precisely, if we start with another parabolic subgroup  $Q'$  of  $G$  with Levi part  $L$  and with the same  $X' \rightarrow \bar{O}'$ , then we can make a similar diagram for  $G \times^{Q'} (\mathfrak{r}' \times X')$ . We claim that the same  $Z_{\mathfrak{l}, X'}$  is obtained as the Stein factorization. The claim follows from the next lemma. In fact, by the lemma,  $G \times^Q (\mathfrak{r} \times X')$  and  $G \times^{Q'} (\mathfrak{r}' \times X')$  have the same function field, and  $Z_{\mathfrak{l}, X'}$  is nothing but the normalization of  $Y_{\mathfrak{l}, O'}$  in this field.

**Lemma 2.** *Let  $Q$  and  $Q'$  be parabolic subgroups of  $G$  containing  $L$  as the Levi parts. Then there is a commutative diagram*

$$\begin{array}{ccc}
G \times^Q (\mathfrak{r} \times X') & \dashrightarrow & G \times^{Q'} (\mathfrak{r}' \times X') \\
\downarrow & & \downarrow \\
G \times^Q (\mathfrak{r} + \bar{O}') & \dashrightarrow & G \times^{Q'} (\mathfrak{r}' + \bar{O}') \\
\downarrow & & \downarrow \\
\mathfrak{k} \times_{\mathfrak{l}/W} G \cdot (\mathfrak{r} + \bar{O}') & = & \mathfrak{k} \times_{\mathfrak{l}/W} G \cdot (\mathfrak{r}' + \bar{O}') \\
\downarrow & & \downarrow \\
\mathfrak{k} & = & \mathfrak{k}
\end{array}$$

where the 1-st and the 2-nd horizontal maps are both birational maps.

*Proof.* The fiber of the map  $G \times^Q (\mathfrak{r} + \bar{O}') \rightarrow \mathfrak{k}$  over  $t \in \mathfrak{k}^{reg}$  is  $G \times^Q (t + \mathfrak{n} + \bar{O}')$ . By Lemma 1,

$$(*) \quad G \times^Q (t + \mathfrak{n} + \bar{O}') = G \times^Q (\cup_{y \in \bar{O}'} Q \cdot (t + y)) = \cup_{y \in \bar{O}'} G \times^Q Q \cdot (t + y)$$

Denote by  $Q_{t+y}$  the stabilizer group of  $t + y$ . Note that  $Q_{t+y} \subset L$ . In fact, assume that  $q \in Q_{t+y}$ . Put  $z = t + x$ . Since  $t \in \mathfrak{k}$  and  $y \in \mathfrak{l}$ , we have  $[t, y] = 0$ ; hence,  $z = t + y$  is the Jordan-Chevalley decomposition of  $z$ . Then  $Ad_q(z) = Ad_q(t) + Ad_q(y)$  is also the Jordan-Chevalley decomposition of  $z = Ad_q(z)$ . This means that  $Ad_q(t) = t$  and  $Ad_q(x) = x$ . In particular,  $q$  is contained in the centralizer  $Z(t)$  of  $t$  (in  $Q$ ).  $Z(t) = L$  because  $t \in \mathfrak{k}^{reg}$ . Therefore  $Q_{t+y} \subset L$ . We have

$$Q \cdot (t + y) = Q \times^{Q_{t+y}} (t + y) = Q \times^L L \times^{Q_{t+y}} (t + y) = Q \times^L L \cdot (t + y).$$

The right hand side of  $(*)$  can be written as

$$\cup_{y \in \bar{O}'} G \times^Q Q \cdot (t + y) = \cup_{y \in \bar{O}'} G \times^Q (Q \times^L L \cdot (t + y)) = \cup_{y \in \bar{O}'} G \times^L L \cdot (t + y) = G \times^L (t + \bar{O}').$$

Changing  $Q$  by  $Q'$ , we similarly get

$$G \times^{Q'} (t + \mathfrak{n}' + \bar{O}') = G \times^L (t + \bar{O}').$$

As a consequence we have

$$G \times^Q (t + \mathfrak{n} + \bar{O}') = G \times^{Q'} (t + \mathfrak{n}' + \bar{O}').$$

This means that the birational map  $G \times^Q (\mathfrak{r} + \bar{O}') \dashrightarrow G \times^{Q'} (\mathfrak{r}' + \bar{O}')$  is an isomorphism over any point of  $\mathfrak{k}^{reg}$ .

We next look at a fiber of the map  $G \times^Q (\mathfrak{r} \times X') \rightarrow \mathfrak{k}$  over  $t \in \mathfrak{k}^{reg}$ . The fibre is  $G \times^Q ((t + \mathfrak{n}) \times X')$ . We can write

$$(t + \mathfrak{n}) \times X' = \cup_{\tilde{y} \in X'} Q \cdot (t, \tilde{y})$$

because  $t + \mathfrak{n} + \bar{O}' = \cup_{y \in \bar{O}'} Q \cdot (t + y)$ . For  $\tilde{y} \in X'$ , we denote by  $y \in \bar{O}'$  its image by the map  $X' \rightarrow \bar{O}$ . Let  $Q_{(t, \tilde{y})}$  be the stabilizer group of  $(t, \tilde{y})$ . Then  $Q_{(t, \tilde{y})} \subset Q_{t+y}$ . In particular,  $Q_{(t, \tilde{y})} \subset L$ . Then, by the same argument as above, we have

$$G \times^Q ((t + \mathfrak{n}) \times X') = G \times^L (t \times X').$$

Since the right hand side depends only on  $L$  and  $X'$ , we conclude that

$$G \times^Q ((t + \mathfrak{n}) \times X') = G \times^{Q'} ((t + \mathfrak{n}') \times X').$$

Therefore we have a birational map

$$G \times^Q (\mathfrak{r} \times X') \dashrightarrow G \times^{Q'} (\mathfrak{r}' \times X'),$$

which is an isomorphism over any point of  $\mathfrak{k}^{reg}$ .  $\square$

By definition the map  $\mathfrak{m}'_Q : G \times^Q (\mathfrak{r} \times X') \rightarrow Z_{\mathfrak{l}, X'}$  is a birational map. For  $t \in \mathfrak{k}$  it induces a map  $\mathfrak{m}'_{Q, t} : G \times^Q ((t + \mathfrak{n}) \times X') \rightarrow Z_{\mathfrak{l}, X', t}$  between fibers.

**Lemma 3.** *The map  $\mathfrak{m}'_{Q, t}$  is a birational morphism for every  $t \in \mathfrak{k}$ . For  $t \in \mathfrak{k}^{reg}$ , the map  $\mathfrak{m}'_{Q, t}$  is an isomorphism. In particular,  $\mathfrak{m}'_Q$  is an isomorphism in codimension one.*

*Proof.* When  $t = 0$ , the map  $\mathfrak{m}_Q : G \times^Q (\mathfrak{r} \times \tilde{O}') \rightarrow Y_{\mathfrak{l}, \tilde{O}'}$  induces a birational morphism  $\mathfrak{m}_{Q, 0} : G \times^Q (\mathfrak{n} \times \tilde{O}') \rightarrow (G \cdot (\mathfrak{n} + \tilde{O}'))^s$ . Here  $(G \cdot (\mathfrak{n} + \tilde{O}'))^s$  is the Stein factorization of the generalized Springer map  $\mu : G \times^Q (\mathfrak{n} \times \tilde{O}') \rightarrow G \cdot (\mathfrak{n} + \tilde{O}')$ . We have a commutative diagram

$$\begin{array}{ccc} G \times^Q (\mathfrak{n} \times X') & \xrightarrow{\mathfrak{m}'_{Q, 0}} & Z_{\mathfrak{l}, X', 0} \\ \pi' \downarrow & & \downarrow \\ G \times^Q (\mathfrak{n} \times \tilde{O}') & \xrightarrow{\mathfrak{m}_{Q, 0}} & (G \cdot (\mathfrak{n} + \tilde{O}'))^s \end{array} \quad (10)$$

The composite  $\mathfrak{m}_{Q, 0} \circ \pi'$  is a generically finite surjective map. Since  $\mathfrak{m}'_Q$  has only connected fibers,  $\mathfrak{m}'_{Q, 0}$  also has connected fibers. This means that  $\mathfrak{m}'_{Q, 0}$  is a birational map. The space  $\mathfrak{r} + \tilde{O}'$  has a natural scaling  $\mathbf{C}^*$ -action. Let  $s : \mathbf{C}^* \rightarrow \text{Aut}(\mathfrak{r} + \tilde{O}')$  be the corresponding homomorphism. We define a new  $\mathbf{C}^*$ -action, say  $\sigma$ , on  $\mathfrak{r} + \tilde{O}'$  by the composite of the map  $\mathbf{C}^* \rightarrow \mathbf{C}^* (t \rightarrow t^2)$  and  $s$ . Then  $\sigma$  lifts to a  $\mathbf{C}^*$ -action on  $\mathfrak{r} \times X'$ . Thus  $G \times^Q (\mathfrak{r} \times X')$  has a  $\mathbf{C}^*$ -action. We also introduce a  $\mathbf{C}^*$ -action on  $\mathfrak{k}$  by the composite of  $\mathbf{C}^* \rightarrow \mathbf{C}^* (t \rightarrow t^2)$  and the scaling action  $\mathbf{C}^* \rightarrow \text{Aut}(\mathfrak{k})$ . Then the map  $G \times^Q (\mathfrak{r} \times X') \rightarrow \mathfrak{k}$  is  $\mathbf{C}^*$ -equivariant. By this  $\mathbf{C}^*$ -action we see that  $\mathfrak{m}'_{Q, t}$  is a birational map for every  $t \in \mathfrak{k}$ .

We next look at the fibers over  $t \in \mathfrak{k}^{reg}$ . We then have a commutative diagram

$$\begin{array}{ccc} G \times^Q ((t + \mathfrak{n}) \times X') & \xrightarrow{\mathfrak{m}'_{Q, t}} & Z_{\mathfrak{l}, X', t} \\ \downarrow & & \downarrow \\ G \times^Q (t + \mathfrak{n} \times \tilde{O}') & \xrightarrow{\mathfrak{m}_{Q, t}} & Y_{\mathfrak{l}, O', t} \end{array} \quad (11)$$

By [Na 3], Proposition 3.6,  $\mathfrak{m}_{Q, t}$  is an isomorphism. Moreover, by Lemma 2,  $G \times^Q (t + \mathfrak{n} + \tilde{O}') = G \times^L (t + \tilde{O}')$ . Taking the normalizations of both sides, we have  $G \times^Q ((t +$



$\mathfrak{n}) \times \tilde{O}') = G \times^L (t \times \tilde{O}')$ . Moreover we have  $G \times^Q ((t + \mathfrak{n}) \times X') = G \times^L (t \times X')$ . Hence  $Z_{\mathfrak{l}, O', t}$  is the Stein factorization of the finite surjective map  $G \times^L (t \times X') \rightarrow G \times^L (t \times \tilde{O}')$ . Then we see that  $\mathfrak{m}'_{Q, t}$  is an isomorphism.  $\square$

**Proposition 4.** *The maps  $Z_{\mathfrak{l}, X'} \rightarrow \mathfrak{k}$  and  $Y_{\mathfrak{l}, O'} \rightarrow \mathfrak{k}$  are both flat. Their central fibers  $Z_{\mathfrak{l}, X', 0}$  and  $Y_{\mathfrak{l}, O', 0}$  are both normal varieties. Moreover,  $Z_{\mathfrak{l}, X', 0} = X$ .*

*Proof.* This is similar to [Na 2], Lemma 2.1.  $\square$

Let  $\tilde{O}'$  be the normalization of  $\bar{O}'$ . We first introduce a Poisson structure on  $G \times^Q (\mathfrak{r} \times \tilde{O}')$  over  $\mathfrak{k}$ . In order to do this, we consider the  $G$ -equivariant map

$$\mu_t : G \times^Q (t + \mathfrak{n} \times \tilde{O}') \rightarrow \mathfrak{g}, \quad [g, t + x + y] \mapsto Ad_g(t + x + y)$$

for each  $t \in \mathfrak{k}$ . Then the image of this map coincides with the closure of an adjoint orbit on  $\mathfrak{g}$ , say  $O_{\mu_t}$ . The pull-back  $\mu_t^* \omega_t$  of the Kostant-Kirillov form  $\omega_t$  on  $O_{\mu_t}$  gives a symplectic 2-form on  $G \times^Q (t + \mathfrak{n} \times \tilde{O}')$  and  $\{\mu_t^* \omega_t\}_{t \in \mathfrak{k}}$  determines a relative symplectic 2-form on  $G \times^Q (\mathfrak{r} \times \tilde{O}')$  over  $\mathfrak{k}$  (cf. [Na 3], Proposition 4.2). This relative symplectic 2-form determines a Poisson structure on  $G \times^Q (\mathfrak{r} \times \tilde{O}')$  and the Poisson structure uniquely extends to a Poisson structure on  $G \times^Q (\mathfrak{r} \times \tilde{O}')$ . We next introduce a Poisson structure on  $G \times^Q (\mathfrak{r} \times X')$  over  $\mathfrak{k}$ . There is an etale finite map

$$G \times^Q (\mathfrak{r} \times (X')^0) \rightarrow G \times^Q (\mathfrak{r} \times O').$$

If we pull back the relative symplectic 2-form on  $G \times^Q (\mathfrak{r} \times O')$  by this map, we get a relative symplectic 2-form on  $G \times^Q (\mathfrak{r} \times (X')^0)$ . This relative symplectic 2-form determines a Poisson structure on  $G \times^Q (\mathfrak{r} \times (X')^0)$ . Moreover, such a Poisson structure uniquely extends to a Poisson structure on  $G \times^Q (\mathfrak{r} \times X')$ . Finally these Poisson structures respectively induce Poisson structures on  $Y_{\mathfrak{l}, O'}$  and  $Z_{\mathfrak{l}, X'}$  because

$$Y_{\mathfrak{l}, O'} = \text{Spec } \Gamma(G \times^Q (\mathfrak{r} \times O'), \mathcal{O}), \quad Z_{\mathfrak{l}, X'} = \text{Spec } \Gamma(G \times^Q (\mathfrak{r} \times X'), \mathcal{O}).$$

Therefore we have:

**Proposition 5.**  $G \times^Q (\mathfrak{r} \times X') \rightarrow \mathfrak{k}$  is a Poisson deformation of  $G \times^Q (\mathfrak{n} \times X')$ .  $Z_{\mathfrak{l}, X'} \rightarrow \mathfrak{k}$  is a Poisson deformation of  $X$ .

By assumption,  $N_G(L, T)$  acts on  $O'$ . For each  $w \in N_G(L, T)$ , the automorphism  $Ad_w : O' \rightarrow O'$  lifts to an automorphism  $\theta_w : (X')^0 \rightarrow (X')^0$ . In fact, take a point  $z \in O'$  and write  $O' = L/L^z$ . By the assumption  $Ad_w(z) \in O'$ .  $Ad_w : L \rightarrow L$  sends  $L^z$  onto  $L^{Ad_w(z)}$ . Hence  $Ad_w$  sends the identity component  $(L^z)^0$  of  $L^z$  to the identity component  $(L^{Ad_w(z)})^0$  of  $L^{Ad_w(z)}$ . In particular,  $Ad_w$  induces an isomorphism  $L/(L^z)^0 \rightarrow L/(L^{Ad_w(z)})^0$ . Recall that  $(X')^0 = L/(L^z)^0$  as an  $L$ -variety. Take points  $\tilde{z}, Ad_w(z) \in (X')^0$  lying over  $z$ ,  $Ad_w(z) \in O'$ . Then the isomorphism determines an automorphism  $\theta_w : (X')^0 \rightarrow (X')^0$  which sends  $\tilde{z}$  to  $Ad_w(z)$ . Note that, for a covering transformation  $\sigma \in \text{Aut}((X')^0/O')$ , the composite  $\sigma \circ \theta_w$  also gives a lifting of  $Ad_w$ . For

each  $w \in N_G(L, T)$ , we consider all liftings of  $\theta_w$  of  $Ad_w$ . Then these form a group  $\widetilde{N_G(L, T)}$  and one has an exact sequence

$$1 \rightarrow \text{Aut}((X')^0/O') \rightarrow \widetilde{N_G(L, T)} \rightarrow N_G(L, T) \rightarrow 1.$$

$\widetilde{N_G(L, T)}$  acts on  $O'$  by means of the map  $\widetilde{N_G(L, T)} \rightarrow N_G(L, T)$ . In general, the  $N_G(L, T)$ -action on  $O'$  does not lift to an action on  $(X')^0$ , but the  $\widetilde{N_G(L, T)}$ -action on  $O'$  lifts to an action on  $(X')^0$ . This  $\widetilde{N_G(L, T)}$ -action extends to an action on  $X'$ . For  $\tilde{w} \in \widetilde{N_G(L, T)}$  we denote by  $w \in N_G(L, T)$  its image by the map  $\widetilde{N_G(L, T)} \rightarrow N_G(L, T)$ . We define an isomorphism

$$\phi_{\tilde{w}} : G \times^Q (\mathfrak{r} \times X') \rightarrow G \times^{Ad_w(Q)} (Ad_w(\mathfrak{r}) \times X'), \quad [g, (y, \tilde{z})] \mapsto [gw^{-1}, (Ad_w(y), \tilde{w}(\tilde{z}))].$$

As is already remarked,  $N_G(L, T)$  (or its quotient group  $W'$ ) acts on  $Y_{\mathfrak{l}, O'}$ . We have a commutative diagram

$$\begin{array}{ccc} G \times^Q (\mathfrak{r} \times X') & \xrightarrow{\phi_{\tilde{w}}} & G \times^{Ad_w(Q)} (Ad_w(\mathfrak{r}) \times X') \\ \downarrow & & \downarrow \\ Y_{\mathfrak{l}, O'} & \xrightarrow{w} & Y_{\mathfrak{l}, O'} \end{array} \quad (12)$$

By Lemma 2,  $G \times^Q (\mathfrak{r} \times X')$  and  $G \times^{Ad_w(Q)} (Ad_w(\mathfrak{r}) \times X')$  are birational (not by  $\phi_{\tilde{w}}$ , but as varieties over  $Y_{\mathfrak{l}, O'}$ ), hence they have the same function field. By  $\phi_{\tilde{w}}$ , the group  $\widetilde{N_G(L, T)}$  acts on the function field. We have defined  $Z_{\mathfrak{l}, X'}$  as the normalization of  $Y_{\mathfrak{l}, O'}$  in this field. Hence  $\widetilde{N_G(L, T)}$  acts on  $Z_{\mathfrak{l}, X'}$  and the map  $Z_{\mathfrak{l}, X'} \rightarrow \mathfrak{k}$  is  $\widetilde{N_G(L, T)}$ -equivariant. Note that  $\widetilde{N_G(L, T)}$ -action on  $Z_{\mathfrak{l}, X'}$  is not effective. Since  $N_L(T)$ -action on  $O'$  lifts to an action on  $(X')^0$ ,  $N_L(T)$  is regarded as a subgroup of  $\widetilde{N_G(L, T)}$  and it is characterized as the subgroup consisting of the elements which act trivially on  $Z_{\mathfrak{l}, X'}$ . Define

$$\widetilde{W'} := \widetilde{N_G(L, T)} / N_L(T).$$

Then we have an exact sequence

$$1 \rightarrow \text{Aut}((X')^0/O') \rightarrow \widetilde{W'} \rightarrow W' \rightarrow 1$$

and  $\widetilde{W'}$  acts on  $Z_{\mathfrak{l}, X'}$ . Let us consider the central fiber  $Z_{\mathfrak{l}, X', 0}$  of the map  $Z_{\mathfrak{l}, X'} \rightarrow \mathfrak{k}$ . Then  $Z_{\mathfrak{l}, X', 0} = X$  and  $\widetilde{W'}$  acts on  $X$  as a covering transformation of  $X \rightarrow \bar{O}$ . Therefore we have a homomorphism

$$\rho_X : \widetilde{W'} \rightarrow \text{Aut}(X/\bar{O}).$$

We define  $W_X$  as the kernel of  $\rho_X$ :

$$W_X := \text{Ker}(\rho_X).$$

Let us consider the composite  $\text{Aut}((X')^0/O') \rightarrow \widetilde{W}' \xrightarrow{\rho_X} \text{Aut}(X/\bar{O})$ . This is an injection. In fact, an element  $\tau \in \text{Aut}((X')^0/O')$  determines an automorphism

$$\phi_\tau : G \times^Q (\mathfrak{n} \times X') \rightarrow G \times^Q (\mathfrak{n} \times X'), [g, (x, \tilde{y})] \mapsto [g, (x, \tau \tilde{y})],$$

which induces an automorphism of  $X$  over  $\bar{O}$ . If  $\rho_X(\tau) = id$ , then we get a commutative diagram

$$\begin{array}{ccc} G \times^Q (\mathfrak{n} \times X') & \xrightarrow{\phi_\tau} & G \times^Q (\mathfrak{n} \times X') \\ \downarrow & & \downarrow \\ X & \xrightarrow{id_X} & X \end{array} \quad (13)$$

This means that  $\phi_\tau = id$  and, hence  $\tau = id$ . We regard  $\text{Aut}((X')^0/O')$  as a subgroup of  $\text{Aut}(X/\bar{O})$  in this way. Then we have a commutative diagram of exact sequences

$$\begin{array}{ccccccc} & & 1 & & 1 & & \\ & & \downarrow & & \downarrow & & \\ & 1 & \longrightarrow & W_X & \longrightarrow & \pi(W_X) & \longrightarrow 1 \\ & \downarrow & & \downarrow & & \downarrow & \\ 1 & \longrightarrow & \text{Aut}((X')^0/O') & \longrightarrow & \widetilde{W}' & \xrightarrow{\pi} & W' \longrightarrow 1 \\ & id \downarrow & & \rho_X \downarrow & & \downarrow & \\ 1 & \longrightarrow & \text{Aut}((X')^0/O') & \longrightarrow & \text{Im}(\rho_X) & \longrightarrow & \text{Im}(\rho_X)/\text{Aut}((X')^0/O') \longrightarrow 1 \end{array} \quad (14)$$

Therefore,  $W_X$  can be regarded as a subgroup of  $W'$ .

*Twisting* (cf. [Na 1], pp.386-388) Define

$$\mathcal{S}(L) := \{Q \subset G \mid Q \text{ is a parabolic subgroup which contains } L \text{ as Levi part}\}.$$

Take an element  $Q$  from  $\mathcal{S}(L)$ . Then we obtain a new element  $Q' \in \mathcal{S}(L)$  from  $Q$  by making an operation called a *twisting*. We shall explain this. Fix a maximal torus  $T$  of  $L$ . Put  $\mathfrak{t} = \text{Lie}(T)$  and  $\mathfrak{l} = \text{Lie}(L)$ . Let  $\Phi \subset \mathfrak{t}^*$  be the root system determined by  $(\mathfrak{g}, \mathfrak{t})$ . If we take a basis  $\Delta$  of  $\Phi$ , then we can make a Dynkin diagram  $D$  attaching an element (= simple root) of  $\Delta$  to each vertex. Moreover, if we mark some vertices of  $D$  with back, then it determines a parabolic subalgebra  $\mathfrak{q}$  of  $\mathfrak{g}$ . More precisely, let  $I \subset \Delta$  be the set of simple roots corresponding to unmarked vertices. Let  $\Phi_I$  be the root subsystem of  $\Phi$  generated by  $I$ , and put  $\Phi_I^- := \Phi_I \cap \Phi^-$ , where  $\Phi^-$  is the set of negative roots. Then

$$\mathfrak{q} = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi_I^-} \mathfrak{g}_\alpha \oplus \bigoplus_{\alpha \in \Phi^+} \mathfrak{g}_\alpha$$

is a parabolic subalgebra of  $\mathfrak{g}$ . For  $\beta \in \Delta - I$ , we make a new marked Dynkin diagram  $\bar{D}$  from  $D$  by erasing the marking of  $\beta$ . By definition, the set  $\bar{I}$  of unmarked simple roots for  $\bar{D}$  is  $I \cup \{\beta\}$ . The new marked Dynkin diagram  $\bar{D}$  determines a parabolic subalgebra

$\bar{q}$ . The unmarked vertices of  $\bar{D}$  determines a Dynkin subdiagram, which is decomposed into the disjoint sum of the connected component containing  $\beta$  and the union of other components. Correspondingly, we have a decomposition  $\bar{I} = I_\beta \cup I'_\beta$  with  $\beta \in I_\beta$ . Let  $\bar{\mathfrak{l}}$  be the Levi part of  $\bar{q}$  containing  $\mathfrak{t}$ , and let  $\mathfrak{z}(\bar{\mathfrak{l}})$  be its center. Then  $\bar{\mathfrak{l}}/\mathfrak{z}(\bar{\mathfrak{l}})$  is decomposed into the direct sum of simple factors. Let  $\bar{\mathfrak{l}}_\beta$  be the simple factor corresponding to  $I_\beta$ , and let  $\bar{\mathfrak{l}}'_\beta$  be the direct sum of other simple factors. Then

$$\bar{l}/\mathfrak{d}(\bar{l}) = \bar{l}_\beta \oplus \bar{l}'_\beta.$$

Put  $\mathfrak{q}_\beta := \mathfrak{q} \cap \bar{\mathfrak{l}}_\beta$ . Then  $\mathfrak{q}_\beta$  is a parabolic subalgebra of  $\bar{\mathfrak{l}}_\beta$ . Let  $\pi : \bar{\mathfrak{l}} \rightarrow \bar{\mathfrak{l}}/\mathfrak{z}(\bar{\mathfrak{l}})$  be the quotient map. Note that

$$\begin{aligned}\bar{\mathfrak{q}} &= \bar{\mathfrak{l}} \oplus \mathfrak{n}(\bar{\mathfrak{q}}), \\ \mathfrak{q} &= \pi^{-1}(\mathfrak{q}_\beta \oplus \bar{\mathfrak{l}}'_\beta) \oplus \mathfrak{n}(\bar{\mathfrak{q}}).\end{aligned}$$

Let  $\Phi_{I_\beta}$  be the root subsystem of  $\Phi$  generated by  $I_\beta$ . We put  $\mathfrak{t}_\beta := \mathfrak{t}/\mathfrak{z}(\bar{\mathfrak{l}}) \cap \bar{\mathfrak{l}}_\beta$ . Then  $\Phi_{I_\beta}$  is the root system for  $(\bar{\mathfrak{l}}_\beta, \mathfrak{t}_\beta)$ . The root system  $\Phi_{I_\beta}$  has a standard involution  $-1$ . There is a unique involution  $\phi_\beta$  of  $\bar{\mathfrak{l}}_\beta$  which stabilizes  $\mathfrak{t}_\beta$  and which acts on  $\Phi_{I_\beta}$  as  $-1$ . We define

$$\mathfrak{q}' := \pi^{-1}(\phi_\beta(\mathfrak{q}_\beta) \oplus \bar{\mathfrak{l}}'_\beta) \oplus \mathfrak{n}(\bar{\mathfrak{q}}).$$

Then  $\mathfrak{q}'$  is a parabolic subalgebra of  $\mathfrak{g}$ , which contains  $\mathfrak{l}$  as the Levi part. We say that  $\mathfrak{q}'$  is a *twist* of  $\mathfrak{q}$  by  $\beta$ .

**Example 6.** (i) Assume that  $\mathfrak{q}$  is determined by the following marked Dynkin diagram  $D$

$$\cdots \text{---} \bullet_{\alpha_0} \text{---} \circ_{\alpha_1} \text{---} \circ_{\alpha_2} \text{---} \cdots \quad \cdots \text{---} \bullet_{\alpha_k} \text{---} \cdots \quad \cdots \text{---} \circ_{\alpha_{n-1}} \text{---} \circ_{\alpha_n} \text{---} \bullet_{\alpha_{n+1}} \text{---} \cdots$$

We make a new marked Dynkin diagram  $\bar{D}$  by erasing the marking of  $\alpha_k$ . The unmarked vertices of  $\bar{D}$  determine a non-connected Dynkin subdiagram. Its connected component containing  $\alpha_k$  is of type  $A_n$ . The twist  $\mathbf{q}'$  of  $\mathbf{q}$  by  $\alpha_k$  is then determined by the marked Dynkin diagram  $D'$ :

$$\begin{array}{ccccccc} \text{---} & \bullet & \text{---} & \overset{-\alpha_n}{\circ} & \text{---} & \overset{-\alpha_{n-1}}{\circ} & \text{---} & \text{---} & \bullet & \text{---} & \overset{-\alpha_k}{\circ} & \text{---} & \text{---} & \overset{-\alpha_2}{\circ} & \text{---} & \overset{-\alpha_1}{\circ} & \text{---} & \bullet & \text{---} & \text{---} \\ \alpha_0 + \sum_{1 \leq i < n} \alpha_i & & & & & & & & & & & & & & & & & & & & & \alpha_{n+1} + \sum_{1 \leq i < n} \alpha_i \end{array}$$

Other simple roots in  $D$  different from  $\{\alpha_i\}_{0 \leq i \leq n+1}$  do not have any effect by the twist.

(ii) Assume that  $\mathfrak{g}$  is determined by the marked Dynkin diagram

The twist  $\mathfrak{g}'$  of  $\mathfrak{g}$  by  $\alpha_k$  is then determined by the marked Dynkin diagram

$$\begin{array}{c}
\cdots \text{---} \bullet \text{---} \overset{-\alpha_n}{\circ} \text{---} \overset{-\alpha_{n-1}}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{-\alpha_2}{\circ} \begin{array}{l} \nearrow \overset{-\alpha_1}{\circ} \\ \searrow \bullet \end{array} \\
\alpha_0 + \sum_{1 \leq i \leq n} \alpha_i \qquad \qquad \qquad \alpha_1 + 2 \sum_{2 \leq i \leq n-1} \alpha_i + \alpha_n + \alpha_{n+1}
\end{array}$$

(iii) Assume that  $\mathfrak{q}$  is determined by the marked Dynkin diagram

$$\cdots \text{---} \bullet \text{---} \overset{\alpha_1}{\circ} \text{---} \overset{\alpha_2}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{\alpha_{n-1}}{\circ} \text{---} \overset{\alpha_n}{\circ} \Rightarrow \bullet \text{---} \overset{\alpha_{n+1}}{\circ}$$

The twist  $\mathfrak{q}'$  of  $\mathfrak{q}$  by  $\alpha_k$  is then determined by the marked Dynkin diagram

$$\begin{array}{c}
\cdots \text{---} \bullet \text{---} \overset{-\alpha_n}{\circ} \text{---} \overset{-\alpha_{n-1}}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{-\alpha_2}{\circ} \text{---} \overset{-\alpha_1}{\circ} \Rightarrow \bullet \\
\alpha_0 + \sum_{1 \leq i \leq n} \alpha_i \qquad \qquad \qquad \sum_{1 \leq i \leq n} \alpha_i + \alpha_{n+1}
\end{array}$$

(iv) Assume that  $\mathfrak{q}$  is determined by the marked Dynkin diagram

$$\cdots \text{---} \bullet \text{---} \overset{\alpha_1}{\circ} \text{---} \overset{\alpha_2}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{\alpha_{n-1}}{\circ} \text{---} \overset{\alpha_n}{\circ} \Leftarrow \bullet \text{---} \overset{\alpha_{n+1}}{\circ}$$

The twist  $\mathfrak{q}'$  of  $\mathfrak{q}$  by  $\alpha_k$  is then determined by the marked Dynkin diagram

$$\begin{array}{c}
\cdots \text{---} \bullet \text{---} \overset{-\alpha_n}{\circ} \text{---} \overset{-\alpha_{n-1}}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{-\alpha_2}{\circ} \text{---} \overset{-\alpha_1}{\circ} \Leftarrow \bullet \\
\alpha_0 + \sum_{1 \leq i \leq n} \alpha_i \qquad \qquad \qquad 2 \sum_{1 \leq i \leq n} \alpha_i + \alpha_{n+1}
\end{array}$$

(v) Assume that  $\mathfrak{q}$  is determined by the marked Dynkin diagram

$$\cdots \text{---} \bullet \text{---} \overset{\alpha_1}{\circ} \text{---} \overset{\alpha_2}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{\alpha_{n-1}}{\circ} \Rightarrow \overset{\alpha_n}{\circ}$$

The twist  $\mathfrak{q}'$  of  $\mathfrak{q}$  by  $\alpha_k$  is then determined by the marked Dynkin diagram

$$\begin{array}{c}
\cdots \text{---} \bullet \text{---} \overset{-\alpha_1}{\circ} \text{---} \overset{-\alpha_2}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{-\alpha_{n-1}}{\circ} \Rightarrow \overset{-\alpha_n}{\circ} \\
\alpha_0 + 2 \sum_{1 \leq i \leq n} \alpha_i
\end{array}$$

(vi) Assume that  $\mathfrak{q}$  is determined by the marked Dynkin diagram

$$\cdots \text{---} \bullet \text{---} \overset{\alpha_1}{\circ} \text{---} \overset{\alpha_2}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{\alpha_{n-1}}{\circ} \Leftarrow \overset{\alpha_n}{\circ}$$

The twist  $\mathfrak{q}'$  of  $\mathfrak{q}$  by  $\alpha_k$  is then determined by the marked Dynkin diagram

$$\begin{array}{c} \cdots \text{---} \bullet \text{---} \overset{-\alpha_1}{\circ} \text{---} \overset{-\alpha_2}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{-\alpha_{n-1}}{\circ} \text{---} \overset{-\alpha_n}{\circ} \\ \alpha_0 + 2 \sum_{1 \leq i \leq n-1} \alpha_i + \alpha_n \end{array}$$

(vii) Let  $n$  be an odd number and let  $k$  be an integer such that  $1 \leq k \leq n-2$ . Assume that  $\mathfrak{q}$  is determined by the marked Dynkin diagram

$$\begin{array}{c} \cdots \text{---} \bullet \text{---} \overset{\alpha_1}{\circ} \text{---} \overset{\alpha_2}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{\alpha_{n-2}}{\circ} \begin{array}{l} \nearrow \overset{\alpha_{n-1}}{\circ} \\ \searrow \overset{\alpha_n}{\circ} \end{array} \end{array}$$

Then the twist  $\mathfrak{q}'$  of  $\mathfrak{q}$  by  $\alpha_k$  is determined by the marked Dynkin diagram

$$\begin{array}{c} \cdots \text{---} \bullet \text{---} \overset{-\alpha_1}{\circ} \text{---} \overset{-\alpha_2}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{-\alpha_{n-2}}{\circ} \begin{array}{l} \nearrow \overset{-\alpha_n}{\circ} \\ \searrow \overset{-\alpha_{n-1}}{\circ} \end{array} \\ \alpha_0 + 2 \sum_{1 \leq i \leq n-2} \alpha_i + \alpha_{n-1} + \alpha_n \end{array}$$

On the other hand, when  $k = n-1$ ,  $\mathfrak{q}$  is given by the marked Dynkin diagram

$$\begin{array}{c} \cdots \text{---} \bullet \text{---} \overset{\alpha_1}{\circ} \text{---} \overset{\alpha_2}{\circ} \text{---} \cdots \text{---} \circ \text{---} \cdots \text{---} \overset{\alpha_{n-2}}{\circ} \begin{array}{l} \nearrow \bullet \overset{\alpha_{n-1}}{\circ} \\ \searrow \overset{\alpha_n}{\circ} \end{array} \end{array}$$

Then the twist  $\mathfrak{q}'$  of  $\mathfrak{q}$  by  $\alpha_{n-1}$  is determined by

$$\begin{array}{c} \cdots \text{---} \bullet \text{---} \overset{-\alpha_1}{\circ} \text{---} \overset{-\alpha_2}{\circ} \text{---} \cdots \text{---} \circ \text{---} \cdots \text{---} \overset{-\alpha_{n-2}}{\circ} \begin{array}{l} \nearrow \overset{-\alpha_n}{\circ} \\ \searrow \bullet \overset{-\alpha_{n-1}}{\circ} \end{array} \\ \alpha_0 + 2 \sum_{1 \leq i \leq n-2} \alpha_i + \alpha_{n-1} + \alpha_n \end{array}$$

(viii) Let  $n$  be an even integer, and let  $k$  be an integer such that  $1 \leq k \leq n-1$ . Assume  $\mathfrak{q}$  is determined by the marked Dynkin diagram

$$\begin{array}{c} \cdots \text{---} \bullet \text{---} \overset{\alpha_1}{\circ} \text{---} \overset{\alpha_2}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{\alpha_{n-2}}{\circ} \begin{array}{l} \nearrow \overset{\alpha_{n-1}}{\circ} \\ \searrow \overset{\alpha_n}{\circ} \end{array} \end{array}$$

The twist  $\mathfrak{q}'$  of  $\mathfrak{q}$  by  $\alpha_k$  is then determined by the marked Dynkin diagram

$$\begin{array}{c} \cdots \text{---} \bullet \text{---} \overset{-\alpha_1}{\circ} \text{---} \overset{-\alpha_2}{\circ} \text{---} \cdots \text{---} \bullet \text{---} \cdots \text{---} \overset{-\alpha_{n-2}}{\circ} \begin{array}{l} \nearrow \overset{-\alpha_{n-1}}{\circ} \\ \searrow \overset{-\alpha_n}{\circ} \end{array} \\ \alpha_0 + 2 \sum_{1 \leq i \leq n-2} \alpha_i + \alpha_{n-1} + \alpha_n \end{array}$$

**Theorem 7.** (1) Every element  $Q \in \mathcal{S}(L)$  is obtained from a fixed element  $Q_0 \in \mathcal{S}(L)$  by a finite succession of the twists. Every  $\mathbf{Q}$ -factorial terminalization of  $Z_{\mathfrak{l}, X'}$  has the form  $G \times^Q (\mathfrak{r} \times X')$  with  $Q \in \mathcal{S}(L)$ . Conversely, for any  $Q \in \mathcal{S}(L)$ ,  $G \times^Q (\mathfrak{r} \times X')$  gives a  $\mathbf{Q}$ -factorial terminalization of  $Z_{\mathfrak{l}, X'}$ . Moreover, there is a one-one correspondence between the set of all  $\mathbf{Q}$ -factorial terminalizations of  $Z_{\mathfrak{l}, X'}$  and  $\mathcal{S}(L)$ .

(2) Every  $\mathbf{Q}$ -factorial terminalization of  $X$  has the form  $G \times^Q (\mathfrak{n} \times X')$  with  $Q \in \mathcal{S}(L)$ . Conversely, for any  $Q \in \mathcal{S}(L)$ ,  $G \times^Q (\mathfrak{n} \times X')$  gives a  $\mathbf{Q}$ -factorial terminalization of  $X$ .

**Remark 8.**

(i) In (2) different two parabolic subgroups  $Q, Q' \in \mathcal{S}(L)$  may possibly give the same  $\mathbf{Q}$ -factorial terminalization of  $X$ .

(ii) The pair  $(X', O')$  is uniquely determined by  $X$ . We call  $(X', O')$  the *core* of  $X$ .

*Proof.* (1) The first statement is proved in [Na 1], Theorem 1.3. For each  $Q \in \mathcal{S}(L)$ , we have isomorphisms

$$\mathrm{Pic}(G \times^Q (\mathfrak{r} \times X')) \cong \mathrm{Pic}(G/Q) \cong \mathrm{Hom}_{\mathrm{alg. gp}}(L, \mathbf{C}^*).$$

The nef cone  $\overline{\mathrm{Amp}}(G \times^Q (\mathfrak{r} \times X')) \subset \mathrm{Pic}(G \times^Q (\mathfrak{r} \times X')) \otimes_{\mathbf{Z}} \mathbf{R}$  is identified with a cone in  $\mathrm{Hom}_{\mathrm{alg. gp}}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R}$ . By [Na 1], Remark 1.6 we have

$$\mathrm{Hom}_{\mathrm{alg. gp}}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R} = \bigcup_{Q \in \mathcal{S}(L)} \overline{\mathrm{Amp}}(G \times^Q (\mathfrak{r} \times X'))$$

Fix  $Q_0 \in \mathcal{S}(L)$  and identify  $\mathrm{Pic}(G \times^{Q_0} (\mathfrak{r}_0 \times X'))$  with  $\mathrm{Hom}_{\mathrm{alg. gp}}(L, \mathbf{C}^*)$ . Then, by Lemma 3, we have

$$\mathrm{Hom}_{\mathrm{alg. gp}}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R} = \mathrm{Mov}(G \times^{Q_0} (\mathfrak{r}_0 \times X'))$$

For each  $Q \in \mathcal{S}(L)$ , we can identify the nef cone  $\overline{\mathrm{Amp}}(G \times^Q (\mathfrak{r} \times X'))$  as a cone inside  $\mathrm{Mov}(G \times^{Q_0} (\mathfrak{r}_0 \times X'))$  by the birational map

$$G \times^Q (\mathfrak{r} \times X') \dashrightarrow G \times^{Q_0} (\mathfrak{r}_0 \times X').$$

As in [Na 1], Observation 2 we can check that this identification is the same one explained above.

(2) Once (1) is proved, (2) is proved in the same way as in [Na 1], Corollary 3.4.  $\square$

**Theorem 9.** The following commutative diagram gives the universal Poisson deformations of  $G \times^Q (\mathfrak{n} \times X')$  and  $X$ :

$$\begin{array}{ccc} G \times^Q (\mathfrak{r} \times X') & \longrightarrow & Z_{\mathfrak{l}, X'} / W_X \\ \downarrow & & \downarrow \\ \mathfrak{k} & \longrightarrow & \mathfrak{k} / W_X \end{array} \quad (15)$$

*Proof.* Put  $0_X = \pi^{-1}(0)$  for  $\pi : X \rightarrow \bar{O}$ . Then  $G \times^Q (\mathfrak{r} + 0_X) \subset G \times^Q (\mathfrak{r} \times X')$  is a closed Poisson subscheme. Moreover,  $G \times^Q (\mathfrak{r} + 0_X) \rightarrow \mathfrak{k}$  is a Poisson deformation of  $T^*(G/Q) = G \times^Q (\mathfrak{n} + 0_X)$ . Let

$$p\kappa_{T^*(G/Q)} : \mathfrak{k} \rightarrow H^2(T^*(G/Q), \mathbf{C})$$

be the Poisson Kodaira Spencer map for this Poisson deformation at  $0 \in \mathfrak{k}$ . Let

$$p\kappa_{G \times^Q (\mathfrak{n} \times X')} : \mathfrak{k} \rightarrow H^2(G \times^Q (\mathfrak{n} \times X'), \mathbf{C})$$

be the Poisson Kodaira Spencer map for  $G \times^Q (\mathfrak{r} \times X') \rightarrow \mathfrak{k}$  at  $0 \in \mathfrak{k}$ . Then  $p\kappa_{T^*(G/Q)}$  factorizes as

$$p\kappa_{T^*(G/Q)} : \mathfrak{k} \xrightarrow{p\kappa_{G \times^Q (\mathfrak{n} \times X')}} H^2(G \times^Q (\mathfrak{n} \times X'), \mathbf{C}) \xrightarrow{res} H^2(T^*(G/Q)).$$

By [Na 2], Proposition 2.7,  $p\kappa_{T^*(G/Q)}$  is an isomorphism. Then  $p\kappa_{G \times^Q (\mathfrak{n} \times X')}$  is an injection. Since  $\dim \mathfrak{k} = \dim H^2(G \times^Q (\mathfrak{n} \times X'), \mathbf{C})$ , we see that  $p\kappa_{G \times^Q (\mathfrak{n} \times X')}$  is an isomorphism, which implies the versality of the Poisson deformation. But, since the Poisson deformation functor  $\text{PD}_{G \times^Q (\mathfrak{n} \times X')}$  is pro-representable by [Na 6], Corollary 2.5, we see that  $G \times^Q (\mathfrak{r} \times X') \rightarrow \mathfrak{k}$  is the universal Poisson deformation of  $G \times^Q (\mathfrak{n} \times X')$ . Then the Weyl group  $W(X)$  of  $X$  acts linearly on  $\mathfrak{k}$  and there is a universal Poisson deformation

$$\mathcal{X} \rightarrow \mathfrak{k}/W(X)$$

of  $X$  (cf. [Na 4], Main Theorem). The Poisson deformation  $Z_{\mathfrak{l}, X'} \rightarrow \mathfrak{k}$  is nothing but the pull back of the universal Poisson deformation by the quotient map ([ibid])

$$\begin{array}{ccc} Z_{\mathfrak{l}, X'} & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ \mathfrak{k} & \longrightarrow & \mathfrak{k}/W(X) \end{array} \quad (16)$$

Then  $W(X)$  acts on  $Z_{\mathfrak{l}, X'}$  as  $G$ -equivariant Poisson automorphisms, and  $\mathcal{X} = Z_{\mathfrak{l}, X'}/W(X)$ .

The next proposition completes the proof of Theorem 9.

**Proposition 10.**  $W(X) = W_X$ .

*Proof.* We first show that  $W(X) \subset W_X$ . The Poisson structure on  $G \times^Q (\mathfrak{r} \times X')$  determines a symplectic 2-form  $\omega_t$  on the regular part of  $G \times^Q ((t + \mathfrak{n}) \times X')$  for each  $t \in \mathfrak{k}$ . Each symplectic 2-form  $\omega_t$  determines a 2-nd cohomology class  $[\omega_t] \in H^2(G \times^Q ((t + \mathfrak{n}) \times X'), \mathbf{C})$  (not only a 2-nd cohomology class on the regular part). The restriction maps of 2-nd cohomology are both isomorphisms:

$$H^2(G \times^Q ((t + \mathfrak{n}) \times X'), \mathbf{C}) \xleftarrow{\text{res}_t} H^2(G \times^Q (\mathfrak{r} \times X'), \mathbf{C}) \xrightarrow{\text{res}_0} H^2(G \times^Q (\mathfrak{n} \times X'), \mathbf{C}).$$

We define the period map  $p : \mathfrak{k} \rightarrow H^2(G \times^Q (\mathfrak{n} \times X'), \mathbf{C})$  by  $p(t) := \text{res}_0 \circ \text{res}_t^{-1}([\omega_t])$ . Since  $G \times^Q (\mathfrak{r} \times X') \rightarrow \mathfrak{k}$  is the universal Poisson deformation of  $G \times^Q (\mathfrak{n} \times X')$ , the period map  $p$  is an isomorphism. We identify  $\mathfrak{k}$  with  $H^2(G \times^Q (\mathfrak{r} \times X'), \mathbf{C})$  by  $\text{res}_0^{-1} \circ p$ . The  $W(X)$ -action on  $H^2(G \times^Q (\mathfrak{r} \times X'), \mathbf{C})$  is defined over  $\mathbf{Q}$ . In particular,  $W(X)$  acts on  $H^2(G \times^Q (\mathfrak{r} \times X'), \mathbf{R})$ . By [Na 4], Main Theorem,  $H^2(G \times^Q (\mathfrak{r} \times X'), \mathbf{R})$  is divided into finite number of chambers by finite linear spaces  $\{H_i\}_{i \in I}$  of codimension 1, and each chamber corresponds to an ample cone of  $G \times^{Q'} (\mathfrak{r}' \times X')$  for  $Q' \in \mathcal{S}(L)$ . There is a finite subset  $J$  of  $I$  such that  $W(X)$  is generated by the reflections  $\sigma_i$  with respect to  $H_i$  with



$i \in J$ . Take  $i \in J$  and put  $H := H_i$  and  $\sigma := \sigma_i$ . Take a chamber which has (an non-empty open set of)  $H$  as a codimension 1 face. This chamber is  $\text{Amp}(G \times^P (\mathfrak{r}(\mathfrak{p}) \times X'))$  for some  $P \in \mathcal{S}(L)$ . The element  $\sigma$  induces an automorphism  $\sigma_{\mathfrak{k}} : \mathfrak{k} \rightarrow \mathfrak{k}$ . The fiber product  $(G \times^P (\mathfrak{r}(\mathfrak{p}) \times X')) \times_{\mathfrak{k}} \mathfrak{k}$  fits in the commutative diagram

$$\begin{array}{ccc}
(G \times^P (\mathfrak{r}(\mathfrak{p}) \times X')) \times_{\mathfrak{k}} \mathfrak{k} & \xrightarrow{\phi_{\sigma}} & G \times^P (\mathfrak{r}(\mathfrak{p}) \times X') \\
\downarrow & & \downarrow \\
Z_{\mathfrak{l}, X'} & \longrightarrow & Z_{\mathfrak{l}, X'} \\
\downarrow & & \downarrow \\
\mathfrak{k} & \xrightarrow{\sigma_{\mathfrak{k}}} & \mathfrak{k}
\end{array} \tag{17}$$

Here each square is the Cartesian square. Since  $(G \times^P (\mathfrak{r}(\mathfrak{p}) \times X')) \times_{\mathfrak{k}} \mathfrak{k}$  is a  $\mathbf{Q}$ -factorial terminalization of  $Z_{\mathfrak{l}, X'}$ , one can write it as  $G \times^{P'} (\mathfrak{r}(\mathfrak{p}') \times X')$  for some  $P' \in \mathcal{S}(L)$ .

**Claim 1.** *There is an element  $w \in W'$  such that  $P = \text{Ad}_w(P')$ .*

*Proof.* In the commutative diagram above, let us consider the central fibers. Then we have

$$\begin{array}{ccc}
G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X') & \xrightarrow{\cong} & G \times^P (\mathfrak{n}(\mathfrak{p}) \times X') \\
\mathfrak{m}'_{P',0} \downarrow & & \mathfrak{m}'_{P,0} \downarrow \\
X & \xrightarrow{id} & X \\
\downarrow & & \downarrow \\
\{0\} & \xrightarrow{\sigma} & \{0\}
\end{array} \tag{18}$$

The isomorphism  $G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X') \cong G \times^P (\mathfrak{n}(\mathfrak{p}) \times X')$  is  $G$ -equivariant. Recall that  $\pi^{-1}(0)$  is a point for  $\pi : X \rightarrow \bar{O}$ . We put  $0_X := \pi^{-1}(0)$ . Then the  $G$ -equivariant isomorphism induces a  $G$ -equivariant isomorphism between  $\mathfrak{m}'_{P',0}{}^{-1}(0_X) = G/P'$  and  $\mathfrak{m}'_{P,0}{}^{-1}(0_X) = G/P$ . If the isomorphism sends  $\bar{1} \in G/P'$  to  $g_0 \in G/P$ , then by the  $G$ -equivariance, it sends  $\bar{g}$  to  $\overline{gg_0}$  for  $g \in G$ . Similarly it sends  $\overline{gp'}$  to  $\overline{gp'g_0}$  for  $p' \in P'$ . Since  $\bar{g} = \overline{gp'}$  in  $G/P'$ , we have  $\overline{gg_0} = \overline{gp'g_0}$  in  $G/P$ . This means that one can write  $gp'g_0 = gg_0p$  with a suitable  $p \in P$ . Then  $p' = g_0pg_0^{-1}$ . Therefore  $P'$  is conjugate to  $P$ . Since  $P$  and  $P'$  are contained in  $\mathcal{S}(L)$ , one can write  $P = \text{Ad}_w(P')$  with a suitable element  $w$  of  $W'$  by [Na 1], Proposition 2.2. Moreover, by the same proposition, this  $w$  is uniquely determined.  $\square$

By the definition of  $W'$ ,  $w$  acts on  $\mathfrak{k}$ . We denote by  $w_{\mathfrak{k}}$  the automorphism of  $\mathfrak{k}$  induced by  $w$ . We take a lift  $\tilde{w} \in \tilde{W}'$  of  $w$  and define an isomorphism

$$\phi_{\tilde{w}} : G \times^{P'} (\mathfrak{r}(\mathfrak{p}') \times X') \rightarrow G \times^P (\mathfrak{r}(\mathfrak{p}) \times X')$$

by  $[g, (x, \tilde{y})] \mapsto [gw^{-1}, (\text{Ad}_w(x), \tilde{w}(\tilde{y}))]$  for  $g \in G$ ,  $x \in \mathfrak{r}(\mathfrak{p}')$  and  $\tilde{y} \in X'$ . There is a

commutative diagram

$$\begin{array}{ccc}
G \times^{P'} (\mathfrak{r}(\mathfrak{p}') \times X') & \xrightarrow{\phi_{\tilde{w}}} & G \times^P (\mathfrak{r}(\mathfrak{p}) \times X') \\
\downarrow & & \downarrow \\
\mathfrak{k} & \xrightarrow{w_{\mathfrak{k}}} & \mathfrak{k}
\end{array} \tag{19}$$

**Claim 2.**  $w_{\mathfrak{k}} = \sigma_{\mathfrak{k}}$ .

*Proof.* By using the period map  $p$  for  $G \times^P (\mathfrak{r} \times X') \rightarrow \mathfrak{k}$ , we identify  $\mathfrak{k}$  with  $H^2(G \times^P (\mathfrak{n}(\mathfrak{p}) \times X'), \mathbf{C})$ . The isomorphism  $\phi_{\tilde{w}}$  induces an isomorphism of the 2-nd cohomology

$$(\phi_{\tilde{w}})^* : H^2(G \times^P (\mathfrak{r}(\mathfrak{p}) \times X'), \mathbf{C}) \rightarrow H^2(G \times^{P'} (\mathfrak{r}(\mathfrak{p}') \times X'), \mathbf{C}).$$

On the other hand, since both  $G \times^P (\mathfrak{r}(\mathfrak{p}) \times X')$  and  $G \times^{P'} (\mathfrak{r}(\mathfrak{p}') \times X')$  give  $\mathbf{Q}$ -factorial terminalizations of  $Z_{l,X'}$ , there is a natural identification

$$(\mathfrak{m}_P^{-1} \circ \mathfrak{m}_{P'})_* : H^2(G \times^{P'} (\mathfrak{r}(\mathfrak{p}') \times X'), \mathbf{C}) \rightarrow H^2(G \times^P (\mathfrak{r}(\mathfrak{p}) \times X'), \mathbf{C}).$$

Take  $t \in \mathfrak{k}$  general. Then, as seen in the proof of Lemma 2, the birational map  $\mathfrak{m}_P^{-1} \circ \mathfrak{m}_{P'}$  identifies  $G \times^P ((w_{\mathfrak{k}}(t) + \mathfrak{n}(\mathfrak{p})) \times X')$  with  $G \times^{P'} (w_{\mathfrak{k}}(t) + \mathfrak{n}(\mathfrak{p}')) \times X')$ . We then have the commutative diagram

$$\begin{array}{ccc}
H^2(G \times^P (\mathfrak{n}(\mathfrak{p}) \times X'), \mathbf{C}) & \longleftarrow & H^2(G \times^P (\mathfrak{n}(\mathfrak{p}) \times X'), \mathbf{C}) \\
\cong \text{res}_0 \uparrow & & \cong \text{res}_0 \uparrow \\
H^2(G \times^P (\mathfrak{r}(\mathfrak{p}) \times X'), \mathbf{C}) & \xleftarrow{(\mathfrak{m}_P^{-1} \circ \mathfrak{m}_{P'})_* \circ (\phi_{\tilde{w}})^*} & H^2(G \times^P (\mathfrak{r}(\mathfrak{p}) \times X'), \mathbf{C}) \\
\cong \text{res}_{w_{\mathfrak{k}}(t)} \downarrow & & \cong \text{res}_t \downarrow \\
H^2(G \times^P ((w_{\mathfrak{k}}(t) + \mathfrak{n}(\mathfrak{p})) \times X'), \mathbf{C}) & \xleftarrow{(\phi_{\tilde{w},t})^*} & H^2(G \times^P ((t + \mathfrak{n}(\mathfrak{p})) \times X'), \mathbf{C})
\end{array} \tag{20}$$

By the diagram, we have

$$\begin{array}{ccc}
w_{\mathfrak{k}}(t) & \longrightarrow & t \\
\text{res}_0 \circ \text{res}_{w_{\mathfrak{k}}(t)}^{-1} \uparrow & & \text{res}_0 \circ \text{res}_t^{-1} \uparrow \\
[\omega_{w_{\mathfrak{k}}(t)}] & \longrightarrow & [\omega_t]
\end{array} \tag{21}$$

Hence we have

$$w_{\mathfrak{k}}^{-1} = \text{res}_0 \circ (\mathfrak{m}_P^{-1} \circ \mathfrak{m}_{P'})_* \circ (\phi_{\tilde{w}})^* \circ (\text{res}_0)^{-1}.$$

Similarly, for  $\sigma$ , we have already constructed an isomorphism

$$\phi_{\sigma} : G \times^{P'} (\mathfrak{r}(\mathfrak{p}') \times X') \rightarrow G \times^P (\mathfrak{r}(\mathfrak{p}) \times X')$$

in the commutative diagram just before Claim 1.

This induces an isomorphism of 2-nd cohomology

$$(\phi_{\sigma})^* : H^2(G \times^P (\mathfrak{r}(\mathfrak{p}) \times X'), \mathbf{C}) \rightarrow H^2(G \times^{P'} (\mathfrak{r}(\mathfrak{p}') \times X'), \mathbf{C}).$$

Then we have

$$\sigma_{\mathfrak{t}}^{-1} = \text{res}_0 \circ (\mathfrak{m}_P^{-1} \circ \mathfrak{m}_{P'})_* \circ (\phi_\sigma)^* \circ (\text{res}_0)^{-1}.$$

It is sufficient to prove that  $(\phi_{\tilde{w}})^*$  and  $(\phi_\sigma)^*$  coincide. If we restrict  $\phi_{\tilde{w}}$  and  $\phi_\sigma$  to the fibers over  $0 \in \mathfrak{l}$ , then we respectively have isomorphisms  $\phi_{\tilde{w},0} : G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X') \rightarrow G \times^P (\mathfrak{n}(\mathfrak{p}) \times X')$  and  $\phi_{\sigma,0} : G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X') \rightarrow G \times^P (\mathfrak{n}(\mathfrak{p}) \times X')$ . Since there is a deformation retract of  $G \times^P (\mathfrak{r}(\mathfrak{p}) \times X')$  (resp.  $G \times^{P'} (\mathfrak{r}(\mathfrak{p}') \times X')$ ) to  $G \times^P (\mathfrak{n}(\mathfrak{p}) \times X'), \mathbf{C}$ ) (resp.  $G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X')$ ), it is enough to prove that the two maps

$$(\phi_{\tilde{w},0})^* : H^2(G \times^P (\mathfrak{n}(\mathfrak{p}) \times X'), \mathbf{C}) \rightarrow H^2(G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X'), \mathbf{C})$$

and

$$(\phi_{\sigma,0})^* : H^2(G \times^P (\mathfrak{n}(\mathfrak{p}) \times X'), \mathbf{C}) \rightarrow H^2(G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X'), \mathbf{C})$$

coincide. Recall that  $\phi_{\tilde{w},0}$  induces an automorphism  $\rho_X(\tilde{w}) \in \text{Aut}(X/\bar{O})$ :

$$\begin{array}{ccc} G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X') & \xrightarrow{\phi_{\tilde{w},0}} & G \times^P (\mathfrak{n}(\mathfrak{p}) \times X') \\ \mathfrak{m}'_{P',0} \downarrow & & \mathfrak{m}'_{P,0} \downarrow \\ X & \xrightarrow{\rho_X(\tilde{w})} & X \end{array} \quad (22)$$

By definition  $X$  has a  $G$ -action and  $\rho_X(\tilde{w})$  is  $G$ -equivariant. The  $G$ -action on  $X$  lifts to a  $G$ -action on  $G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X')$  (resp.  $G \times^P (\mathfrak{n}(\mathfrak{p}) \times X')$ ) by  $\mathfrak{m}'_{P',0}$  (resp.  $\mathfrak{m}'_{P,0}$ ). Hence  $\phi_{\tilde{w}}$  is also  $G$ -equivariant. Recall that, for the covering map  $\pi : X \rightarrow \bar{O}$ , the inverse image  $\pi^{-1}(0)$  consists of one point  $0_X$ . Since  $\rho_X(\tilde{w})(0_X) = 0_X$ ,  $\phi_{\tilde{w},0}$  induces a  $G$ -equivariant isomorphism between  $\mathfrak{m}'_{P',0}{}^{-1}(0_X) = G/P'$  and  $\mathfrak{m}'_{P,0}{}^{-1}(0_X) = G/P$ . Let  $N_G(P)$  be the normalizer of  $P$  in  $G$ . Fix an element  $g_0 \in G$  such that  $P' = g_0 P g_0^{-1}$ . Then any  $G$ -equivariant isomorphism  $G/P' \rightarrow G/P$  is written as  $\bar{g} \mapsto \bar{g} \bar{g}'_0$ ,  $\bar{g} \in G/P'$  for an element  $g'_0 \in g_0 N_G(P)$ . Since  $P$  is a parabolic subgroup of  $G$ , we have  $N_G(P) = P$ . This means that such a  $G$ -equivariant isomorphism is uniquely determined and we denote it by  $f_{P',P}$ . The inclusion map  $\mathfrak{m}'_{P',0}{}^{-1}(0_X) \rightarrow G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X')$  induces an isomorphism  $H^2(G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X'), \mathbf{C}) \cong H^2(G/P', \mathbf{C})$ . Similarly the inclusion map  $\mathfrak{m}'_{P,0}{}^{-1}(0_X) \rightarrow G \times^P (\mathfrak{n}(\mathfrak{p}) \times X')$  induces an isomorphism  $H^2(G \times^P (\mathfrak{n}(\mathfrak{p}) \times X'), \mathbf{C}) \cong H^2(G/P, \mathbf{C})$ . Then we have a commutative diagram

$$\begin{array}{ccc} H^2(G \times^P (\mathfrak{n}(\mathfrak{p}) \times X'), \mathbf{C}) & \xrightarrow{(\phi_{\tilde{w},0})^*} & H^2(G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X'), \mathbf{C}) \\ \cong \downarrow & & \cong \downarrow \\ H^2(G/P, \mathbf{C}) & \xrightarrow{(f_{P',P})^*} & H^2(G/P', \mathbf{C}) \end{array} \quad (23)$$

On the other hand,  $\phi_{\sigma,0}$  induces the identity map of  $X$  by definition. Hence we get a commutative diagram

$$\begin{array}{ccc} G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X') & \xrightarrow{\phi_{\sigma,0}} & G \times^P (\mathfrak{n}(\mathfrak{p}) \times X') \\ \mathfrak{m}'_{P',0} \downarrow & & \mathfrak{m}'_{P,0} \downarrow \\ X & \xrightarrow{id} & X \end{array} \quad (24)$$

The diagram is  $G$ -equivariant and it induces a  $G$ -equivariant isomorphism between  $\mathfrak{m}'_{P',0}(0_X) = G/P'$  and  $\mathfrak{m}'_{P,0}(0_X) = G/P$ . As explained above, this  $G$ -equivariant isomorphism is  $f_{P',P}$ . We then have a commutative diagram

$$\begin{array}{ccc} H^2(G \times^P (\mathfrak{n}(\mathfrak{p}) \times X'), \mathbf{C}) & \xrightarrow{(\phi_{\sigma,0})^*} & H^2(G \times^{P'} (\mathfrak{n}(\mathfrak{p}') \times X'), \mathbf{C}) \\ \cong \downarrow & & \cong \downarrow \\ H^2(G/P, \mathbf{C}) & \xrightarrow{(f_{P',P})^*} & H^2(G/P', \mathbf{C}) \end{array} \quad (25)$$

By the two diagrams of 2-nd cohomology, we conclude that  $(\phi_{\tilde{w},0})^* = (\phi_{\sigma,0})^*$ ; hence  $(\phi_{\tilde{w}})^* = (\phi_{\sigma})^*$ .  $\square$

**Claim 3.** *One can lift the element  $w \in W'$  to an element  $\tilde{w} \in \widetilde{W}'$  so that  $\phi_{\sigma} = \phi_{\tilde{w}}$ . In particular,  $\tilde{w} \in W_X$ .*

*Proof.* Take an arbitrary lift  $\tilde{w} \in \widetilde{W}'$  of  $w \in W$ . By Claim 2 we have a commutative diagram

$$\begin{array}{ccc} G \times^P (\mathfrak{r}(\mathfrak{p}) \times X') & \xrightarrow{\phi_{\tilde{w}} \circ \phi_{\sigma}^{-1}} & G \times^P (\mathfrak{r}(\mathfrak{p}) \times X') \\ \downarrow & & \downarrow \\ Z_{\mathfrak{l}, X'} & \xrightarrow{\tilde{w}_Z \circ \sigma_Z^{-1}} & Z_{\mathfrak{l}, X'} \\ \downarrow & & \downarrow \\ \mathfrak{k} & \xrightarrow{id} & \mathfrak{k} \end{array} \quad (26)$$

Take  $t \in \mathfrak{k}^{reg}$ . By the proof of Lemma 2, the fiber of  $G \times^P (\mathfrak{r}(\mathfrak{p}) \times X') \rightarrow \mathfrak{k}$  over  $t$  is isomorphic to  $G \times^L (t \times X')$ . Therefore the map  $\phi_{\tilde{w}} \circ \phi_{\sigma}^{-1}$  induces a  $G$ -equivariant Poisson automorphism

$$\phi_{\tilde{w},t} \circ \phi_{\sigma,t}^{-1} : G \times^L (t \times X') \cong G \times^L (t \times X').$$

Let us consider the map  $\mu_t : G \times^L (t \times X') \rightarrow G \cdot (t + \bar{O}') \subset \mathfrak{g}$ . Take  $x \in G \times^L (t \times X')$  general. Then the  $G$ -orbit  $G \cdot x$  is open dense in  $G \times^L (t \times X')$ . Moreover, let  $O_{\mu_t(x)} \subset \mathfrak{g}$  be the adjoint orbit containing  $\mu_t(x)$ . Then  $\mu_t|_{G \cdot x} : G \cdot x \rightarrow O_{\mu_t(x)} \subset \mathfrak{g}$  is the moment map for  $G \cdot x$ . Note that  $O_{\mu_t(x)}$  is also contained in  $G \times^L (t + \bar{O}')$  as the open dense  $G$ -orbit.  $\phi_{\tilde{w},t} \circ \phi_{\sigma,t}^{-1}$  induces a symplectic automorphism of  $G \cdot x$ . By [B-K], Theorem 6, it must be a deck transformation of  $G \cdot x \rightarrow O_{\mu_t(x)}$ . Let  $\tilde{O}'$  be the normalization of  $\bar{O}'$ . Then this implies that

$$\phi_{\tilde{w},t} \circ \phi_{\sigma,t}^{-1} \in \text{Aut}(G \times^L (t \times X') / G \times^L (t \times \tilde{O}')).$$

Therefore, we have

$$\phi_{\tilde{w}} \circ \phi_{\sigma}^{-1} \in \text{Aut}(Z_{\mathfrak{l}, X'} / Y_{\mathfrak{l}, O'}).$$

By taking a suitable lift  $\tilde{w} \in \widetilde{W}'$  of  $w \in W'$ , we have

$$\phi_{\tilde{w}} \circ \phi_{\sigma}^{-1} = id,$$

which means that  $\phi_\sigma = \phi_{\tilde{w}}$ . By definition,  $\phi_{\sigma,0}$  induces the identity map of  $Z_{\mathfrak{l},X',0}(= X)$ . Hence we have  $\tilde{w} \in W_X$ .  $\square$

Claim 2 and Claim 3 imply that  $W(X) \subset W_X$ .

We next prove that  $W_X \subset W(X)$ . We assume that  $W(X) \subsetneq W_X$  and derive a contradiction. Take an element  $w \in W_X - W(X)$ . For simplicity we write  $C_Q$  for the ample cone  $\text{Amp}(G \times^Q (\mathfrak{n} \times X'))$ . Since  $\text{Mov}(G \times^Q (\mathfrak{n} \times X'))$  is a fundamental domain for the  $W(X)$ -action, we can take an element  $w_0 \in W(X)$  so that  $w_0(w \cdot C_Q) = C_{w_0 w(Q)} \subset \text{Mov}(G \times^Q (\mathfrak{n} \times X'))$ . By changing  $w$  by  $w_0 w$ , we may assume, from the first, that

$$C_{w(Q)} \subset \text{Mov}(G \times^Q (\mathfrak{n} \times X')).$$

Since  $w \in W_X$ , we can take a lift  $\tilde{w} \in \tilde{W}'$  so that the the following diagram commutes

$$\begin{array}{ccc} G \times^Q (\mathfrak{n} \times X') & \xrightarrow{\phi_{\tilde{w},0}} & G \times^{Ad_w(Q)} (Ad_w(\mathfrak{n}) \times X') \\ \mathfrak{m}'_{Q,0} \downarrow & & \mathfrak{m}'_{w(Q),0} \downarrow \\ X & \xrightarrow{id} & X \end{array} \quad (27)$$

By the diagram  $\mathfrak{m}'_{Q,0}$  and  $\mathfrak{m}'_{w(Q),0}$  give the same  $\mathbf{Q}$ -factorization of  $X$ . This means that  $C_Q$  and  $C_{w(Q)}$  are the same chambers inside  $\text{Mov}(G \times^Q (\mathfrak{n} \times X'))$ . On the other hand, since  $w \in W'$  and  $w \neq id$ , we see that  $C_Q$  and  $C_{w(Q)}$  are different chambers. This is a contradiction.  $\square$

As in Proposition 10, we identify  $\mathfrak{k}$  with  $H^2(G \times^Q (\mathfrak{r} \times X'), \mathbf{C})$  and put  $\mathfrak{k}_{\mathbf{R}} = H^2(G \times^Q (\mathfrak{r} \times X'), \mathbf{R})$  by the identification. By Theorem 7, (1),  $\mathfrak{k}_{\mathbf{R}}$  is decomposed into finite number of chambers.

**Corollary 11.** *The Weyl group  $W_X$  acts on  $\mathfrak{k}_{\mathbf{R}}$  so that each chamber is sent to another chamber. The movable cone  $\text{Mov}(G \times^Q (\mathfrak{n} \times X'))$  is a fundamental domain of the  $W_X$ -action.  $\text{Mov}(G \times^Q (\mathfrak{n} \times X'))$  contains exactly  $\sharp \mathcal{S}(L)/|W_X|$  chambers. In particular,  $X$  has exactly  $\sharp \mathcal{S}(L)/|W_X|$  different  $\mathbf{Q}$ -factorial terminalizations.*

*Proof.* These follow from [Na 4], Main Theorem. We remark that the second statement was first pointed out in [BPW], Proposition 2.19.  $\square$

**Remark 12.**

We can compute the number  $N$  of the conjugacy classes in  $\mathcal{S}(L)$  by using the equivalence relation [Na 5], Definition 1, (ii) p. 95 in the set of marked Dynkin diagrams. Then, by [Na 1], Proposition 2.2, we have

$$\sharp \mathcal{S}(L) = N \cdot |W'|,$$

where  $W'$  is the finite group defined by the Equation (8). The group  $W'$  was studied in [Ho].

## § 2. The map $\rho_X$

**Proposition 13.** *The map  $\rho_X : \tilde{W}' \rightarrow \text{Aut}(X/\bar{O})$  is surjective.*

*Proof.* The group  $\text{Aut}(X/\bar{O})$  acts on the universal Poisson deformation  $Z_{\mathfrak{l},X'}/W_X \rightarrow \mathfrak{k}/W_X$  equivariantly.  $\text{Aut}(X/\bar{O})$  does not necessarily act effectively on the base space  $\mathfrak{k}/W_X$ . Let  $K \subset \text{Aut}(X/\bar{O})$  be the subgroup consisting of the elements which act trivially on  $\mathfrak{k}/W_X$ . Let us consider the surjection  $\pi : \tilde{W}' \rightarrow W'$ . By  $\pi$ ,  $W_X \subset \tilde{W}'$  is mapped isomorphically onto  $\pi(W_X) \subset W'$ . Then  $W'/\pi(W_X)$  acts on  $\mathfrak{k}/W_X$ . The injection  $\tilde{W}'/W_X \rightarrow \text{Aut}(X/\bar{O})$  induces a map  $W'/\pi(W_X) \rightarrow \text{Aut}(X/\bar{O})/K$ . We shall prove that this is an isomorphism. This is analogous to [Lo], Proposition 4.7. If we identify  $\mathfrak{g}$  with  $\mathfrak{g}^*$  by the Killing form,  $G \cdot (\mathfrak{r} + \bar{O}')$  is a  $\mathbf{C}^*$ -invariant subvariety of  $\mathfrak{g}^*$ . Look at the moment map

$$Z_{\mathfrak{l},X'}/W_X \rightarrow G \cdot (\mathfrak{r} + \bar{O}') \subset \mathfrak{g}^*$$

for the Poisson  $G$ -variety  $Z_{\mathfrak{l},X'}/W_X$ . The moment map is unique up to an element  $(\mathfrak{g}^* \otimes_{\mathbf{C}} \mathbf{C}[\mathfrak{k}/W_X])^G$ , where  $G$  acts trivially on  $\mathbf{C}[\mathfrak{k}/W_X]$ . Since  $(\mathfrak{g}^*)^G = \{0\}$ , we see that the moment map is unique. Since the  $\text{Aut}(X/\bar{O})$  action on  $Z_{\mathfrak{l},X'}/W_X$  commutes with the  $G$ -action, the moment map induces an injection

$$\mathfrak{g} \rightarrow \mathbf{C}[Z_{\mathfrak{l},X'}/W_X]^{\text{Aut}(X/\bar{O})}.$$

Let  $\bar{t}, \bar{t}' \in \mathfrak{k}^{reg}/W_X$  be  $\text{Aut}(X/\bar{O})/K$ -conjugate, and let  $t, t' \in \mathfrak{k}^{reg}$  be their lifts. Look at the map  $Z_{\mathfrak{l},X'}/W_X \rightarrow \mathfrak{k}/W_X$ . Then the fibers of the map over  $\bar{t}, \bar{t}'$  are respectively identified with  $Z_{\mathfrak{l},X',t}$  and  $Z_{\mathfrak{l},X',t'}$ . The  $\text{Aut}(X/\bar{O})$ -action on  $Z_{\mathfrak{l},X'}/W_X$  induces a  $G$ -equivariant Poisson automorphism  $Z_{\mathfrak{l},X',t} \cong Z_{\mathfrak{l},X',t'}$ . By composing the inclusion  $\mathfrak{g} \subset \mathbf{C}[Z_{\mathfrak{l},X'}/W_X]$  with the surjection to  $\mathbf{C}[Z_{\mathfrak{l},X',t}]$ , we get a map  $\mathfrak{g} \rightarrow \mathbf{C}[Z_{\mathfrak{l},X',t}]$ . Similarly we have a map  $\mathfrak{g} \rightarrow \mathbf{C}[Z_{\mathfrak{l},X',t'}]$ . Since  $\text{Aut}(X/\bar{O})$  acts on  $\mathfrak{g}$  trivially, we have a commutative diagram

$$\begin{array}{ccc} \mathbf{C}[Z_{\mathfrak{l},X',t}] & \longrightarrow & \mathbf{C}[Z_{\mathfrak{l},X',t'}] \\ \uparrow & & \uparrow \\ \mathfrak{g} & \xrightarrow{id} & \mathfrak{g} \end{array} \quad (28)$$

This induces a commutative diagram of the moment maps

$$\begin{array}{ccc} Z_{\mathfrak{l},X',t'} & \longrightarrow & Z_{\mathfrak{l},X',t} \\ \mu_{t'} \downarrow & & \mu_t \downarrow \\ \mathfrak{g}^* & \xrightarrow{id} & \mathfrak{g}^* \end{array} \quad (29)$$

In particular,  $\mu_t(Z_{\mathfrak{l},X',t})$  and  $\mu_{t'}(Z_{\mathfrak{l},X',t'})$  are the closure of the same coadjoint orbit of  $\mathfrak{g}^*$ . This means that  $t$  and  $t'$  are  $W'$ -conjugate.

We next prove that  $K = \text{Aut}((X')^0/O')$ . Since  $K$  acts on  $\mathfrak{k}/W_X$  trivially,  $K$  acts on  $Z_{\mathfrak{l},X'} = \mathfrak{k} \times_{\mathfrak{k}/W_X} Z_{\mathfrak{l},X'}/W_X$  in such a way that it acts on  $\mathfrak{k}$  trivially. Then an element of  $K$  determines a  $G$ -equivariant Poisson automorphism of every fiber of the map  $Z_{\mathfrak{l},X'} \rightarrow \mathfrak{k}$ . By the same argument as in Claim 3 in the proof of Proposition 10, we see that

$$K = \text{Aut}(Z_{\mathfrak{l},X'}/Y_{\mathfrak{l},O'}) = \text{Aut}((X')^0/O').$$

Finally we look at the commutative diagram

$$\begin{array}{ccc}
1 & & 1 \\
\downarrow & & \downarrow \\
\mathrm{Aut}((X')^0/O') & \longrightarrow & K \\
\downarrow & & \downarrow \\
\tilde{W}'/W_X & \longrightarrow & \mathrm{Aut}(X/\bar{O}) \\
\downarrow & & \downarrow \\
W'/\pi(W_X) & \longrightarrow & \mathrm{Aut}(X/\bar{O})/K \\
\downarrow & & \downarrow \\
1 & & 1
\end{array} \tag{30}$$

The 1-st horizontal map and the 3-rd horizontal map are both isomorphisms by the observations above. Then the middle horizontal map is an isomorphism.  $\square$

By Corollary 11, Remark 12 and Proposition 13, we have:

**Theorem 14.** *Let  $N$  be the number of conjugacy classes in  $\mathcal{S}(L)$ . Then the number of different  $\mathbf{Q}$ -factorial terminalizations of  $X$  is*

$$\frac{N \cdot |\mathrm{Aut}(X/\bar{O})|}{|\mathrm{Aut}((X')^0/O')|}.$$

We put  $\mathfrak{k}_{\mathbf{R}} := H^2(G \times^Q (\mathfrak{r} \times X'), \mathbf{R})$ . Then  $\mathfrak{k}_{\mathbf{R}} \otimes \mathbf{C} = \mathfrak{k}$ .  $\mathfrak{k}_{\mathbf{R}}$  is divided into finite number of chambers by finite linear spaces  $\{H_i\}_{i \in I}$  of codimension 1 (= hyperplanes), and each chamber corresponds to an ample cone of  $G \times^{Q'} (\mathfrak{r}' \times X')$  for  $Q' \in \mathcal{S}(L)$ . An element  $w$  of  $W'$  is a reflection if  $w$  has order 2 and the fixed points set of  $w$  on  $\mathfrak{k}_{\mathbf{R}}$  is a hyperplane. Such a hyperplane is contained in the set  $\{H_i\}_{i \in I}$ . Recall that there is an exact sequence

$$1 \rightarrow \mathrm{Aut}((X')^0/O') \rightarrow \tilde{W}' \rightarrow W' \rightarrow 1$$

The composite  $\mathrm{Aut}((X')^0/O') \rightarrow \tilde{W}' \xrightarrow{\rho_X} \mathrm{Aut}(X/\bar{O})$  is an injection. Put  $Y := Y_{\mathfrak{t}, O', 0}$ . Then  $\pi : X \rightarrow \bar{O}$  factorizes as  $X \rightarrow Y \rightarrow \bar{O}$  and  $\mathrm{Aut}((X')^0/O')$  is identified with  $\mathrm{Aut}(X/Y) \subset \mathrm{Aut}(X/\bar{O})$  by the injection. Take a reflection  $w \in W'$  and its lift  $\tilde{w} \in \tilde{W}'$ . Consider the subgroup  $G(w) \subset \tilde{W}'$  generated by  $\tilde{w}$  and  $\mathrm{Aut}((X')^0/O')$ . Note that  $G(w)$  is independent of the choice of the lift  $\tilde{w}$ . There are two cases.

- (a)  $G(w) \cap W_X \neq \{1\}$ .
- (b)  $G(w) \cap W_X = \{1\}$ .

In the case (a), we have  $\rho_X(G(w)) = \mathrm{Aut}(X/Y)$ .

**Lemma 15.** *Assume that  $\mathfrak{g}$  is classical and assume that  $X \rightarrow \bar{O}$  is the finite covering associated with the universal covering of  $O$ . In the case (b),  $\rho_X(G(w)) = \mathrm{Aut}(X/Y_w)$  for a finite covering  $Y_w \rightarrow \bar{O}$  with the following properties.*

$\pi$  is factorized into intermediate coverings

$$X \rightarrow Y \xrightarrow{f_w} Y_w \rightarrow \bar{O}$$

with  $\deg(f_w) = 2$ .

*Proof.* Let  $H \subset \mathfrak{k}_{\mathbf{R}}$  be the fixed points set of  $w$ . Take two chambers which share a codimension 1 face  $F$  lying on  $H$ . These chambers correspond to  $\text{Amp}(G \times^Q (\mathfrak{n} \times X'))$  and  $\text{Amp}(G \times^{Q'} (\mathfrak{n}' \times X'))$  for some  $Q, Q' \in \mathcal{S}(L)$ . The two parabolic subgroups are related by  $Q = \text{Ad}_w(Q')$ . The face  $F$  determines two birational morphisms respectively from  $G \times^Q (\mathfrak{n} \times X')$  and  $G \times^{Q'} (\mathfrak{n}' \times X')$  to the same target space. The target space is described as follows. Take a parabolic subgroup  $\bar{Q}$  of  $G$  in such a way that  $Q \subset \bar{Q}$ ,  $Q' \subset \bar{Q}$  and  $b_2(G/\bar{Q}) = b_2(G/Q) - 1 = b_2(G/Q') - 1$ . Let  $\bar{L}$  be the Levi part of  $\bar{Q}$  such that  $T \subset \bar{L}$ . We put  $\bar{l} := \text{Lie}(\bar{L})$  and  $\bar{\mathfrak{n}} := \mathfrak{n}(\bar{\mathfrak{q}})$ . Define  $O_{\bar{l}} := \text{Ind}_{\bar{l}}^{\bar{L}}(O')$ . Let  $X_{\bar{l}}$  be the Stein factorization of the composite

$$\bar{L} \times^{\bar{L} \cap Q} (\mathfrak{n}(\bar{l} \cap \mathfrak{q}) \times X') \rightarrow \bar{L} \times^{\bar{L} \cap Q} (\mathfrak{n}(\bar{l} \cap \mathfrak{q}) + \bar{O}') \rightarrow \bar{O}_{\bar{l}}.$$

Then the target space is  $G \times^{\bar{Q}} (\bar{\mathfrak{n}} \times X_{\bar{l}})$ .

There is a commutative diagram

$$\begin{array}{ccccc} G \times^{Q'} (\mathfrak{n}' \times X') & \xrightarrow{\mathbf{m}'_{Q', \bar{Q}, 0}} & G \times^{\bar{Q}} (\bar{\mathfrak{n}} \times X_{\bar{l}}) & \xleftarrow{\mathbf{m}'_{\bar{Q}, \bar{Q}, 0}} & G \times^Q (\mathfrak{n} \times X') \\ \downarrow & & \downarrow & & \downarrow \\ G \times^{Q'} (\mathfrak{n}' + \bar{O}') & \xrightarrow{\mathbf{m}_{Q', \bar{Q}, 0}} & G \times^{\bar{Q}} (\bar{\mathfrak{n}} + \bar{O}_{\bar{l}}) & \xleftarrow{\mathbf{m}_{\bar{Q}, \bar{Q}, 0}} & G \times^Q (\mathfrak{n} + \bar{O}') \end{array} \quad (31)$$

such that  $\mathbf{m}'_{Q', \bar{Q}, 0}$  and  $\mathbf{m}'_{\bar{Q}, \bar{Q}, 0}$  are the birational morphisms determined by  $F$ .

Note that  $w \in W(\bar{L})$  for the Levi part  $\bar{L}$  of  $\bar{Q}$ . Then, since  $w \in \bar{Q}$ , we have a commutative diagram

$$\begin{array}{ccc} G \times^{Q'} (\mathfrak{n}' \times X') & \xrightarrow{\phi_{\bar{w}, 0}} & G \times^Q (\mathfrak{n} \times X') \\ \downarrow & & \downarrow \\ G \times^{\bar{Q}} (\bar{\mathfrak{n}} + \bar{O}_{\bar{l}})^n & \xrightarrow{id} & G \times^{\bar{Q}} (\bar{\mathfrak{n}} + \bar{O}_{\bar{l}})^n \end{array} \quad (32)$$

We put here

$$Y_w := \text{Spec } \Gamma(G \times^{\bar{Q}} (\bar{\mathfrak{n}} + \bar{O}_{\bar{l}})^n, \mathcal{O})$$

Since  $Y = \text{Spec } \Gamma(G \times^Q (\mathfrak{n} + \bar{O}')^n, \mathcal{O})$ , the map  $\mathbf{m}_{Q, \bar{Q}, 0}$  induces a map  $f_w : Y \rightarrow Y_w$  and the finite cover  $Y \rightarrow \bar{O}$  factors through  $Y_w$ . The map  $\mathbf{m}_{Q, \bar{Q}, 0}$  is a generalized Springer map determined by an *induction* of type I or of type II (see [Na, Part I, §2, §3]). Then  $\deg(\mathbf{m}_{Q, \bar{Q}, 0}) = 1$  or  $\deg(\mathbf{m}_{Q, \bar{Q}, 0}) = 2$  by [ibid, (2.3.1), (2.3.2), (3.6.1) and (3.6.2)]. We used here the assumption that  $\mathfrak{g}$  is classical. In particular,  $\deg(f_w) = 1, 2$ . The map  $\phi_{\bar{w}, 0} : G \times^{Q'} (\mathfrak{n}' \times X') \rightarrow G \times^P (\mathfrak{n} \times X')$  induces automorphisms of  $X, Y$  and  $Y_w$ . Hence



we get a commutative diagram

$$\begin{array}{ccc}
G \times^{Q'} (\mathfrak{n}' \times X') & \xrightarrow{\phi_{\tilde{w},0}} & G \times^Q (\mathfrak{n} \times X') \\
\downarrow & & \downarrow \\
X & \xrightarrow{\tilde{w}_{Z,0}} & X \\
\downarrow & & \downarrow \\
Y & \xrightarrow{\tilde{w}_{Y,0}} & Y \\
\downarrow & & \downarrow \\
Y_w & \xrightarrow{id} & Y_w
\end{array} \tag{33}$$

If  $\tilde{w}_{Y,0} = id$ , then  $\tilde{w}_{Z,0}$  is an element of  $\text{Aut}(X/Y)$ . This means that  $G(w) \cap W_X \neq \{1\}$ , which contradicts the assumption. Therefore  $\tilde{w}_{Y,0} \neq id$ . Then  $\rho_X(\tilde{w}) \in \text{Aut}(X/Y_w)$ , but  $\rho_X(\tilde{w}) \notin \text{Aut}(X/Y)$ . In particular,  $\text{Aut}(X/Y_w) \neq \text{Aut}(X/Y)$ ; hence,  $f_w : Y \rightarrow Y_w$  is a double cover. This means that  $\rho_X(G(w)) = \text{Aut}(X/Y_w)$ .  $\square$

**Remark 16.**

(1) The proof of Lemma 15 shows that if Case (b) occurs, then  $\deg \mathfrak{m}_{Q,\bar{Q},0} = 2$ .

(2) If  $\mathfrak{m}'_{Q,\bar{Q},0}$  is divisorial, then Case (a) occurs. If  $\mathfrak{m}'_{Q,\bar{Q},0}$  is a small birational map (i.e.  $\text{Codim Exc}(\mathfrak{m}'_{Q,\bar{Q},0}) \geq 2$ ), then Case (b) occurs.

In fact, if  $\mathfrak{m}'_{Q,\bar{Q},0}$  is divisorial, then  $\mathfrak{m}_{Q,\bar{Q},0}$  is birational by the next Proposition 17. Then  $Y_w = Y$ ; hence  $\tilde{w}_{Y,0} = id$ . If  $\mathfrak{m}'_{Q,\bar{Q},0}$  is small, then  $\mathfrak{m}'_{Q',\bar{Q},0}$  is also small. Therefore  $\text{Amp}(G \times^{Q'} (\mathfrak{n}' \times X'))$  is contained in  $\text{Mov}(G \times^Q (\mathfrak{n} \times X'))$ . Since  $Q = \text{Ad}_w(Q')$  and  $w \neq 1$ , the two cones  $\text{Amp}(G \times^Q (\mathfrak{n} \times X'))$  and  $\text{Amp}(G \times^{Q'} (\mathfrak{n}' \times X'))$  determine different chambers inside  $\text{Mov}(G \times^Q (\mathfrak{n} \times X'))$ . We prove that  $\tilde{w}_{Y,0} \neq id$  for any lift  $\tilde{w}$  of  $w$ . In fact, if  $\tilde{w}_{Y,0} = id$ , then one can assume that  $\tilde{w}_{Z,0} = id$  by replacing  $\tilde{w}$  by a suitable lift of  $w$ . Then we have a commutative diagram

$$\begin{array}{ccc}
G \times^{Q'} (\mathfrak{n}' \times X') & \xrightarrow{\phi_{\tilde{w},0}} & G \times^Q (\mathfrak{n} \times X') \\
\downarrow & & \downarrow \\
X & \xrightarrow{id} & X
\end{array} \tag{34}$$

By the diagram  $G \times^{Q'} (\mathfrak{n}' \times X')$  and  $G \times^Q (\mathfrak{n} \times X')$  give the same  $\mathbf{Q}$ -factorial terminalization of  $X$ . This means that

$$\text{Amp}(G \times^Q (\mathfrak{n} \times X')) = \text{Amp}(G \times^{Q'} (\mathfrak{n}' \times X'))$$

This is a contradiction.  $\square$

Take a reflection  $w \in W'$ . Let  $H \subset \mathfrak{t}_{\mathbf{R}}$  be the fixed hyperplane of  $w$ . As in the proof of Lemma 15, we find  $Q \in \mathcal{S}(L)$  in such a way that, if we put  $Q' := \text{Ad}_w(Q)$ , then the

ample cones  $\text{Amp}(G \times^Q (\mathfrak{n} \times X'))$  and  $\text{Amp}(G \times^{Q'} (\mathfrak{n}' \times X'))$  share a codimension 1 face  $F$  lying on  $H$ . The codimension 1 face  $F$  determines a birational contraction map of  $G \times^Q (\mathfrak{n} \times X')$ . The birational map is described as follows. There is a parabolic subgroup  $\bar{Q}$  of  $G$  so that  $Q \subset \bar{Q}$  and  $b_2(G/\bar{Q}) = b_2(G/Q) - 1$ . Let  $\bar{\mathfrak{l}}$  be the Levi part of  $\bar{\mathfrak{q}}$ . There are a nilpotent orbit  $O_{\bar{\mathfrak{l}}} \subset \bar{\mathfrak{l}}$ , a finite covering  $X_{\bar{\mathfrak{l}}} \rightarrow \bar{O}_{\bar{\mathfrak{l}}}$  and a commutative diagram

$$\begin{array}{ccc} G \times^Q (\mathfrak{n} \times X') & \xrightarrow{\mathfrak{m}'_{Q,\bar{Q},0}} & G \times^{\bar{Q}} (\bar{\mathfrak{n}} \times X_{\bar{\mathfrak{l}}}) \\ \pi'_Q \downarrow & & \bar{\pi} \downarrow \\ G \times^Q (\mathfrak{n} + \bar{O}') & \xrightarrow{\mathfrak{m}_{Q,\bar{Q},0}} & G \times^{\bar{Q}} (\bar{\mathfrak{n}} + \bar{O}_{\bar{\mathfrak{l}}}) \end{array} \quad (35)$$

such that  $\mathfrak{m}'_{Q,\bar{Q},0}$  is a birational map. The map  $\mathfrak{m}'_{Q,\bar{Q},0}$  is the desired birational contraction map. Note that  $\mathfrak{m}_{Q,\bar{Q},0}$  is not necessarily a birational map.

**Proposition 17.** *Assume that  $\mathfrak{g}$  is a classical simple Lie algebra and assume that  $X \rightarrow \bar{O}$  is the finite covering associated with the universal covering of  $O$ .*

- (1) *If  $\mathfrak{m}'_{Q,\bar{Q},0}$  is a divisorial birational map, then  $\mathfrak{m}_{Q,\bar{Q},0}$  is a divisorial birational map.*
- (2) *If  $\mathfrak{m}'_{Q,\bar{Q},0}$  is a small birational map, then  $\deg \mathfrak{m}_{Q,\bar{Q},0} = 2$ .*

*Proof.* (2) follows from Remark 16. In fact, if  $\mathfrak{m}'_{Q,\bar{Q},0}$  is a small birational map, then Case (b) occurs by Remark 16, (2). By Remark 16, (1) we then have  $\deg \mathfrak{m}_{Q,\bar{Q},0} = 2$ . In the remainder we prove (1). To prove (1) we only have to show that  $\mathfrak{m}_{Q,\bar{Q},0}$  is birational if  $\mathfrak{m}'_{Q,\bar{Q},0}$  is a divisorial birational map. In fact, if  $\mathfrak{m}'_{Q,\bar{Q},0}$  and  $\mathfrak{m}_{Q,\bar{Q},0}$  are both birational, then one is divisorial if and only if the other is divisorial.

When  $\mathfrak{g} = \mathfrak{sl}(n)$ , the map  $\mathfrak{m}_{Q,\bar{Q},0}$  is always birational. When  $\mathfrak{g} = \mathfrak{sp}(2n)$  or  $\mathfrak{so}(n)$ , we prove that  $\text{Codim Exc}(\mathfrak{m}'_{Q,\bar{Q},0}) \geq 2$  if  $\deg(\mathfrak{m}_{Q,\bar{Q},0}) \geq 2$ .

Case (a):  $\mathfrak{g} = \mathfrak{sp}(2n)$

In [Part 1], §2, we constructed a  $\mathbf{Q}$ -factorial terminalization of  $X$ . For such a  $\mathbf{Q}$ -factorial terminalization,  $O'$  is contained in a simple factor  $\mathfrak{l}_0$  of the Levi part  $\bar{\mathfrak{l}}$ , and  $\mathfrak{l}_0$  is of type  $C_k$ . The Jordan type  $\mathbf{p} = [d^{r_d}, (d-1)^{r_{d-1}}, \dots, 2^{r_2}, 1^{r_1}]$  of  $O'$  is then a partition of  $2k$  which satisfies the following conditions

- (i)  $r_i$  is even for each odd  $i$ ,
- (ii)  $r_i \neq 0$  for each even  $i$ , and
- (iii)  $r_i \neq 2$  for each even  $i$ .

Moreover,  $X' \rightarrow \bar{O}'$  is the finite cover associated with the universal covering of  $O'$ . Since the core  $(X', O')$  is independent of the choice of a  $\mathbf{Q}$ -factorial terminalization of  $X$ , we can start with this  $(X', O')$ .

There is a unique simple factor  $\bar{\mathfrak{l}}_0$  of the Levi part  $\bar{\mathfrak{l}}$  such that  $\mathfrak{l}_0 \subset \bar{\mathfrak{l}}_0$ . Then  $\bar{\mathfrak{l}}_0$  is again of type  $C$  and

$$O_{\bar{\mathfrak{l}}} = \text{Ind}_{\bar{\mathfrak{l}}_0}^{\bar{\mathfrak{l}}}(O').$$

If  $\deg(\mathfrak{m}_{Q,\bar{Q},0}) \geq 2$ , then it is an induction of type II (cf. [Part 1], §2) and  $\deg(\mathfrak{m}_{Q,\bar{Q},0}) = 2$ . Let  $\bar{\mathbf{p}} := [\bar{d}^{\bar{r}_{\bar{d}}}, (\bar{d}-1)^{\bar{r}_{\bar{d}-1}}, \dots, 2^{\bar{r}_2}, 1^{\bar{r}_1}]$  be the Jordan type of  $O_{\bar{\mathfrak{l}}}$ . Two partitions  $\mathbf{p}$  and  $\bar{\mathbf{p}}$  are related as follows. There is an even member  $i$  of  $\bar{\mathbf{p}}$  with  $\bar{r}_i = 2$ , and  $\bar{\mathbf{p}}$  can be written as

$$\mathbf{p} = [(\bar{d}-2)^{\bar{r}_{\bar{d}}}, \dots, i^{\bar{r}_i+2}, (i-1)^{\bar{r}_{i+1}+2+\bar{r}_{i-1}}, (i-2)^{\bar{r}_{i-2}}, \dots, 1^{\bar{r}_1}].$$

It is checked that  $\bar{\mathbf{p}}$  satisfies the conditions (i) and (ii) by using the fact that  $\mathbf{p}$  satisfies (i), (ii) and (iii). Let  $b$  be the number of distinct even members of  $\mathbf{p}$ . Recall that  $\pi_1(O') = (\mathbf{Z}/2\mathbf{Z})^{\oplus b}$ . This means that  $\deg \pi'_Q = 2^b$ . Since  $\deg(\mathbf{m}_{Q,\bar{Q},0}) = 2$  and  $\mathbf{m}'_{Q,\bar{Q},0}$  is birational, we see that  $\deg \bar{\pi} = 2^{b+1}$ . Let  $\bar{\mathbf{p}}$  be the number of distinct even members of  $\bar{\mathbf{p}}$ . Then  $\pi_1(O_{\bar{l}}) = (\mathbf{Z}/2\mathbf{Z})^{\oplus \bar{b}}$ . By the description of  $\mathbf{p}$  above, we have  $\bar{b} = b + 1$ . These facts imply that  $X_{\bar{l}} \rightarrow \bar{O}_{\bar{l}}$  is the finite covering associated with the universal covering of  $O_{\bar{l}}$ . Since the Jordan type  $\bar{\mathbf{p}}$  of  $O_{\bar{l}}$  satisfies the conditions (i) and (ii), we see that  $\text{Codim}_{X_{\bar{l}}} \text{Sing}(X_{\bar{l}}) \geq 4$  by [Part 1], Proposition (2,3). This means that  $\text{Codim Exc}(\mathbf{m}'_{Q,\bar{Q},0}) \geq 2$ .

Case (b):  $\mathfrak{g} = \mathfrak{so}(n)$

In [Part 1], §3, we constructed a  $\mathbf{Q}$ -factorial terminalization of  $X$ . Assume that  $\mathfrak{l}$  has simple factors  $\mathfrak{so}(k)$ ,  $\mathfrak{sl}(k_1)$ , ..., and  $\mathfrak{sl}(k_s)$ . There are three cases for  $(X', O')$ . The first case is when  $O$  is rather odd, and the remaining cases are when  $O$  is not rather odd.

Case (b-1):  $k_1, \dots, k_s$  are all even and

$$O' = O_{\mathbf{p}} \times O_{\lfloor \frac{k_1}{2} \rfloor} \times \dots \times O_{\lfloor \frac{k_s}{2} \rfloor} \subset \mathfrak{so}(k) \oplus \mathfrak{sl}(k_1) \oplus \dots \oplus \mathfrak{sl}(k_s)$$

Here  $\mathbf{p} := [d^{r_d}, (d-1)^{r_{d-1}}, \dots, 2^{r_2}, 1^{r_1}]$  is a *rather odd* partition of  $k$  with the following conditions.

- (i)  $r_i$  is even for each even  $i$
  - (ii) For any couple  $(i, j)$  of adjacent members,  $|i - j| \leq 4$ . Moreover  $|i - j| = 4$  occurs only when  $i$  and  $j$  are both odd. The smallest member of  $\mathbf{p}$  is smaller than 4.
- Moreover, if we put  $m_i := r_d + r_{d-1} + \dots + r_i$ , then  $\frac{k_1}{2}, \dots, \frac{k_s}{2}$  respectively coincide with some  $m_i$ 's.

We shall prove that  $\deg(\mathbf{m}_{Q,\bar{Q},0}) \geq 2$  never happens. In fact, there are two possibilities for  $\bar{\mathfrak{l}}$ .

$$\bar{\mathfrak{l}} = \mathfrak{so}(k + 2k_t) \oplus \mathfrak{sl}(k_1) \oplus \dots \oplus \mathfrak{sl}(k_{t-1}) \oplus \mathfrak{sl}(k_{t+1}) \oplus \dots \mathfrak{sl}(k_s) \oplus \mathfrak{z}(\bar{\mathfrak{l}})$$

$$\bar{\mathfrak{l}} = \mathfrak{so}(k) \oplus \mathfrak{sl}(k_i + k_j) \oplus \mathfrak{sl}(k_1) \oplus \dots \oplus \mathfrak{sl}(\check{k}_i) \oplus \dots \oplus \mathfrak{sl}(\check{k}_j) \oplus \dots \oplus \mathfrak{sl}(k_s) \oplus \mathfrak{z}(\bar{\mathfrak{l}}).$$

In the first case, one can write  $k_t = 2m_i$  for some  $i$ . If we put

$$\bar{\mathbf{p}} := [(d+4)^{r_d}, (d+3)^{r_{d-1}}, \dots, (i+4)^{r_i}, (i-1)^{r_{i-1}}, \dots, 2^{r_2}, 1^{r_1}],$$

then

$$O_{\bar{\mathfrak{l}}} = O_{\bar{\mathbf{p}}} \times O_{\lfloor \frac{k_1}{2} \rfloor} \times \dots \times O_{\lfloor \frac{k_{t-1}}{2} \rfloor} \times O_{\lfloor \frac{k_{t+1}}{2} \rfloor} \times \dots \times O_{\lfloor \frac{k_s}{2} \rfloor}.$$

In the second case,

$$O_{\bar{\mathfrak{l}}} = O_{\mathbf{p}} \times O_{\lfloor \frac{k_i+k_j}{2} \rfloor} \times O_{\lfloor \frac{k_1}{2} \rfloor} \times \dots \times \check{O}_{\lfloor \frac{k_i}{2} \rfloor} \times \dots \times \check{O}_{\lfloor \frac{k_j}{2} \rfloor} \times \dots \times O_{\lfloor \frac{k_s}{2} \rfloor}.$$

One can check that, in both cases,  $\mathbf{m}_{Q,\bar{Q},0}$  is birational.

Case (b-2) :  $O' = O_{\mathbf{p}} \subset \mathfrak{so}(k)$ . Here  $\mathbf{p} := [d^{r_d}, (d-1)^{r_{d-1}}, \dots, 2^{r_2}, 1^{r_1}]$  is a partition of  $k$  with the following conditions.

- (i)  $r_i$  is even for each even  $i$ ,
- (ii)  $r_i \neq 0$  for each odd  $i$ ,

- (iii)  $r_i \neq 2$  for each odd  $i$ , and
- (iv)  $r_i \geq 3$  for some odd  $i$ .

Moreover,  $X' \rightarrow \bar{O}'$  is the finite cover associated with the universal covering of  $O$ .

For some  $t$ , we have

$$\bar{\mathbf{l}} = so(k + 2k_t) \oplus sl(k_1) \oplus \dots \oplus sl(k_{t-1}) \oplus sl(k_{t+1}) \oplus \dots \oplus sl(k_s) \oplus \mathfrak{z}(\bar{\mathbf{l}})$$

and

$$O_{\bar{\mathbf{l}}} = \text{Ind}_{so(k)}^{so(k+2k_t)}(O') \subset so(k + 2k_t).$$

If  $\deg(\mathbf{m}_{Q, \bar{Q}, 0}) \geq 2$ , then it is an induction of type II and  $\deg(\mathbf{m}_{Q, \bar{Q}, 0}) = 2$ . Let  $\bar{\mathbf{p}} := [\bar{d}^{\bar{r}_{\bar{d}}}, (\bar{d} - 1)^{\bar{d}_{\bar{d}-1}}, \dots, 2^{\bar{r}_2}, 1^{\bar{r}_1}]$  be the Jordan type of  $O_{\bar{\mathbf{l}}}$ . Two partitions  $\mathbf{p}$  and  $\bar{\mathbf{p}}$  are related as follows. There is an odd member  $i$  of  $\bar{\mathbf{p}}$  with  $\bar{r}_i = 2$ , and  $\mathbf{p}$  can be written as

$$\mathbf{p} = [(\bar{d} - 2)^{\bar{r}_{\bar{d}}}, \dots, i^{\bar{r}_i+2}, (i - 1)^{\bar{r}_{i+1}+2+\bar{r}_{i-1}}, (i - 2)^{\bar{r}_{i-2}}, \dots, 1^{\bar{r}_1}].$$

Here  $k_t = \bar{r}_{\bar{d}} + \bar{r}_{\bar{d}-1} + \dots + \bar{r}_{i+1} + 1$ . It is checked that  $\bar{\mathbf{p}}$  satisfies the conditions (i), (ii) and (iv) by using the fact that  $\mathbf{p}$  satisfies (i), (ii), (iii) and (iv). Let  $a$  be the number of distinct odd members of  $\mathbf{p}$ . Since  $\mathbf{p}$  is not rather odd and  $a > 0$  by the condition (iv), we have  $\pi_1(O') = (\mathbf{Z}/2\mathbf{Z})^{\oplus a-1}$ . This means that  $\deg \pi'_Q = 2^{a-1}$ . Since  $\deg(\mathbf{m}_{Q, \bar{Q}, 0}) = 2$  and  $\mathbf{m}'_{Q, \bar{Q}, 0}$  is birational, we see that  $\deg \bar{\pi} = 2^a$ . Let  $\bar{a}$  be the number of distinct odd members of  $\bar{\mathbf{p}}$ . By the description of  $\bar{\mathbf{p}}$  above,  $\bar{a} = a + 1$ . Then  $\pi_1(O_{\bar{\mathbf{l}}}) = (\mathbf{Z}/2\mathbf{Z})^{\oplus \bar{a}-1}$  because  $\bar{\mathbf{p}}$  is not rather odd. These facts imply that  $X_{\bar{\mathbf{l}}} \rightarrow \bar{O}_{\bar{\mathbf{l}}}$  is the finite covering associated with the universal covering of  $O_{\bar{\mathbf{l}}}$ . Since  $\bar{\mathbf{p}}$  is not rather odd and satisfies the conditions (i) and (ii), we have  $\text{Codim}_{X_{\bar{\mathbf{l}}}} \text{Sing}(X_{\bar{\mathbf{l}}}) \geq 4$  by [Part 1], Proposition (3.4). This means that  $\text{Codim Exc}(\mathbf{m}'_{Q, \bar{Q}, 0}) \geq 2$ .

Case (b-3):  $O' = O_{\mathbf{p}} \subset so(k)$ . Here  $\mathbf{p} := [d^{r_d}, (d - 1)^{r_{d-1}}, \dots, 2^{r_2}, 1^{r_1}]$  is a partition of  $k$  with the following conditions.

- (i)  $r_i$  is even for each even  $i$ , and
- (ii)  $r_i = 1$  for each odd  $i$

In particular,  $\mathbf{p}$  is rather odd. Let  $a$  be the number of distinct odd members of  $\mathbf{p}$ . Then we have an exact sequence

$$1 \rightarrow \mathbf{Z}/2\mathbf{Z} \rightarrow \pi_1(O') \rightarrow (\mathbf{Z}/2\mathbf{Z})^{\oplus \min(a-1, 0)} \rightarrow 1.$$

The finite cover  $X' \rightarrow \bar{O}'$  is induced from the etale covering of  $O'$  determined by the surjection  $\pi_1(O') \rightarrow (\mathbf{Z}/2\mathbf{Z})^{\oplus \min(a-1, 0)}$ . In particular, we have  $\deg(\pi'_Q) = 2^{\min(a-1, 0)}$ . For some  $t$ , we have

$$\bar{\mathbf{l}} = so(k + 2k_t) \oplus sl(k_1) \oplus \dots \oplus sl(k_{t-1}) \oplus sl(k_{t+1}) \oplus \dots \oplus sl(k_s) \oplus \mathfrak{z}(\bar{\mathbf{l}})$$

and

$$O_{\bar{\mathbf{l}}} = \text{Ind}_{so(k)}^{so(k+2k_t)}(O') \subset so(k + 2k_t).$$

If  $\deg(\mathbf{m}_{Q, \bar{Q}, 0}) \geq 2$ , then it is an induction of type II and  $\deg(\mathbf{m}_{Q, \bar{Q}, 0}) = 2$ . Let  $\bar{\mathbf{p}} := [\bar{d}^{\bar{r}_{\bar{d}}}, (\bar{d} - 1)^{\bar{d}_{\bar{d}-1}}, \dots, 2^{\bar{r}_2}, 1^{\bar{r}_1}]$  be the Jordan type of  $O_{\bar{\mathbf{l}}}$ . Two partitions  $\mathbf{p}$  and  $\bar{\mathbf{p}}$  are related as follows. There is an odd member  $i$  of  $\bar{\mathbf{p}}$  with  $\bar{r}_i = 2$ , and  $\mathbf{p}$  can be written as

$$\mathbf{p} = [(\bar{d} - 2)^{\bar{r}_{\bar{d}}}, \dots, i^{\bar{r}_i+2}, (i - 1)^{\bar{r}_{i+1}+2+\bar{r}_{i-1}}, (i - 2)^{\bar{r}_{i-2}}, \dots, 1^{\bar{r}_1}].$$

Here  $k_t = \bar{r}_{\bar{d}} + \bar{r}_{\bar{d}-1} + \dots + \bar{r}_{i+1} + 1$ . It is checked that  $\bar{\mathbf{p}}$  satisfies the following conditions

- (i)  $\bar{r}_i$  is even for each even  $i$ , and
- (iii)  $\bar{r}_i \neq 0$  for each odd  $i$ .

We may assume that  $a > 0$ . In fact, if  $a = 0$ , then  $\mathbf{p} = [0]$  because  $\mathbf{p}$  satisfies the condition (ii). In such a case  $\mathbf{m}_{Q,\bar{Q},0}$  is birational by [Part 1], Claim (3.6.2). Therefore,  $\deg(\pi'_Q) = 2^{\min(a-1,0)} = 2^{a-1}$ . Since  $\deg(\mathbf{m}_{Q,\bar{Q},0}) = 2$  and  $\mathbf{m}'_{Q,\bar{Q},0}$  is birational, we see that  $\deg \bar{\pi} = 2^a$ . Let  $\bar{a}$  be the number of distinct odd members of  $\bar{\mathbf{p}}$ . By the description of  $\bar{\mathbf{p}}$  above,  $\bar{a} = a + 1$ . Then  $\pi_1(O_{\bar{\mathbf{l}}}) = (\mathbf{Z}/2\mathbf{Z})^{\oplus \bar{a}-1}$  because  $\bar{\mathbf{p}}$  is not rather odd (because  $\bar{r}_i = 2$  for some odd  $i$ ). These facts imply that  $X_{\bar{\mathbf{l}}} \rightarrow \bar{O}_{\bar{\mathbf{l}}}$  is the finite covering associated with the universal covering of  $O_{\bar{\mathbf{l}}}$ . Since  $\bar{\mathbf{p}}$  is not rather odd and satisfies the conditions (i) and (iii), we have  $\text{Codim}_{X_{\bar{\mathbf{l}}}} \text{Sing}(X_{\bar{\mathbf{l}}}) \geq 4$  by [Part 1], Proposition (3.4). This means that  $\text{Codim Exc}(\mathbf{m}'_{Q,\bar{Q},0}) \geq 2$ .  $\square$

The homomorphism  $\rho_X : \tilde{W}' \rightarrow \text{Aut}(X/\bar{O})$  induces a homomorphism

$$\overline{\rho_X} : W' \rightarrow \text{Aut}(X/\bar{O})/\text{Aut}((X')^0/O').$$

**Corollary 18.** *Assume that  $\mathfrak{g}$  is classical and assume that  $X \rightarrow \bar{O}$  is the finite covering associated with the universal covering of  $O$ . For a reflection  $w \in W'$ , the map  $\mathbf{m}_{Q,\bar{Q},0}$  is a divisorial birational map or a generically finite surjective map of degree 2. In the first case,  $\overline{\rho_X}(w) = 1$ , and in the latter case  $\overline{\rho_X}(w) \neq 1$ .*

When  $\mathfrak{g}$  is classical,  $\deg(\mathbf{m}_{Q,\bar{Q},0}) = 1$  or  $\deg(\mathbf{m}_{Q,\bar{Q},0}) = 2$ . Moreover, one can check which case occurs by using the criterion [Part 1], (2.3.1), (2.3.2), (3.6.1), (3.6.2). Then the Weyl group  $W_X$  is the subgroup of  $W'$  generated by all reflections  $w$  for which  $\deg(\mathbf{m}_{Q,\bar{Q},0}) = 1$ .

### § 3. Examples

**Example 19.** ([Part 1], Example (2.5), (1)) :

Let us consider the nilpotent orbit  $O_{[6^2,4^2]} \subset sp(20)$  and let  $\pi : X \rightarrow \bar{O}_{[6^2,4^2]}$  be the finite covering associated with the universal covering of  $O_{[6^2,4^2]}$ . We have  $\deg(\pi) = 4$  because the partition  $[6^2, 4^2]$  has exactly two distinct even members. We can construct a  $\mathbf{Q}$ -factorial terminalization of  $X$  by the induction step

$$([1^4], sp(4)) \xrightarrow{\text{Type II}} (O_{[3^2,2^2]}, sp(10)) \xrightarrow{\text{Type II}} (O_{[4^2,2^2]}, sp(12)) \xrightarrow{\text{Type I}} (O_{[6^2,4^2]}, sp(20)).$$

We consider the three generalized Springer maps

$$Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + \bar{O}_{[4^2,2^2]}) \rightarrow \bar{O}_{[6^2,4^2]},$$

$$Sp(20) \times^{Q_{4,1,10,1,4}} (\mathbf{n}(\mathbf{q}_{4,1,10,1,4}) + \bar{O}_{[3^2,2^2]}) \rightarrow Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + \bar{O}_{[4^2,2^2]}),$$

and

$$Sp(20) \times^{Q_{4,1,3,4,3,1,4}} \mathbf{n}(\mathbf{q}_{4,1,3,4,3,1,4}) \rightarrow Sp(20) \times^{Q_{4,1,10,1,4}} (\mathbf{n}(\mathbf{q}_{4,1,10,1,4}) + \bar{O}_{[3^2,2^2]}).$$

Composing these 3 maps together, we have a generically finite map of degree 4

$$Sp(20) \times^{Q_{4,1,3,4,3,1,4}} \mathbf{n}(\mathbf{q}_{4,1,3,4,3,1,4}) \rightarrow \bar{O}_{[6^2, 4^2]}.$$

This map factors through  $X$  and  $Sp(20) \times^{Q_{4,1,3,4,3,1,4}} \mathbf{n}(\mathbf{q}_{4,1,3,4,3,1,4})$  gives a crepant resolution of  $X$ . We shall calculate how many different crepant resolutions  $X$  has.

The parabolic subgroup  $Q_{4,1,3,4,3,1,4}$  corresponds to the marked Dynkin diagram

$$\circ - \circ - \circ - \bullet - \bullet - \circ - \circ - \bullet - \circ \Leftarrow \circ$$

Let  $L$  be the Levi part of  $Q_{4,1,3,4,3,1,4}$ . One can read from the marked Dynkin diagram that the semi simple part of  $\mathfrak{l}$  is  $A_3 + A_2 + C_2$ . By [Ho], p.71, we see that

$$W' = (\mathbf{Z}/2\mathbf{Z})^{\oplus 3}.$$

$\mathcal{S}(L)$  contains exactly 6 conjugacy classes. Other 5 conjugacy classes correspond to the following marked Dynkin diagrams (cf. Remark 11)

$$\bullet - \circ - \circ - \circ - \bullet - \circ - \circ - \bullet - \circ \Leftarrow \circ$$

$$\bullet - \circ - \circ - \bullet - \circ - \circ - \circ - \bullet - \circ \Leftarrow \circ$$

$$\circ - \circ - \bullet - \bullet - \circ - \circ - \circ - \bullet - \circ \Leftarrow \circ$$

$$\circ - \circ - \bullet - \circ - \circ - \circ - \bullet - \bullet - \circ \Leftarrow \circ$$

$$\circ - \circ - \circ - \bullet - \circ - \circ - \bullet - \bullet - \circ \Leftarrow \circ$$

It follows from this fact that

$$N = 6, \quad \#\mathcal{S}(L) = 6|W'| = 48.$$

In our case  $X' = \bar{O}' = \{0\}$ . On the other hand,  $\text{Aut}(X/\bar{O}_{[6^2, 4^2]}) = (\mathbf{Z}/2\mathbf{Z})^{\oplus 2}$  (cf. [Part 1], Proposition (2.1)). Then, by Theorem 13, we have

$$\#\{\text{crepant resolutions of } X\} = N \cdot |\text{Aut}(X/\bar{O}_{[6^2, 4^2]})| = 6 \cdot 4 = 24.$$

We next closely study the map  $\rho_X : \tilde{W}' \rightarrow \text{Aut}(X/\bar{O}_{[6^2, 4^2]})$ . Look at the first marked Dynkin diagram

$$\circ - \circ - \circ - \bullet - \bullet - \circ - \circ - \underset{\alpha}{\bullet} - \circ \Leftarrow \circ$$

Let  $Q'_{4,1,3,4,3,1,4}$  be a parabolic subgroup of  $Sp(20)$  obtained from  $Q_{4,1,3,4,3,1,4}$  by twisting at  $\alpha$ . Then  $Q_{4,1,3,4,3,1,4} = \text{Ad}_{w_1}(Q'_{4,1,3,4,3,1,4})$  for a (uniquely determined) element  $w_1 \in W'$ . Let  $\bar{Q}_{4,1,10,1,4}$  be the parabolic subgroup of  $Sp(20)$  with flag type  $(4, 1, 10, 1, 4)$  such that  $Q_{4,1,3,4,3,1,4} \subset \bar{Q}_{4,1,10,1,4}$  and  $Q'_{4,1,3,4,3,1,4} \subset \bar{Q}_{4,1,10,1,4}$ . Put  $Q_I = Q_{4,1,3,4,3,1,4}$ ,

$Q'_I = Q'_{4,1,3,4,3,1,4}$  and  $\bar{Q}_I = \bar{Q}_{4,1,10,1,4}$ . Let  $\mathbf{n}_I$ ,  $\mathbf{n}'_I$  and  $\bar{\mathbf{n}}_I$  be respectively the nilradicals of  $Q_I$ ,  $Q'_I$  and  $\bar{Q}_I$ . Then we have a commutative diagram

$$\begin{array}{ccccc} Sp(20) \times^{Q'_I} \mathbf{n}'_I & \xrightarrow{\mathfrak{m}'_{Q'_I, \bar{Q}_I, 0}} & Sp(20) \times^{\bar{Q}_I} (\bar{\mathbf{n}}_I \times X_{[3^2, 2^2]}) & \xleftarrow{\mathfrak{m}'_{Q_I, \bar{Q}_I, 0}} & Sp(20) \times^{Q_I} \mathbf{n}_I \\ \downarrow & & \downarrow & & \downarrow \\ Sp(20) \times^{Q'_I} \mathbf{n}'_I & \xrightarrow{\mathfrak{m}_{Q'_I, \bar{Q}_I, 0}} & Sp(20) \times^{\bar{Q}_I} (\bar{\mathbf{n}}_I + \bar{O}_{[3^2, 2^2]}) & \xleftarrow{\mathfrak{m}_{Q_I, \bar{Q}_I, 0}} & Sp(20) \times^{Q_I} \mathbf{n}_I \end{array} \quad (36)$$

The horizontal maps on the first row are birational maps, and the horizontal maps on the second row have degree 2 because they are determined by a type II induction. Then the birational maps on the first row are both small by Proposition 17. We put

$$Y_{w_1} := \text{Spec } \Gamma(Sp(20) \times^{\bar{Q}_I} (\bar{\mathbf{n}}_I + \bar{O}_{[3^2, 2^2]}), \mathcal{O}).$$

$Y_{w_1}$  is a double covering of  $\bar{O}_{[6^2, 4^2]}$ .

Next let  $Q_{4,3,1,4,1,3,4}$  be the parabolic subgroup corresponding to the marked Dynkin diagram

$$\circ \text{---} \circ \text{---} \circ \text{---} \bullet \text{---} \circ \text{---} \circ \text{---} \bullet \text{---} \bullet \text{---} \circ \text{---} \circ \text{---} \beta$$

Let  $Q'_{4,3,1,4,1,3,4}$  be a parabolic subgroup of  $Sp(20)$  obtained from  $Q_{4,3,1,4,1,3,4}$  by twisting at  $\beta$ . Then  $Q_{4,3,1,4,1,3,4} = Ad_{w_2}(Q'_{4,3,1,4,1,3,4})$  for a (uniquely determined) element  $w_2 \in W'$ . Let  $\bar{Q}_{4,3,6,3,4}$  be the parabolic subgroup of  $Sp(20)$  with flag type  $(4, 3, 6, 3, 4)$  such that  $Q_{4,3,1,4,1,3,4} \subset \bar{Q}_{4,3,6,3,4}$  and  $Q'_{4,3,1,4,1,3,4} \subset \bar{Q}_{4,3,6,3,4}$ . Put  $Q_{II} = Q_{4,3,1,4,1,3,4}$ ,  $Q'_{II} = Q'_{4,3,1,4,1,3,4}$  and  $\bar{Q}_{II} = \bar{Q}_{4,3,6,3,4}$ . Let  $\mathbf{n}_{II}$ ,  $\mathbf{n}'_{II}$  and  $\bar{\mathbf{n}}_{II}$  be respectively the nilradicals of  $Q_{II}$ ,  $Q'_{II}$  and  $\bar{Q}_{II}$ . Then we have a commutative diagram

$$\begin{array}{ccccc} Sp(20) \times^{Q'_{II}} \mathbf{n}'_{II} & \xrightarrow{\mathfrak{m}'_{Q'_{II}, \bar{Q}_{II}, 0}} & Sp(20) \times^{\bar{Q}_{II}} (\bar{\mathbf{n}}_{II} \times X_{[2^2, 1^2]}) & \xleftarrow{\mathfrak{m}'_{Q_{II}, \bar{Q}_{II}, 0}} & Sp(20) \times^{Q_{II}} \mathbf{n}_{II} \\ \downarrow & & \downarrow & & \downarrow \\ Sp(20) \times^{Q'_{II}} \mathbf{n}'_{II} & \xrightarrow{\mathfrak{m}_{Q'_{II}, \bar{Q}_{II}, 0}} & Sp(20) \times^{\bar{Q}_{II}} (\bar{\mathbf{n}}_{II} + \bar{O}_{[2^2, 1^2]}) & \xleftarrow{\mathfrak{m}_{Q_{II}, \bar{Q}_{II}, 0}} & Sp(20) \times^{Q_{II}} \mathbf{n}_{II} \end{array} \quad (37)$$

The horizontal maps on the first row are birational maps, and the horizontal maps on the second row have degree 2 because they are determined by a type II induction. Then the birational maps on the first row are both small by Proposition 17. We put

$$Y_{w_2} := \text{Spec } \Gamma(Sp(20) \times^{\bar{Q}_{II}} (\bar{\mathbf{n}}_{II} + \bar{O}_{[2^2, 1^2]}), \mathcal{O}).$$

$Y_{w_2}$  is a double covering of  $\bar{O}_{[6^2, 4^2]}$ .

In our case,  $Y = X$  and  $W' = W'$ . By the next lemma  $\text{Aut}(X/Y_{w_1})$  and  $\text{Aut}(X/Y_{w_2})$  are mutually different, index 2, subgroups of  $\text{Aut}(X/\bar{O}_{[6^2, 4^2]}) = (\mathbf{Z}/2\mathbf{Z})^{\oplus 2}$ . By Lemma 14,  $\text{Im}(\rho_X)$  contains both  $\text{Aut}(X/Y_{w_i})$ . This reproves that

$$\text{Im}(\rho_X) = \text{Aut}(X/\bar{O}_{[6^2, 4^2]}).$$

In particular,

$$W_X = \text{Ker}(\rho_X) = \mathbf{Z}/2\mathbf{Z}.$$

**Lemma 20.** *The double coverings  $\pi_1 : Y_{\omega_1} \rightarrow \bar{O}_{[6^2, 4^2]}$  and  $\pi_2 : Y_{\omega_2} \rightarrow \bar{O}_{[6^2, 4^2]}$  are different coverings in the function field  $\mathbf{C}(X)$  of  $X$ .*

*Proof.*  $\bar{O}_{[6^2, 4^2]}$  has  $A_3$ -surface singularity along the orbit  $O_{[6^2, 4, 2^2]}$ . This means that the transverse slice  $S \subset \bar{O}_{[6^2, 4^2]}$  for  $O_{[6^2, 4, 2^2]}$  is a complex analytic germ of an  $A_3$ -singularity. We claim that  $\pi_1^{-1}(S)$  is isomorphic to a disjoint union of two copies of  $S$ , but  $\pi_2^{-1}(S)$  is isomorphic to an  $A_1$ -germ. We first look at the generalized Springer map

$$f : Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + \bar{O}_{[4^2, 2^2]}) \rightarrow \bar{O}_{[6^2, 4^2]}$$

Since this map is determined by the induction of type I, it is a birational map. The closed subvariety  $Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + \bar{O}_{[4^2, 2, 1^2]}) \subset Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + \bar{O}_{[4^2, 2^2]})$  is mapped onto the subvariety  $\bar{O}_{[6^2, 4, 2^2]} \subset \bar{O}_{[6^2, 4^2]}$  by  $f$ . The map

$$Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + \bar{O}_{[4^2, 2, 1^2]}) \rightarrow \bar{O}_{[6^2, 4, 2^2]}$$

is a generically finite map of degree 2 because it is determined by the type II induction. Note that  $Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + \bar{O}_{[4^2, 2^2]})$  has  $A_1$ -surface singularity along  $Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + O_{[4^2, 2, 1^2]})$ . Therefore we conclude that  $f^{-1}(S)$  is a crepant partial resolution of the  $A_3$ -singularity with one exceptional curve  $C \cong \mathbf{P}^1$  and  $f^{-1}(S)$  has two  $A_1$ -singularities at two points  $p_+, p_- \in C$ . We next look at the generalized Springer maps

$$g_1 : Sp(20) \times^{\bar{Q}_{4,1,10,1,4}} (\mathbf{n}(\bar{\mathbf{q}}_{4,1,10,1,4}) + \bar{O}_{[3^2, 2^2]}) \rightarrow Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + \bar{O}_{[4^2, 2^2]})$$

$$g_2 : Sp(20) \times^{\bar{Q}_{4,3,6,3,4}} (\mathbf{n}(\bar{\mathbf{q}}_{4,3,6,3,4}) + \bar{O}_{[2^2, 1^2]}) \rightarrow Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + \bar{O}_{[4^2, 2^2]})$$

and observe  $g_1^{-1}(p)$  and  $g_2^{-1}(p)$  for a point  $p \in C$ . The map  $g_1$  can be described as

$$\begin{aligned} Sp(20) \times^{\bar{Q}_{4,1,10,1,4}} (\mathbf{n}(\bar{\mathbf{q}}_{4,1,10,1,4}) + \bar{O}_{[3^2, 2^2]}) &= Sp(20) \times^{Q_{4,12,4}} (Q_{4,12,4} \times^{\bar{Q}_{4,1,10,1,4}} (\mathbf{n}(\bar{\mathbf{q}}_{4,1,10,1,4}) + \bar{O}_{[3^2, 2^2]})) \\ &= Sp(20) \times^{Q_{4,12,4}} \{ \mathbf{n}(\mathbf{q}_{4,12,4}) \times (L(Q_{4,12,4}) \times^{L(Q_{4,12,4}) \cap \bar{\mathbf{q}}_{4,1,10,1,4}} (\mathbf{n}(\mathbf{l}(\mathbf{q}_{4,12,4}) \cap \bar{\mathbf{q}}_{4,1,10,1,4}) + \bar{O}_{[3^2, 2^2]})) \} \\ &\longrightarrow Sp(20) \times^{Q_{4,12,4}} \{ \mathbf{n}(\mathbf{q}_{4,12,4}) \times L(Q_{4,12,4}) \cdot (\mathbf{n}(\mathbf{l}(\mathbf{q}_{4,12,4}) \cap \bar{\mathbf{q}}_{4,1,10,1,4}) + \bar{O}_{[3^2, 2^2]}) \} \\ &= Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) + \bar{O}_{[4^2, 2^2]}) \end{aligned}$$

Therefore  $g_1$  is obtained from the map

$$\begin{aligned} L(Q_{4,12,4}) \times^{L(Q_{4,12,4}) \cap \bar{\mathbf{q}}_{4,1,10,1,4}} (\mathbf{n}(\mathbf{l}(\mathbf{q}_{4,12,4}) \cap \bar{\mathbf{q}}_{4,1,10,1,4}) + \bar{O}_{[3^2, 2^2]}) \\ \rightarrow L(Q_{4,12,4}) \cdot (\mathbf{n}(\mathbf{l}(\mathbf{q}_{4,12,4}) \cap \bar{\mathbf{q}}_{4,1,10,1,4}) + \bar{O}_{[3^2, 2^2]}) \end{aligned}$$

by taking  $Sp(20) \times^{Q_{4,12,4}} (\mathbf{n}(\mathbf{q}_{4,12,4}) \times \cdot)$ . This map is nothing but the generalized Springer map

$$\bar{g}_1 : Sp(12) \times^{Q_{1,10,1}} (\mathbf{n}(\mathbf{q}_{1,10,1}) + \bar{O}_{[3^2, 2^2]}) \rightarrow \bar{O}_{[4^2, 2^2]}.$$

Similarly,  $g_2$  is obtained from the generalized Springer map

$$\bar{g}_2 : Sp(12) \times^{Q_{3,6,3}} (\mathbf{n}(\mathbf{q}_{3,6,3}) + \bar{O}_{[2^2, 1^2]}) \rightarrow \bar{O}_{[4^2, 2^2]}.$$



Write  $p \in Sp(20) \times^{Q_{4,12,4}} (\mathfrak{n}(\mathfrak{q}_{4,12,4}) + \bar{O}_{[4^2,2^2]})$  as  $p = [h, z + x]$  with  $h \in Sp(20)$ ,  $z \in \mathfrak{n}(\mathfrak{q}_{4,12,4})$  and  $x \in \bar{O}_{[4^2,2^2]}$ . Then

$$g_1^{-1}(p) = [h, z + \bar{g}_1^{-1}(x)] \cong \bar{g}_1^{-1}(x), \quad g_2^{-1}(p) = [h, z + \bar{g}_2^{-1}(x)] \cong \bar{g}_2^{-1}(x).$$

**Case 1:**  $p = p_+$  or  $p = p_-$ .

In this case we have  $x \in O_{[4^2,2,1^2]}$ . In order to see the fibers  $\bar{g}_1^{-1}(x)$  and  $\bar{g}_2^{-1}(x)$ , we introduce a basis  $\{e(i, j)\}$  of the symplectic vector space  $\mathbf{C}^{12}$  so that

(a)  $\{e(i, j)\}$  is a Jordan basis for  $x$ , i.e.  $x \cdot e(i, j) = e(i - 1, j)$  for  $i > 1$  and  $x \cdot e(1, j) = 0$ .

(b)  $\langle e(i, j), e(q, r) \rangle \neq 0$  if and only if  $q = d_j - i + 1$  and  $r = \beta(j)$ . Here  $\beta$  is a permutation of  $\{1, 2, \dots, s\}$  such that  $\beta^2 = id$ ,  $d_{\beta(j)} = d_j$ , and  $\beta(j) \neq j$  if  $d_j$  is odd (cf. [S-S], p.259, see also [?], 5.1).

In our case  $d_1 = d_2 = 4$ ,  $d_3 = 2$  and  $d_4 = d_5 = 1$ . Hence  $\beta(4) = 5$ ,  $\beta(5) = 4$  and  $\beta(3) = 3$ . After a suitable basis change of the subspace  $\Sigma_{1 \leq i \leq 4, 1 \leq j \leq 2} \mathbf{C}e(i, j)$ , we may also assume that  $\beta(1) = 2$  and  $\beta(2) = 1$ .

e(1,5)			
e(1,4)			
e(1,3)	e(2,3)		
e(1,2)	e(2,2)	e(3,2)	e(4,2)
e(1,1)	e(2,1)	e(3,1)	e(4,1)

If we put  $F_0 = \mathbf{C}e(1, 1)$ , then  $F_0 \subset F_0^\perp$  is an isotropic flag such that  $x \cdot F_0 = 0$ ,  $x \cdot \mathbf{C}^{12} \subset F_0^\perp$ , and  $x$  is an endomorphism of  $F_0^\perp/F_0$  with Jordan type  $[3^2, 2, 1^2]$ . On the other hand,  $F'_0 := \mathbf{C}e(1, 2)$  and  $(F'_0)^\perp$  also satisfy the same property. These two flags are only those flags of type  $(1, 10, 1)$  which have the properties

- (i)  $x \cdot F = 0$ ,  $x \cdot \mathbf{C}^{12} \subset F^\perp$ , and
- (ii)  $x$  is an endomorphism of  $F^\perp/F = \mathbf{C}^{10}$  with  $x \in \bar{O}_{[3^2,2^2]}$ .

This means that  $\bar{g}_1^{-1}(p)$  consists of two points. Moreover, these two points are lying on the locus

$$Sp(12) \times^{Q_{1,10,1}} (\mathfrak{n}(\mathfrak{q}_{1,10,1}) + O_{[3^2,2,1^2]}) \subset Sp(12) \times^{Q_{1,10,1}} (\mathfrak{n}(\mathfrak{q}_{1,10,1}) + \bar{O}_{[3^2,2^2]})$$

Note that  $Sp(12) \times^{Q_{1,10,1}} (\mathfrak{n}(\mathfrak{q}_{1,10,1}) + \bar{O}_{[3^2,2^2]})$  has  $A_1$  singularities along  $Sp(12) \times^{Q_{1,10,1}} (\mathfrak{n}(\mathfrak{q}_{1,10,1}) + O_{[3^2,2,1^2]})$ .

We next consider the fiber  $\bar{g}_2^{-1}(p)$ . If we put  $F_0 := \mathbf{C}e(1, 1) \oplus \mathbf{C}e(1, 2) \oplus \mathbf{C}e(1, 3)$ , then  $F_0 \subset F_0^\perp$  is an isotropic flag such that  $x \cdot F_0 = 0$ ,  $x \cdot \mathbf{C}^6 \subset F_0^\perp$ , and  $x$  is an endomorphism of  $F_0^\perp/F_0$  with Jordan type  $[2^2, 1^2]$ . This is the only isotropic flag of type  $(3, 6, 3)$  which has the properties

- (i)  $x \cdot F = 0$ ,  $x \cdot \mathbf{C}^6 \subset F^\perp$ , and
- (ii)  $x$  is an endomorphism of  $F^\perp/F = \mathbf{C}^6$  with  $x \in \bar{O}_{[2^2,1^2]}$ .

Therefore  $\bar{g}_2^{-1}(p)$  consists of one point.

**Case 2:**  $p \neq p_+, p_-$ .

In this case  $x \in O_{[4^2, 2^2]}$ . Since  $\bar{g}_1$  and  $\bar{g}_2$  are both generalized Springer map made from type II inductions, we conclude that  $\bar{g}_1^{-1}(p)$  and  $\bar{g}_2^{-1}(p)$  respectively consist of two points.

By these observations, we see that

1.  $(f \circ g_1)^{-1}(S)$  is isomorphic to a disjoint union of two copies of  $f^{-1}(S)$ .
2.  $(f \circ g_2)^{-1}(S)$  is isomorphic to the minimal resolution of an  $A_1$ -singularity. The map  $(f \circ g_2)^{-1}(S) \rightarrow f^{-1}(S)$  is a double covering only ramified at  $p_+$  and  $p_-$ . Here  $g_2^{-1}(C)$  corresponds to the exceptional curve of the minimal resolution of an  $A_1$ -singularity.

Let us consider the commutative diagrams

$$\begin{array}{ccc}
Sp(20) \times^{\bar{Q}_{4,1,10,1,4}} (\mathfrak{n}(\bar{\mathfrak{q}}_{4,1,10,1,4}) + \bar{O}_{[3^2, 2^2]}) & \xrightarrow{g_1} & Sp(20) \times^{Q_{4,12,4}} (\mathfrak{n}(\mathfrak{q}_{4,12,4}) + \bar{O}_{[4^2, 2^2]}) \\
f_1 \downarrow & & f \downarrow \\
Y_{\omega_1} & \xrightarrow{\pi_1} & \bar{O}_{[6^2, 4^2]}
\end{array} \tag{38}$$

$$\begin{array}{ccc}
Sp(20) \times^{\bar{Q}_{4,3,6,3,4}} (\mathfrak{n}(\bar{\mathfrak{q}}_{4,3,6,3,4}) + \bar{O}_{[2^2, 1^2]}) & \xrightarrow{g_2} & Sp(20) \times^{Q_{4,12,4}} (\mathfrak{n}(\mathfrak{q}_{4,12,4}) + \bar{O}_{[4^2, 2^2]}) \\
f_2 \downarrow & & f \downarrow \\
Y_{\omega_2} & \xrightarrow{\pi_2} & \bar{O}_{[6^2, 4^2]}
\end{array} \tag{39}$$

Here  $(f \circ g_1)^{-1}(S)$  is mapped to a disjoint union of two copies of an  $A_3$ -singularity by  $f_1$ . On the other hand,  $(f \circ g_2)^{-1}(S)$  is mapped to an  $A_1$ -singularity by  $f_2$ .  $\square$

**Example 21.** ([Part 1], Example (3.10), (2)):


Let  $X$  be a finite covering of  $\bar{O}_{[11^3, 3^2, 1]} \subset so(40)$  associated with the universal covering of  $O_{[11^3, 3^2, 1]}$ . By inductions of type I and of type II

$$[11^3, 3^2, 1] \xleftarrow{Type I} [9^3, 3^2, 1] \xleftarrow{Type I} [7^3, 3^2, 1] \xleftarrow{Type I} [5^3, 3^2, 1] \xleftarrow{Type II} [3^3, 2^2, 1]$$

we finally get a partition  $[3^3, 2^2, 1]$  of 14. Let  $\bar{Q} \subset SO(40)$  be a parabolic subgroup stabilizing an isotropic flag of type  $(3, 3, 3, 4, 14, 4, 3, 3, 3)$  and put  $Q := \rho_{40}^{-1}(\bar{Q})$  for  $\rho_{40} : Spin(40) \rightarrow SO(40)$ . Let us consider the nilpotent orbit  $O_{[3^3, 2^2, 1]} \subset so(14)$ . Let  $X_{[3^3, 2^2, 1]} \rightarrow \bar{O}_{[3^3, 2^2, 1]}$  be a finite covering associated with the universal covering of  $O_{[3^3, 2^2, 1]}$ . Then

$$Spin(40) \times^Q (\mathfrak{n} \times X_{[3^3, 2^2, 1]})$$

is a  $\mathbf{Q}$ -factorial terminalization of  $X$ . Fix a Borel subgroup  $B$  of  $Spin(40)$  with  $B \subset Q$  and a maximal torus  $T \subset B$ . We denote by  $\{\alpha_i\}_{1 \leq i \leq 20}$  the simple roots determined by  $B$ . Then the parabolic subgroup  $Q$  corresponds to the marked Dynkin diagram of type  $D_{20}$ :



Put  $\mathfrak{t} := \text{Lie}(T)$  and define  $\mathfrak{t}_{\mathbf{R}}^* \subset \mathfrak{t}^*$  to be the real vector space spanned by all simple roots. Let  $I := \{\alpha_i\}_{1 \leq i \leq 20, i \neq 3, 6, 9, 13}$  and denote by  $\Phi_I$  the root subsystem of  $\Phi$  generated by  $I$ . The semisimple part of the Levi part  $L$  of  $\text{Spin}(40)$  is  $3A_2 + A_3 + D_7$  and its root system is nothing but  $\Phi_I$ . By [Ho, p.71] we have

$$W' = W(B_3) \times W(B_1).$$


We shall explicitly describe  $W'$  in terms of twists.

We have

$$H^2(\text{Spin}(40)/Q, \mathbf{R}) \cong \text{Hom}_{\text{alg. gp}}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R} = \{x \in \mathfrak{t}_{\mathbf{R}}^*; (x, \alpha_i) = 0 \text{ for all } \alpha_i \in I\},$$

where  $(\cdot, \cdot)$  is the Killing form on  $\mathfrak{t}_{\mathbf{R}}^*$ . By the Killing form we identify  $\mathfrak{t}_{\mathbf{R}}^*$  with  $\mathfrak{t}_{\mathbf{R}}$  (= dual space of  $\mathfrak{t}_{\mathbf{R}}^*$ ). Thus  $\text{Hom}_{\text{alg. gp}}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R}$  can be identified with the subspace of  $\mathfrak{t}_{\mathbf{R}}$  consisting of the elements which kill all  $\alpha_i \in I$ .

We twist the  $Q$  by  $\alpha_3$ :



Then the twist by  $\alpha_3$  induces an automorphism  $T_3$  of  $\mathfrak{t}_{\mathbf{R}}^*$ . In particular, we have

$$T_3(\alpha_3) = -\alpha_3, \quad T_3(\alpha_6) = \sum_{1 \leq i \leq 6} \alpha_i, \quad T_3(\alpha_9) = \alpha_9, \quad T_3(\alpha_{13}) = \alpha_{13}$$

We define an automorphism  $T_3^t$  of  $\mathfrak{t}_{\mathbf{R}}$  by

$$\langle T_3^t(\phi), \cdot \rangle = \langle \phi, T_3(\cdot) \rangle, \quad \phi \in \mathfrak{t}_{\mathbf{R}}$$

Since  $T_3(\sum_{\alpha_i \in I} \mathbf{R}\alpha_i) = \sum_{\alpha_i \in I} \mathbf{R}\alpha_i$ , the automorphism  $T_3^t$  induces an automorphism of  $\text{Hom}_{\text{alg. gp}}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R} \subset \mathfrak{t}_{\mathbf{R}}$ . If we write

$$\mathfrak{t}_{\mathbf{R}}^* / \sum_{\alpha_i \in I} \alpha_i = \mathbf{R}\alpha_3 \oplus \mathbf{R}\alpha_6 \oplus \mathbf{R}\alpha_9 \oplus \mathbf{R}\alpha_{13},$$

then

$$\text{Hom}_{\text{alg. gp}}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R} = \mathbf{R}\alpha_3^* \oplus \mathbf{R}\alpha_6^* \oplus \mathbf{R}\alpha_9^* \oplus \mathbf{R}\alpha_{13}^*.$$

Then we have

$$T_3^t(\alpha_3^*) = -\alpha_3^* + \alpha_6^*, \quad T_3^t(\alpha_6^*) = \alpha_6^*, \quad T_3^t(\alpha_9^*) = \alpha_9^*, \quad T_3^t(\alpha_{13}^*) = \alpha_{13}^*.$$

We next twist  $Q$  by  $\alpha_6$ :

$$\begin{array}{c}
\text{---} \bullet \xrightarrow{\alpha_3} \circ \xrightarrow{\alpha_4} \circ \xrightarrow{\alpha_5} \bullet \xrightarrow{\alpha_6} \circ \xrightarrow{\alpha_7} \circ \xrightarrow{\alpha_8} \bullet \text{---} \\
\text{---} \bullet \xrightarrow{\sum_{3 \leq i \leq 8} \alpha_i} \circ \xrightarrow{-\alpha_8} \circ \xrightarrow{-\alpha_7} \bullet \xrightarrow{-\alpha_6} \circ \xrightarrow{-\alpha_5} \circ \xrightarrow{-\alpha_4} \bullet \xrightarrow{\sum_{4 \leq i \leq 9} \alpha_i} \text{---}
\end{array}$$

Let  $T_6$  be the automorphism of  $\mathfrak{t}_{\mathbf{R}}^*$  determined by the twist by  $\alpha_6$ . Then

$$T_6(\alpha_3) = \sum_{3 \leq i \leq 8} \alpha_i, \quad T_6(\alpha_6) = -\alpha_6, \quad T_6(\alpha_9) = \sum_{4 \leq i \leq 9} \alpha_i, \quad T_6(\alpha_{13}) = \alpha_{13}.$$

By using this we see that the automorphism  $T_6^t$  of  $\text{Hom}_{alg.gp}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R}$  is given by

$$T_6^t(\alpha_3^*) = \alpha_3^*, \quad T_6^t(\alpha_6^*) = \alpha_3^* - \alpha_6^* + \alpha_9^*, \quad T_6^t(\alpha_9^*) = \alpha_9^*, \quad T_6^t(\alpha_{13}^*) = \alpha_{13}^*.$$

We finally twist  $Q$  by  $\alpha_{13}$ :

$$\begin{array}{c}
\text{---} \bullet \xrightarrow{\alpha_9} \circ \xrightarrow{\alpha_{10}} \circ \xrightarrow{\alpha_{11}} \circ \xrightarrow{\alpha_{12}} \bullet \xrightarrow{\alpha_{13}} \circ \xrightarrow{\alpha_{14}} \circ \xrightarrow{\alpha_{15}} \circ \xrightarrow{\alpha_{16}} \circ \xrightarrow{\alpha_{17}} \circ \xrightarrow{\alpha_{18}} \circ \begin{array}{l} \nearrow \alpha_{19} \\ \searrow \alpha_{20} \end{array} \\
\text{---} \bullet \xrightarrow{-\alpha_{10}} \circ \xrightarrow{-\alpha_{11}} \circ \xrightarrow{-\alpha_{12}} \circ \xrightarrow{-\alpha_{13}} \bullet \xrightarrow{-\alpha_{14}} \circ \xrightarrow{-\alpha_{15}} \circ \xrightarrow{-\alpha_{16}} \circ \xrightarrow{-\alpha_{17}} \circ \begin{array}{l} \nearrow -\alpha_{20} \\ \searrow -\alpha_{18} \\ \searrow -\alpha_{19} \end{array} \\
\alpha_9 + 2 \sum_{10 \leq i \leq 18} \alpha_i + \alpha_{19} + \alpha_{20}
\end{array}$$

Let  $T_{13}$  be the automorphism of  $\mathfrak{t}_{\mathbf{R}}^*$  determined by the twist by  $\alpha_{13}$ . Then

$$\begin{aligned}
T_{13}(\alpha_3) &= \alpha_3, \quad T_{13}(\alpha_6) = \alpha_6, \\
T_{13}(\alpha_9) &= \alpha_9 + 2 \sum_{10 \leq i \leq 18} \alpha_i + \alpha_{19} + \alpha_{20}, \quad T_{13}(\alpha_{13}) = -\alpha_{13}.
\end{aligned}$$


By using this we see that the automorphism  $T_{13}^t$  of  $\text{Hom}_{alg.gp}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R}$  is given by

$$T_{13}^t(\alpha_3^*) = \alpha_3^*, \quad T_{13}^t(\alpha_6^*) = \alpha_6^*, \quad T_{13}^t(\alpha_9^*) = \alpha_9^*, \quad T_{13}^t(\alpha_{13}^*) = 2\alpha_9^* - \alpha_{13}^*.$$


We need one more twisting transformation to describe  $W'$ . In order to find the twisting, we must start with the parabolic subgroup  $Q'$  obtained from  $Q$  by the twist at  $\alpha_9$ . The parabolic subgroup  $Q'$  corresponds to the marked Dynkin diagram

$$\circ \text{---} \circ \text{---} \bullet \xrightarrow{\alpha_3} \circ \text{---} \circ \text{---} \bullet \xrightarrow{\sum_{6 \leq i \leq 12} \alpha_i} \circ \text{---} \circ \text{---} \bullet \xrightarrow{-\alpha_9} \circ \text{---} \bullet \xrightarrow{\sum_{7 \leq i \leq 13} \alpha_i} \begin{array}{l} \nearrow \circ \\ \searrow \circ \end{array}$$

We illustrate the marked Dynkin diagram in more details around the vertex  $\sum_{7 \leq i \leq 13} \alpha_i$ .



If we twist  $Q'$  by  $\sum_{7 \leq i \leq 13} \alpha_i$ , then we get



Let  $T_{9+13}$  be the automorphism of  $\mathfrak{t}_{\mathbf{R}}^*$  determined by the twist by  $\sum_{7 \leq i \leq 13} \alpha_i$ . Then

$$T_{9+13}(\alpha_3) = \alpha_3, \quad T_{9+13}\left(\sum_{6 \leq i \leq 12} \alpha_i\right) = \sum_{6 \leq i \leq 12} \alpha_i,$$

$$T_{9+13}(-\alpha_9) = \alpha_9 + 2 \sum_{10 \leq i \leq 18} \alpha_i + \alpha_{19} + \alpha_{20}, \quad T_{9+13}\left(\sum_{7 \leq i \leq 13} \alpha_i\right) = - \sum_{7 \leq i \leq 13} \alpha_i.$$

By using this we see that the automorphism  $T_{9+13}^t$  of  $\text{Hom}_{alg, gp}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R}$  is given by

$$T_{9+13}^t(\alpha_3^*) = \alpha_3^*, \quad T_{9+13}^t(\alpha_6^*) = \alpha_6^*, \quad T_{9+13}^t(\alpha_9^*) = 2\alpha_6^* - \alpha_9^*, \quad T_{9+13}^t(\alpha_{13}^*) = 2\alpha_6^* - 2\alpha_9^* + \alpha_{13}^*.$$

We introduce here a new basis  $\alpha_3^*$ ,  $\alpha_6^*$ ,  $\alpha_9^*$  and  $\alpha_9^* - \alpha_{13}^*$  on  $\text{Hom}_{alg, gp}(L, \mathbf{C}^*) \otimes_{\mathbf{Z}} \mathbf{R}$ . With respect to this basis, we have the following matrix representation.

$$T_3^t = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad T_6^t = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad T_{9+13}^t = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$


and


$$T_{13}^t = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}.$$

One can easily read off that

$$\langle T_3^t, T_6^t, T_{9+13}^t \rangle = W(B_3), \quad \langle T_{13}^t \rangle = W(B_1), \quad \langle T_3^t, T_6^t, T_{9+13}^t, T_{13}^t \rangle = W(B_3) \times W(B_1)$$

Let us compute how many  $\mathbf{Q}$ -factorial terminalizations  $X$  has.  $\mathcal{S}(L)$  contains 4 conjugacy classes. Two of them are those classes which respectively contain  $Q$  and  $Q'$ . Other two classes are given by the marked Dynkin diagrams (cf. Remark 11):





Therefore

$$\#\mathcal{S}(L) = 4 \cdot |W'| = 4 \cdot 96 = 384$$

Note that  $X_{[3^3, 2^2, 1]} \rightarrow \bar{O}_{[3^3, 2^2, 1]}$  is a double covering. Recall that there is an exact sequence

$$1 \rightarrow W_X \rightarrow W' \xrightarrow{\bar{\rho}_X} \text{Aut}(X/\bar{O})/\text{Aut}(X_{[3^3, 2^2, 1]}/\bar{O}_{[3^3, 2^2, 1]})(= \mathbf{Z}/2\mathbf{Z})$$

Let  $\omega_3$  (resp.  $\omega_6, \omega_{9+13}, \omega_{13}$ ) be an element of  $W'$  determined by the twist  $T_3^t$  (resp.  $T_6^t, T_{9+13}^t, T_{13}^t$ ). Then

$$\bar{\rho}_X(\omega_3) = \bar{\rho}_X(\omega_6) = \bar{\rho}_X(\omega_{9+13}) = 0, \quad \bar{\rho}_X(\omega_{13}) = \bar{1}$$

This means that

$$W_X = W(B_3).$$

As a consequence, we have

$$\#\{\mathbf{Q}\text{-factorial terminalizations of } X\} = \#\mathcal{S}(L)/|W_X| = 8.$$

□

## References

- [B-K] Brylinski, R., Kostant, B.: The variety of all invariant symplectic structures on a homogeneous space and normalizers of isotropy subgroups, in Symplectic geometry and Mathematical Physics (edited by P. Donato, C. Duval, J. Elhada, G.M. Tuynman), Progress in Math. **99**, (1991) Birkhauser, 80-113
- [BPW] Braden, T., Proudfoot, N., Webster, B.: Quantizations of conical symplectic resolutions I: local and global structures, Astérisque **384** (2016) 1-73
- [C-M] Collingwood, D. , McGovern, W. : Nilpotent orbits in semi-simple Lie algebras, van Nostrand Reinhold, Math. Series, 1993
- [Ho] Howlett, R.: Normalizers of parabolic subgroups of reflection groups, J. London Math. Soc. **21** (1980) 62-80
- [Lo] Losev, I.: Deformations of symplectic singularities and Orbit method for semisimple Lie algebras, arXiv: 1605.00592, Selecta Math. **28** (2022) online available
- [LMM] Losev, I., Mason-Brown, L., Matvieievskyi, D.: Unipotent ideals and Harish-Chandra bimodules, arXiv: 2108.03453
- [Ma] Matvieievskyi, D.: On the affinization of a nilpotent orbit cover, arXiv : 2003.09356

- [Na 1] Namikawa, Y.: Birational geometry and deformations of nilpotent orbits, *Duke Math. J.* **143** (2008) 375- 405
- [Na 2] Namikawa, Y.: Poisson deformations of affine symplectic varieties, II, *Kyoto J. Math.* **50**, No 4. (2010) 727-752
- [Na 3] Namikawa, Y.: Birational geometry for nilpotent orbits, in *Handbook of Moduli* (edited by G. Farkas and I. Morrison), Vol. III, *Advanced lectures in Mathematics* **26**, International Press, (2013) 1-38
- [Na 4] Namikawa, Y.: Poisson deformations and birational geometry, *J. Math. Sci. Univ. Tokyo*, **22** (2015) 339-359
- [Na 5] Namikawa, Y.: Birational geometry of symplectic resolutions of nilpotent orbits, *Advanced Studies in Pure Mathematics* **45**, (2006), in *Moduli Spaces and Arithmetic Geometry* (Kyoto, 2004), pp. 75-116
- [Na 6] Namikawa, Y.: Poisson deformations of affine symplectic varieties, *Duke Math. J.* **156** (2011) 51-85
- [Part 1] Namikawa, Y.: Birational geometry for the covering of a nilpotent orbit closure, *arXiv*: 1907.07812
- [S-S] Springer, T. A.; Steinberg, R.: Conjugacy classes. 1970 Seminar on Algebraic Groups and Related Finite Groups (The Institute for Advanced Study, Princeton, N.J., 1968/69) pp. 167–266 *Lecture Notes in Mathematics*, Vol. 131 Springer, Berlin

Research Institute for Mathematical Science, Kyoto University, Japan  
namikawa@kurims.kyoto-u.ac.jp