Enhanced Edelstein effect and interdimensional effects in an electron gas with Rashba spin-orbit coupling interface

A. C. Zulkoskey,^{*} R. Dick,[†] and K. Tanaka[‡]

and Centre for Quantum Topology and Its Applications (quanTA),

University of Saskatchewan, 116 Science Place, Saskatoon, Saskatchewan, Canada S7N 5E2

(Dated: January 6, 2022)

We examine the bound-state and free-state contributions to the density of states in a threedimensional electron gas with a two-dimensional interface with Rashba spin-orbit coupling. Confinement of electrons to the interface is achieved through the inclusion of an attractive potential in the interface. Motivation for our research comes from interest in heterostructure materials that exhibit the Edelstein and inverse Edelstein effects on surfaces or interfaces due to large Rashba spinorbit coupling. By modifying the Hamiltonian of a three-dimensional free electron gas to include an interface with Rashba spin-orbit coupling and an attractive potential, we are able to calculate the bound-state and free-state wavefunctions and corresponding density of states analytically. We find that one of the spin-split energy bands in the interface has an upper bound, resulting in an enhancement of the Edelstein and inverse Edelstein effect.

I. INTRODUCTION

The propagation properties of particles (or quasiparticles) affected by the presence of a surface or an interface in a three-dimensional material can be described using low-dimensional quantum mechanics. Analytic models can be constructed to include extra substructure terms, which affect propagation properties of electrons through a change in the effective $mass^1$ or confinement in the form of a quantum well². In both cases the Hamiltonian is constructed as a linear superposition of a free three-dimensional electron gas and a low-dimensional substructure contribution describing the effects of a surface or an interface. The density of states inside the lowdimensional structure that allows for calculation of, e.g., the number of charge carriers and thermal conductivity can be obtained analytically for these types of Hamiltonians and is thus a powerful tool for studying the interdimensional properties of electrons in a material with substructure.

The system of a two-dimensional quantum well immersed in a three-dimensional bulk is described by the Hamiltonian, 2

$$H = \frac{\mathbf{p}^2}{2m} - \frac{\hbar^2 \kappa}{m} \delta(z - z_0), \qquad (1)$$

for a particle of mass m. The quantum well exhibits confining properties through the binding energy, $B = \hbar^2 \kappa^2 / 2m$, with an inverse penetration depth κ . The corresponding density of states per volume at the location of the quantum-well structure ($z = z_0$) is given as a function of energy E by

$$\varrho(E, z_0) = \kappa \varrho_{d=2} (E + (\hbar^2 \kappa^2 / 2m)) + \varrho_{d=3}(E) \\ \times \left[1 - \frac{\hbar \kappa}{\sqrt{2mE}} \arctan\left(\frac{\sqrt{2mE}}{\hbar \kappa}\right) \right], \quad (2)$$

where

$$\varrho_d(E) = 2\Theta(E) \sqrt{\frac{m}{2\pi}}^d \frac{\sqrt{E}^{d-2}}{\Gamma(d/2)\hbar^d} \tag{3}$$

is the density of states for a free particle of mass m in d spatial dimensions and the particle is assumed to have spin 1/2 as an electron. Integrating the density of states $\varrho(E, z_0)$ over energy yields the relation between the Fermi energy and the particle density inside the quantum well at zero temperature:²

$$n(z_{0})\Big|_{-B < E_{F} < 0} = \frac{\kappa m}{\pi \hbar^{2}} \left(E_{F} + \frac{\hbar^{2} \kappa^{2}}{2m} \right) = \kappa n_{2}\Big|_{E_{2,F} = K_{2,F}},$$

$$n(z_{0})\Big|_{E_{F} > 0} = \frac{\kappa}{2\pi^{2}\hbar^{2}}$$

$$\times \left[\hbar \kappa \sqrt{2mE_{F}} - (\hbar^{2}\kappa^{2} + 2mE_{F}) \arctan\left(\frac{\sqrt{2mE_{F}}}{\hbar \kappa}\right) \right]$$

$$+ \frac{\kappa m}{\pi \hbar^{2}} \left(E_{F} + \frac{\hbar^{2}\kappa^{2}}{2m} \right) + \frac{1}{3\pi^{2}} \left(\frac{\sqrt{2mE_{F}}}{\hbar}\right)^{3}, \qquad (4)$$

where

$$n_d = \frac{2}{\hbar^d \Gamma((d+2)/2)} \sqrt{\frac{mE_F}{2\pi}}^d \tag{5}$$

is the density of particles in d dimensions, and $K_{2,F} = E_F + \hbar^2 \kappa^2 / 2m$ is the kinetic energy inside the quantum well. The analytic results for the density of states and the particle density inside the quantum well smoothly transitions from two-dimensional to three-dimensional behaviour as the inverse penetration depth κ approaches zero. Both results demonstrate that bound states exist for $E \geq -B$, and that particles confined to the quantum well contribute a two-dimensional density term, made dimensionally correct through the factor κ , reflecting the three-dimensional nature of the system.

Department of Physics and Engineering Physics,

Rashba spin-orbit coupling (RSOC) arises as a result of bulk inversion asymmetry (BIA), e.g., in the zinc blende structure³, as well as structure inversion asymmetry (SIA) in semiconductors⁴. The Rashba spin-orbit interaction, first analysed for a two-dimensional electron gas,⁴ stems from the nonrelativistic approximation of the Dirac equation⁵. The Hamiltonian and dispersion relation for a two-dimensional electron gas including RSOC is given by^{4,5}

$$H = \frac{\boldsymbol{p}_{\parallel}^{2}}{2m} + \alpha \left[\boldsymbol{\sigma} \times \boldsymbol{p}_{\parallel}/\hbar\right] \cdot \hat{z} ,$$

$$E_{\pm}(k_{\parallel}) = \frac{\hbar^{2}k_{\parallel}^{2}}{2m} \pm \alpha |\boldsymbol{k}_{\parallel}| , \qquad (6)$$

where *m* is the effective mass of an electron, $\alpha = e\hbar E_z(z)/4m^2c^2$ is the Rashba coefficient, $E_z(z)$ is an electric field in the direction \hat{z} perpendicular to the electron gas, $\boldsymbol{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ are the Pauli matrices, $\boldsymbol{k}_{\parallel} = (k_x, k_y)$ is the two-dimensional wave vector, and $k_{\parallel} = |\boldsymbol{k}_{\parallel}|$. RSOC causes momentum and spin to be "locked in" such that the E_+ and E_- branches in Eq. (6) have clockwise and counterclockwise winding of spin, respectively, as one goes around the Fermi surface. In Fig. 1, E_{\pm} are plotted as a function of k_x for $k_y = 0$, where spin points in either +y or -y direction. The density of states per unit area per spin,

$$\varrho(E) = \frac{1}{2\pi} \frac{k_{\parallel}(E)}{|dE/dk_{\parallel}|},\tag{7}$$

for $E = E_{\pm}$ using Eq. (6) is given by⁶

$$\varrho(E_{\pm}) = \frac{m}{2\pi\hbar^2} \left(1 \mp \frac{b}{\sqrt{b^2 + 2Em/\hbar^2}} \right), \quad E \ge 0, \\
\varrho(E_{-}) = \frac{mb}{\pi\hbar^2\sqrt{b^2 + 2mE/\hbar^2}}, \quad E < 0, \quad (8)$$

where $b = m\alpha/\hbar^2$.

Recent efforts have been put forth in the area of spintronics, which utilizes the spin degree of freedom for information storage and processing.⁷ In candidate materials for spintronics, strong RSOC induces novel properties on surfaces or interfaces⁸ such as the Edelstein effect⁹ or the inverse Edelstein effect, where conversion between charge and spin currents occurs.^{8,10} We study this kind of effects on the density of states for Hamiltonians which are a linear combination of three-dimensional kinetic terms and a two-dimensional RSOC term, in order to model systems with large RSOC on an interface or a surface. Materials which necessitate this description include topological insulators,¹¹ interfaces between metallic layers, e.g., $Bi/Ag^{12,13}$ or Cu/Bi^{14} interfaces, and conducting interfaces between LaAlO₃/SrTiO₃ insulating oxide layers.¹⁵ Heterostructures involving metaloxide interfaces¹⁶ as well as graphene,^{17,18} in which RSOC is enhanced by proximity to, e.g., transition metal dichalcogenides,^{8,10} also present systems in which RSOC




FIG. 1. (Colour online) The spin-split dispersion relation for a two-dimensional electron gas with RSOC for $k_y = 0$ in arbitrary units. The red (green) curve corresponds to E_+ (E_-) in Eq. (6) with a minimum energy of $E_{\min} = -m\alpha^2/2\hbar^2$. Up and down arrows correspond to +y and -y spin alignment for $k_y = 0$.

is prominent along an interface or a surface. In this paper we study the low-energy physics on a surface or an interface in such materials. The remaining sections are laid out as follows. In Sec. II we calculate the bound-state and free-state wavefunctions of our interdimensional model. In Sec. III the enhancement of the Edelstein effect for electrons bound to the interface is demonstrated. In Sec. IV we present the analytic results for the boundstate and free-state density of states and discuss the interdimensional behaviour. Our findings are summarised in Sec. V.

II. INTERDIMENSIONAL EFFECTS OF ELECTRONS WITH RSOC INTERFACE

Motivated by materials which exhibit novel features on interfaces or surfaces as a result of RSOC, as well as heterostructures where RSOC in the interface is enhanced by the neighbouring substrate^{8,10}, we construct a Hamiltonian as a superposition of a three-dimensional free electron gas and a two-dimensional interface with RSOC and an attractive potential at $z = z_0$. We extend the work of Ref. 1 to include a RSOC term⁴ in the interface,

$$H_{SO} = \alpha \left[\boldsymbol{\sigma} \times \boldsymbol{k}_{\parallel} \right] \cdot \hat{z} \,. \tag{9}$$

The construction of the Hamiltonian for electrons subject to spin-orbit coupling in the interface is as follows: We assume that the wavenumber component orthogonal to the interface is small compared to the inverse thickness of the interface L_{\perp}^{-1} , i.e., $|k_{\perp}L_{\perp}| \ll 1$. This implies that




FIG. 2. (Colour online) The shifted Fermi surfaces of a twodimensional electron gas with RSOC due to an applied electric field in the x direction. The red (green) circle corresponds to the E_+ (E_-) branch for the applied electric field $\mathbf{E} = 0$ and the maroon (olive) circle corresponds to the E_+ (E_-) branch for $\mathbf{E} \neq 0$.

the wavefunction in the direction orthogonal to the interface can be approximated as constant. This formulation yields a second-quantized Hamiltonian,

$$H = \int d^{3}\boldsymbol{x} \, \frac{\hbar^{2}}{2m} \boldsymbol{\nabla} \psi^{\dagger}(\boldsymbol{x}) \cdot \boldsymbol{\nabla} \psi(\boldsymbol{x}) - \int d^{2}\boldsymbol{x}_{\parallel} \left(i\alpha L_{\perp} \psi^{\dagger}(\boldsymbol{x}_{\parallel}, z_{0}) \left(\boldsymbol{\sigma}_{\parallel} \times \boldsymbol{\nabla}_{\parallel}\right) \cdot \hat{z} \psi(\boldsymbol{x}_{\parallel}, z_{0}) + V_{0} \psi^{\dagger}(\boldsymbol{x}_{\parallel}, z_{0}) \cdot \psi(\boldsymbol{x}_{\parallel}, z_{0}) \right),$$
(10)

where $\boldsymbol{x} = (\boldsymbol{x}_{\parallel}, z)$ and $\boldsymbol{\sigma}_{\parallel} = (\sigma_x, \sigma_y)$. The eigenvalues and eigenfunctions of Eq. (10) are separated into states which are bound to the interface $(E < \hbar^2 k_{\parallel}^2/2m, \kappa > 0)$,

$$\psi_{\boldsymbol{k}_{\parallel},\kappa_{\pm},+}(\boldsymbol{x}_{\parallel},z)_{\pm} = \langle \boldsymbol{x}_{\parallel}, z | \boldsymbol{k}_{\parallel},\kappa,+\rangle_{\pm} \\ = \frac{\exp(i\boldsymbol{k}_{\parallel}\cdot\boldsymbol{x}_{\parallel})}{2\pi}\sqrt{\kappa_{\pm}}\exp(-\kappa_{\pm}|z-z_{0}|)\varphi(\boldsymbol{k}_{\parallel})_{\pm}, \quad (11)$$

$$E_{\pm} = \frac{\hbar^2}{2m} (\mathbf{k}_{\parallel}^2 - \kappa_{\pm}^2) = \frac{\hbar^2 k_{\parallel}^2}{2m} - \frac{m L_{\perp}^2}{2\hbar^2} (V_0 \pm \alpha k_{\parallel})^2, \quad (12)$$

where $\varphi(\mathbf{k}_{\parallel})_{\pm} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ \pm ik_{+}/k_{\parallel} \end{pmatrix}$, $k_{\pm} = k_{x} \pm ik_{y}$, and $\kappa_{\pm} = (mL_{\perp}/\hbar^{2})(V_{0} \pm \alpha k_{\parallel})$, and two sets of orthogonal, transversely free states $(k_{\perp} \geq 0, E = \hbar^{2}(\mathbf{k}_{\parallel}^{2} + k_{\perp}^{2}))$

written as even (+) and odd (-) parity eigenstates,

$$\psi_{\boldsymbol{k}_{\parallel},\boldsymbol{k}_{\perp},+}(\boldsymbol{x}_{\parallel},z)_{\pm} = \langle \boldsymbol{x}_{\parallel}, z | \boldsymbol{k}_{\parallel}, \boldsymbol{k}_{\perp},+\rangle_{\pm}$$

$$= \frac{\exp(i\boldsymbol{k}_{\parallel} \cdot \boldsymbol{x}_{\parallel})}{2\sqrt{\pi^{3}}\sqrt{1 + (mL_{\perp}/\hbar^{2}k_{\perp})^{2}(V_{0} \pm \alpha \boldsymbol{k}_{\parallel})^{2}}} \left(\cos(k_{\perp}(z-z_{0})) - \frac{mL_{\perp}}{\hbar^{2}k_{\perp}}(V_{0} \pm \alpha \boldsymbol{k}_{\parallel})\sin(k_{\perp}|z-z_{0}|)\right)\varphi(\boldsymbol{k}_{\parallel})_{\pm}, \quad (13)$$

$$\psi_{\boldsymbol{k}_{\parallel},\boldsymbol{k}_{\perp},-}(\boldsymbol{x}_{\parallel},z) = \langle \boldsymbol{x}_{\parallel}, z | \boldsymbol{k}_{\parallel}, \boldsymbol{k}_{\perp}, -\rangle$$
$$= \frac{\exp(i\boldsymbol{k}_{\parallel} \cdot \boldsymbol{x}_{\parallel})}{2\pi} \frac{1}{\sqrt{\pi}} \sin(\boldsymbol{k}_{\perp}(z-z_{0})) \begin{pmatrix} \cos\xi\\ e^{i\chi}\sin\xi \end{pmatrix}, \quad (14)$$

where χ and ξ are arbitrary real numbers. Note that the interface bound states (11) have even (+) parity. In order for the ground-state energy to exist as a lower bound in Eq. (12) we require $\eta^2 \leq 1$ where $\eta = m\alpha L_{\perp}/\hbar^2$. Without this restriction $E \to -\infty$ as $k_{\parallel} \to \infty$. Thus, the Hamiltonian in Eq. (10) yields the dispersion relation,

$$E = E(|\boldsymbol{k}_{\parallel}|, \kappa_{\pm}) \equiv E_{\pm}(|\boldsymbol{k}_{\parallel}|)$$
(15)

for bound states $\{|\mathbf{k}_{\parallel}, \kappa, +\rangle_{\pm}\}$ with $E < \hbar^2 \mathbf{k}_{\parallel}^2/2m$, and

$$E = E(|\boldsymbol{k}_{\parallel}|, k_{\perp}) \tag{16}$$

for the unbound states $\{|\mathbf{k}_{\parallel}, k_{\perp}, \pm\rangle_{\pm}\}$ with $E \geq \hbar^{2} \mathbf{k}_{\parallel}^{2}/2m$. The corresponding density of states is given by¹

$$\varrho(E, \boldsymbol{x}) = \sum_{\pm} \int d^2 \boldsymbol{k}_{\parallel} \left(\delta(E - E_{\pm}(|\boldsymbol{k}_{\parallel}|)) |\langle \boldsymbol{x} | \boldsymbol{k}_{\parallel}, \kappa \rangle_{\pm} |^2 + \Theta(E - \hbar^2 \boldsymbol{k}_{\parallel}^2 / 2m) \left| \frac{\partial k_{\perp}(E, \boldsymbol{k}_{\parallel})}{\partial E} \right| |\langle \boldsymbol{x} | \boldsymbol{k}_{\parallel}, k_{\perp} \rangle_{\pm} |^2 \right),$$
(17)

where the sum over even- and odd-parity unbound states is implicitly assumed, and as such + and - for parity have been removed from the eigenvectors. Hence,

$$\varrho(E, \boldsymbol{x}) = \sum_{\pm} \left(\int_{0}^{2\pi} d\theta \, k_{\parallel} \left| \frac{\partial k_{\parallel}(E, \kappa_{\pm})}{\partial E} \right| |\langle \boldsymbol{x} | \boldsymbol{k}_{\parallel}, \kappa \rangle_{\pm} |^{2} \right. \\ \left. + \int d^{2} \boldsymbol{k}_{\parallel} \, \Theta(E - \hbar^{2} \boldsymbol{k}_{\parallel}^{2} / 2m) \left| \frac{\partial k_{\perp}(E, \boldsymbol{k}_{\parallel})}{\partial E} \right| |\langle \boldsymbol{x} | \boldsymbol{k}_{\parallel}, k_{\perp} \rangle_{\pm} |^{2} \right).$$

$$(18)$$

III. BOUND-STATE DISPERSION RELATION AND ENHANCED EDELSTEIN EFFECT

In a purely two-dimensional system with RSOC, an applied electric field along +x causes electrons to move in the -x direction and populate states with $k_x < 0$ at the expense of states with $k_x > 0$. Figure 2 illustrates




FIG. 3. (Colour online) The spin-split bound-state energy $(2mL_{\perp}^2/\hbar^2)E_{\pm}$ as a function of k_xL_{\perp} for $k_y = 0$. The red (green) curve corresponds to E_- (E_+) in Eq. (12) with a minimum dimensionless energy of $2mE_{\min}L_{\perp}^2/\hbar^2 = -\beta^2/(1-\eta^2)$, where $\beta = mV_0L_{\perp}^2/\hbar^2$ and $\eta = m\alpha L_{\perp}/\hbar^2$.

the shift of the inner and outer Fermi circles due to an applied electric field in the x direction. An increase in -y and +y spin polarization states for $k_y = 0$ creates a net -y spin polarization, as the outer Fermi circle dominates over the inner one with a larger number of states. This is the well-known Edelstein effect,⁸⁻¹⁰ where a charge current is converted to an accumulation of spin in the transverse direction. Likewise, the inverse Edelstein effect is the conversion of a spin current to a transverse charge current.^{8,10,12}

In our three-dimensional model we find that the Edelstein effect⁹ (and inverse Edelstein effect) is affected through the restriction $\kappa_{-} > 0$. The spin-split boundstate bands in Eq. (12), made dimensionless with a factor $2mL_{\perp}^2/\hbar^2$, are plotted as a function of $k_x L_{\perp}$ for $k_y = 0$ in Fig. 3. Note that $E_{-} \geq E_{+}$ for $k_{\parallel} \geq 0$. The requirement $\kappa_{-} > 0$ implies $k_{\parallel} < V_0/\alpha$ in order for a bound-state solution to exist. This restricts the energy E_{-} in Eq. (12) to a maximum value of $E_{-} = \hbar^2 V_0^2 / 2m\alpha^2$. Consider the situation of $E_F \lesssim \hbar^2 V_0^2 / 2m\alpha^2$. Åpplying an electric field along the interface results in a net spin polarization in the -y direction due to the Edelstein effect. However, the states belonging to the E_{-} branch can only be populated up to the maximum dimensionless energy of $V_0^2 L_\perp^2 / \alpha^2$. The contribution to the net spin polarization from the increase in +y spin alignment in the $k_x < 0$ region of the E_{-} branch therefore has an upper bound. The E_+ branch has no restriction owing to $\kappa_+ > 0$; $\forall k_{\parallel}$, which leads to a larger net spin polarization in the -ydirection compared to the purely two-dimensional model and hence an enhanced Edelstein effect. The dimensionless energy bands $(2mL_{\perp}^2/\hbar^2)E_{\pm}$ are plotted as a function of $k_x L_{\perp}$ for $k_y = 0$ in Fig. 4 along with the maximum allowed E_{-} for the bound states. The energy cap on E_{-}




FIG. 4. (Colour online) The spin-split bound-state energy $(2mL_{\perp}^2/\hbar^2)E_{\pm}$ as a function of k_xL_{\perp} for $k_y = 0$. The red (green) curve corresponds to E_- (E_+) in Eq. (12). Due to the requirement $\kappa_- > 0$, E_- is restricted to a maximum dimensionless energy of $(V_0L_{\perp}/\alpha)^2 = \beta^2/\eta^2$, where $\beta = mV_0L_{\perp}^2/\hbar^2$ and $\eta = m\alpha L_{\perp}/\hbar^2$, indicated by the blue line.

results in shifting of the Fermi circles to elliptical shape, as illustrated in Fig. 5.

IV. BOUND-STATE AND FREE-STATE DENSITY OF STATES AT $z = z_0$

The restriction on the energy range for the E_{\perp} branch also impacts the bound-state density of states at $z = z_0$. Substituting Eq. (11) in Eq. (18) yields the density of states of the spin-split energy bands in three energy ranges. For $E_{\min} \leq E < -mV_0^2 L_{\perp}^2/2\hbar^2$ only the E_+ branch contributes to the density of states and is given by

$$\varrho(E_+, z_0) = \frac{m^2 \eta \left(V_0^2 L_\perp^2 + f(E, \eta, V_0) \right)}{\pi \hbar^4 \sqrt{f(E, \eta, V_0)} (1 - \eta^2)^2}, \qquad (19)$$

where $E_{\rm min} = -mV_0^2 L_{\perp}^2/2\hbar^2(1-\eta^2)$, $f(E,\eta,V_0) = V_0^2 L_{\perp}^2 + (2E\hbar^2/m)(1-\eta^2)$ and $\eta = m\alpha L_{\perp}/\hbar^2$. Equation (19) shows that the density of states for $E_{\rm min} \leq E \leq -mV_0^2 L_{\perp}^2/2\hbar^2$ contains a van Hove singularity at $E = E_{\rm min}$. This singularity occurs at the bottom of the E_+ branch, which is analogous to the van Hove singularity at $E = -m\alpha^2/2\hbar^2$ in the purely two-dimensional model.⁴ Shown in Fig. 6 is the density of states in the energy range $E_{\rm min} \leq E \leq -mV_0^2 L_{\perp}^2/2\hbar^2$. The density of states of the spin-split bound states in the energy range $-mV_0^2 L_{\perp}^2/2\hbar^2 \leq E \leq \hbar^2 V_0^2/2m\alpha^2$ is displayed in Fig. 7

-70

-60

-50 -40 = -30

-20 --10




FIG. 5. (Colour online) The shifted Fermi surfaces due to the enhanced Edelstein effect. The red and green circles correspond to E_{-} and E_{+} , respectively, in Eq. (12), while the maroon and olive circles correspond to the situation shown in Fig. 4.

FIG. 6. (Colour online) The spin-split and total bound-state density of states at $z = z_0$ for $E_{\min} \leq E \leq -mV_0^2 L_\perp^2/2\hbar^2$ in units of $m/(2\pi\hbar^2 L_\perp)$. The green curve is the total bound-state density of states, while the blue (red) curve is the contribution from the + (-) sign choice in the solutions for k_{\parallel} in E_+ .

 $\frac{2 mEL^2}{\hbar^2}$

-26.8 -26.6

-27.6 -27.4 -27.2 -27

and is given by

$$\varrho(E_{\pm}, z_0) = \frac{mL_{\pm}}{2\pi\hbar^2} \left[\left(\frac{\pm mV_0 \alpha L_{\pm}^2 / \hbar^2 + \sqrt{f(E, \eta, V_0)}}{(\sqrt{f(E, \eta, V_0)})(\hbar^2 / m)(1 - \eta^2)} \right) \times \left(V_0 \pm \alpha \frac{\pm mV_0 \alpha L_{\pm}^2 / \hbar^2 + \sqrt{f(E, \eta, V_0)}}{(\hbar^2 / m)(1 - \eta^2)} \right) \right] \\
\times \Theta(E + mV_0^2 L_{\pm}^2 / 2\hbar^2).$$
(20)

The total density of states corresponding to Eq. (20) is

$$\varrho(E, z_0) = \frac{\varrho_{d=2}(K_2)\beta}{L_{\perp}(1 - \eta^2)^2}, \qquad (21)$$

where $K_2 = E + mV_0^2 L_\perp^2/2\hbar^2$ is the kinetic energy of electrons whose wavefunctions are exponentially suppressed perpendicular to the interface, and $\beta = mV_0L_\perp^2/\hbar^2$. Equation (21) demonstrates that the total density of states in the energy range where both spin-split bands contribute is proportional to the free two-dimensional density of states scaled by the interface thickness L_\perp , to reflect the three-dimensional nature of the system. For $E > \hbar^2 V_0^2/2m\alpha^2$ the E_- branch no longer contributes and the bound-state density of states is given by $\varrho(E_+, z_0)$ in Eq. (20). This is shown in Fig. 7. The freestate density of states is calculated by inserting Eq. (13) into Eq. (18),




FIG. 7. (Colour online) The spin-split and total boundstate density of states at $z = z_0$ for $-mV_0^2 L_\perp^2/2\hbar^2 \leq E \leq \hbar^2 V_0^2/2m\alpha^2$ in units of $m/(2\pi\hbar^2 L_\perp)$. The green curve is the total bound-state density of states, while the blue and red curves correspond to, respectively, $\varrho(E_+, z_0)$ and $\varrho(E_-, z_0)$ in Eq. (20).






FIG. 8. (Colour online) The spin-split free-state density of states at $z = z_0$ in units of $m/(2\pi^2\hbar^2L_{\perp})$. The green and red curves correspond to, respectively, $\varrho(E, z_0)_{-\alpha x}$ and $\varrho(E, z_0)_{+\alpha x}$ in Eq. (22). At $2mEL_{\perp}^2/\hbar^2 = \beta^2/\eta^2$, $\varrho(E, z_0)_{-\alpha x}$ exhibits a van Hove type singularity.

$$\varrho(E, z_0) = \frac{m\Theta(E)}{2\pi^2 \hbar^2 L_{\perp}} \int_0^{\sqrt{2mEL_{\perp}^2/\hbar}} dx \\
\left(\frac{x\sqrt{2mEL_{\perp}^2/\hbar^2 - x^2}}{2mEL_{\perp}^2/\hbar^2 - x^2 + (mL_{\perp}/\hbar^2)^2(-V_0 + \alpha x)^2} + \frac{x\sqrt{2mEL_{\perp}^2/\hbar^2 - x^2}}{2mEL_{\perp}^2/\hbar^2 - x^2 + (mL_{\perp}/\hbar^2)^2(-V_0 - \alpha x)^2}\right) \\
\equiv \varrho(E, z_0)_{+\alpha x} + \varrho(E, z_0)_{-\alpha x} .$$
(22)

The contributions to the density of states from the two bands are signified by the sign of the αx terms in Eq. (22), and correspond to the $b_{\pm}(E,\beta,\eta)$ terms in the total density of states,

$$\varrho(E, z_0) = \frac{m\Theta(E)}{2\pi^2 \hbar^2 L_{\perp}(1 - \eta^2)\sqrt{d(E, \beta, \eta)}} \times \left[b_+(E, \beta, \eta) \left(\sqrt{\frac{2mEL_{\perp}^2}{\hbar^2}} - c_+(E, \beta, \eta) \arctan\left(\frac{\sqrt{2mEL_{\perp}^2/\hbar^2}}{c_+(E, \beta, \eta)}\right) \right) + b_-(E, \beta, \eta) \left(\sqrt{\frac{2mEL_{\perp}^2}{\hbar^2}} - c_-(E, \beta, \eta) \arctan\left(\frac{\sqrt{2mEL_{\perp}^2/\hbar^2}}{c_-(E, \beta, \eta)}\right) \right) \right],$$
(23)

where

$$b_{\pm}(E,\beta,\eta) = \pm \eta\beta + \sqrt{d(E,\beta,\eta)},$$

$$c_{\pm}(E,\beta,\eta) = \sqrt{h_{\pm}(E,\beta,\eta) - 2mEL_{\perp}^{2}/\hbar^{2}},$$

$$d(E,\beta,\eta) = \beta^{2} + \frac{2mEL_{\perp}^{2}}{\hbar^{2}}(1-\eta^{2}),$$

$$h_{\pm}(E,\beta,\eta) = \left(\frac{\sqrt{d(E,\beta,\eta)} \pm \eta\beta}{1-\eta^{2}}\right)^{2}.$$
(24)

The spin-split density of states, $\rho(E, z_0)_{+\alpha x}$ and $\rho(E, z_0)_{-\alpha x}$ in Eq. (22), is displayed in Fig. 8. Lowenergy contributions to the free-state density of states are dominated by the arctan terms in Eq. (23) with the threedimensional \sqrt{E} behaviour appearing for $E \gg E_{\min}$. The $\rho(E, z_0)_{-\alpha x}$ term in Eq. (22) contains a van Hove singularity at $2mEL_{\perp}^2/\hbar^2 = \beta^2/\eta^2$, which corresponds to $k_{\parallel} = V_0/\alpha, k_{\perp} = 0$ and the energy cap on the E_{\perp} branch. This singularity originates in the $\partial k_{\perp}/\partial E \sim 1/k_{\perp}$ term in Eq. (18). $k_{\perp} = 0$ does not necessarily generate a singularity in the density of states due to $|\langle \boldsymbol{x} | \boldsymbol{k}_{\parallel}, k_{\perp} \rangle_{\pm}|^2$ multiplied to $|\partial k_{\perp}/\partial E|$ in Eq. (18), with the wavefunction given in Eq. (13) vanishing sufficiently quickly. However, when the bound states cease to exist in the E_{\perp} branch, the $1/k_{\perp}$ divergence is no longer compensated, resulting in the van Hove singularity.

V. CONCLUSIONS

We have constructed a model Hamiltonian for electrons in a three-dimensional system with an interface, in which the electrons can be trapped by an attractive potential and experience RSOC. We have analytically obtained the bound-state and free-state wavefunctions, and the density of states at the location of the interface. We find that the density of states in the interface exhibits twodimensional behaviour due to the bound states as well as three-dimensional behaviour in the high-energy limit, $E \gg E_{\min} = -mV_0^2 L_{\perp}^2/2\hbar^2(1-\eta^2)$. The requirement of $\eta^2 = (m\alpha L_{\perp}/\hbar^2)^2 \leq 1$ for a bound state to exist results in a maximum allowed energy of $\hbar^2 V_0^2/2m\alpha^2$ for one of the spin-split bands. This restriction impacts the allowed populations of each energy band and enhances the Edelstein and inverse Edelstein effects. Furthermore, the existence of the upper bound in one of the energy bands in the interface leads to a van Hove type singularity in the free-state density of states. Our results for the density of states are analytical and readily applicable to interfaces and surfaces with RSOC.

VI. ACKNOWLEDGEMENTS

The research was supported by the Natural Sciences and Engineering Research Council of Canada.

- * Corresponding author, acz621@mail.usask.ca
- [†] rainer.dick@usask.ca
- ‡ kat
221@campus.usask.ca
- ¹ R. Dick, Physica E **40**, 2973 (2008); Nano. Res. Lett. **5**, 1546 (2010).
- ² R. Dick, Nanoscale Res. Lett. **7**, 581 (2012).
- ³ G. Dresselhaus, Phys. Rev. **100**, 580-586 (1955).
- ⁴ Yu. A. Bychkov, E. I. Rashba, JETP Lett. **39**, 78 (1984).
- ⁵ R. Dick, Advanced Quantum Mechanics: Materials and Photons, Second Edition, Springer, Switzerland, (2016).
- ⁶ R. Winkler, Spin-Orbit Coupling Effects in Two-Dimensional Electron and Hole Systems, Springer, Berlin (2003).
- ⁷ S. A. Wolf et al., Science 294, 1488-1495 (2001); I. Zutic, J. Fabian and S. Das Sarma, Rev. Mod. Phys. 76, 323-410 (2004).
- ⁸ A. Soumyanarayanan, N. Reyren, A. Fert and C. Panagopoulos, Nature **539**, 509 (2016).
- ⁹ V.M. Edelstein, Solid State Commun. **73**, 233-235 (1990).
- ¹⁰ W. Han, Y.-C. Otani and S. Maekawa, npj Quantum Mat.
 3, 27 (2018).
- ¹¹ M. Z. Hasan, C. L. Kane, Rev. Mod. Phys. 82, 3045 (2010); X.-L. Qi, S.-C. Zhang, Rev. Mod. Phys. 83, 1057 (2011).
- ¹² J. C. R. Sánchez et al., Nat. Comm. 4, 2944 (2013); W. Zhang, M. B. Jungfleisch, W. Jiang, J. E. Pearson and A. Hoffmann, Appl. Phys. **117**, 17C727 (2015).
- ¹³ H. J. Zhang, S. Yamamoto, B. Gu, H. Li, M. Maekawa, Y. Fukaya, and A. Kawasuso, Phys. Rev. Lett. **114**, 166602

(2015).

- ¹⁴ M. Isasa et al., Phys. Rev. B **93**, 014420 (2016).
- ¹⁵ A. Ohtomo and H.Y. Hwang, Nature **427**, 423-426 (2004);
 H.Y. Hwang et al., Nat. Mater. **11**, 103-113 (2012);
 J. Mannhart and D.G. Schlom, Science **327**, 1607-1611 (2010);
 S. Okamoto and A.J. Millis, Nature **428**, 630-633 (2004);
 S.A. Chambers, Surf. Sci. **605**, 1133-1140 (2011);
 G. Herranz et al., Phys. Rev. Lett. **98**, 216803 (2007);
 A.D. Caviglia, M. Gabay, S. Gariglio, N. Reyren, C. Cancellieri and J.-M. Triscone, Phys. Rev. Lett. **104**, 126803 (2010).
- ¹⁶ S. Karube, K. Kondou and Y.-C. Otani, Appl. Phys. Expr. 9, 033001 (2016); J. Kim, Y.-T. Chen, S. Karube, S. Takahashi, K. Kondou, G. Tatara and Y.-C. Otani, Phys. Rev. B 96, 140409(R) (2017).
- ¹⁷ K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, A. A. Firsov, Nature **438**, 197 (2005); A. H. Castro Neto, F. Guinea, N. M. R. Peres, K.S. Novoselov, A. K. Geim, Rev. Mod. Phys. **81**, 109 (2009); D. S. L. Abergel, V. Apalkov, J. Berashevich, K. Ziegler, T. Chakraborty, Adv. Mat. **59**, 261 (2010).
- ¹⁸ G. W. Semenoff, Phys. Rev. Lett. **53**, 2449 (1984).
- ¹⁹ L. van Hove, Phys. Rev. **89**, 1189-1193 (1953).
- ²⁰ H. Zhang et al., Nat. Phys. 5, 438-442 (2009); P. D. C. King, R. C. Hatch, M. Bianchi, R. Ovsyannikov, C. Lupulescu, G. Landolt et al., Phys. Rev. Lett. 107, 096802 (2011).