

Extremal case of parabolic differential equations having discontinuous unbounded coefficients.

Existence of fundamental solution for an initial Cauchy problem. Parametrix method

M.R.Formica, E.Ostrovsky, L.Sirota

Università degli Studi di Napoli Parthenope, via Generale Parisi 13, Palazzo Pacanowsky, 80132, Napoli, Italy.

e-mail: mara.formica@uniparthenope.it

Israel, Bar-Ilan University, department of Mathematic and Statistics, 59200,

e-mail:eugostrovsky@list.ru

e-mail:sirota3@bezeqint.net

Abstract

We prove in this short report the existence of a fundamental solution (F.S.) for the Cauchy initial boundary problem on the whole space for the parabolic differential equation having at origin the point of non-integrable unbounded discontinuity for coefficient before a first order derivative.

We give also the non-asymptotic rapidly decreasing at infinity estimate for these function.

We extend the classical parametrix method offered by E.E.Levi.

Key words and phrases

Parabolic differential equation (PDE), parametrix method, fundamental solution (F.S.), degenerate diffusion, random variable (r.v.), random process (r.p.), Gamma, Beta-functions; Bessel's and variable Bessel's random process, modified Bessel's function, reflection at the origin, singular heat equation, subgaussian random variables and processes, Brownian motion, Markov random process, transfer density of probability for diffusion random processes, Mittag-Leffler function, non-asymptotic estimate, Chapman-Kolmogorov equation, Neuman series, Hölder's continuity, Volterra's integral equation, initial Cauchy problem.

1 Definitions. Notations. Statement of problem.

We consider in this report the Partial Differential Equation (PDE) of (uniform) parabolic type (singular heat equation)

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2} + \frac{1 + 2b(t, x)}{x} \frac{\partial u}{\partial x}, \quad (1)$$

$u = u(t, x)$, where $t \geq 0$, $x \in R_+$, and the coefficient $b = b(t, x)$ is continuous bounded function such that in general case $b(t, 0) \neq 0$, common with the condition of (homogeneous) reflection at the origin

$$\lim_{x \rightarrow 0^+} x^{1-2b(t, x)} u'_x(t, x) = 0 \quad (2)$$

and initial value problem (Cauchy statement)

$$\lim_{t \rightarrow s^+} u(t, x) = f(x), \quad u = u(t, x) = u[f](t, x), \quad s \geq 0, \quad (3)$$

where $f(\cdot)$ is certain measurable function satisfying some growth condition at $x \rightarrow \infty$; for definiteness one can suppose its continuity and boundedness: $\text{vraisup}_x |f(x)| < \infty$.

Briefly, $L_{t,x}u = L[b]_{t,x}u = 0$, where $L[b]_{t,x}$ is a linear parabolic partial differential operator of a parabolic type with singular not integrable coefficients of the form

$$L[b]_{t,x} = L_{t,x} \stackrel{\text{def}}{=} \frac{\partial}{\partial t} - \frac{1}{2} \frac{\partial^2}{\partial x^2} - \frac{1 + 2b(t, x)}{x} \frac{\partial}{\partial x} \quad (4)$$

in the domain $t, x > 0$.

Let us recall the classical definition.

Definition 1.1. The (measurable) function $p = p(t, x, s, y)$, $0 < s < t$; $(x, y) \in (R_+)^2$ as ordinary is said to be a fundamental solution (F.S.) for the equation (1) common with the initial condition (3) and the reflection condition (2), iff for all the fixed values (s, y) , $0 < s < t$, $x, y \in R_+$ it satisfies the equation (1) as well as the initial condition (3) and the reflection condition (2) and for all the bounded continuous function $f = f(x)$

$$\lim_{t \rightarrow s^+} \int_{R_+} p(t, x, s, y) f(y) dy = f(x), \quad x \in R_+, \quad (5)$$

if of course there exists.

In this case the F.S. $p(t, x, s, y)$ may be interpreted as the transfer density of probability for diffusion random non-homogeneous in the time as well as in the space Markov process, see e.g. [4], [11]. As a consequence:

$$p(t, x, s, y) \geq 0; \quad \int_0^\infty p(t, x, s, y) dy = 1$$

and it satisfies also the famous Chapman-Kolmogorov equation

$$p(t, x, s, y) = \int_0^\infty p(t, x, v, z) p(v, z, s, y) dz, \quad 0 < s < v < t. \quad (6)$$

On the other words, the (unique) solution $u = u(t, x)$ of the Cauchy problem (1), (3), (2) in the whole space R_+ may be written as follows

$$u(t, x) = \int_{R_+} p(t, x, s, y) f(y) dy. \quad (7)$$

For instance, the famous Gauss function

$$Z(t, x, s, y) = Z(t - s, x - y) := (2\pi(t - s))^{-1/2} \exp \left\{ -\frac{1}{2} \frac{(x - y)^2}{(t - s)} \right\} \quad (8)$$

is the F.S. for the ordinary heat equation only with initial condition in the whole space $x \in R$

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2}.$$

The case of the reflected Brownian motion will be considered further.

Another example, important for us. Consider the following infinitesimal operator for diffusion random Markov process $\xi = \xi(t)$

$$L_a = \frac{1}{2} \frac{\partial^2}{\partial x^2} + \frac{1 + 2a}{x} \frac{\partial}{\partial x}, \quad a = \text{const} \in (-1, 0), \quad x \in R_+,$$

with reflection at the origin

$$f \in D(L_a) \Rightarrow \left\{ f \in C^2(0, \infty), \lim_{x \rightarrow 0^+} x^{1-2a} f'(x) = 0 \right\}$$

and corresponding PDE

$$\frac{\partial u}{\partial t} = L_a u$$

with ordinary initial condition.

If $a \geq 0$ then the point $x = 0$ is unattainable; when $a \leq -1$ it is absorbing point, see [42]. *Therefore we consider in this report only the case $-1 < a < 0$ and correspondingly*

$$-1 < b(t, x) < 0.$$

The fundamental solution has a form $p(t, x, s, y) = p_a(t, x, s, y) \stackrel{\text{def}}{=}$

$$(t-s)^{-1} x^{-a} y^{a+1} \exp\left\{-\frac{x^2+y^2}{2(t-s)}\right\} I_a\left(\frac{xy}{t-s}\right), \quad (9)$$

$x, y > 0$, $t > s$, where $I_a(\cdot)$ denotes an usually *modified* Bessel's function of order a :

$$I_a(z) \stackrel{def}{=} \sum_{k=0}^{\infty} \frac{(z/2)^{a+2k}}{k! \Gamma(a+k+1)}, \quad (10)$$

see [42], [44].

The case $a = -1/2$ correspondent to the ordinary reflected at the origin Brownian motion:

$$p_{-1/2}(t, x, s, y) = (2\pi(t-s))^{-1/2} \left\{ \exp\left[\frac{(x+y)^2}{2(t-s)}\right] + \exp\left[\frac{(x-y)^2}{2(t-s)}\right] \right\},$$

$x, y > 0$, $0 < s < t$.

Note by the way as long as

$$\int_0^{\infty} p(t, x, s, y) dy = 1,$$

$$\int_0^{\infty} y^{a+1} e^{-y^2/2t} I_a(xy/t) dy = t x^a v e^{x^2/2t}, \quad x, y, t > 0, \quad a \in (-1, 0),$$

or equally for at the same values of parameters

$$\int_0^{\infty} z^{a+1} e^{-wz^2/2} I_a(z) dz = w^{-a-1} e^{1/2w}, \quad w > 0. \quad (11)$$

Further, it follows from the Chapman-Kolmogorov's equation (6) that

$$\begin{aligned} & \int_0^{\infty} (t-v)^{-1} x^{-a} z^{a+1} \exp\left\{-\frac{x^2+z^2}{2(t-v)}\right\} I_a\left(\frac{xz}{t-v}\right) \times \\ & (v-s)^{-1} z^{-a} y^{a+1} \exp\left\{-\frac{z^2+y^2}{2(v-s)}\right\} I_a\left(\frac{zy}{v-s}\right) dz = \\ & (t-s)^{-1} x^{-a} y^{a+1} \exp\left\{-\frac{x^2+y^2}{2(t-s)}\right\} I_a\left(\frac{xy}{t-s}\right), \quad 0 < s < v < t < \infty, \quad (12) \end{aligned}$$

or equally

$$\begin{aligned} & \int_0^{\infty} t^{-1} z \exp\left\{-\frac{x^2+z^2}{2t}\right\} I_a\left(\frac{xz}{t}\right) \times \\ & s^{-1} \exp\left\{-\frac{z^2+y^2}{2s}\right\} I_a\left(\frac{zy}{s}\right) dz = \end{aligned}$$

$$(t+s)^{-1} \exp \left\{ -\frac{x^2+y^2}{2(t+s)} \right\} I_a \left(\frac{xy}{t+s} \right), 0 < s, t < \infty. \quad (13)$$

We will emphasize that the condition $a \in (-1, 0)$ is very important for the last fourth relations.

Our claim in this preprint is to prove the existence of F.S. for our equation (1) with initial condition (3) and reflection condition (2) under suitable conditions.

We give also the exact bilateral estimates for this fundamental solution.

The existence of F.S. for *strictly* parabolic PDE with smooth coefficients, for instance, bounded and belonging to the Hölder space, is explained in many works, e.g. [4], [14], [12], [18], [19], [20], [25], [33], [34], [36], [37]. At the same problem for parabolic systems is considered in [17], [52]. The case of parabolic equations with discontinuous coefficients is investigated in [12], [24], [28], [31], [48], [53] etc.

The recent results about the parabolic equations with singular coefficients may be found in articles [8], [46], [54], [55] and so one.

We will essentially follow the interest article of Thomas Deck and Susanne Kruse [12], in which was considered the case of an equation

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2} + b_1(t, x) \frac{\partial u}{\partial x}, \quad (14)$$

in which the coefficient $b_1 = b_1(t, x)$ is Hölder continuous and may be unbounded at $x \rightarrow \pm\infty$ in the following sense

$$|b_1(t, x)| \leq C \left(1 + |x|^\beta \right), \quad C, \beta = \text{const} < \infty.$$

The uniqueness of the F.S. for parabolic PDE is proved, e.g. in [13], [14], [20], [28], [33], [48].

In the probability theory, more precisely, in the theory of diffusion random processes the F.S. $p(t, x, s, y)$ represent the density of transition of probability. The equations of the form (1) appears in particular in [42], [46]; see also [4], [11].

2 Main result. Existence of the F.S.

Theorem 2.1. Suppose in the equation (1) the coefficient $b = b(t, x)$ is Hölder continuous:

$$\exists \alpha = \text{const} \in (0, 1], \quad H = \text{const} \in (0, \infty) \Rightarrow$$

$$|b(t, x) - b(s, y)| \leq H[|t - s| + |x - y|]^\alpha. \quad (15)$$

Suppose also that this function is bilateral bounded as follows

$$\beta \stackrel{def}{=} \inf_{t,x} b(t, x) > -1, \quad (16)$$

$$\beta_+ \stackrel{def}{=} \sup_{t,x} b(t, x) < 0. \quad (17)$$

Then there exists a F.S. $p(t, x, s, y)$ for the equation (1), common with initial and reflection conditions, which satisfies in addition the following estimates (38), (41).

Proof.

1. We will use the classical parametrix method, see [36], [37], [14], [12], [19], [20], [25], [33], [34], etc., applying as the initial approximation the function $p_{b(s,y)}(t, x, s, y)$ instead the well - known approximation $Z(t, x, s, y)$.

Note that the initial approximation of the form $Z(t, x, s, y)$. was used also in the recent article [18] for the parabolic PDE having discontinuous but local integrable coefficient

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2} + \frac{1 + 2b(t, x)}{|x|^\gamma} \frac{\partial u}{\partial x}, \quad (18)$$

$\gamma = \text{const} \in (0, 1)$, $x \in \mathbb{R}$.

We will follow the article [25], down to notations.
Introduce for this purpose the following functions.

$$w = w(t, x, \tau, \xi) = w_{\tau, \xi}(t, x, \tau, \xi) := p_{b(\tau, \xi)}(t, x, \tau, \xi),$$

$$w^{(2)} = w^{(2)}(t, x, \tau, \xi) = w_{t, x}(t, x, \tau, \xi) := p_{b(t, x)}(t, x, \tau, \xi),$$

where (here and throughout whole report)

$$t, x, \tau, \xi > 0, \tau < t.$$

The F.S. $p(t, x, \tau, \xi)$ for source equation (1) may be found from the representation

$$p(t, x, \tau, \xi) := w_{\tau, \xi}(t, x, \tau, \xi) + \int_{\tau}^t d\theta \int_0^{\infty} w_{\theta, \zeta}(t, x, \theta, \zeta) \cdot \Phi(\theta, \zeta, \tau, \xi) d\zeta, \quad (19)$$

see [25], relation 4.18, where the function $\Phi = \Phi(\theta, \zeta, \tau, \xi)$ may be found from the following linear integral equation

$$\Phi(t, x, \tau, \xi) = L_{t, x} (w_{\tau, \xi}(t, x, \tau, \xi)) + \quad (20)$$

$$\int_{\tau}^t d\theta \int_0^{\infty} L_{t,x}(w_{\theta,\zeta}(t,x,\theta,\zeta)) \cdot \Phi(\theta,\zeta,\tau,\xi) d\zeta. \quad (21)$$

see [25], relation 4.54.

2. Denote for convenience

$$K = K(t,x,\tau,\xi) := L_{t,x}(w_{\tau,\xi}(t,x,\tau,\xi)), \quad t,x,\tau,\xi > 0;$$

and define for brevity following the authors of [12] the two-dimensional (generalized) convolution

$$f * g(t,x,s,y) \stackrel{def}{=} \int_s^t dr \int_0^{\infty} f(t,x,r,z)g(r,z,s,y) dz, \quad (22)$$

then the equation (21) takes the form $\Phi(t,x,s,y) = K(t,x,s,y) +$

$$\int_s^t dr \int_0^{\infty} dz K(t,x,r,z) \cdot \Phi(r,z,s,y), \quad 0 < s \leq t < \infty, \quad x,y > 0. \quad (23)$$

or briefly

$$\Phi = K + K * \Phi.$$

The expression for the kernel $K(t,x,s,y)$, as well as for the "right-hand" side of (23) has a form

$$K(t,x,s,y) = \frac{b(t,x) - b(s,y)}{x} \cdot \frac{\partial p_{b(s,y)}(t,x,s,y)}{\partial x}. \quad (24)$$

3. The last two equations are well-known Volterra's equations. Its (unique) solution may be represented by the form

$$\Phi(t,x,s,y) = K(t,x,s,y) + \sum_{m=1}^{\infty} \Phi_m(t,x,s,y), \quad (25)$$

where

$$\Phi_1(t,x,s,y) = K * K(t,x,s,y)$$

and recursively

$$\Phi_{m+1}(t,x,s,y) = \Phi_m * K(t,x,s,y), \quad m = 1, 2, \dots$$

Define formally $\Phi_0(t,x,s,y) := K(t,x,s,y)$; so that

$$\Phi(t,x,s,y) = \sum_{m=0}^{\infty} \Phi_m(t,x,s,y), \quad (26)$$

It remains to ground the convergence of the series (25) or (26) and made some quantitative estimates.

4. Note first of all that it follows from the condition (15) the estimate

$$|K(t, x, s, y)| \leq H \frac{(|x - y| + |t - s|)^\alpha}{x} \left| \frac{\partial p_{b(s,y)}(t, x, s, y)}{\partial x} \right|. \quad (27)$$

Further, we will use the following well known properties of the modified Bessel's functions: the power series in (10) and the following asymptotical expression

$$z \rightarrow \infty \Rightarrow I_a(z) \sim (2\pi z)^{-1/2} \exp(z). \quad (28)$$

We will use also some estimations relative (modified) Bessel's and gamma functions from the works [1], [2], [3], [5], [6], [22], [29].

Let δ be arbitrary fixed number from an open interval $(0, 1) : \delta \in (0, 1)$. We get synthesizing both the estimates for Bessel's functions (28) and (10) analogously to the work [25]

$$|K(t, x, s, y)| \leq C(\beta, \beta_+) (t - s)^{-1+\alpha/2} \times \frac{H}{\sqrt{e \delta}} p_\beta(t, (1 - \delta)x, s, (1 - \delta)y), 0 < s < t; x, y > 0. \quad (29)$$

5. We have after simple calculations using convolution identity (12)

$$|\Phi_1(t, x, s, y)| \leq C^2(\beta, \beta_+) (1 - \delta)^{-1} (e \delta)^{-1} \frac{H^2 \Gamma^2(\alpha/2)}{\Gamma(\alpha)} \times (t - s)^{\alpha-1} p_\beta(t, (1 - \delta)x, s, (1 - \delta)y),$$

where as ordinary $\Gamma(\cdot)$ denotes the Gamma function.

6. We find by means of an induction

$$|\Phi_m(t, x, s, y)| \leq C^m(\beta, \beta_+) H^{m+1} (1 - \delta)^{-m} (e \delta)^{-m} \frac{\Gamma^{m+1}(\alpha/2)}{\Gamma((m + 1)\alpha/2)} \times (t - s)^{(m+1)\alpha/2-1} p_\beta(t, (1 - \delta)x, s, (1 - \delta)y), m = 0, 1, 2, \dots \quad (30)$$

Therefore, for all the values $\delta \in (0, 1) \Rightarrow$

$$|\Phi(t, x, s, y)| \leq H (t - s)^{-1+\alpha/2} \Gamma(\alpha/2) p_\beta(t, x, s, y) \times \quad (31)$$

$$\sum_{m=0}^{\infty} \frac{H^m C^m(\beta, \beta_+) \Gamma^m(\alpha/2) (1 - \delta)^{-m} (e\delta)^{-m} (t - s)^{m \alpha/2}}{\Gamma((m + 1)\alpha/2)}. \quad (32)$$

Notice that the series in (32) convergent, following the F.S. $p(t, x, s, y)$ for source equation there exists.

Let us estimate the obtained before function $\Phi(t, x, s, y)$. Recall for this purpose briefly the classical definition of the so - called (generalized) Mittag-Leffler functions [40], [41], [51]

$$E_{A,B}(z) \stackrel{def}{=} \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(Ak + B)}.$$

Here $A = \text{const} > 0$, $B \geq 0$. We need only further the following particular case on these functions

$$g_{\alpha}(z) \stackrel{def}{=} \sum_{m=0}^{\infty} \frac{z^m}{\Gamma(m\alpha/2 + \alpha/2)} = E_{\alpha/2, \alpha/2}(z), \quad \alpha \in (0, 1]. \quad (33)$$

We need to use as a values z in (33) only real non - negative values $z \in [0, \infty)$, but the expression in (33) is also defined for all the complex values z .

The asymptotic as well as non-asymptotic behavior of these functions as $|z| \rightarrow \infty$ is investigated in [51], see also [16]. Namely, as $z \rightarrow \infty$, $z \in R_+$

$$E_{A,B}(z) \sim A^{-1} \exp\left(z^{1/A}\right).$$

As a particular case

$$g_{\alpha}(z) \sim \frac{2}{\alpha} \exp\left(z^{2/\alpha}\right), \quad z \rightarrow \infty. \quad (34)$$

Put now

$$v = v_{\delta}(t - s) := H C(\beta, \beta_1) \Gamma(\alpha/2) (1 - \delta)^{-1} (e\delta)^{-1} (t - s)^{\alpha/2},$$

then

$$\tilde{v} := v_{1/2}(t - s) = H C(\beta, \beta_1) \Gamma(\alpha/2) 4 e^{-1} (t - s)^{\alpha/2},$$

$$|\Phi(t, x, s, y)| \leq H (t - s)^{-1+\alpha/2} \Gamma(\alpha/2) p_{\beta}(t, x, s, y) \cdot g_{\alpha}(v_{\delta}(t - s)). \quad (35)$$

Of course

$$|\Phi(t, x, s, y)| \leq \inf_{\delta \in (0,1)} H (t - s)^{-1+\alpha/2} \Gamma(\alpha/2) p_{\beta}(t, x, s, y) \cdot g_{\alpha}(v_{\delta}(t - s)) \quad (36)$$

and

$$|\Phi(t, x, s, y)| \leq H (t - s)^{-1+\alpha/2} \Gamma(\alpha/2) p_{\beta}(t, x, s, y) \cdot g_{\alpha}(v_{1/2}(t - s)). \quad (37)$$

7. Finally, it remains to apply the representation (19) for the fundamental solution $p(t, x, \tau, \xi)$. We estimate

$$\forall \delta \in (0, 1) \Rightarrow p(t, x, \tau, \xi) \leq p^{(\delta)}(t, x, \tau, \xi),$$

where

$$p^{(\delta)}(t, x, \tau, \xi) \stackrel{def}{=} w_{\tau, \xi}(t, x, \tau, \xi) + \int_{\tau}^t d\theta \int_0^{\infty} w_{\theta, \zeta}(t, x, \theta, \zeta) \cdot \Phi(\theta, \zeta, \tau, \xi) d\zeta =$$

$$p_{b(\tau, \xi)}(t, x, \tau, \xi) + \int_{\tau}^t d\theta \int_0^{\infty} p_{b(\theta, \zeta)}(t, x, \theta, \zeta) \cdot \Phi(\theta, \zeta, \tau, \xi) d\zeta \leq$$

$$p_{b(\tau, \xi)}(t, x, \tau, \xi) + C_2(\beta, \beta_+) H (t - s)^{-1+\alpha/2} \Gamma(\alpha/2) p_{\beta}(t, x, \tau, \xi) \cdot g_{\alpha}(v_{\delta}(t - s)). \quad (38)$$

Of course,

$$p(t, x, \tau, \xi) \leq \bar{p}(t, x, \tau, \xi) \stackrel{def}{=} \inf_{\delta \in (0, 1)} p^{(\delta)}(t, x, \tau, \xi), \quad (39)$$

and following

$$p(t, x, \tau, \xi) \leq p^{(1/2)}(t, x, \tau, \xi). \quad (40)$$

On the other hand, $\forall \delta \in (0, 1) \Rightarrow p(t, x, \tau, \xi) \geq$

$$p_{b(\tau, \xi)}(t, x, \tau, \xi) - C_3(\beta, \beta_+) H (t - s)^{-1+\alpha/2} \Gamma(\alpha/2) p_{\beta}(t, x, \tau, \xi) \cdot g_{\alpha}(v_{\delta}(t - s)) \quad (41)$$

Recall that

$$p_{b(\tau, \xi)}(t, x, \tau, \xi) = (t - \tau)^{-1} x^{-b(\tau, \xi)} \xi^{b(\tau, \xi)+1} \exp \left\{ -\frac{x^2 + \xi^2}{2(t - \tau)} \right\} \times$$

$$I_{b(\tau, \xi)} \left(\frac{x\xi}{t - \tau} \right), \quad x, \xi > 0, \quad t > \tau > 0.$$

One can establish analogously from the relation (19) also the estimations for the suitable derivatives $\partial p(t, x, s, y)/\partial t$, $\partial p(t, x, s, y)/\partial x$, $\partial^2 p(t, x, s, y)/\partial x^2$ etc. to ground that the obtained function solved the source problem.

3 Solving of several partial differential equations.

1. Let us return to the source problem (1), (2), and (3).

Proposition 3.1. We suppose in the sequel that the coefficient $b = b(t, x)$ satisfies all the conditions of theorem 2.1. Suppose also that the initial function $f = f(x)$, $x \geq 0$ is continuous and such that

$$\exists \Delta \in (0, 1) \forall x \geq 0 \Rightarrow |f(x)| \leq \exp\left((1 - \Delta)x^2/2\right). \quad (42)$$

Then the (unique) solution of the considered problem has a form

$$u(t, x) = \int_0^\infty p(t, x, s, y) f(y) dy. \quad (43)$$

Proof. Namely, the convergence of the integral in (43) common with its derivatives follows immediately from (47) and from the estimate (38), if we choose for instance $\delta = \Delta/2$.

2. Let us consider now the non - homogeneous equation of the form

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2} + \frac{1 + 2b(t, x)}{x} \frac{\partial u}{\partial x} + g(t, x), \quad (44)$$

$u = u(t, x)$, where $t \geq 0$, $x \in R_+$, under reflection restriction

$$\lim_{x \rightarrow 0^+} x^{1-2b(t, x)} u'_x(t, x) = 0 \quad (45)$$

and zero initial value

$$\lim_{t \rightarrow s^+} u(t, x) = 0. \quad (46)$$

One can state analogously the following assertion.

Proposition 3.2. We suppose that the coefficient $b = b(t, x)$ again satisfies all the conditions of theorem 2.1. Suppose also that the right - hand side function $g = g(t, x)$, $x \geq 0$ is continuous and such that

$$\exists \Delta \in (0, 1) \forall x \geq 0 \Rightarrow |g(t, x)| \leq \exp\left((1 - \Delta)x^2/2\right). \quad (47)$$

Then the (unique) solution of the considered problem has a form

$$u(t, x) = \int_s^t d\theta \int_0^\infty p(t, x, \theta, y) g(\theta, y) dy. \quad (48)$$

4 Variable Bessel's random processes.

Recall, see [7], [39], [50], that the Markov diffusion separable random process $\xi = \xi(t)$, $t, \xi(t) \geq 0$ having an infinitesimal operator (generator) of the form

$$L_a u = \frac{1}{2} \frac{\partial^2}{\partial x^2} + \frac{1+2a}{x} \frac{\partial}{\partial x}, \quad a = \text{const} \in (-1, 0), \quad x \in \mathbf{R}_+, \quad a = \text{const}$$

with or without reflection at the origin, is named as (ordinary) *Bessel's process*.

More information about these processes may be found in the articles [7], [39], [50] etc. A very interest applications of these processes is represented in a recent article [54].

An interest example. Let $w = w(t)$, $t \geq 0$, $w(t) \in \mathbf{R}^d$, $d = 1, 2, \dots$ be a multivariate Brownian motion. Define for Euclidean norm $\|x\|$, $x \in \mathbf{R}^d$ the random process $\eta = \eta(t)$ as the (Euclidean) norm of the Brownian motion $\eta(t) = \|w(t)\|$. Then $\eta(t)$ is the Bessel's random process with correspondent generator

$$M_n u = \frac{1}{2} \frac{\partial^2 u}{\partial x^2} + \frac{n-1}{2x} \frac{\partial u}{\partial x}. \quad (49)$$

Definition 4.1. The random diffusion process $\theta(t)$, $\theta(s) = x_0 = \text{const} > 0$, $x \in \mathbf{R}_+$, having appeared before (4) the infinitesimal operator

$$Q[b]_{t,x} u \stackrel{\text{def}}{=} \frac{\partial^2 u}{\partial x^2} + \frac{1+2b(t,x)}{x} \frac{\partial u}{\partial x} \quad (50)$$

in the domain $t, x > 0$ will be named as *variable Bessel's random process*.

We can and will assume that the coefficient $b = b(t, x)$ satisfies all the conditions of theorem 2.1., therefore $\theta(t)$ there exists and obeys the transfer density of certain random Markov's diffusion process $\xi = \xi(t)$, $t, \xi(t) > 0$; defined on the suitable sufficiently rich probability space $(\Omega = \{\omega\}, B, \mathbf{P})$, and investigated before.

We intend to prove in this section that this r.p. $\xi = \xi(t)$ is *subgaussian* and continuous almost everywhere (i.e., with probability one).

Let us recall at first some facts from the theory of subgaussian r.v. and r.p. More information may be found in a works [9], [10], [26], [27], [30], [35], [45] and so one.

The numerical valued r.v. $X = X(\omega)$ is said to be subgaussian, if there exists positive finite constants C, v , such that

$$\forall u \geq 0 \Rightarrow \mathbf{P}(|X| > u) \leq C e^{-v u^2}. \quad (51)$$

These r.v. forms a so-called Birnbaum-Orlicz Banach complete space relative the classical Luxemburg norm

$$\|X\|_{\psi(2)} \stackrel{\text{def}}{=} \inf \left\{ s, s > 0, \mathbf{E} \exp(|X/s|^2) \leq 2 \right\}.$$

Equivalent (up to multiplicative constant) definition:

$$\|X\|_{\psi(2)} \stackrel{def}{=} \sup_{p \geq 1} \left[\frac{[\mathbf{E}|X|^p]^{1/p}}{\sqrt{p}} \right].$$

By definition the random process $X = X(t)$, $t \geq 0$ is named subgaussian, iff for all the values $t \geq 0$

$$\|X(t)\|_{\psi(2)} < \infty.$$

Alike, the random process $X = X(t)$, $t \geq 0$ is named uniform subgaussian, iff

$$\sup_{t \geq 0} \|X(t)\|_{\psi(2)} < \infty.$$

Analogously, the random process $X = X(t)$, $t \geq 0$ is named local subgaussian, iff

$$\forall T \in (0, \infty) \sup_{t \in [0, T]} \|X(t)\|_{\psi(2)} < \infty.$$

Theorem 4.1. Let a deterministic variable T be sufficiently great: $T > e$. Suppose in addition to the conditions of theorem 2.1 that the "initial value" for $\theta(\cdot)$, say, $\theta(0)$ is subgaussian; for instance, may be a deterministic constant. The described in definition 4.1 (50) random process $\theta = \theta(t)$, $t \geq 0$ is local subgaussian and has alike the Levi's Brownian motion module of continuity:

$$\exists \nu = \nu(\omega) \in G\psi(2), |\theta(t) - \theta(s)| \leq \nu \cdot \sqrt{|t - s| \ln(2 + 1/|t - s|)}, t, s \in (0, 1). \quad (52)$$

Further, there exists a subgaussian positive r.v. τ , $\|\tau\|_{\psi(2)} = 1$ such that

$$\max_{t \in [0, T]} |\theta(t)| \leq C_1(\beta) \tau \sqrt{T \ln T}. \quad (53)$$

Proof. Suppose for beginning that $\|\theta(s)\|_{\psi(2)} < \infty$. We apply the obtained estimate (38) and obtain after simple calculations

$$\|\theta(t) - \theta(s)\|_{\psi(2)} \leq C_2(\beta) \sqrt{|t - s|}. \quad (54)$$

It remains to use the results of the sections 3.12 and 3.14 of the monograph [45], devoting namely the subgaussian random fields.

The results (52) and (53) are essentially non-improvable, for example, for the classical Bessel's processes (49).

Acknowledgement. The first author has been partially supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale

di Alta Matematica (INdAM) and by Università degli Studi di Napoli Parthenope through the project “sostegno alla Ricerca individuale” (triennio 2015-2017).

The second author is very grateful to Molchanov S.A. for the statement of the considered problem and valuable guidance.

References

- [1] **M. Abamowitz and I. A. Stegun (Eds.)**. *Handbook of Mathematical Functions*. Dover, New York, 1965, pp. 3748, 4435. Google Scholar
- [2] **Abramowitz, M. and Stegun, I. A. (Eds.)**. ”Modified Bessel Functions I and K.” 9.6 in *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, 9th printing. New York: Dover, pp. 374-377, 1972.
- [3] **Arfken, G.** *Modified Bessel Functions, $I_n u(x)$ and $K_n u(x)$* . Section 11.5 in *Mathematical Methods for Physicists*, 3rd ed. Orlando, FL: Academic Press, pp. 610-616, 1985.
- [4] **Vlad Bally and Arturo Kohatsu-Higa**. *A probabilistic interpretation of the parametrix method*. *The Annals of Applied Probability*, 2015, Vol. 25, No. 6, 3095-3138, DOI: 10.1214/14-AAP1068.
- [5] **Barkat Ali Bhao**. *Onn functional inequalities for the psi function*. arXiv:1210.0046v2 [math.CA] 7 Sep 2015
- [6] **H.Bateman at all**. *Higer transcendental functions*. Volume 2. Mc GRAW HILL book company, New York, London, Toronto, 1953.
- [7] **Kamil Bogus, Jacek Maleck**. *Sharp Estimates of Transition Probability Density for Bessel Process in Half-Line*. *Potential Anal* (2015) 43:122 DOI 10.1007/s11118-015-9461-x
- [8] **Sig Byun, Jung-Tae Park, and Pilsoo Shin**. *Global regularity for degenerate/singular parabolic equation involving measure data*. arXiv:1903.07843v1 [math.AP] 19 Mar 2019
- [9] **V.V. Buldygin and Yu.V. Kozachenko**. *Subgaussian random variables*. *Ukrainian Math. J.* **32**, (1980), 483-489.
- [10] **V.V. Buldygin and Y. Kozachenko**. *Metric Characterization of Random Variables and Random Processes*. Volume 188 of *Translations of Mathematical Monographs*. American Mathematical Society, 2000
- [11] **Corielli, F., Foschi, P. and Pascucci, A.** (2010). *Parametrix approximation of diffusion transition densities*. *SIAM J. Financial Math.*, **1**, 833-867. MR2736914

- [12] **Deck Thomas and Susanne Kruse.** *Parabolic Differential Equations with Unbounded Coefficients A Generalization of the Parametrix Method.* Acta Applicandae Mathematicae 74: 71-91, 2002. Kluwer Academic Publishers. Printed in the Netherlands.
- [13] **F.Dressel.** *The fundamental solution of the parabolic equation.* Duke Math. J., 7, No. 4, 186-203 (1940).
- [14] **F.Dressel.** *The fundamental solution of the parabolic equation. II.* Duke Math. J., 13, No. 1, 61-70 (1946).
- [15] **J.F. van Diejen, L. Vinet.** *Calogero-Sutherland-Moser Models.* CRM Series in Mathematical Physics, Springer, Berlin, 2000.
- [16] **M.A.Evgrafov.** *Asymptotic estimates and entire functions.* Gordon and Breach, 1961
- [17] **S. D. Eydelman.** *On fundamental solutions of parabolic systems. I-II,* Mat. Sb., 53 (95), No. 1, 73-136, (1961).
- [18] **Maria Rosaria Formica, Eugeny Ostrovsky, Leonid Sirota.** *Fundamental solution for Cauchy initial value problem for parabolic PDEs with discontinuous unbounded first-order coefficient at the origin. Extension of the classical parametrix method.*
arXiv:1906.05880v1 [math.AP] 13 Jun 2019
- [19] **Friedman, A.** (1964). *Partial Differential Equations of Parabolic Type.* Prentice-Hall, Englewood Cliffs, NJ. MR0181836
- [20] **E. Gagliardo.** *Teoremi di esistenza e di unicita per problemi al contorno relativi ad equazioni paraboliche lineari e quasi lineari in n variabili,* Ricerche Matem., 5, 239-257 (1956).
- [21] **Glagoleva, R. J.** *A priori estimate of the Hölder norm and the Harnack inequality for the solution of a second order linear parabolic equation with discontinuous coefficients.* Mat. Sb. (N.S.) 76 (118), (1968), 167-185.
- [22] **I. S. Gradshteyn and I. M. Ryzhik.** *Table of Integrals, Series and Products.* Academic, New York, 1965, p. 307, Eq. 3.222.
- [23] **Gruber.** *M.Harnack inequalities for solutions of general second order parabolic equations and estimates of their Hölder constants.* Math. Z., 185, 1, (1984), 23-43.
- [24] **A. M. Ilyin.** *Degenerate elliptic and parabolic equations,* Mat. Sb., 50 (94), No. 4, 443-498 (1960).
- [25] **A. M. Ilyin, A. S. Kalashnikov, and O. A. Oleynik.** *Linear second-order partial differential equations of the parabolic type.* Journal of Mathematical Sciences, Vol. 108, No. 4, 2002.

- [26] **J.P. Kahane.** *Propri et les locales des fonctions a s ries de Fourier aleatoires.* Stud. Math.19(1960), 1 - 25.
- [27] **J.P. Kahane.** *Some random series of functions.* 2nd ed. Cambridge University Press, London, 1985.
- [28] **L.I.Kamynin and V.N.Maslennikov.** *On the maximum principle for a parabolic equation with discontinuous coefficients.* Sib. Mat. Zh., 2, No. 3, 384-399 (1961).
- [29] **M. Kazeminia, M. Mehrjoo.** *On the Derivatives of Bessel and Modified Bessel Functions with Respect to the Order and the Argumant.* International Research Journal of Applied and Basic Sciences, 2013, Available online at www.irjabs.comISSN 2251-838X / Vol, 4 (12):4127-4133 Science Explorer Publications.
- [30] **Yu.V. Kozachenko, E.I. Ostrovskii.** *Banach spaces of random variables of sub-Gaussian type.* Teor. Veroyatnost. i Mat. Statist., **32**, 42-53, **77**, 1985.
- [31] **M. A. Krasnoselskiy and S. G. Kreyn.** *On differential equations with nonbounded operators in Banach spaces.* Dokl. Akad. Nauk SSSR, 111, No. 1, 19-22, (1956).
- [32] **Krylov, N. V.** *Lectures on elliptic and parabolic equations in Hölder spaces.* Vol. 12 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 1996.
- [33] **O. A. Ladyzhenskaya, V. A. Solonnikov and N. N. Uraltseva.** *Linear and Quasilinear Equations of Parabolic Type.* Nauka, Moscow, 1967.
- [34] **P. Lax and A. Milgram.** *Parabolic equations.* Ann. Math. Studies, 33, 167-190, (1954).
- [35] **Ledoux, Michel; Talagrand, Michel.** (1991). *Probability in Banach Spaces.* Springer-Verlag.
- [36] **E.E. Levi.** *Sulle equazioni lineari alle derivate parziali totalmente ellittiche.* Rend. R. Acc. Lincei, Classe Sci. (V) , **16**, (1907).
- [37] **E.E.Levi.** *Sulle equazioni lineari totalmente ellittiche alle derivate parziali.* Rend. Circ. Mat. Palermo , 24 (1907) pp. 275-317.
- [38] **Lieberman, G.M.** (1996). *Second Order Parabolic Partial Differential Equations.* World Scientific Publishing House, New-York, 1996.
- [39] **H.P.McKean,Jr.** *The Bessel motion and a singular integral equation.* Mem.Coll.Sci.Univ.Kyoto,SerA,Math.,33(1960),317-322
- [40] **Mittag- Leffler G.** *Sur la nouvelle fonction Ea.* C.R. Acad Sci Paris., 1903; 137, 554-558.

- [41] **Magnus G. Mittag-Leffler.** *Sopra la funzione $Ea(x)$.* Rend. Accad. Lincei, 13(5): 3 - 5, 1904.
- [42] **S.A.Molchanov, E.Ostrovskii.** *Symmetric stable processes as traces of degenerate diffusion processes.* Teor. Veroyatnost. i Primenen., 1969, Volume 14, Issue 1, 127 - 130.
- [43] **Nash, J.** (1958). *Continuity of solutions of parabolic and elliptic equations.* Amer. J. Math., 80: 931-954.
- [44] **Ashkan Nikeghbali** *Some random times and martingales associated with $BES_0(\delta)$ processes ($0 < \delta < 2$).*
arXiv:math/0505423v2 [math.PR] 3 Aug 2007
- [45] **Ostrovsky E.I.** *Exponential Estimations for Random Fields.* Moscow-Obninsk, OINPE, 1999 (Russian).
- [46] **Paul-Eric Chaudru de Raynal and Stephane Menozzi Enrico Priola.** *Schauder estimates for drifted fractional operators in the supercritical case.* Internet preprint, February 2019.
- [47] **Revuz, D.; Yor, M.** (1999). *Continuous Martingales and Brownian Motion.* Berlin: Springer. ISBN 3-540-52167-4.
- [48] **A.A.Samarskiy.** *Equations of the parabolic type with discontinuous coefficients.* Dokl. Akad. Nauk SSSR, 121, No. 2, 255-228 (1958).
- [49] **Sachkov V.N.** *Combinatorial Methods in Discrete Mathematics.* Cambridge University Press, (1996), 2010, Online ISBN: 9780511666186 <https://doi.org/10.1017/CBO9780511666186>
- [50] **Tokuzo Shiga and Shinzo Watanabe.** *Bessel diffusions as a one-parameter family of diffusion processes.* Z. Wahrscheinlichkeitstheorie und Verw. Gebiete 27 (1973), 37-46. MR 0368192
- [51] **Shukla A.K., Prajapati J.C.** *On a generalization of Mittag-Leffler function and its properties.* J. Math Anal Appl. 2007; **336(2)**, 797-811. doi: 10.1016/j.jmaa.2007.03.018. [CrossRef].
- [52] **L.N.Slobodetskiy.** *On the fundamental solution and the Cauchy problem for a parabolic system.* Mat. Sb., 46 (88), No. 2, 229-258, (1958).
- [53] **P. E. Sobolevskiy.** *On equations of parabolic type with nonbounded operators in Banach spaces.* Trudy MMO, 10, 297-350, (1961).
- [54] **Michael Voit, Jeannette H.C.Woerner.** *Functional Central Limit Theorem in the Freezing Regime.* arXiv:1901.08390v1 [math.PR] 24 Jan 2019.

- [55] **Qi.S.Zhang.** *Local estimates on two linear parabolic equations with singular coefficients.*
arXiv:1905.13329 v1 [math.AP] 30 May 2019