

ON PARITY UNIMODALITY OF q -CATALAN POLYNOMIALS

GUOCE XIN* AND YUEMING ZHONG

ABSTRACT. A polynomial $A(q) = \sum_{i=0}^n a_i q^i$ is said to be unimodal if $a_0 \leq a_1 \leq \dots \leq a_k \geq a_{k+1} \geq \dots \geq a_n$. We investigate the unimodality of rational q -Catalan polynomials, which is defined to be $C_{m,n}(q) = \frac{1}{[n+m]} \begin{bmatrix} m+n \\ n \end{bmatrix}_q$ for a coprime pair of positive integers (m, n) . We conjecture that they are unimodal with respect to parity, or equivalently, $(1+q)C_{m+n}(q)$ is unimodal. By using generating functions and the constant term method, we verify our conjecture for $m \leq 5$ in a straightforward way.

Mathematics Subject Classifications: 05A15, 05A20, 05E05

Keywords: rational Dyck paths; rational q -Catalan polynomials; unimodal sequences.

1. INTRODUCTION

We will consider the unimodality of some symmetric polynomials. A sequence $a_0, a_1, \dots, a_n$ is said to be symmetric if $a_i = a_{n-i}$ for all i . It is said to be unimodal if there is a k such that $a_0 \leq a_1 \leq \dots \leq a_k \geq a_{k+1} \geq \dots \geq a_n$. It is said to be unimodal with respect to parity if $a_0, a_2, \dots$ and $a_1, a_3, \dots$ are both unimodal. A polynomial $P(q) = a_0 + a_1 q + \dots + a_n q^n$ of degree n is said to be symmetric (resp., unimodal) if its coefficient sequence $a_0, a_1, \dots, a_n$ is symmetric (resp., unimodal).

Stanley gave a nice survey [22] on various methods for showing that a sequence is unimodal (or log-concave which we will not discuss here). A classical example is the following.

Theorem 1. *The Gaussian polynomial $G_{m,n}(q)$ is symmetric and unimodal, where*

$$G_{m,n}(q) = \begin{bmatrix} m+n \\ m \end{bmatrix}_q = \frac{[m+n]!}{[m]![n]},$$

with the classical q -notation $[n] = \frac{1-q^{n+1}}{1-q}$, $[n]! = [n][n-1] \cdots [1]$.

This important result has many proofs. See [4, 6, 7, 17, 19, 22, 29, 30]. It is an outstanding open question to find an explicit order matching proof for the unimodality. Though the Gaussian polynomials have been extensively studied, the closely related q -Catalan polynomials are less studied.

The q -Catalan polynomials (or numbers) $C_n(q)$ we are discussing here are defined by

$$C_n(q) = \frac{1}{[n+1]} \begin{bmatrix} 2n \\ n \end{bmatrix}_q = \frac{[2n]!}{[n+1]![n]}.$$

It starts with $C_0(q) = C_1(q) = 1$, $C_2(q) = 1 + q^2$, $C_3(q) = 1 + q^2 + q^3 + q^4 + q^6$, $C_4(q) = 1 + q^2 + q^3 + 2q^4 + q^5 + 2q^6 + q^7 + 2q^8 + q^9 + q^{10} + q^{12}$. Clearly, one sees that $C_n(q)$ is symmetric, but not unimodal. However, we have the following conjecture.

Conjecture 2. *The q -Catalan polynomials $C_n(q)$ are unimodal with respect to parity.*

We find that Conjecture 2 can be extended for rational q -Catalan polynomials. For a pair (m, n) of positive integers, define

$$C_{m,n}(q) = \frac{1}{[n+m]} \begin{bmatrix} m+n \\ n \end{bmatrix}_q = \frac{1}{[m]} \begin{bmatrix} m+n-1 \\ n \end{bmatrix}_q.$$

When m and n are coprime to each other, i.e., $\gcd(m, n) = 1$, $C_{m,n}(q)$ is known to be in $\mathbb{N}[q]$ (a polynomial with nonnegative coefficients), and is called the (m, n) rational q -Catalan polynomials (or q -Catalan numbers). See, e.g., [13].

Conjecture 3. *For a coprime pair of positive integers (m, n) , the (m, n) -rational q -Catalan polynomials $C_{m,n}(q)$ are unimodal with respect to parity.*

When $\gcd(m, n) > 1$, $C_{m,n}(q)$ is usually not a polynomial, while it has been shown that $\bar{C}_{m,n}(q) = [\gcd(m, n)]C_{m,n}(q)$ is in $\mathbb{N}[q]$. See, e.g., [1, 9, 12].

Conjecture 4. *For a pair of positive integers (m, n) , the polynomial $\bar{C}_{m,n}(q)$ is unimodal with respect to parity.*

Conjecture 4 includes Conjecture 3 as a special case, since $\bar{C}_{m,n}(q)$ reduces to $C_{m,n}(q)$ when $\gcd(m, n) = 1$. We state the latter separately because the combinatorial meaning of $C_{m,n}(q)$ is much more elegant as we will explain later in Section 5. Conjecture 3 includes Conjecture 2 as a special case, since $C_{n+1,n}(q) = C_n(q)$. We state the latter separately because $C_n(q)$ has a different combinatorial interpretation. See Section 5.

We have verified Conjecture 4 for $m, n \leq 180$ by Maple. Observe that $\bar{C}_{m,km}(q) = [m]C_{m,km} = \begin{bmatrix} km+m-1 \\ km \end{bmatrix}_q$, which by Theorem 1 is indeed unimodal. We will prove this conjecture for $m \leq 5$. Our method is to compute the corresponding generating functions by means of the constant term method. It turns out that for $m \leq 5$, the positivity is transparent in view of their generating functions.

This paper is organized as follows. Section 1 is this introduction. Section 2 introduces the basic concepts and idea for our proof. The unimodality conjectures are translated by using their generating functions. Section 3 tries direct computation, which already becomes complicated for $m = 4$. Section 4 uses the constant term method to compute the corresponding generating functions, whose positivity is transparent and hence proves Conjecture 3 for $m \leq 5$. In Section 5, we discuss possible representation approach for settling these conjectures. We also introduce the combinatorial interpretations of q -Catalan polynomials.

2. PRELIMINARY

A Laurent polynomial $L(q)$ is said to be symmetric if $L(q^{-1}) = L(q)$, and is said to be anti-symmetric if $L(q^{-1}) = -L(q)$.

Define the normalization of a polynomial $P(q)$ of degree n by

$$\mathcal{N}P(q) := \mathcal{N}(P(q)) = P(q^2)q^{-n}, \quad \mathcal{N}(0) = 0.$$

Then the symmetry of $P(q)$ (i.e., $P(q) = q^n P(q^{-1})$) is transformed to the more natural symmetry of the Laurent polynomial $\mathcal{N}P(q)$. The following properties are easy to verify and will be used without mentioning:

- (1) For polynomials $P(q)$ and $Q(q)$, we have $\mathcal{N}(P(q)Q(q)) = \mathcal{N}(P(q))\mathcal{N}(Q(q))$.
- (2) The product of two symmetric Laurent polynomials is still symmetric.
- (3) If $L_1(q^{-1}) = L_1(q)$ and $L_2(q^{-1}) = -L_2(q)$, then $L_1(q)L_2(q)|_{q=q^{-1}} = -L_1(q)L_2(q)$.

We will also use the following linear operators on Laurent polynomials in $\mathbb{Q}[q, q^{-1}]$.

$$\begin{aligned} \text{PT}_q \sum_i a_i q^i &= \sum_{i>0} a_i q^i, & (\text{extracting the positive exponent terms}) \\ \text{CT}_q \sum_i a_i q^i &= a_0, & (\text{extracting the constant term}) \\ \text{NT}_q \sum_i a_i q^i &= \sum_{i<0} a_i q^i, & (\text{extracting the negative exponent terms}) \end{aligned}$$

These operators clearly extend to $\mathbb{Q}[q, q^{-1}][[x]]$, the ring of power series in x with coefficients Laurent polynomials in q . Indeed, they act coefficient wise in x .

The following lemma is transparent.

Lemma 5. *Suppose $P(q)$ is a symmetric polynomial of degree n . Then*

- (1) $P(q)$ is unimodal if and only if $PT_q \mathcal{N}((q-1)P(q)) \in \mathbb{N}[q]$.
- (2) $P(q)$ is unimodal with respect to parity if and only if

$$PT \mathcal{N}(q^2 - 1)P(q) = PT(q^2 - q^{-2})\mathcal{N}P(q) \in \mathbb{N}[q].$$

Thus Conjecture 3 can be restated as follows.

Conjecture 3^a. *For a coprime pair of positive integers (m, n) , the polynomial*

$$\begin{aligned} PT \mathcal{N}(q^2 - 1)C_{m,n}(q) &= PT(q^2 - q^{-2})C_{m,n}(q^2)q^{-(m-1)(n-1)} \\ (1) \quad &= PT(q^2 - q^{-2}) \frac{(1 - q^{2m+2})(1 - q^{2m+4}) \cdots (1 - q^{2m+2n-2})}{(1 - q^4)(1 - q^6) \cdots (1 - q^{2m})} q^{-(m-1)(n-1)} \end{aligned}$$

has nonnegative integer coefficients.

Our point is that it is usually easier to consider generating functions. Let

$$(2) \quad F_m(x, q) := \sum_{n \geq 0} (q^2 - q^{-2}) \mathcal{N}C_{m,n}(q) x^n.$$

Note that the coefficients in x are not always Laurent polynomials in q . We take $F_m(x, q)$ as an element in $\mathbb{Q}((q))[[x]]$, the ring of power series in x with coefficients Laurent series in q .

For integers $m > r \geq 0$, let $X_{m,r}$ be the linear operator acting on $\mathbb{Q}(q)[[x]]$ by

$$(3) \quad X_{m,r} \sum_{n \geq 0} a_n(q) x^n = \sum_{k \geq 0} a_{km+r}(q) x^k. \quad (\text{extracting the terms with special exponents})$$

When $\gcd(m, r) = 1$, $X_{m,r}F_m(x, q) \in \mathbb{Q}[q, q^{-1}][[x]]$.

Then Conjecture 3^a is transformed as follows.

Conjecture 3^b. Let $m > r$ be positive integers with $\gcd(m, r) = 1$. Then

$$PT_q X_{m,r} F_m(x, q) = X_{m,r} PT_q F_m(x, q)$$

is a power series in x with coefficients in $\mathbb{N}[q]$.

The case $\gcd(m, r) = d$ is a little complicated. We need to consider the generating function

$$\begin{aligned} \sum_{k \geq 0} \mathcal{N}((q^2 - 1)[d]C_{m, km+r}(q))x^k &= \sum_{k \geq 0} \frac{q^d - q^{-d}}{q - q^{-1}}(q^2 - q^{-2})\mathcal{N}(C_{m, km+r}(q))x^k \\ &= \frac{q^d - q^{-d}}{q - q^{-1}}X_{m,r} \sum_{n \geq 0} (q^2 - q^{-2})\mathcal{N}(C_{m,n}(q))x^n \\ &= \frac{q^d - q^{-d}}{q - q^{-1}}X_{m,r}F_m(x, q). \end{aligned}$$

Thus Conjecture 4 can be transformed as follows.

Conjecture 4^b. Let $m > r$ be nonnegative integers with $\gcd(m, r) = d$. Then

$$PT_q X_{m,r} \frac{q^d - q^{-d}}{q - q^{-1}}F_m(x, q) = X_{m,r} PT_q \frac{q^d - q^{-d}}{q - q^{-1}}F_m(x, q)$$

is a power series in x with coefficients in $\mathbb{N}[q]$.

We remark that $\frac{q^d - q^{-d}}{q - q^{-1}}F_m(x, q)$ is a power series in x with coefficient Laurent series in q , so we need to extend the PT_q operator. See Section 4.

To our surprise, $F_m(x, q)$ has a product formula as follows.

Proposition 6. For any positive integer m , we have

$$(4) \quad F_m(x, q) = (q^2 - q^{-2}) \frac{(q - q^{-1})}{(q^m - q^{-m})} \prod_{i=0}^{m-1} \frac{1}{(1 - xq^{1-m} \cdot q^{2i})}$$

Proof. The proposition is indeed a consequence of the following well-known identity. See, e.g., [23, pp. 272].

$$(5) \quad \frac{1}{(1-x)(1-xq)(1-xq^2) \cdots (1-xq^m)} = \sum_{n \geq 0} \begin{bmatrix} m+n \\ m \end{bmatrix}_q x^n$$

We have

$$\begin{aligned} &\sum_{n \geq 0} q^{\frac{-(n-1)(m-1)-2}{2}}(q^2 - 1)C_{m,n}(q)x^n \\ &= \sum_{n \geq 0} q^{\frac{-(n-1)(m-1)-2}{2}}(q^2 - 1) \frac{1-q}{1-q^m} \begin{bmatrix} m+n-1 \\ m-1 \end{bmatrix}_q x^n \\ &= q^{\frac{m-3}{2}}(q^2 - 1) \frac{1-q}{1-q^m} \sum_{n \geq 0} \begin{bmatrix} m+n-1 \\ m-1 \end{bmatrix}_q (q^{\frac{-(m-1)}{2}}x)^n \\ &= q^{\frac{m-3}{2}}(q^2 - 1) \frac{1-q}{1-q^m} \prod_{i=0}^{m-1} \frac{1}{(1 - xq^{\frac{-(m-1)}{2}}q^i)}. \end{aligned}$$

We can get

$$\begin{aligned}
F_m(x, q) &= \sum_{n>0} (q^2 - q^{-2}) \mathcal{N}C_{m,n}(q) x^n \\
&= \sum_{n>0} (q^2 - q^{-2}) C_{m,n}(q^2) q^{-(m-1)(n-1)} x^n \\
&= \sum_{n>0} q^{-(n-1)(m-1)-2} (q^4 - 1) C_{m,n}(q^2) x^n \\
&= q^{m-3} (q^4 - 1) \frac{1 - q^2}{1 - q^{2m}} \prod_{i=0}^{m-1} \frac{1}{(1 - xq^{1-m} \cdot q^{2i})} \\
&= (q^2 - q^{-2}) \frac{(q - q^{-1})}{(q^m - q^{-m})} \prod_{i=0}^{m-1} \frac{1}{(1 - xq^{1-m} \cdot q^{2i})}.
\end{aligned}$$

■

3. DIRECT COMPUTATION

Conjecture 3 can be verified directly for $m = 3$, but already becomes complicated for $m = 4$.

3.1. **The case $m = 3$.** In this case, we have the following explicit expansion.

Proposition 7. *We have*

$$(1) \quad (q^2 - 1)C_{3,n}(q) = \begin{cases} q^{3k+1}(-\sum_{i=0}^k q^{-(3i+1)} + \sum_{i=0}^k q^{3i+1}), & \text{if } n = 3k + 1; \\ q^{3k+2}(-\sum_{i=0}^k q^{-(3i+2)} + \sum_{i=0}^k q^{3i+2}), & \text{if } n = 3k + 2. \end{cases}$$

Consequently, Conjecture 3 holds true for $m = 3$.

Proof. By direct computation, we have

$$\begin{aligned}
(q^2 - 1)C_{3,n}(q) &= \frac{(q^2 - 1)(1 - q)}{1 - q^{n+3}} \frac{(1 - q^{n+1})(1 - q^{n+2})(1 - q^{n+3})}{(1 - q)(1 - q^2)(1 - q^3)} \\
&= -\frac{(1 - q^{n+1})(1 - q^{n+2})}{(1 - q^3)}
\end{aligned}$$

When $n = 3k + 1$, we have

$$\begin{aligned}
(q^2 - 1)C_{3,3k+1}(q) &= -\frac{(1 - q^{3k+2})(1 - q^{3k+3})}{(1 - q^3)} \\
&= (q^{3k+2} - 1)(1 + q^3 + q^6 + \cdots + q^{3k}) \\
&= q^{3k+1} \left(-\sum_{i=0}^k q^{-(3i+1)} + \sum_{i=0}^k q^{3i+1} \right).
\end{aligned}$$

This proves Conjecture 3 for $(m, n) = (3, 3k + 1)$.

When $n = 3k + 2$, we have

$$\begin{aligned}
 (q^2 - 1)C_{3,3k+2}(q) &= -\frac{(1 - q^{3k+3})(1 - q^{3k+4})}{(1 - q^3)} \\
 &= (q^{3k+4} - 1)(1 + q^3 + q^6 + \cdots + q^{3k}) \\
 &= q^{3k+2} \left(-\sum_{i=0}^k q^{-(3i+2)} + \sum_{i=0}^k q^{3i+2} \right).
 \end{aligned}$$

This proves Conjecture 3 for $(m, n) = (3, 3k + 2)$. ■

3.2. The case $m = 4$. This case is already complicated. We have

$$(q^2 - 1)C_{4,n}(q) = -\frac{(1 - q^{n+1})(1 - q^{n+2})(1 - q^{n+3})}{(1 - q^3)(1 - q^4)}.$$

We can have explicit polynomial representation, but that will not help to prove our conjecture. For instance, if $n = 12k + 1$, then

$$\begin{aligned}
 (q^2 - 1)C_{4,12k+1}(q) &= -\frac{(1 - q^{12k+2})(1 - q^{12k+3})(1 - q^{12k+4})}{(1 - q^3)(1 - q^4)} \\
 &= (q^{12k+2} - 1) \cdot \sum_{i=0}^{4k} q^{3i} \cdot \sum_{j=0}^{3k} q^{4j}.
 \end{aligned}$$

Now it is unclear why its coefficients in q^r is negative for $r \leq \frac{3n-1}{2}$.

4. THE GENERATING FUNCTION METHOD

4.1. Basic idea. We illustrate the idea by redoing the case $m = 3$, $n = 3k + 1$. Consider the generating function

$$\begin{aligned}
 \sum_{k \geq 0} (q - q^{-1})q^{-3k}C_{3,3k+1}(q)x^k &= \sum_{k \geq 0} -q^{-3k-1} \frac{(1 - q^{3k+2})(1 - q^{3k+3})}{(1 - q^3)} x^k \\
 &= \frac{1}{1 - q^3} \sum_{k \geq 0} (-q^{-3k-1} + q + q^2 - q^{3k+4}) x^k \\
 &= \frac{1}{1 - q^3} \left(-\frac{q^{-1}}{1 - q^{-3}x} + \frac{q^2 + q}{1 - x} - \frac{q^4}{1 - q^3x} \right) \\
 &= \frac{q(1 - q)(1 + q)(x + q + q^2x)}{(1 - x)(x - q^3)(1 - q^3x)}.
 \end{aligned}$$

By taking partial fraction decompositions in q , we obtain:

$$\frac{q(1 - q)(1 + q)(x + q + q^2x)}{(1 - x)(x - q^3)(1 - q^3x)} = \frac{q^2}{(q^3 - x)(x - 1)} + \frac{q}{(q^3x - 1)(x - 1)}.$$

When expanding as a power series in x , the first term has only negative powers in q and the second term has only positive powers in q :

$$\begin{aligned}\frac{q^2}{(q^3 - x)(x - 1)} &= \frac{-q^{-1}}{(1 - q^{-3}x)(1 - x)} = -q^{-1} \left(\sum_{i \geq 0} x^i \right) \left(\sum_{i \geq 0} (q^{-3}x)^i \right), \\ \frac{q}{(q^3x - 1)(x - 1)} &= q \left(\sum_{i \geq 0} x^i \right) \left(\sum_{i \geq 0} (q^3x)^i \right).\end{aligned}$$

Thus by extracting positive powers in q , we obtain

$$\sum_{k \geq 0} \text{PT}_q(q - q^{-1})q^{-3k}C_{3,3k+1}(q)x^k = \frac{q}{(q^3x - 1)(x - 1)} \in \mathbb{N}[q][[x]].$$

Conjecture 3^b thus holds for the case $(m, r) = (3, 1)$.

The case $(m, r) = (3, 2)$ can be done similarly.

Extracting positive powers in q of a general class of rational series can be done systematically by the constant term method. The resulting rational function turns out to be trivially positive for $m \leq 5$.

4.2. The constant term method. Constant term extraction or residue computation has a long history. See, e.g., [26] for further references. The fundamental problem we are concerned here is to compute the constant term of in a set of variables of a formal series in the field of iterated Laurent series $K = \mathbb{Q}((x_n)) \cdots ((x_1))$, which is called the working field. The reader is referred to [9] for the original development of the field of iterated Laurent series. Here we only recall that K defines a total ordering $0 < x_1 < x_2 < \cdots < x_n < 1$ on the variables (more precisely, a total group order on its monomials), which can be formally treated as $0 \ll x_1 \ll x_2 \ll \cdots \ll x_n \ll 1$. Every rational function has a unique series expansion in K . We will focus on the class of Elliott-rational functions, which are rational functions whose denominators are the product of binomials. It is known that the constant term of an Elliott-rational function is still an Elliott-rational function. Efficient algorithms have been developed to evaluate this type of constant terms, such as the Omega Mathematica package [2, 3], Ell Maple package [27] developed from Algebraic Combinatorics. See [25] for further references on algorithmic development from Computational Geometry and Algebraic Combinatorics.

We will use the first author's (updated) Ell2 Maple package. We use a list $xvar = [x_1, x_2, \dots, x_n]$ to specify the working field $\mathbb{Q}((x_n)) \cdots ((x_1))$. Let $var = [x_{i_1}, \dots, x_{i_s}]$ be a list of variables to be eliminated, then the constant term of an Elliott-rational function $F(x_1, \dots, x_n)$

$$\text{CT}_{x_{i_1}, \dots, x_{i_s}} F(x_1, \dots, x_n)$$

can be evaluated by the command $E.OeqW(F, xvar, var)$ after loading the Ell2 package.

In what follows, we always specify the working field K by letting $0 < x < q < \lambda < 1$. This K includes all the rings, such as $\mathbb{Q}((q))[[x]]$, appear below as a subring. Firstly, we shall explain how to realize the PT_q and $X_{m,r}$ operators by the constant term operator.

For anti-symmetric Laurent polynomials $L(q)$, we have $\text{PT}_q L(q) = -\text{NT}_q L(q) \Big|_{q=q^{-1}}$. For anti-symmetric $F(x, q) \in \mathbb{Q}((q))[[x]]$, $\text{PT}_q F(x, q)$ is not in $\mathbb{Q}[q][[x]]$, but $\text{NT}_q F(x, q)$

belongs to $\mathbb{Q}[q^{-1}][[x]]$. It is convenient for us to use

$$\text{PT}'_q F(x, q) = -\text{NT}_q F(x, q) \Big|_{q=q^{-1}}$$

to replace $\text{PT}_q F(x, q)$, since they agree when $F(x, q) \in \mathbb{Q}[q, q^{-1}][[x]]$. We have

$$(1) \quad \text{PT}'_q F(x, q) = -\text{NT}_q F(x, q) \Big|_{q=q^{-1}} = -\text{CT}_\lambda \frac{\lambda q}{1 - \lambda q} F(x, \lambda),$$

$$(2) \quad X_{m,r} F(x, q) = \text{CT}_\lambda \frac{\lambda^{-r}}{1 - x\lambda^{-m}} F(\lambda, q).$$

The above identities are easily verified for $F(x, q) = q^i x^j$ and then extended by linearity for arbitrary $F(x, q)$.

Let us redo the $m = 3$ case for the sake of clarity. The cases $n = 3k + 1$ and $n = 3k + 2$ can be done simultaneously. By starting with the formula

$$F_3(x, q) = \frac{q^2 (1 - q^2) (1 - q^4)}{(1 - x) (x - q^2) (1 - q^2 x) (1 - q^6)},$$

we compute

$$G_3(x, q) = \text{PT}'_q F_3(x, q) = \frac{q^2 x}{(1 - x^3) (1 - q^2 x)}.$$

This clearly belongs to $\mathbb{N}[q][[x]]$, and hence reprove Conjecture 3^b for $(m, r) = (3, 1)$ and $(m, r) = (3, 2)$. Indeed, a further step gives

$$X_{3,1} G_3(x, q) = \frac{q^2}{(1 - x) (1 - q^6 x)} \in \mathbb{N}[q][[x]],$$

$$X_{3,2} G_3(x, q) = \frac{q^4}{(1 - x) (1 - q^6 x)} \in \mathbb{N}[q][[x]].$$

The case $n = 3k$ is a little different. We need to compute

$$H_3^0(x, q) = \text{PT}'_q \frac{q^3 - q^{-3}}{q - q^{-1}} F_3(x, q) = \frac{q^2}{(1 - x) (1 - q^2 x)},$$

which clearly belongs to $\mathbb{N}[q][[x]]$. This implies the positivity of $X_{3,0} H_3^0(x, q)$ and hence reproves Conjecture 4^b for $(m, r) = (3, 0)$. Indeed, we have

$$X_{3,0} H_3^0(x, q) = \frac{q^2 (1 + q^2 x + q^4 x)}{(1 - x) (1 - q^6 x)}.$$

4.3. The case $m = 4$. We shall establish the following result.

Proposition 8. *Conjecture 4^b holds true for $m = 4$.*

Proof. We start with the formula

$$F_4(x, q) = -\frac{q^5 (1 - q^2) (1 - q^4)}{(x - q) (1 - qx) (x - q^3) (1 - q^3 x) (1 - q^8)}.$$

By constant term extraction, we obtain

$$G_4(x, q) = \text{PT}'_q F_4(x, q) = -\frac{xq(-q + x^3 - qx^2 + q^2x - qx^4)}{(1 - qx) (1 - xq^3) (1 - x^8)},$$

which do not show the positivity directly. By applying $X_{4,r}$ for $r = 1, 3$, we obtain

$$\begin{aligned} X_{4,1}G_4(x, q) &= \frac{q^2(1 + q^6x + q^6x^2 + q^{12}x^2)}{(1 - x^2)(1 - q^4x)(1 - q^{12}x)}, \\ X_{4,3}G_4(x, q) &= \frac{q^2(1 + q^6 + q^6x + q^{12}x^2)}{(1 - x^2)(1 - q^4x)(1 - q^{12}x)}. \end{aligned}$$

This proves Conjecture 3^b for $(m, r) = (4, 1), (4, 3)$.

For the case $(m, r) = (4, 2)$, we need to compute

$$\begin{aligned} H_4^2(x, q) &= \text{PT}'_q \frac{q^2 - q^{-2}}{q - q^{-1}} F_4(x, q) = \frac{xq(1 - x^4)(1 - qx + q^2 - q^3x + q^3x^3)}{(1 - x^2)(1 - qx)(1 - q^3x)(1 - x^8)}, \\ X_{4,2}H_4^2(x, q) &= \frac{q^4(1 + q^2)(1 + xq^6)}{(1 - x)(1 - q^4x)(1 - q^{12}x)}, \end{aligned}$$

which is clearly in $\mathbb{N}[q][[x]]$.

For the case $(m, r) = (4, 0)$, we need to compute

$$\begin{aligned} H_4^0(x, q) &= \text{PT}'_q \frac{q^4 - q^{-4}}{q - q^{-1}} F_4(x, q) = \frac{q^2}{(1 - x^2)(1 - qx)(1 - q^3x)}, \\ X_{4,0}H_4^0(x, q) &= \frac{q^2(1 + q^2x + q^4x + 2xq^6 + q^8x + q^{10}x + q^{12}x^2)}{(1 - x)(1 - q^4x)(1 - q^{12}x)}, \end{aligned}$$

which is clearly in $\mathbb{N}[q][[x]]$. ■

4.4. The case $m = 5$.

Proposition 9. *Conjecture 4^b holds true for $m = 5$.*

Proof. We start with the formula

$$F_5(x, q) = -\frac{q^8(1 - q^2)(1 - q^4)}{(1 - x)(x - q^2)(1 - q^2x)(x - q^4)(1 - q^4x)(1 - q^{10})}.$$

By the constant term method, we obtain

$$G_5(x, q) = \text{PT}'_q F_5(x, q) = \frac{xq^2(1 - x)(1 + x - x^3 - q^2x + q^2x^3 + q^2x^4)}{(1 - x^2)(1 - x^3)(1 - q^2x)(1 - x^5)(1 - q^4x)},$$

which do not show the positivity directly. By applying $X_{4,r}$ for $r = 1, 2, 3, 4$, we obtain

$$X_{5,r}G_5(x, q) = \frac{q^2P_{5,r}(x, q)}{(1 - x^2)(1 - x^3)(1 - q^{10}x)(1 - q^{20}x)},$$

where

$$\begin{aligned} P_{5,1}(x, q) &= 1 + q^2x + q^2x^2 + q^4x + q^4x^2 + q^4x^3 + q^6x^2 + 2q^6x^3 + q^8x + q^6x^4 \\ &\quad + 2q^8x^2 + q^8x^3 + q^8x^4 + 2q^{10}x^2 + 2q^{10}x^3 + q^{12}x + q^{10}x^4 + q^{12}x^2 \\ &\quad + 2q^{12}x^3 + q^{14}x + q^{12}x^4 + q^{14}x^2 + q^{14}x^3 + q^{14}x^4 + 2q^{16}x^2 + q^{14}x^5 \\ &\quad + 2q^{16}x^3 + q^{16}x^4 + q^{18}x^2 + 2q^{18}x^3 + 2q^{18}x^4 + q^{20}x^2 + q^{20}x^3 + q^{20}x^4 \\ &\quad + q^{20}x^5 + q^{22}x^3 + q^{22}x^4 + q^{22}x^5 + q^{24}x^3 + q^{24}x^4 + q^{26}x^2, \end{aligned}$$

$$\begin{aligned}
P_{5,2}(x, q) = & x + q^2x^2 + q^4 + q^2x^3 + q^4x + q^4x^2 + q^6x + q^6x^2 + q^6x^3 + q^8x \\
& + q^6x^4 + 2q^8x^2 + 2q^8x^3 + q^{10}x + q^{10}x^2 + 2q^{10}x^3 + q^{12}x + q^{10}x^4 \\
& + 2q^{12}x^2 + q^{12}x^3 + q^{12}x^4 + 2q^{14}x^2 + 2q^{14}x^3 + q^{16}x + q^{14}x^4 + q^{16}x^2 \\
& + 2q^{16}x^3 + q^{18}x + q^{16}x^4 + q^{18}x^2 + q^{18}x^3 + q^{18}x^4 + q^{20}x^2 + q^{18}x^5 + 2q^{20}x^3 \\
& + q^{20}x^4 + q^{22}x^2 + q^{22}x^3 + q^{22}x^4 + q^{24}x^4 + q^{24}x^5 + q^{26}x^3,
\end{aligned}$$

$$\begin{aligned}
P_{5,3}(x, q) = & x^2 + q^2 + q^2x + q^4x + q^4x^2 + q^4x^3 + q^6x + 2q^6x^2 + q^8 + q^6x^3 + q^8x \\
& + q^8x^2 + q^8x^3 + q^{10}x + q^8x^4 + 2q^{10}x^2 + q^{10}x^3 + q^{12}x + q^{10}x^4 + 2q^{12}x^2 \\
& + 2q^{12}x^3 + q^{14}x + q^{14}x^2 + 2q^{14}x^3 + q^{16}x + q^{14}x^4 + 2q^{16}x^2 + q^{16}x^3 + q^{16}x^4 \\
& + 2q^{18}x^2 + 2q^{18}x^3 + q^{20}x + q^{18}x^4 + q^{20}x^2 + q^{20}x^3 + q^{20}x^4 + q^{22}x^3 + q^{22}x^4 \\
& + q^{24}x^2 + q^{22}x^5 + q^{24}x^3 + q^{26}x^4,
\end{aligned}$$

$$\begin{aligned}
P_{5,4}(x, q) = & x^3 + q^2x + q^2x^2 + q^4 + q^4x + q^4x^2 + q^6 + q^6x + q^6x^2 + q^6x^3 + 2q^8x + 2q^8x^2 \\
& + q^8x^3 + q^{10}x + 2q^{10}x^2 + q^{12} + 2q^{10}x^3 + q^{12}x + q^{12}x^2 + q^{12}x^3 + q^{14}x + q^{12}x^4 \\
& + 2q^{14}x^2 + q^{14}x^3 + q^{16}x + q^{14}x^4 + 2q^{16}x^2 + 2q^{16}x^3 + q^{18}x + q^{18}x^2 + 2q^{18}x^3 \\
& + q^{20}x + q^{18}x^4 + 2q^{20}x^2 + q^{20}x^3 + q^{22}x^2 + q^{22}x^3 + q^{22}x^4 + q^{24}x^3 + q^{24}x^4 + q^{26}x.
\end{aligned}$$

This proves Conjecture 3^b for $m = 5$ and $r = 1, 2, 3, 4$.

For the case $n = 5k$, we have

$$H_5^0(x, q) = \text{PT}'_q \frac{q^5 - q^{-5}}{q - q^{-1}} F_5(x, q) = \frac{q^2(1 + q^2x^2)}{(1 - x^2)(1 - x^3)(1 - q^2x)(1 - q^4x)},$$

which clearly implies the positivity for $X_{5,0}H_5^0(x, q)$. Indeed, we have

$$X_{5,0}H_5^0(x, q) = \frac{q^2P_{5,0}(x, q)}{(1 - x^2)(1 - x^3)(1 - q^{10}x)(1 - q^{20}x)},$$

where $P_{5,0}(x, q) \in \mathbb{N}[q, x]$ contains 64 terms. This reproves Conjecture 4^b for $(m, r) = (5, 0)$. $\blacksquare$

4.5. The cases $m \geq 6$. When we calculated the case $m \geq 6$ in a similar way, we are not able to prove the positivity in a straightforward way as before. Let us explain the problem by working with the $m = 6$ case. We start with the formula

$$F_6(x, q) = \frac{q^{12}(1 - q^2)(1 - q^4)}{(x - q)(1 - qx)(x - q^3)(1 - q^3x)(x - q^5)(1 - q^5x)(1 - q^{12}x)}.$$

By the constant term method, we obtain

$$G_6(x, q) = \text{PT}'_q F_6(x, q) = -\frac{xqM_6}{(1 - qx)(1 - q^3x)(1 - x^6)(1 - q^5x)(1 - x^8)(1 - x^{12})},$$

where M_6 is a polynomial of many terms that does not show positivity. We can apply $X_{6,r}$ for $r = 1, 5$, corresponding to the $\gcd(m, r) = 1$ case. Neither of the two cases shows the positivity directly.

For the $\gcd(m, r) = 2$ case, i.e., $r = 2, 4$, we need to compute

$$H_6^2(x, q) = \text{PT}'_q \frac{q^2 - q^{-2}}{q - q^{-1}} F_6(x, q) = \frac{\text{A lengthy polynomial}}{(1 - x)^2(1 - x^2)(1 - x^4)(1 - q^6x)(1 - q^{18}x)(1 - q^{30}x)},$$

We can apply $X_{6,r}$ for $r = 2, 4$. Neither of the two cases shows the positivity directly.

For the $\gcd(m, r) = 3$ case, i.e., $r = 3$, we need to compute

$$H_6^3(x, q) = \text{PT}'_q \frac{q^3 - q^{-3}}{q - q^{-1}} F_6(x, q) = \frac{\text{A lengthy polynomial}}{(1-x)(1-x^2)(1-x^4)(1-q^6x)(1-q^{18}x)(1-q^{30}x)},$$

We can apply $X_{6,3}$, and the result does not show the positivity directly.

For the case $r = 0$, we need to compute

$$H_6^0(x, q) = \text{PT}'_q \frac{q^6 - q^{-6}}{q - q^{-1}} F_6(x, q) = \frac{\text{A lengthy polynomial}}{(1-x)^2(1-x^4)(1-q^6x)(1-q^{18}x)(1-q^{30}x)},$$

We can apply $X_{6,0}$, and the result does not show the positivity directly.

4.6. An extension. The computation of the case $m = 6$ suggests that we need to consider different cases for proving Conjecture 4^b. However, we find a possible unified way to attack the conjecture. More precisely, let

$$(3) \quad G_m(x, q) = \text{PT}'_q F_m(x, q) \text{ for } m \geq 3,$$

or equivalently,

$$(4) \quad [q^i]G_m(x, q) = \text{CT}_q - q^i F_m(x, q) = \text{CT}_q - q^i (q^2 - q^{-2}) \frac{(q - q^{-1})}{(q^m - q^{-m})} \prod_{i=0}^{m-1} \frac{1}{(1 - xq^{1-m} \cdot q^{2i})}.$$

Then $G_m(x, q) \in \mathbb{Q}[q][[x]]$, and it is easy to verify that:

$$[q^0]G_m(x, q) = 0,$$

$$[qx^n]G_m(x, q) = [q]\mathcal{N}(q^2 - 1)C_{m,n}(q) \neq 0 \text{ only when } m, n \text{ are both even,}$$

$$[q]G_4(x, q) = -\frac{1}{1 - x^4},$$

$$\begin{aligned} [q]G_6(x, q) &= -\frac{x^6(1 - x^2 + x^4 - x^6 - x^8)}{(1 - x^6)(1 - x^8)(1 - x^{12})} \\ &= -(x^6 + x^{10} + 2x^{18} + x^{22} + 2x^{30} + x^{34} + 2x^{42} + x^{54}) + \text{positive terms,} \end{aligned}$$

$$[q]G_{10}(x, q) = -(x^6q + x^{10}q) + \text{positive terms.}$$

We summarized all the other cases in the following conjecture.

Conjecture 10. *Let $G_m(x, q)$ be as in (3). Then $G_m(x, q)$ is almost positive for $m \geq 3$. More precisely, besides the above formula, we have*

- (1) *For every $k \geq 3$, $[q]G_{2k}(x, q)$ has only finitely many negative terms.*
- (2) *For every $i \geq 2$, $[q^i]G_m(x, q) \in \mathbb{N}[[x]]$.*

A unified approach for $m = 4$ and $r = 1, 3$ can be given as follows. We have

$$\begin{aligned}
G_4(x, q) &= -\frac{xq(-q + x^3 - x^2q + q^2x - x^4q)}{(1 - qx)(1 - q^3x)(1 - x^8)} \\
&= \frac{q^2x(1 + x^2 - qx - q^2x^4 + q^3x^5)}{(1 - qx)(1 - q^3x)(1 - x^8)} - \frac{x^4q}{1 - x^8} \\
&= \frac{q^2x(1 - qx + x^2(1 - q^2x^2))}{(1 - qx)(1 - q^3x)(1 - x^8)} + \frac{q^2x(q^3x^5)}{(1 - qx)(1 - q^3x)(1 - x^8)} - \frac{x^4q}{1 - x^8} \\
&= \frac{q^2x(1 + x^2 + qx^3)}{(1 - q^3x)(1 - x^8)} + \frac{q^5x^6}{(1 - qx)(1 - q^3x)(1 - x^8)} - \frac{x^4q}{1 - x^8}
\end{aligned}$$

It follows that

$$X_{4,r}G_4(x, q) = X_{4,r}\frac{q^2x(1 + x^2 + qx^3)}{(1 - q^3x)(1 - x^8)} + X_{4,r}\frac{q^5x^6}{(1 - qx)(1 - q^3x)(1 - x^8)},$$

which is clearly positive.

Generally, for odd m , we need to show the positivity of $G_m(x, q)$. It is possible to decompose $G_m(x, q)$ as a sum of trivially positive rational functions. The decomposition is nontrivial even for $G_5(x, q)$. The even m case needs a minor modification. We succeeded in doing this type of decomposition in [20], and hence decomposition of $G_m(x, q)$ for small m , at least for $m \leq 6$, should be possible. This suggested us to reconsider the following problem in the near future.

Problem: Given an Elliott rational function Q , how to decompose $Q = \sum_i Q_i$ with Q_i all trivially positive if possible.

5. COMBINATORIAL MODEL

5.1. Combinatorial interpretation of $C_{m,n}(q)$. In this section (m, n) is always a co-prime pair of positive integers, unless specified otherwise. The general case $\gcd(m, n) = d > 1$ can be considered but is much more complicated.

We should mention that representation theory maybe suitable for settling our conjectures. For instance, Conjecture 3 can be proved if we can find (usually hard to find) an $sl(2)$ module whose character is $q^{-(m-1)(n-1)/2}C_{m,n}(q)$. This is based on the following well-known result. See, e.g., [22, Theorem 15].

Theorem 11. *Let $\psi : sl(2) \mapsto gl(n)$ be a representation of $sl(2)$ with*

$$\text{char } \psi = \sum_i b_i q^i.$$

Then the sequence $\dots, b_{-2}, b_{-1}, b_0, b_1, b_2, \dots$ is symmetric and unimodal with respect to parity.

Let $\mathcal{D}_{m,n}$ be the set of Dyck paths in the $m \times n$ lattice rectangle, i.e., paths from $(0, 0)$ to (m, n) with unit North step and unit East step, that never go below the diagonal line $y = nx/m$. The rational q, t Catalan polynomials are defined by

$$C_{m,n}(q, t) = \sum_{D \in \mathcal{D}_{m,n}} q^{\text{area}(D)} t^{\text{dinv}(D)},$$

where the sum is over Dyck paths in the $m \times n$ lattice rectangle, $\text{area}(D)$ gives the number of lattice squares between the path and the diagonal, and $\text{dinv}(D)$ is a Dyck path statistic that can also be given a relatively simple geometric construction. There is also an equivalent interpretation in terms of simultaneous core partitions. See [5, 24].

The rational q -Catalan polynomials are specializations of the q, t Catalan polynomials introduced by Garsia and Haiman [10]. They have the following combinatorial interpretation:

$$C_{m,n}(q) = q^{(m-1)(n-1)/2} C_{m,n}(q, q^{-1}) = \sum_{D \in \mathcal{D}_{m,n}} q^{\text{coarea}(D) + \text{dinv}(D)}.$$

It seems hard to show the parity unimodality of $C_{m,n}(q)$ by this model, because the dinv statistic is still hard to understand.

A mysterious property of $C_{m,n}(q, t)$ is its symmetry in q and t , i.e., $C_{m,n}(q, t) = C_{m,n}(t, q)$. As a symmetric polynomial, $C_{m,n}(q, t)$ has a Schur expansion

$$C_{m,n}(q, t) = \sum_{\lambda} c_{\lambda} s_{\lambda}[q, t],$$

where λ can have only two parts, say $\lambda = (\lambda_1, \lambda_2)$, and

$$s_{\lambda_1 \lambda_2}[q, t] = (qt)^{\lambda_2} [\lambda_1 - \lambda_2]_{q,t}, \quad \text{where } [k]_{q,t} = q^{k-1} + q^{k-2}t + \cdots + t^{k-1}.$$

Then the (q, t) Schur positivity of $C_{m,n}(q, t)$ (i.e., $c_{\lambda} \geq 0$ for all λ) implies the unimodality of $C_{m,n}(q)$ with respect to parity.

The symmetry of $C_{m,n}(q, t)$ is a consequence of the rational shuffle conjecture, which can be written as

$$Q_{m,n}(-1)^n = H_{m,n}(X; q, t),$$

where $H_{m,n}(X; q, t)$ is the Hikita polynomial that has combinatorial interpretation as a sum over rational parking functions [14], and $Q_{m,n}$ is a symmetric function operator introduced by Gosky and Negut [11]. The rational Shuffle conjecture was proved by Mellit [18]. Detailed definitions are too involved. The reader is referred to [21] for further information on this topic. We should mention that no combinatorial proof of this symmetry is known up to now.

As a symmetric function in X , we can write

$$H_{m,n}(X; q, t) = \sum_{\lambda \vdash n} [s_{\lambda}]_{m,n} s_{\lambda}[X].$$

Then $C_{m,n}(q, t)$ is just $[s_{1^n}]_{m,n}$. From the algebraic side, $H_{m,n}(X; q, t)$ is easily seen to be q, t symmetric, so is its coefficients $[s_{\lambda}]_{m,n}$. It is then natural to conjecture that $[s_{\lambda}]_{m,n}(q, t)$ is (q, t) Schur positive. The positivity (though not stated this way) has been proved for the case $n = 2$ by Leven [15] and for the case $n = 3$ by Qiu and Remmel [21].

5.2. Combinatorial interpretation of $C_n(q)$. Since $C_n(q) = C_{n+1,n}(q)$, we have a combinatorial interpretation of $C_n(q)$. Indeed, let $\mathcal{D}_n$ be short for $\mathcal{D}_{n,n}$. Then we have

$$C_n(q) = \frac{1}{[n+1]_q} \begin{bmatrix} 2n \\ n \end{bmatrix}_q = \sum_{D \in \mathcal{D}_n} q^{\text{coarea}(D) + \text{dinv}(D)} = \sum_{D \in \mathcal{D}_n} q^{\text{coarea}(D) + \text{bounce}(D)}.$$

The second equality follows by the symmetry of $C_n(q)$ and by application of the zeta map, which is a bijection from $\mathcal{D}_n$ to itself that takes dinv to area and area to bounce .

See [24, 16]. Usually we think the statistic bounce is simpler than dinv in this case, (while for $D \in \mathcal{D}_{m,n}$, the $\text{dinv}(D)$ is known but the $\text{bounce}(D)$ is not).

Currently the simplest way to compute $\text{coarea}(D)$ and $\text{bounce}(D)$ might be as follows (see [28]). Firstly, there is a easy way to convert $D \in \mathcal{D}_n$ to a standard Young tableau $T(D)$ of shape (n, n) . Then $\text{coarea}(D)$ is just the sum of the first row entries minus $\binom{n}{2}$, and $\text{bounce}(D)$ is the sum of the first row ranks, where the ranks of the entries of T can be computed in a simple way: i) $r(1) = 0$; ii) $r(i) = r(i-1)$ if i is in the first row; iii) $r(j) = r(i) + 1$ if j is under i . For example, Figure 1 illustrates these statistics for the case $n = 3$.

	1 2 3	1 2 4	1 2 5	1 3 5	1 3 4
	4 5 6	3 5 6	3 4 6	2 4 6	2 5 6
	6 ₌₁₊₂₊₃	7 ₌₁₊₂₊₄	8 ₌₁₊₂₊₅	9 ₌₁₊₃₊₅	8 ₌₁₊₄₊₄
coarea	0	1	2	3	2
	0 0 0	0 0 1	0 0 1	0 1 2	0 1 1
	1 1 1	1 1 2	1 1 2	1 2 3	1 2 2
bounce	0	1	1	3	2
$q^{\text{coarea}+\text{bounce}}$	q^0	q^2	q^3	q^6	q^4

$$C_3(q) = 1 + q^2 + q^3 + q^4 + q^6$$

Figure 1. Bounce and coarea for $\mathcal{D}_3$: The top row gives the 5 standard Young tableaux. The bottom row gives the corresponding rank tableaux.

There is a better interpretation found earlier than the above statistics. See, e.g., [23].

$$C_n(q) = \frac{1}{[n+1]} \left[\begin{matrix} 2n \\ n \end{matrix} \right]_q = \sum_{D \in \mathcal{D}_n} q^{\text{maj}(D)}$$

where $\text{maj}(D)$ is the major index of D , usually defined as the sum of the descent positions. (Here a descent corresponds to a EN turn). The major index is also defined for standard Young tableaux. We only exhibit the major index for $C_3(q)$.

1 2 3	1 2 4	1 2 5	1 3 5	1 3 4
4 5 6	3 5 6	3 4 6	2 4 6	2 5 6
q^3	q^6	q^2	q^0	q^4

$$C_3(q) = 1 + q^2 + q^3 + q^4 + q^6$$

Figure 2. Major index by standard Young tableaux of shape (n, n) . Entry i is a descent if $i+1$ appears to the left of i .

There are also two closely related results. One is the following [22, p. 523].

Theorem 12. *The polynomial $K_n(q) = \frac{1+q}{1+q^n} C_n(q)$ is symmetric and unimodal.*




Figure 3. Major index by Dyck paths in $\mathcal{D}_n$. Descents appear at the EN turns.

For instance, $K_0(q) = K_1(q) = 1$, $K_2(q) = 1 + q$, $K_3(q) = 1 + q + q^2 + q^3 + q^4$, $K_4(q) = 1 + q + q^2 + 2q^3 + 2q^4 + 2q^5 + 2q^6 + q^7 + q^8 + q^9$. This is also a q -analogue of the Catalan number $C_n = \frac{1}{n+1} \binom{2n}{n}$. The degree of $K_n(q)$ is $(n-1)^2$.

It is not hard to see that the unimodality of $K_n(q)$ implies that $(1+q)C_n(q) = (q^n + 1)K_n(q)$ is almost unimodal. Indeed, if we let $K_n(q) = \sum_i k_i q^i$ and $(1+q)C_n(q) = \sum_i c_i q^i$, then $c_i = k_i + k_{i+n}$. Consider

$$c_{i+1} - c_i = (k_{i+1} + k_{i+1+n}) - (k_i + k_{i+n}) = (k_{i+1} - k_i) + (k_{i+1+n} - k_{i+n}).$$

Thus the unimodality of $K_n(q)$ (Theorem 12) implies that $c_{i+1} - c_i \geq 0$ for $0 < i < \frac{n^2-4n-1}{2}$, while the desired positivity $c_{i+1} - c_i \geq 0$ is for $1 \leq i < \frac{n^2-3n-1}{2}$.

The other one is the following conjecture. See [8].

Conjecture 13. Write $C_n(q) = \sum_k m_n(k)q^k$. The sequence $(m_n(1), m_n(2), \dots, m_n(n(n-1) - 1))$ is unimodal when n is sufficiently large. (Seem to hold for $n \geq 16$.)

If this conjecture is true for $n \geq 16$, then Conjecture 3 is also true because the $n \leq 16$ cases are easily verified to be true.

Acknowledgements:

The authors would like to thank Bill Chen and Arthur Yang for helpful discussions.

REFERENCES

- [1] Andrews, George E, The Friedman-Joichi-Stanton monotonicity conjecture at primes, Unusual Applications of Number Theory (M. Nathanson, ed.), DIMACS Ser. Discrete Math. Theor. Comp. Sci., vol. 64, Amer. Math. Soc., Providence, R.I., 2004, pp. 9–15.
- [2] Andrews, George E. and Paule, Peter and Riese, Axel, MacMahons partition analysis. VI. A new reduction algorithm, Ann. Comb. 2001, 5, 251–270, Dedicated to the memory of Gian-Carlo Rota (Tianjin, 1999).
- [3] Andrews, George E. and Peter Paule and Axel Riese, MacMahons partition analysis: the Omega package, European J. Combin. 2001, 22(7), 887–904.
- [4] Andrews, George E., On the difference of successive Gaussian polynomials, Journal of Statistical Planning and Inference, 1993, 34(1), 19–22.
- [5] Armstrong, Drew and Hanusa, Christopher R. H. and Jones, Brant C., Results and Conjectures on Simultaneous Core Partitions, European J. Combin., 2014, 41, 205–220.
- [6] Bressou, David M., Unimodality of Gaussian polynomials. Discrete Mathematics, 1992, 99(1-3), 17–24.
- [7] Butler, Lynne M., A unimodality result in the enumeration of subgroups of a finite abelian group, Proceedings of the American Mathematical Society, 1987, 101(4), 771–775.
- [8] Chen, William Y. C and Wang, Carol J and Wang, Larry X. W, The limiting distribution of the coefficients of the q -Catalan numbers, Proceedings of the American Mathematical Society, 2008, 136(11), 3759–3767.
- [9] Garsia, Adriano and Leven, Emily Sergel and Wallach, Nolan and Xin, Guoce, A new plethystic symmetric function operator and the rational compositional shuffle conjecture at $t = 1/q$, Journal of Combinatorial Theory, Series A, 2017, 145, 57–100.

- [10] Garsia, Adriano M. and Haiman, Mark D., A remarkable q, t -Catalan sequence and q -Lagrange inversion, *J. Algebraic Combinatorics*, 1996, 5, 191–244.
- [11] Gorsky, Eugene and Negut, Andrei, Refined knot invariants and Hilbert schemes, *Journal de Mathématiques Pures et Appliquées*, 2015, 104(3), 403–435.
- [12] Guo, Victor J. W. and Krattenthaler, Christian, Some divisibility properties of binomial and q -binomial coefficients, *J. Number Theory*, 2014, 135, 167–184.
- [13] Haiman, Mark D., Conjectures on the quotient ring by diagonal invariants, *J. Algebraic Combin.*, 1994, 3, 17–76.
- [14] Hikita, Tatsuyuki, Affine Springer fibers of type A and combinatorics of diagonal coinvariants, *Advances in Mathematics*, 2014, 263, 88–122.
- [15] Leven, Emily, Two special cases of the Rational Shuffle Conjecture, *Discrete Mathematics & Theoretical Computer Science Proceedings*, 2014, 01, 789–800.
- [16] Loehr, Nicholas A, Conjectured statistics for the higher q, t -Catalan sequences, *Electron. J. Combin.*, 2005, 12 research paper R9; 54 pages (electronic).
- [17] Macdonald, G., An elementary proof of a q -Binomial identity, in D. Stanton ed., *q-Series and Partitions*, IMA Vol. Math. Appl. 18 (Springer, New York 1989), 73–75.
- [18] Mellit, Anton, Toric braids and (m, n) -parking functions, preprint, 2016, arXiv:1604.07456.
- [19] O’Hara, Kathleen M, Unimodality of gaussian coefficients: a constructive proof. *Journal of Combinatorial Theory*, 1990, 53(1), 29–52.
- [20] Qiu, Dun and Remmel, Jeffrey B. and Sergel, Emily and Xin, Guoce, On the Schur positivity of $\Delta_{e_2} e_n[X]$, *Electron. J. Combin.*, 2018, 25(4), P4.20.
- [21] Qiu, Dun and Remmel, Jeffrey, Schur function expansions and the rational shuffle conjecture, *Proceedings of the 29th Conference on Formal Power Series and Algebraic Combinatorics*, London (2017).
- [22] Stanley, Richard P, Log-concave and unimodal sequences in algebra, combinatorics, and geometry. *Annals of the New York Academy of Sciences*, 2010, 576(1), 500–535.
- [23] Stanley, Richard P, *Enumerative Combinatorics*, Vol. 2, Cambridge University Press, Cambridge, 1999.
- [24] Xin, Guoce, A note on rank complement of rational Dyck paths and conjugation of (m, n) -core partitions, *J. of Combin.*, 2017, 8(no.4), 704–726.
- [25] Xin, Guoce, A Euclid style algorithm for MacMahon’s partition analysis, *J. Combin. Theory A.*, 2015, 131, 32–60.
- [26] Xin, Guoce, The Ring of Malcev-Neumann Series and The Residue Theorem, Ph.D. thesis, Brandeis University, 2004, arXiv:math.CO/0405133.
- [27] Xin, Guoce, A fast algorithm for MacMahon’s partition analysis, *Electron. J. Combin.*, 2004, 11, R58. (20 pages).
- [28] Xin, Guoce and Zhang, Yingrui, On the Sweep Map for $\vec{k}$ -Dyck Paths, arXiv:1811.07475.
- [29] Zeilberger, Doron. Kathy O’Hara’s constructive proof of the unimodality of the gaussian polynomials. *American Mathematical Monthly*, 1989, 96(7), 590–602.
- [30] Zeilberger, Doron. A one-line high school algebra proof of the unimodality of the gaussian polynomials $\begin{bmatrix} n \\ k \end{bmatrix}_q$ for $k < 20$, in Dennis Stanton (ed.) *q-Series and partitions*, IMA Volumes in Mathematics and its Applications, Volume # 18, Springer-Verlag, New York(1989). pp. 67–72.

¹SCHOOL OF MATHEMATICAL SCIENCES, CAPITAL NORMAL UNIVERSITY, BEIJING 100048, PR CHINA, ²SCHOOL OF MATHEMATICAL SCIENCES, CAPITAL NORMAL UNIVERSITY, BEIJING 100048, PR CHINA

E-mail address: *guoce.xin@gmail.com &zhongyueming107@gmail.com