

Entropy-Based Formulation of Thermodynamics in Arbitrary Quantum Evolution

S. Alipour,¹ A. T. Rezakhani,² A. Chenu,^{3,4,5} A. del Campo,^{3,4,5,6} and T. Ala-Nissila^{1,7}

¹*QTF Center of Excellence, Department of Applied Physics,
Aalto University, P. O. Box 11000, FI-00076 Aalto, Espoo, Finland*

²*Department of Physics, Sharif University of Technology, Tehran 14588, Iran*

³*Department of Physics and Materials Science, University of Luxembourg, L-1511 Luxembourg, G.D. Luxembourg*

⁴*Donostia International Physics Center, E-20018 San Sebastián, Spain*

⁵*IKERBASQUE, Basque Foundation for Science, E-48013 Bilbao, Spain*

⁶*Department of Physics, University of Massachusetts, Boston, MA 02125, USA*

⁷*Interdisciplinary Centre for Mathematical Modelling and Department of Mathematical Sciences,
Loughborough University, Loughborough, Leicestershire LE11 3TU, UK*

Given the evolution of an arbitrary open quantum system, we formulate a general and unambiguous method to separate the internal energy change of the system into an entropy-related contribution and a part causing no entropy change, identified as heat and work, respectively. We also demonstrate that heat and work admit geometric and dynamical descriptions by developing a universal dynamical equation for the given trajectory of the system. The dissipative and coherent parts of this equation contribute exclusively to heat and work, where the specific role of a work contribution from a counterdiabatic drive is underlined. Next we define an expression for the irreversible entropy production of the system which does not have explicit dependence on the properties of the ambient environment; rather, it depends on a set of the system's observables excluding its Hamiltonian and is independent of internal energy change. We illustrate our results with three examples.

Quantum thermodynamics holds a central stage at the interface of quantum information science, statistical mechanics, and quantum technologies, and it has shed new light on the laws of thermodynamics in the quantum regime [1–15]. Apart from uncovering a plethora of new phenomena, quantum thermodynamics has also motivated efforts to engineer efficient quantum machines in the laboratory [16–20].

Yet, several fundamental issues remain to be clarified. Most notably, except for particular regimes (e.g., weak-coupling, Markovian dynamics with slow driving [21]), an unequivocal definition of heat and work in arbitrary open-quantum system dynamics has been elusive. One problem is that such thermodynamic variables are not observables described by Hermitian operators [22]; rather, they are trajectory-dependent quantities [23–27]. Existing definitions mostly incorporate generators of the dynamics (besides the state) and dynamical master equations with *coherent* and *dissipative* parts [10, 21, 28–32]. For instance, in the widely-used *conventional* framework [21, 33] the internal energy change due to the dissipative (coherent) part of the master equation has been called heat (work) change. An alternative strategy uses the potential of mean force that amounts to pre-averaging the total partition function over the environmental degrees of freedom [28–32]. In addition, there are *semiclassical* approaches where a coarse-grained version of the system state in the energy eigenbasis and semiclassical definitions of heat and work have been employed [34–36].

Despite extensive efforts, some of the existing approaches neither are consistent with each other nor do they always reproduce expected results according to thermodynamics. Although there exist attempts to overcome such inconsistencies through ad hoc methods [32, 37, 38], the issue remains unresolved.

To further clarify the issue, we note the following drawbacks of the conventional framework. (i) Its formulation has been originally devised only for particular conditions, e.g., a Markovian dynamics with a constant or relatively slow-varying Hamiltonian, when the dissipative part does not explicitly de-

pend on the physical system Hamiltonian. Despite this subtlety, the conventional definitions of heat and work have been applied almost arbitrarily in the literature to more general scenarios. In fact, dissipative part of a master equation may explicitly depend on the physical Hamiltonian of the system [39, 40]. Hence in such physical scenarios, external driving may also contribute to the heat exchanged. This aspect introduces further ambiguities and hinders a clear and thermodynamically consistent assignment of the heat and work concepts. (ii) Since the description of the dynamics in the form of the conventional Lindblad master equation is not unique [41], different energy values can be assigned to differently chosen generators of the same dynamics. This indicates that part of the energy assigned to the dissipative part of the master equation may not necessarily lead to an entropy change, and should be identified as work. Specifically, for initial states of the decoherence-free subspace type [42, 43], the dynamics leads to no dissipation along the corresponding trajectories and all the energy change, if any, would be of the work type. Recently, a scheme to partially remedy the shortcomings of the conventional definitions has been proposed in Ref. [44], where heat has been defined by subtracting *ergotropy* (maximum extractable work from a system in a cyclic unitary process) [45] from the dissipative part of the internal energy change by considering a *virtual* instantaneous unitary transformation. Nevertheless, an extension of ergotropy to open systems is still elusive, and since this extra transformation does not rely on the real physical dynamics, trajectory-dependent quantities such as heat can be fictitious. For an alternative approach, separating the internal energy based on conserved quantities of the dynamics, see Ref. [46].

In this Letter, on pure thermodynamic grounds supplemented with dynamical and geometric arguments, we circumvent the issue and ambiguities discussed above and put forward a set of universal definitions for heat, work, and entropy production based explicitly on trajectories in the state space of a quantum system (note the difference with quantum jump *tra*-

jectories arising in stochastic dynamics of monitored systems [47, 48]). Our framework is general and independent of the dynamics which generates the trajectory, and it is thus applicable to any time-continuous evolution.

In particular, following standard thermodynamics wherein energy change in an isentropic process is identified as work [49, 50], we define heat solely as the part of the internal energy of the system which can be associated with entropy change. While the first law of thermodynamics treats heat and work on an equal footing, the second law distinguishes them. Essentially heat is a form of disorganized energy, and some disorganization (entropy) will flow with it [51]. Hence, heat is introduced as the part of the energy change which can be accompanied by an entropy change, whereas the energy transfer in the form of work definitely does not lead to any entropy change. We use this key feature to identify heat and work and accordingly obtain a computable expression for the irreversible entropy production. We show that this quantity varies due to the change in variables of the system other than its energy. This clarifies why in general scenarios heat and entropy changes are not necessarily monotonic with respect to each other.

Consider the evolution of the system in a time interval $t \in [0, t_f]$ described by a time-dependent density matrix $\varrho(t)$, expressed in its spectral decomposition as

$$\varrho(t) = \sum_{k=1}^D r_k(t) |r_k(t)\rangle \langle r_k(t)|. \quad (1)$$

One can consider $\{\varrho(t)\}$ as a *trajectory* in the state space of the system, starting from given $\{\varrho(0)\}$ (for brevity, we drop all time dependence henceforth unless necessary). Note that the change in ϱ can be decomposed as $d\varrho = \bar{d}\varrho^{(\text{ev})} + \bar{d}\varrho^{(\text{ep})}$, where $\bar{d}\varrho^{(\text{ev})} = \sum_k dr_k |r_k\rangle \langle r_k|$ is the change due to the variation of the eigenvalues and $\bar{d}\varrho^{(\text{ep})} = \sum_k r_k (|dr_k\rangle \langle r_k| + |r_k\rangle \langle dr_k|)$ is due to the variation in the eigenprojectors, where \bar{d} denotes inexact differential. In addition, we observe that the entropy $S(\varrho) = -\text{Tr}[\varrho \ln \varrho]$ changes only when the eigenvalues vary,

$$dS = -\sum_k dr_k \ln r_k = -\text{Tr}[\bar{d}\varrho^{(\text{ev})} \ln \varrho]. \quad (2)$$

In a system with the physical Hamiltonian H , the system internal energy $U = \text{Tr}[\varrho H]$ changes along the trajectory with $t \rightarrow t + dt$ as $U \rightarrow U + dU$, with $dU = \text{Tr}[d\varrho H] + \text{Tr}[\varrho dH]$. Here, the contribution associated with $\bar{d}\varrho^{(\text{ev})}$ is solely related to the change of the eigenvalues, which is in line with how the entropy changes. Hence, we define the heat change as

$$\bar{d}Q = \text{Tr}[\bar{d}\varrho^{(\text{ev})} H] = \sum_k dr_k \langle r_k | H | r_k \rangle, \quad (3)$$

and assign the remaining variations to work change,

$$\begin{aligned} \bar{d}W &= \text{Tr}[\varrho dH] + \text{Tr}[\bar{d}\varrho^{(\text{ep})} H] \\ &= \sum_k r_k (\langle r_k | dH | r_k \rangle + \langle dr_k | H | r_k \rangle + \langle r_k | H | dr_k \rangle), \end{aligned} \quad (4)$$

such that

$$dU = \bar{d}Q + \bar{d}W \quad (5)$$

encompasses the first law of thermodynamics.

Dynamical analysis of the heat and work definitions.—We use a recently proposed *trajectory-based shortcut to adiabaticity* (TB-STA) framework for open-system dynamics, which

identifies a particular dynamical equation of motion that generates any given time-dependent density matrix or trajectory [52]. This technique, based on the general Lindblad-like equation described later, is not limited by the system-environment coupling strength or initial correlation. Since the TB-STA equation describes a given trajectory, its coherent and dissipative parts are directly related to the coherences and dissipation in the course of the evolution, which contrast with the conventional Markovian Lindblad master equation. Our main result shows that, among all possible dynamical equations describing a given trajectory, TB-STA allows us to unambiguously separate the change of the internal energy into heat and work for any open system. We prove that energy change assigned to the (so-called) dissipative part of the equation is accompanied by entropy change, and it is therefore *heat*. The coherent part, which does not involve entropy change, corresponds to a *dissipative work*, which for its relation to nonunitarity of the dynamics entails implicit existence of some ambient environment.

Having a trajectory $\{\varrho(t)\}$ at hand, an associated differential equation describing its dynamics is given by the following Lindblad-like equation [52]:

$$\dot{\varrho} = -i[\mathfrak{h}, \varrho] + \mathbb{D}_{\text{cd}}[\varrho], \quad (6)$$

where

$$\mathfrak{h} = i \sum_k (|\dot{r}_k\rangle \langle r_k| - \langle r_k | \dot{r}_k \rangle |r_k\rangle \langle r_k|), \quad (7)$$

$$\mathbb{D}_{\text{cd}}[\varrho] = \sum_{k,j} \mathfrak{C}_{kj} (\mathbb{L}_{kj} \varrho \mathbb{L}_{kj}^\dagger - \frac{1}{2} \{ \mathbb{L}_{kj}^\dagger \mathbb{L}_{kj}, \varrho \}), \quad (8)$$

$$\mathbb{L}_{kj} = |r_k\rangle \langle r_j|, \quad \mathfrak{C}_{kj} = (1 - \delta_{r_j 0}) \dot{r}_k / (Dr_j), \quad (9)$$

with the dot denoting time differentiation. Several remarks are in order: (i) \mathfrak{h} operates as a *counterdiabatic* (CD) Hamiltonian which alone generates a parallel transport $|r_k(0)\rangle \rightarrow |r_k(t)\rangle \forall k$ (cf. CD Hamiltonians in the energy sense [53, 54]); (ii) in the CD dissipator \mathbb{D}_{cd} the anticommutator identically vanishes, whence the dissipator reduces exclusively to jumps in the instantaneous eigenbasis of ϱ ; (iii) the dynamical equation (6) is universal, irrespective of any physical setting (system Hamiltonian and the environment) in which the system has obtained this trajectory. In the standard technical sense [55, 56], this dynamical equation for a trajectory is different from the master equation of the system. The same physical system prepared in a different initial condition with the same physical setting can yield a different trajectory—hence a different dynamical equation associated with the new trajectory; and (iv) a Lindblad-like master equation for general open-system dynamics has also been introduced recently [57].

Comparison with the conventional definitions.—Along the trajectory, the conventional definitions are $Q = \int_0^t dt \text{Tr}[\dot{\varrho} H]$ for heat change and $W = \int_0^t dt \text{Tr}[\varrho \dot{H}]$ for work exchange, hence $dU = \bar{d}Q + \bar{d}W$ [cf. Eq. (5)]. Now rewrite $\bar{d}Q$ as

$$\bar{d}Q = \sum_k dr_k \langle r_k | H | r_k \rangle + \sum_k r_k (\langle dr_k | H | r_k \rangle + \langle r_k | H | dr_k \rangle). \quad (10)$$

The first sum originates from a change in the eigenvalues of the state, contributing when not all dr_k s vanish (similar to the reason for the change in S); the second sum is solely associated with the change of the eigenvectors. The first sum can be

rewritten as $\text{Tr}[\mathbb{D}_{\text{cd}}[\varrho]H]dt$, which is related to the *dissipative* part of the trajectory dynamical equation (6); whereas the second sum can be recast as $-i\text{Tr}[[H, \mathbb{h}]\varrho]dt$, which is associated with the *coherent* part of the dynamical equation—hence dissipative *work*. From this perspective, we can consider the following relations as a justification of our definitions:

$$dQ = dQ - dW_{\text{cd}} = \text{Tr}[\mathbb{D}_{\text{cd}}[\varrho]H]dt, \quad (11)$$

$$dW = dW + dW_{\text{cd}} = \text{Tr}[\varrho(\dot{H} - i[H, \mathbb{h}])]dt, \quad (12)$$

where dW_{cd} is an environment-induced dissipative work,

$$\begin{aligned} dW_{\text{cd}} &= \sum_k r_k (\langle dr_k | H | r_k \rangle + \langle r_k | H | dr_k \rangle) \\ &= -i\text{Tr}[[H, \mathbb{h}]\varrho]dt, \end{aligned} \quad (13)$$

and is generated due to the CD evolution. In this framework the work is due to both driving the system through varying its physical Hamiltonian (dW) and the CD evolution along the trajectory due to the environment (dW_{cd}). Importantly, unlike in Ref. [53], the CD term here is not related to an external physical CD control of the Hamiltonian; rather, it corresponds to the natural CD evolution along the state trajectory.

Relation to the semiclassical definitions.—In the *semiclassical* formulation of thermodynamics for quantum systems, heat and work are defined differently [36]. Using the instantaneous eigenbasis of the system Hamiltonian $H = \sum_n E_n |E_n\rangle\langle E_n|$, we obtain $U = \sum_n p_n E_n$, where $p_n = \langle E_n | \varrho | E_n \rangle$ is the population of the energy eigenstate $|E_n\rangle$. From the identity $dU = \sum_n dp_n E_n + \sum_n p_n dE_n$, one can read the semiclassical heat and work variations as

$$dq = \sum_n dp_n E_n, \quad (14)$$

$$dw = \sum_n p_n dE_n. \quad (15)$$

However, if instead of using the instantaneous eigenbasis of H , we evaluate the trace in U in the instantaneous eigenbasis of ϱ [Eq. (1)], then $U = \sum_k r_k H_k$ and $dU = \sum_k dr_k H_k + \sum_k r_k dH_k$, where $H_k = \langle r_k | H | r_k \rangle$. As in the semiclassical setting, one can now read the first term as the heat change and the second one as the work change. We observe that these are equivalent to the TB-STA definitions [Eqs. (11) and (12)], $dQ = \sum_k dr_k H_k$ and $dW = \sum_k r_k dH_k$. One can argue that the instantaneous eigenbasis of ϱ is preferred in calculating the energy contributions [58]. A closer inspection of dq reveals that it includes the energy (not heat) change dW_{cd} which we have assigned to the CD Hamiltonian in the form of work,

$$dq = dQ + dW_{\text{cd}} + \sum_n E_n (\langle dE_n | \varrho | E_n \rangle + \langle E_n | \varrho | dE_n \rangle). \quad (16)$$

Thus, since dq has contributions from both heat and work exchanges (in our sense), we conclude that in general the semiclassical definitions of heat and work fail to properly account for various contributions to the internal energy change.

Irreversible/Internal entropy production.—The entropy change dS [Eq. (2)], together with the heat change dQ [Eq. (11)] give the change in the irreversible entropy \mathcal{S} as

$$d\mathcal{S} \equiv dS - \beta dQ = \sum_k dr_k \langle r_k | (\mathbb{H} - \beta H) | r_k \rangle, \quad (17)$$

where $\mathbb{H} = -\log \varrho$ and β is the nonequilibrium, instantaneous inverse temperature of the system given by (if $k_B \equiv 1$) [1]

$$\beta = (\partial S / \partial U)_{x_2, x_3, \dots} = \text{Cov}(H, \mathbb{H}) / \text{Cov}(H, H). \quad (18)$$

Here $\{x_i\}$ are a set of independent variables obtained from the expectation values of a complete set of traceless orthonormal observables $\{O_i\}_{i=2}^{D^2-1}$, with $O_0 = \mathbb{1}/\sqrt{D}$ (the normalized identity operator of dimension D) and $O_1 = (H - \text{Tr}[H]/D)/\sqrt{D \text{Cov}(H, H)}$. In addition, $\text{Cov}(X, Y) \equiv \text{Tr}[XY]/D - \text{Tr}[X]\text{Tr}[Y]/D^2$. In terms of the $\{x_i\}$ variables we have $d\varrho^{(\text{ev})} = [D \text{Cov}(H, H)]^{-1/2} dQ H + \sum_{i \geq 2} dx_i O_i$, where $dx_i = \text{Tr}[d\varrho^{(\text{ev})} O_i]$. As a result, we obtain $dS = \beta dQ + \sum_{i \geq 2} \text{Tr}[O_i \mathbb{H}] dx_i$, whence

$$d\mathcal{S} = \sum_{i \geq 2} \text{Tr}[O_i \mathbb{H}] dx_i, \quad (19)$$

which is indeed independent of the choice of $\{x_i\}$ [60]. We observe that $d\mathcal{S}$ is determined only in terms of the system variables, independent of the existence of an environment in a thermal equilibrium, and the system Hamiltonian does not play any explicit role therein; it is an irrelevant observable—see Ref. [60] for further discussion. We remark that, besides using dQ in our definition of irreversible entropy production (17), unlike the earlier literature [61] here β is associated to the system (not to a large environment or heat bath). A discussion of why β is associated with the system can be found in Supplemental Material [60]. Another alternative expression for $d\mathcal{S}$ can be obtained in terms of the relative entropy $S(\varrho || \varrho_{\text{eq}}) = \text{Tr}[\varrho \ln \varrho - \varrho \ln \varrho_{\text{eq}}]$ between the state ϱ and the instantaneous canonical Gibbs state $\varrho_{\text{eq}} \equiv e^{-\beta H} / \text{Tr}[e^{-\beta H}]$ as [60] $d\mathcal{S} = -dS(\varrho || \varrho_{\text{eq}}) + \beta dW_{\text{cd}} + \text{Tr}[(\varrho - \varrho_{\text{eq}})d(\beta H)]$. We also note that interestingly from Eq. (19) an inequality can also be derived as [60]

$$d\mathcal{S} + b(\mathbb{O}) dB \geq 0, \quad (20)$$

which is reminiscent of a generalized Clausius inequality [62, 63] with all quantities depending explicitly on the system. Here $b(\mathbb{O})$ is the spread of $\mathbb{O} = \sum_{i \geq 2} O_i \otimes O_i$ and $dB = -\sum_k \ln r_k \sum_{k' | dr_{k'} \geq 0} dr_{k'}$.

For dynamically *closed quantum systems* (where $\dot{\varrho} = -i[H, \varrho]$, assuming $\hbar \equiv 1$), we have $\dot{r}_k = 0 \forall k$, hence $dU = dW = dQ$ and $d\mathcal{S} = dS = dQ = dQ = dW_{\text{cd}} = 0$. This case also naturally includes phase-space-preserving cooling processes [64]. For dynamically *open quantum systems* weakly coupled to a large environment, when H is constant or slowly driven the dynamics obeys a Markovian Lindblad master equation $\dot{\varrho} = \mathcal{L}[\varrho]$, where $\mathcal{L}[\varrho] = -i[H + H_{\text{Lamb}}, \varrho] + \mathcal{D}[\varrho]$, with H_{Lamb} being the environment-induced Lamb-shift correction and $\mathcal{D}[\varrho] = \sum_\alpha c_\alpha (L_\alpha \varrho L_\alpha^\dagger - (1/2)\{L_\alpha^\dagger L_\alpha, \varrho\})$ ($c_\alpha > 0$) being the quantum dissipator [55, 56] [cf Eq. (8)]. Additionally, if the system starts at $\varrho_{\text{eq}}(0)$ and also $\varrho_{\text{eq}}(t)$ is the unique instantaneous steady state of the dynamics [65, 66], one can prove that $\varrho(t) \approx \varrho_{\text{eq}}(t) \forall t$ [67], namely the trajectory remains near the steady state, and also $dS - \beta dQ \geq 0$ [10, 21, 33]. This implies that $d\mathcal{S} - \beta dW_{\text{cd}} \geq 0$, where dW_{cd} [Eq. (13)] is an energetic cost associated to how different the real trajectory is from the quasistatic one—cf. the




FIG. 1. Rates of internal energy (solid orange), heat (solid blue), and work (dashed red) changes vs. time (in natural units $\hbar \equiv k_B \equiv 1$), for a qubit in a heat bath as given in example I, with $\gamma = 0.1$ and $\omega_0 = 1$. The integrated values are $U = -0.25$, $Q = -0.138$, and $W = -0.112$, respectively, where $X = \int_0^{10} dt \dot{X}$ with $X \in \{U, Q, W\}$. Inset compares \dot{S} and \dot{Q} .

general relation (20). This relation can be considered as a manifestation of the Clausius inequality [62, 63]. Note that in general $dW \neq d\mathcal{W}$ and $dQ \neq d\mathcal{Q} = \text{Tr}[\mathcal{D}[\varrho]H]dt$ [cf. Eq. (11)]; whereas in the *quasistatic* regime $\varrho(t) = \varrho_{\text{eq}}(t)$, we have $|r_k\rangle = |E_k\rangle$, $\mathbb{H} = \beta H$, and thus $dW_{\text{cd}} = 0$, $dQ = d\mathcal{Q}$, $dS = 0$, and $dS = \beta dQ$. The latter is a manifestation of the thermodynamic reversibility [49].

Example I: Qubit in a Markovian environment.—Consider a qubit with $H = \omega_0 \sigma_z$ that weakly interacts with a Markovian environment such that $\dot{\varrho} = -i[\omega_0 \sigma_z, \varrho] + \gamma(\sigma_x \varrho \sigma_x - \varrho)$, where ω_0 and γ are positive constants, and $\sigma_x = |0\rangle\langle 1| + |1\rangle\langle 0|$ and $\sigma_z = |0\rangle\langle 0| - |1\rangle\langle 1|$ are the x - and z -Pauli operators [55]. The quantity $\dot{U} = \text{Tr}[H\dot{\varrho}] = -2\gamma\omega_0 \text{Tr}[\sigma_z \varrho]$ is fully due to the dissipative part of the dynamics. We consider three different initial states (for details, see Ref. [60]). (i) Thermal state $\varrho_{\text{eq}}(0)$, with $\beta(0)$ being the initial inverse temperature of the system (this can differ from that of the environment, and the latter need not be in a thermal state). In this case, we obtain $\dot{Q} = \dot{U} = 2\gamma\omega_0 e^{-2\gamma t} \tanh[\beta(0)\omega_0]$, $\dot{q} = \dot{Q}$, and $\dot{w} = \dot{W}$. (ii) Pure state $|\psi(0)\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$. One can see that $\dot{U} = \dot{Q} = \dot{W} = \dot{q} = \dot{w} = 0$. (iii) Starting from $\varrho(0) = (\mathbb{1} + [\sigma_x + \sigma_z]/2)/2$, we can obtain the heat, work, and internal energy change in the system—Fig. 1. It can be shown that $\dot{q} = \dot{U} = -(\gamma/2)e^{-2\gamma t}$ and $\dot{w} = 0$.

Example II: Qubit in a dephasing environment.—Consider the dephasing master equation for a qubit, $\dot{\varrho} = -i[\omega_0 \sigma_z, \varrho] + \gamma(\sigma_z \varrho \sigma_z - \varrho)$. Generally, in this process the system energy is preserved and there is no net energy exchange between the system and the environment. However, the system and the environment can still exchange heat and work. This can be contrasted with Landauer's principle, where the environment does work to erase phase information and the system releases heat back into the environment. Let $\omega_0 = 1$ and $\gamma = 0.1$ and consider two different cases: (i) The initial state is $|\psi(0)\rangle = (|0\rangle +$

$\sqrt{2}|1\rangle)/\sqrt{3}$: we obtain that $\dot{Q} = -\dot{W}_{\text{cd}} = (8/15)/[8 + e^{2t/5}]$ and $\dot{S} = 4 \ln[(1 + 2\Delta)/(1 - 2\Delta)]/(15e^{t/5}\sqrt{8 + e^{2t/5}})$, where $\Delta = (1/6)e^{-t/5}\sqrt{8 + e^{2t/5}}$. This process can be compared with an isothermal process in ideal gases, where the energy is constant and the heat exchanged is equal to the work exchanged with the opposite sign. However, since this dynamics does not exactly model an ideal gas (its internal energy is not proportional to its temperature), β varies and thus the process is not isothermal. (ii) The initial state is $|\psi(0)\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$: in this case $dU = dQ = dW_{\text{cd}} = 0$. However, $dS \neq 0$, which means that the whole entropy change is due to the irreversible entropy production. This can be compared with a free expansion process in classical (nonideal) gases. For a further example of a driven qubit, see Ref. [68].

Another example, a quantum damped harmonic oscillator in an environment of oscillators, has been worked out in Ref. [60]. It appears that at all times the state of the oscillator is a Gibbs state with a time-dependent inverse temperature and a constant Hamiltonian. Because only the eigenvalues of the state change in time and the eigenvectors remain constant, dW_{cd} vanishes. Hence heat and work here reduce to the values obtained from the conventional definitions.

Summary.—We have revisited the assignment of thermodynamic quantities to an open quantum system strongly coupled to an environment. In general, there is no unique way of separating the system internal energy from that of the environment. Despite this fundamental issue, by introducing a dissipative work, we have shown that it is possible to consistently split the internal energy change into work change (causing no entropy change) and heat change (which can cause entropy change). The key ingredient is to use the trajectory-based description of the state of the system and its associated equation of motion, which is universally valid for any coupling strength and yields a spectral decomposition of the system density matrix, separating the changes in the eigenvalues from those of the eigenvectors. We have compared our entropy-based definitions with the conventional and semiclassical ones and have argued that these two approaches are inadequate. More importantly, by using a definition of nonequilibrium temperature in quantum systems, we have obtained the irreversible entropy production in a general dynamical process. We have demonstrated that the irreversible entropy production is a function of the system variables other than the energy and does not have any explicit dependence on the environmental degrees of freedom. We have also derived an inequality which can be contrasted with the generalized Clausius inequality. The consistency of our formalism has been illustrated in paradigmatic scenarios.

Acknowledgement.—We thank F. J. Gómez-Ruiz and M. Zahidy for discussions. This work has been supported in part by the Academy of Finland's Center of Excellence QTF Project 312298 and Sharif University of Technology's Office of Vice President for Research and Technology.

Note added.—Recently, we learned about another independent related study [69].

- [2] F. Binder, L. A. Correa, C. Gogolin, J. Anders, and G. Adesso, eds., *Thermodynamics in the Quantum Regime: Fundamental Aspects and New Directions* (Springer International, Cham, Switzerland, 2018).
- [3] S. Deffner and S. Campbell, *Quantum Thermodynamics – An Introduction to the Thermodynamics of Quantum Information* (Morgan & Claypool, San Rafael, CA, 2019).
- [4] R. Kosloff, Quantum thermodynamics: A dynamical viewpoint, *Entropy* **15**, 2100 (2013).
- [5] J. Goold, M. Huber, A. Riera, L. del Rio, and P. Skrzypczyk, The role of quantum information in thermodynamics – a topical review, *J. Phys. A: Math. Theor.* **49**, 143001 (2016).
- [6] T. D. Kieu, The Second Law, Maxwell’s Demon, and Work Derivable from Quantum Heat Engines, *Phys. Rev. Lett.* **93**, 140403 (2004).
- [7] F. Brandão, M. Horodecki, N. Ng, J. Oppenheim, and S. Wehner, The second laws of quantum thermodynamics, *Proc. Natl. Acad. Sci. U.S.A.* **112**, 3275 (2015).
- [8] L. Masanes and J. Oppenheim, A general derivation and quantification of the third law of thermodynamics, *Nature Commun.* **8**, 14538 (2017).
- [9] H. Hossein-Nejad, E. J. O’Reilly, and A. Olaya-Castro, Work, heat and entropy production in bipartite quantum systems, *New J. Phys.* **17**, 075014 (2015).
- [10] S. Alipour, F. Benatti, F. Bakhshinezhad, M. Afsary, S. Marcantoni, and A. T. Rezakhani, Correlations in quantum thermodynamics: Heat, work, and entropy production, *Sci. Rep.* **6**, 35568 (2016).
- [11] S. Marcantoni, S. Alipour, F. Benatti, R. Floreanini, and A. T. Rezakhani, Entropy production and non-Markovian dynamical maps, *Sci. Rep.* **7**, 12447 (2017).
- [12] M. Esposito, U. Harbola, and S. Mukamel, Nonequilibrium fluctuations, fluctuation theorems, and counting statistics in quantum systems, *Rev. Mod. Phys.* **81**, 1665 (2009).
- [13] M. Campisi, P. Hänggi, and P. Talkner, Quantum fluctuation relations: Foundations and applications, *Rev. Mod. Phys.* **83**, 771 (2011).
- [14] K. Funo, M. Ueda, and T. Sagawa, Quantum fluctuation theorems, in *Thermodynamics in the Quantum Regime: Fundamental Aspects and New Directions*, edited by F. Binder, L. A. Correa, C. Gogolin, J. Anders, and G. Adesso (Springer International, Cham, Switzerland, 2018).
- [15] F. Bakhshinezhad, F. Clivaz, G. Vitagliano, P. Erker, A. Reza-khani, M. Huber, and N. Friis, Thermodynamically optimal creation of correlations, *J. Phys. A: Math. Theor.* **52**, 465303 (2019).
- [16] J. Roßnagel, S. T. Dawkins, K. N. Tolazzi, O. Abah, E. Lutz, F. Schmidt-Kaler, and K. Singer, A single-atom heat engine, *Science* **352**, 325 (2016).
- [17] S. Deng, A. Chenu, P. Diao, F. Li, S. Yu, I. Coulamy, A. del Campo, and H. Wu, Superadiabatic quantum friction suppression in finite-time thermodynamics, *Sci. Adv.* **4**, eaar5909 (2018).
- [18] G. Maslennikov, S. Ding, R. Häblützel, J. Gan, A. Roulet, S. Nimmrichter, J. Dai, V. Scarani, and D. Matsukevich, Quantum absorption refrigerator with trapped ions, *Nature Commun.* **10**, 202 (2019).
- [19] J. P. S. Peterson, T. B. Batalhão, M. Herrera, A. M. Souza, R. S. Sarthour, I. S. Oliveira, and R. M. Serra, Experimental Characterization of a Spin Quantum Heat Engine, *Phys. Rev. Lett.* **123**, 240601 (2019).
- [20] D. von Lindenfels, O. Gräb, C. T. Schmiegelow, V. Kaushal, J. Schulz, M. T. Mitchison, J. Goold, F. Schmidt-Kaler, and U. G. Poschinger, A Spin Heat Engine Coupled to a Harmonic Oscillator Flywheel, *Phys. Rev. Lett.* **123**, 080602 (2019).
- [21] R. Alicki, The quantum open system as a model of the heat engine, *J. Phys. A: Math. Gen.* **12**, L103 (1979).
- [22] P. Talkner, E. Lutz, and P. Hänggi, Fluctuation theorems: Work is not an observable, *Phys. Rev. E* **75**, 050102 (2007).
- [23] J. M. G. Vilar and J. M. Rubi, Failure of the Work-Hamiltonian Connection for Free-Energy Calculations, *Phys. Rev. Lett.* **100**, 020601 (2008).
- [24] D. Gelbwaser-Klimovsky and A. Aspuru-Guzik, On thermodynamic inconsistencies in several photosynthetic and solar cell models and how to fix them, *Chem. Sci.* **8**, 1008 (2017).
- [25] W. Niedenzu, M. Huber, and E. Boukobza, Concepts of work in autonomous quantum heat engines, *Quantum* **3**, 195 (2019).
- [26] R. Sampaio, S. Suomela, T. Ala-Nissila, J. Anders, and T. G. Philbin, Quantum work in the Bohmian framework, *Phys. Rev. A* **97**, 012131 (2018).
- [27] C. Elouard, D. A. Herrera-Martí, M. Clusel, and A. Auffèves, The role of quantum measurement in stochastic thermodynamics, *npj Quantum Info.* **3**, 9 (2017).
- [28] C. Jarzynski, Nonequilibrium work theorem for a system strongly coupled to a thermal environment, *J. Stat. Mech.: Theo. Exp.* **2004**, P09005 (2004).
- [29] M. F. Gelin and M. Thoss, Thermodynamics of a subensemble of a canonical ensemble, *Phys. Rev. E* **79**, 051121 (2009).
- [30] M. Campisi, P. Talkner, and P. Hänggi, Fluctuation Theorem for Arbitrary Open Quantum Systems, *Phys. Rev. Lett.* **102**, 210401 (2009).
- [31] H. J. D. Miller and J. Anders, Energy-temperature uncertainty relation in quantum thermodynamics, *Nature Commun.* **9**, 2203 (2018).
- [32] Á. Rivas, Strong Coupling Thermodynamics of Open Quantum Systems, *Phys. Rev. Lett.* **124**, 160601 (2020).
- [33] H. Spohn, Entropy production for quantum dynamical semi-groups, *J. Math. Phys.* **19**, 1227 (1978).
- [34] C. M. Bender, D. C. Brody, and B. K. Meister, Quantum mechanical Carnot engine, *J. Phys. A: Math. Gen.* **33**, 4427 (2000).
- [35] S. Abe, Maximum-power quantum-mechanical Carnot engine, *Phys. Rev. E* **83**, 041117 (2011).
- [36] A. Polkovnikov, Microscopic diagonal entropy and its connection to basic thermodynamic relations, *Ann. Phys. (N.Y.)* **326**, 486 (2011).
- [37] Y.-H. Shi, H.-L. Shi, X.-H. Wang, M.-L. Hu, S.-Y. Liu, W.-L. Yang, and H. Fan, Quantum coherence in a quantum heat engine, *J. Phys. A: Math. Theor.* **53**, 085301 (2020).
- [38] S. Su, J. Chen, Y. Ma, J. Chen, and C. Sun, The heat and work of quantum thermodynamic processes with quantum coherence, *Chin. Phys. B* **27**, 060502 (2018).
- [39] R. Dann, A. Tobalina, and R. Kosloff, Shortcut to Equilibration of an Open Quantum System, *Phys. Rev. Lett.* **122**, 250402 (2019).
- [40] Z. N. Qaleh and A. T. Rezakhani, Enhancing energy transfer in quantum systems via periodic driving: Floquet master equations, *arXiv:2105.14358* (2021).
- [41] K. Funo, N. Shiraishi, and K. Saito, Speed limit for open quantum systems, *New J. Phys.* **21**, 013006 (2019).
- [42] D. A. Lidar and K. B. Whaley, Decoherence-free subspaces and subsystems, in *Irreversible Quantum Dynamics*, edited by F. Benatti and R. Floreanini (Springer, Berlin, 2003) p. 83.
- [43] R. Blume-Kohout, H. K. Ng, D. Poulin, and L. Viola, Characterizing the Structure of Preserved Information in Quantum Processes, *Phys. Rev. Lett.* **100**, 030501 (2008).
- [44] W. Niedenzu, V. Mukherjee, A. Ghosh, A. G. Kofman, and G. Kurizki, Quantum engine efficiency bound beyond the second law of thermodynamics, *Nature Commun.* **9**, 165 (2018).
- [45] A. E. Allahverdyan, R. Balian, and T. M. Nieuwenhuizen, Maximal work extraction from finite quantum systems, *Europhys. Lett.*

- 67, 565 (2004).
- [46] G. Manzano, Squeezed thermal reservoir as a generalized equilibrium reservoir, *Phys. Rev. E* **98**, 042123 (2018).
- [47] J. M. Horowitz and J. M. R. Parrondo, Entropy production along nonequilibrium quantum jump trajectories, *New J. Phys.* **15**, 085028 (2013).
- [48] J. J. Alonso, E. Lutz, and A. Romito, Thermodynamics of Weakly Measured Quantum Systems, *Phys. Rev. Lett.* **116**, 080403 (2016).
- [49] H. B. Callen, *Thermodynamics and Introduction to Thermostatistics* (John Wiley & Sons, New York, 1985).
- [50] H. Weimer, M. J. Henrich, F. Rempp, H. Schröder, and G. Mahler, Local effective dynamics of quantum systems: A generalized approach to work and heat, *Europhys. Lett.* **83**, 30008 (2008).
- [51] Y. A. Çengel and M. A. Boles, *Thermodynamics: An Engineering Approach* (McGraw-Hill, New York, 2007).
- [52] S. Alipour, A. Chenu, A. T. Rezakhani, and A. del Campo, Shortcuts to adiabaticity in driven open quantum systems: Balanced gain and loss and non-Markovian evolution, *Quantum* **4**, 336 (2020).
- [53] M. Demirplak and S. A. Rice, Adiabatic population transfer with control fields, *J. Phys. Chem. A* **107**, 9937 (2003).
- [54] M. V. Berry, Transitionless quantum driving, *J. Phys. A: Math. Theor.* **42**, 365303 (2009).
- [55] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, New York, 2007).
- [56] Á. Rivas and S. F. Huelga, *Open Quantum Systems* (Springer, Berlin, 2012).
- [57] S. Alipour, A. T. Rezakhani, A. P. Babu, K. Mølmer, M. Möttönen, and T. Ala-Nissila, Correlation-Picture Approach to Open Quantum-System Dynamics, *Phys. Rev. X* **10**, 041024 (2020).
- [58] Heat is associated with entropy change. Note that $S(\varrho) = \min_{\{\Pi_k\}} S(\{l_k\})$, where $\{\Pi_k\}$ is a resolution of the identity, $\Pi_k \geq 0$ and $\sum_k \Pi_k = \mathbb{1}$, $\{l_k = \text{Tr}[\Pi_k \varrho]\}$ constitutes probabilities, and $S(\{l_k\}) = -\sum_k l_k \log l_k$ is the Shannon entropy [V. Vedral, Classical Correlations and Entanglement in Quantum Measurements, *Phys. Rev. Lett.* **90**, 050401]. The optimal measurement minimizing the Shannon entropy is given by the eigenvectors of ϱ ; $\Pi_k = |r_k\rangle\langle r_k|$.
- [59] S. Alipour, M. Afsary, F. Bakhshinezhad, M. Ramezani, F. Benatti, T. Ala-Nissila, and A. T. Rezakhani, Temperature in nonequilibrium quantum systems, *arXiv:2105.11915* (2021).
- [60] See Supplemental Material.
- [61] G. T. Landi and M. Paternostro, Irreversible entropy production: From classical to quantum, *Rev. Mod. Phys.* (to be published).
- [62] R. Uzdin and S. Rahav, Global Passivity in Microscopic Thermodynamics, *Phys. Rev. X* **8**, 021064 (2018).
- [63] M. Esposito, K. Lindenberg, and C. Van den Broeck, Entropy production as correlation between system and reservoir, *New J. Phys.* **12**, 013013 (2010).
- [64] X. Chen, A. Ruschhaupt, S. Schmidt, A. del Campo, D. Guéry-Odelin, and J. G. Muga, Fast Optimal Frictionless Atom Cooling in Harmonic Traps: Shortcut to Adiabaticity, *Phys. Rev. Lett.* **104**, 063002 (2010).
- [65] H. Spohn, An algebraic condition for the approach to equilibrium of an open N -level system, *Lett. Math. Phys.* **2**, 33 (1977).
- [66] V. Bach, J. Fröhlich, and I. M. Sigal, Return to equilibrium, *J. Math. Phys.* **41**, 3985 (2000).
- [67] V. Cavina, A. Mari, and V. Giovannetti, Slow Dynamics and Thermodynamics of Open Quantum Systems, *Phys. Rev. Lett.* **119**, 050601 (2017).
- [68] A. Juan-Delgado and A. Chenu, First law of quantum thermodynamics in a driven open two-level system, *arXiv:2104.10691* (2021).
- [69] B. Ahmadi, S. Salimi, and A. S. Khorashad, Refined definitions of heat and work in quantum thermodynamics, *arXiv:1912.01983* (2019).

Supplemental Material

Appendix A: Irreversible entropy production

Irreversible entropy production is defined as the difference between the entropy dS and the reversible entropy. Using the temperature of the system T as introduced in Ref. [1] and Eq. (18) of the main text the irreversible entropy production is given by $d\mathcal{S} = dS - \beta d\mathcal{Q}$. By expanding $d\varrho^{(\text{ev})}$ in terms of the set of orthonormal traceless observables $\{O_i\}$ (described in the main text) as

$$d\varrho^{(\text{ev})} = \sum_i dx_i O_i, \quad (\text{A1})$$

where $dx_i = \text{Tr}[d\varrho^{(\text{ev})} O_i]$ we can obtain from the entropy change $dS = -\text{Tr}[d\varrho^{(\text{ev})} \ln \varrho]$ that

$$\begin{aligned} dS &= -\text{Tr}[d\varrho^{(\text{ev})} O_1] \text{Tr}[O_1 \ln \varrho] - \sum_{i \geq 2} \text{Tr}[d\varrho^{(\text{ev})} O_i] \text{Tr}[O_i \ln \varrho] \\ &= -\frac{1}{\sqrt{\text{Cov}(H, H)}} \text{Tr}[O_1 \ln \varrho] d\mathcal{Q} - \sum_{i \geq 2} \text{Tr}[d\varrho^{(\text{ev})} O_i] \text{Tr}[O_i \ln \varrho], \\ &= \beta d\mathcal{Q} - \sum_{i \geq 2} \text{Tr}[d\varrho^{(\text{ev})} O_i] \text{Tr}[O_i \ln \varrho], \\ &= \beta d\mathcal{Q} - \sum_{i \geq 2} \sum_j dr_j \langle r_j | O_i | r_j \rangle \text{Tr}[O_i \ln \varrho], \end{aligned} \quad (\text{A2})$$

where we have used the definition of inverse temperature β [Eq. (18) of the main text]. The expression for the irreversible entropy production is obtained as

$$\begin{aligned} d\mathcal{S} &= -\sum_{i \geq 2} \text{Tr}[d\varrho^{(\text{ev})} O_i] \text{Tr}[O_i \ln \varrho] \\ &= -\sum_{i \geq 2} \sum_j dr_j \langle r_j | O_i | r_j \rangle \text{Tr}[O_i \ln \varrho]. \end{aligned} \quad (\text{A3})$$

It should be noted that despite what may seem at first, $d\mathcal{S}$ is independent of the choice of the basis operators $\{O_i\}_{i \geq 2}$. This can be inferred from the fact that the entropy change $dS = -\text{Tr}[d\varrho \ln \varrho]$ is independent of the choice of basis, and $d\mathcal{Q}$ (heat depends only on O_1). Thus, one can conclude that the choice of observables $\{O_{i \geq 2}\}$ does not affect the irreversible entropy production. We also prove this observation in a different manner in the following.

To see that $d\mathcal{S}$ is independent of our choice of basis operators $\{O_i\}_{i=0}^{D^2-1}$, let $\{O'_i\}_{i=0}^{D^2-1}$ be another set of orthonormal operators such that still $O'_0 = O_0$ and $O'_1 = O_1$. These bases must be related through an orthogonal matrix \mathcal{U} as $O_i = \sum_{k=2}^{D^2-1} \mathcal{U}_{ik} O'_k$, for $i \geq 2$. From Eq. (A3) one finds

$$\begin{aligned} d\mathcal{S} &= -\sum_{i=2}^{D^2-1} \text{Tr}[O_i \ln \varrho] \text{Tr}[O_i d\varrho^{(\text{ev})}] \\ &= -\sum_{i=2}^{D^2-1} \sum_{k=2}^{D^2-1} \sum_{l=2}^{D^2-1} \mathcal{U}_{ik} \mathcal{U}_{il} \text{Tr}[O'_k \ln \varrho] \text{Tr}[O'_l d\varrho^{(\text{ev})}] \\ &= -\sum_{k=2}^{D^2-1} \sum_{l=2}^{D^2-1} (\sum_{i=2}^{D^2-1} (\mathcal{U}^T)_{ki} (\mathcal{U})_{il}) \text{Tr}[O'_k \ln \varrho] \text{Tr}[O'_l d\varrho^{(\text{ev})}] \\ &= -\sum_{k=2}^{D^2-1} \text{Tr}[O'_k \ln \varrho] \text{Tr}[O'_k d\varrho^{(\text{ev})}], \end{aligned}$$

which proves $d\mathcal{S}$ is independent of the choice of basis.

Remark.—One can observe that [1] β appears naturally as a variable of the state ϱ as

$$\varrho = (1/Z) e^{-\beta H + \sum_{i \geq 2} c_i O_i}, \quad (\text{A4})$$

where $Z = e^{-\text{Tr}[\beta H]}$ and $c_i = -\text{Tr}[O_i H]$.

Appendix B: Further justification for $d\mathcal{S}$

Here, to argue that $d\mathcal{S}$, as in Eq. (19) of the main text, can well describe the irreversible entropy production, we revisit two classic processes in thermodynamics: free expansion and Joule's experiment [2]—Fig. 2.

1. Free expansion

Assume an isolated rigid container of an ideal gas such that the gas is initially confined in one half of the container, separated by an almost massless partition from the other half—no energy exchange with the environment. The partition is then removed



FIG. 2. (Left) Schematic of the free expansion. An ideal gas is confined in a half of an adiabatic container by using a partition. The partition is then removed and the gas fills the whole container. (Right) Schematic of Joule's experiment. By rotating paddles we perform work on a fluid inside an adiabatic container.

suddenly and the gas can fill the whole volume of the container. This process does not involve any energy change, since the temperature is constant (both heat and work are zero). However, pressure and volume are two other variables of the system—one of which can be considered as an independent variable since the state equation connects these variables. Although no energy change occurs in the system, because the change of pressure (and volume) causes irreversible entropy production in the system.

2. Joule's experiment

Assume a fluid (a liquid or a nonideal gas) inside a container with adiabatic (i.e., thermally isolated) walls. By rotating paddles from outside of the container it is possible to perform work on the fluid, which leads to an increase in the internal energy. At the same time rotating paddles also leads to a change in other independent variables and degrees of freedom of the system, such as angular momentum, density, pressure, which in turn yield an increase in the disorder in the system and hence an increase in the irreversible entropy production.

Appendix C: A Clausius inequality for the irreversible entropy production $d\mathcal{S}$

By using the identity $\text{Tr}[X] \text{Tr}[Y] = \text{Tr}[X \otimes Y]$, for any pair of linear operators X and Y , we can rewrite Eq. (19) of the main text as

$$d\mathcal{S} = \text{Tr}[(\mathbb{H} \otimes d\varrho^{(\text{ev})})\mathbb{O}], \quad (\text{C1})$$

where $\mathbb{O} = \sum_{i \geq 2} O_i \otimes O_i$. Note that \mathbb{H} is a positive operator and $\mathbb{H} \otimes d\varrho^{(\text{ev})}$ and both \mathbb{O} are traceless Hermitian operators. We also use the fact that any Hermitian operator X can be written as $X = X^{(+)} - X^{(-)}$ where both $X^{(\pm)}$ are positive operators, defined in the spectral form as $X^{(+)} = \sum_{k|x_k \geq 0} x_k |x_k\rangle\langle x_k|$ and $X^{(-)} = -\sum_{k|x_k < 0} x_k |x_k\rangle\langle x_k|$. If such an operator is also traceless, then we have $\text{Tr}[X^{(+)}] = \text{Tr}[X^{(-)}]$. As a result, it follows that

$$\begin{aligned} d\mathcal{S} &= \text{Tr}[\{(\mathbb{H} \otimes d\varrho^{(\text{ev})})^{(+)} - (\mathbb{H} \otimes d\varrho^{(\text{ev})})^{(-)}\}\mathbb{O}] \\ &= \sum_I \phi_I \langle \phi_I | (\mathbb{H} \otimes d\varrho^{(\text{ev})})^{(+)} | \phi_I \rangle - \sum_I \phi_I \langle \phi_I | (\mathbb{H} \otimes d\varrho^{(\text{ev})})^{(-)} | \phi_I \rangle \\ &\geq \phi_{\min} \sum_I \langle \phi_I | (\mathbb{H} \otimes d\varrho^{(\text{ev})})^{(+)} | \phi_I \rangle - \phi_{\max} \sum_I \langle \phi_I | (\mathbb{H} \otimes d\varrho^{(\text{ev})})^{(-)} | \phi_I \rangle \\ &= -(\phi_{\max} - \phi_{\min}) \text{Tr}[(\mathbb{H} \otimes d\varrho^{(\text{ev})})^{(+)}]. \end{aligned}$$

Thus,

$$d\mathcal{S} + b(\mathbb{O}) dB \geq 0, \quad (\text{C2})$$

where $b(\mathbb{O}) = |\phi_{\max} - \phi_{\min}|$ is spread of the operator \mathbb{O} , which somehow encompasses the information of all non-Hamiltonian system observables needed to describe the state of the system. This quantity is independent of ϱ . In addition, $dB = \text{Tr}[(\mathbb{H} \otimes d\varrho^{(\text{ev})})^{(+)}] = -(\sum_k \ln r_k)(\sum_{k'} dr_{k'}^{(+)}) \geq 0$, which fully depends on the state ϱ .

The above general inequality reminisces a *generalized Clausius inequality*—see Ref. [62] of the main text. We remark that all quantities appeared here are explicitly system or trajectory dependent, without need for explicit parameters from an environment

or heat bath. Of course when the trajectory is not unitary, existence of an environment is inevitable, hence dB can be considered to have contributions from the environment.

Appendix D: Alternative form for the irreversible entropy production

Consider the relative entropy

$$S(\varrho||\varrho_{\text{eq}}) = \text{Tr}[\varrho \ln \varrho - \varrho \ln \varrho_{\text{eq}}], \quad (\text{D1})$$

where ϱ_{eq} is a reference equilibrium state which is a Gibbs state with the temperature β of the system in the form $\varrho_{\text{eq}} = e^{-\beta H}/Z_{\text{eq}}$, where H is the physical Hamiltonian of the system and $Z_{\text{eq}} = \text{Tr}[e^{-\beta H}]$. Using $\varrho = \sum_k r_k |r_k\rangle\langle r_k|$, it follows that

$$\begin{aligned} S(\varrho||\varrho_{\text{eq}}) &= \sum_k r_k (\ln r_k - \langle r_k | \ln \varrho_{\text{eq}} | r_k \rangle) \\ &= \sum_k r_k \ln r_k + \beta \text{Tr}[\varrho H] + \ln Z_{\text{eq}}. \end{aligned} \quad (\text{D2})$$

Differentiation with respect to time yields

$$dS(\varrho||\varrho_{\text{eq}}) = \sum_k dr_k (\ln r_k - \langle r_k | \ln \varrho_{\text{eq}} | r_k \rangle) - \sum_k r_k (\langle dr_k | \ln \varrho_{\text{eq}} | r_k \rangle - \beta \langle r_k | dH | r_k \rangle - d\beta \langle r_k | H | r_k \rangle - dZ_{\text{eq}}/Z_{\text{eq}} + \langle r_k | \ln \varrho_{\text{eq}} | dr_k \rangle). \quad (\text{D3})$$

Using the relations

$$dS = -\sum_k dr_k \ln r_k, \quad (\text{D4})$$

$$d\mathcal{W}_{\text{cd}} = \sum_k r_k (\langle dr_k | H | r_k \rangle + \langle r_k | H | dr_k \rangle), \quad (\text{D5})$$

$$d\mathcal{Q} = \sum_k dr_k \langle r_k | H | r_k \rangle, \quad (\text{D6})$$

$$dZ_{\text{eq}}/Z_{\text{eq}} = -d\beta \text{Tr}[H \varrho_{\text{eq}}] - \beta \text{Tr}[dH \varrho_{\text{eq}}], \quad (\text{D7})$$

it follows that

$$d\mathcal{S} = -dS(\varrho||\varrho_{\text{eq}}) + \beta d\mathcal{W}_{\text{cd}} + \beta \text{Tr}[(\varrho - \varrho_{\text{eq}}) dH] + d\beta \text{Tr}[(\varrho - \varrho_{\text{eq}}) H], \quad (\text{D8})$$

where we have used the definition of irreversible entropy change $d\mathcal{S} = dS - \beta d\mathcal{Q}$.

However, under general conditions of a Markovian dynamics with a time-dependent steady state of ϱ_{eq} where both β and H can be time-dependent, from $dS(\varrho||\varrho_{\text{eq}}) \leq 0$ it can be concluded that

$$d\mathcal{S} \geq \beta d\mathcal{W}_{\text{cd}} - \beta \text{Tr}[(\varrho - \varrho_{\text{eq}}) dH] - d\beta \text{Tr}[(\varrho - \varrho_{\text{eq}}) H]. \quad (\text{D9})$$

Appendix E: Temperature in the irreversible entropy production and the Clausius inequality

The entropy production itself originates from the Clausius inequality. A widespread assumption in thermodynamics is that the temperature $T = 1/\beta$ in the Clausius inequality $\oint dQ/T \leq 0$, and hence the one in the irreversible entropy production ($d\mathcal{S} = dS - \beta d\mathcal{Q}$), is the temperature of the ambient bath. However, it can be argued that in general scenarios this assignment may be misleading. In fact, the discussion about the meaning of T in dQ/T even in standard classical thermodynamics is far from settled. For example, see the discussion in Ref. [3]. By revisiting the proof of the Clausius inequality (see, e.g., Ref. [2]) it becomes clear that the temperature that appears in this inequality is the temperature of an *auxiliary* environment which can provide the system with $d\mathcal{Q}$ (the heat needed for proceeding with the given thermodynamics process). Hence there is some freedom in this temperature as far as the auxiliary environment can provide the necessary heat for the process. Even if there is a real physical environment around the system, there is no need to put the temperature of this environment into the entropy production to have a valid Clausius inequality.

Given this freedom, the temperature in the Clausius inequality can instead be assigned to the system itself as its instantaneous (and generally nonequilibrium) temperature. Clausius himself had indicated in his 1856 paper: “ T is a function of the temperature of the changing body in that moment in which it takes in the element of heat dQ or, if the body has different temperatures in its different parts, of the temperature of that part of it which takes in dQ ” [4] (see also Ref. [5]). Along similar lines we also think that the correct way to describe the entropy production is when the temperature therein is the system temperature. This is in accordance with a system-based description of the thermodynamic properties and is in complete accordance with the laws of thermodynamics. As we have shown in the main text and also in Sec. A (of this Supplemental Material) the system temperature—in the way that we have derived—appears naturally when expanding entropy in terms of a set of relevant

independent variables. Thus, with our definition, entropy production has a reasonable and natural physical interpretation as the contribution of the other properties and physical variables of the system but heat (or energy) in entropy change of the system. In fact, if in general scenarios we put the temperature of the environment into the definition of the entropy production, the term dQ/T would not be a genuine part of the entropy change of the *system*. It seems more reasonable that entropy production should be a *system* property not depending on hypothetical situations by which a particular process can be realized.

Appendix F: Details of Example I

1. Case (i)

The initial state of the atom is a thermal state $\varrho(0) = e^{-\beta(0)\omega_0\sigma_z}/\text{Tr}[e^{-\beta(0)\omega_0\sigma_z}]$ with initial inverse temperature $\beta(0)$. In this case, the trajectory becomes

$$\varrho(t) = (1/2)\sum_{k=0,1} (1 + (-1)^{k+1}e^{-2\gamma t} \tanh(\beta(0)\omega_0)) |r_k\rangle\langle r_k|, \quad (\text{F1})$$

where $\{|r_0\rangle = |0\rangle, |r_1\rangle = |1\rangle\}$. Since $\varrho(t)$ is diagonal in a time-independent basis, its change is only due to the change in the eigenvalues and thus the whole internal energy change is due to the heat exchange with the bath, $\dot{Q}(t) = \dot{U}(t) = 2\gamma\omega_0 e^{-2\gamma t} \tanh[\beta(0)\omega_0]$. A direct calculation also shows that $\dot{q}(t) = \dot{Q}(t)$ and $\dot{w}(t) = \dot{W}(t)$.

2. Case (ii)

The initial state of the atom is a pure state $\varrho(0) = (1/2)(\mathbb{1} + \sigma_x)$, which is a steady state for the dissipative part of the dynamics $\mathcal{D}[\varrho(0)] = 0$. However, the time evolution of the state, i.e., $\varrho(t)$ is not an steady state of the dissipative part and is given by

$$\varrho(t) = (1/2)\mathbb{1} + c(t)|0\rangle\langle 1| + c^*(t)|1\rangle\langle 0|, \quad (\text{F2})$$

where

$$c(t) = e^{-\gamma t} (\Delta \cosh(t\Delta) + (\gamma - 2i\omega_0) \sinh(t\Delta)) / (2\Delta), \quad (\text{F3})$$

and $\Delta = \sqrt{\gamma^2 - 4\omega_0^2}$. From the expression for $\dot{U}(t)$ in the main text,

$$\dot{U}(t) = -2\gamma\omega_0 \text{Tr}[\sigma_z \varrho(t)], \quad (\text{F4})$$

it is evident that although the state of the system varies in time, the system internal energy does not change, $\dot{U}(t) = 0$.

By calculating the eigenvalues and the eigenvectors of $\varrho(t)$ as follows:

$$r_{\pm}(t) = \frac{1}{2} \pm \frac{e^{-\gamma t}}{2\Delta} \left(\gamma^2 \cosh(2t\Delta) + \gamma\Delta \sinh(2t\Delta) - 4\omega_0^2 \right)^{1/2}, \quad (\text{F5})$$

$$\sqrt{N_{\pm}(t)} |r_{\pm}(t)\rangle = \frac{(\gamma^2 \cosh(2t\Delta) + \gamma\Delta \sinh(2t\Delta) - 4\omega_0^2)^{1/2}}{\Delta \cosh(t\Delta) + (\gamma + 2i\omega_0) \sinh(t\Delta)} |0\rangle \pm |1\rangle, \quad (\text{F6})$$

where N_{\pm} is the normalization factor for $|r_{\pm}(t)\rangle$, it can be shown that $\dot{Q}(t) = \dot{W}(t) = \dot{q}(t) = \dot{w}(t) = 0$.

3. Case (iii)

Starting from $\varrho(0) = (\mathbb{1} + [\sigma_x + \sigma_z]/2)/2$ and obtaining the instantaneous state of the system, we can obtain the heat, work, and internal energy change in the system. Figure 1 of the main text illustrates the results. It can be shown that $\dot{q}(t) = \dot{U}(t) = -(\gamma/2)e^{-2\gamma t}$ and $\dot{w}(t) = 0$.

Appendix G: Example III: Damped harmonic oscillator within a bath of oscillators

The main manuscript discusses Examples I and II. We next consider a quantum harmonic oscillator interacting with a bath of oscillators with the total Hamiltonian

$$H_{\text{SB}} = \omega_0 \hat{a}^\dagger \hat{a} + \sum_{k=1}^M \omega_k \hat{b}_k^\dagger \hat{b}_k + \sum_{k=1}^M g_k (\hat{a}^\dagger \hat{b}_k + \hat{a} \hat{b}_k^\dagger), \quad (\text{G1})$$




FIG. 3. Numerical simulation for TB-STA heat exchange rate (green) and entropy rate (orange) vs. time, for a harmonic oscillator in a bath of harmonic oscillators as given in Example III, for $M = 255$, $g_k = \sqrt{\omega_k/\pi}e^{-\omega_k/(2\omega_c)}$, $\omega_k = 0.1k$, $\omega_0 = 1$, and $\omega_c = 5$.

where M is the number of the bath oscillators. For simplicity of the analysis, we assume the initial system-bath state $|1\rangle_S \otimes |0\rangle_B^{\otimes M}$, where $|i\rangle$ denotes the eigenstate of the corresponding number operator with eigenvalue i . Using the exact numerical simulation, we obtain the time derivatives of heat and entropy; see Fig. 3. Interestingly, we see that in this special case, $d\mathcal{S}$ vanishes. Hence the process is thermodynamically reversible. A closer look at the dynamics shows that the system evolves quasistatically and its state at any time is given by a Gibbs state $\varrho_S(t) = \varrho_{eq}(t) = e^{-\beta(t)H_S}/\text{Tr}[e^{-\beta(t)H_S}]$ and $\beta(t)$ is obtained from Eq. (18) of the main text. Thus, only the eigenvalues of the state change in time and the eigenvectors remain unchanged, indicating that dW_{cd} vanishes. Hence heat and work here are consistent with the conventional definitions.

-
- [1] S. Alipour, M. Afsary, F. Bakhshinezhad, M. Ramezani, F. Benatti, T. Ala-Nissila, and A. T. Rezakhani, Temperature in nonequilibrium quantum systems, [arXiv:2105.11915](#).
 - [2] C. B. P. Finn, *Thermal Physics* (CRC Press, Boca Raton, FL, 1993).
 - [3] D. J. Evans, D. J. Searles, and S. R. Williams, *Fundamentals of Classical Statistical Thermodynamics: Dissipation, Relaxation and Fluctuation Theorems* (Wiley-VCH, Weinheim, 2016), p. 2.
 - [4] M. M. Pradas and J. L. G. Ribelles, Poincaré's Proof of Clausius' Inequality, *Philosophia Scientiae* **1**, 135 (1996).
 - [5] R. Clausius, *The Mechanical Theory of Heat, with its Applications to the Steam-Engine and to the Physical Properties of the Bodies* (Taylor and Francis, London, 1864), pp. 141-142.