Exact solution of a anisotropic $J_1 - J_2$ spin chain with antiperiodic boundary condition

Yi Qiao^{a,b,c}, Jian Wang^{b,d}, Junpeng Cao^{b,d,e1} and Wen-Li Yang^{a,c,f2}

^a Institute of Modern Physics, Northwest University, Xian 710127, China

^b Beijing National Laboratory for Condensed Matter Physics, Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

^c Shaanxi Key Laboratory for Theoretical Physics Frontiers, Xian 710127, China

^d School of Physical Sciences, University of Chinese Academy of Sciences, Beijing, China

^e Songshan Lake Materials Laboratory, Dongguan, Guangdong 523808, China

^f School of Physics, Northwest University, Xian 710127, China

Abstract

The exact solution of an integrable anisotropic Heisenberg spin chain with nearestneighbour, next-nearest-neighbour and scalar chirality couplings is studied, where the boundary condition is the antiperiodic one. The detailed construction of Hamiltonian and the proof of integrability are given. The antiperiodic boundary condition breaks the U(1)-symmetry of the system and we use the off-diagonal Bethe Ansatz to solve it. The energy spectrum is characterized by the inhomogeneous T - Q relations and the contribution of the inhomogeneous term is studied. The ground state energy and the twisted boundary energy in different regions are obtained. We also find that the Bethe roots at the ground state form the string structure if the coupling constant J = -1although the Bethe Ansatz equations are the inhomogeneous ones.

PACS: 75.10.Pq, 03.65.Vf, 71.10.Pm Keywords: Quantum spin chain; Bethe Ansatz; Yang-Baxter equation.

 $^{^1\}mathrm{Corresponding}$ author: junpengcao@iphy.ac.cn

²Corresponding author: wlyang@nwu.edu.cn

1 Introduction

The Heisenberg model is a typical system to describe the quantum magnetism, where the spin exchanging interaction is the nearest-neighbor (NN) one. A nontrivial generalization of the Heisenberg model is the $J_1 - J_2$ model, where the NN and the next-nearest-neighbor (NNN) interactions are involved [1, 2, 3, 4, 5, 6, 7]. Many interesting phenomena have been found in the $J_1 - J_2$ model. For example, at the point of $J_2/J_1 = 0.241$, the $J_1 - J_2$ model has a topological phase transition [8, 9]. At the Majumdar-Ghosh point, $J_2/J_1 = 0.5$, the model Hamiltonian degenerates into a projector operator and only the ground state can be obtained exactly [10].

Although the $J_1 - J_2$ model can not be solved exactly, people find that the $J_1 - J_2$ model with some additional terms is integrable. For example, Popkov and Zvyagin proposed the integrable two-chain and multichain quantum spin model [11, 12, 13, 14]. Frahm and Rödenbeck constructed an integrable model of two coupled Heisenberg chains by taking the derivative of the logarithm of product of two transfer matrices with different spectral parameters [15, 16, 17]. Using the samilar idea, Ikhlef, Jacobsen and Saleur constructed the Z_2 staggered vertex model [18, 19]. These models are equivalent to the $J_1 - J_2$ model with some spin chirality terms, where the extra scalar chirality terms are introduced to ensure the integrability. Tavares and Ribeiro studied the thermodynamic properties of this kind of models by using the quantum transfer matrix method [20, 21]. Recently, the models with chirality terms have attracted lot of interest in the context of quantum spin liquids [22, 23].

The quantization condition used in the above references are the periodic boundary condition. In this paper, we study the integrable anisotropic $J_1 - J_2$ spin chain with antiperiodic boundary condition. We note that in this case, the U(1) symmetry of the system is broken and the traditional Bethe ansatz method does not work due to the lack of reference state. The antiperiodic (twisted) boundary condition is tightly related to the recent study on the topological states of matter. The model Hamiltonian considered in this paper is

$$H = J \sum_{j=1}^{2N} \left\{ \cosh(2a) (\sigma_{j}^{x} \sigma_{j+1}^{x} + \sigma_{j}^{y} \sigma_{j+1}^{y}) + \cosh \eta \sigma_{j}^{z} \sigma_{j+1}^{z} - \frac{\sinh^{2}(2a) \cosh \eta}{2 \sinh^{2} \eta} \vec{\sigma}_{j} \cdot \vec{\sigma}_{j+2} + \frac{(-1)^{j} i \sinh(2a)}{2 \sinh \eta} \left\{ \cosh \eta \vec{\sigma}_{j+1} \cdot (\vec{\sigma}_{j} \times \vec{\sigma}_{j+2}) + [\cosh(2a) - \cosh \eta] \sigma_{j+1}^{z} (\sigma_{j}^{x} \sigma_{j+2}^{y} - \sigma_{j}^{y} \sigma_{j+2}^{x}) \right\} \right\},$$
(1.1)

where $\vec{\sigma}_j \equiv (\sigma_j^x, \sigma_j^y, \sigma_j^z)$ are the Pauli matrices at site j, a and η are the generic constants describing the coupling strengths, and the boundary condition is the antiperiodic one

$$\sigma_{2N+n}^{\alpha} = \sigma_n^x \sigma_n^{\alpha} \sigma_n^x, \quad n = 1, 2, \quad \alpha = x, y, z.$$
(1.2)

In the Hamiltonian (1.1), the first two terms describe an anisotropic NN interaction, the third term is an isotropic NNN interaction and the last one corresponds to an anisotropic chiral three-spin interaction. We note that the hermitian of the Hamiltonian (1.1) requires that a must be real if η is imaginary (gapped regime), and a must be imaginary if η is real (gapless regime). We use the off-diagonal Bethe Ansatz (ODBA) [24, 25] to solve the model.

The paper is organized as follows. In the next section, we prove that the model (1.1) is integrable. In section 3, we derive the exact energy spectrum and the Bethe Ansatz equations. Ground state and twisted boundary energy with J = 1 are given in section 4 and the corresponding results with J = -1 are discussed in section 5. Section 6 is attributed to the concluding remarks.

2 Integrability

Throughout, V denotes a two-dimensional linear space and let $\{|m\rangle, m = 0, 1\}$ be an orthogonal basis of it. We shall adopt the standard notations: for any matrix $A \in \text{End}(V)$, A_j is an embedding operator in the tensor space $V \otimes V \otimes \cdots$, which acts as A on the *j*-th space and as identity on the other factor spaces. For $B \in \text{End}(V \otimes V)$, B_{ij} is an embedding operator of B in the tensor space, which acts as identity on the factor spaces except for the *i*-th and *j*-th ones.

Let us introduce the *R*-matrix $R_{0,j}(u) \in \text{End}(V_0 \otimes V_j)$

$$R_{0,j}(u) = \frac{1}{2} \left[\frac{\sinh(u+\eta)}{\sinh\eta} (1+\sigma_0^z \sigma_j^z) + \frac{\sinh u}{\sinh\eta} (1-\sigma_0^z \sigma_j^z) \right] + \frac{1}{2} (\sigma_0^x \sigma_j^x + \sigma_0^y \sigma_j^y) \\ = \frac{1}{\sinh\eta} \left(\begin{array}{ccc} \sinh(u+\eta) & 0 & 0 & 0 \\ 0 & \sinh u & \sinh\eta & 0 \\ 0 & \sinh\eta & \sinh u & 0 \\ 0 & 0 & 0 & \sinh(u+\eta) \end{array} \right),$$
(2.1)

where u is the spectral parameter. The *R*-matrix (2.1) satisfies the following relations

Initial condition :
$$R_{0,j}(0) = P_{0,j}$$
,
Unitary relation : $R_{0,j}(u)R_{j,0}(-u) = \phi(u) \times id$,
Crossing relation : $R_{0,j}(u) = -\sigma_0^y R_{0,j}^{t_0}(-u-\eta)\sigma_0^y$,
PT-symmetry : $R_{0,j}(u) = R_{j,0}(u) = R_{0,j}^{t_0 t_j}(u)$, (2.2)

where $\phi(u) = -\sinh(u+\eta)\sinh(u-\eta)/\sinh^2\eta$, t_0 (or t_j) denotes the transposition in the space V_0 (or V_j) and $P_{0,j}$ is the permutation operator possessing the property

$$R_{j,k}(u) = P_{0,j}R_{0,k}(u)P_{0,j}.$$
(2.3)

The *R*-matrix (2.1) satisfies the Yang-Baxter equation (YBE)

$$R_{1,2}(u_1 - u_2)R_{1,3}(u_1 - u_3)R_{2,3}(u_2 - u_3)$$

= $R_{2,3}(u_2 - u_3)R_{1,3}(u_1 - u_3)R_{1,2}(u_1 - u_2).$ (2.4)

We define the monodromy matrices as

$$T_{0}(u) = \sigma_{0}^{x} R_{0,1}(u+a) R_{0,2}(u-a) \cdots R_{0,2N-1}(u+a) R_{0,2N}(u-a),$$

$$\hat{T}_{0}(u) = \sigma_{0}^{x} R_{0,2N}(u+a) R_{0,2N-1}(u-a) \cdots R_{0,2}(u+a) R_{0,1}(u-a),$$
 (2.5)

where V_0 is the auxiliary space, $V_1 \otimes V_2 \otimes \cdots \otimes V_{2N}$ is the physical or quantum space, 2N is the number of sites and a is the inhomogeneous parameter. From the YBE (2.4) and the fact

$$[R_{1,2}(u), \sigma_1^x \sigma_2^x] = 0, (2.6)$$

one can prove that the monodromy matrix T(u) satisfies the Yang-Baxter relation

$$R_{1,2}(u-v)T_1(u)T_2(v) = T_2(v)T_1(u)R_{1,2}(u-v).$$
(2.7)

The transfer matrices are the trace of monodromy matrices in the auxiliary space

$$t(u) = tr_0 T_0(u), \quad \hat{t}(u) = tr_0 \hat{T}_0(u).$$
 (2.8)

Using the crossing symmetry in Eq.(2.2), we obtain the relations between transfer matrices t(u) and $\hat{t}(u)$

$$t(u) = -\hat{t}(-u - \eta), \quad \hat{t}(u) = -t(-u - \eta).$$
(2.9)

From the Yang-Baxter relation (2.7) and Eq.(2.9), one can prove that the transfer matrices t(u) [or $\hat{t}(u)$] with different spectral parameters commute with each other. Meanwhile, the transfer matrices t(u) and $\hat{t}(u)$ also commute with each other

$$[t(u), t(v)] = [\hat{t}(u), \hat{t}(v)] = [t(u), \hat{t}(u)] = 0.$$
(2.10)

Therefore, both t(u) and $\hat{t}(u)$ serve as the generating functions of all the conserved quantities of the system, and the transfer matrices t(u) and $\hat{t}(u)$ can be diagonalized simultaneously.

Using the initial condition of the R-matrix (2.2), we obtain

$$\hat{t}(-a) = R_{2N,2N-1}(-2a)\cdots R_{2N,2}(0)R_{2N,1}(-2a)\sigma_{2N}^{x},$$
$$\hat{t}(a) = \sigma_{1}^{x}R_{1,2N}(2a)R_{1,2N-1}(0)\cdots R_{1,2}(2a).$$
(2.11)

Taking the derivative of transfer matrix t(u) with respect to u and consider the values at the point of u = a, we have

$$\frac{\partial t(u)}{\partial u}\Big|_{u=a} = \sum_{j=1}^{N-1} \sigma_{2N}^{x} R_{2N,1}(2a) R_{2N,2}(0) [R'_{2N,2j-1}(2a) R_{2N,2j}(0) + R_{2N,2j-1}(2a) \\ \times R'_{2N,2j}(0)] \cdots R_{2N,2N-1}(2a) + \sigma_{2N}^{x} R_{2N,1}(2a) R_{2N,2}(0) \cdots R'_{2N,2N-1}(2a) \\ \times + R_{2,3}(2a) R_{2,4}(0) \cdots R_{2,2N-1}(2a) R'_{2,2N}(0) \sigma_{2}^{x} R_{2,1}(2a),$$
(2.12)

where $R'_{i,j}(u) = \frac{\partial}{\partial u} R_{i,j}(u)$. Similarly the derivative of t(u) at the point of u = -a is

$$\frac{\partial t(u)}{\partial u}\Big|_{u=-a} = \sum_{j=1}^{N} R_{1,2}(-2a) \cdots [R'_{1,2j-1}(0)R_{1,2j}(-2a) + R_{1,2j-1}(0)R'_{1,2j}(-2a)] \cdots$$

$$\times R_{1,2N-1}(0)R_{1,2N}(-2a)\sigma_2^x + R'_{1,2}(-2a)R_{1,3}(0) \cdots R_{1,2N}(-2a)\sigma_1^x$$

$$\times + R_{2N-1,2N}(-2a)\sigma_{2N-1}^x R_{2N-1,1}(0)R_{2N-1,2}(-2a) \cdots R_{2N-1,2N-2}(-2a). \quad (2.13)$$

The integrable Hamiltonian can be constructed from the transfer matrices t(u) and $\hat{t}(u)$ as

$$H = \frac{JN\cosh\eta[\cosh^2(2a) - \cosh(2\eta)]}{\sinh^2\eta} + J\phi^{1-N}(2a)\sinh\eta$$
$$\times J\left\{\hat{t}(-a)\frac{\partial t(u)}{\partial u}\Big|_{u=a} + \hat{t}(a)\frac{\partial t(u)}{\partial u}\Big|_{u=-a}\right\}.$$
(2.14)

Substituting the relations (2.11)-(2.13) into above expression (2.14), we obtain

$$H = \sinh \eta \left\{ \sum_{j=1}^{N-1} \left[R_{2j,2j-1}(-2a) R'_{2j,2j-1}(2a) + R_{2j+1,2j}(2a) R'_{2j+1,2j}(-2a) + R_{2j+2,2j+1}(-2a) P_{2j+2,2j} R'_{2j+2,2j}(0) R_{2j+2,2j+1}(2a) + R_{2j+1,2j}(2a) + R_{2j+1,2j-1} R'_{2j+1,2j-1}(0) R_{2j+1,2j}(-2a) \right] + R_{2N,2N-1}(-2a) R'_{2N,2N-1}(2a) + \sigma_1^x R_{1,2N}(2a) R'_{2j+1,2j}(-2a) \sigma_1^x + R_{2,1}(-2a) \sigma_2^x P_{2,2N} R'_{2,2N}(0) \sigma_2^x R_{2,1}(2a) + \sigma_1^x R_{1,2N}(2a) + \sigma_1^x R_{1,2N}(2a) + S_{1,2N-1} R'_{1,2N-1}(0) R_{1,2N}(-2a) \sigma_1^x + \frac{N \cosh \eta [\cosh^2(2a) - \cosh(2\eta)]}{\sinh^2 \eta}.$$
(2.15)

The derivative of the R-matrix reads

$$R'_{0,j}(u) = \frac{1}{2} \left[\frac{\cosh(u+\eta)}{\sinh\eta} (1 + \sigma_0^z \sigma_j^z) + \frac{\cosh u}{\sinh\eta} (1 - \sigma_0^z \sigma_j^z) \right].$$
 (2.16)

The commutative relation between the permutation operators is

$$[P_{2,1}, P_{2,0}] = \frac{1}{4} [(1 + \vec{\sigma}_2 \cdot \vec{\sigma}_1), (1 + \vec{\sigma}_2 \cdot \vec{\sigma}_0)] \\ = \frac{i}{2} \vec{\sigma}_2 \cdot (\vec{\sigma}_1 \times \vec{\sigma}_0).$$
(2.17)

Substituting Eqs.(2.16) and (2.17) into (2.15) and after some tedious calculations, we arrive at the Hamiltonian (1.1). From the construction (2.14) and the commutation relation (2.10) of generating functions t(u) and $\hat{t}(u)$, we conclude that the model (1.1) with the antiperiodic boundary condition is integrable.

3 Exact solution

We first introduce the inhomogeneous monodromy matrix

$$T_0^g(u) = \sigma_0^x R_{0,1}(u - \theta_1) R_{0,2}(u - \theta_2) \cdots R_{0,2N-1}(u - \theta_{2N-1}) R_{0,2N}(u - \theta_{2N}),$$
(3.1)

where the $\{\theta_j, j = 1, \dots, N\}$ are the inhomogeneous parameters. The matrix form of monodromy matrix $T_0^g(u)$ in the auxiliary space is

$$T_0^g(u) = \begin{pmatrix} C(u) & D(u) \\ A(u) & B(u) \end{pmatrix}.$$
(3.2)

where A(u), B(u), C(u) and D(u) are the operators acting in the quantum space. We denote the all spins aligning up state as the vacuum state $|0\rangle$,

$$|0\rangle = \begin{pmatrix} 1\\0 \end{pmatrix}_1 \otimes \dots \otimes \begin{pmatrix} 1\\0 \end{pmatrix}_{2N}.$$
(3.3)

The elements of the monodromy matrix $T_0^g(u)$ acting on the vacuum state gives

$$A(u)|0\rangle = \tilde{a}(u)|0\rangle, \quad B(u)|0\rangle \neq 0,$$

$$C(u)|0\rangle = 0, \qquad D(u)|0\rangle = \tilde{d}(u)|0\rangle, \qquad (3.4)$$

where

$$\tilde{a}(u) = \prod_{j=1}^{2N} \frac{\sinh(u-\theta_j+\eta)}{\sinh\eta}, \quad \tilde{d}(u) = \prod_{j=1}^{2N} \frac{\sinh(u-\theta_j)}{\sinh\eta}.$$

The transfer matrix t(u) defined as

$$t^{g}(u) = tr_{0}T_{0}^{g}(u) = B(u) + C(u).$$
(3.5)

Suppose $|\Phi\rangle$ is the eigenstate of the transfer matrix $t^g(u)$ and the corresponding eigenvalue is $\Lambda^g(u)$,

$$t^{g}(u)|\Phi\rangle = \Lambda^{g}(u)|\Phi\rangle. \tag{3.6}$$

According to the results given in [25], we known that $\Lambda^{g}(u)$ satisfies following functional relations

$$\Lambda^{g}(\theta_{j})\Lambda^{g}(\theta_{j}-\eta) = -\tilde{a}(\theta_{j})\tilde{d}(\theta_{j}-\eta), \quad j=1,\cdots,N.$$
(3.7)

Meanwhile, $\Lambda^{g}(u)$ is a polynomial of e^{u} with the degree 2N - 1 and satisfies the periodicity property

$$\Lambda^{g}(u+i\pi) = (-1)^{2N-1} \Lambda^{g}(u).$$
(3.8)

The constraints (3.7) and (3.8) show that the eigenvalue $\Lambda^{g}(u)$ can be parameterized as the following inhomogeneous T - Q relation

$$\Lambda^{g}(u) = e^{u}\tilde{a}(u)\frac{Q(u-\eta)}{Q(u)} - e^{-u-\eta}\tilde{d}(u)\frac{Q(u+\eta)}{Q(u)} - \frac{\tilde{c}(u)\tilde{a}(u)\tilde{d}(u)}{Q(u)},$$
(3.9)

where Q(u) is a trigonometric polynomial of the type

$$Q(u) = \prod_{j=1}^{2N} \frac{\sinh(u - \lambda_j)}{\sinh\eta},$$
(3.10)

and $\tilde{c}(u)$ is given by

$$\tilde{c}(u) = e^{u - 2N\eta + \sum_{l=1}^{2N} (\theta_l - \lambda_l)} - e^{-u - \eta - \sum_{l=1}^{2N} (\theta_l - \lambda_l)}.$$
(3.11)

The singularity of $\Lambda^{g}(u)$ requires that the Bethe roots $\{\lambda_{j}\}$ in Eq.(3.9) should satisfy the Bethe ansatz equations (BAEs)

$$e^{\lambda_j}\tilde{a}(\lambda_j)Q(\lambda_j - \eta) - e^{-\lambda_j - \eta}\tilde{d}(\lambda_j)Q(\lambda_j + \eta) - \tilde{c}(\lambda_j)\tilde{a}(\lambda_j)\tilde{d}(\lambda_j) = 0,$$

$$j = 1, \cdots, N.$$
(3.12)

Put $\theta_{2j-1} = -a$ and $\theta_{2j} = a$ for $j = 1, \dots, N$, we obtain the eigenvalue $\Lambda(u)$ of the transfer matrix t(u)

$$\Lambda(u) = e^{u}a(u)\frac{Q(u-\eta)}{Q(u)} - e^{-u-\eta}d(u)\frac{Q(u+\eta)}{Q(u)} - \frac{c(u)a(u)d(u)}{Q(u)},$$
(3.13)

where

$$a(u) = \frac{\sinh^{N}(u+a+\eta)\sinh^{N}(u-a+\eta)}{\sinh^{2N}\eta},$$

$$d(u) = \frac{\sinh^{N}(u+a)\sinh^{N}(u-a)}{\sinh^{2N}\eta},$$

$$c(u) = e^{u-2N\eta - \sum_{l=1}^{2N}\lambda_{l}} - e^{-u-\eta + \sum_{l=1}^{2N}\lambda_{l}},$$

(3.14)

and the BAEs read

$$e^{\lambda_j}a(\lambda_j)Q(\lambda_j - \eta) - e^{-\lambda_j - \eta}d(\lambda_j)Q(\lambda_j + \eta) - c(\lambda_j)a(\lambda_j)d(\lambda_j) = 0,$$

$$j = 1, \cdots, N.$$
(3.15)

From Eqs.(2.9), (2.14) and (3.13), the energy spectrum of the Hamiltonian (1.1) is

$$E = \sum_{j=1}^{2N} J \sinh \eta \,\phi(2a) \bigg\{ \coth(\lambda_j - a) - \coth(\lambda_j - a + \eta) + \coth(\lambda_j + a) \\ - \coth(\lambda_j + a + \eta) \bigg\} + 2J \sinh \eta \,\phi(2a) - \frac{JN \cosh \eta [\cosh^2(2a) - \cosh(2\eta)]}{\sinh^2 \eta}.$$
(3.16)

Next, we check above results numerically. Numerical solutions of the BAEs and exact diagonalization of the Hamiltonian (1.1) are performed for the case of 2N = 4 with randomly choosing of model parameters. The results are listed in Table 1. We see that the eigenvalues obtained by solving the BAEs are exactly the same as those obtained by the exact diagonalization of the Hamiltonian (1.1). Meanwhile, the expression (3.16) gives the complete spectrum of the system.

Table 1: Numerical solutions of the BAEs (3.15) for real η case, where J = 1, $\eta = 1$, b = 1, 2N = 4, n indicates the number of the energy levels and E_n is the corresponding energy. The energy E_n calculated from the Bethe roots is exactly the same as that from the exact diagonalization of the Hamiltonian (1.1).

λ_1	λ_2	λ_3	λ_4	E_n	n
-1.0873 - 1.5708i	-0.5000 - 0.4634i	-0.5000 + 0.4634i	0.0873 - 1.5708i	-5.8897	1
-1.7292 - 1.5708i	-0.5000 - 1.2559i	-0.5000 + 1.2559i	0.7292 - 1.5708i	-5.8897	1
-1.4553 + 1.0461i	-0.5000 - 1.1872i	-0.5000 + 0.8274i	0.4553 + 1.0461i	-3.9899	2
-1.4553 - 1.0461i	-0.5000 - 0.8274i	-0.5000 + 1.1872i	0.4553 - 1.0461i	-3.9899	2
-1.7487 + 1.3618i	-0.5000 + 0.7332i	-0.5000 + 1.4005i	0.7487 + 1.3618i	-3.9899	2
-1.7487 - 1.3618i	-0.5000 - 1.4005i	-0.5000 - 0.7332i	0.7487 - 1.3618i	-3.9899	2
-1.6037 + 1.2858i	-0.5000 - 0.4949i	-0.5000 + 1.1518i	0.6037 + 1.2858i	-3.0796	3
-1.6037 - 1.2858i	-0.5000 - 1.1518i	-0.5000 + 0.4949i	0.6037 - 1.2858i	-3.0796	3
-1.7809 + 0.6635i	-0.9098 + 0.3143i	-0.0902 + 0.3143i	0.7809 + 0.6635i	3.9899	4
-1.7809 - 0.6635i	-0.9098 - 0.3143i	-0.0902 - 0.3143i	0.7809 - 0.6635i	3.9899	4
-2.0629 - 1.2897i	-0.9988 - 1.1945i	-0.0012 - 1.1945i	1.0629 - 1.2897i	3.9899	4
-2.0629 + 1.2897i	-0.9988 + 1.1945i	-0.0012 + 1.1945i	1.0629 + 1.2897i	3.9899	4
-1.4107 + 0.0000i	-1.2598 - 0.0000i	0.2598 + 0.0000i	0.4107 - 0.0000i	4.2251	5
-2.0830 - 1.5708i	-1.0046 - 1.5708i	0.0046 - 1.5708i	1.0830 - 1.5708i	4.2251	5
-2.0016 - 0.9831i	-1.0025 - 0.8073i	0.0025 - 0.8073i	1.0016 - 0.9831i	4.7442	6
-2.0016 + 0.9831i	-1.0025 + 0.8073i	0.0025 + 0.8073i	1.0016 + 0.9831i	4.7442	6

4 Ground state and twisted boundary energy

In this section, we consider the case of J = 1. We first analyze the contribution of the third term in the inhomogeneous T - Q relation (3.13). For simplicity, we constrain that η is real and a is imaginary. We first introduce following homogeneous T - Q relation

$$\Lambda_h(u) = e^u a(u) \frac{Q_h(u-\eta)}{Q_h(u)} - e^{-u-\eta} d(u) \frac{Q_h(u+\eta)}{Q_h(u)},$$
(4.1)

where $Q_h(u)$ is a trigonometric polynomial of the type

$$Q_h(u) = \prod_{j=1}^M \frac{\sinh(u - \lambda_j)}{\sinh\eta}, \quad M = 1, \cdots, 2N.$$
(4.2)

The period of Bethe roots $\{\lambda_j\}$ is 2π , thus we fix the real part of Bethe roots in the interval $[-\pi, \pi)$. For convenience, we put $\lambda_j = iu_j/2 - \eta/2$ and a = ib. The singularity of Eq.(4.1) gives

$$e^{iu_j} \left[\frac{\sin\frac{1}{2}(u_j - 2b - i\eta)\sin\frac{1}{2}(u_j + 2b - i\eta)}{\sin\frac{1}{2}(u_j - 2b + i\eta)\sin\frac{1}{2}(u_j + 2b + i\eta)} \right]^N = \prod_{l=1}^M \frac{\sin\frac{1}{2}(u_j - u_l - 2\eta i)}{\sin\frac{1}{2}(u_j - u_l + 2\eta i)},$$

$$j = 1, \cdots, M.$$
 (4.3)

Define

$$E_{h} = \frac{JN\cosh\eta[\cosh^{2}(2a) - \cosh(2\eta)]}{\sinh^{2}\eta} - J\phi^{1-N}(2a)\sinh\eta$$
$$\times J\left\{\Lambda_{h}(a-\eta)\frac{\partial\Lambda_{h}(u)}{\partial u}\Big|_{u=a} + \Lambda_{h}(-a-\eta)\frac{\partial\Lambda_{h}(u)}{\partial u}\Big|_{u=-a}\right\}$$
$$= -\sum_{j=1}^{M} 4\pi\sinh\eta\phi(2a)[a_{1}(u_{j}+2b) + a_{1}(u_{j}-2b)] + 2\sinh\eta\phi(2a)$$
$$-\frac{N\cosh\eta[\cosh^{2}(2a) - \cosh(2\eta)]}{\sinh^{2}\eta}, \qquad (4.4)$$

where the function $a_n(x)$ is given by

$$a_n(x) = \frac{1}{2\pi} \frac{\sinh(n\eta)}{\cosh(n\eta) - \cos x},\tag{4.5}$$

and the Bethe roots $\{u_j\}$ satisfy the BAEs (4.3). Taking the logarithm of Eq.(4.3), we have

$$u_j + N[\theta_1(u_j + 2b) + \theta_1(u_j - 2b)] = 2\pi I_j + \sum_{k=1}^M \theta_2(u_j - u_k),$$

$$j = 1, \dots, M,$$
(4.6)

where the quantum number $\{I_j\}$ are certain integers (or half odd integers) if M is even (or odd),

$$\theta_n(x) = 2 \arctan \frac{\tan(x/2)}{\tan(n\eta/2)} + 2\pi \lfloor \frac{x}{2\pi} + \frac{1}{2} \rfloor,$$

and $\lfloor \rfloor$ is the Gauss mark.

From the analysis of Eq.(4.6) and numerical calculations, we know that the energy E_h arrives at its minimum E_h^g when M = N. Meanwhile, all the Bethe roots $\{u_j\}$ are real and the corresponding quantum numbers are

$$I_j = -\frac{N}{2} + 1, -\frac{N}{2} + 2, \cdots, \frac{N}{2}.$$
(4.7)

From Eq.(4.7), we see that there is a hole in the real axis which can be put at the boundary $x_0 = -\pi$ to minimize the energy.

Denote the true ground state energy of Hamiltonian (1.1) as E^{g} . We use the physical quantity

$$E_{inh}^g = E_h^g - E^g, (4.8)$$

to characterize the contribution of the inhomogeneous term in the T - Q relation (3.13) at the ground state. By using the density matrix renormalization group (DMRG) method, we obtain the ground state energy E^g of the Hamiltonian (1.1). By solving the BAEs (4.6) and substituting the values of Bethe roots into the Eq.(4.4), we obtain the energy E_h^g . Substituting E^g and E_h^g into Eq.(4.8), we obtain the values of E_{inh}^g and the results are shown as the dots in Fig.1. The fitting of the data gives that the contribution of the inhomogeneous term tends to zero when the system-size tends to infinity. Then, we conclude that the Eq.(4.4) is a suitable approximation of the energy of the system (1.1) in the thermodynamic limit. In the following, we use E_h^g to quantity the ground state energy of the model (1.1).




Figure 1: The contribution of the inhomogeneous term to the ground state energy with $\eta = 2$ and (a) a = 0.3; (b) a = 0.75. The data can be fitted as (a) $E_{inh}^g = 10.96N^{-1.673} - 0.0021$; (b) $E_{inh}^g = 6.237N^{-1.738} - 0.0043$. When the system-size tends to infinity, the contribution of the inhomogeneous term can be neglected.

Define

$$Z(u) = \frac{1}{4\pi} \left[\frac{u}{N} + \theta_1(u+2b) + \theta_1(u-2b) - \frac{1}{N} \sum_{k=1}^M \theta_2(u-u_k) \right].$$
 (4.9)

Obviously, $Z(u_j) = I_j/2N$ corresponds to the Eq.(4.6). In the thermodynamic limit, $N \to \infty$, $M \to \infty$ and N/M finite, the counting function Z(u) becomes a continue function of u. Taking the derivative of Eq.(4.9) with respect to u, we obtain

$$\frac{dZ(u)}{du} = \frac{1}{4\pi N} + \frac{1}{2} \left[a_1(u+2b) + a_1(u-2b) \right] - \int_{-\pi}^{\pi} a_2(u-\lambda)\rho(\lambda)d\lambda$$
$$\equiv \rho(u) + \rho^h(u), \tag{4.10}$$

where $\rho(x)$ and $\rho^h(x)$ are the densities of particles and holes, respectively. The Fourier transformation of $a_n(x)$ is

$$\tilde{a}_n(\omega) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i\omega x} \frac{\sinh(n\eta)}{\cosh(n\eta) - \cos x} dx = e^{-n\eta|\omega|}.$$
(4.11)

At the ground state, there is a hole at the point of $x_0 = -\pi$. Thus the density of holes reads

$$\rho_g^h(x) = \frac{\delta(x - x_0)}{2N},$$
(4.12)

and the corresponding Fourier transformation is

$$\tilde{\rho}_g^h(w) = \frac{e^{-i\omega x_0}}{2N}.\tag{4.13}$$

Taking the Fourier transformation of Eq.(4.10), we obtain

$$\tilde{\rho}_g(\omega) = -\frac{e^{i\omega\pi}}{2N(1+e^{-2\eta|\omega|})} + \frac{\delta_{\omega,0}}{2N(1+e^{-2\eta|\omega|})} + \frac{\cos(2b\,\omega)}{2\cosh(\eta\,\omega)}.\tag{4.14}$$

The ground state energy of model (1.1) is

$$E^{g} = -4N \sinh \eta \,\phi(2bi) \sum_{\omega=-\infty}^{\infty} \frac{e^{-\eta|\omega|} \cos^{2}(2b\omega)}{\cosh(\eta\omega)} - \frac{N \cosh \eta [\cos^{2}(2b) - \cosh(2\eta)]}{\sinh^{2} \eta} + 2 \sinh \eta \,\phi(2bi) \sum_{\omega=-\infty}^{\infty} \frac{e^{i\pi\omega} \cos(2b\omega)}{\cosh(\eta\omega)}.$$

$$(4.15)$$

The twisted boundary energy is defined as

$$E_b = E^g - E_p^g, (4.16)$$

where E_p^g is the ground state energy for the Hamiltonian (1.1) with periodic boundary condition [26]

$$E_p^g = -4N\sinh\eta\,\phi(2bi)\sum_{\omega=-\infty}^{\infty}\frac{e^{-\eta|\omega|}\cos^2(2b\omega)}{\cosh(\eta\omega)} - \frac{N\cosh\eta[\cos^2(2b) - \cosh(2\eta)]}{\sinh^2\eta}.$$
 (4.17)

Substitute Eqs.(4.15) and (4.17) into (4.16), we obtain the value of twisted boundary energy as

$$E_b = 2\sinh\eta\,\phi(2bi)\sum_{\omega=-\infty}^{\infty}\frac{e^{i\pi\omega}\cos(2b\omega)}{\cosh(\eta\omega)}.$$
(4.18)

5 Bethe roots of inhomogeneous BAEs

In this section, we consider the case J = -1. In this case, the ground state spin configuration and the solution of Bethe roots in BAEs (3.15) are different from those with J = 1. Again, η is set as real and a is set as imaginary. For convenience, put $\lambda_j = iu_j - \eta/2$. The BAEs (3.15) become

$$\frac{e^{iu_j}\prod_{l=1}^{2N}\sin(u_j - u_l + i\eta)}{\sin^N(u_j + b + \frac{1}{2}i\eta)\sin^N(u_j - b + \frac{1}{2}i\eta)} = \frac{e^{-iu_j}\prod_{l=1}^{2N}\sin(u_j - u_l - i\eta)}{\sin^N(u_j + b - \frac{1}{2}i\eta)\sin^N(u_j - b - \frac{1}{2}i\eta)} + 2i\,e^{-N\eta}\sin\left(u_j - \sum_{l=1}^{2N}u_l\right), \qquad j = 1, \cdots, 2N.$$
(5.1)

By careful analysis and numerical check, we find the Bethe roots in Eq.(5.1) form the 2Nstrings at the ground state

$$u_j = x_0 + \left(\frac{N+1}{2} - j\right)i\eta + o(2N), \quad j = 1, \cdots, 2N,$$
(5.2)

where x_0 is real and o(2N) stands for a small correction which is related with 2N and i is the imaginary unit. The numerical results of Bethe roots at the ground state with 2N = 8 is shown in Fig.2. From which, the string structure of Bethe roots can been seen very clearly. Substituting the string hypothesis (5.2) into the energy expression (3.16) and neglecting the small correction, we obtain the energy for the 2N-string states

$$E_s = -2\sinh\eta\,\phi(2a) \left[\frac{\sinh(2N\eta)}{\cos(2x_0 + 2b) - \cosh(2N\eta)} + \frac{\sinh(2N\eta)}{\cos(2x_0 - 2b) - \cosh(2N\eta)} \right]$$
$$-2\sinh\eta\,\phi(2a) + \frac{N\cosh\eta[\cosh^2(2a) - \cosh(2\eta)]}{\sinh^2\eta}.$$
(5.3)




Figure 2: The distributions of Bethe roots at the ground state, where $\eta = 1, a = 0.3i$ and 2N = 8. The string structure can be seen very clearly from the data.

When system size 2N is very large, the energy for the 2N-string (5.3) turns to

$$E_s = 4\sinh\eta\,\phi(2a)\tanh(2N\eta) - 2\sinh\eta\,\phi(2a) + \frac{N\cosh\eta[\cosh^2(2a) - \cosh(2\eta)]}{\sinh^2\eta}.$$
 (5.4)

We see that the 2N-string energy (5.4) is irrelevant with the position x_0 of string.

In order to show the correction of Eq.(5.4), we calculate the ground state energy of the system (1.1) by exactly diagonalizing the Hamiltonian up to 2N = 18 and compare the results with those obtained from Eq.(5.4) in Fig.3. From it, we see that the energy difference ΔE , which comes from the finite-size effect of the string (5.2), exponentially tends to zero with the increasing 2N. The data satisfy the scaling law, $\Delta E \propto e^{\alpha 2N}$, $\alpha < 0$. Thus in the thermodynamic limit, the expression (5.4) gives the exact value of the ground state energy.

Now, we calculate the twisted boundary energy. The ground state for the Hamiltonian (1.1) with periodic boundary condition is the ferromagnetic state if J = -1. It is easy to obtain the ground state energy as

$$E^{p} = -2N\cosh\eta + \frac{N\cosh\eta\sinh^{2}(2a)}{\sinh^{2}\eta}.$$
(5.5)

From the definition of twisted boundary energy

$$E_t = E_s - E^p, (5.6)$$

and substituting Eqs. (5.4) and (5.5) into (5.6), we obtain

$$E_t = 4 \phi(2a) \sinh \eta \, \tanh(2N\eta) - 2 \sinh \eta \, \phi(2a). \tag{5.7}$$




Figure 3: The energy difference $\Delta E = E_s - E^{ED}$, where E_s is the ground state energy calculated by the string solution (5.4) and E^{ED} is the ground state energy calculated by the exact diagonalization. The ΔE is exponentially decreasing with the increasing system-size 2N and the data can be fitted as (a) $\Delta E = 46.9e^{-1.068*(2N)}$ (b) $\Delta E = 53.7e^{-1.32*(2N)}$. Thus, the energy difference ΔE will be zero in the thermodynamic limit.

In the thermodynamic limit, the twisted boundary energy arrives at

$$E_t = 2\sinh\eta\,\phi(2a).\tag{5.8}$$

6 Conclusion

In this paper, we study an integrable anisotropic $J_1 - J_2$ spin chain with antiperiodic boundary condition. By means of the off-diagonal Bethe Ansatz, we obtain the exact solution of the system. We show that the contribution of inhomogeneous term in the T - Q relation can be neglected when the system-size tends to infinity. Based on it, we discuss the ground state energy and the twisted boundary energy. We find the string structure of Bethe roots at the ground state if the coupling constant J = -1 although the corresponding Bethe Ansatz equations are the inhomogeneous ones.

Acknowledgments

We would like to thank Prof. Y. Wang for his valuable discussions and continuous encouragements. The financial supports from the National Program for Basic Research of MOST (Grant Nos. 2016YFA0300600 and 2016YFA0302104), the National Natural Science Foundation of China (Grant Nos. 11934015, 11434013, 11425522, 11547045, 11774397, 11775178 and 11775177), the Major Basic Research Program of Natural Science of Shaanxi Province (Grant Nos. 2017KCT-12, 2017ZDJC-32), Australian Research Council (Grant No. DP 190101529), the Strategic Priority Research Program of the Chinese Academy of Sciences and the Double First-Class University Construction Project of Northwest University are gratefully acknowledged.

References

- C. Zeng and J.B. Parkinson, Spatial periodicity of the spin-1/2 heisenberg antiferromagnetic chain with competing interactions, Phys. Rev. B,51 (17):11609, 1995.
- [2] S.R. White and I. Affleck, Dimerization and incommensurate spiral spin correlations in the zigzag spin chain: Analogies to the kondo lattice, Phys. Rev. B, 54:9862-9869, 1996.
- S. Eggert, Numerical evidence for multiplicative logarithmic corrections from marginal operators, Phys. Rev. B, 54:R9612-R9615, 1996.
- [4] B.S. Shastry and B. Sutherland, Excitation spectrum of a dimerized next-neighbor antiferromagnetic chain, Phys. Rev. Lett., 47:964-967, 1981.
- [5] K. Okamoto and K. Nomura, Fluid-dimer critical point in s = 1/2 antiferromagnetic heisenberg chain with next nearest neighbor interactions, Phys. Lett. A, 169(6):433-437, 1992.
- [6] R. Jafari and A. Langari, Second order quantum renormalisation group of xxz chain with next-nearest neighbour interactions, Physica A, 364:213-222, 2006.
- [7] Z.I. Djoufack, E. Tala-Tebue, J.P. Nguenang, and A. Kenfack-Jiotsa, Quantum soliton in 1d heisenberg spin chains with dzyaloshinsky-moriya and next-nearest-neighbor interactions, Chaos, 26(10):103110, 2016.
- [8] R. Bursill, G.A. Gehring, D.J.J Farnell, J.B. Parkinson, T. Xiang, and C. Zeng, Numerical and approximate analytical results for the frustrated spin-1/2 quantum spin chain, J. Phys.: Condens. Mat., 7(45):8605, 1995.
- [9] R. Jafari and A. Langari, Phase diagram of the one-dimensional s=1/2 xxz model with ferromagnetic nearest-neighbor and antiferromagnetic next-nearest-neighbor interactions, Phys. Rev. B, 76:014412, 2007.

- [10] C.K. Majumdar and D.K. Ghosh, On next-nearest-neighbor interaction in linear chain I, J. Math. Phys., 10(8):1388-1398, 1969.
- [11] V.Y. Popkov and A.A. Zvyagin, "antichiral" exactly solvable effectively two-dimensional quantum spin model, Phys. Lett. A, 175(5):295-298, 1993.
- [12] A.A. Zvyagin, Thermodynamics of the exactly solvable two-chain and multichain quantum spin models, Phys. Rev. B, 51:12579-12584, 1995.
- [13] A.A. Zvyagin, Exactly solvable multichain supersymmetric t-J model, Phys. Rev. B, 52:15050-15053, 1995.
- [14] A.A. Zvyagin, Bethe ansatz solvable multi-chain quantum systems, J. Phys. A, 34(41):R21-R53, 2001.
- [15] H. Frahm and C. Rödenbeck, Integrable models of coupled heisenberg chains, Europhys. Lett., 33(1):47-52, 1996.
- [16] H. Frahm and C. Rödenbeck, Properties of the chiral spin liquid state in generalized spin ladders, J. Phys. A, 30(13):4467-4479, 1997.
- [17] H. Frahm and A. Seel, The staggered six-vertex model: Conformal invariance and corrections to scaling, Nucl. Phys. B, 879:382-406, 2014.
- [18] Y. Ikhlef, J.L. Jacobsen, and H. Saleur, A staggered six-vertex model with non-compact continuum limit, Nucl. Phys. B, 789(3):483-524, 2008.
- [19] Y. Ikhlef, J.L. Jacobsen, and H. Saleur, The z₂ staggered vertex model and its applications, J. Phys. A, 43(22):225201, 2010.
- [20] T.S. Tavares and G.A.P. Ribeiro, Thermodynamics of quantum spin chains with competing interactions, J. Stat. Mech., 2013(09):P09007, 2013.
- [21] T.S. Tavares and G.A.P. Ribeiro, Magnetocaloric effect in the spin-1/2 chain with competing interactions, J. Stat. Mech., 2014(11):P11026, 2014.
- [22] G. Gorohovsky, R.G. Pereira, and E. Sela, Chiral spin liquids in arrays of spin chains, Phys. Rev. B, 91:245139, 2015.

- [23] J.-H. Chen, C. Mudry, C. Chamon, and A. M. Tsvelik, Model of chiral spin liquids with abelian and non-abelian topological phases, Phys. Rev. B, 96:224420, 2017.
- [24] J. Cao, W.-L. Yang, K. Shi, and Y. Wang, Off-diagonal bethe ansatz and exact solution of a topological spin ring, Phys. Rev. Lett., 111:137201, 2013.
- [25] Y. Wang, W.-L. Yang, J. Cao, and K. Shi, Off-diagonal Bethe ansatz for exactly solvable models, Springer, 2015.
- [26] Y. Qiao, P. Sun, Z. Xin, J. Cao, and W.-L. Yang, Exact solution of an integrable anisotropic $J_1 J_2$ spin chain model, arXiv:1902.00688, 2019.