

Large Higgs quartic coupling and (A)DM from extended Bosonic Technicolor

Mads T. Frandsen* and Wei-Chih Huang†

CP³-Origins, University of Southern Denmark, Campusvej 55 DK-5230 Odense M, Denmark

We propose novel bosonic Technicolor models augmented by an $SU(2)_R$ gauge group and scalar doublet. Dynamical breaking of $SU(2)_R$ induced by technifermion condensation triggers $SU(2)_L$ breaking via a portal coupling. The scale of the new strong interactions is as high as that of composite Higgs models, and the vacuum stability challenge confronting ordinary bosonic Technicolor models is avoided. Thermal or asymmetric dark matter, whose stability is ensured by a $U(1)_{TB}$ technibaryon symmetry, can be realized. In the latter case, the correct relic density can be reproduced for a wide range of dark matter mass via leptogenesis.

Preprint: CP³-Origins-2019-46 DNRf90

INTRODUCTION

Extensions of Technicolor (TC) [1, 2] and Composite Higgs (CH) [3] models with dynamical SM fermion mass generations are challenging and complex [4–7].

In bosonic Technicolor (bTC) [8–13] and partially composite Higgs (pCH) [3, 14, 15] models, the dynamical fermion condensates are instead coupled to an elementary $SU(2)_L$ scalar doublet H_L via Yukawa interactions. SM fermion masses are generated via four-fermion operators by integrating out H_L [8] or via an induced vacuum expectation value (vev) of H_L [3]. In the latter case the electroweak (EW) boson masses originate from both the fermion condensate and the vev. As a result, the EW scale is $v_{EW} = (v_L^2 + f^2 \sin^2 \theta)^{1/2} = 246$ GeV where v_L is the vev of H_L , f is the Goldstone-boson decay constant of the composite sector, and the angle θ parameterizes the vacuum misalignment with $\sin \theta = 1$ being bTC.

In bTC the scale f is therefore below the EW scale such that new resonances from the strong dynamics can significantly modify EW precision observables and hence be severely constrained [16, 17]. Furthermore, in bTC models the Higgs quartic coupling at the EW scale is typically smaller than that of the SM (due to additional bTC contributions to the Higgs mass), but the top Yukawa coupling becomes larger. These two effects combined will usually turn the running quartic coupling negative below the bTC cut-off scale, leading to an issue of low-scale vacuum instability [16]. These challenges can be alleviated in pCH models because of the high compositeness scale f [14, 18–20].

On the other hand, another motivation for TC and bTC models was asymmetric technibaryon dark matter (DM), connecting the baryon and DM densities [21]. The lightest composite technibaryon is stable due to a $U(1)_{TB}$ asymmetry associated with technibaryon number TB . Similar to the lepton and baryon numbers L and B , TB is preserved up to anomalous $SU(2)_L$ sphalerons, yielding a relation $\alpha L + \beta B + \gamma TB = 0$, where the coefficients depend on particles involved in the sphalerons. In this case, sphalerons can transfer asymmetry among L , B and TB [22]. But in the (p)CH models the vacuum explicitly breaks the $U(1)_{TB}$ symmetry.

In this work, we propose a new class of bTC models, denoted by RbTC, with an augmented EW sector $SU(2)_L \times SU(2)_R \times U(1)_{Y'}$ in addition to the strongly-interacting gauge group G_{TC} . An $SU(2)_R$ doublet H_R is introduced and couples to technifermions (charged under

G_{TC} and $SU(2)_R$), the condensation of which induces a vev of H_R , breaking the $SU(2)_R$ symmetry. Then via a portal coupling $-\lambda_{LR} (H_L^\dagger H_L)(H_R^\dagger H_R)$, a negative mass for H_L (identified as the SM Higgs doublet) is generated, breaking the $SU(2)_L$ symmetry. All in all, we have

$$SU(2)_L \times SU(2)_R \times U(1)_{Y'} \xrightarrow{f, v_R} SU(2)_L \times U(1)_Y \xrightarrow{v_L = v_{EW}} U(1)_Q. \quad (1)$$

Different from conventional bTC models, where technifermions couple directly to H_L and thus can modify EW observables significantly because of $f < v_{EW}$, in RbTC the scale f is not directly related to v_L and can be above TeV: $v_L (= v_{EW}) < v_R < f$ in regions of interest. Furthermore, contrary to the positive contribution from technifermions to the SM Higgs mass in bTC, the SM Higgs boson receives a negative mass contribution via the portal coupling and mixes with the neutral component of H_R , leading to a larger quartic coupling and a smaller top-quark Yukawa coupling. Consequently, the issue of vacuum instability is solved.

The RbTC vacuum still preserves a global $U(1)_{TB}$ symmetry and the lightest composite state of technifermions charged under this $U(1)_{TB}$ can be electrically neutral and therefore a DM candidate. If SM fermions are also charged under $SU(2)_R$, the sphalerons of $SU(2)_{L,R}$ can transfer asymmetry among L , B and TB . As we shall see below both thermal DM and asymmetric DM (ADM) candidates arise in the RbTC framework.

RBTC MODELS FOR $SU(2)_L \times SU(2)_R \times U(1)_{Y'}$

The SM gauge group is extended with an $SU(2)_R$ and a strongly-coupled G_{TC} gauge groups,

$$SU(3)_{QCD} \times G_{TC} \times SU(2)_L \times SU(2)_R \times U(1)_{Y'}, \quad (2)$$

where the SM hypercharge is given by $Y = T_R^3 + Y'$. We restrict to minimal TC sectors with an $SU(2)_R$ doublet (C_R, S_R) in the representation $\mathcal{R}$ under G_{TC} and $SU(2)_R$ singlets $\tilde{C}_R, \tilde{S}_R$ in the conjugated representation. The global symmetry in the TC sector is $SU(2) \times SU(2) \times U(1)_{TB}$ if $\mathcal{R}$ is complex and is enlarged to $SU(4)$, acting on the vector Q consisting of four Weyl spinors $Q = (C_R, S_R, \tilde{C}_R, \tilde{S}_R)$, if $\mathcal{R}$ is (pseudo-)real.

By virtue of minimality, we choose $G_{TC} = SU(2)_{TC}$. In the first model discussed below, $\mathcal{R}$ is the pseudo-real fundamental representation of $SU(2)_{TC}$. In the second model, $\mathcal{R}$ is the real adjoint representation under $SU(2)_{TC}$. It is straightforward to generalize to other (pseudo-)real or complex representations.

In the pseudo-real and real $\mathcal{R}$, the condensation of $\langle Q_i^T Q_j \rangle$ is a linear combination of $SU(2)_R$ -breaking vacuum $E_{B_{\mp}}$ (so-called TC vacuum) and $SU(2)_R$ -preserving one E_- : $E = \sin \theta E_{B_{\mp}} + \cos \theta E_-$ with

$$E_{B_{\mp}} = \begin{pmatrix} 0 & 1 \\ \mp 1 & 0 \end{pmatrix}, \quad E_- = \begin{pmatrix} i\sigma_2 & 0 \\ 0 & -i\sigma_2 \end{pmatrix}, \quad (3)$$

where $E_{B_{+}}$ is for the real representation while $E_{B_{-}}$ is for the pseudo-real case. As DM stability is ensured by $U(1)_{TB}$ which remains unbroken only under $E_{B_{\mp}}$, it is paramount that $\theta = \pi/2$ is dynamically realized.

Model 1 - Thermal technibaryon DM

In this model, only the TC fermions and the H_R doublet are charged under the $SU(2)_R$ while the SM fields including the H_L doublet are gauged as in the SM. The particle contents of interest are summarized in Table I. The relevant Lagrangian describing the new strong sec-

	$SU(2)_{TC}$	$SU(2)_L$	$SU(2)_R$	$U(1)_{Y'}$	$U(1)_{TB}$
(C_R, S_R)	$\mathcal{R}$	1	$\square$	0	1
$\tilde{C}_R$	$\mathcal{R}$	1	1	-1/2	-1
$\tilde{S}_R$	$\mathcal{R}$	1	1	+1/2	-1
H_R	1	1	$\square$	+1/2	0
H_L	1	$\square$	1	+1/2	0

TABLE I: Field contents and quantum numbers of the first RbTC model for $SU(2)_L \times SU(2)_R \times U(1)_{Y'}$ electroweak symmetries. All of them are singlets under $SU(3)_{QCD}$.

tor and the elementary doublets consist of three parts: kinetic terms, Higgs potential and Yukawa couplings between Q and H_R

$$\mathcal{L}_{\text{RbTC}} = \mathcal{L}_{\text{kin}} - V(H_R, H_L) + \mathcal{L}_{Y_{CS}}. \quad (4)$$

Below the condensation scale, $\Lambda_{\text{RbTC}} \sim 4\pi f$, the global symmetry G breaks down to a subgroup H and we parameterise the composite Goldstone bosons Π_a in the coset G/H by

$$\Sigma = \exp \left(\sum_{a=1}^{\dim(G/H)} \frac{2\sqrt{2}i}{f} \Pi_a X_a \right) E, \quad (5)$$

where X_a are the broken generators. In our case, $G = SU(4)$ and $H = Sp(4), SO(4)$ with $\dim(G/H) = 5, 9$ for the pseudo-real and real representations, respectively. The generators are listed explicitly in Appendix. In terms of Σ and the gauge fields, the kinetic terms read

$$\begin{aligned} \mathcal{L}_{\text{kin}} = & \frac{f^2}{8} \text{Tr}[(D_\mu \Sigma_R)^\dagger D^\mu \Sigma_R] \\ & + (D_\mu H_R)^\dagger D^\mu H_R + (D_\mu H_L)^\dagger D^\mu H_L, \end{aligned} \quad (6)$$

where

$$D_\mu \Sigma_R = \partial_\mu \Sigma_R - i(G_\mu \Sigma_R + \Sigma_R G_\mu^T), \quad (7)$$

containing the gauge fields

$$G_\mu = g_R W_{\mu R}^i T_R^i + g_{Y'} B'_\mu T_{Y'}, \quad (8)$$

with

$$T_R^i = \frac{1}{2} \begin{pmatrix} \sigma_i & 0 \\ 0 & 0 \end{pmatrix}, \quad T_{Y'} = \frac{1}{2} \begin{pmatrix} 0 & 0 \\ 0 & -\sigma_3 \end{pmatrix}, \quad (9)$$

where $i = (1, 2, 3)$ and σ_i are the Pauli matrices. Note that T_R^i only act on the first two elements of Q , namely (C_R, S_R) , as they are embedded in the $SU(2)_R$ doublet. For the covariant derivative of $H_{L,R}$, it reads

$$D_\mu P = \partial_\mu - i \frac{1}{2} (g_P W_{\mu P}^i \sigma_i + g_{Y'} B'_\mu), \quad (10)$$

where $P = L, R$. In light of the $SU(2)_{L,R}$ gauge symmetry, the most general renormalizable potential of $H_{L,R}$ is

$$\begin{aligned} V(H_R, H_L) = & m_R^2 H_R^\dagger H_R + m_L^2 H_L^\dagger H_L + \lambda_R (H_R^\dagger H_R)^2 \\ & + \lambda_L (H_L^\dagger H_L)^2 - \lambda_{LR} H_L^\dagger H_L H_R^\dagger H_R, \end{aligned} \quad (11)$$

where $H_A = \left(h_A^+ \frac{h_A + i\tilde{h}_A}{\sqrt{2}} \right)^T$ with $A = (L, R)$, and $\lambda_{LR} > 0$. As mentioned above, the $H_{L,R}$ mixing term with a negative coefficient can induce a vev of H_L after H_R develops a vev, even if the H_L has a positive mass term, $m_L^2 > 0$. Composite dynamics inducing a vev for H_L via a second scalar multiplet is also studied in a scale invariant extension of the SM [23], where the additional scalar is a singlet, rather than our $SU(2)_R$ doublet. Our elementary scalar sector is instead similar to the Gauged Two Higgs Doublet Model [24] where the vev of $SU(2)_L$ is induced by a vev or condensation from another sector.

Finally, Yukawa couplings of Q to H_R are included:

$$\mathcal{L}_{Y_{CS}} = -y_C \epsilon^{\alpha\beta} H_{R\alpha} (Q^T P_\beta Q) + y_S H_{R\alpha}^* (Q^T \tilde{P}_\alpha Q) + \text{h.c.}, \quad (12)$$

where α and β are the $SU(2)_{L,R}$ indices with $\epsilon^{\alpha\beta} = \delta^{\alpha 1} \delta^{\beta 2} - \delta^{\alpha 2} \delta^{\beta 1}$, $2(P_\alpha)_{ij} = \delta_{i\alpha} \delta_{j3} \mp \delta_{i3} \delta_{j\alpha}$ and $2(\tilde{P}_\alpha)_{ij} = \delta_{i\alpha} \delta_{j4} \mp \delta_{i4} \delta_{j\alpha}$ [19]. The $- (+)$ sign corresponds to pseudo-real (real) representations of $\mathcal{R}$. It is clear that the condensation results in a linear term in H_R , implying a nonzero vev of H_R and thus breaking $SU(2)_R$ regardless of the mass term m_R^2 . Below the condensation scale, the interactions lead to an effective potential [15]

$$\begin{aligned} V_{\text{eff}}^0 = & 4\pi f^3 Z_2 \left(y_C \epsilon^{\alpha\beta} H_{R\alpha} \text{Tr} [P_\beta \Sigma^T] + y_S \tilde{H}_{R\alpha}^* \text{Tr} [\tilde{P}_\alpha \Sigma^T] \right. \\ & \left. + \text{h.c.} \right), \end{aligned} \quad (13)$$

where Z_2 is a non-perturbative $\mathcal{O}(1)$ constant [25]. For simplicity, we set $y_C = y_S \equiv y_{CS}/2$.

So far, the scalar potential contains two components: $V(H_L, H_R)$ and contributions from the previous Yukawa coupling. However, due to the fact the $SU(2)_R \times U(1)_{Y'}$

gauge symmetry explicitly breaks the global symmetry group $SU(4)$, there exists another contribution to the effective potential which can destabilize the TC vacuum [26, 27]. The corresponding gauge contribution is

$$V_{\text{eff}}^{1-loop} \supset -\frac{1}{2}\tilde{C}_g Z_2^2 f^4 c_\theta^2, \quad (14)$$

with $\tilde{C}_g \equiv C_g(3g_R^2 + g_{Y'}^2)$ for the pseudo-real representation and $\tilde{C}_g = 0$ for the real one. The C_g is a loop factor, assumed to be of $\mathcal{O}(1)$ here. In the following computation on minimization and Higgs masses, we study the pseudo-real case but results of the real $\mathcal{R}$ can be obtained simply by $E_{B-} \rightarrow E_{B+}$. The total effective scalar potential becomes

$$V_{\text{eff}} = -4\sqrt{2}\pi f^3 Z_2 y_{CS} \sin\theta h_R - \frac{1}{2}\tilde{C}_g Z_2^2 f^4 c_\theta^2 + V(h_R, h_L), \quad (15)$$

where the expression is written in terms of real, neutral components of the doublets, $h_{L,R}$.

The condition of the vacuum being a minimum is the vanishing of the first derivatives $\partial V_{\text{eff}}/\partial x_i$ with respect to $x_i = h_L, h_R$, and θ , evaluated at $(v_L, v_R, \pi/2)$, respectively. It yields

$$\begin{aligned} 2v_R (m_R^2 + \lambda_R v_R^2) &= 8\sqrt{2}\pi y_{CS} Z_2 f^3 + \lambda_{LR} v_L^2 v_R \\ m_L^2 + \lambda_L v_L^2 &= \frac{1}{2}\lambda_{LR} v_R^2, \end{aligned} \quad (16)$$

while $0 = \partial V_{\text{eff}}/\partial\theta|_{\theta=\pi/2}$ is automatically satisfied. Moreover, this minimum is stable if eigenvalues of the matrix of the second derivatives (the Hessian) $\partial^2 V_{\text{eff}}/\partial x_i \partial x_j$ are positive and if the potential is bounded from below for large field values in all directions (e.g., Ref. [28]). In addition, for the minimum to be the TC vacuum we require $s_\theta = 1$, which is non-trivial for the pseudo-real representations. All in all, we have the following constraints

$$\begin{aligned} 4\sqrt{2}\pi y_{CS} v_R &> \tilde{C}_g Z_2 f, \quad \lambda_L > 0, \\ 4\lambda_R + 8\sqrt{2}\pi y_{CS} Z_2 \frac{f^3}{v_R^3} &> \frac{\lambda_{LR}^2}{\lambda_L}, \quad 4\lambda_L \lambda_R > \lambda_{LR}^2, \end{aligned} \quad (17)$$

where the first criterion ensures $\theta = \pi/2$ is a stable minimum, i.e., the TC vacuum with unbroken $U(1)_{\text{TB}}$.

The mass matrix of the CP-even scalars $h_{L,R}$ is

$$M_h^2 = \begin{pmatrix} 2\lambda_L v_L^2 & -\lambda_{LR} v_L v_R \\ -\lambda_{LR} v_L v_R & M^2 \end{pmatrix} \quad (18)$$

where $M^2 = 4\sqrt{2}\pi Z_2 y_{CS} \frac{f^3}{v_R} + 2\lambda_R v_R^2$, and $2\lambda_L v_L^2$ would be the SM Higgs mass expression given $v_L = 246$ GeV with $\lambda_L = 0.13$. The resulting mass eigenstates are

$$h_1 = c_\alpha h_L + s_\alpha h_R, \quad h_2 = s_\alpha h_L - c_\alpha h_R, \quad (19)$$

with

$$t_{2\alpha} = \frac{2\lambda_{LR} v_L v_R}{M^2 - 2\lambda_L v_L^2}. \quad (20)$$

In the limit of small α , the masses of $h_{1,2}$ are

$$m_{h_1}^2 \simeq 2\lambda_L v_L^2 - \frac{t_{2\alpha}}{2}\lambda_{LR} v_L v_R, \quad m_{h_2}^2 \simeq M^2, \quad (21)$$

where h_1 is identified as the 125 GeV Higgs boson. Therefore, the value of λ_L is larger than that of the SM in order to compensate the negative contribution from the mixing, while the Higgs-fermion and Higgs-gauge couplings are reduced by a factor of c_α . In this case, a very SM-like Higgs boson and vacuum stability to a high scale is easily attained unlike ordinary bTC models.

In Fig. 1, assuming $(v_L, v_R, f, \lambda_R, y_{CS}, s_\alpha) = (246 \text{ GeV}, 3 \text{ TeV}, 5 \text{ TeV}, 0.5, 1, < 0.05)$, the purple region satisfies Eq. (17) and $m_{h_1} = 125$ GeV. The blue region is further constrained by $m_L^2, m_R^2 > 0$. That is, both $SU(2)_R$ and $SU(2)_L$ symmetry breaking are induced by strong dynamics. In the two regions, the value of λ_L can be much larger than the SM value marked by the red dashed line.




FIG. 1: The purple region of the λ_L - λ_{LR} plane satisfies the vacuum stability constraints in Eq. (17) and reproduces a 125 GeV SM-like Higgs. The blue subregion further satisfies $m_L^2, m_R^2 > 0$ such that both the $SU(2)_R$ and $SU(2)_L$ symmetries are dynamically broken.

The Goldstone bosons $h_L^\pm$ and $\tilde{h}_L$ are eaten by the $W^\pm, Z$ bosons, and the masses of the gauge bosons are the exactly same as in the SM at tree level since $v_L = v_{\text{EW}}$. On the other hand, as can be seen from Eq. (6) the $SU(2)_R$ $W_R^\pm$ and Z_R absorb linear combinations of $h_R^\pm, \tilde{h}_R$ and $\Pi_{1,2,3}$ and become massive:

$$m_{W_R^\pm}^2 = \frac{g_R^2}{4}(f^2 + v_R^2), \quad m_{Z_R}^2 = \frac{g_R^2 + g_{Y'}^2}{4}(f^2 + v_R^2). \quad (22)$$

The identification of absorbed Goldstone bosons, the mass spectrum of physical scalars, and the mixing between the neutral gauge bosons are discussed in Appendix.

To demonstrate that the correct DM relic density can be obtained, we study the DM annihilation cross-section in the pseudo-real $\mathcal{R}$. The DM candidate is the neutral complex Goldstone boson carrying a $U(1)_{\text{TB}}$ charge of 2

$$\Pi_{CS} \equiv (\Pi_4 - i\Pi_5)/\sqrt{2}. \quad (23)$$

It receives a mass from gauge interactions and the Yukawa interaction in Eq. (13)

$$m_{\Pi_{CS}}^2 = -\frac{3(3g_R^2 + g_{Y'}^2)}{64\pi^2}\tilde{m}^2 + 4\sqrt{2}\pi Z_2 y_{CS} f v_R, \quad (24)$$

where $\tilde{m}$ is $\mathcal{O}(f)$ [26, 29].

The kinetic term in Eq. (6) gives rise to a contact interaction of Π_{CD} with the $SU(2)_R$ gauge bosons: $-(g_R^2/2)W_{R\mu}^+W_R^{-\nu}\Pi_{CS}\bar{\Pi}_{CS}$, implying the DM annihilation cross-section is

$$\langle\sigma v\rangle\approx\frac{g_R^4}{256\pi m_{\Pi_{CS}}^2}(8t^2-12t+9), \quad (25)$$

where $t=m_{\Pi_{CS}}^2/m_{W_R}^2$. The desired cross-section of $3\times 10^{-26}\text{cm}^3/\text{sec}$ for the correct DM density can be easily attained, given $g_R\sim\mathcal{O}(1)$ and $m_{\Pi}\gtrsim m_{W_R}\sim\text{TeV}$.

Model 2 - asymmetric technibaryon DM

In this model, the SM right-handed charged leptons ℓ_R and additional right-handed neutrinos ν_R form $SU(2)_R$ doublets, $l_R=(\nu_\alpha\ell_\alpha)^T_R$ with $\alpha=(e,\mu,\tau)$. The particle contents and quantum numbers are summarized in Table II. The technifermions are in the adjoint representation and the model is free from the gauge and Witten anomalies.

	$SU(3)_{QCD}$	$SU(2)_{TC}$	$SU(2)_L$	$SU(2)_R$	$U(1)_{Y'}$
q_L	$\square$	1	$\square$	1	$+1/6$
u_R	$\square$	1	1	1	$+2/3$
d_R	$\square$	1	1	1	$-1/3$
l_L	1	1	$\square$	1	$-1/2$
l_R	1	1	1	$\square$	$-1/2$
(C_R, S_R)	1	Adj	1	$\square$	$1/2$
$\tilde{C}_R$	1	Adj	1	1	-1
$\tilde{S}_R$	1	Adj	1	1	0
$N^0(N^\pm)$	1	1	1	1	$0(\pm 1)$
H_R	1	1	1	$\square$	$+1/2$
H_L	1	1	$\square$	1	$+1/2$

TABLE II: Field contents and quantum numbers of the second RbTC model. The $U(1)_{TB}$ charge assignment of technifermions is the same as in Table I.

The quarks obtain masses via Yukawa couplings of H_L , analogous to the SM. In contrast, lepton masses and couplings to the Higgs h_1 are realized via Yukawa couplings with new charged and neutral vector-like massive fermions, $N^\pm$ and N^0 , respectively:

$$\begin{aligned} \mathcal{L}_{\text{Yuk}} = & -y_L\bar{l}_L\tilde{H}_LN^0 - y_R\bar{l}_R\tilde{H}_RN^0 \\ & -y'_L\bar{l}_L H_L N^- - y'_R\bar{l}_R H_R N^- + \text{h.c.}, \end{aligned} \quad (26)$$

where the flavor indices are suppressed and $\tilde{H}=\epsilon H^*$. By integrating out the heavy $N^\pm$ and N^0 fermions, one obtains the lepton masses:

$$m_\ell = \frac{|y'_L y'_R|}{2m_N^\pm} v_L v_R, \quad m_\nu = \frac{|y_L y_R|}{2m_N^0} v_L v_R. \quad (27)$$

Since $m_{\ell,\nu}\ll v_L$, $m_N^\pm$ and m_N^0 can be much larger than v_R , given $\mathcal{O}(y'_{(L,R)})\sim 1$, which justifies integrating out $N^\pm$ and N^0 .

If both the $SU(2)_L$ and $SU(2)_R$ sphalerons are in equilibrium at temperatures $T>f$, one has:

$$L_L + B = 0, \quad L_R + \frac{1}{2}TB = 0, \quad (28)$$

where

$$\begin{aligned} L_{L(R)} &= \sum_{e,\mu,\tau} 2\mu_{l_{L(R)}} \quad , \quad B = 3(2\mu_{q_L} + \mu_{u_R} + \mu_{d_R}) \quad , \\ TB &= \frac{3}{2}(\mu_{C_R} + \mu_{S_R} - \mu_{\tilde{C}_R} - \mu_{\tilde{S}_R}) \quad , \end{aligned} \quad (29)$$

refer to asymmetries in the lepton, baryon and TC sectors, respectively.

After taking into account the sphalerons, Yukawa interactions and $U(1)_{Y'}$ neutrality conditions with $\mu_W = \mu_{W_R} = 0$, all potentials can be rewritten in terms of two unconstrained chemical potentials, chosen to be μ_{l_L} and μ_{l_R} . That is the reason why both L_L and L_R are needed in Eq. (28). As demonstrated in Appendix, the final TB and B have different dependence on the initial values (denoted by the superscript i)

$$B = \frac{2}{5}(B^i - L_L^i) \quad , \quad TB = \frac{1}{3}(TB^i - 2L_R^i) \quad , \quad (30)$$

implying that final B and TB can be uncorrelated. To generate an initial asymmetry, one can resort to leptogenesis [30] by having N^0 be a Majorana fermion (instead of being vector-like) and decay asymmetrically and out of equilibrium into both $l_L + H_L$ ($L_L^i \neq 0$) and $l_R + H_R$ ($L_R^i \neq 0$). The asymmetries are controlled by the Yukawa couplings y_L and y_R in Eq. (26), respectively. In this case, one can obtain the correct relic density of ADM for any mass by adjusting y_R . In contrast, to achieve the correct relic density in simple TC scenarios, one usually has to rely on the Boltzmann suppression to reduce the number density of heavy ADM, given $\mu_B \sim \mu_{TB}$ [22].

CONCLUSIONS

We have proposed a novel class of extended bTC models, featuring $SU(2)_L \times SU(2)_R \times U(1)_{Y'}$ EW symmetry with an $SU(2)_R$ doublet and an $SU(2)_L$ doublet scalars, identified as the SM Higgs doublet. The technifermions are charged only under the $SU(2)_R \times U(1)_{Y'}$ and the condensation triggers $SU(2)_R$ breaking, which in turn renders $SU(2)_L$ symmetry broken. In this scenario, the compositeness scale is much larger than the EW scale, and the Higgs quartic coupling can be much larger than the SM value. That implies the models do not suffer from the problems of vacuum stability which plague ordinary bTC models.

In this framework we obtain DM candidates whose stability is ensured by an unbroken $U(1)_{TB}$ symmetry. In the first model we considered where the SM fermions are not gauged under $SU(2)_R$, the DM relic density can be thermal. In the second model where the SM leptons are also charged under $SU(2)_R$, the $SU(2)_{L,R}$ sphalerons can transfer particle asymmetries among leptons, baryons

and technifermions. Asymmetric DM can then be realized via leptogenesis.

Here we briefly comment on constraints from DM direct detection and collider Z' resonance searches. In Model 1, the DM candidate is a pure singlet under all gauge groups. Hence, it couples to SM fermions only through the small H_L - H_R mixing and in turn is suppressed. By contrast, in Model 2 both the DM candidate and SM fermions couple to the heavy neutral boson Z' , leading to DM-nucleon interactions (dominated by the proton) that can be approximated as $\sigma \approx \frac{18(g_{Y'}^2 - g'^2)^2}{\pi g_{Y'}^4} \frac{\mu^2}{(f^2 + v_R^2)^2}$, where μ is the DM-nucleon reduced mass and g' is the SM $U(1)_{Y'}$ coupling. The latest XENON1T result [31] implies $\sqrt{f^2 + v_R^2} \gtrsim 15$ TeV, depending on the DM mass. It implies the Z' is heavier than 5 TeV, assuming $g_R \sim g_L$. Therefore by satisfying the direct search bound the model also avoids the recent constraints, e.g., from di-lepton resonance searches [32], that are relevant as Z' couples to both SM quarks and leptons. The latest high-mass resonances on W' and Z' can be found in Refs [32–38] where the bounds are around multiple TeV, depending on the underlying models. These experimental bounds also require the DM mass to be heavier than TeV since the mass is proportional to f and v_R as shown in Eqs. (24) and (44). Moreover, since all technifermions are singlets under $SU(2)_L$, there are no contributions to the oblique parameters (S , T , U) [39, 40], as illustrated in, e.g., Ref [41]. Because the fermions do carry $U(1)_{Y'}$ charges there is a contribution to the Y parameter [42], which can be estimated either from fermion loops in the underlying technifermion theory or using the effective resonance Lagrangian to be $Y \ll v_L^2/f^2 \lesssim 10^{-4}$ [43] that is well below the bound, $Y \lesssim 10^{-3}$, from the electroweak precision data [42].

Lastly, the large Higgs quartic coupling can be tested in next-generation colliders and in the second model, the $SU(2)_R$ and DM particles can be potentially probed by future e^+e^- colliders, such as ILC [44], FCC-ee (formerly known as TLEP [45]) and CEPC [46]. We leave for future work detailed phenomenology studies as well as other possible charge assignments of SM fermions under $SU(2)_R$ and different strongly-interacting sectors.

Acknowledgments

MTF and WCH acknowledge partial funding from the Independent Research Fund Denmark, grant number DFF 6108-00623. The CP3-Origins centre is partially funded by the Danish National Research Foundation, grant number DNRF90.

Appendix

$SU(4)/Sp(4)$

Here we explicitly give the $SU(4)$ generators, e.g. [47, 48] in terms of the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (31)$$

and the 2-by-2 identity matrix, denoted by $\mathbb{1}$. The ten unbroken operators with respect to the vev E_- in Eq. (3) are

$$S_{1,2,3} = \frac{1}{2} \begin{pmatrix} \sigma_{1,2,3} & 0 \\ 0 & 0 \end{pmatrix}, \quad S_{4,5,6} = \frac{1}{2} \begin{pmatrix} 0 & 0 \\ 0 & -\sigma_{1,2,3}^T \end{pmatrix}, \quad S_{7,8,9} = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & i\sigma_{1,2,3} \\ -i\sigma_{1,2,3} & 0 \end{pmatrix}, \quad S_{10} = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}, \quad (32)$$

while the five unbroken generators are

$$X'_{1,3,4} = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & \sigma_{3,1,2} \\ \sigma_{3,1,2} & 0 \end{pmatrix}, \quad X'_2 = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & i\mathbb{1} \\ -i\mathbb{1} & 0 \end{pmatrix}, \quad X'_5 = \frac{1}{2\sqrt{2}} \begin{pmatrix} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{pmatrix}. \quad (33)$$

In case of the vev of $E = \cos \theta E_- + \sin \theta E_{B_-}$, the broken generators become:

$$X_i = \cos \theta X'_i \mp \frac{\sin \theta}{\sqrt{2}} (S_i - S_{i+3}), \quad X_4 = X'_4, \quad X_5 = \cos \theta X'_5 - \sin \theta S_8, \quad (34)$$

where $i = (1, 2, 3)$, the “−” sign for $i = 1$ and “+” for $i = (2, 3)$ in the first equation.

From the kinetic term in Eq. (6) and the definition of Σ , we can identify the Goldstone bosons that are absorbed by the $W_R^\pm$ and Z_R through

$$\mathcal{L} \supset \frac{g_R}{2} \sqrt{f^2 + v_R^2} \left(W_R^{+\mu} \partial_\mu G^- + h.c. + \sqrt{1 + \frac{g_{Y'}^2}{g_R^2}} Z_R^\mu \partial_\mu G^0 \right) \quad (35)$$

where

$$G^\pm = \frac{f\Pi^\pm \pm i v_R h_R^\pm}{\sqrt{f^2 + v_R^2}}, \quad G^0 = \frac{-f\Pi_3 + v_R \tilde{h}_R}{\sqrt{f^2 + v_R^2}}, \quad (36)$$

with $\Pi^\pm = (\Pi_1 \pm i\Pi_2)/\sqrt{2}$. The corresponding physical states are

$$S^\pm = \frac{v_R \Pi^\pm \mp i f h_R^\pm}{\sqrt{f^2 + v_R^2}}, \quad S^0 = \frac{v_R \Pi_3 + f \tilde{h}_R}{\sqrt{f^2 + v_R^2}}. \quad (37)$$

Therefore, the remaining degrees of freedom in Π_a are Π_4 and Π_5 which comprise DM, Π_{CS} in Eq. (23). Finally the mass terms of the physical states of $S^\pm$ and S^0 induced by the Yukawa interactions in Eq. (13) read

$$m_{S^\pm}^2 = m_{S^0}^2 = 4\sqrt{2} \pi y_{CS} Z_2 \left(\frac{2f^2 + v_R^2}{f^2 + v_R^2} \right) f v_R. \quad (38)$$

$SU(4)/SO(4)$

The six unbroken generators under the vev of E_{B+} are, e.g. [49, 50]

$$S_i = \frac{1}{2\sqrt{2}} \begin{pmatrix} \sigma_i & 0 \\ 0 & -\sigma_i^T \end{pmatrix}, \quad S_{5,6} = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & \mathbf{B}_{5,6} \\ (\mathbf{B}_{5,6})^\dagger & 0 \end{pmatrix}, \quad (39)$$

where $i = (1, 2, 3, 4)$ with $\sigma_4 = \mathbb{1}$, $\mathbf{B}_5 = \sigma_2$ and $\mathbf{B}_6 = i\sigma_2$. The nine broken generators are

$$X_{1,2,3} = \frac{1}{2\sqrt{2}} \begin{pmatrix} \sigma_{1,2,3} & 0 \\ 0 & \sigma_{1,2,3}^T \end{pmatrix}, \quad X_i = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & \mathbf{D}_i \\ (\mathbf{D}_i)^\dagger & 0 \end{pmatrix}, \quad (40)$$

where $i = (4, \dots, 9)$ with $\mathbf{D}_4 = \mathbb{1}$, $\mathbf{D}_5 = i\mathbb{1}$, $\mathbf{D}_6 = \sigma_3$, $\mathbf{D}_7 = i\sigma_3$, $\mathbf{D}_8 = \sigma_1$ and $\mathbf{D}_9 = i\sigma_1$.

The combinations of $\Pi_{1,2,3}$, $h_R^\pm$ and $\tilde{h}_R$ absorbed by $W_R^\pm$ and Z_R are

$$G^\pm = \frac{f\Pi^\pm \mp iv_R h_R^\pm}{\sqrt{f^2 + v_R^2}}, \quad G^0 = \frac{-f\Pi_3 + v_R \tilde{h}_R}{\sqrt{f^2 + v_R^2}}, \quad (41)$$

with $\Pi^\pm = (\Pi_1 \mp i\Pi_2)/\sqrt{2}$. Note that there exist sign differences on the equations between the $Sp(4)$ and $SO(4)$ cases due to the different definitions of $SU(4)$ operators. The three uneaten states are

$$S^\pm = \frac{v_R \Pi^\pm \pm if h_R^\pm}{\sqrt{f^2 + v_R^2}}, \quad S^0 = \frac{v_R \Pi_3 + f \tilde{h}_R}{\sqrt{f^2 + v_R^2}}, \quad (42)$$

and also six degrees of freedom from $\Pi_{4,\dots,9}$:

$$\begin{aligned} \Pi_{CC} &= \frac{\Pi_4 + i\Pi_5 + (\Pi_6 + i\Pi_7)}{2}, \quad \Pi_{CS} = \frac{\Pi_8 + i\Pi_9}{\sqrt{2}}, \\ \Pi_{SS} &= \frac{\Pi_4 + i\Pi_5 - (\Pi_6 + i\Pi_7)}{2}, \quad \Pi_{SS}^*, \Pi_{CC}^*, \Pi_{CS}^*, \end{aligned} \quad (43)$$

where all of them carry two units of $U(1)_{TB}$ charge and Π_{SS} is the neutral DM candidate.

The mass contributions from the Yukawa interactions to $S^\pm$ and S^0 are exactly the same as the $Sp(4)$ case shown in Eq. (38). The other $U(1)_{TB}$ charged particles have

$$\begin{aligned} m_{CC}^2 &= 4\sqrt{2}\pi y_{CS} Z_2 f v_R + \frac{3}{64\pi^2} (g_R^2 + 9g_{Y'}^2) \tilde{m}^2, \\ m_{CS}^2 &= 4\sqrt{2}\pi y_{CS} Z_2 f v_R + \frac{3}{64\pi^2} (g_R^2 + 3g_{Y'}^2) \tilde{m}^2, \\ m_{SS}^2 &= 4\sqrt{2}\pi y_{CS} Z_2 f v_R + \frac{3}{64\pi^2} (g_R^2 + g_{Y'}^2) \tilde{m}^2, \end{aligned} \quad (44)$$

where $\tilde{m}^2$ is $\mathcal{O}(f^2)$ [26, 29]. Clearly, Π_{SS} is the lightest one and hence the DM candidate.

Neutral gauge boson mixing and couplings to fermions

The kinetic terms in Eq. (6) induces the mixing among the neutral gauge bosons and the mass matrix reads

$$\mathcal{M}_0^2 = \frac{1}{4} \begin{pmatrix} g_{Y'}^2 (f^2 + v_R^2 + v_L^2) & -g_{Y'} g_L v_L^2 & -g_{Y'} g_R (f^2 + v_R^2) \\ -g_{Y'} g_L v_L^2 & g_L^2 v_L^2 & 0 \\ -g_{Y'} g_R (f^2 + v_R^2) & 0 & g_R^2 (f^2 + v_R^2) \end{pmatrix}, \quad (45)$$

in the basis of (B', W_L^3, W_R^3) . That results in three mass eigenstates: the massless photon, the SM Z and an additional heavy neutral Z' . In regions of interest where v_R and f are above the TeV scale ($v_L \ll v_R, f$), the Z mass is the same as in the SM, $m_Z^2 \approx (g'^2 + g_L^2) v_L^2/4$, and for Z' we have $m_{Z'}^2 \approx (g'^2 + g_R^2) (f^2 + v_R^2)/4$, where $1/g'^2 = 1/g_{Y'}^2 + 1/g_R^2$ with g' being the SM $U(1)_Y$ coupling. Note that the fundamental and adjoint cases yield the same matrix matrix.

One can diagonalize the matrix with a rotation matrix $\mathcal{R}$:

$$\mathcal{R}^T \mathcal{M}_0^2 \mathcal{R} \quad (46)$$

which is the product of three rotation matrices R^{12} , R^{13} and R^{23}

$$\mathcal{R} = R^{12}(\theta_{12}) \cdot R^{13}(\theta_{13}) \cdot R^{23}(\theta_{23}), \quad (47)$$

where

$$\begin{aligned} [R^{ij}(\theta)]_{\alpha\beta} &= \cos \theta (\delta_{\alpha i} \delta_{\beta j} + \delta_{\alpha j} \delta_{\beta i}) \\ &\quad + \sin \theta (\delta_{\alpha i} \delta_{\beta j} - \delta_{\alpha j} \delta_{\beta i}) + \delta_{\alpha\beta} \epsilon_{ij\alpha}, \end{aligned} \quad (48)$$

where $\epsilon_{ij\alpha}$ is the Levi-Civita symbol in three dimensions. The three rotation angles are

$$\begin{aligned} \tan \theta_{12} &= \frac{g'}{g_L}, \quad \tan \theta_{13} = \frac{g_{Y'}}{g_R}, \\ \tan 2\theta_{23} &\approx 2g_{Y'}^2 \frac{\sqrt{g_{Y'}^2 g_R^2 + g_L^2 (g_{Y'}^2 + g_R^2)}}{(g_{Y'}^2 + g_R^2)^2} \left(\frac{v_L^2}{f^2 + v_R^2} \right), \end{aligned} \quad (49)$$

and the flavor and mass eigenstates are connected via the mixing matrix $\mathcal{R}$ as $(B', W_L^3, W_R^3)_\alpha^T = \mathcal{R}_{\alpha\beta}(\gamma, Z, Z')_\beta^T$.

It is straightforward to show that, up to a small correction characterized by θ_{23} , fermions couplings to γ and Z are the same as in the SM that are determined by the electric charge Q_f ($\equiv Q_{Y'} + T_L^3 + T_R^3$) and the weak isospin T_L^3 . On the other hand, the fermion coupling to Z' is $(-g_{Y'}^2 Q_{Y'} + g'^2 (Q_{Y'} + T_R^3)) / \sqrt{g_{Y'}^2 - g'^2}$.

Chemical equilibrium conditions

We here follow the formalism employed in Ref. [51] to perform the analysis on the chemical potentials of equilibrium above the phase transition scale. That is, the potentials of W and W_R are zero, and particles embedded

in an $SU(2)_L$ or $SU(2)_R$ doublet have the same chemical potential, denoted by the potential of the doublet; e.g., $\mu_{u_L} = \mu_{d_L} \equiv \mu_{q_L}$.

Moreover, due to the CKM mixing matrix among quark generations and the common origin of the lepton mass in Eq. (26), chemical potentials are the same among different generations. The Yukawa coupling interactions imply

$$\begin{aligned} \mu_{q_L} + \mu_{H_L} - \mu_{u_R} &= 0, \quad \mu_{q_L} - \mu_{H_L} - \mu_{d_R} = 0, \\ \mu_{l_L} - \mu_{H_L} + \mu_{u_{N^+}} &= 0, \quad \mu_{l_R} - \mu_{H_R} + \mu_{u_{N^+}} = 0, \\ -\mu_{C_R} + \mu_{H_R} - \mu_{\tilde{C}_R} &= 0, \quad \mu_{C_R} + \mu_{H_R} + \mu_{\tilde{S}_R} = 0, \end{aligned} \quad (50)$$

while the neutrality condition of $U(1)_Y$ charges dictates

$$\begin{aligned} 0 = & 9 \left(\frac{1}{3} \mu_{q_L} + \frac{2}{3} \mu_{u_R} - \frac{1}{3} \mu_{d_R} \right) - 3 (\mu_{l_L} + \mu_{l_R}) \\ & + 3 (\mu_{C_R} - \mu_{\tilde{S}_R}) + 2 (\mu_{H_L} + \mu_{H_R}) + 2 \mu_{N^+}. \end{aligned} \quad (51)$$

Note that we do not take into account Yukawa interactions of y_L and y_R in Eq. (26) as N^0 is assumed to be heavy and the couplings are required to be small to realize a neutrino mass of eV such that $l_L + H_L \leftrightarrow l_R + H_R$ mediated by N^0 is not in equilibrium.

On the other hand, the $SU(2)_{L,R}$ sphalerons yields:

$$3\mu_{q_L} + \mu_{l_L} = 0, \quad \mu_{C_R} + \mu_{l_R} = 0. \quad (52)$$

In light of the above constraints, all chemical potentials can be expressed as functions of two unconstrained chemical potentials chosen to be μ_{l_L} and μ_{l_R} . From Eq. (29), we can obtain the asymmetry of TB and L normalized to that of B as

$$\frac{TB}{B} = \frac{3\mu_{l_R}}{2\mu_{l_L}}, \quad \frac{L}{B} = \frac{L_L + L_R}{B} = -\frac{3}{2} \left(1 + \frac{\mu_{l_R}}{\mu_{l_L}} \right) \quad (53)$$

with $B = -4\mu_{l_L}$.

Two conserved quantities, directions perpendicular to $SU(2)_{L,R}$ sphalerons in Eq. (28), denoted as C_1 and C_2 are

$$C_1 = -L_L + B, \quad C_2 = -2L_R + TB, \quad (54)$$

which are invariant under two sphaleron processes. Thus, one can express the two unconstrained parameters μ_{l_L} and μ_{l_R} in terms of initial values of $L_{L,R}$, B and TB by $C_{1,2}^{\text{initial}} = C_{1,2}^{\text{final}}$. The final asymmetry reads

$$\begin{aligned} L_L &= -\frac{3}{5} (B^i - L_L^i), \quad L_R = -\frac{1}{3} (TB^i - 2L_R^i), \\ B &= \frac{2}{5} (B^i - L_L^i), \quad TB = \frac{1}{3} (TB^i - 2L_R^i), \end{aligned} \quad (55)$$

where the superscript i refers to the initial values and it is clear that $-L_L + B$ and $-2L_R + TB$ are conserved.

Note that in the context of ADM, the ratio of the number density of technibaryons (assuming a degenerate mass m_{TB}) to baryons at temperature T is linked to ratio of the chemical potentials as

$$\frac{Y_{TB}}{Y_B} = \frac{\mu_{TB}}{\mu_B} \zeta\left(\frac{m_{TB}}{T}\right). \quad (56)$$

where the function $\zeta(z)$ is given by

$$\zeta(z) = \frac{6}{\pi^2} \int_z^\infty dx x \sqrt{x^2 - z^2} \frac{e^x}{(e^x - \eta_i)^2}, \quad (57)$$

with $\eta_i = 1$ (-1) for a boson (fermion). For a relativistic boson (fermion) with $z \ll 1$, we have $\zeta(z) \approx 2$ (1). In case of $m_{TB} \gg m_B$ and $\mu_{TB} \sim \mu_B$, a large suppression from ζ is needed to obtain comparable energy densities: $\Omega_{TB} \sim \Omega_B$.

* Electronic address: frandsen@cp3.sdu.dk

† Electronic address: huang@cp3.sdu.dk

- [1] S. Weinberg, Phys.Rev. **D13**, 974 (1976).
- [2] L. Susskind, Phys.Rev. **D20**, 2619 (1979).
- [3] D. B. Kaplan and H. Georgi, Phys.Lett. **B136**, 183 (1984).
- [4] E. Eichten and K. D. Lane, Phys. Lett. **B90**, 125 (1980).
- [5] S. Dimopoulos and L. Susskind, Nucl. Phys. **B155**, 237 (1979).
- [6] D. B. Kaplan, H. Georgi, and S. Dimopoulos, Phys.Lett. **B136**, 187 (1984).
- [7] D. B. Kaplan, Nucl. Phys. **B365**, 259 (1991).
- [8] G. 't Hooft, NATO Sci. Ser. B **59**, 135 (1980).
- [9] E. H. Simmons, Nucl.Phys. **B312**, 253 (1989).
- [10] A. Kagan and S. Samuel, Phys. Lett. **B252**, 605 (1990).
- [11] S. Samuel, Nucl. Phys. **B347**, 625 (1990).
- [12] C. D. Carone and E. H. Simmons, Nucl. Phys. B **397**, 591 (1993), hep-ph/9207273.
- [13] C. D. Carone and H. Georgi, Phys. Rev. D **49**, 1427 (1994), hep-ph/9308205.
- [14] J. Galloway, A. L. Kagan, and A. Martin, Phys. Rev. **D95**, 035038 (2017), 1609.05883.
- [15] T. Alanne *et al.*, JHEP **01**, 051 (2018), 1711.10410.
- [16] C. D. Carone, Phys.Rev. **D86**, 055011 (2012), 1206.4324.
- [17] T. Alanne, S. Di Chiara, and K. Tuominen, JHEP **01**, 041 (2014), 1303.3615.
- [18] A. Agugliaro, O. Antipin, D. Becciolini, S. De Curtis, and M. Redi, Phys. Rev. **D95**, 035019 (2017), 1609.07122.
- [19] T. Alanne, M. T. Frandsen, and D. Buarque Franzosi, Phys. Rev. **D94**, 071703 (2016), 1607.01440.
- [20] T. Alanne, D. Buarque Franzosi, and M. T. Frandsen, Phys. Rev. **D96**, 095012 (2017), 1709.10473.
- [21] S. Nussinov, Phys. Lett. **165B**, 55 (1985).
- [22] S. M. Barr, R. S. Chivukula, and E. Farhi, Phys. Lett. **B241**, 387 (1990).
- [23] T. Hur and P. Ko, Phys. Rev. Lett. **106**, 141802 (2011), 1103.2571.
- [24] W.-C. Huang, Y.-L. S. Tsai, and T.-C. Yuan, JHEP **04**, 019 (2016), 1512.00229.
- [25] R. Arthur *et al.*, Phys. Rev. **D94**, 094507 (2016), 1602.06559.
- [26] M. E. Peskin, Nucl. Phys. **B175**, 197 (1980).
- [27] J. Preskill, Nucl. Phys. **B177**, 21 (1981).
- [28] G. C. Branco *et al.*, Phys. Rept. **516**, 1 (2012), 1106.0034.
- [29] D. D. Dietrich and M. Jarvinen, Phys. Rev. **D79**, 057903 (2009), 0901.3528.
- [30] M. Fukugita and T. Yanagida, Phys. Lett. **B174**, 45 (1986).
- [31] XENON, E. Aprile *et al.*, Phys. Rev. Lett. **121**, 111302 (2018), 1805.12562.
- [32] ATLAS, G. Aad *et al.*, Phys. Lett. B **796**, 68 (2019), 1903.06248.
- [33] ATLAS, (2018), ATL-PHYS-PUB-2018-044.
- [34] CMS, A. M. Sirunyan *et al.*, JHEP **06**, 120 (2018), 1803.06292.

- [35] CMS, A. M. Sirunyan *et al.*, JHEP **06**, 128 (2018), 1803.11133.
- [36] ATLAS, G. Aad *et al.*, JHEP **03**, 145 (2020), 1910.08447.
- [37] CMS, A. M. Sirunyan *et al.*, JHEP **05**, 033 (2020), 1911.03947.
- [38] ATLAS, G. Aad *et al.*, JHEP **06**, 151 (2020), 2002.11325.
- [39] M. E. Peskin and T. Takeuchi, Phys. Rev. Lett. **65**, 964 (1990).
- [40] M. E. Peskin and T. Takeuchi, Phys. Rev. D **46**, 381 (1992).
- [41] L. Lavoura and J. P. Silva, Phys. Rev. D **47**, 2046 (1993).
- [42] R. Barbieri, A. Pomarol, R. Rattazzi, and A. Strumia, Nucl. Phys. B **703**, 127 (2004), hep-ph/0405040.
- [43] R. Foadi, M. T. Frandsen, and F. Sannino, Phys. Rev. D **77**, 097702 (2008), 0712.1948.
- [44] H. Baer *et al.*, (2013), 1306.6352.
- [45] TLEP Design Study Working Group, M. Bicer *et al.*, JHEP **01**, 164 (2014), 1308.6176.
- [46] M. Ahmad *et al.*, (2015), IHEP-CEPC-DR-2015-01, IHEP-TH-2015-01, IHEP-EP-2015-01.
- [47] T. Appelquist, P. Rodrigues da Silva, and F. Sannino, Phys. Rev. **D60**, 116007 (1999), hep-ph/9906555.
- [48] G. Cacciapaglia and F. Sannino, JHEP **04**, 111 (2014), 1402.0233.
- [49] S. B. Gudnason, C. Kouvaris, and F. Sannino, Phys. Rev. **D74**, 095008 (2006), hep-ph/0608055.
- [50] R. Foadi, M. T. Frandsen, T. A. Rytlov, and F. Sannino, Phys. Rev. **D76**, 055005 (2007), 0706.1696.
- [51] J. A. Harvey and M. S. Turner, Phys. Rev. **D42**, 3344 (1990).