

Lie point symmetries and conservation laws for a Gardner type system

Valter Aparecido Silva Junior

*Instituto Federal de Educação, Ciência e
Tecnologia de São Paulo - IFSP
Avenida Marginal, 585, Fazenda Nossa Senhora Aparecida do Jaguari
13871-298 - São João da Boa Vista/SP, Brasil
E-mail: valtersjunior@ifsp.edu.br*

*Instituto de Física “Gleb Wataghin” - IFGW
Universidade Estadual de Campinas - UNICAMP
Rua Sérgio Buarque de Holanda, 777, Cidade Universitária
13083-859 - Campinas/SP, Brasil
E-mail: valtersjunior@gmail.com*

Abstract

We determine the Lie point symmetries of a Gardner type system and establish its nonlinear self-adjointness. We then construct conservation laws via Ibragimov's Theorem.

2010 AMS Mathematics Classification numbers:

76M60, 58J70, 70G65

Keywords: Gardner type system, Lie point symmetries, nonlinear self-adjointness, conservation laws

1 Introduction

The well-known KdV, mKdV and Gardner equations [4, 13, 15], given respectively by

$$u_t + auu_x + cu_{xxx} = 0, \quad u_t + bu^2u_x + cu_{xxx} = 0$$

and

$$u_t + (au + bu^2)u_x + cu_{xxx} = 0,$$

are special cases of

$$u_t + (au^p + bu^{2p})u_x + cu_{xxx} = 0, \quad a, b, c \in \mathbb{R}, \quad p \in \mathbb{Z}, \quad (1)$$

a recurrent nonlinear evolution equation that, in the framework of hydrodynamics, models internal gravity waves in a density-stratified ocean [2]. Of great physical importance for also describing a variety of wave phenomena in plasma, solid state and quantum physics, its integrability has been investigated by several authors under different approaches. Explicit solutions (including solitons) and a detailed discussion of their properties (like stability) can be found in [6, 14, 22] and references therein.

The situations covered by the above equations tend to be idealized. This occurs, among some reasons, because such models are derived without considering the possible effects of one wave on another. However, since the pioneer works of Hirota and Satsuma [7, 18], where they present a system to study the interactions of two long waves with distinct dispersion relations, an increasing number of coupled KdV equations have been introduced in the literature with a substantially larger and more realistic range of applications. See for example [17, 20, 21]. In particular, starting from [1, 19]

$$\begin{cases} u_t + a[(u^2 + v^2)u]_x + u_{xxx} = 0 \\ v_t + a[(u^2 + v^2)v]_x + v_{xxx} = 0 \end{cases}$$

and

$$\begin{cases} u_t + (u^p v^{p+1})_x + u_{xxx} = 0 \\ v_t + (u^{p+1} v^p)_x + v_{xxx} = 0 \end{cases},$$

we propose in this paper the new class of system

$$\begin{cases} F_1 \equiv u_t + a[(u^p + v^p)u]_x + b[(uv)^p v]_x + cu_{xxx} = 0 \\ F_2 \equiv v_t + a[(u^p + v^p)v]_x + b[(uv)^p u]_x + cv_{xxx} = 0 \end{cases}, \quad (2)$$

a two-component generalization¹ of the equation (1), with the objective of determining the Lie point symmetries (Section 2). Established its nonlinear self-adjointness (Section 3), we then construct conservation laws via the recent Ibragimov's Theorem (Section 4), an extension of the celebrated Noether's Theorem to problems with no variational structure. The main references used are [3, 5, 8, 9, 10, 11, 12, 16].

In what follows, we assume that $\{ac, bc\} \neq \{0\}$ and $p \geq 1$. All constants c_i 's are arbitrary and all functions are smooth.

¹If $u = v$, the system (2) is reduced to equation (1).

2 Lie Point Symmetries Classification

To begin with, we denote

$$f(u, v) \equiv a(u^p + v^p)u + b(uv)^p v, \quad g(u, v) \equiv a(u^p + v^p)v + b(uv)^p u.$$

Applying the standard algorithm presented in [3] and [16], a differential operator

$$X = \mathcal{T}(t, x, u, v) \frac{\partial}{\partial t} + \mathcal{X}(t, x, u, v) \frac{\partial}{\partial x} + \mathcal{U}(t, x, u, v) \frac{\partial}{\partial u} + \mathcal{V}(t, x, u, v) \frac{\partial}{\partial v}$$

generates the Lie point symmetries of the system (2) if the conditions of invariance

$$\begin{aligned} (D_t W^u + \mathcal{T} u_{tt} + \mathcal{X} u_{tx}) + f_u (D_x W^u + \mathcal{T} u_{tx} + \mathcal{X} u_{xx}) + f_v (D_x W^v + \mathcal{T} v_{tx} + \mathcal{X} v_{xx}) + \\ + f_{uu} \mathcal{U} u_x + f_{uv} (\mathcal{V} u_x + \mathcal{U} v_x) + f_{vv} \mathcal{V} v_x + c (D_x^3 W^u + \mathcal{T} u_{txxx} + \mathcal{X} u_{xxxx}) = M F_1 + N F_2 \end{aligned} \quad (3)$$

and

$$\begin{aligned} (D_t W^v + \mathcal{T} v_{tt} + \mathcal{X} v_{tx}) + g_u (D_x W^u + \mathcal{T} u_{tx} + \mathcal{X} u_{xx}) + g_v (D_x W^v + \mathcal{T} v_{tx} + \mathcal{X} v_{xx}) + \\ + g_{uu} \mathcal{U} u_x + g_{uv} (\mathcal{V} u_x + \mathcal{U} v_x) + g_{vv} \mathcal{V} v_x + c (D_x^3 W^v + \mathcal{T} v_{txxx} + \mathcal{X} v_{xxxx}) = P F_1 + Q F_2, \end{aligned} \quad (4)$$

where

$$W^u = \mathcal{U} - \mathcal{T} u_t - \mathcal{X} u_x, \quad W^v = \mathcal{V} - \mathcal{T} v_t - \mathcal{X} v_x,$$

are satisfied for some set of coefficients M , N , P and Q to be determined. From (3) and (4), the determining equations can be written as

$$\begin{aligned} M = \mathcal{U}_u - \mathcal{T}_t, \quad N = \mathcal{U}_v, \quad P = \mathcal{V}_u, \quad Q = \mathcal{V}_v - \mathcal{T}_t, \\ \mathcal{T}_x = \mathcal{T}_u = \mathcal{T}_v = \mathcal{T}_t - 3\mathcal{X}_x = 0, \\ \mathcal{X}_t = \mathcal{X}_u = \mathcal{X}_v = 0, \\ \mathcal{U}_t = \mathcal{U}_x = \mathcal{U}_{uv} = \mathcal{U} - (\mathcal{U}_u u + \mathcal{U}_v v) = 0, \\ \mathcal{V}_t = \mathcal{V}_x = \mathcal{V}_{uv} = \mathcal{V} - (\mathcal{V}_u u + \mathcal{V}_v v) = 0, \\ a[2\mathcal{X}_x + p\mathcal{U}_u + (2-p)\mathcal{V}_u] = 0, \\ b(\mathcal{X}_x + p\mathcal{U}_u) = b\mathcal{U}_v = b\mathcal{V}_u = 0, \\ (1-p)(\mathcal{U}_v + \mathcal{V}_u) = (1-p)(2-p)\mathcal{U}_v = 0, \\ \mathcal{U}_u - \mathcal{V}_v = (2-p)(\mathcal{U}_v - \mathcal{V}_u). \end{aligned}$$

It's easy to see therefore that

$$\begin{aligned} \mathcal{T} = 3c_1 t + c_2, \quad \mathcal{X} = c_1 x + c_3, \\ \mathcal{U} = c_4 v + [c_6 + (2-p)c_4]u, \quad \mathcal{V} = [c_6 + (2-p)c_5]v + c_5 u \end{aligned}$$

with

$$\begin{aligned} a(2c_1 + c_4 + c_5 + pc_6) = 0, \\ bc_4 = bc_5 = b(c_1 + pc_6) = 0, \\ (1-p)(2-p)c_4 = (1-p)(c_4 + c_5) = 0. \end{aligned}$$

Proposition 1. *The Lie point symmetries of the system (2) are summarized in Table 1, where*

$$\begin{aligned} X_1 &= 3t \frac{\partial}{\partial t} + x \frac{\partial}{\partial x} - \frac{1}{p} \left(u \frac{\partial}{\partial u} + v \frac{\partial}{\partial v} \right), \\ Y_1 &= 3t \frac{\partial}{\partial t} + x \frac{\partial}{\partial x} - \frac{2}{p} \left(u \frac{\partial}{\partial u} + v \frac{\partial}{\partial v} \right), \quad X_2 = \frac{\partial}{\partial t}, \quad X_3 = \frac{\partial}{\partial x}, \\ X_4 &= v \left(\frac{\partial}{\partial u} - \frac{\partial}{\partial v} \right), \quad Y_4 = v \frac{\partial}{\partial u} - u \frac{\partial}{\partial v}, \quad X_5 = u \left(\frac{\partial}{\partial u} - \frac{\partial}{\partial v} \right). \end{aligned}$$

$a = 0$	$b = 0$	$ab \neq 0$
X_1, X_2	Y_1, X_2, X_3, X_4 ($p = 1$)	X_2, X_3
X_3	Y_4 ($p = 2$), X_5 ($p = 1$)	

Table 1.

3 Self-Adjointness Classification

For the system (2), the formal Lagrangian is

$$\mathcal{L} = \bar{u}F_1 + \bar{v}F_2.$$

Here \bar{u} and \bar{v} are the new dependent variables (so-called nonlocal variables). Calculated the adjoint equations

$$F_1^* \equiv -\frac{\delta \mathcal{L}}{\delta u} = \bar{u}_t + f_u \bar{u}_x + g_u \bar{v}_x + c \bar{u}_{xxx} = 0$$

and

$$F_2^* \equiv -\frac{\delta \mathcal{L}}{\delta v} = \bar{v}_t + f_v \bar{u}_x + g_v \bar{v}_x + c \bar{v}_{xxx} = 0,$$

where $\delta/\delta u$ and $\delta/\delta v$ are Euler-Lagrange operators, we impose that

$$F_1^*|_{(\bar{u}, \bar{v})=(\varphi, \psi)} = MF_1 + NF_2, \quad F_2^*|_{(\bar{u}, \bar{v})=(\varphi, \psi)} = PF_1 + QF_2. \quad (5)$$

Again M, N, P and Q is a set of coefficients to be determined and

$$\varphi = \varphi(t, x, u, v), \quad \psi = \psi(t, x, u, v) \quad (6)$$

two functions that not vanish simultaneously.

As

$$F_1^*|_{(\bar{u}, \bar{v})=(\varphi, \psi)} = D_t \varphi + f_u D_x \varphi + g_u D_x \psi + c D_x^3 \varphi$$

and

$$F_2^*|_{(\bar{u}, \bar{v})=(\varphi, \psi)} = D_t \psi + f_v D_x \varphi + g_v D_x \psi + c D_x^3 \psi,$$

from (5) it's possible to conclude that $M = \varphi_u$, $N = \varphi_v$, $P = \psi_u$, $Q = \psi_v$ and

$$\begin{aligned}\varphi_t - \psi_t &= \varphi_x - \psi_x = \varphi_u - \psi_v = \varphi_v - \psi_u = 0, \\ \varphi_t + 2a(u+v)\varphi_x &= b\varphi_x = (p-1)\varphi_x = 0, \\ a(p-2)(\varphi_u - \varphi_v) &= b\varphi_v = (p-1)\varphi_v = 0, \\ \varphi_{xx} = \varphi_{xu} = \varphi_{xv} &= \varphi_{uu} = \varphi_{uv} = \varphi_{vv} = 0.\end{aligned}$$

Hence

$$\begin{cases} \varphi = (2ac_1t + c_2)v + (2ac_1t + c_3)u - c_1x + c_4 \\ \psi = (2ac_1t + c_3)v + (2ac_1t + c_2)u - c_1x + c_5 \end{cases},$$

with

$$\begin{aligned}(p-1)c_1 &= bc_1 = (p-1)c_2 = bc_2 = 0, \\ a(p-2)(c_2 - c_3) &= 0.\end{aligned}$$

Proposition 2. *The system (2) is nonlinearly self-adjoint. The substitutions (6) are as follows.*

i) If $a = 0$ or $p = 2$,

$$\varphi = c_3u + c_4, \quad \psi = c_3v + c_5;$$

ii) if $b = 0$ and $p = 1$,

$$\varphi = (2ac_1t + c_2)(v + u) - c_1x + c_4, \quad \psi = (2ac_1t + c_2)(v + u) - c_1x + c_5;$$

iii) in all other cases,

$$\varphi = c_4, \quad \psi = c_5.$$

Remark 1. Actually, the system (2) is quasi self-adjoint. In particular, it's strictly self-adjoint iff $a = 0$ or $p = 2$.

Remark 2. Indeed, any system of the form

$$\begin{cases} u_t + r_x(u, v) + cu_{xxx} = 0 \\ v_t + s_x(u, v) + cv_{xxx} = 0 \end{cases}$$

is strictly self-adjoint whenever $r_v = s_u$.

4 Conservation Laws

According to the theorem demonstrated by Ibragimov [9], the components of the conserved vector $C = (C^t, C^x)$ associated to X , a Lie point symmetry admitted by the system (2), are given by

$$C^t = \varphi W^u + \psi W^v \quad (7)$$

and

$$C^x = \varphi(f_u W^u + f_v W^v) + c(\varphi D_x^2 - D_x \varphi D_x + D_x^2 \varphi) W^u + \psi(g_u W^u + g_v W^v) + c(\psi D_x^2 - D_x \psi D_x + D_x^2 \psi) W^v. \quad (8)$$

In view of Proposition 2, the next result takes into account all the generators of Table 1. In most cases, however, the expressions (7) and (8) lead us to trivial conservation laws or to the vectors

$$C^t = u, \quad C^x = f(u, v) + cu_{xx}$$

and

$$C^t = v, \quad C^x = g(u, v) + cv_{xx}$$

that can be readily obtained from the system (2) by direct integration. Below we only bring the unobvious cases.

Proposition 3. i) *Let $a = 0$. For*

$$\varphi = u, \quad \psi = v,$$

X_1 provides

$$C^t = (p+1)(u^2 + v^2), \\ C^x = 2b(2p+1)(uv)^{p+1} + c(p+1)[(u^2 + v^2)_{xx} - 3(u_x^2 + v_x^2)].$$

ii) *Let $b = 0$.*

ii.a) *Taking*

$$\varphi = \psi = u + v$$

for Y_1 and

$$\varphi = \psi = 2at(u + v) - x$$

for X_2 , we obtain

$$C^t = 3(u + v)^2, \\ C^x = 4a(u + v)^3 + 3c\{[(u + v)^2]_{xx} - 3(u_x + v_x)^2\}$$

when $p = 1$.

ii.b) *Furthermore, for*

$$\varphi = u, \quad \psi = v,$$

Y_1 also provides

$$\begin{aligned}C^t &= 2(u^2 + v^2), \\C^x &= 3a(u^2 + v^2)^2 + 2c[(u^2 + v^2)_{xx} - 3(u_x^2 + v_x^2)]\end{aligned}$$

when $p = 2$.

References

- [1] Alarcon E, Angulo J, Montenegro JF. Stability and instability of solitary waves for a nonlinear dispersive system. *Nonlinear Anal Theory Methods Appl* 1999;36:1015–35.
- [2] Biswas A, Konar S. Soliton perturbation theory for the compound KdV equation. *Int J Theor Phys* 2007;46:237–43.
- [3] Bluman GW, Kumei S. Symmetries and differential equations. New York: Springer; 1989.
- [4] Fu Z, Liu S, Liu S. New kinds of solutions to Gardner equation. *Chaos Solitons Fractals* 2004;20:301–9.
- [5] Gandarias ML. Weak self-adjoint differential equations. *J Phys A Math Theor* 2011;44:262001 (6 pp).
- [6] Hamdi S, Morse B, Halphen B, Schiesser W. Analytical solutions of long nonlinear internal waves: Part I. *Nat Hazards* 2011;57:597–607.
- [7] Hirota R, Satsuma J. Soliton solutions of a coupled Korteweg–de Vries equation. *Phys Lett A* 1981;85:407–8.
- [8] Ibragimov NH. Integrating factors, adjoint equations and Lagrangians. *J Math Anal Appl* 2006;318:742–57.
- [9] Ibragimov NH. A new conservation theorem. *J Math Anal Appl* 2007;333:311–28.
- [10] Ibragimov NH. Quasi-self-adjoint differential equations. *Arch Alga* 2007;4:55–60.
- [11] Ibragimov NH. Nonlinear self-adjointness and conservation laws. *J Phys A Math Theor* 2011;44:432002 (8 pp).
- [12] Ibragimov NH. Nonlinear self-adjointness in constructing conservation laws. *Arch Alga* 2011;7/8:1–90.
- [13] Lin L, Zhu S, Xu Y, Shi Y. Exact solutions of Gardner equations through tanh-coth method. *Appl Math* 2016;7:2374–81.

- [14] Minying T, Ruiqi W, Zhujun J. Solitary waves and their bifurcations of KdV like equation with higher order nonlinearity. *Sci China Ser A Math* 2002;45:1255–67.
- [15] Mohamad MNB. Exact solutions to the combined KdV and MKdV equation. *Math Methods Appl Sci* 1992;15:73–8.
- [16] Olver PJ. Applications of Lie groups to differential equations. New York: Springer; 1986.
- [17] Sakovich, SY. Coupled KdV equations of Hirota–Satsuma type. *J Nonlin Math Phys* 1999;6:255–62.
- [18] Satsuma J, Hirota R. A coupled KdV equation is one case of the four-reduction of the KP hierarchy. *J Phys Soc Jpn* 1982;51:3390–7.
- [19] Triki H, Ismail MS. Solitary wave solutions for a coupled pair of mKdV equations. *Appl Math Comput* 2010;217:1540–8.
- [20] Triki H, Jovanoski Z, Biswas A. Dynamics of two-layered shallow water waves with coupled KdV equations. *Rom Rep Phys* 2014;66:251–61.
- [21] Wright JD, Scheel A. Solitary waves and their linear stability in weakly coupled KdV equations. *Z Angew Math Phys* 2007;58:535–70.
- [22] Zhang WG, Li HW, Bu XS, Bian LY. Orbital stability of solitary waves of compound KdV–type equation. *Acta Math Appl Sin* 2015;31:1033–42.