

On the electron-polaron–electron-polaron scattering and Landau levels in pristine graphene-like quantum electrodynamics

W.B. De Lima*

*Centro Brasileiro de Pesquisas Físicas (CBPF),
Rua Dr. Xavier Sigaud 150 - 22290-180 - Urca - RJ - Brazil.*

O.M. Del Cima[†] and E.S. Miranda[‡]

*Universidade Federal de Viçosa (UFV),
Departamento de Física - Campus Universitário,
Avenida Peter Henry Rolfs s/n - 36570-900 - Viçosa - MG - Brazil.*

The parity-preserving $U(1) \times U(1)$ massless QED₃ is proposed as a pristine graphene-like planar quantum electrodynamics model. The spectrum content, the degrees of freedom, spin, masses and charges of the quasiparticles (electron-polaron, hole-polaron, photon and Néel quasiparticles) which emerge from the model are discussed. The four-fold broken degeneracy of the Landau levels, similar as the one experimentally observed in pristine graphene submitted to high applied external magnetic fields, is obtained. Furthermore, the model exhibits zero-energy Landau level indicating a kind of anomalous quantum Hall effect. The electron-polaron–electron-polaron scattering potentials in s - and p -wave states mediated by photon and Néel quasiparticles are computed and analyzed. Finally, the model foresees that two electron-polarons (s -wave state) belonging to inequivalent \mathbf{K} and \mathbf{K}' points in the Brillouin zone might exhibit attractive interaction, while two electron-polarons (p -wave state) lying both either in \mathbf{K} or in \mathbf{K}' points experience repulsive interaction.

I. INTRODUCTION

The quantum electrodynamics in three dimensional space-time (QED₃) has drawn attention since the groundbreaking works by Schonfeld, Jackiw, Templeton and Deser [1] owing to the viability of taking planar quantum electrodynamics models as theoretical foundation for quasiplanar condensed matter phenomena, such as high- T_c superconductors [2], quantum Hall effect [3], topological insulators [4], topological superconductors [5] and graphene [6]. Since then, planar quantum electrodynamics models have been investigated in many physical arrangements, namely, small (perturbative) and large (non perturbative) gauge transformations, Abelian and non-Abelian gauge groups, fermions families, odd and even under parity, compact space-times, space-times with boundaries, curved space-times, discrete (lattice) space-times, external fields and finite temperatures.

The pristine graphene, a monolayer of pure graphene [6], is a gapless quasibidimensional system behaving like a half-filling semimetal where the quasiparticles charge carriers are described by massless charged Dirac fermions. The electron-electron interactions in graphene [7] include electron-polarons [8] scattering processes [9], where the quasiparticle electron-polaron (or hole-polaron) is formed by a bound state of electron (or hole) and phonon [10].

In this work a pristine graphene-like planar quantum electrodynamics model, the parity-preserving $U(1) \times U(1)$ massless QED₃, is proposed and introduced as follows. In

Section II, the model defined by its discrete and continuous symmetries is presented and its spectrum – degrees of freedom, spin, masses and charges of all the quanta particles content of the model – is discussed. In graphene the interactions among the massless fermion quasiparticles (electron-polaron and hole-polaron) are nonconfining, so the vector meson mediated quasiparticles contained in the model, namely the photon and the Néel quasiparticles, must be massive – massless mediated quanta in three space-time dimensions yield logarithm-type (confining) interaction potentials [11] – consequently the asymptotic states for the massless fermion quasiparticles might be determined. Also, similar effect as the four-fold broken degeneracy of the Landau levels experimentally observed in pristine graphene under high applied external magnetic fields is noticed. Next, in Section III, the s - and p -wave Møller (electron-polaron–electron-polaron) scattering amplitudes are computed and their respective interaction potentials obtained and analyzed. Conclusions and final comments are left to Section IV.

II. THE MODEL

The proposed Lorentz invariant model for a pristine graphene-like planar quantum electrodynamics, the parity-even $U(1)_A \times U(1)_a$ massless QED₃, is defined by the action:

$$S = \int d^3x \left\{ i\bar{\psi}_+ \not{D}\psi_+ + i\bar{\psi}_- \not{D}\psi_- + \right. \\ \left. - \frac{1}{4} F^{\mu\nu} F_{\mu\nu} - \frac{1}{4} f^{\mu\nu} f_{\mu\nu} + \mu \epsilon^{\mu\rho\nu} A_\mu \partial_\rho a_\nu + \right. \\ \left. - \frac{1}{2\alpha} (\partial^\mu A_\mu)^2 - \frac{1}{2\beta} (\partial^\mu a_\mu)^2 \right\}, \quad (1)$$

*Electronic address: wellissonblima@cbpf.br

[†]Electronic address: oswaldo.delcima@ufv.br

[‡]Electronic address: emerson.s.miranda@ufv.br

where $\mathcal{D}\psi_{\pm} \equiv (\not{\partial} + ie\not{A} \pm ig\not{\phi})\psi_{\pm}$, and any object $X \equiv X^{\mu}\gamma_{\mu}$. The coupling constants e (electric charge) and g (chiral charge) carry mass dimension $\frac{1}{2}$. The field strengths, $F_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$ and $f_{\mu\nu} = \partial_{\mu}a_{\nu} - \partial_{\nu}a_{\mu}$, are associated to the electromagnetic field (A_{μ}) and the Néel field (a_{μ}), respectively. The spinors ψ_{+} and ψ_{-} are two kinds of massless fermions, each of them representing electron-polaron (electron-phonon) and hole-polaron (hole-phonon) quasiparticles, where the subscripts $+$ (sublattice **A**) and $-$ (sublattice **B**) are related to their coupling to the Néel gauge field, or alternatively, to the two inequivalent **K** and **K'** points in the Brillouin zone of a monolayer graphene. Also, the gamma matrices are fixed by $\gamma^{\mu} = (\sigma_z, -i\sigma_x, i\sigma_y)$.

A. The symmetries: parity and $U(1) \times U(1)$

The CPT-even action (1) is invariant under the following discrete and continuous symmetries:

1. parity symmetry (P):

$$\begin{aligned} x_{\mu} &\xrightarrow{P} x_{\mu}^P = (x_0, -x_1, x_2), \\ \psi_{\pm} &\xrightarrow{P} \psi_{\pm}^P = -i\gamma^1\psi_{\mp}, \quad \bar{\psi}_{\pm} \xrightarrow{P} \bar{\psi}_{\pm}^P = i\bar{\psi}_{\mp}\gamma^1, \\ A_{\mu} &\xrightarrow{P} A_{\mu}^P = (A_0, -A_1, A_2), \\ a_{\mu} &\xrightarrow{P} a_{\mu}^P = (-a_0, a_1, -a_2). \end{aligned} \quad (2)$$

2. gauge $U(1)_A \times U(1)_a$ symmetry (δ_g):

$$\begin{aligned} \delta_g\psi_{\pm}(x) &= i[\theta(x) \pm \omega(x)]\psi_{\pm}(x), \\ \delta_g\bar{\psi}_{\pm}(x) &= -i[\theta(x) \pm \omega(x)]\bar{\psi}_{\pm}(x), \\ \delta_g A_{\mu}(x) &= -\frac{1}{e}\partial_{\mu}\theta(x), \\ \delta_g a_{\mu}(x) &= -\frac{1}{g}\partial_{\mu}\omega(x). \end{aligned} \quad (3)$$

B. The spectrum: charges, spin, Landau levels, masses and degrees of freedom

The free Dirac equations associated to massless spinors, ψ_{+} and ψ_{-} , derived from the action (1), read:

$$i\not{\partial}\psi_{+} = 0 \quad \text{and} \quad i\not{\partial}\psi_{-} = 0. \quad (4)$$

Thus, expanding the spinor field operators ψ_{+} and ψ_{-} in terms of the c -number plane wave solutions of the Dirac equations, with operator-valued amplitudes, a_{+} , b_{+} , a_{-} and b_{-} (annihilation operators), and a_{+}^{\dagger} , b_{+}^{\dagger} , a_{-}^{\dagger} and b_{-}^{\dagger}

(creation operators), it follows that:

$$\begin{aligned} \psi_{+}(x) &= \int \frac{d^2\vec{p}}{(2\pi)\sqrt{2E}} \{a_{+}(p)u_{+}(p)e^{-ipx} + \\ &\quad + b_{+}^{\dagger}(p)v_{+}(p)e^{ipx}\}; \end{aligned} \quad (5)$$

$$\begin{aligned} \psi_{-}(x) &= \int \frac{d^2\vec{p}}{(2\pi)\sqrt{2E}} \{a_{-}(p)u_{-}(p)e^{-ipx} + \\ &\quad + b_{-}^{\dagger}(p)v_{-}(p)e^{ipx}\}, \end{aligned} \quad (6)$$

where $\bar{\psi}_{\pm} = \psi_{\pm}^{\dagger}\gamma^0$. Consequently, taking into account (4) and (5)-(6), and by adopting $p^{\mu} = (E, p_x, p_y)$ where $E = \sqrt{p_x^2 + p_y^2}$, since $p^{\mu}p_{\mu} = 0$, the wave functions, u_{+} , v_{+} , u_{-} and v_{-} , are given by:

$$u_{+}(p) = \frac{\not{p}}{\sqrt{E}}u'_{+}, \quad v_{+}(p) = \frac{\not{p}}{\sqrt{E}}u'_{-}; \quad (7)$$

$$v_{+}(p) = \frac{-\not{p}}{\sqrt{E}}v'_{+}, \quad v_{-}(p) = \frac{-\not{p}}{\sqrt{E}}v'_{-}, \quad (8)$$

fulfilling the conditions below:

$$u_{+}^{\dagger}(p)u_{+}(p) = v_{+}^{\dagger}(p)v_{+}(p) = 2E; \quad (9)$$

$$u_{-}^{\dagger}(p)u_{-}(p) = v_{-}^{\dagger}(p)v_{-}(p) = 2E; \quad (10)$$

$$\bar{u}_{\pm}(p)u_{\pm}(p) = \bar{v}_{\pm}(p)v_{\pm}(p) = 0, \quad (11)$$

where

$$u'_{+} = u'_{-} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{and} \quad v'_{+} = v'_{-} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (12)$$

From the microcausality conditions for the massless fermions ψ_{+} and ψ_{-} :

$$\{\psi_{\pm}(x), \psi_{\pm}^{\dagger}(y)\}_{x^0=y^0} = \delta^2(\vec{x} - \vec{y}), \quad (13)$$

together with the Dirac equations (4) and the normalization conditions (9)-(10), it stems that:

$$\{a_{\pm}(p), a_{\pm}^{\dagger}(k)\} = \delta^2(\vec{p} - \vec{k}), \quad (14)$$

$$\{b_{\pm}(p), b_{\pm}^{\dagger}(k)\} = \delta^2(\vec{p} - \vec{k}), \quad (15)$$

where all other anticommutators vanish and, for the vacuum state $|0\rangle$, $a_{\pm}(k)|0\rangle = b_{\pm}(k)|0\rangle = 0$.

1. Charges

The quantum operators associated with the internal $U(1)_A \times U(1)_a$ symmetry, namely electric charge (Q_{\pm}) and Néel (chiral) charge (q_{\pm}) operators, are

$$Q_{\pm} = -e \int d^2\vec{x} : \psi_{\pm}^{\dagger}(x)\psi_{\pm}(x) : \quad (16)$$

$$= -e \int d^2\vec{p} \{a_{\pm}^{\dagger}(p)a_{\pm}(p) - b_{\pm}^{\dagger}(p)b_{\pm}(p)\},$$

$$q_{\pm} = \mp g \int d^2\vec{x} : \psi_{\pm}^{\dagger}(x)\psi_{\pm}(x) : \quad (17)$$

$$= \mp g \int d^2\vec{p} \{a_{\pm}^{\dagger}(p)a_{\pm}(p) - b_{\pm}^{\dagger}(p)b_{\pm}(p)\},$$

respectively. Therefore, the electric charges and chiral charges of the asymptotic massless fermion (antifermion) states, $|f_{\uparrow}^{-}\rangle$ ($|f_{\uparrow}^{+}\rangle$) and $|f_{\downarrow}^{-}\rangle$ ($|f_{\downarrow}^{+}\rangle$) read:

$$\begin{aligned} Q_{+}|f_{\uparrow}^{-}\rangle &= -e|f_{\uparrow}^{-}\rangle, & Q_{+}|f_{\downarrow}^{+}\rangle &= +e|f_{\downarrow}^{+}\rangle; \\ Q_{-}|f_{\downarrow}^{-}\rangle &= -e|f_{\downarrow}^{-}\rangle, & Q_{-}|f_{\uparrow}^{+}\rangle &= +e|f_{\uparrow}^{+}\rangle; \end{aligned} \quad (18)$$

$$\begin{aligned} q_{+}|f_{\uparrow}^{-}\rangle &= -g|f_{\uparrow}^{-}\rangle, & q_{+}|f_{\downarrow}^{+}\rangle &= +g|f_{\downarrow}^{+}\rangle; \\ q_{-}|f_{\downarrow}^{-}\rangle &= +g|f_{\downarrow}^{-}\rangle, & q_{-}|f_{\uparrow}^{+}\rangle &= -g|f_{\uparrow}^{+}\rangle; \end{aligned} \quad (19)$$

where

$$|f_{\uparrow}^{-}\rangle = a_{+}^{\dagger}(k)|0\rangle, \quad |f_{\downarrow}^{+}\rangle = b_{+}^{\dagger}(k)|0\rangle, \quad (20)$$

$$|f_{\downarrow}^{-}\rangle = a_{-}^{\dagger}(k)|0\rangle, \quad |f_{\uparrow}^{+}\rangle = b_{-}^{\dagger}(k)|0\rangle, \quad (21)$$

meaning that the creation operators a_{\pm}^{\dagger} and a_{\pm}^{\dagger} create a fermion (electron-polaron) whereas the creation operators b_{\pm}^{\dagger} and b_{\pm}^{\dagger} creates an antifermion (hole-polaron). The electron-polaron and hole-polaron electric and chiral charges are displayed in TAB. I.

2. (Pseudo)spin

Whenever dealing with massless particles in three space-time dimensions, since there is no rest frame, spin cannot be straightforwardly defined, nonetheless, spin (in the generalized sense of a quantum number labelling the representation of the little group [12]) is still a fundamental quantum number. In the present case of massless fermions, it is verified that

$$\left[H_{\pm}^{(0)}, L + \frac{1}{2}\sigma_z \right] = 0, \quad (22)$$

where $H_{\pm}^{(0)} = \vec{\alpha} \cdot \vec{p}$ (with $\vec{\alpha} = \gamma^0 \vec{\gamma}$) is the free Hamiltonian operator associated to the massless spinors, ψ_{+} and ψ_{-} , and $L = xp_y - yp_x$ is the angular momentum operator. Thus, accordingly to (22) it shall be concluded that $\frac{1}{2}\sigma_z$ is the (pseudo)spin operator.

3. Landau levels

The issue of the (pseudo)spin can also be investigated by computing the quantum Landau levels of the model. Therefore, subjecting this pristine graphene-like system to high external and static magnetic field, $B = \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y}$ and $A^{\mu} = (0, \vec{A})$, somehow inducing within the bulk of the system a static magnetic-chiral field, $b = \frac{\partial a_y}{\partial x} - \frac{\partial a_x}{\partial y}$ and $a^{\mu} = (0, \vec{a})$, the hamiltonians for the both massless spinors, ψ_{+} and ψ_{-} , are respectively given by:

$$H_{+} = \vec{\alpha} \cdot (\vec{p} - e\vec{A} - g\vec{a}); \quad (23)$$

$$H_{-} = \vec{\alpha} \cdot (\vec{p} - e\vec{A} + g\vec{a}), \quad (24)$$

whose spectrum, the quantum Landau levels, reads as follows¹:

$$E_{+} = \pm \sqrt{2(eB + gb)} \sqrt{n + \frac{1}{2} \underbrace{\pm \frac{1}{2}}_s}, \quad (25)$$

$$E_{-} = \pm \sqrt{2(eB - gb)} \sqrt{n + \frac{1}{2} \underbrace{\pm \frac{1}{2}}_s}, \quad (26)$$

with n being a non-negative integer number, $n \in \mathbb{N}$ ($n = 0, 1, 2, \dots$), where E_{+} and E_{-} are the Landau levels associated to ψ_{+} (at sublattice **A**) and ψ_{-} (at sublattice **B**) for electron-polarons (if + sign in E_{+} and E_{-}) or hole-polarons (if - sign in E_{+} and E_{-}) and $s = \pm \frac{1}{2}$ are the (pseudo)spin eigenvalues. It shall be stressed that the result displayed in (25)-(26) mimics the four-fold broken degeneracy effect of the Landau levels experimentally observed in pristine graphene under high applied magnetic fields [14]. Also, it yields from the equations (25)-(26) that the lowest Landau level ($n = 0$) appears at $E_{+} = E_{-} = 0$ and accommodates electron-polarons or hole-polarons with only one ($s = -\frac{1}{2}$) pseudospin eigenvalue, signaling a possible anomalous-type quantum Hall effect. All other levels $n \geq 1$ are occupied by electron-polarons or hole-polarons with both ($s = \pm \frac{1}{2}$) pseudospin eigenvalues. Therefore, this implies that for the lowest Landau level $n = 0$ the degeneracy is half of that for any other $n \geq 1$, likewise, all Landau levels ($n \geq 1$) have the same degeneracy (a number of electron-polaron or hole-polaron states with a given energy) but the zero-energy ($n = 0$) Landau level is shared equally by electron-polarons and hole-polarons, *i.e.* depending on the sign of the applied magnetic field there is only sublattice **A** or sublattice **B** states which contribute to the zero-energy (lowest) Landau level.

4. Masses and degrees of freedom

For further computation on the electron-polaron-electron-polaron scattering amplitudes, the tree-level propagators in momenta space for all the fields have to be obtained and this can be achieved by switching off the coupling constants e and g in action (1). Thus, it can be

¹ In [13], the pristine graphene quantum electrodynamics model is introduced adopting the International System of Units, the quantum Landau levels computations are discussed in details, and the results compared with the experimental data of [14].

verified that:

$$\begin{aligned}\Delta_{++}(k) &= \Delta_{--}(k) = i \frac{k}{k^2}; \quad (27) \\ \Delta_{AA}^{\mu\nu}(k) &= -i \left\{ \frac{1}{k^2 - \mu^2} \left(\eta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} \right) + \frac{\alpha}{k^2} \frac{k^\mu k^\nu}{k^2} \right\}, \\ \Delta_{aa}^{\mu\nu}(k) &= -i \left\{ \frac{1}{k^2 - \mu^2} \left(\eta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} \right) + \frac{\beta}{k^2} \frac{k^\mu k^\nu}{k^2} \right\}, \\ \Delta_{Aa}^{\mu\nu}(k) &= \Delta_{aA}^{\mu\nu}(k) = \frac{\mu}{k^2(k^2 - \mu^2)} \epsilon^{\mu\rho\nu} k_\rho. \quad (28)\end{aligned}$$

From the propagators, Δ_{++} and Δ_{--} (27), $\Delta_{AA}^{\mu\nu}$, $\Delta_{aa}^{\mu\nu}$ and $\Delta_{Aa}^{\mu\nu}$ (28), the tree-level unitarity of the model, so as its spectrum, shall be verified by coupling them to conserved external currents $\mathcal{J}_{\Phi_i} = (\mathcal{J}_+, \mathcal{J}_-, \mathcal{J}_A^\mu, \mathcal{J}_a^\mu)$ compatible with the symmetries of the model. Thereby, the current-current transition amplitudes in momentum space are written as $\mathcal{A}_{\Phi_i \Phi_j} = \mathcal{J}_{\Phi_i}^*(k) \langle \Phi_i(k) \Phi_j(k) \rangle \mathcal{J}_{\Phi_j}(k)$. Furthermore, picking up the imaginary part of the residues of the current-current amplitudes ($\mathcal{A}_{\Phi_i \Phi_j}$) at the poles, it can be verified the necessary conditions for unitarity of the S -matrix at the tree-level – positive imaginary part of the residues of the transition amplitudes, $\Im \text{Res } \mathcal{A}_{\Phi_i \Phi_j} > 0$ at all the poles – besides the degrees of freedom counting of all the quantum fields presented in the model, $\Phi_i = (\psi_+, \psi_-, A_\mu, a_\mu)$. Briefly, it has been concluded [15] that the two spinors, ψ_+ and ψ_- , hold two degrees of freedom – the electron-polaron $|f_\uparrow^- \rangle$ (u_+) and the hole-polaron $|f_\downarrow^+ \rangle$ (v_+) associated to the spinor ψ_+ , and the electron-polaron $|f_\downarrow^- \rangle$ (u_-) and the hole-polaron $|f_\uparrow^+ \rangle$ (v_-) associated to the spinor ψ_- . Moreover, the gauge fields, the electromagnetic field (A_μ) and the Néel field (a_μ), carry each one two massive degrees of freedom with mass μ . Also, the single massless mode in $\Delta_{Aa}^{\mu\nu}$ (28) presented in the interaction sector does not propagate, it decouples. Summarizing all results presented above, it is concluded that the parity-even $U(1)_A \times U(1)_a$ massless QED₃, a pristine graphene-like planar quantum electrodynamics model, is free from tachyons and ghosts at the classical level. Notwithstanding, to guarantee the unitarity at tree-level, it is still necessary to investigate the behaviour of the scattering cross sections by testing the fulfillment of the Froissart-Martin bound [16] in the ultraviolet regime, as also in the infrared limit due to the presence of massless quantum fermions.

state	wave function	electric charge	chiral charge	quasiparticle
$ f_\uparrow^- \rangle$	u_+	$-e$	$-g$	electron-polaron
$ f_\downarrow^- \rangle$	u_-	$-e$	$+g$	electron-polaron
$ f_\downarrow^+ \rangle$	v_+	$+e$	$+g$	hole-polaron
$ f_\uparrow^+ \rangle$	v_-	$+e$	$-g$	hole-polaron

TABLE I: The electron-polaron and hole-polaron electric and chiral charges.

III. THE MØLLER SCATTERING: ELECTRON-POLARON-ELECTRON-POLARON

In order to calculate the scattering amplitudes, so as to obtain the scattering potentials, use has been made of the vertex Feynman rules associated to the interaction vertices $-e\bar{\psi}_\pm A\psi_\pm$ and $\mp g\bar{\psi}_\pm a\psi_\pm$: $\Upsilon_{\pm\pm}^\mu = ie\gamma^\mu$ and $v_{\pm\pm}^\mu = \pm ig\gamma^\mu$, respectively.



FIG. 1: e^- -polaron- e^- -polaron (Møller) t -channel scattering mediated by electromagnetic (A_μ) and Néel (a_μ) quantum fields.

The t -channel in s - and p -wave states of electron-polaron-electron-polaron scattering amplitudes owing to electromagnetic and Néel quanta exchange (see FIG. 1) are given by:

$$-i\mathcal{M}_{\pm A \mp} = \bar{u}_\pm(p'_1) [\Upsilon_{\pm\pm}^\mu] u_\pm(p_1) \Delta_{\mu\nu}^{AA}(k) \bar{u}_\mp(p'_2) [\Upsilon_{\mp\mp}^\nu] u_\mp(p_2), \quad (29)$$

$$-i\mathcal{M}_{\pm a \mp} = \bar{u}_\pm(p'_1) [v_{\pm\pm}^\mu] u_\pm(p_1) \Delta_{\mu\nu}^{aa}(k) \bar{u}_\mp(p'_2) [v_{\mp\mp}^\nu] u_\mp(p_2), \quad (30)$$

$$-i\mathcal{M}_{\pm A \pm} = \bar{u}_\pm(p'_1) [\Upsilon_{\pm\pm}^\mu] u_\pm(p_1) \Delta_{\mu\nu}^{AA}(k) \bar{u}_\pm(p'_2) [\Upsilon_{\pm\pm}^\nu] u_\pm(p_2), \quad (31)$$

$$-i\mathcal{M}_{\pm a \pm} = \bar{u}_\pm(p'_1) [v_{\pm\pm}^\mu] u_\pm(p_1) \Delta_{\mu\nu}^{aa}(k) \bar{u}_\pm(p'_2) [v_{\pm\pm}^\nu] u_\pm(p_2), \quad (32)$$

with $\mathcal{M}_{\pm A \mp}$ and $\mathcal{M}_{\pm a \mp}$ being the scattering amplitudes in s -wave state, whereas $\mathcal{M}_{\pm A \pm}$ and $\mathcal{M}_{\pm a \pm}$ being those in p -wave state.

The three-momenta configuration of the two scattered massless electron-polarons in the center of momenta (CM) reference frame; the incoming momenta, p_1 and p_2 ; the outgoing momenta, p'_1 and p'_2 , as well as the momentum transfer, k , are defined below

$$p_1 = (E, p, 0), \quad p'_1 = (E, p \cos \varphi, p \sin \varphi); \quad (33)$$

$$p_2 = (E, -p, 0), \quad p'_2 = (E, -p \cos \varphi, -p \sin \varphi); \quad (34)$$

$$k = p_1 - p'_1 = (0, p(1 - \cos \varphi), -p \sin \varphi) = (0, \vec{k}), \quad (35)$$

where φ is the CM scattering angle, defined as the angle among the directions in the CM frame of the two incoming (initial state) and outgoing (final state) massless electron-polarons.

Assuming the momenta configuration above (33)-(35) and taking into consideration the conditions (7)-(12) on the wave functions u_+ and u_- , the total s - and p -wave Møller scattering amplitudes can be obtained from the partial ones (29)-(32), $\mathcal{M}_{\pm\mp}$ ($|\uparrow\rangle + |\downarrow\rangle \rightarrow |\uparrow\rangle + |\downarrow\rangle$) and $\mathcal{M}_{\pm\pm}$ ($|\uparrow\rangle + |\uparrow\rangle \rightarrow |\uparrow\rangle + |\uparrow\rangle$ or $|\downarrow\rangle + |\downarrow\rangle \rightarrow |\downarrow\rangle + |\downarrow\rangle$), then it

follows that:

$$\mathcal{M}_{\pm\mp} = -(e^2 - g^2) \left[\left(\frac{s-u}{t-\mu^2} \right) - (u \leftrightarrow t) \right] ; \quad (36)$$

$$\mathcal{M}_{\pm\pm} = (e^2 + g^2)e^{\pm i\varphi} \left[\left(\frac{s-u}{t-\mu^2} \right) + (u \leftrightarrow t) \right] , \quad (37)$$

where, s , t and u are the Lorentz invariant Mandelstam variables evaluated at the CM frame:

$$\begin{aligned} s &= (p_1 + p_2)^2 = 4E^2 , \\ t &= (p_1 - p'_1)^2 = -2p^2(1 - \cos \varphi) = -4p^2 \sin^2 \left(\frac{\varphi}{2} \right) , \\ u &= (p_1 - p'_2)^2 = -2p^2(1 + \cos \varphi) = -4p^2 \cos^2 \left(\frac{\varphi}{2} \right) . \end{aligned}$$

A. The scattering potentials

The two-particle interaction potential for two distinguishable electron-polarons (fermions), 1 and 2, in the tree approximation [22] at the CM frame, reads:

$$V(\vec{r}) = \int \frac{d^2\vec{k}}{(2\pi)^2} e^{i\vec{k}\cdot\vec{r}} \beta_1 \beta_2 F(\vec{k}) , \quad (38)$$

with the product $\beta_1 \beta_2$ being a spinorial factor in the space of the electron-polarons 1 and 2, where $\beta_1 = \gamma_1^0$ and $\beta_2 = \gamma_2^0$. Also, in addition to that, taking into account only the t -channel part of the electron-polaron–electron-polaron total scattering amplitude (\mathcal{M}) evaluated at the CM frame, it follows that

$$\mathcal{M} = \bar{u}_1(p'_1) \bar{u}_2(p'_2) F(k) u_1(p_1) u_2(p_2) . \quad (39)$$

Moreover, in accordance to (29)-(32), (38) and (39), the scattering potentials among two electron-polarons, firstly, in s -wave state – one situated at \mathbf{K} point and the other at \mathbf{K}' point in the Brillouin zone – and secondly, in p -wave state – the both located at either \mathbf{K} point or \mathbf{K}' point in the Brillouin zone – mediated by photon and Néel quasiparticles, can be respectively written as:

$$V_s(r) = (1 - \vec{\alpha}_1 \cdot \vec{\alpha}_2) \frac{(e^2 - g^2)}{2\pi} K_0(\mu r) , \quad (40)$$

$$V_p(r) = (1 - \vec{\alpha}_1 \cdot \vec{\alpha}_2) \frac{(e^2 + g^2)}{2\pi} K_0(\mu r) , \quad (41)$$

where $\vec{\alpha} = \gamma^0 \vec{\gamma}$. Thereafter, from (41) it is concluded that, regardless the values of the electromagnetic (e) and the chiral coupling (g) constants, the electron-polaron–electron-polaron interaction in p -wave state ($|\mathbf{K}\rangle + |\mathbf{K}'\rangle$ or $|\mathbf{K}'\rangle + |\mathbf{K}\rangle$) is invariably repulsive. Nevertheless, drawing attention to (40) it takes in evidence about the possibility of attractive electron-polaron–electron-polaron interaction in s -wave state ($|\mathbf{K}\rangle + |\mathbf{K}'\rangle$) provided $g^2 > e^2$. Notwithstanding, in spite of attractive potential be a necessary condition, although not a sufficient one, for the existence of s -wave (\mathbf{K} - \mathbf{K}') massless bipolarons bound

states, it still remains to verify relativistic conditions similar as the non relativistic massive case [17] – where the non relativistic s -wave attractive scattering potential [18] satisfies the Kato condition [19], the Newton-Setô and the Bargmann upper bounds [20, 21] – in which s -wave (\mathbf{K} - \mathbf{K}') massive bipolarons bound states might be arisen.

It shall be also mentioned that the presence of Breit-Darwin-type term $\vec{\alpha}_1 \cdot \vec{\alpha}_2$ in (40) and (41), where in the non-zero gap (mass-gap) graphene-type [17] is of order $\frac{v_1 v_2}{c^2}$ (in real graphene units $\frac{v_1 v_2}{v_F^2}$) and can be therefore neglected at low energies (low speeds), here in the pristine (gapless) graphene-type case, wherein the electron-polarons and hole-polarons are massless, cannot. In summary, it has yet to be investigated in relativistic regime if the attractive ($g^2 > e^2$) s -wave state potential (40) favours massless bipolarons, namely, electron-polaron–electron-polaron or hole-polaron–hole-polaron s -wave bound states. Furthermore, in order to avoid the continuum dissolution problem [22], the Dirac equation for the wave function (Ψ) associated to a two-particle-interaction system, $H_D \Psi = E \Psi$, has to be rewritten as:

$$H_D \Psi = H_{D1} \Psi + H_{D2} \Psi + V_{++} \Psi = E \Psi , \quad (42)$$

where $V_{++} = \Lambda_{++} V \Lambda_{++}$, with $\Lambda_{++} = \Lambda(1) \Lambda(2)$ being the product of Casimir-type positive energy projection operators $\Lambda(i) = \frac{1}{2} (\mathbb{I} + \frac{H_{Di}}{E})$ with $H_{Di} = \vec{\alpha}_i \cdot \vec{p}_i$ ($i = 1, 2$). Besides that, bearing in mind that at the CM frame $\vec{p}_1 = -\vec{p}_2 \equiv \vec{p}$, thus

$$\vec{\alpha}_1 \cdot \vec{p} \Psi - \vec{\alpha}_2 \cdot \vec{p} \Psi + \Lambda_{++} V \Lambda_{++} \Psi = E \Psi . \quad (43)$$

Finally, the question if whether or not, whenever $g^2 > e^2$, the attractive s -wave state potential (40) will favour s -wave massless bipolarons (two-fermion bound states) shall be definitively answered by investigating into details the Dirac equation (43), together with necessary and sufficient conditions which guarantee relativistic two-particle bound states [23].

IV. CONCLUSIONS

The model proposed, a gapless pristine graphene-like planar quantum electrodynamics model, the parity-preserving $U(1) \times U(1)$ massless QED₃, exhibits two-fermion scattering short range non confining potentials originated by two massive vector-mediated quanta, the photon (electric charge) and the Néel (chiral charge) quasiparticle, both stemming from the gauging of the $U(1) \times U(1)$ global symmetry. At the tree-level, the absence in the spectrum of tachyons ($k^2 < 0$) and ghosts ($\langle \psi | \psi \rangle < 0$) assures respectively, causality and unitarity, at this level. Additionally, the charges of the quasiparticles (electron-polaron, hole-polaron, photon and Néel quasiparticles), their masses, degrees of freedom and (pseudo)spin are determined and discussed. As a by-product, it is obtained the four-fold broken degeneracy of the Landau levels, reminding those experimentally

observed in pristine graphene subjected to high external magnetic fields [14], moreover, the system presents zero-energy Landau level suggesting a kind of anomalous quantum Hall effect – detailing results and discussions shall appear further [13].

The p -wave state fermion–fermion (or antifermion–antifermion) scattering potential shows to be repulsive (41) whatever the values of the electric (e) and chiral (g) charges. Nevertheless, for s -wave scattering of fermion–fermion (or antifermion–antifermion), the interaction potential (40) might be attractive provided $g^2 > e^2$. In summary, if two electron-polarons (or hole-polarons) lie in the inequivalent \mathbf{K} and \mathbf{K}' points in the Brillouin zone the interaction might be attractive, otherwise the interaction is always repulsive if those two electron-polarons (or hole-polarons) rest both either in \mathbf{K} or in \mathbf{K}' points.

In view of possible applications of this quantum electrodynamics three space-time dimensional model to pristine (gapless) graphene, or any other planar system, the orders of magnitude of some theoretical parameters have to be estimated. The typical energy-scale in graphene – for instance $E = v_F |\vec{p}|$, $C_s = \frac{1}{2\pi}(e^2 - g^2)$ or $C_p = \frac{1}{2\pi}(e^2 + g^2)$ – is around meV [6], while the length-scale interaction $\lambda = \frac{2\pi\hbar}{\mu c}$ – the reduced Compton wavelength of the quantum-mediated photon and Néel massive quasiparticles – is orders of magnitude in nm [24].

To end this conclusions, it has already been in progress

the prove whether the attractive s -wave state potential (40), provided $g^2 > e^2$, could favour s -wave massless bipolarons [23]. The possible emergence of such a kind of Cooper-type electron-polaron–electron-polaron (hole-polaron–hole-polaron) condensate draws attention to superconductivity in graphene [25].

Acknowledgments

The authors thank J.A. Helayël-Neto and D.H.T. Franco for helpful comments and suggestions. O.M.D.C. dedicates this work to his father (Oswaldo Del Cima, *in memoriam*), mother (Victoria M. Del Cima, *in memoriam*), daughter (Vittoria) and son (Enzo). In spite of the heavy budget cuts on science, scientific research and education promoted this year by the Brazilian Ministry of Education (MEC), CAPES-Brazil is acknowledged for invaluable financial help. FAPEMIG is unworthy owing to the extinction of the scientific initiation fellowship program for undergraduate students executed by the Minas Gerais State government.

Author contribution statement

All authors have been contributed equally.

-
- [1] J.F. Schonfeld, Nucl. Phys. B185 (1981) 157; R. Jackiw and S. Templeton, Phys. Rev. D23 (1981) 2291; S. Deser, R. Jackiw and S. Templeton, Ann. Phys. (NY) 140 (1982) 372, Phys. Rev. Lett. 48 (1982) 975 and Ann. Phys. (NY) 281 (2000) 409.
- [2] M. Franz, Z. Tešanović and O. Vafek, Phys. Rev. B66 (2002) 054535; I.F. Herbut, Phys. Rev. B66 (2002) 094504; H.R. Christiansen, O.M. Del Cima, M.M. Ferreira Jr and J.A. Helayël-Neto, Int. J. Mod. Phys. A18 (2003) 725.
- [3] R.B. Laughlin, Phys. Rev. Lett. 50 (1983) 1395; A.M.J. Schakel, Phys. Rev. D43 (1991) 1428; A. Raya and E.D. Reyes, J. Phys. A: Math. Theor. 41 (2008) 355401.
- [4] M.Z. Hasan and C.L. Kane, Rev. Mod. Phys. 82 (2010) 3045; C.L. Kane and J.E. Moore, Physics World 24 (2011) 32; J.M. Fonseca, W.A. Moura-Melo and A.R. Pereira, J. Appl. Phys. 111 (2012) 064913.
- [5] M. Leijnse and K. Flensberg, Semicond. Sci. Technol. 27 (2012) 124003; S. Yonezawa, arXiv:1604.07930v3, AAPPS Bulletin 26 No.3 (2016) 3; M. Sato and Y. Ando, Rep. Prog. Phys. 80 (2017) 076501.
- [6] V.P. Gusynin, V.A. Miransky and I.A. Shovkovy, Phys. Rev. Lett. 73 (1994) 3499; K.S. Novoselov, A.K. Geim, S.V. Morozov, D. Jiang, M.I. Katsnelson, I.V. Grigorieva, S.V. Dubonos, A.A. Firsov, Nature 438 (2005) 197; V.P. Gusynin, S.G. Sharapov and J.P. Carbotte, Int. J. Mod. Phys. B21 (2007) 4611; R. Jackiw and S.-Y. Pi, Phys. Rev. Lett. 98 (2007) 266402; R. Jackiw and S.-Y. Pi, Phys. Rev. B78 (2008) 132104; M.I. Katsnelson and K.S. Novoselov, Solid State Comm. 143 (2007) 3; A.H. Castro Neto, F. Guinea, N.M.R. Peres, K.S. Novoselov and A.K. Geim, Rev. Mod. Phys. 81 (2009) 109; A. Shytov, M. Rudner, N. Gu, M.I. Katsnelson and L. Levitov, Solid State Commun. 149 (2009) 1087; V.P. Gusynin, “Graphene and Quantum Electrodynamics”, talk given at the International Jubilee Seminar “Current problems in Solid State Physics” (2011), Kharkov, Ukraine; E.M.C. Abreu, M.A. De Andrade, L.P.G. De Assis, J.A. Helayël-Neto, A.L.M.A. Nogueira and R.C. Paschoal, J. High Energy Phys. 1105 (2011) 001; I. Fialkovsky and D.V. Vassilevich, Eur. Phys. J. B85 (2012) 384.
- [7] G.Q. Hai, L. Cândido, B.G.A. Brito and F.M. Peeters, J. Phys. Commun. 2 (2018) 035017.
- [8] C. Chen, J. Avila, S. Wang, Y. Wang, C. Shen, R. Yang, B. Nosarzewski, T.P. Devreux, G. Zhang and M.C. Asensio, Nano Lett. 18 (2018) 1082.
- [9] J.T. Devreese, Z. Phys. B 104 (1997) 601; J.T. Devreese and A.S. Alexandrov, Rep. Prog. Phys. 72 (2009) 066501; D.A. Siegel, C. Hwang, A.V. Fedorov and A. Lanzara, New J. Phys. 14 (2012) 095006; J.C. Johannsen, S. Ulstrup, M. Bianchi, R. Hatch, D. Guan, F. Mazzola, L. Hornekær, F. Fromm, C. Raidel, T. Seyller and P. Hofmann, J. Phys.: Condens. Matter 25 (2013) 094001; C. Chen, J. Avila, E. Frantzeskakis, A. Levy and M.C. Asensio, Nature Commun. 6 (2015) 8585.
- [10] L.D. Landau, Phys.Z.Sowjetunion 3 (1933) 644.
- [11] P. Maris, Phys. Rev. D52 (1995) 6087.
- [12] B. Binegar, J. Math. Phys. 23 (1982) 1511.
- [13] W.B. De Lima, O.M. Del Cima and E.S. Miranda, “Landau-level four-fold splitting in pristine graphene

- quantum electrodynamics”, in preparation.
- [14] Y. Zhang, Z. Jiang, J. P. Small, M.S. Purewal, Y.-W. Tan, M. Fazlollahi, J.D. Chudow, J.A. Jaszczak, H.L. Stormer and P. Kim, *Phys. Rev. Lett.* 96 (2006) 136806.
- [15] W.B. De Lima, “Møller scattering of massless fermions in a parity-even Maxwell-Chern-Simons QED₃”, M.Sc. Thesis - UFV - (2019).
- [16] M. Chaichian, J. Fischer, Yu.S. Vernov, *Nucl. Phys.* B383 (1992) 152; O.M. Del Cima, *Mod. Phys. Lett.* A9 (1994) 1695; K. Chadan, N.N. Khuri, A. Martin, T.T. Wu, *Phys. Rev.* D58 (1998) 025014.
- [17] O.M. Del Cima and E.S. Miranda, *Eur. Phys. J.* B91 (2018) 212.
- [18] W.B. De Lima, O.M. Del Cima, D.H.T. Franco and B.C. Neves, “On the two-dimensional Schrödinger operator with an attractive potential of the Bessel-Macdonald type”, arXiv:1810.02008.
- [19] T. Kato, “Perturbation theory for linear operators” (Springer-Verlag, Heidelberg, 1976).
- [20] N. Setô, *Publ. of RIMS* 9 (1974) 429; B. Simon, *Ann. Phys. (NY)* 97 (1976) 279; R.G. Newton, “Scattering Theory of Waves and Particles”, (Springer-Verlag, New York, 1982).
- [21] V. Bargmann, *Proc. Nat. Acad. Sci. U.S.A.* 38 (1952) 961; B. Simon, “On the number of bound states of two-body Schrödinger operators: A review”, in *Studies in Mathematical Physics, “Essays in Honour of Valentine Bargmann”* (Princeton University Press, Princeton, 1976) 305.
- [22] G. Brown and D.G. Ravenhall, *Proc. R. Soc. London* A208 (1951) 552; J. Sucher, *Phys. Rev.* A22 (1980) 348; *Phys. Rev. Lett.* 55 (1985) 1033; “Potentials from field theory: non-uniqueness, gauge dependence and all that”, *AIP Conference Proceedings* 189 (1989) 337; “The concept of potential in quantum field theory”, proceedings of “Quantum Systems: New Trends and Methods (QS 94)” (1994), Minsk, Belarus; *Adv. Quant. Chem.* 30 (1998) 433.
- [23] O.M. Del Cima and D.H.T. Franco, “On the stability and the spectrum of the modified Brown-Ravenhall operator in two-dimensional space”, in preparation.
- [24] M. Kim, D. Jeong, G.-H. Lee, Y.-S. Shin, H.-W. Lee and H.-J. Lee, *Sci. Rep.* 5 (2015) 8715.
- [25] S. Ichinokura, K. Sugawara, A. Takayama, T. Takahashi and S. Hasegawa, *ACS Nano* 10 (2016) 2761; Y. Cao, V. Fatemi, S. Fang, K. Watanabe, T. Taniguchi, E. Kaxiras and P. Jarillo-Herrero, *Nature* 556 (2018) 43; Y. Cao, V. Fatemi, A. Demir, S. Fang, S.L. Tomarken, J.Y. Luo, J.D. Sanchez-Yamagishi, K. Watanabe, T. Taniguchi, E. Kaxiras, R.C. Ashoori and P. Jarillo-Herrero, *Nature* 556 (2018) 80.