

# Construction of $s$ -extremal optimal unimodular lattices in dimension 52

Masaaki Harada\*

November 30, 2021

**In memory of Masaaki Kitazume**

## Abstract

An  $s$ -extremal optimal unimodular lattice in dimension 52 is constructed for the first time. This lattice is constructed from a certain self-dual  $\mathbb{F}_5$ -code by Construction A. In addition, as neighbors of the lattice, two more  $s$ -extremal optimal unimodular lattices are constructed.

## 1 Introduction

A (Euclidean) integral lattice  $L$  in dimension  $n$  is *unimodular* if  $L = L^*$ , where  $L^*$  denotes the dual lattice of  $L$ . A unimodular lattice with the largest minimum norm among all unimodular lattices in that dimension is called *optimal* [2]. The largest possible minimum norms for dimension  $n \leq 40$  are given in [2, Table 1]. This work was extended in [10]. For example, the largest minimum norms for dimensions 52 and 54 are 5 (see [10]).

Let  $L$  be an odd unimodular lattice in dimension  $n$ . The *shadow*  $S(L)$  of  $L$  is defined to be  $S(L) = L_0^* \setminus L$ , where  $L_0$  denotes the even sublattice of  $L$  [2]. It was shown in [6] that  $8 \min(L) + 4 \min(S(L)) \leq 8 + n$ , unless  $n = 23$  and  $\min(L) = 3$ , where  $\min(L)$  and  $\min(S(L))$  denote the minimum norms of  $L$  and  $S(L)$ , respectively. A unimodular lattice meeting the above bound with equality is called  *$s$ -extremal*. For minimum norms  $\mu \leq 4$ ,  $s$ -extremal

---

\*Research Center for Pure and Applied Mathematics, Graduate School of Information Sciences, Tohoku University, Sendai 980-8579, Japan. email: mharada@tohoku.ac.jp.

unimodular lattices have been widely studied (see e.g. [6], [7] and the references therein). For minimum norm  $\mu \geq 5$ , only a few  $s$ -extremal unimodular lattices are currently known. More precisely, three nonisomorphic  $s$ -extremal odd unimodular lattices in dimension 48 having minimum norm 5 were constructed in [9], and an  $s$ -extremal optimal unimodular lattice in dimension 54 was constructed in [6]. The aim of this note is to establish the following theorem.

**Theorem 1.** *There are at least three nonisomorphic  $s$ -extremal optimal unimodular lattices in dimension 52.*

For dimension 52, no  $s$ -extremal optimal unimodular lattice was previously known.

## 2 Preliminaries

### 2.1 Unimodular lattices

A (Euclidean) lattice  $L \subset \mathbb{R}^n$  in dimension  $n$  is *unimodular* if  $L = L^*$ , where the dual lattice  $L^*$  of  $L$  is defined as  $\{x \in \mathbb{R}^n \mid (x, y) \in \mathbb{Z} \text{ for all } y \in L\}$  under the standard inner product  $(x, y)$ . A unimodular lattice  $L$  is *even* if the norm  $(x, x)$  of every vector  $x$  of  $L$  is even, and *odd* otherwise. An even unimodular lattice in dimension  $n$  exists if and only if  $n \equiv 0 \pmod{8}$ , while an odd unimodular lattice exists for every dimension. The *minimum norm*  $\min(L)$  of a unimodular lattice  $L$  is the smallest norm among all nonzero vectors of  $L$ . The *kissing number* of  $L$  is the number of vectors of minimum norm in  $L$ . Two lattices  $L$  and  $L'$  are *isomorphic*, if there is an orthogonal matrix  $A$  with  $L' = L \cdot A$ , where  $L \cdot A = \{xA \mid x \in L\}$ . The *automorphism group* of  $L$  is the group of all orthogonal matrices  $A$  with  $L = L \cdot A$ .

Let  $L$  be an odd unimodular lattice in dimension  $n$ . The *shadow*  $S(L)$  of  $L$  is defined to be  $S(L) = L_0^* \setminus L$ , where  $L_0$  denotes the even sublattice of  $L$ . Shadows of odd unimodular lattices appeared in [2] (see also [3, p. 440]), and shadows play an important role in the study of odd unimodular lattices. It was shown in [6] that

$$8 \min(L) + 4 \min(S(L)) \leq 8 + n, \quad (1)$$

unless  $n = 23$  and  $\min(L) = 3$ , where  $\min(S(L))$  denotes the minimum norm of  $S(L)$ , that is, the smallest norm among all nonzero vectors of  $S(L)$ . An odd unimodular lattice meeting the bound (1) with equality is called  *$s$ -extremal*.

The *theta series* of an odd unimodular lattice  $L$  and its shadow  $S(L)$  are the formal power series  $\theta_L(q) = \sum_{x \in L} q^{(x,x)}$  and  $\theta_{S(L)}(q) = \sum_{x \in S(L)} q^{(x,x)}$ , respectively. Conway and Sloane [2] showed that when the theta series of an odd unimodular lattice  $L$  in dimension  $n$  is written as:

$$\sum_{j=0}^{\lfloor n/8 \rfloor} a_j \theta_3(q)^{n-8j} \Delta_8(q)^j, \quad (2)$$

the theta series of the shadow  $S(L)$  is written as:

$$\sum_{j=0}^{\lfloor n/8 \rfloor} \frac{(-1)^j}{16^j} a_j \theta_2(q)^{n-8j} \theta_4(q^2)^{8j}, \quad (3)$$

where  $\Delta_8(q) = q \prod_{m=1}^{\infty} (1 - q^{2m-1})^8 (1 - q^{4m})^8$ , and  $\theta_2(q), \theta_3(q), \theta_4(q)$  are the Jacobi theta series [3].

## 2.2 Self-dual $\mathbb{F}_5$ -codes and Construction A

Let  $\mathbb{F}_5$  denote the finite field of order 5. An  $\mathbb{F}_5$ -code  $C$  of length  $n$  is a vector subspace of  $\mathbb{F}_5^n$ . The *dual* code  $C^\perp$  of an  $\mathbb{F}_5$ -code  $C$  of length  $n$  is defined as  $C^\perp = \{x \in \mathbb{F}_5^n \mid x \cdot y = 0 \text{ for all } y \in C\}$ , where  $x \cdot y$  is the standard inner product. A code  $C$  is *self-dual* if  $C = C^\perp$ .

An  $n \times n$  negacirculant matrix has the following form:

$$\begin{pmatrix} r_0 & r_1 & \cdots & r_{n-2} & r_{n-1} \\ -r_{n-1} & r_0 & \cdots & r_{n-3} & r_{n-2} \\ \vdots & \ddots & \ddots & & \vdots \\ -r_2 & & \ddots & \ddots & r_1 \\ -r_1 & -r_2 & \cdots & -r_{n-1} & r_0 \end{pmatrix}.$$

Let  $A$  and  $B$  be  $n \times n$  negacirculant matrices. Consider a matrix of the following form:

$$\begin{pmatrix} I_{2n} & A & B \\ & -B^T & A^T \end{pmatrix}, \quad (4)$$

where  $I_{2n}$  denotes the identity matrix of order  $2n$ , and  $A^T$  denotes the transpose of a matrix  $A$ . The codes with generator matrices of the form (4) are called *four-negacirculant* [8].

Let  $C$  be a self-dual  $\mathbb{F}_5$ -code of length  $n$ . Then the following lattice

$$A_5(C) = \frac{1}{\sqrt{5}} \{(x_1, \dots, x_n) \in \mathbb{Z}^n \mid (x_1 \bmod 5, \dots, x_n \bmod 5) \in C\}$$

is a unimodular lattice in dimension  $n$ . This construction of lattices is well known as Construction A.

### 3 On $s$ -extremal optimal unimodular lattice in dimension 52

The largest minimum norm among unimodular lattices in dimension 52 is 5 [10]. By using (2) and (3), we determine the possible theta series  $\theta_{52}^L(q)$  and  $\theta_{52}^S(q)$  of an optimal odd unimodular lattice and its shadow, respectively, as follows. Since the minimum norm is 5, we have that

$$a_0 = 1, a_1 = -104, a_2 = 3016, a_3 = -22464, a_4 = 19656,$$

in (2). By considering the coefficients of  $q$  and  $q^3$  in (3),  $a_5$  and  $a_6$  are written as:

$$a_5 = -256\alpha, a_6 = 1048576\beta,$$

by using integers  $\alpha$  and  $\beta$ . Hence, we have that

$$\begin{aligned} \theta_{52}^L(q) &= 1 + (157248 - 256\alpha)q^5 + (15462720 + 4096\alpha + 1048576\beta)q^6 \\ &\quad + (729181440 - 21504\alpha - 41943040\beta)q^7 + \dots, \\ \theta_{52}^S(q) &= \beta q + (\alpha - 92\beta)q^3 + (314496 - 68\alpha + 4134\beta)q^5 \\ &\quad + (1458362880 + 2226\alpha - 120888\beta)q^7 + \dots. \end{aligned}$$

The first optimal unimodular lattice in dimension 52 was constructed in [5], and the lattice has theta series  $\theta_{52}^L(q)$  with  $(\alpha, \beta) = (104, 0)$ .

An easy observation on  $\theta_{52}^L(q)$  and  $\theta_{52}^S(q)$  gives the following lemma.

**Lemma 2.** *Let  $L$  be an optimal odd unimodular lattice in dimension 52. Then, the kissing number of  $L$  is 157248 if and only if  $L$  is  $s$ -extremal.*

*Proof.* Follows from the coefficient of  $q^5$  in  $\theta_{52}^L(q)$  and the coefficients of  $q$  and  $q^3$  in  $\theta_{52}^S(q)$ .  $\square$

For an odd unimodular lattice  $L$ , there are cosets  $L_1, L_2, L_3$  of  $L_0$  such that  $L_0^* = L_0 \cup L_1 \cup L_2 \cup L_3$ , where  $L = L_0 \cup L_2$  and  $S(L) = L_1 \cup L_3$ . Two lattices  $L$  and  $L'$  are *neighbors* if both lattices contain a sublattice of index 2 in common. If  $L$  is an odd unimodular lattice in dimension divisible by 4, then there are two unimodular lattices containing  $L_0$ , which are rather than  $L$ , namely,  $L_0 \cup L_1$  and  $L_0 \cup L_3$  (see [4]). We denote the two unimodular neighbors by

$$N_1(L) = L_0 \cup L_1 \text{ and } N_2(L) = L_0 \cup L_3.$$

**Lemma 3.** *If  $L$  is an  $s$ -extremal optimal unimodular lattice in dimension 52, then the lattices  $N_1(L)$  and  $N_2(L)$  are also  $s$ -extremal optimal unimodular lattices in that dimension.*

*Proof.* Since  $L$  has shadow of minimum norm 5, both  $N_1(L)$  and  $N_2(L)$  have minimum norm 5. The shadow of  $N_1(L)$  is  $L_2 \cup L_3$  and the shadow of  $N_2(L)$  is  $L_2 \cup L_1$ . Hence, both  $N_1(L)$  and  $N_2(L)$  have shadows of minimum norm 5, that is,  $s$ -extremal.  $\square$

## 4 Three examples of $s$ -extremal optimal unimodular lattices in dimension 52

By considering four-negacirculant codes, we found a certain self-dual  $\mathbb{F}_5$ -code  $C_{52}$  of length 52. The self-duality of  $C_{52}$  was verified by the MAGMA function `IsSelfDual` (see [1] for MAGMA). The code  $C_{52}$  has generator matrix (4), where the first rows of the negacirculant matrices  $A$  and  $B$  in (4) are as follows:

$$(1, 2, 3, 3, 4, 3, 2, 0, 3, 1, 0, 3, 1) \text{ and } (4, 4, 0, 3, 1, 0, 4, 2, 0, 1, 3, 3, 1),$$

respectively. In addition, we verified that  $A_5(C_{52})$  has minimum norm 5 and kissing number 157248. The minimum norm and the kissing number of  $A_5(C_{52})$  were calculated by the MAGMA functions `Minimum` and `KissingNumber`, respectively. By Lemma 2,  $A_5(C_{52})$  is an  $s$ -extremal optimal unimodular lattice in dimension 52. By Lemma 3,  $N_1(A_5(C_{52}))$  and  $N_2(A_5(C_{52}))$  are  $s$ -extremal optimal unimodular lattices in dimension 52. In addition, we verified by the MAGMA function `IsIsomorphic` that  $A_5(C_{52})$ ,  $N_1(A_5(C_{52}))$  and  $N_2(A_5(C_{52}))$  are nonisomorphic. Hence, we have established Theorem 1.

Finally, we verified that  $A_5(C_{52})$ ,  $N_1(A_5(C_{52}))$  and  $N_2(A_5(C_{52}))$  have automorphism groups of order 52. This was verified by the MAGMA function `AutomorphismGroup`.

**Acknowledgment.** This work was supported by JSPS KAKENHI Grant Number 19H01802.

## References

- [1] W. Bosma, J. Cannon and C. Playoust, The Magma algebra system I: The user language, *J. Symbolic Comput.* **24** (1997), 235–265.
- [2] J.H. Conway and N.J.A. Sloane, A note on optimal unimodular lattices, *J. Number Theory* **72** (1998), 357–362.
- [3] J.H. Conway and N.J.A. Sloane, *Sphere Packing, Lattices and Groups* (3rd ed.), Springer-Verlag, New York, 1999.
- [4] S.T. Dougherty, M. Harada and P. Solé, Shadow lattices and shadow codes, *Discrete Math.* **219** (2000), 49–64.
- [5] P. Gaborit, Construction of new extremal unimodular lattices, *European J. Combin.* **25** (2004), 549–564.
- [6] P. Gaborit, A bound for certain  $s$ -extremal lattices and codes, *Arch. Math.* **89** (2007), 143–151.
- [7] M. Harada, Construction of some unimodular lattices with long shadows, *Int. J. Number Theory* **7** (2011), 1345–1358.
- [8] M. Harada, W. Holzmann, H. Kharaghani and M. Khorvash, Extremal ternary self-dual codes constructed from negacirculant matrices, *Graphs Combin.* **23** (2007), 401–417.
- [9] M. Harada, M. Kitazume, A. Munemasa and B. Venkov, On some self-dual codes and unimodular lattices in dimension 48, *European J. Combin.* **26** (2005), 543–557.

- [10] G. Nebe and N.J.A. Sloane, Unimodular lattices, together with a table of the best such lattices, in *A Catalogue of Lattices*, published electronically at <http://www.math.rwth-aachen.de/~Gabriele.Nebe/LATTICES/>.