

UNIFORM BASES FOR IDEAL ARRANGEMENTS

MAKOTO ENOKIZONO, TATSUYA HORIGUCHI, TAKAHIRO NAGAOKA,
AND AKIYOSHI TSUCHIYA

ABSTRACT. In this paper we introduce and study uniform bases for the ideal arrangements. In particular, explicit uniform bases are presented on each Lie type. Combining the explicit uniform bases with the work of Abe-Horiguchi-Masuda-Murai-Sato, we also obtain explicit presentations of the cohomology rings of regular nilpotent Hessenberg varieties in all Lie types.

CONTENTS

1. Introduction	1
2. Ideal arrangements	3
3. Uniform bases	6
4. Main theorem	10
5. The invertible matrices associated with uniform bases	13
6. Uniform bases for the ideal arrangements in a root subsystem	50
7. The cohomology rings of regular nilpotent Hessenberg varieties	53
References	56

1. INTRODUCTION

In this paper we study bases of the logarithmic derivation modules of the ideal arrangements. Ideal arrangements are subarrangements of the Weyl arrangement which are free arrangements from the work of Abe, Barakat, Cuntz, Hoge, and Terao ([3]). Here, a (central) hyperplane arrangement \mathcal{A} is free if its logarithmic derivation module $D(\mathcal{A})$ (geometrically a polynomial vector fields tangent to \mathcal{A}) is a free module. To prove freeness for the ideal arrangements, they first provided the multiple addition theorem (MAT) and applied MAT to the ideal arrangements. In the proof of MAT they gave a method to construct a basis of the logarithmic derivation module $D(\mathcal{A})$ from a basis of $D(\mathcal{A}')$ for suitable arrangements $\mathcal{A} \supset \mathcal{A}'$.

For a construction of explicit bases of the logarithmic derivation modules of the ideal arrangements for each Lie type, Barakat, Cuntz, and Hoge provided ones for types E and F by computer when the work of [3] was in progress. Also, Terao and Abe worked for types A and B , respectively. In [4] explicit and uniform bases were constructed for types A, B, C, G . Motivated by this, we introduce the notion of uniform bases for the ideal arrangements. We then prove the existence of uniform bases by using the method to construct bases in MAT of [3].

Key words and phrases. ideal arrangements, logarithmic derivation modules, Hessenberg varieties.

We now describe our uniform bases for the ideal arrangements. Let \mathfrak{t} be a real euclidean space and Φ an irreducible root system of rank n on \mathfrak{t}^* , the dual space of \mathfrak{t} . The set of positive roots is denoted by Φ^+ . To $\alpha \in \Phi^+$ we assign the hyperplane $H_\alpha := \ker \alpha$ in \mathfrak{t} . The set of hyperplanes H_α ($\alpha \in \Phi^+$) is called the **Weyl arrangement**. Let $I \subset \Phi^+$ be a lower ideal and the set of hyperplanes H_α ($\alpha \in I$) is called the **ideal arrangement**, denoted by \mathcal{A}_I . Let $\mathcal{R} = \text{Sym}(\mathfrak{t}^*)$ be the symmetric algebra of the dual space \mathfrak{t}^* and we consider the set of \mathbb{R} -derivations of \mathcal{R} , denoted by $\text{Der } \mathcal{R}$. The **logarithmic derivation module** of an ideal arrangement \mathcal{A}_I is defined as

$$D(\mathcal{A}_I) = \{\theta \in \text{Der } \mathcal{R} \mid \theta(\alpha) \in \mathcal{R}\alpha \ (\forall \alpha \in I)\}.$$

By the work of [3] the logarithmic derivation module $D(\mathcal{A}_I)$ is a free \mathcal{R} -module, namely \mathcal{A}_I is a free arrangement. To explain our uniform bases for the ideal arrangement, we need a “good” decomposition of positive roots $\Phi^+ = \coprod_{i=1}^n \Phi_i^+$ (see Section 3 for the details). We fix such a decomposition and define the **Hessenberg function** $h_I : \{1, \dots, n\} \rightarrow \mathbb{Z}_{\geq 0}$ **associated with a lower ideal** I . Note that the Hessenberg function h_I suitably converts a lower ideal I into numerical values. A set of derivations $\{\psi_{i,j} \in \text{Der } \mathcal{R} \mid 1 \leq i \leq n, i \leq j \leq i + |\Phi_i^+|\}$ forms **uniform bases for the ideal arrangements** if derivations $\{\psi_{i,h_I(i)} \mid 1 \leq i \leq n\}$ form an \mathcal{R} -basis of $D(\mathcal{A}_I)$ for any lower ideal I . Our main theorem states that there exist uniform bases for the ideal arrangements (Theorem 4.1). Furthermore, our uniform bases are inductively constructed by invertible matrices P_m for $0 \leq m \leq \text{ht}(\Phi^+)$. More precisely, the initial data $\psi_{1,1}, \dots, \psi_{n,n}$ are the dual basis of the simple roots $\alpha_1, \dots, \alpha_n$ up to a non-zero scalar multiplication. (Note that P_0 is the diagonal matrix whose diagonal entries are the non-zero scalars.) For the next step $\psi_{1,2}, \dots, \psi_{n,n+1}$, each $\psi_{i,i+1}$ is defined by a linear combination of $\alpha_1\psi_{1,1}, \dots, \alpha_n\psi_{n,n}$ as follows:

$$[\psi_{i,i+1}]_{1 \leq i \leq n} = P_1[\alpha_j\psi_{j,j}]_{1 \leq j \leq n}$$

for some invertible matrix P_1 . Proceeding inductively, $\psi_{i,i+m}$ is defined by a linear combination of the products $\alpha_{j,j+m}\psi_{j,j+m-1}$ where P_m denotes the array of the coefficients in the linear combinations. Here, the derivations $\psi_{j,j+m-1}$ are defined in the previous step, and the notation $\alpha_{j,j+m}$ means a positive root in Φ_j^+ with height m . Hence, our uniform bases are determined by the invertible matrices $P_0, P_1, \dots, P_{\text{ht}(\Phi^+)}$ in order. We also prove that when we determined P_0, P_1, \dots, P_{m-1} , the invertible matrix P_m is uniquely determined up to an equivalence defined by the special elementary row operations (Theorem 4.2). Moreover, we give explicitly the invertible matrices $P_0, P_1, \dots, P_{\text{ht}(\Phi^+)}$ on each Lie type (Section 5). In particular, for type D an explicit formula for uniform bases is described. (For types A, B, C, G an explicit formula for uniform bases were given in [4].)

Surprisingly, the logarithmic derivation module of an ideal arrangement determines the cohomology ring of the regular nilpotent Hessenberg variety from the work of [4]. Hessenberg varieties are subvarieties of the full flag variety which was introduced by De Mari, Procesi, and Shayman around 1990 ([7, 6]). This subject is relatively new, and it has been found that geometry, combinatorics, and representation theory interact nicely on Hessenberg varieties (cf. [2]). As one of the interactions, the cohomology ring of a regular nilpotent Hessenberg variety can be described in terms of the logarithmic derivation module of the ideal arrangement ([4]). Originally, explicit presentations of the cohomology rings of regular nilpotent Hessenberg varieties in type A are given by [1] from the point of view of the localization technique. Motivated by this work, in [4] explicit

presentations of the cohomology rings of regular nilpotent Hessenberg varieties for types A, B, C, G are given from the perspective of surprising connection with the logarithmic derivation modules of the ideal arrangements. From our explicit uniform bases for the ideal arrangements, we obtain explicit presentations of the cohomology rings of regular nilpotent Hessenberg varieties in all Lie types (Corollary 7.2) which generalize the result of [8].

The paper is organized as follows. After briefly reviewing some background and terminology on ideal arrangements in Section 2, we introduce the notion of uniform bases and state a key proposition for uniform bases in Section 3. The proofs of the main theorems (Theorems 4.1 and 4.2) for the existence and “uniqueness” of uniform bases are achieved in Section 4. We construct explicit uniform bases on each Lie type in Section 5. In Section 6, we discuss uniform bases for the ideal arrangements in a root subsystem. In particular, uniform bases for the ideal arrangements of types E_6, E_7 can be obtained from that of type E_8 . In Section 7, we explain the connection between ideal arrangements and Hessenberg varieties given in [4] and give explicit presentations of the cohomology rings of regular nilpotent Hessenberg varieties in all Lie types (Corollary 7.2).

Acknowledgements.

We are grateful to Takuro Abe for his historical and valuable comments on bases of the logarithmic derivation modules of the ideal arrangements. The first author was partially supported by JSPS Grant-in-Aid for Research Activity Start-up: 19K23407. The second author was partially supported by JSPS Grant-in-Aid for JSPS Research Fellow: 17J04330. He was also partially supported by JSPS Grant-in-Aid for Young Scientists: 19K14508. The third author was partially supported by JSPS Grant-in-Aid for JSPS Research Fellow: 19J11207. The fourth author was partially supported by JSPS Grant-in-Aid for JSPS Research Fellow: 19J00312. He was also partially supported by JSPS Grant-in-Aid for Young Scientists: 19K14505.

2. IDEAL ARRANGEMENTS

In this section we first refer some terminologies of hyperplane arrangements (for a general reference, see [10]). Then we explain ideal-free theorem proved by [3]. More specifically, we explain the method to construct bases of the logarithmic derivation modules of the ideal arrangements.

Let V be a finite dimensional real vector space. A **hyperplane arrangement** \mathcal{A} in V is a finite set of linear hyperplanes in V . Let $\mathcal{R} = \text{Sym}(V^*)$ be the symmetric algebra of V^* , where V^* is the dual space of V . A map $\theta : \mathcal{R} \rightarrow \mathcal{R}$ is an **\mathbb{R} -derivation** if it satisfies

- (1) θ is \mathbb{R} -linear,
- (2) $\theta(f \cdot g) = \theta(f) \cdot g + f \cdot \theta(g)$ for all $f, g \in \mathcal{R}$.

We denote the set of \mathbb{R} -derivations by $\text{Der } \mathcal{R}$. Note that $\text{Der } \mathcal{R}$ is an \mathcal{R} -module. We can naturally regard an element of V as an \mathbb{R} -derivation, so we have the following identification:

$$\text{Der } \mathcal{R} = \mathcal{R} \otimes V.$$

If we take a basis x_1, \dots, x_n of V^* , then the \mathcal{R} -module $\text{Der } \mathcal{R}$ can be expressed as $\bigoplus_{i=1}^n \mathcal{R} \frac{\partial}{\partial x_i}$ where $\frac{\partial}{\partial x_i}$ denotes the partial derivative with respect to x_i .

A non-zero element $\theta \in \text{Der } \mathcal{R}$ is **homogeneous of (polynomial) degree d** if $\theta = \sum_{k=1}^{\ell} f_k \otimes v_k$ ($f_k \in \mathcal{R}$, $v_k \in V$) and all non-zero f_k 's are of degree d . In this case, we denote $\deg(\theta) = d$ as usual.

For each hyperplane $H \in \mathcal{A}$, we denote the defining linear form of H by $\alpha_H \in V^*$. The **logarithmic derivation module** $D(\mathcal{A})$ of a hyperplane arrangement \mathcal{A} is defined as

$$D(\mathcal{A}) := \{\theta \in \text{Der } \mathcal{R} \mid \theta(\alpha_H) \in \mathcal{R}\alpha_H \ (\forall H \in \mathcal{A})\}.$$

Note that $D(\mathcal{A})$ is a graded \mathcal{R} -submodule, but not a free \mathcal{R} -module in general. We say that a hyperplane arrangement \mathcal{A} is **free with the exponents** $\exp(\mathcal{A}) = (d_1, \dots, d_n)$ if $D(\mathcal{A})$ is a free \mathcal{R} -module with homogeneous basis $\theta_1, \dots, \theta_n$ of degree d_1, \dots, d_n .

Let $\alpha_1, \dots, \alpha_n$ be a basis of V^* . Given derivations $\theta_1, \dots, \theta_n$, we define a matrix $M(\theta_1, \dots, \theta_n)$ by

$$M(\theta_1, \dots, \theta_n) = (\theta_i(\alpha_j))_{1 \leq i, j \leq n}.$$

It is convenient to write $f \doteq g$ for $f, g \in \mathcal{R}$ if $f = cg$ for some $c \in \mathbb{R} \setminus \{0\}$. The following criterion for bases of the logarithmic derivation modules is known.

Theorem 2.1 (Saito's criterion, [11], see also [10]). *Let \mathcal{A} be a hyperplane arrangement in an n -dimensional real vector space V . Let $\theta_1, \dots, \theta_n \in D(\mathcal{A})$ be homogeneous derivations. Then the following arguments are equivalent:*

- (1) $\theta_1, \dots, \theta_n$ form an \mathcal{R} -basis for $D(\mathcal{A})$;
- (2) $\theta_1, \dots, \theta_n$ are linearly independent over \mathcal{R} and $\sum_{i=1}^n \deg \theta_i = |\mathcal{A}|$;
- (3) $\det M(\theta_1, \dots, \theta_n) \doteq (\prod_{H \in \mathcal{A}} \alpha_H)$.

We now explain ideal arrangements which are the main objects of the paper. Let \mathfrak{t} be a (real) euclidean space. Let $\Phi \subset \mathfrak{t}^*$ be an irreducible root system of rank n . We denote the set of positive roots by Φ^+ . We fix simple roots $\alpha_1, \dots, \alpha_n$ and define a partial order \preceq on Φ^+ ; $\alpha \preceq \beta$ if and only if $\beta - \alpha \in \sum_{i=1}^n \mathbb{Z}_{\geq 0} \alpha_i$. A **lower ideal** $I \subset \Phi^+$ is a collection of positive roots such that if $\alpha \in \Phi^+$ and $\beta \in I$ with $\alpha \preceq \beta$, then $\alpha \in I$. The **ideal arrangement** \mathcal{A}_I associated with a lower ideal I is defined as

$$\mathcal{A}_I := \{\ker \alpha \mid \alpha \in I\}.$$

If we take $I = \Phi^+$, then \mathcal{A}_{Φ^+} is called the **Weyl arrangement**. Recall that $\mathcal{R} = \text{Sym}(\mathfrak{t}^*)$ and the logarithmic derivation module of the ideal arrangement \mathcal{A}_I is

$$D(\mathcal{A}_I) = \{\theta \in \text{Der } \mathcal{R} = \mathcal{R} \otimes \mathfrak{t} \mid \theta(\alpha) \in \mathcal{R}\alpha \ (\forall \alpha \in I)\}.$$

The height of a root $\alpha = \sum_{i=1}^n k_i \alpha_i$ is defined by $\text{ht}(\alpha) = \sum_{i=1}^n k_i$. The **height distribution** in I is a sequence (i_1, i_2, \dots, i_m) , where i_j is the number of positive roots of height j in I , and m is the maximum of the height of positive roots in I . Also, the **dual partition** of the height distribution (i_1, i_2, \dots, i_m) in I is the sequence of n elements given by $((0)^{i_0-i_1}, (1)^{i_1-i_2}, \dots, (m-1)^{i_{m-1}-i_m}, (m)^{i_m})$, where $i_0 = n$ and $(i)^j$ denotes the j -copies of i . We denote the dual partition of the height distribution in I by $\mathcal{DP}(I)$.

Theorem 2.2 (Ideal-free theorem, [3, Theorem 1.1]). *Any ideal arrangement \mathcal{A}_I is free with the exponents $\mathcal{DP}(I)$.*

To prove Theorem 2.2, Abe, Barakat, Cuntz, Hoge, and Terao provided the multiple addition theorem (MAT) ([3, Theorem 3.1]). In the proof of MAT they gave a method to construct an \mathcal{R} -basis of $D(\mathcal{A})$ from that of $D(\mathcal{A}')$ for suitable arrangements $\mathcal{A} \supset \mathcal{A}'$. For

the rest of this section, we briefly explain the method to construct an \mathcal{R} -basis under the circumstances of ideal arrangements.

For a lower ideal I we define the height of I by $\text{ht}(I) = \max\{\text{ht}(\alpha) \mid \alpha \in I\}$. Suppose that $\text{ht}(I) = m + 1$ with $m \geq 0$. Let I' be a lower ideal defined by

$$I' := \{\alpha \in I \mid \text{ht}(\alpha) \leq m\}$$

and we set $I \setminus I' = \{\beta_1, \dots, \beta_q\}$. For each $j = 1, \dots, q$, we define a hyperplane arrangement \mathcal{A}_j'' in H_j by $\mathcal{A}_j'' = \{H \cap H_j \mid H \in \mathcal{A}_{I'}\}$ where H_j is a hyperplane defined by the linear function β_j . For each $j = 1, \dots, q$, we fix a map

$$\nu_j : \mathcal{A}_j'' \rightarrow \mathcal{A}_{I'}$$

such that $\nu_j(X) \cap H_j = X$, and define a homogeneous polynomial

$$b_{\nu_j} := \frac{\prod_{H \in \mathcal{A}_{I'}} \alpha_H}{\prod_{X \in \mathcal{A}_j''} \alpha_{\nu_j(X)}} = \frac{\prod_{\alpha \in I'} \alpha}{\prod_{X \in \mathcal{A}_j''} \alpha_{\nu_j(X)}}.$$

Proposition 2.3 ([12], see also [10, p.114, Proposition 4.41]). *For any $\theta \in D(\mathcal{A}_{I'})$,*

$$\theta(\beta_j) \in \mathcal{R}(\beta_j, b_{\nu_j}).$$

Proposition 2.4 ([3, Proposition 4.2]). *One has $\deg(b_{\nu_j}) = m$.*

Let $\theta_1, \dots, \theta_n$ be an \mathcal{R} -basis of $D(\mathcal{A}_{I'})$ with $d_i := \deg(\theta_i)$ such that $d_1 \leq \dots \leq d_{n-p} < d_{n-p+1} = \dots = d_n = m$ for some p . Then, the degree of $\theta_1, \dots, \theta_{n-p}$ is strictly less than m , so we have $\theta_1, \dots, \theta_{n-p} \in D(\mathcal{A}_I)$ from Propositions 2.3 and 2.4. We put $\varphi_i := \theta_{n-i+1}$ for $1 \leq i \leq p$. By Proposition 2.3, for $1 \leq i \leq p$ and $1 \leq j \leq q$ we can write

$$\varphi_i(\beta_j) \equiv c_{ij}^{(\nu_j)} b_{\nu_j} \pmod{\beta_j}$$

for some rational number $c_{ij}^{(\nu_j)}$ ($1 \leq i \leq p$, $1 \leq j \leq q$). The following is a key of the proof of MAT ([3, Theorem 3.1]).

Proposition 2.5 ([3]). *The $(p \times q)$ -matrix $C := (c_{ij}^{(\nu_j)})_{\substack{1 \leq i \leq p \\ 1 \leq j \leq q}}$ has rank q .*

Hence by Proposition 2.5, there exists $P = (p_{ik})_{1 \leq i, k \leq p} \in \text{GL}(p, \mathbb{Q})$ such that

$$PC = \begin{bmatrix} E_q \\ O \end{bmatrix}.$$

We put

$$\psi_i := \sum_{k=1}^p p_{ik} \varphi_k$$

for $1 \leq i \leq p$. One can see that $\beta_1 \psi_1, \dots, \beta_q \psi_q, \psi_{q+1}, \dots, \psi_p \in D(\mathcal{A}_I)$. It is clear that $\theta_1, \dots, \theta_{n-p}, \beta_1 \psi_1, \dots, \beta_q \psi_q, \psi_{q+1}, \dots, \psi_p$ are linearly independent over \mathcal{R} and the sum of their degree is equal to $|\mathcal{A}_I|$, so they form an \mathcal{R} -basis of $D(\mathcal{A}_I)$ from Theorem 2.1.

Theorem 2.6 ([3]). *The derivations $\theta_1, \dots, \theta_{n-p}, \beta_1 \psi_1, \dots, \beta_q \psi_q, \psi_{q+1}, \dots, \psi_p$ form an \mathcal{R} -basis of $D(\mathcal{A}_I)$.*

Hence by Theorem 2.6 we can construct an \mathcal{R} -basis of $D(\mathcal{A}_I)$ for any lower ideal I from that of $D(\mathcal{A}_{I'})$ for the smaller lower ideal $I' \subset I$. Our motivation is to construct these bases uniformly. In next section we introduce the notion of *uniform bases*. Then, we construct uniform bases inductively by using the method of Abe, Barakat, Cuntz, Hoge, and Terao explained above.

3. UNIFORM BASES

In this section we first introduce the notion of Hessenberg functions h_I associated with lower ideals I for all Lie types. Hessenberg functions convert the lower ideals into numerical values. Then we define uniform bases and state a key proposition for uniform bases. In what follows, we frequently use the symbol

$$[n] := \{1, 2, \dots, n\}.$$

Let e_1, e_2, \dots, e_n be the exponents of the Weyl group W . (For the list of exponents e_1, \dots, e_n , see for example [9, p.59 Table 1 and p.81 Theorem 3.19].) We define a decomposition of all positive roots Φ^+ as follows. Let Φ_i^+ be a set of positive roots $\alpha_{i,i+1}, \alpha_{i,i+2}, \dots, \alpha_{i,i+e_i}$ for $1 \leq i \leq n$ such that

$$(3.1) \quad \alpha_{i,i+1} = \alpha_i \quad \text{the simple root}$$

$$(3.2) \quad \alpha_{i,j} \triangleleft \alpha_{i,j+1} \quad \text{for any } j \text{ with } i < j < i + e_i$$

Here, we denote the covering relation by the symbol \triangleleft , namely, $\alpha_{i,j} \triangleleft \alpha_{i,j+1}$ and there is no element $\beta \in \Phi^+$ such that $\alpha_{i,j} \triangleleft \beta \triangleleft \alpha_{i,j+1}$. The sets Φ_i^+ ($1 \leq i \leq n$) give disjoint n maximal chains. In particular, we have a decomposition of the positive roots $\Phi^+ = \coprod_{i=1}^n \Phi_i^+$. Note that such a decomposition is not unique. We fix such a decomposition $\Phi^+ = \coprod_{i=1}^n \Phi_i^+$ and define the **Hessenberg function** $h_I : [n] \rightarrow \mathbb{Z}_{\geq 0}$ **associated with a lower ideal I** by the following formula:

$$(3.3) \quad h_I(i) := \begin{cases} \max\{j \mid \alpha_{i,j} \in I \cap \Phi_i^+\} & \text{if } I \cap \Phi_i^+ \neq \emptyset, \\ i & \text{if } I \cap \Phi_i^+ = \emptyset \end{cases}$$

for $1 \leq i \leq n$.

Example 3.1. In type A_{n-1} we set the exponents $e_i = n - i$ for $i = 1, \dots, n - 1$ and positive roots $\Phi_{A_{n-1}}^+ = \{x_i - x_j \mid 1 \leq i < j \leq n\}$. Let $\alpha_{i,j} = x_i - x_j$ for $1 \leq i < j \leq n$. Then, $\Phi_i^+ = \{\alpha_{i,i+1}, \dots, \alpha_{i,n}\}$ satisfies the conditions (3.1) and (3.2). In this setting, for example, the Hessenberg function¹ $h_I : [n-1] \rightarrow \mathbb{Z}_{\geq 0}$ associated with the lower ideal $I = \{\alpha_1, \dots, \alpha_{n-1}\}$ is given by $h_I(i) = i + 1$ for $i \in [n-1]$.

Definition 3.2. We fix a decomposition $\Phi^+ = \coprod_{i=1}^n \Phi_i^+$ satisfying (3.1) and (3.2). A set of derivations $\{\psi_{i,j} \in \text{Der } \mathcal{R} \mid 1 \leq i \leq n, i \leq j \leq i + e_i\}$ forms **uniform bases for the ideal arrangements** (or simply **uniform bases**) if derivations $\{\psi_{i,h_I(i)} \mid 1 \leq i \leq n\}$ form an \mathcal{R} -basis of $D(\mathcal{A}_I)$ for any lower ideal I .

¹A Hessenberg function h for type A_{n-1} is usually defined on the set $[n]$. More specifically, $h : [n] \rightarrow [n]$ is a Hessenberg function for type A_{n-1} if h is weakly increasing and $h(i) \geq i$ for $i \in [n]$. We have $h(n) = n$ by the definition, so we may omit the n -th value of h .

Noting that $\text{ht}(\alpha_{i,j}) = j - i$, a set $\{h_I(i) - i \mid 1 \leq i \leq n\}$ is the dual partition of the height distribution in I . From this together with Theorem 2.2 we have

$$(3.4) \quad \exp(\mathcal{A}_I) = \{h_I(i) - i \mid 1 \leq i \leq n\}.$$

In particular, we have

$$\text{ht}(I) = \max\{h_I(i) - i \mid 1 \leq i \leq n\}.$$

For each lower ideal I we define a subset Λ_I of $[n]$ as follows:

$$\Lambda_I := \{i \in [n] \mid h_I(i) - i = \text{ht}(I)\}.$$

Let us denote the cardinality of Λ_I by $\lambda_I = |\Lambda_I|$.

Example 3.3. We consider the setting of Example 3.1. Let $n = 5$ and we take

$$I = \{x_1 - x_2, x_1 - x_3, x_2 - x_3, x_3 - x_4, x_3 - x_5, x_4 - x_5\}.$$

Then, the associated Hessenberg function h_I is given by $h_I(1) = 3, h_I(2) = 3, h_I(3) = 5, h_I(4) = 5$ and we have $\Lambda_I = \{1, 3\}$.

For each integer m with $0 \leq m \leq \text{ht}(\Phi^+)$, we define the lower ideal I_m by

$$(3.5) \quad I_m := \{\alpha \in \Phi^+ \mid \text{ht}(\alpha) \leq m\}.$$

We write the Hessenberg function associated with the lower ideal I_m by h_m . For simplicity, we denote Λ_{I_m} and λ_{I_m} by Λ_m and λ_m , respectively. Namely,

$$(3.6) \quad \Lambda_m = \{i \in [n] \mid h_m(i) - i = m\}.$$

We note that

$$\{(i, j) \mid 1 \leq i \leq n \text{ and } i \leq j \leq i + e_i\} = \{(i, i + m) \mid 0 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m\}.$$

Let R be a commutative ring. For two subsets $S, T \subset [n]$ we denote by $M(S, T; R)$ the set of matrices $[a_{s,t}]_{\substack{s \in S \\ t \in T}}$ with entries $a_{s,t} \in R$. We also denote by $\text{GL}(S; R)$ the set of invertible matrices $[a_{s,t}]_{s,t \in S}$ with entries $a_{s,t} \in R$. That is,

$$\begin{aligned} M(S, T; R) &= \{(a_{s,t})_{\substack{s \in S \\ t \in T}} \text{ matrices} \mid a_{s,t} \in R\}, \\ \text{GL}(S; R) &= \{(a_{s,t})_{s,t \in S} \text{ invertible matrices} \mid a_{s,t} \in R\}. \end{aligned}$$

The following is a key proposition.

Proposition 3.4. Assume that derivations $\{\psi_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m\}$ satisfy the following four conditions:

- (1) For any $i \in [n]$, $\psi_{i,i} = p_i \alpha_i^*$ for some non-zero rational number p_i where $\alpha_1^*, \dots, \alpha_n^*$ is the dual basis of the simple roots $\alpha_1, \dots, \alpha_n$;
- (2) For any $m \geq 1$ there exists $P_m \in \text{GL}(\Lambda_m; \mathbb{Q})$ such that

$$[\psi_{i,i+m}]_{i \in \Lambda_m} = P_m [\alpha_{i,i+m} \psi_{i,i+m-1}]_{i \in \Lambda_m};$$

- (3) For any $m \geq 0$, $\{\psi_{i,h_m(i)} \mid i \in [n]\} \subset D(\mathcal{A}_{I_m})$;
- (4) For any $m \geq 0$ and any $(i, j) \in \Lambda_m \times \Lambda_{m+1}$ with $i \neq j$,

$$\psi_{i,i+m}(\alpha_{j,j+m+1}) \in \mathcal{R} \alpha_{j,j+m+1}.$$

Then, derivations $\{\psi_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m\}$ form uniform bases.

Proof. One can see that $\deg(\psi_{i,j}) = j - i$ from the conditions (1) and (2). By (3.4) we have $\sum_{i=1}^n \deg(\psi_{i,h_I(i)}) = |\mathcal{A}_I|$. From this together with Theorem 2.1 it is enough to prove the following two claims:

Claim 1 For any lower ideal I , $\psi_{i,h_I(i)} \in D(\mathcal{A}_I)$ for each $i \in [n]$.

Claim 2 For any lower ideal I , the derivations $\{\psi_{i,h_I(i)} \mid i \in [n]\}$ are linearly independent over \mathcal{R} .

Proof of Claim 1. We prove this by induction on $\text{ht}(I)$. The base case $\text{ht}(I) = 0$ is clear since $I = \emptyset$ and $D(\mathcal{A}_I) = \text{Der}(\mathcal{R})$. Now we assume that $m > 0$ and Claim 1 holds for any lower ideal I' with $\text{ht}(I') = m - 1$. For a lower ideal I with $\text{ht}(I) = m$, we define a lower ideal I' by $I' := I \cap I_{m-1}$. Namely, $I' = \{\alpha \in I \mid \text{ht}(\alpha) \leq m - 1\}$ by the definition (3.5). One can see that

$$h_{I'}(i) = \begin{cases} h_I(i) - 1 & \text{if } i \in \Lambda_I, \\ h_I(i) & \text{if } i \notin \Lambda_I. \end{cases}$$

Case 1 Suppose that $i \notin \Lambda_I$. In this case we have $h_I(i) = h_{I'}(i)$, so we prove that $\psi_{i,h_{I'}(i)} \in D(\mathcal{A}_I)$. Since $\text{ht}(I') = m - 1$, we have $\psi_{i,h_{I'}(i)} \in D(\mathcal{A}_{I'})$ by inductive assumption. Noting that $I = I' \cup \{\alpha_{j,j+m} \mid j \in \Lambda_I\}$, it is enough to prove that $\psi_{i,h_{I'}(i)}(\alpha_{j,j+m}) \in \mathcal{R}\alpha_{j,j+m}$ for any $j \in \Lambda_I$. Let $H_{i,j}$ be the hyperplane defined by a positive root $\alpha_{i,j}$. For each $j \in \Lambda_I$ we define a hyperplane arrangement \mathcal{A}_j'' in $H_{j,j+m}$ by $\mathcal{A}_j'' = \{H \cap H_{j,j+m} \mid H \in \mathcal{A}_{I'}\}$ and we take a map

$$\nu_j : \mathcal{A}_j'' \rightarrow \mathcal{A}_{I'}$$

such that $\nu_j(X) \cap H_{j,j+m} = X$. The homogeneous polynomial

$$b_{\nu_j} := \frac{\prod_{H \in \mathcal{A}_{I'}} \alpha_H}{\prod_{X \in \mathcal{A}_j''} \alpha_{\nu_j(X)}} = \frac{\prod_{\alpha \in I'} \alpha}{\prod_{X \in \mathcal{A}_j''} \alpha_{\nu_j(X)}}$$

has degree $m - 1$ by Proposition 2.4, and we obtain

$$(3.7) \quad \psi_{i,h_{I'}(i)}(\alpha_{j,j+m}) \in \mathcal{R}(\alpha_{j,j+m}, b_{\nu_j})$$

from Proposition 2.3. If $i \notin \Lambda_{I'}$, then we have $\deg(\psi_{i,h_{I'}(i)}) = h_{I'}(i) - i < \text{ht}(I') = m - 1 = \deg(b_{\nu_j})$. Hence by (3.7) we obtain $\psi_{i,h_{I'}(i)}(\alpha_{j,j+m}) \in \mathcal{R}(\alpha_{j,j+m})$ for $i \notin \Lambda_{I'}$. If $i \in \Lambda_{I'}$, then we have $h_{I'}(i) = i + m - 1$. Note that $i \in \Lambda_{I'} \subset \Lambda_{m-1}$ and $j \in \Lambda_I \subset \Lambda_m$. We also have $i \neq j$ because $i \notin \Lambda_I$ and $j \in \Lambda_I$. Thus, it follows from the condition (4) that $\psi_{i,h_{I'}(i)}(\alpha_{j,j+m}) = \psi_{i,i+m-1}(\alpha_{j,j+m}) \in \mathcal{R}(\alpha_{j,j+m})$.

Case 2 Suppose that $i \in \Lambda_I$. Then, we have $h_I(i) = i + m = h_m(i)$. From the condition (3) and $\mathcal{A}_I \subset \mathcal{A}_{I_m}$, we obtain $\psi_{i,h_I(i)} = \psi_{i,h_m(i)} \in D(\mathcal{A}_{I_m}) \subset D(\mathcal{A}_I)$.

Case 1 and Case 2 show Claim 1.

Proof of Claim 2. In order to prove Claim 2, we prove Claim 2' as follows:

Claim 2' For any lower ideal I with $\text{ht}(I) = m$ and any matrix $F = [f_{ij}]_{\substack{i \in \Lambda_I \\ j \in \Lambda_m}} \in M(\Lambda_I, \Lambda_m; \mathcal{R})$ such that row vectors $\mathbf{f}_i = [f_{ij}]_{j \in \Lambda_m}$ ($i \in \Lambda_I$) are linearly independent over \mathcal{R} , we put $[\xi_i^F]_{i \in \Lambda_I} := F[\psi_{i,i+m}]_{i \in \Lambda_m}$. Then, the derivations $\{\xi_i^F \mid i \in \Lambda_I\} \cup \{\psi_{i,h_I(i)} \mid i \notin \Lambda_I\}$ are linearly independent over \mathcal{R} .

We prove Claim 2' by induction on $\text{ht}(I)$. The base case $\text{ht}(I) = 0$ is clear since $I = \emptyset$ and the derivations $\{\psi_{i,i} = p_i \alpha_i^* \mid i \in [n]\}$ are \mathcal{R} -basis of $D(\mathcal{A}_I) = \text{Der}(\mathcal{R})$.

Now we assume that $m > 0$ and Claim 2' holds for any lower ideal I' with $\text{ht}(I') = m - 1$ and any matrix $F' \in M(\Lambda_{I'}, \Lambda_{m-1}; \mathcal{R})$ such that row vectors in F' are linearly independent

over \mathcal{R} . For a lower ideal I with $\text{ht}(I) = m$ and any matrix $F \in M(\Lambda_I, \Lambda_m; \mathcal{R})$ such that row vectors in F are linearly independent over \mathcal{R} , we define the lower ideal $I' := I \cap I_{m-1}$. From the condition (2) we have

$$\begin{aligned} \begin{bmatrix} [\xi_i^F]_{i \in \Lambda_I} \\ [\psi_{i, h_I(i)}]_{i \notin \Lambda_I} \end{bmatrix} &= \begin{bmatrix} F[\psi_{i, i+m}]_{i \in \Lambda_m} \\ [\psi_{i, h_I(i)}]_{i \notin \Lambda_I} \end{bmatrix} = \begin{bmatrix} FP_m[\alpha_{i, i+m} \psi_{i, i+m-1}]_{i \in \Lambda_m} \\ [\psi_{i, h_I(i)}]_{i \notin \Lambda_I} \end{bmatrix} \\ &= \begin{bmatrix} FP_m \text{diag}(\alpha_{i, i+m})[\psi_{i, i+m-1}]_{i \in \Lambda_m} \\ [\psi_{i, h_I(i)}]_{i \notin \Lambda_I} \end{bmatrix} \\ &= \begin{bmatrix} FP_m \text{diag}(\alpha_{i, i+m})[\psi_{i, i+m-1}]_{i \in \Lambda_m} \\ [\psi_{i, i+m-1}]_{i \in \Lambda_{I'} \setminus \Lambda_I} \\ [\psi_{i, h_I(i)}]_{i \notin \Lambda_{I'}} \end{bmatrix} \\ &= \begin{bmatrix} F'[\psi_{i, i+m-1}]_{i \in \Lambda_{m-1}} \\ [\psi_{i, h_I(i)}]_{i \notin \Lambda_{I'}} \end{bmatrix} = \begin{bmatrix} [\xi_i^{F'}]_{i \in \Lambda_{I'}} \\ [\psi_{i, h_I(i)}]_{i \notin \Lambda_{I'}} \end{bmatrix} \end{aligned}$$

where $[\xi_i^{F'}]_{i \in \Lambda_{I'}} := F'[\psi_{i, i+m-1}]_{i \in \Lambda_{m-1}}$ and $F' = [f'_{ij}]_{\substack{i \in \Lambda_{I'} \\ j \in \Lambda_{m-1}}}$ is the matrix given by

$$F' = \underbrace{\left[\begin{array}{c|c} FP_m \text{diag}(\alpha_{i, i+m}) & O \\ \hline O & \delta_{ij} \end{array} \right]_{\Lambda_I} }_{\substack{\Lambda_m \\ \Lambda_{m-1}}} \Bigg\}_{\Lambda_{I'}}$$

More precisely,

$$f'_{ij} = \begin{cases} (i, j)\text{-entry of } [FP_m \text{diag}(\alpha_{i, i+m})] & \text{if } (i, j) \in \Lambda_I \times \Lambda_m, \\ \delta_{ij} & \text{if } (i, j) \in (\Lambda_{I'} \setminus \Lambda_I) \times (\Lambda_{m-1} \setminus \Lambda_m), \\ 0 & \text{otherwise.} \end{cases}$$

It is clear that row vectors in F' are linearly independent over \mathcal{R} . Hence by the inductive assumption, $\{\xi_i^{F'} \mid i \in \Lambda_{I'}\} \cup \{\psi_{i, h_I(i)} \mid i \notin \Lambda_{I'}\}$ are linearly independent over \mathcal{R} , so is $\{\xi_i^F \mid i \in \Lambda_I\} \cup \{\psi_{i, h_I(i)} \mid i \notin \Lambda_I\}$. Applying Claim 2' to $F = [\delta_{ij}]_{\substack{i \in \Lambda_I \\ j \in \Lambda_m}}$, we obtain Claim 2.

Therefore, we proved Claim 1 and Claim 2. This completes the proof. \square

Remark 3.5. Assume that the derivations $\{\psi_{i, j} \in \text{Der}(\mathcal{R}) \mid 1 \leq i \leq n \text{ and } i \leq j \leq i + e_i\}$ satisfy the following conditions:

- (1) For any $i \in [n]$ we can write $\psi_{i, i} = p_i \alpha_i^*$ for some non-zero rational number p_i ;
- (2) For any $m \geq 1$ we can write

$$[\psi_{i, i+m}]_{i \in \Lambda_m} = P_m[\alpha_{i, i+m} \psi_{i, i+m-1}]_{i \in \Lambda_m}$$

for some invertible matrix $P_m \in \text{GL}(\Lambda_m; \mathbb{Q})$;

- (3) For any lower ideal $I \subset \Phi^+$, $\{\psi_{i, h_I(i)} \mid i \in [n]\} \subset D(\mathcal{A}_I)$.

Then the derivations $\{\psi_{i, j} \in \text{Der}(\mathcal{R}) \mid 1 \leq i \leq n \text{ and } i \leq j \leq i + e_i\}$ form uniform bases because the conditions (3) and (4) in Proposition 3.4 mean that $\{\psi_{i, h_I(i)} \mid i \in [n]\} \subset D(\mathcal{A}_I)$ for the special lower ideal $I = I_m, I_m \cup \{\alpha_{j, j+m+1}\}_{\substack{j \in \Lambda_{m+1} \\ j \neq i}}$.

4. MAIN THEOREM

In this section we prove the existence of uniform bases which are inductively constructed by using invertible matrices (Theorem 4.1). Moreover, we prove that the invertible matrices associated with our uniform bases are unique in some sense (Theorem 4.2).

Theorem 4.1. *For arbitrary decomposition $\Phi^+ = \coprod_{i=1}^n \Phi_i^+$ satisfying (3.1) and (3.2), there exist uniform bases $\{\psi_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m\}$ of the following form. The initial data are of the form*

$$\psi_{i,i} = p_i \alpha_i^* \quad (i \in [n])$$

where p_i is arbitrary non-zero rational number and $\alpha_1^*, \dots, \alpha_n^*$ is the dual basis of the simple roots $\alpha_1, \dots, \alpha_n$. For any m with $1 \leq m \leq \text{ht}(\Phi^+)$

$$\psi_{i,i+m} = \sum_{j \in \Lambda_m} p_{ij}^{(m)} \alpha_{j,j+m} \psi_{j,j+m-1} \quad (i \in \Lambda_m)$$

for some rational numbers $p_{ij}^{(m)}$.

Proof. It suffices to construct derivations $\{\psi_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m\}$ satisfying the four conditions in Proposition 3.4. For the construction, we apply the method of Abe, Barakat, Cuntz, Hoge, and Terao explained in Section 2 to the sequence of the lower ideals in (3.5):

$$I_1 \subset I_2 \subset \dots \subset I_{\text{ht}(\Phi^+)}.$$

For m with $0 \leq m < \text{ht}(\Phi^+)$, we construct inductively derivations $\{\psi_{i,i+m}\}_{i \in \Lambda_m}$ and $\{\theta_{i,i+m+1}\}_{i \in \Lambda_{m+1}}$ with $\deg(\psi_{i,j}) = \deg(\theta_{i,j}) = j - i$ so that the derivations $\{\theta_{i,i+m+1} \mid i \in \Lambda_{m+1}\} \cup \{\psi_{i,i+m} \mid i \in \Lambda_m \setminus \Lambda_{m+1}\} \cup \dots \cup \{\psi_{i,i+1} \mid i \in \Lambda_1 \setminus \Lambda_2\}$ form an \mathcal{R} -basis for $D(\mathcal{A}_{I_{m+1}})$ as follows. As the base case, when $m = 0$, for any non-zero rational number p_i we define

$$(4.1) \quad \psi_{i,i} = p_i \alpha_i^* \quad \text{for } i \in \Lambda_0 = [n],$$

and

$$\theta_{i,i+1} = \alpha_{i,i+1} \psi_{i,i} \quad \text{for } i \in \Lambda_1 = [n].$$

For general m with $0 < m < \text{ht}(\Phi^+)$ we proceed inductively as follows. Let $H_{i,j}$ be the hyperplane defined by a positive root $\alpha_{i,j}$. Then we have

$$\mathcal{A}_{I_{m+1}} = \mathcal{A}_{I_m} \cup \{H_{j,j+m+1} \mid j \in \Lambda_{m+1}\}.$$

For each $j \in \Lambda_{m+1}$ we define a hyperplane arrangement \mathcal{A}_j'' in $H_{j,j+m+1}$ by $\mathcal{A}_j'' = \{H \cap H_{j,j+m+1} \mid H \in \mathcal{A}_{I_m}\}$ and we take a map

$$\nu_j : \mathcal{A}_j'' \rightarrow \mathcal{A}_{I_m}$$

such that $\nu_j(X) \cap H_{j,j+m+1} = X$. The homogeneous polynomial

$$b_{\nu_j} = \frac{\prod_{H \in \mathcal{A}_{I_m}} \alpha_H}{\prod_{X \in \mathcal{A}_j''} \alpha_{\nu_j(X)}} = \frac{\prod_{\alpha \in I_m} \alpha}{\prod_{X \in \mathcal{A}_j''} \alpha_{\nu_j(X)}}$$

has degree m by Proposition 2.4, and we obtain

$$\theta_{i,i+m}(\alpha_{j,j+m+1}) \in \mathcal{R}(\alpha_{j,j+m+1}, b_{\nu_j})$$

for $i \in \Lambda_m$ and $j \in \Lambda_{m+1}$ from Proposition 2.3. Hence, we can write

$$(4.2) \quad \theta_{i,i+m}(\alpha_{j,j+m+1}) \equiv c_{ij}^{(\nu_j)} b_{\nu_j} \pmod{\alpha_{j,j+m+1}}$$

for some rational numbers $c_{ij}^{(\nu_j)}$ ($i \in \Lambda_m$, $j \in \Lambda_{m+1}$). By Proposition 2.5, the matrix $C_m := (c_{ij}^{(\nu_j)})_{\substack{i \in \Lambda_m \\ j \in \Lambda_{m+1}}}$ has full rank λ_{m+1} . Thus, there exists $P_m = (p_{ij}^{(m)})_{i,j \in \Lambda_m} \in \text{GL}(\Lambda_m, \mathbb{Q})$ such that

$$(4.3) \quad P_m C_m = (\delta_{ij})_{\substack{i \in \Lambda_m \\ j \in \Lambda_{m+1}}}.$$

We define

$$(4.4) \quad \psi_{i,i+m} := \sum_{j \in \Lambda_m} p_{ij}^{(m)} \theta_{j,j+m}$$

for $i \in \Lambda_m$ and

$$(4.5) \quad \theta_{i,i+m+1} := \alpha_{i,i+m+1} \psi_{i,i+m}$$

for $i \in \Lambda_{m+1}$. From the inductive assumption together with (4.4) we see that

$$(4.6) \quad \psi_{i,h_m(i)} \in D(\mathcal{A}_{I_m}) \quad \text{for } i \in [n].$$

By the inductive argument, it follows from Theorem 2.6 that the derivations $\{\theta_{i,i+m+1} \mid i \in \Lambda_{m+1}\} \cup \{\psi_{i,i+m} \mid i \in \Lambda_m \setminus \Lambda_{m+1}\} \cup \dots \cup \{\psi_{i,i+1} \mid i \in \Lambda_1 \setminus \Lambda_2\}$ form an \mathcal{R} -basis for $D(\mathcal{A}_{I_{m+1}})$.

Finally, when $m = \text{ht}(\Phi^+)$, we define

$$\psi_{i,i+m} = \theta_{i,i+m}$$

for $i \in \Lambda_m$. Note that $|\Lambda_m| = 1$ whenever $m = \text{ht}(\Phi^+)$ because the root with the highest height is uniquely determined. Hence, if $m = \text{ht}(\Phi^+)$, then we have $[\psi_{i,i+m}]_{i \in \Lambda_m} = P_m [\theta_{i,i+m}]_{i \in \Lambda_m}$ where $P_m = (1) \in \text{GL}(\Lambda_m, \mathbb{Q})$.

Now we check that the derivations $\{\psi_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m\}$ satisfy the conditions (1), (2), (3), (4) in Proposition 3.4. The condition (1) is exactly the definition in (4.1). The condition (2) follows from (4.4) and (4.5). The condition (3) is nothing but (4.6). We check the condition (4). If $m = 0$, then $\psi_{i,i}(\alpha_{j,j+1}) = p_i \alpha_i^*(\alpha_j) = 0$ for $i \in \Lambda_0 = [n]$ and $j \in \Lambda_1 = [n]$ with $i \neq j$. If $m \geq 1$, then we have for $i \in \Lambda_m$ and $j \in \Lambda_{m+1}$ with $i \neq j$

$$\begin{aligned} \psi_{i,i+m}(\alpha_{j,j+m+1}) &= \sum_{k \in \Lambda_m} p_{ik}^{(m)} \theta_{k,k+m}(\alpha_{j,j+m+1}) \quad (\text{from (4.4)}) \\ &\equiv \sum_{k \in \Lambda_m} p_{ik}^{(m)} c_{kj}^{(\nu_j)} b_{\nu_j} \quad \text{mod } \alpha_{j,j+m+1} \quad (\text{by (4.2)}) \\ &= \delta_{ij} b_{\nu_j} \quad (\text{from (4.3)}) \\ &= 0 \quad (\text{because } i \neq j). \end{aligned}$$

Therefore, we conclude that the derivations $\{\psi_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m\}$ form uniform bases from Proposition 3.4. \square

We obtain from Theorem 4.1 uniform bases $\{\psi_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m\}$ by the recursive description. For any non-zero rational numbers p_1, \dots, p_n , we fix the initial data

$$\psi_{i,i} = p_i \alpha_i^* \quad (i \in [n]).$$

Proceeding inductively,

$$(4.7) \quad \psi_{i,i+m} = \sum_{j \in \Lambda_m} p_{ij}^{(m)} \alpha_{j,j+m} \psi_{j,j+m-1} \quad (1 \leq m \leq \text{ht}(\Phi^+), i \in \Lambda_m)$$

for some rational numbers $p_{ij}^{(m)}$. Then we define matrices P_m ($0 \leq m \leq \text{ht}(\Phi^+)$) by

$$(4.8) \quad \begin{aligned} P_0 &:= \text{diag}(p_1, \dots, p_n), \\ P_m &:= (p_{ij}^{(m)})_{i,j \in \Lambda_m} \quad \text{for } m > 0. \end{aligned}$$

As seen in the proof of Theorem 4.1, P_m is invertible for all m . We call the matrices $\{P_m \mid 0 \leq m \leq \text{ht}(\Phi^+)\}$ **the invertible matrices associated with uniform bases** $\{\psi_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m\}$. These invertible matrices are not unique. In fact, we can multiply $\psi_{i,i+m}$ by a non-zero scalar. Also, $\psi_{j,j+m}$ is an element of $D(\mathcal{A}_{\Phi^+})$ for any $j \in \Lambda_m \setminus \Lambda_{m+1}$, so we can replace $\psi_{i,i+m}$ with the $\psi_{i,i+m}$ plus a constant times $\psi_{j,j+m}$ for arbitrary $j \in \Lambda_m \setminus \Lambda_{m+1}$ with $j \neq i$. These correspond to the following two types of elementary row operations for matrices in $\text{GL}(\Lambda_m; \mathbb{Q})$:

- (1) multiply one row of the matrix by a non-zero scalar constant;
- (2) replace an i -th row with the i -th row plus a constant times j -th row for $j \in \Lambda_m \setminus \Lambda_{m+1}$ with $j \neq i$.

For $P_m, P'_m \in \text{GL}(\Lambda_m; \mathbb{Q})$, P_m is **equivalent** to P'_m if P'_m is obtained from P_m by finitely many elementary row operations (1) and (2) above. The following theorem states that P_m is uniquely determined up to the equivalence when we determined the invertible matrices P_0, \dots, P_{m-1} .

Theorem 4.2. *Let $\Phi^+ = \coprod_{i=1}^n \Phi_i^+$ be a decomposition satisfying (3.1) and (3.2). Let $\{P_m \mid 0 \leq m \leq \text{ht}(\Phi^+)\}$ and $\{P'_m \mid 0 \leq m \leq \text{ht}(\Phi^+)\}$ be two sets of the invertible matrices associated with uniform bases $\{\psi_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+), i \in \Lambda_m\}$ and $\{\psi'_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+), i \in \Lambda_m\}$ respectively. Assume that $P_r = P'_r$ for any $r = 0, 1, \dots, m-1$. Then P_m is equivalent to P'_m .*

Proof. Let $\{P_m \mid 0 \leq m \leq \text{ht}(\Phi^+)\}$ be the invertible matrices associated with uniform bases $\Psi := \{\psi_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+), i \in \Lambda_m\}$. Then, we can write from (4.7)

$$(4.9) \quad \psi_{i,i+m} = \sum_{k \in \Lambda_m} p_{ik}^{(m)} \alpha_{k,k+m} \psi_{k,k+m-1} = \sum_{k \in \Lambda_m} p_{ik}^{(m)} \theta_{k,k+m}$$

where $P_m = (p_{ij}^{(m)})_{i,j \in \Lambda_m}$ and $\theta_{k,k+m} := \alpha_{k,k+m} \psi_{k,k+m-1}$ for $k \in \Lambda_m$. For $k \in \Lambda_m$, we define a lower ideal $I_m^{(k)}$ by $I_m^{(k)} = I_m \setminus \{\alpha_{k,k+m}\}$. Since the set of derivations Ψ forms uniform bases, we have $\psi_{k,k+m-1} \in D(\mathcal{A}_{I_m^{(k)}})$ which implies that $\theta_{k,k+m} = \alpha_{k,k+m} \psi_{k,k+m-1} \in D(\mathcal{A}_{I_m})$.

We write $\mathcal{A}_j'' = \{H \cap H_{j,j+m+1} \mid H \in \mathcal{A}_{I_m}\}$ where $H_{j,j+m+1}$ is the hyperplane defined by the positive root $\alpha_{j,j+m+1}$. We fix a map $\nu_j : \mathcal{A}_j'' \rightarrow \mathcal{A}_{I_m}$ such that $\nu_j(X) \cap H_{j,j+m+1} = X$, and define the homogeneous polynomial $b_{\nu_j} = (\prod_{H \in \mathcal{A}_{I_m}} \alpha_H) / (\prod_{X \in \mathcal{A}_j''} \alpha_{\nu_j(X)})$ which has degree m from Proposition 2.4. It follows from Proposition 2.3 that for $k \in \Lambda_m$ and $j \in \Lambda_{m+1}$ we can write

$$\theta_{k,k+m}(\alpha_{j,j+m+1}) \equiv c_{kj}^{(\nu_j)} b_{\nu_j} \pmod{\alpha_{j,j+m+1}}$$

for some rational numbers $c_{kj}^{(\nu_j)}$. From this together with (4.9) we have

$$(4.10) \quad \psi_{i,i+m}(\alpha_{j,j+m+1}) \equiv b_{\nu_j} \left(\sum_{k \in \Lambda_m} p_{ik}^{(m)} c_{kj}^{(\nu_j)} \right) \pmod{\alpha_{j,j+m+1}}.$$

We consider a lower ideal $I := I_m \cup \{\alpha_{j,j+m+1}\}$, and one see that $\psi_{i,i+m} \in D(\mathcal{A}_I)$ for $i \in \Lambda_m$ and $j \in \Lambda_{m+1}$ with $i \neq j$ since Ψ form uniform bases. In particular, $\psi_{i,i+m}(\alpha_{j,j+m+1}) \in \mathcal{R}\alpha_{j,j+m+1}$. From this together with (4.10), we obtain

$$\sum_{k \in \Lambda_m} p_{ik}^{(m)} c_{kj}^{(\nu_j)} = 0 \quad \text{for } i \in \Lambda_m, j \in \Lambda_{m+1} \text{ with } i \neq j.$$

Let $C_m := (c_{ij}^{(\nu_j)})_{\substack{i \in \Lambda_m \\ j \in \Lambda_{m+1}}}$. The matrix C_m has full rank λ_{m+1} by Proposition 2.5. Therefore, we obtain

$$(4.11) \quad P_m C_m = (q_j \delta_{ij})_{\substack{i \in \Lambda_m \\ j \in \Lambda_{m+1}}}$$

for some non-zero rational numbers q_j ($j \in \Lambda_{m+1}$).

We take another invertible matrices $\{P'_m \mid 0 \leq m \leq \text{ht}(\Phi^+)\}$ associated with uniform bases $\{\psi'_{i,i+m} \in \text{Der}(\mathcal{R}) \mid 0 \leq m \leq \text{ht}(\Phi^+), i \in \Lambda_m\}$. From the assumption $P_r = P'_r$ for $r = 0, \dots, m-1$, we have $\psi_{s,s+r} = \psi'_{s,s+r}$ for $0 \leq r \leq m-1$ and $s \in \Lambda_r$. Noting that $\theta'_{k,k+m} := \alpha_{k,k+m} \psi'_{k,k+m-1} = \alpha_{k,k+m} \psi_{k,k+m-1} = \theta_{k,k+m}$ for $k \in \Lambda_m$, by similar argument we have

$$(4.12) \quad P'_m C_m = (q'_j \delta_{ij})_{\substack{i \in \Lambda_m \\ j \in \Lambda_{m+1}}}$$

for some non-zero rational numbers q'_j ($j \in \Lambda_{m+1}$). From (4.11) and (4.12), we have

$$(4.13) \quad A_m C_m = (\delta_{ij})_{\substack{i \in \Lambda_m \\ j \in \Lambda_{m+1}}} = A'_m C_m$$

where $A_m = (q_j^{-1} \delta_{ij})_{\substack{i \in \Lambda_m \\ j \in \Lambda_{m+1}}} \cdot P_m$ and $A'_m = (q'^{-1}_j \delta_{ij})_{\substack{i \in \Lambda_m \\ j \in \Lambda_{m+1}}} \cdot P'_m$. Let \mathbf{a}_i and \mathbf{a}'_i be i -th row vectors of A_m and A'_m respectively for $i \in \Lambda_m$. Then, the difference $\mathbf{a}'_i - \mathbf{a}_i$ belongs to the kernel of C_m by (4.13). For any $j \in \Lambda_m \setminus \Lambda_{m+1}$ the j -th row vector \mathbf{p}_j of P_m is an element of $\ker C_m$ by (4.11), and these row vectors \mathbf{p}_j are linearly independent because P_m is invertible. Since the matrix C_m has rank λ_{m+1} , the row vectors \mathbf{p}_j ($j \in \Lambda_m \setminus \Lambda_{m+1}$) form a basis of $\ker C_m$. Hence, the difference $\mathbf{a}'_i - \mathbf{a}_i$ can be written as a linear combination of \mathbf{p}_j ($j \in \Lambda_m \setminus \Lambda_{m+1}$), which means that P_m is equivalent to P'_m , as desired. \square

If we want to find uniform bases, then it is enough to determine the invertible matrices $P_0, P_1, \dots, P_{\text{ht}(\Phi^+)}$ in order. In next section, giving a decomposition $\Phi^+ = \coprod_{i=1}^n \Phi_i^+$ satisfying (3.1) and (3.2), we will describe explicitly uniform bases and the associated invertible matrices P_m on each Lie type.

5. THE INVERTIBLE MATRICES ASSOCIATED WITH UNIFORM BASES

In this section we fix a decomposition $\Phi^+ = \coprod_{i=1}^n \Phi_i^+$ satisfying (3.1) and (3.2), and describe explicitly uniform bases and the associated invertible matrices P_m in (4.8) on each Lie type. For a construction of an explicit \mathcal{R} -basis of $D(\mathcal{A}_I)$ for each Lie type, Barakat, Cuntz, and Hoge provided ones for types E and F by computer when the work of [3] was in progress. Also, Terao and Abe worked for the special lower ideal I_m in (3.5) of types A and B , respectively. Uniform bases were constructed for types A, B, C, G in

[4, Proposition 10.3, Theorem 10.9, Theorem 10.14, and Theorem 10.17]. For the other types, we construct explicit uniform bases. We also describe the associated invertible matrices P_m on each Lie type.

Throughout this section, V is an n -dimensional real euclidean space, and x_1, \dots, x_n is an orthonormal basis of V^* . We denote the partial derivatives by $\partial_i = \frac{\partial}{\partial x_i}$ for all $i = 1, \dots, n$. The euclidean space \mathfrak{t} in which the ideal arrangements are realized will be defined to be a certain subspace of V or V itself.

5.1. Type A_{n-1} ($n \geq 2$). Let \mathfrak{t} be the hyperplane in V defined by the linear function $x_1 + \dots + x_n$. Then we have

$$\mathcal{R} = \text{Sym}(\mathfrak{t}^*) = \mathbb{R}[x_1, \dots, x_n]/(x_1 + \dots + x_n).$$

We take the set of positive roots of type A_{n-1} as

$$\Phi_{A_{n-1}}^+ = \{x_i - x_j \in \mathfrak{t}^* \mid 1 \leq i < j \leq n\}$$

and set the exponents e_1, \dots, e_{n-1} as

$$e_i = n - i \quad \text{for } 1 \leq i \leq n - 1.$$

We arrange all positive roots in a strict upper triangular $n \times n$ matrix shown in Figure 1.




FIGURE 1. The arrangement of all positive roots for type A_{n-1} .

In Figure 1 the partial order \preceq on $\Phi_{A_{n-1}}^+$ is defined as follows:

- (1) if a root α is left-adjacent to a root β , then $\alpha \preceq \beta$;
- (2) if a root α is lower-adjacent to a root β , then $\alpha \preceq \beta$.

For two positive roots α, β , we define $\alpha \preceq \beta$ if there exist positive roots $\gamma_0, \dots, \gamma_N$ such that $\alpha = \gamma_0 \preceq \gamma_1 \preceq \dots \preceq \gamma_N = \beta$. We denote positive roots in $\Phi_{A_{n-1}}^+$ by

$$(5.1) \quad \alpha_{i,j} = x_i - x_j \quad \text{for all } 1 \leq i < j \leq n.$$

Note that $\alpha_{i,j} \preceq \alpha_{k,\ell}$ iff $i \geq k$ and $j \leq \ell$. Recall that $\Phi_i^+ = \{\alpha_{i,j} \mid i < j \leq i + e_i\}$, namely it is the set of all positive roots in the i -th row in Figure 1. A **Hessenberg function for type A_{n-1}** is defined to be a function $h : [n-1] \rightarrow [n-1]$ satisfying the following two conditions

- (1) $h(1) \leq h(2) \leq \dots \leq h(n-1)$,
- (2) $h(i) \geq i$ for $i \in [n-1]$.

Under the decomposition $\Phi_{A_{n-1}}^+ = \coprod_{i=1}^{n-1} \Phi_i^+$, one can see that the set of lower ideals $I \subset \Phi^+$ and the set of Hessenberg functions h for type A_{n-1} are in one-to-one correspondence which sends I to h_I in (3.3).

Now we explain the uniform bases given by [4]. Let $\bar{\partial} := \partial_1 + \cdots + \partial_n$. For $1 \leq i \leq n-1$ and $i \leq j \leq n$, we define

$$\psi_{i,j}^{A_{n-1}} := \sum_{k=1}^i \left(\prod_{\ell=i+1}^j (x_k - x_\ell) \right) \left(\partial_k - \frac{1}{n} \bar{\partial} \right) \in \text{Der } \mathcal{R} = \mathcal{R} \otimes \mathfrak{t}$$

with the convention $\prod_{\ell=i+1}^j (x_k - x_\ell) = 1$ whenever $j = i$. Note that ∂_k is not an element of $\text{Der } \mathcal{R}$ but $\partial_k - \frac{1}{n} \bar{\partial}$ is, because $\mathcal{R} = \text{Sym}(\mathfrak{t}^*) = \mathbb{R}[x_1, \dots, x_n]/(x_1 + \cdots + x_n)$.

Theorem 5.1. ([4, Proposition 10.3]) *The derivations $\{\psi_{i,j}^{A_{n-1}} \mid 1 \leq i \leq n-1, i \leq j \leq n\}$ form uniform bases for the ideal arrangements of type A_{n-1} .*

As explained in [4], the derivations $\psi_{i,j}^{A_{n-1}}$ satisfy the following recursive formula:

$$\begin{aligned} \psi_{i,i}^{A_{n-1}} &= \partial_1 + \cdots + \partial_i - \frac{i}{n} \bar{\partial} \quad \text{for } i \in [n-1]; \\ \psi_{i,j}^{A_{n-1}} &= \psi_{i-1,j-1}^{A_{n-1}} + (x_i - x_j) \psi_{i,j-1}^{A_{n-1}} \quad \text{for } (i, j) \text{ with } i < j \leq n, \end{aligned}$$

where we take the convention $\psi_{0,*}^{A_{n-1}} = 0$ for any $*$. Note that $\text{ht}(\Phi_{A_{n-1}}^+) = n-1$ and the set Λ_m in (3.6) is given by

$$\Lambda_m = \begin{cases} [n-1] & \text{if } m = 0, \\ [n-m] & \text{if } 1 \leq m \leq n-1. \end{cases}$$

One can see from the recursive formula above that

$$\begin{aligned} \psi_{i,i}^{A_{n-1}} &= \alpha_i^* \quad \text{for } i \in [n-1], \\ \psi_{i,i+m}^{A_{n-1}} &= \sum_{j=1}^i \alpha_{j,j+m} \psi_{j,j+m-1}^{A_{n-1}} \quad \text{for } 1 \leq m \leq n-1 \text{ and } 1 \leq i \leq n-m. \end{aligned}$$

Therefore, the invertible matrices $P_m^{A_{n-1}}$ ($0 \leq m \leq n-1$) associated with the uniform bases $\{\psi_{i,i+m}^{A_{n-1}} \mid 0 \leq m \leq n-1, i \in \Lambda_m\}$ are given by

$$P_0^{A_{n-1}} = \begin{pmatrix} 1 & & \\ & \ddots & \\ & & 1 \end{pmatrix}, \quad P_m^{A_{n-1}} = \begin{pmatrix} 1 & & \\ \vdots & \ddots & \\ 1 & \cdots & 1 \end{pmatrix} \quad (1 \leq m \leq n-1).$$

5.2. Type B_n ($n \geq 2$). Let $\mathfrak{t} = V$. Then we have

$$\mathcal{R} = \text{Sym}(\mathfrak{t}^*) = \mathbb{R}[x_1, \dots, x_n].$$


We take the set of positive roots of type B_n as

$$\Phi_{B_n}^+ = \{x_i \pm x_j \in \mathfrak{t}^* \mid 1 \leq i < j \leq n\} \cup \{x_i \in \mathfrak{t}^* \mid 1 \leq i \leq n\}$$

and set the exponents e_1, \dots, e_n as

$$e_i = 2(n-i) + 1 \quad \text{for } 1 \leq i \leq n.$$

We arrange all positive roots of B_n as shown in Figure 2.

FIGURE 2. The arrangement of all positive roots for type B_n .

In Figure 2 the partial order \preceq on $\Phi_{B_n}^+$ is defined as follows:

- (1) if a root α is left-adjacent to a root β , then $\alpha \prec \beta$;
- (2) if a root α is lower-adjacent to a root β , then $\alpha \prec \beta$.

For two positive roots α, β , we define $\alpha \preceq \beta$ if there exist positive roots $\gamma_0, \dots, \gamma_N$ such that $\alpha = \gamma_0 \prec \gamma_1 \prec \dots \prec \gamma_N = \beta$. For each $i = 1, \dots, n$, we denote positive roots in $\Phi_{B_n}^+$ by

$$\alpha_{i,j} = \begin{cases} x_i - x_j & \text{if } i+1 \leq j \leq n, \\ x_i & \text{if } j = n+1, \\ x_i + x_{2n+2-j} & \text{if } n+2 \leq j \leq 2n+1-i. \end{cases}$$

Note that $\alpha_{i,j} \preceq \alpha_{k,\ell}$ iff $i \geq k$ and $j \leq \ell$. Similarly to type A , $\Phi_i^+ = \{\alpha_{i,j} \mid i < j \leq i + e_i\}$ is the set of all positive roots in the i -th row in Figure 2. A **Hessenberg function for type B_n** is defined to be a function $h : [n] \rightarrow [2n]$ satisfying the following three conditions

- (1) $i \leq h(i) \leq 2n+1-i$ for $i \in [n]$,
- (2) if $h(i) \neq 2n+1-i$, then $h(i) \leq h(i+1)$ for $i = 1, \dots, n-1$,
- (3) if $h(i) = 2n+1-i$, then $h(i+1) = 2n+1-(i+1)$ for $i = 1, \dots, n-1$.

Under the decomposition $\Phi_{B_n}^+ = \coprod_{i=1}^n \Phi_i^+$, one can see that the set of lower ideals $I \subset \Phi_{B_n}^+$ and the set of Hessenberg functions h for type B_n are in one-to-one correspondence which sends I to h_I in (3.3).

In [4], the derivation $\psi_{i,j}^{B_n}$ for $1 \leq i \leq n$ and $i \leq j \leq 2n+1-i$ is defined as

$$\psi_{i,j}^{B_n} := \sum_{k=1}^i \left(\prod_{\ell=i+1}^j \alpha_{k,\ell} \right) \partial_k \in \text{Der } \mathcal{R} = \mathcal{R} \otimes \mathfrak{t},$$

where we take the convention $\prod_{\ell=i+1}^j \alpha_{k,\ell} = 1$ whenever $j = i$.

Theorem 5.2. ([4, Theorem 10.9]) *The derivations $\{\psi_{i,j}^{B_n} \mid 1 \leq i \leq n, i \leq j \leq 2n+1-i\}$ form uniform bases for the ideal arrangements of type B_n .*

The derivations $\psi_{i,j}^{B_n}$ satisfy the following recursive formula ([4])

$$\begin{aligned} \psi_{i,i}^{B_n} &= \partial_1 + \dots + \partial_i \quad \text{for } i \in [n], \\ \psi_{i,j}^{B_n} &= \psi_{i-1,j-1}^{B_n} + \alpha_{i,j} \psi_{i,j-1}^{B_n} \quad \text{for } (i,j) \text{ with } i < j \leq 2n+1-i, \end{aligned}$$

where we take the convention $\psi_{0,*}^{B_n} = 0$ for any $*$. Note that $\text{ht}(\Phi_{B_n}^+) = 2n - 1$ and the set Λ_m in (3.6) is given by

$$\Lambda_m = [n - k] \quad \text{if } m = 2k, 2k + 1.$$

From the recursive formula above we obtain

$$\begin{aligned} \psi_{i,i}^{B_n} &= \alpha_i^* \quad \text{for } 1 \leq i \leq n, \\ \psi_{i,i+m}^{B_n} &= \sum_{j=1}^i \alpha_{j,j+m} \psi_{j,j+m-1}^{B_n} \quad \text{for } 1 \leq m \leq 2n - 1 \text{ and } i \in \Lambda_m. \end{aligned}$$

Hence, the invertible matrices $P_m^{B_n}$ ($0 \leq m \leq 2n - 1$) associated with the uniform bases $\{\psi_{i,i+m}^{B_n} \mid 0 \leq m \leq 2n - 1, i \in \Lambda_m\}$ are given by

$$P_0^{B_n} = \begin{pmatrix} 1 & & \\ & \ddots & \\ & & 1 \end{pmatrix}, \quad P_m^{B_n} = \begin{pmatrix} 1 & & \\ \vdots & \ddots & \\ 1 & \cdots & 1 \end{pmatrix} \quad (1 \leq m \leq 2n - 1).$$

5.3. **Type C_n** ($n \geq 2$). Let $\mathfrak{t} = V$ and we have

$$\mathcal{R} = \text{Sym}(\mathfrak{t}^*) = \mathbb{R}[x_1, \dots, x_n].$$

We take the set of positive roots of type C_n as

$$\Phi_{C_n}^+ = \{x_i \pm x_j \in \mathfrak{t}^* \mid 1 \leq i < j \leq n\} \cup \{2x_i \in \mathfrak{t}^* \mid 1 \leq i \leq n\}$$

and set the exponents e_1, \dots, e_n as

$$e_i = 2(n - i) + 1 \quad \text{for } 1 \leq i \leq n.$$

We arrange all positive roots of C_n as shown in Figure 3.




FIGURE 3. The arrangement of all positive roots for type C_n .

In Figure 3 the partial order \preceq on $\Phi_{C_n}^+$ is defined as follows:

- (1) if a root α is left-adjacent to a root β , then $\alpha \preceq \beta$;
- (2) if a root α is lower-adjacent to a root β , then $\alpha \preceq \beta$.

For two positive roots α, β , we define $\alpha \preceq \beta$ if there exist positive roots $\gamma_0, \dots, \gamma_N$ such that $\alpha = \gamma_0 \preceq \gamma_1 \preceq \dots \preceq \gamma_N = \beta$. For each $i = 1, \dots, n$, we denote positive roots² in $\Phi_{C_n}^+$

²We remark that the notation $\alpha_{i,2n+1-i}$ is used for x_i in [4].

by

$$\alpha_{i,j} = \begin{cases} x_i - x_j & \text{if } i+1 \leq j \leq n, \\ x_i + x_{2n+1-j} & \text{if } n+1 \leq j \leq 2n-i, \\ 2x_i & \text{if } j = 2n+1-i. \end{cases}$$

Note that $\alpha_{i,j} \preceq \alpha_{k,\ell}$ iff $i \geq k$ and $j \leq \ell$. Similarly to type A , $\Phi_i^+ = \{\alpha_{i,j} \mid i < j \leq i+e_i\}$ is the set of all positive roots in the i -th row in Figure 3. A **Hessenberg function for type C_n** is defined to be the same one as type B_n . Under the decomposition $\Phi_{C_n}^+ = \coprod_{i=1}^n \Phi_i^+$, one can see that the set of lower ideals $I \subset \Phi_{C_n}^+$ and the set of Hessenberg functions h for type C_n are in one-to-one correspondence which sends I to h_I in (3.3).

Like the derivation defined in [4], we define the derivation³ $\psi_{i,j}^{C_n}$ for $1 \leq i \leq n$ and $i \leq j \leq 2n+1-i$ by

$$(5.2) \quad \psi_{i,j}^{C_n} := \sum_{k=1}^i \left(\prod_{\ell=i+1}^j \alpha_{k,\ell} \right) \partial_k \in \text{Der } \mathcal{R} = \mathcal{R} \otimes \mathfrak{t},$$

where we take the convention $\prod_{\ell=i+1}^i \alpha_{k,\ell} = 1$.

Theorem 5.3. ([4, Theorem 10.14]) *The derivations $\{\psi_{i,j}^{C_n} \mid 1 \leq i \leq n, i \leq j \leq 2n+1-i\}$ form uniform bases for the ideal arrangements of type C_n .*

Using the recursive formula in [4], we have

$$\begin{aligned} \psi_{i,i}^{C_n} &= \partial_1 + \cdots + \partial_i \quad \text{for } i = 1, \dots, n, \\ \psi_{i,j}^{C_n} &= \psi_{i-1,j-1}^{C_n} + \alpha_{i,j} \psi_{i,j-1}^{C_n} \quad \text{for } (i,j) \text{ with } i < j < 2n+1-i, \\ \psi_{i,2n+1-i}^{C_n} &= 2\psi_{i-1,2n-i}^{C_n} + \alpha_{i,2n+1-i} \psi_{i,2n-i}^{C_n} \quad \text{for } i = 1, \dots, n, \end{aligned}$$

where we take the convention $\psi_{0,*}^{C_n} = 0$ for any $*$. Note that $\text{ht}(\Phi_{C_n}^+) = 2n-1$ and the set Λ_m in (3.6) is given by

$$\Lambda_m = [n-k] \quad \text{if } m = 2k, 2k+1.$$

From the recursive formula above we obtain

$$\begin{aligned} \psi_{i,i}^{C_n} &= \alpha_i^* \quad \text{for } 1 \leq i \leq n-1, \\ \psi_{n,n}^{C_n} &= 2\alpha_n^*, \\ \psi_{i,i+m}^{C_n} &= \sum_{j=1}^i \alpha_{j,j+m} \psi_{j,j+m-1}^{C_n} \quad \text{for } 1 \leq m \leq 2n-1 \text{ and } i \in \Lambda_m \setminus \{\max_{\ell \in \Lambda_m} \ell\}. \\ \psi_{i,i+m}^{C_n} &= \sum_{j=1}^{i-1} 2\alpha_{j,j+m} \psi_{j,j+m-1}^{C_n} + \alpha_{i,i+m} \psi_{i,i+m-1}^{C_n} \quad \text{for } 1 \leq m \leq 2n-1 \text{ and } i = \max_{\ell \in \Lambda_m} \ell. \end{aligned}$$

³The derivation $\psi_{i,2n+1-i}^{C_n}$ in (5.2) is $\psi_{i,2n+1-i}^C$ defined in [4] up to scalar multiplication.

Thus, the invertible matrices $P_m^{C_n}$ ($0 \leq m \leq 2n - 1$) associated with the uniform bases $\{\psi_{i,i+m}^{C_n} \mid 0 \leq m \leq 2n - 1, i \in \Lambda_m\}$ are given by

$$P_0^{C_n} = \begin{pmatrix} 1 & & & \\ & \ddots & & \\ & & 1 & \\ & & & 2 \end{pmatrix}, \quad P_m^{C_n} = \begin{pmatrix} 1 & & & \\ \vdots & \ddots & & \\ 1 & \cdots & 1 & \\ 2 & \cdots & 2 & 1 \end{pmatrix} \quad (1 \leq m \leq 2n - 1).$$

5.4. Type D_n ($n \geq 4$).⁴

Let $\mathfrak{t} = V$ and we have

$$\mathcal{R} = \text{Sym}(\mathfrak{t}^*) = \mathbb{R}[x_1, \dots, x_n].$$

We take the set of positive roots of type D_n as

$$\Phi_{D_n}^+ = \{x_i \pm x_j \in \mathfrak{t}^* \mid 1 \leq i < j \leq n\}$$

and set the exponents e_1, \dots, e_n as

$$\begin{aligned} e_i &= 2(n - i) - 1 & \text{for } 1 \leq i \leq n - 1, \\ e_n &= n - 1. \end{aligned}$$

We arrange all positive roots of D_n as shown in Figure 4.




FIGURE 4. The arrangement of all positive roots for type D_n .

In Figure 4 the partial order \preceq on $\Phi_{D_n}^+$ is defined as follows:

- (1) if a root α is left-adjacent to a root β , then $\alpha \preceq \beta$, except for $(\alpha, \beta) = (x_i - x_n, x_i + x_n)$ ($1 \leq i \leq n - 1$) which are pictorially divided by a dotted line;
- (2) $x_i - x_{n-1} \preceq x_i + x_n$ and $x_i - x_n \preceq x_i + x_{n+1}$ for $1 \leq i \leq n - 1$;
- (3) if a root α is lower-adjacent to a root β , then $\alpha \preceq \beta$.

For two positive roots α, β , we define $\alpha \preceq \beta$ if there exist positive roots $\gamma_0, \dots, \gamma_N$ such that $\alpha = \gamma_0 \preceq \gamma_1 \preceq \dots \preceq \gamma_N = \beta$. We denote positive roots in $\Phi_{D_n}^+$ by

$$\alpha_{i,j} = \begin{cases} x_i - x_j & \text{if } i + 1 \leq j \leq n, \\ x_i + x_{2n-j} & \text{if } n + 1 \leq j \leq 2n - i - 1 \end{cases}$$

for each $i = 1, \dots, n - 1$, and

$$\alpha_{n,j} = x_{2n-j} + x_n \quad \text{if } n + 1 \leq j \leq 2n - 1.$$

⁴Note that statements in this subsection valid for $n \geq 2$.

Note that Φ_i^+ is the set of all positive roots in the i -th row except for the root $x_i + x_n$ for $i = 1, \dots, n-1$, and Φ_n^+ is the set of all positive roots in the $(n+1)$ -th column in Figure 4. Motivated by this, we define the coordinate in type D_n as shown in Figure 5.




FIGURE 5. The coordinate in type D_n .

We define a **Hessenberg function for type D_n** as a function $h : [n] \rightarrow [2n-1]$ satisfying the following conditions

- (1) $i \leq h(i) \leq 2n-1-i$ for $i = 1, \dots, n-1$,
- (2) $n \leq h(n) \leq 2n-1$,
- (3) if $h(i) \neq 2n-1-i$, then $h(i) \leq h(i+1)$ for $i = 1, \dots, n-2$,
- (4) if $h(i) = 2n-1-i$, then $h(i+1) = 2n-1-(i+1)$ for $i = 1, \dots, n-2$,
- (5) if $h(i) \geq n+1$, then $h(n) \geq 2n-i$ for $i = 1, \dots, n-2$,⁵
- (6) if $h(n) \geq 2n-i$, then $h(i) \geq n-1$ for $i = 1, \dots, n-2$.

Under the decomposition $\Phi_{D_n}^+ = \coprod_{i=1}^n \Phi_i^+$, one can see that the set of lower ideals $I \subset \Phi_{D_n}^+$ and the set of Hessenberg functions h for type D_n are in one-to-one correspondence which sends I to h_I in (3.3). We write a Hessenberg function by listing its values in sequence, namely, $h = (h(1), h(2), \dots, h(n))$.

It is useful to express a Hessenberg function h pictorially by drawing a configuration of boxes on a square grid with the coordinate in Figure 5 whose shaded boxes correspond to the roots of the lower ideal I associated with h and (i, i) -boxes (see for example [1] for type A). For example the picture of the Hessenberg function $h = (3, 5, 4, 7)$ is shown in Figure 6.

(1,1)	(1,2)	(1,3)	(1,4)	(4,7)	(1,5)	(1,6)
	(2,2)	(2,3)	(2,4)	(4,6)	(2,5)	
		(3,3)	(3,4)	(4,5)		
				(4,4)		

FIGURE 6. The picture for the Hessenberg function $h = (3, 5, 4, 7)$.

Now we define the derivations $\{\psi_{i,j}^{D_n} \mid 1 \leq i \leq n-1, i \leq j \leq 2n-i-1\} \cup \{\psi_{n,j}^{D_n} \mid n \leq j \leq 2n-1\}$ by a recursive formula which is a different flavor than that of types A, B, C .

⁵The condition (5) is true for $i = n-1$ because $h(n-1) = n-1$ or $h(n-1) = n$.

We begin with the case when $j = i$. In this case we make the following definition

$$\begin{aligned}\psi_{i,i}^{D_n} &= \partial_1 + \cdots + \partial_i, \quad \text{for } 1 \leq i \leq n-2, \\ \psi_{n-1,n-1}^{D_n} &= \partial_1 + \cdots + \partial_{n-1} - \partial_n, \\ \psi_{n,n}^{D_n} &= \partial_1 + \cdots + \partial_{n-1} + \partial_n.\end{aligned}$$

Now we proceed inductively for the rest of the $\psi_{i,j}^{D_n}$ as follows:

$$(5.3) \quad \psi_{i,j}^{D_n} = \psi_{i-1,j-1}^{D_n} + \alpha_{i,j} \psi_{i,j-1}^{D_n} + \delta_{j,n-1} \xi_i^{D_n} \quad \text{for } 1 \leq i \leq n-1, i+1 \leq j \leq 2n-1-i,$$

$$(5.4) \quad \psi_{n,j}^{D_n} = \alpha_{n,j} \psi_{n,j-1}^{D_n} + (-1)^{j-n} \psi_{2n-j,n}^{D_n} \quad \text{for } n+1 \leq j \leq 2n-1,$$

where $\psi_{0,j}^{D_n}$ is defined to be the following

$$(5.5) \quad \psi_{0,j}^{D_n} = \begin{cases} 0 & \text{for } 1 \leq j \leq n-2, \\ (-1)^n (\sum_{k=1}^n x_1 \cdots \widehat{x}_k \cdots x_n \partial_k) & \text{for } j = n-1, \\ -x_{2n-j} x_{2n-j+1} \cdots x_n \psi_{0,n-1}^{D_n} & \text{for } n \leq j \leq 2n-3, \end{cases}$$

and $\xi_i^{D_n}$ is defined as

$$(5.6) \quad \begin{aligned} \xi_i^{D_n} &= \sum_{k=1}^i ((x_k - x_{i+1}) \cdots (x_k - x_{n-1}) x_n + (-1)^{n-i} x_{i+1} \cdots x_n) x_k^{-1} \partial_k \\ &\quad + (-1)^{n-i} \sum_{k=i+1}^n x_{i+1} \cdots \widehat{x}_k \cdots x_n \partial_k, \end{aligned}$$

where the caret sign $\widehat{}$ over x_j means that the entry x_j is to be omitted. Note that $((x_j - x_{i+1}) \cdots (x_j - x_{n-1}) x_n + (-1)^{n-i} x_{i+1} \cdots x_{n-1} x_n)$ is divisible by x_j . Thus, $\xi_i^{D_n}$ is an element of $\text{Der } \mathcal{R} = \mathcal{R} \otimes \mathfrak{t}$, so is $\psi_{i,j}^{D_n}$. Also, we define $\xi_i^{D_n}$ for $i = 0$ as

$$\xi_0^{D_n} := \psi_{0,n-1}^{D_n}.$$

Note that we need the derivations $\xi_i^{D_n}$ only for $0 \leq i \leq n-2$. In fact, if we take $i = n-1$ in (5.3), then one can see that $j = n$ and $\xi_{n-1}^{D_n}$ does not appear in the right hand side of (5.3).

Example 5.4. In type D_4 , the positive roots $\alpha_{i,j}$ are given by

$$\begin{aligned} \alpha_{1,2} &= x_1 - x_2, \quad \alpha_{1,3} = x_1 - x_3, \quad \alpha_{1,4} = x_1 - x_4, \quad \alpha_{1,5} = x_1 + x_3, \quad \alpha_{1,6} = x_1 + x_2, \\ \alpha_{2,3} &= x_2 - x_3, \quad \alpha_{2,4} = x_2 - x_4, \quad \alpha_{2,5} = x_2 + x_3, \\ \alpha_{3,4} &= x_3 - x_4, \\ \alpha_{4,5} &= x_3 + x_4, \quad \alpha_{4,6} = x_2 + x_4, \quad \alpha_{4,7} = x_1 + x_4, \end{aligned}$$

and we arrange the derivations $\psi_{i,j}^{D_4}$ in the coordinate shown in Figure 5 as follows:

$\psi_{1,1}^{D_4}$	$\psi_{1,2}^{D_4}$	$\psi_{1,3}^{D_4}$	$\psi_{1,4}^{D_4}$	$\psi_{4,7}^{D_4}$	$\psi_{1,5}^{D_4}$	$\psi_{1,6}^{D_4}$
	$\psi_{2,2}^{D_4}$	$\psi_{2,3}^{D_4}$	$\psi_{2,4}^{D_4}$	$\psi_{4,6}^{D_4}$	$\psi_{2,5}^{D_4}$	
		$\psi_{3,3}^{D_4}$	$\psi_{3,4}^{D_4}$	$\psi_{4,5}^{D_4}$		
				$\psi_{4,4}^{D_4}$		

Then the derivations $\psi_{i,j}^{D_4} \in \text{Der } \mathcal{R} = \mathcal{R} \otimes \mathfrak{t}$ are computed as follows:

$$\begin{aligned}
\psi_{1,1}^{D_4} &= \partial_1, \\
\psi_{1,2}^{D_4} &= (x_1 - x_2)\partial_1, \\
\psi_{1,3}^{D_4} &= \frac{((x_1 - x_2)(x_1 - x_3)(x_1 + x_4) - x_2x_3x_4)}{x_1}\partial_1 - x_3x_4\partial_2 - x_2x_4\partial_3 - x_2x_3\partial_4, \\
\psi_{1,4}^{D_4} &= \frac{((x_1 - x_2)(x_1 - x_3)(x_1 - x_4)(x_1 + x_4) + x_2x_3x_4^2)}{x_1}\partial_1 + x_3x_4^2\partial_2 + x_2x_4^2\partial_3 + x_2x_3x_4\partial_4, \\
\psi_{1,5}^{D_4} &= \frac{((x_1 - x_2)(x_1 - x_3)(x_1 - x_4)(x_1 + x_4)(x_1 + x_3) + x_2x_3^2x_4^2)}{x_1}\partial_1 \\
&\quad + x_3^2x_4^2\partial_2 + x_2x_3x_4^2\partial_3 + x_2x_3^2x_4\partial_4, \\
\psi_{1,6}^{D_4} &= \frac{((x_1 - x_2)(x_1 - x_3)(x_1 - x_4)(x_1 + x_4)(x_1 + x_3)(x_1 + x_2) + x_2^2x_3^2x_4^2)}{x_1}\partial_1 \\
&\quad + x_2x_3^2x_4^2\partial_2 + x_2^2x_3x_4^2\partial_3 + x_2^2x_3^2x_4\partial_4, \\
\psi_{2,2}^{D_4} &= \partial_1 + \partial_2, \\
\psi_{2,3}^{D_4} &= \frac{((x_1 - x_3)(x_1 + x_4) + x_3x_4)}{x_1}\partial_1 + \frac{((x_2 - x_3)(x_2 + x_4) + x_3x_4)}{x_2}\partial_2 + x_4\partial_3 + x_3\partial_4, \\
\psi_{2,4}^{D_4} &= \frac{((x_1 - x_3)(x_1 - x_4)(x_1 + x_4) - x_3x_4^2)}{x_1}\partial_1 + \frac{((x_2 - x_3)(x_2 - x_4)(x_2 + x_4) - x_3x_4^2)}{x_2}\partial_2 \\
&\quad - x_4^2\partial_3 - x_3x_4\partial_4, \\
\psi_{2,5}^{D_4} &= \frac{((x_1 - x_3)(x_1 - x_4)(x_1 + x_4)(x_1 + x_3) - x_3^2x_4^2)}{x_1}\partial_1 \\
&\quad + \frac{((x_2 - x_3)(x_2 - x_4)(x_2 + x_4)(x_2 + x_3) - x_3^2x_4^2)}{x_2}\partial_2 \\
&\quad - x_3x_4^2\partial_3 - x_3^3x_4\partial_4, \\
\psi_{3,3}^{D_4} &= \partial_1 + \partial_2 + \partial_3 - \partial_4, \\
\psi_{3,4}^{D_4} &= x_1\partial_1 + x_2\partial_2 + x_3\partial_3 + x_4\partial_4, \\
\psi_{4,4}^{D_4} &= \partial_1 + \partial_2 + \partial_3 + \partial_4, \\
\psi_{4,5}^{D_4} &= -\frac{((x_1 - x_3)(x_1 - x_4) - x_3x_4)}{x_1}\partial_1 - \frac{((x_2 - x_3)(x_2 - x_4) - x_3x_4)}{x_2}\partial_2 + x_4\partial_3 + x_3\partial_4, \\
\psi_{4,6}^{D_4} &= \frac{((x_1 - x_2)(x_1 - x_3)(x_1 - x_4) + x_2x_3x_4)}{x_1}\partial_1 + x_3x_4\partial_2 + x_2x_4\partial_3 + x_2x_3\partial_4, \\
\psi_{4,7}^{D_4} &= x_2x_3x_4\partial_1 + x_1x_3x_4\partial_2 + x_1x_2x_4\partial_3 + x_1x_2x_3\partial_4.
\end{aligned}$$

It is straightforward from the definition of $\psi_{i,j}^{D_n}$ to see the following explicit formula for $\psi_{i,j}^{D_n}$. For $1 \leq i \leq n-1$ we obtain

$$(5.7) \quad \psi_{i,j}^{D_n} = \sum_{k=1}^i (x_k - x_{i+1}) \cdots (x_k - x_j) \partial_k \quad \text{for } i \leq j \leq n-2 \ (i \neq n-1),$$

$$(5.8) \quad \psi_{i,n-1}^{D_n} = \sum_{k=1}^i ((x_k - x_{i+1}) \cdots (x_k - x_{n-1})(x_k + x_n) + (-1)^{n-i} x_{i+1} \cdots x_n) x_k^{-1} \partial_k \\ + (-1)^{n-i} \sum_{k=i+1}^n x_{i+1} \cdots \widehat{x_k} \cdots x_n \partial_k,$$

$$(5.9) \quad \psi_{i,n+j}^{D_n} = \sum_{k=1}^i ((x_k - x_{i+1}) \cdots (x_k - x_n)(x_k + x_n) \cdots (x_k + x_{n-j}) \\ + (-1)^{n-i+1} x_{i+1} \cdots x_{n-1-j} x_{n-j}^2 \cdots x_n^2) x_k^{-1} \partial_k \\ + (-1)^{n-i+1} x_{n-j} \cdots x_n \sum_{k=i+1}^n x_{i+1} \cdots \widehat{x_k} \cdots x_n \partial_k \quad \text{for } 0 \leq j \leq n-1-i,$$

and

$$(5.10) \quad \psi_{n,2n-1-r}^{D_n} = \sum_{k=1}^r ((-1)^{n-r+1} (x_k - x_{r+1}) \cdots (x_k - x_{n-1})(x_k - x_n) + x_{r+1} \cdots x_n) x_k^{-1} \partial_k \\ + \sum_{k=r+1}^n x_{r+1} \cdots \widehat{x_k} \cdots x_n \partial_k \quad \text{for } 0 \leq r \leq n-1.$$

Proposition 5.5. *Let I be a lower ideal in $\Phi_{D_n}^+$ and h_I the associated Hessenberg function in (3.3). Then, $\psi_{i,h_I(i)}^{D_n}$ is an element of $D(\mathcal{A}_I)$ for $i = 1, \dots, n$.*

Proof. We fix i and put $j = h_I(i)$. We consider a Hessenberg function h such that $h(i) = j$, and choose the maximal Hessenberg function $h^{(i,j)}$ from such Hessenberg functions. The lower ideal associated with $h^{(i,j)}$ is denoted by $I^{(i,j)}$. Note that the lower ideal $I^{(i,j)}$ is the maximal lower ideal among lower ideals I such that $\alpha_{i,j+1} \notin I$ with respect to inclusion. It is enough to prove that $\psi_{i,j}^{D_n}$ belong to $D(\mathcal{A}_{I^{(i,j)}})$ because $D(\mathcal{A}_{I^{(i,j)}}) \subset D(\mathcal{A}_I)$.

Case 1 Suppose that $1 \leq i \leq n-2$ and $i \leq j \leq n-2$. Then the Hessenberg function $h^{(i,j)}$ is given by

$$h^{(i,j)}(t) = \begin{cases} j & \text{if } 1 \leq t \leq i, \\ 2n-1-t & \text{if } i+1 \leq t \leq n-1, \\ 2n-1-i & \text{if } t = n. \end{cases}$$


The picture of the Hessenberg function $h^{(i,j)}$ is shown in Figure 7. Here, $d_t = t + e_t = 2n-1-t$ for $1 \leq t \leq n-1$.

Let $\alpha \in I^{(i,j)}$. Since all shaded boxes except for the boxes (r, r) with $1 \leq r \leq n$ in Figure 7 correspond to coordinates of all positive roots in $I^{(i,j)}$, α is one of the following forms

$$(5.11) \quad x_k - x_\ell \quad (1 \leq k < \ell \leq i),$$

$$(5.12) \quad x_k - x_\ell \quad (1 \leq k \leq i < \ell \leq j),$$

$$(5.13) \quad x_k \pm x_\ell \quad (i < k < \ell \leq n).$$

FIGURE 7. The Hessenberg function $h^{(i,j)}$ for $1 \leq i \leq n-2$ and $i \leq j \leq n-2$.

From the formula (5.7), we have


$$\psi_{i,j}^{D_n}(\alpha) = \begin{cases} (x_k - x_{i+1}) \cdots (x_k - x_j) - (x_\ell - x_{i+1}) \cdots (x_\ell - x_j) & \text{if } \alpha \text{ is of the form (5.11),} \\ (x_k - x_{i+1}) \cdots (x_k - x_j) & \text{if } \alpha \text{ is of the form (5.12),} \\ 0 & \text{if } \alpha \text{ is of the form (5.13).} \end{cases}$$

One can see that $\psi_{i,j}^{D_n}(\alpha) \equiv 0 \pmod{\alpha}$ in both of cases.

Case 2 Suppose that $1 \leq i \leq n-1$ and $j = n-1$. In this case, the Hessenberg function $h^{(i,j)}$ is defined by

$$h^{(i,j)}(t) = \begin{cases} n-1 & \text{if } 1 \leq t \leq i, \\ 2n-1-t & \text{if } i+1 \leq t \leq n-1, \\ 2n-1 & \text{if } t = n. \end{cases}$$

The picture of the Hessenberg function $h^{(i,j)}$ is shown in Figure 8.

FIGURE 8. The Hessenberg function $h^{(i,n-1)}$ with $1 \leq i \leq n-1$.

In this case, $\alpha \in I^{(i,j)}$ is one of the following forms

$$(5.14) \quad x_k - x_\ell \quad (1 \leq k < \ell \leq i),$$

$$(5.15) \quad x_k - x_\ell \quad (1 \leq k \leq i < \ell \leq n-1),$$

$$(5.16) \quad x_k \pm x_\ell \quad (i < k < \ell \leq n),$$

$$(5.17) \quad x_k + x_n \quad (1 \leq k \leq i).$$

Using the formula (5.8), we show that $\psi_{i,j}^{D_n}(\alpha) \equiv 0 \pmod{\alpha}$. If α is of the form (5.14), then we have

$$\begin{aligned} \psi_{i,n-1}^{D_n}(\alpha) &= ((x_k - x_{i+1}) \cdots (x_k - x_{n-1})(x_k + x_n) + (-1)^{n-i} x_{i+1} \cdots x_n) x_k^{-1} \\ &\quad - ((x_\ell - x_{i+1}) \cdots (x_\ell - x_{n-1})(x_\ell + x_n) + (-1)^{n-i} x_{i+1} \cdots x_n) x_\ell^{-1} \\ &\equiv 0 \pmod{x_k - x_\ell}. \end{aligned}$$

If α is of the form (5.15), then

$$\begin{aligned} \psi_{i,n-1}^{D_n}(\alpha) &= ((x_k - x_{i+1}) \cdots (x_k - x_{n-1})(x_k + x_n) + (-1)^{n-i} x_{i+1} \cdots x_n) x_k^{-1} \\ &\quad - (-1)^{n-i} x_{i+1} \cdots \widehat{x}_\ell \cdots x_n \\ &\equiv (-1)^{n-i} x_{i+1} \cdots x_n x_\ell^{-1} - (-1)^{n-i} x_{i+1} \cdots \widehat{x}_\ell \cdots x_n \pmod{x_k - x_\ell} \\ &= 0. \end{aligned}$$

If α is of the form (5.16), then

$$\begin{aligned} \psi_{i,n-1}^{D_n}(\alpha) &= (-1)^{n-i} x_{i+1} \cdots \widehat{x}_k \cdots x_n \pm (-1)^{n-i} x_{i+1} \cdots \widehat{x}_\ell \cdots x_n \\ &\equiv 0 \pmod{x_k \pm x_\ell}. \end{aligned}$$

If α is of the form (5.17), then

$$\begin{aligned} \psi_{i,n-1}^{D_n}(\alpha) &= ((x_k - x_{i+1}) \cdots (x_k - x_{n-1})(x_k + x_n) + (-1)^{n-i} x_{i+1} \cdots x_n) x_k^{-1} \\ &\quad + (-1)^{n-i} x_{i+1} \cdots x_{n-1} \\ &\equiv (-1)^{n-i} x_{i+1} \cdots x_n (-x_n^{-1}) + (-1)^{n-i} x_{i+1} \cdots x_{n-1} \pmod{x_k + x_n} \\ &= 0. \end{aligned}$$

Case 3 Suppose that $1 \leq i \leq n-1$ and $n \leq j \leq 2n-1-i$. Then the Hessenberg function $h^{(i,j)}$ is given by

$$h^{(i,j)}(t) = \begin{cases} j & \text{if } 1 \leq t \leq i, \\ 2n-1-t & \text{if } i+1 \leq t \leq n-1, \\ 2n-1 & \text{if } t = n, \end{cases}$$

and the picture of the Hessenberg function $h^{(i,j)}$ is shown in Figure 9.




FIGURE 9. The Hessenberg function $h^{(i,j)}$ for $1 \leq i \leq n-1, n \leq j \leq 2n-1-i$.

Then, $\alpha \in I^{(i,j)}$ is one of the following forms

$$(5.18) \quad x_k - x_\ell \quad (1 \leq k < \ell \leq i),$$

$$(5.19) \quad x_k - x_\ell \quad (1 \leq k \leq i < \ell \leq n),$$

$$(5.20) \quad x_k + x_\ell \quad (1 \leq k \leq i, 2n - j \leq \ell \leq n),$$

$$(5.21) \quad x_k \pm x_\ell \quad (i < k < \ell \leq n).$$

Using the formula (5.9), we show that $\psi_{i,j}^{D_n}(\alpha) \equiv 0 \pmod{\alpha}$. We put $s = j - n$. Then $0 \leq s \leq n - 1 - i$ and $j = n + s$. If α is of the form (5.18), then we have

$$\begin{aligned} \psi_{i,n+s}^{D_n}(\alpha) &= ((x_k - x_{i+1}) \cdots (x_k - x_n)(x_k + x_n) \cdots (x_k + x_{n-s}) \\ &\quad + (-1)^{n-i+1} x_{i+1} \cdots x_{n-1-s} x_{n-s}^2 \cdots x_n^2) x_k^{-1} \\ &\quad - ((x_\ell - x_{i+1}) \cdots (x_\ell - x_n)(x_\ell + x_n) \cdots (x_\ell + x_{n-s}) \\ &\quad + (-1)^{n-i+1} x_{i+1} \cdots x_{n-1-s} x_{n-s}^2 \cdots x_n^2) x_\ell^{-1} \\ &\equiv 0 \pmod{x_k - x_\ell} \end{aligned}$$

If α is of the form (5.19), then

$$\begin{aligned} \psi_{i,n+s}^{D_n}(\alpha) &= ((x_k - x_{i+1}) \cdots (x_k - x_n)(x_k + x_n) \cdots (x_k + x_{n-s}) \\ &\quad + (-1)^{n-i+1} x_{i+1} \cdots x_{n-1-s} x_{n-s}^2 \cdots x_n^2) x_k^{-1} \\ &\quad - (-1)^{n-i+1} x_{n-s} \cdots x_n (x_{i+1} \cdots \widehat{x}_\ell \cdots x_n) \\ &= ((x_k - x_{i+1}) \cdots (x_k - x_n)(x_k + x_n) \cdots (x_k + x_{n-s}) \\ &\quad + (-1)^{n-i+1} x_{i+1} \cdots x_{n-1-s} x_{n-s}^2 \cdots x_n^2 \\ &\quad - (-1)^{n-i+1} x_{i+1} \cdots x_{n-1-s} x_{n-s}^2 \cdots x_n^2 (x_k x_\ell^{-1})) x_k^{-1} \\ &\equiv 0 \pmod{x_k - x_\ell}. \end{aligned}$$

If α is of the form (5.20), then $i < 2n - j \leq \ell$ because of the condition for Case 3. So, we have

$$\begin{aligned} \psi_{i,n+s}^{D_n}(\alpha) &= ((x_k - x_{i+1}) \cdots (x_k - x_n)(x_k + x_n) \cdots (x_k + x_{n-s}) \\ &\quad + (-1)^{n-i+1} x_{i+1} \cdots x_{n-1-s} x_{n-s}^2 \cdots x_n^2) x_k^{-1} \\ &\quad + (-1)^{n-i+1} x_{n-s} \cdots x_n (x_{i+1} \cdots \widehat{x}_\ell \cdots x_n) \\ &\equiv 0 \pmod{x_k + x_\ell}. \end{aligned}$$


If α is of the form (5.21), then

$$\begin{aligned} \psi_{i,n+s}^{D_n}(\alpha) &= (-1)^{n-i+1} x_{n-s} \cdots x_n (x_{i+1} \cdots \widehat{x}_k \cdots x_n) \pm (-1)^{n-i+1} x_{n-s} \cdots x_n (x_{i+1} \cdots \widehat{x}_\ell \cdots x_n) \\ &\equiv 0 \pmod{x_k \pm x_\ell}. \end{aligned}$$

Case 4 Suppose that $i = n$ and $n \leq j \leq 2n - 1$. We put $r = 2n - 1 - j$. Then $0 \leq r \leq n - 1$ and the Hessenberg function $h^{(i,j)}$ is given as follows

$$h^{(i,j)}(t) = \begin{cases} n & \text{if } 1 \leq t \leq r, \\ 2n - 1 - t & \text{if } r + 1 \leq t \leq n - 1, \\ 2n - 1 - r & \text{if } t = n. \end{cases}$$

The picture of the Hessenberg function $h^{(i,j)}$ is shown in Figure 10.


 FIGURE 10. The Hessenberg function $h^{(n, 2n-1-r)}$ with $0 \leq r \leq n-1$.

Similarly to Case 2, one can see that $\psi_{n, 2n-1-r}^{D_n}(\alpha) \equiv 0 \pmod{\alpha}$ for any $\alpha \in I^{(i, j)}$.

Therefore, it follows from Cases 1, 2, 3, 4 that $\psi_{i, j}^{D_n}$ is an element of $D(\mathcal{A}_{I^{(i, j)}})$, and this completes the proof. \square

Theorem 5.6. *The derivations $\{\psi_{i, j}^{D_n} \mid 1 \leq i \leq n-1, i \leq j \leq 2n-i-1\} \cup \{\psi_{n, j}^{D_n} \mid n \leq j \leq 2n-1\}$ form uniform bases for the ideal arrangements of type D_n .*

Proof. From Proposition 5.5 we have $\{\psi_{i, h_I(i)}^{D_n} \mid i \in [n]\} \subset D(\mathcal{A}_I)$ for any lower ideal I . Since $\deg(\psi_{i, j}^{D_n}) = j - i$, it is enough to prove that $\psi_{1, h(1)}^{D_n}, \dots, \psi_{n, h(n)}^{D_n}$ are linearly independent over \mathcal{R} for any Hessenberg function h for type D_n by Theorem 2.1. We prove this by splitting into two cases. Let

$$(5.22) \quad i_h^{D_n} := \min\{i \in [n-1] \mid h(i) \geq n-1\}.$$

Case 1 Suppose that $h(n) \geq 2n - i_h^{D_n} - 1$. We prove the linear independence of $\psi_{1, h(1)}^{D_n}, \dots, \psi_{n, h(n)}^{D_n}$ over \mathcal{R} by induction on n . As the base case $n = 4$, the claim is straightforward. Now we assume that $n > 4$ and the claim holds for $n-1$. We first note that a Hessenberg function h for type D_n induces the Hessenberg function h' for type D_{n-1} defined by

$$\begin{aligned} h'(i-1) &= h(i) - 1 \quad \text{for } 2 \leq i \leq n-1, \\ h'(n-1) &= \min\{h(n) - 1, 2n-3\}. \end{aligned}$$

The picture of h' is obtained from that of h by removing the 1-st row. One see that $i_{h'}^{D_{n-1}} \geq i_h^{D_n}$, so we have $h'(n-1) \geq 2(n-1) - i_{h'}^{D_{n-1}} - 1$. That is, h' is a Hessenberg function for type D_{n-1} satisfying the condition for Case 1.

We consider the following surjection

$$(5.23) \quad \begin{aligned} \bigoplus_{k=1}^n \mathbb{R}[x_1, \dots, x_n] \partial_k &\longrightarrow \bigoplus_{k=2}^n \mathbb{R}[x_2, \dots, x_n] \partial_k \\ \psi &\longmapsto \sum_{k=1}^n f_k(x_1, \dots, x_n) \partial_k \longmapsto \sum_{k=2}^n f_k(0, x_2, \dots, x_n) \partial_k \end{aligned}$$

and the image of ψ under the map is denoted by $\bar{\psi}$. Then we have

$$(5.24) \quad \overline{\psi_{i, j}^{D_n}} = \psi_{i-1, j-1}^{D_{n-1}}(x_2, \dots, x_n) \quad \text{for } i \in [n-1], i \leq j \leq 2n-1-i,$$

$$(5.25) \quad \overline{\psi_{n, j}^{D_n}} = \begin{cases} \psi_{n-1, j-1}^{D_{n-1}}(x_2, \dots, x_n) & \text{if } i \in [n-1], i \leq j \leq 2n-1-i, \\ 0 & \text{if } j = 2n-1. \end{cases}$$

Suppose that

$$(5.26) \quad \sum_{k=1}^n f_k(x_1, \dots, x_n) \psi_{k, h(k)}^{D_n} = 0$$

for $f_k(x_1, \dots, x_n) \in \mathbb{R}[x_1, \dots, x_n]$ such that all $f_k(x_1, \dots, x_n)$ ($1 \leq k \leq n$) have no common divisor.

Case 1-1 Assume that $h(1) < n - 1$. Then we have $\psi_{0, h(1)-1}^{D_{n-1}} = 0$ and $h(n) < 2n - 1$ by the definition of the Hessenberg function. Applying the map in (5.23) to both sides of (5.26), we obtain from (5.24) and (5.25)

$$\sum_{k=2}^n f_k(0, x_2, \dots, x_n) \psi_{k-1, h'(k-1)}^{D_{n-1}}(x_2, \dots, x_n) = 0.$$

By the inductive assumption, we obtain $f_k(0, x_2, \dots, x_n) = 0$ for $k = 2, \dots, n$. Hence, we can write $f_k(x_1, \dots, x_n) = x_1 f'_k(x_1, \dots, x_n)$ for some $f'_k(x_1, \dots, x_n) \in \mathbb{R}[x_1, \dots, x_n]$. Comparing the coefficient of ∂_1 in (5.26), we have

$$f_1(x_1, \dots, x_n)(x_1 - x_2) \cdots (x_1 - x_{h(1)}) + x_1(f'_2(x_1, \dots, x_n) + \cdots) = 0.$$

Substituting $x_1 = 0$ into the equation above, we obtain

$$f_1(0, x_2, \dots, x_n)(-x_2) \cdots (-x_{h(1)}) = 0.$$

Hence, we have $f_1(0, x_2, \dots, x_n) = 0$ which implies $f_1(x_1, \dots, x_n)$ is divisible by x_1 . However, all $f_k(x_1, \dots, x_n)$ have no common divisor, so $f_k(x_1, \dots, x_n) = 0$ for $k = 1, \dots, n$.

Case 1-2 Assume that $h(1) = n - 1$. Then we have

$$(5.27) \quad \psi_{0, h(1)-1}^{D_{n-1}}(x_2, \dots, x_n) = (-1)^{n-1} \sum_{k=2}^n x_2 \cdots \widehat{x_k} \cdots x_n \partial_k = (-1)^{n-1} \psi_{n-1, 2n-3}^{D_{n-1}}(x_2, \dots, x_n).$$

Since $h(1) = n - 1$, we have $i_h^{D_n} = 1$. Thus, $h(n) \geq 2n - i_h^{D_n} - 1 = 2n - 2$, i.e. $h(n) = 2n - 1$ or $h(n) = 2n - 2$.

If $h(n) = 2n - 1$, then $\overline{\psi_{n, h(n)}^{D_n}} = 0$ from (5.25) and $h'(n - 1) = 2n - 3$. Applying the map in (5.23) to both sides of (5.26), we obtain from (5.27) and (5.24)

$$(-1)^{n-1} f_1(0, x_2, \dots, x_n) \psi_{n-1, h'(n-1)}^{D_{n-1}}(x_2, \dots, x_n) + \sum_{k=2}^{n-1} f_k(0, x_2, \dots, x_n) \psi_{k-1, h'(k-1)}^{D_{n-1}}(x_2, \dots, x_n) = 0.$$

By the inductive assumption, we obtain $f_k(0, x_2, \dots, x_n) = 0$ for $k = 1, \dots, n - 1$. Hence, we can write $f_k(x_1, \dots, x_n) = x_1 f'_k(x_1, \dots, x_n)$ for some $f'_k(x_1, \dots, x_n) \in \mathbb{R}[x_1, \dots, x_n]$. Comparing the coefficient of ∂_1 in (5.26), we have

$$x_1(f'_1(x_1, \dots, x_n) + \cdots) + f_n(x_1, \dots, x_n)x_2 \cdots x_n = 0.$$

Substituting $x_1 = 0$ into the equation above, we obtain $f_n(0, x_2, \dots, x_n) = 0$ which implies $f_n(x_1, \dots, x_n)$ is divisible by x_1 . Therefore, $f_k(x_1, \dots, x_n) = 0$ for $k = 1, \dots, n$ since all $f_k(x_1, \dots, x_n)$ have no common divisor.

If $h(n) = 2n-2$, then $h'(n-1) = 2n-3$. Hence by (5.25), $\overline{\psi_{n,h(n)}^{D_n}} = \psi_{n-1,h'(n-1)}^{D_{n-1}}(x_2, \dots, x_n)$. Applying the map in (5.23) to both sides of (5.26), we obtain from (5.27) and (5.24)

$$\begin{aligned} &((-1)^{n-1}f_1(0, x_2, \dots, x_n) + f_n(0, x_2, \dots, x_n))\psi_{n-1,h'(n-1)}^{D_{n-1}}(x_2, \dots, x_n) \\ &+ \sum_{k=2}^{n-1} f_k(0, x_2, \dots, x_n)\psi_{k-1,h'(k-1)}^{D_{k-1}}(x_2, \dots, x_n) = 0. \end{aligned}$$

By the inductive assumption, we obtain

$$(5.28) \quad f_1(0, x_2, \dots, x_n) = (-1)^n f_n(0, x_2, \dots, x_n),$$

$$(5.29) \quad f_k(0, x_2, \dots, x_n) = 0 \quad \text{for } k = 2, \dots, n-1.$$

Hence, we can write $f_k(x_1, \dots, x_n) = x_1 f'_k(x_1, \dots, x_n)$ for $k = 2, \dots, n-1$. Comparing the coefficient of ∂_1 in (5.26),

$$\begin{aligned} &f_1(x_1, \dots, x_n)((x_1 - x_2) \cdots (x_1 - x_{n-1})(x_1 + x_n) + (-1)^{n-1}x_2 \cdots x_n)x_1^{-1} \\ &+ x_1(f'_2(x_1, \dots, x_n) + \cdots) + f_n(x_1, \dots, x_n)((-1)^n(x_1 - x_2) \cdots (x_1 - x_n) + x_2 \cdots x_n)x_1^{-1} = 0. \end{aligned}$$

Substituting $x_1 = 0$ into the equation above, we obtain

$$\begin{aligned} &f_1(0, x_2, \dots, x_n)\left(\sum_{k=2}^{n-1}(-x_2) \cdots \widehat{(-x_k)} \cdots (-x_{n-1})(x_n) + (-x_2) \cdots (-x_{n-1})\right) \\ &+ (-1)^n f_n(0, x_2, \dots, x_n)\left(\sum_{k=2}^n(-x_2) \cdots \widehat{(-x_k)} \cdots (-x_n)\right) = 0. \end{aligned}$$

Hence by (5.28), we have $2f_1(0, x_2, \dots, x_n)(-x_2) \cdots (-x_{n-1}) = 0$ which implies that $f_1(0, x_2, \dots, x_n) = f_n(0, x_2, \dots, x_n) = 0$. From this together with (5.29), we conclude $f_k(x_1, \dots, x_n) = 0$ for $k = 1, \dots, n$ because all $f_k(x_1, \dots, x_n)$ have no common divisor.

Case 1-3 Assume that $h(1) \geq n$. It follows from similar discussion on Case 1-2 that $f_k(x_1, \dots, x_n) = 0$ for $k = 1, \dots, n$.

Case 2 Suppose that $h(n) \leq 2n - i_h^{D_n} - 1$. We prove the linear independence of $\psi_{1,h(1)}^{D_n}, \dots, \psi_{n,h(n)}^{D_n}$ over \mathcal{R} by induction on $m_h := 2n - i_h^{D_n} - 1 - h(n)$. As the base case $m_h = 0$, namely $h(n) = 2n - i_h^{D_n} - 1$, the claim follows from Case 1. Now we assume that $m_h > 0$, that is,

$$(5.30) \quad h(n) < 2n - i_h^{D_n} - 1$$

and the claim holds for any Hessenberg function \tilde{h} for type D_n with $m_{\tilde{h}} = m_h - 1$.

For the given Hessenberg function h with (5.30), we can define a Hessenberg function \tilde{h} by

$$\tilde{h}(i) = \begin{cases} h(i) & \text{for } i = 1, \dots, n-1, \\ h(n) + 1 & \text{for } i = n. \end{cases}$$

By the definition we have $i_h^{D_n} = i_{\tilde{h}}^{D_n}$, so $m_{\tilde{h}} = m_h - 1$. Hence, $\psi_{1,\tilde{h}(1)}^{D_n}, \dots, \psi_{n,\tilde{h}(n)}^{D_n}$ are linearly independent over \mathcal{R} by the inductive assumption. Suppose that

$$(5.31) \quad \sum_{k=1}^n f_k \psi_{k,h(k)}^{D_n} = 0$$

for $f_k \in \mathbb{R}[x_1, \dots, x_n]$. Since $\psi_{n, \tilde{h}(n)}^{D_n} = \alpha_{n, \tilde{h}(n)} \psi_{n, h(n)}^{D_n} + (-1)^{\tilde{h}(n)-n} \psi_{2n-\tilde{h}(n), n}^{D_n}$ by (5.4), we have

$$(5.32) \quad \sum_{k=1}^{n-1} g_k \psi_{k, \tilde{h}(k)}^{D_n} + f_n \psi_{n, \tilde{h}(n)}^{D_n} = (-1)^{\tilde{h}(n)-n} f_n \psi_{2n-\tilde{h}(n), n}^{D_n},$$

where $g_k = \alpha_{n, \tilde{h}(n)} f_k$ for $k \in [n-1]$. We show that $\psi_{2n-\tilde{h}(n), n}^{D_n}$ can be written as a linear combination of $\psi_{1, \tilde{h}(1)}^{D_n}, \dots, \psi_{n-1, \tilde{h}(n-1)}^{D_n}$ over \mathcal{R} . We first note that $1 \leq 2n - \tilde{h}(n) \leq n$ because $n \leq \tilde{h}(n) \leq 2n-1$ by the definition (2) of a Hessenberg function. If $2n - \tilde{h}(n) = n$, then $n \leq \tilde{h}(n) = \tilde{h}(n) - 1 = n - 1$, yielding a contradiction. If $2n - \tilde{h}(n) = 1$, then $h(n) = \tilde{h}(n) - 1 = 2n - 2$. The assumption (5.30) now implies $i_h^{D_n} < 1$, a contradiction. Hence, we have

$$(5.33) \quad 2 \leq 2n - \tilde{h}(n) \leq n - 1,$$

$$(5.34) \quad 1 \leq 2n - \tilde{h}(n) - 1 \leq n - 2.$$

By the definition (5.22) and assumption (5.30), we have $h(2n - \tilde{h}(n)) \geq h(i_h^{D_n}) \geq n - 1$ and $h(2n - \tilde{h}(n) - 1) \geq h(i_h^{D_n}) \geq n - 1$. Also, by the definition (5) of a Hessenberg function we have $h(2n - \tilde{h}(n)) \leq n$ and $h(2n - \tilde{h}(n) - 1) \leq n$. Thus, both of the values $h(2n - \tilde{h}(n))$ and $h(2n - \tilde{h}(n) - 1)$ are equal to $n - 1$ or n .

If $h(2n - \tilde{h}(n)) = n$, then it follows from (5.33) and the definition of \tilde{h} that

$$\psi_{2n-\tilde{h}(n), n}^{D_n} = \psi_{2n-\tilde{h}(n), h(2n-\tilde{h}(n))}^{D_n} = \psi_{2n-\tilde{h}(n), \tilde{h}(2n-\tilde{h}(n))}^{D_n}.$$

If $h(2n - \tilde{h}(n)) = n - 1$, then we have $h(2n - \tilde{h}(n) - 1) = n - 1$ by the definition (3) of a Hessenberg function. From (5.33), (5.34), and the definition of \tilde{h} together with (5.3), we obtain

$$\begin{aligned} \psi_{2n-\tilde{h}(n), n}^{D_n} &= \psi_{2n-\tilde{h}(n)-1, n-1}^{D_n} + \alpha_{i, n} \psi_{2n-\tilde{h}(n), n-1}^{D_n} = \psi_{2n-\tilde{h}(n)-1, h(2n-\tilde{h}(n)-1)}^{D_n} + \alpha_{i, n} \psi_{2n-\tilde{h}(n), h(2n-\tilde{h}(n))}^{D_n} \\ &= \psi_{2n-\tilde{h}(n)-1, \tilde{h}(2n-\tilde{h}(n)-1)}^{D_n} + \alpha_{i, n} \psi_{2n-\tilde{h}(n), \tilde{h}(2n-\tilde{h}(n))}^{D_n}. \end{aligned}$$

Therefore, $\psi_{2n-\tilde{h}(n), n}^{D_n}$ can be written as a linear combination of $\psi_{1, \tilde{h}(1)}^{D_n}, \dots, \psi_{n-1, \tilde{h}(n-1)}^{D_n}$ over \mathcal{R} . This together with (5.32) implies $f_n = 0$ because $\psi_{1, \tilde{h}(1)}^{D_n}, \dots, \psi_{n, \tilde{h}(n)}^{D_n}$ are linearly independent over \mathcal{R} by the inductive assumption. Hence by (5.31), we have $\sum_{k=1}^{n-1} f_k \psi_{k, \tilde{h}(k)}^{D_n} = 0$. However, since $\psi_{k, \tilde{h}(k)}^{D_n} = \psi_{k, h(k)}^{D_n}$ for $1 \leq k \leq n-1$, from the \mathcal{R} -linearly independence of $\psi_{1, \tilde{h}(1)}^{D_n}, \dots, \psi_{n-1, \tilde{h}(n-1)}^{D_n}$ we obtain $f_k = 0$ for $1 \leq k \leq n-1$.

Therefore, we proved that $\psi_{1, \tilde{h}(1)}^{D_n}, \dots, \psi_{n, \tilde{h}(n)}^{D_n}$ are linearly independent over \mathcal{R} for any Hessenberg function h for type D_n by Cases 1 and 2, and this completes the proof. \square

The derivations $\psi_{i, j}^{D_n}$ are uniform bases and have an explicit formula. Unfortunately, they are not of the form (4.7) because of the form of the derivations $\psi_{0, j}$ for $n \leq j \leq 2n-3$. In order to give uniform bases $\tilde{\psi}_{i, j}^{D_n}$ of the form (4.7), we need to modify $\psi_{i, j}^{D_n}$ a little bit.

As the base case, when $j = i$, we define

$$\tilde{\psi}_{i, i}^{D_n} = \psi_{i, i}^{D_n} \quad \text{for } i \in [n].$$

Proceeding inductively, we define for $i = 1$

$$\begin{aligned}\tilde{\psi}_{1,j}^{D_n} &= \alpha_{1,j} \tilde{\psi}_{1,j-1}^{D_n} & \text{if } 1 < j \leq 2n-2 \text{ with } j \neq n-1, n, \\ \tilde{\psi}_{1,n-1}^{D_n} &= \alpha_{1,n-1} \tilde{\psi}_{1,n-2}^{D_n} + \xi_1^{D_n}, \\ \tilde{\psi}_{1,n}^{D_n} &= \alpha_{1,n} \tilde{\psi}_{1,n-1}^{D_n} + \xi_0^{D_n},\end{aligned}$$

and for $1 < i \leq n-1$

$$\begin{aligned}\tilde{\psi}_{i,j}^{D_n} &= \tilde{\psi}_{i-1,j-1}^{D_n} + \alpha_{i,j} \tilde{\psi}_{i,j-1}^{D_n} & \text{if } i < j \text{ with } j \neq n-1, \\ \tilde{\psi}_{i,n-1}^{D_n} &= \tilde{\psi}_{i-1,n-2}^{D_n} + \alpha_{i,n-1} \tilde{\psi}_{i,n-2}^{D_n} + \xi_i^{D_n} & \text{if } i < n-1,\end{aligned}$$

and for $i = n$

$$\tilde{\psi}_{n,j}^{D_n} = \alpha_{n,j} \tilde{\psi}_{n,j-1}^{D_n} + (-1)^{j-n} \tilde{\psi}_{2n-j,n}^{D_n} \quad \text{if } n+1 \leq j \leq 2n-1.$$

Lemma 5.7. *For all (i, j) , we have*

$$\tilde{\psi}_{i,j}^{D_n} \equiv \psi_{i,j}^{D_n} \pmod{D(\mathcal{A}_{\Phi_{D_n}^+})}.$$

In particular, $\{\psi_{i,h_I(i)}^{D_n} \mid i \in [n]\} \subset D(\mathcal{A}_I)$ for any lower ideal $I \subset \Phi_{D_n}^+$.

Proof. Since $\psi_{0,n-1}$ is an element of $D(\mathcal{A}_{\Phi_{D_n}^+})$, so is $\psi_{0,j}$ for $n \leq j \leq 2n-3$. From this together with the recursive formulas for $\{\psi_{i,j}\}$ and $\{\tilde{\psi}_{i,j}\}$, we obtain that $\tilde{\psi}_{i,j} \equiv \psi_{i,j} \pmod{D(\mathcal{A}_{\Phi_{D_n}^+})}$ for all (i, j) . For the rest, it follows from Proposition 5.5 and $D(\mathcal{A}_{\Phi_{D_n}^+}) \subset D(\mathcal{A}_I)$. \square

Lemma 5.8. *The derivation $\xi_i^{D_n}$ in (5.6) holds*

$$\begin{aligned}\xi_i^{D_n} &= -\frac{1}{2} \alpha_{i+1,n} \psi_{i+1,n-1}^{D_n} + (-1)^{n-i} \frac{1}{2} \alpha_{n,2n-1-i} \psi_{n,2n-2-i}^{D_n} \\ &= -\frac{1}{2} \alpha_{i+1,n} \tilde{\psi}_{i+1,n-1}^{D_n} + (-1)^{n-i} \frac{1}{2} \alpha_{n,2n-1-i} \tilde{\psi}_{n,2n-2-i}^{D_n}\end{aligned}$$

for $0 \leq i \leq n-2$.

Proof. For the second equality, it follows from the recursive formulas for $\{\psi_{i,j}\}$ and $\{\tilde{\psi}_{i,j}\}$ that $\psi_{i+1,n-1}^{D_n} = \tilde{\psi}_{i+1,n-1}^{D_n}$ and $\psi_{n,2n-2-i}^{D_n} = \tilde{\psi}_{n,2n-2-i}^{D_n}$.

We show the first equality. From the formulas (5.8) and (5.10) we have

$$\begin{aligned}\psi_{i+1,n-1}^{D_n} &= \sum_{k=1}^{i+1} ((x_k - x_{i+2}) \cdots (x_k - x_{n-1})(x_k + x_n) + (-1)^{n-i-1} x_{i+2} \cdots x_n) x_k^{-1} \partial_k \\ &\quad + (-1)^{n-i-1} \sum_{k=i+2}^n x_{i+2} \cdots \widehat{x_k} \cdots x_n \partial_k, \\ \psi_{n,2n-2-i}^{D_n} &= \sum_{k=1}^{i+1} ((-1)^{n-i} (x_k - x_{i+2}) \cdots (x_k - x_{n-1})(x_k - x_n) + x_{i+2} \cdots x_n) x_k^{-1} \partial_k \\ &\quad + \sum_{k=i+2}^n x_{i+2} \cdots \widehat{x_k} \cdots x_n \partial_k.\end{aligned}$$

Noting that $\alpha_{i+1,n} = x_{i+1} - x_n$ and $\alpha_{n,2n-1-i} = x_{i+1} + x_n$, we obtain

$$\begin{aligned}
& -\alpha_{i+1,n}\psi_{i+1,n-1}^{D_n} + (-1)^{n-i}\alpha_{n,2n-1-i}\psi_{n,2n-2-i}^{D_n} \\
&= \sum_{k=1}^i ((x_k - x_{i+2}) \cdots (x_k - x_{n-1})x_k \cdot 2x_n - (x_k - x_{i+2}) \cdots (x_k - x_{n-1})x_n \cdot 2x_{i+1} \\
&\quad + (-1)^{n-i}x_{i+2} \cdots x_n \cdot 2x_{i+1})x_k^{-1}\partial_k + ((-1)^{n-i}x_{i+2} \cdots x_n \cdot 2x_{i+1})x_{i+1}^{-1}\partial_{i+1} \\
&\quad + (-1)^{n-i} \sum_{k=i+2}^n x_{i+2} \cdots \widehat{x_k} \cdots x_n \cdot 2x_{i+1}\partial_k \\
&= \sum_{k=1}^i (2(x_k - x_{i+1})(x_k - x_{i+2}) \cdots (x_k - x_{n-1})x_n + (-1)^{n-i}2x_{i+1}x_{i+2} \cdots x_n)x_k^{-1}\partial_k \\
&\quad + (-1)^{n-i}2 \left(\sum_{k=i+1}^n x_{i+1} \cdots \widehat{x_k} \cdots x_n \partial_k \right) \\
&= 2\xi_i^{D_n}
\end{aligned}$$

as desired. \square

For type D_n we have $\text{ht}(\Phi_{D_n}^+) = 2n - 3$. Also, the set Λ_m in (3.6) for $m = 2k, 2k + 1$ is given by

$$(5.35) \quad \Lambda_m = \begin{cases} [n - k - 1] \cup \{n\} & \text{if } 1 \leq m \leq n - 1, \\ [n - k - 1] & \text{if } n \leq m \leq 2n - 3. \end{cases}$$

Proposition 5.9. *The derivations $\{\tilde{\psi}_{i,i+m}^{D_n} \mid 0 \leq m \leq 2n - 3, i \in \Lambda_m\}$ have the following expressions. As the base case $m = 0$,*

$$\begin{aligned}
\tilde{\psi}_{i,i}^{D_n} &= \alpha_i^* & \text{for } i = 1, \dots, n - 2, \\
\tilde{\psi}_{i,i}^{D_n} &= 2\alpha_i^* & \text{for } i = n - 1, n.
\end{aligned}$$

For m with $1 \leq m \leq n - 1$,

$$\begin{aligned}
\tilde{\psi}_{i,i+m}^{D_n} &= \sum_{j=1}^i \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} & \text{for } 1 \leq i < n - m - 1, \\
\tilde{\psi}_{i,i+m}^{D_n} &= \sum_{j=1}^{n-m-1} \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} - \frac{1}{2}\alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} + (-1)^{m+1} \frac{1}{2}\alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n} \\
&& \text{for } i = n - m - 1,
\end{aligned}$$

$$\begin{aligned}
\tilde{\psi}_{i,i+m}^{D_n} &= \sum_{\substack{j=1 \\ j \neq n-m}}^i \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} + \frac{1}{2}\alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} + (-1)^{m+1} \frac{1}{2}\alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n} \\
&& \text{for } n - m \leq i \leq n - 1,
\end{aligned}$$

$$\tilde{\psi}_{n,n+m}^{D_n} = (-1)^m \left(\sum_{j=1}^{n-m-1} \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} + \frac{1}{2}\alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} \right) + \frac{1}{2}\alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n}.$$

For m with $n \leq m \leq 2n - 3$, we have

$$\tilde{\psi}_{i,i+m}^{D_n} = \sum_{j=1}^i \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n}.$$

Proof. When $m = 0$, it is clear that $\tilde{\psi}_{i,i}^{D_n} = \alpha_i^*$ for $i \in [n-2]$ and $\tilde{\psi}_{i,i}^{D_n} = 2\alpha_i^*$ for $i = n-1, n$. We prove the claim for $m > 0$ by induction on i . As the base case $i = 1$, if $1 < j \leq 2n-2$ with $j \neq n-1, n$, then

$$\tilde{\psi}_{1,j}^{D_n} = \alpha_{1,j} \tilde{\psi}_{1,j-1}^{D_n}.$$

If $j = n-1$, then we have

$$\tilde{\psi}_{1,n-1}^{D_n} = \alpha_{1,n-1} \tilde{\psi}_{1,n-2}^{D_n} + \xi_1^{D_n} = \alpha_{1,n-1} \tilde{\psi}_{1,n-2}^{D_n} - \frac{1}{2} \alpha_{2,n} \tilde{\psi}_{2,n-1}^{D_n} + (-1)^{n-1} \frac{1}{2} \alpha_{n,2n-2} \tilde{\psi}_{n,2n-3}^{D_n}$$

by Lemma 5.8. Similarly, if $j = n$, then

$$\tilde{\psi}_{1,n}^{D_n} = \alpha_{1,n} \tilde{\psi}_{1,n-1}^{D_n} + \xi_0^{D_n} = \frac{1}{2} \alpha_{1,n} \tilde{\psi}_{1,n-1}^{D_n} + (-1)^n \frac{1}{2} \alpha_{n,2n-1} \tilde{\psi}_{n,2n-2}^{D_n}.$$

Now we assume that $i > 1$ and the claim holds for $i-1$. We first consider the case $1 \leq m \leq n-1$. If $1 \leq i < n-m-1$, then by the inductive assumption we have

$$\tilde{\psi}_{i,i+m}^{D_n} = \tilde{\psi}_{i-1,i+m-1}^{D_n} + \alpha_{i,j} \tilde{\psi}_{i,i+m-1}^{D_n} = \sum_{j=1}^i \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n}.$$

If $i = n-m-1$, then from Lemma 5.8

$$\begin{aligned} \tilde{\psi}_{i,i+m}^{D_n} &= \tilde{\psi}_{i-1,i+m-1}^{D_n} + \alpha_{i,i+m} \tilde{\psi}_{i,i+m-1}^{D_n} + \xi_i^{D_n} = \tilde{\psi}_{n-m-2,n-2}^{D_n} + \alpha_{n-m-1,n-1} \tilde{\psi}_{n-m-1,n-2}^{D_n} + \xi_{n-m-1}^{D_n} \\ &= \sum_{j=1}^{n-m-1} \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} + \xi_{n-m-1}^{D_n} \\ &= \sum_{j=1}^{n-m-1} \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} - \frac{1}{2} \alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} + (-1)^{m+1} \frac{1}{2} \alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n}. \end{aligned}$$

If $i = n-m$, then we have

$$\begin{aligned} \tilde{\psi}_{i,i+m}^{D_n} &= \tilde{\psi}_{i-1,i+m-1}^{D_n} + \alpha_{i,i+m} \tilde{\psi}_{i,i+m-1}^{D_n} = \tilde{\psi}_{n-m-1,n-1}^{D_n} + \alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} \\ &= \sum_{j=1}^{n-m-1} \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} - \frac{1}{2} \alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} + (-1)^{m+1} \frac{1}{2} \alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n} + \alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} \\ &= \sum_{j=1}^{n-m-1} \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} + \frac{1}{2} \alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} + (-1)^{m+1} \frac{1}{2} \alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n}. \end{aligned}$$

If $n - m < i \leq n - 1$, then by the inductive assumption

$$\begin{aligned}
\tilde{\psi}_{i,i+m}^{D_n} &= \tilde{\psi}_{i-1,i+m-1}^{D_n} + \alpha_{i,j} \tilde{\psi}_{i,i+m-1}^{D_n} \\
&= \sum_{\substack{j=1 \\ j \neq n-m}}^{i-1} \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} + \frac{1}{2} \alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} + (-1)^{m+1} \frac{1}{2} \alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n} + \alpha_{i,j} \tilde{\psi}_{i,i+m-1}^{D_n} \\
&= \sum_{\substack{j=1 \\ j \neq n-m}}^i \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} + \frac{1}{2} \alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} + (-1)^{m+1} \frac{1}{2} \alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n}.
\end{aligned}$$

If $i = n$, then we have

$$\begin{aligned}
\tilde{\psi}_{n,n+m}^{D_n} &= \alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n} + (-1)^m \tilde{\psi}_{n-m,n}^{D_n} \\
&= \alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n} + (-1)^m \left(\sum_{j=1}^{n-m-1} \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} + \frac{1}{2} \alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} \right. \\
&\quad \left. + (-1)^{m+1} \frac{1}{2} \alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n} \right) \\
&= (-1)^m \left(\sum_{j=1}^{n-m-1} \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n} + \frac{1}{2} \alpha_{n-m,n} \tilde{\psi}_{n-m,n-1}^{D_n} \right) + \frac{1}{2} \alpha_{n,n+m} \tilde{\psi}_{n,n+m-1}^{D_n}.
\end{aligned}$$

Next we consider the case $n \leq m \leq 2n - 3$. Then, by the inductive assumption we have

$$\begin{aligned}
\tilde{\psi}_{i,i+m}^{D_n} &= \tilde{\psi}_{i-1,i+m-1}^{D_n} + \alpha_{i,i+m} \tilde{\psi}_{i,i+m-1}^{D_n} \\
&= \sum_{j=1}^i \alpha_{j,j+m} \tilde{\psi}_{j,j+m-1}^{D_n}.
\end{aligned}$$

Therefore, we proved the desired equalities. \square

Motivated by Proposition 5.9, we define a matrix $P_m^{D_n}$ of size $|\Lambda_m| = \lambda_m$ as follows. As the base case, when $m = 0$,

$$P_0^{D_n} = \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & & 2 & \\ & & & & 2 \end{pmatrix}.$$

For m with $1 \leq m < n - 1$, when $m = 2k + 1$ odd, we define

$$P_{2k+1}^{D_n} = \begin{pmatrix} 1 & & & & & & & & \\ \vdots & \ddots & & & & & & & \\ 1 & \cdots & 1 & & & & & & \\ 1 & \cdots & 1 & 1 & -\frac{1}{2} & 0 & \cdots & 0 & \frac{1}{2} \\ 1 & \cdots & 1 & 1 & \frac{1}{2} & 0 & \cdots & 0 & \frac{1}{2} \\ 1 & \cdots & 1 & 1 & \frac{1}{2} & 1 & \ddots & \vdots & \vdots \\ \vdots & & \vdots & \vdots & \vdots & \vdots & \ddots & 0 & \vdots \\ 1 & \cdots & 1 & 1 & \frac{1}{2} & 1 & \cdots & 1 & \frac{1}{2} \\ -1 & \cdots & -1 & -1 & -\frac{1}{2} & 0 & \cdots & 0 & \frac{1}{2} \end{pmatrix},$$

$\underbrace{\hspace{10em}}_{n-2k-3} \qquad \underbrace{\hspace{10em}}_k$

and when $m = 2k + 2$ even, we define

$$P_{2k+2}^{D_n} = \begin{pmatrix} 1 & & & & & & & & \\ \vdots & \ddots & & & & & & & \\ 1 & \cdots & 1 & & & & & & \\ 1 & \cdots & 1 & 1 & -\frac{1}{2} & 0 & \cdots & 0 & -\frac{1}{2} \\ 1 & \cdots & 1 & 1 & \frac{1}{2} & 0 & \cdots & 0 & -\frac{1}{2} \\ 1 & \cdots & 1 & 1 & \frac{1}{2} & 1 & \ddots & \vdots & \vdots \\ \vdots & & \vdots & \vdots & \vdots & \vdots & \ddots & 0 & \vdots \\ 1 & \cdots & 1 & 1 & \frac{1}{2} & 1 & \cdots & 1 & -\frac{1}{2} \\ 1 & \cdots & 1 & 1 & \frac{1}{2} & 0 & \cdots & 0 & \frac{1}{2} \end{pmatrix}.$$

$\underbrace{\hspace{10em}}_{n-2k-4} \qquad \underbrace{\hspace{10em}}_k$

For $m = n - 1$, when $m = 2k + 1$ odd, we define

$$P_{n-1}^{D_n} = \begin{pmatrix} \frac{1}{2} & 0 & \cdots & 0 & \frac{1}{2} \\ \frac{1}{2} & 1 & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & 0 & \vdots \\ \frac{1}{2} & 1 & \cdots & 1 & \frac{1}{2} \\ -\frac{1}{2} & 0 & \cdots & 0 & \frac{1}{2} \end{pmatrix},$$

$\underbrace{\hspace{10em}}_k$

and when $m = 2k + 2$ even, we define

$$P_{n-1}^{D_n} = \begin{pmatrix} \frac{1}{2} & 0 & \cdots & 0 & -\frac{1}{2} \\ \frac{1}{2} & 1 & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & 0 & \vdots \\ \frac{1}{2} & 1 & \cdots & 1 & -\frac{1}{2} \\ \frac{1}{2} & 0 & \cdots & 0 & \frac{1}{2} \end{pmatrix}.$$

$\underbrace{\hspace{10em}}_k$

For m with $n \leq m \leq 2n - 3$, we define

$$P_m^{D_n} = \begin{pmatrix} 1 & & \\ \vdots & \ddots & \\ 1 & \cdots & 1 \end{pmatrix}.$$

From Proposition 5.9 we obtain

$$(5.36) \quad [\tilde{\psi}_{i,i+m}^{D_n}]_{i \in \Lambda_m} = P_m [\alpha_{i,i+m} \tilde{\psi}_{i,i+m-1}^{D_n}]_{i \in \Lambda_m}$$

for any m with $1 \leq m \leq 2n - 3$ where we think of indices for rows and columns of the matrix P_m as the set Λ_m in (5.35), and we arrange them as in increasing order. One can see that $\det P_m = 1$ if $m \neq n - 1$ and $\det P_{n-1} = \frac{1}{2}$, so we obtain $P_m \in \text{GL}(\Lambda_m, \mathbb{Q})$ for all $1 \leq m \leq 2n - 3$. From this together with Lemma 5.7, we obtain the following theorem by Proposition 3.4 (see also Remark 3.5).

Theorem 5.10. *The derivations $\{\tilde{\psi}_{i,i+m}^{D_n} \mid 0 \leq m \leq 2n - 3, i \in \Lambda_m\}$ form uniform bases for the lower ideals of type D_n . Furthermore, $P_m^{D_n}$ ($0 \leq m \leq 2n - 3$) are the invertible matrices associated with the uniform bases $\{\tilde{\psi}_{i,i+m}^{D_n} \mid 0 \leq m \leq 2n - 1, i \in \Lambda_m\}$.*

5.5. Type G_2 . Let \mathfrak{t} be the hyperplane in $V = \mathbb{R}^3$ defined by the linear function $x_1 + x_2 + x_3$. Then we have

$$\mathcal{R} = \text{Sym}(\mathfrak{t}^*) = \mathbb{R}[x_1, x_2, x_3] / (x_1 + x_2 + x_3).$$

We set the exponents e_1, e_2 as

$$e_1 = 5, \quad e_2 = 1$$

We arrange all positive roots of G_2 as shown in Figure 11.

$x_1 - x_2$	$-x_1 + x_3$	$-x_2 + x_3$	$x_1 - 2x_2 + x_3$	$-x_1 - x_2 + 2x_3$
	$-2x_1 + x_2 + x_3$			

FIGURE 11. The arrangement of all positive roots for type G_2 .

In Figure 11 the partial order \preceq on $\Phi_{G_2}^+$ is defined as follows:

- (1) if a root α is left-adjacent to a root β , then $\alpha \prec \beta$,
- (2) if a root α is lower-adjacent to a root β , then $\alpha \prec \beta$.

For two positive roots α, β , we define $\alpha \preceq \beta$ if there exist positive roots $\gamma_0, \dots, \gamma_N$ such that $\alpha = \gamma_0 \prec \gamma_1 \prec \dots \prec \gamma_N = \beta$. We denote positive roots by

$$\begin{aligned} \alpha_{1,2} &= x_1 - x_2, & \alpha_{1,3} &= -x_1 + x_3, & \alpha_{1,4} &= -x_2 + x_3, \\ \alpha_{1,5} &= x_1 - 2x_2 + x_3, & \alpha_{1,6} &= -x_1 - x_2 + 2x_3, \\ \alpha_{2,3} &= -2x_1 + x_2 + x_3. \end{aligned}$$

Note that $\alpha_{i,j} \preceq \alpha_{k,\ell}$ iff $i \geq k$ and $j \leq \ell$. The decomposition $\Phi_{G_2}^+ = \Phi_1^+ \amalg \Phi_2^+$ shows that all positive roots are decomposed by positive roots of each rows in Figure 11. A **Hessenberg function for type G_2** is defined to be a function $h : \{1, 2\} \rightarrow \{1, 2, \dots, 6\}$ satisfying the following conditions

- (1) $1 \leq h(1) \leq 6$ and $2 \leq h(2) \leq 3$,
- (2) if $h(1) \geq 3$, then $h(2) = 3$.

Under the decomposition $\Phi_{G_2}^+ = \Phi_1^+ \amalg \Phi_2^+$, one can see that the set of lower ideals $I \subset \Phi_{G_2}^+$ and the set of Hessenberg functions h for type G_2 are in one-to-one correspondence which sends I to h_I in (3.3).

The uniform bases $\psi_{i,j}^{G_2} \in \text{Der } \mathcal{R}$ given in [4] are defined as follows. Let $\bar{\partial} := \partial_1 + \partial_2 + \partial_3$. Then, we define

$$\begin{aligned}\psi_{1,1}^{G_2} &= -\partial_2 + \partial_3, \\ \psi_{2,2}^{G_2} &= \partial_3 - \frac{1}{3}\bar{\partial}, \\ \psi_{1,j}^{G_2} &= \left(\prod_{\ell=2}^j \alpha_{1,\ell} \right) (-\partial_2 + \partial_3) \quad \text{for } 1 < j \leq 6, \\ \psi_{2,3}^{G_2} &= x_1(\partial_1 - \frac{1}{3}\bar{\partial}) + x_2(\partial_2 - \frac{1}{3}\bar{\partial}) + x_3(\partial_3 - \frac{1}{3}\bar{\partial}).\end{aligned}$$

Theorem 5.11. ([4, Theorem 10.17]) *The derivations $\{\psi_{1,1}^{G_2}, \dots, \psi_{1,6}^{G_2}, \psi_{2,2}^{G_2}, \psi_{2,3}^{G_2}\}$ form uniform bases for the ideal arrangements of type G_2 .*

For type G_2 the derivations $\psi_{i,j}^{G_2}$ satisfy the following recursive formula ([4])

$$\begin{aligned}\psi_{1,1}^{G_2} &= -\partial_2 + \partial_3, \\ \psi_{2,2}^{G_2} &= \partial_3 - \frac{1}{3}\bar{\partial}, \\ \psi_{i,j}^{G_2} &= \psi_{i-1,j-1}^{G_2} + \alpha_{i,j}\psi_{i,j-1}^{G_2} \quad \text{for } (i,j) \text{ with } i < j \leq i + e_i,\end{aligned}$$

where we take the convention $\psi_{0,*}^{G_2} = 0$ for any $*$. Note that $\text{ht}(\Phi_{G_2}^+) = 5$ and the set Λ_m in (3.6) is given by

$$\Lambda_m = \begin{cases} \{1, 2\} & \text{for } m = 0, 1, \\ \{1\} & \text{for } 2 \leq m \leq 5. \end{cases}$$

From the recursive formula above we obtain

$$\begin{aligned}\psi_{i,i}^{G_2} &= \alpha_i^* \quad \text{for } i = 1, 2, \\ \psi_{1,j}^{G_2} &= \alpha_{1,j}\psi_{1,j-1}^{G_2} \quad \text{for } 1 < j \leq 6, \\ \psi_{2,3}^{G_2} &= \alpha_{1,2}\psi_{1,1}^{G_2} + \alpha_{2,3}\psi_{2,2}^{G_2}.\end{aligned}$$

Hence, the invertible matrices $P_m^{G_2}$ associated with the uniform bases $\{\psi_{i,i+m}^{G_2} \mid 0 \leq m \leq 5, i \in \Lambda_m\}$ are given by

$$P_0^{G_2} = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}, \quad P_1^{G_2} = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}, \quad P_m^{G_2} = (1) \quad (2 \leq m \leq 5).$$

5.6. Type F_4 . Let \mathfrak{t} be the euclidean space $V = \mathbb{R}^4$ and we have

$$\mathcal{R} = \text{Sym}(\mathfrak{t}^*) = \mathbb{R}[x_1, x_2, x_3, x_4].$$

We set the exponents e_1, e_2, e_3, e_4 as

$$e_1 = 1, \quad e_2 = 11, \quad e_3 = 7, \quad e_4 = 5.$$

We take the simple roots $\alpha_1 = \frac{1}{2}(x_1 - x_2 - x_3 - x_4), \alpha_2 = x_2 - x_3, \alpha_3 = x_3 - x_4, \alpha_4 = x_4$ so that a labeling of the Dynkin diagram is as follows:




$x_2 - x_3$	$x_2 - x_4$	x_2	$\frac{1}{2}(x_1 + x_2 - x_3 - x_4)$	$x_2 + x_3$	$\frac{1}{2}(x_1 + x_2 + x_3 - x_4)$	$x_1 - x_4$	x_1	$x_1 + x_4$	$x_1 + x_3$	$x_1 + x_2$
	$x_3 - x_4$	x_3	$\frac{1}{2}(x_1 - x_2 + x_3 - x_4)$	$x_2 + x_4$	$\frac{1}{2}(x_1 + x_2 - x_3 + x_4)$	$x_1 - x_3$	$\frac{1}{2}(x_1 + x_2 + x_3 + x_4)$			
		x_4	$\frac{1}{2}(x_1 - x_2 - x_3 + x_4)$	$x_3 + x_4$	$\frac{1}{2}(x_1 - x_2 + x_3 + x_4)$	$x_1 - x_2$				
			$\frac{1}{2}(x_1 - x_2 - x_3 - x_4)$							

FIGURE 12. The arrangement of all positive roots for type F_4 .

We arrange all positive roots of F_4 as shown in Figure 12.

In Figure 12 the partial order \preceq on $\Phi_{F_4}^+$ is defined as follows:

- (1) if a root α is left to a root β , then $\alpha \preceq \beta$, except that α, β are divided by a dotted line;
- (2) if a root α is lower-adjacent to a root β , then $\alpha \preceq \beta$;
- (3) if two roots α, β are divided by a dotted line, and a root γ is immediately to the northwest of α and a root δ is immediately to the southeast of β , then $\alpha \preceq \delta$ and $\gamma \preceq \beta$.



For two positive roots α, β , we define $\alpha \preceq \beta$ if there exist positive roots $\gamma_0, \dots, \gamma_N$ such that $\alpha = \gamma_0 \preceq \gamma_1 \preceq \dots \preceq \gamma_N = \beta$.

Now we fix a decomposition $\Phi_{F_4}^+ = \Phi_1^+ \amalg \Phi_2^+ \amalg \Phi_3^+ \amalg \Phi_4^+$ satisfying (3.1) and (3.2). The positive roots $\alpha_{i,j}$ are defined as

positive roots \ j	2	3	4	5	6	7
$\alpha_{1,j}$	$\frac{1}{2}(x_1 - x_2 - x_3 - x_4)$					
$\alpha_{2,j}$		$x_2 - x_3$	$x_2 - x_4$	x_2	$x_2 + x_4$	$x_2 + x_3$
$\alpha_{3,j}$			$x_3 - x_4$	x_3	$x_3 + x_4$	$\frac{1}{2}(x_1 - x_2 + x_3 + x_4)$
$\alpha_{4,j}$				x_4	$\frac{1}{2}(x_1 - x_2 - x_3 + x_4)$	$\frac{1}{2}(x_1 - x_2 + x_3 - x_4)$

positive roots \ j	8	9	10	11	12	13
$\alpha_{1,j}$						
$\alpha_{2,j}$	$\frac{1}{2}(x_1 + x_2 + x_3 - x_4)$	$\frac{1}{2}(x_1 + x_2 + x_3 + x_4)$	x_1	$x_1 + x_4$	$x_1 + x_3$	$x_1 + x_2$
$\alpha_{3,j}$	$x_1 - x_2$	$x_1 - x_3$	$x_1 - x_4$			
$\alpha_{4,j}$	$\frac{1}{2}(x_1 + x_2 - x_3 - x_4)$	$\frac{1}{2}(x_1 + x_2 - x_3 + x_4)$				

Motivated by this, we define the coordinate in type F_4 as shown in Figure 13.

(2, 2)	(2, 3)	(2, 4)	(2, 5)	(4, 8)	(2, 7)	(2, 8)	(3, 10)	(2, 10)	(2, 11)	(2, 12)	(2, 13)
	(3, 3)	(3, 4)	(3, 5)	(4, 7)	(2, 6)	(4, 9)	(3, 9)	(2, 9)			
		(4, 4)	(4, 5)	(4, 6)	(3, 6)	(3, 7)	(3, 8)				
			(1, 1)	(1, 2)							

FIGURE 13. The coordinate in type F_4 .

Note that $\text{ht}(\Phi_{F_4}^+) = 11$ and the set Λ_m in (3.6) is given by

$$(5.37) \quad \Lambda_m = \begin{cases} \{1, 2, 3, 4\} & \text{for } m = 0, 1, \\ \{2, 3, 4\} & \text{for } 2 \leq m \leq 5, \\ \{2, 3\} & \text{for } m = 6, 7, \\ \{2\} & \text{for } 8 \leq m \leq 11. \end{cases}$$

For the decomposition $\Phi_{F_4}^+ = \Phi_1^+ \coprod \Phi_2^+ \coprod \Phi_3^+ \coprod \Phi_4^+$ above, a **Hessenberg function for type F_4** is defined to be a function $h : \{1, \dots, 4\} \rightarrow \{1, \dots, 13\}$ satisfying the following conditions

- (1) $i \leq h(i) \leq i + e_i$ for $i = 1, 2, 3, 4$,
- (2) if $h(2) \geq k$, then $h(3) \geq k$ for $k = 4, 5, 6, 10$,
- (3) if $h(3) \geq k$, then $h(4) \geq k$ for $k = 5, 7, 9$,
- (4) if $h(4) \geq k$, then $h(1) = 2$ for $k = 6$,
- (5) if $h(4) \geq k$, then $h(3) \geq k - 2$ for $k = 7, 9$,
- (6) if $h(4) \geq k$, then $h(2) \geq k - 3$ for $k = 8, 9$,
- (7) if $h(2) \geq 8$, then $h(4) = 9$,
- (8) if $h(3) \geq 10$, then $h(2) \geq 8$.

Under this decomposition $\Phi_{F_4}^+ = \Phi_1^+ \coprod \Phi_2^+ \coprod \Phi_3^+ \coprod \Phi_4^+$, one can see that the set of lower ideals $I \subset \Phi_{F_4}^+$ and the set of Hessenberg functions h for type F_4 are in one-to-one correspondence which sends I to h_I in (3.3).

Remark 5.12. A Hessenberg function h such that $h(1) = 1, h(2) \leq 7, h(3) \leq 6, h(4) \leq 5$ for type F_4 is exactly the Hessenberg function h such that $h(1) = 1$ for type B_4 which is naturally identified with that of type B_3 .

We find uniform bases $\{\psi_{i,i+m}^{F_4} \in \text{Der } \mathcal{R} \mid 0 \leq m \leq 11, i \in \Lambda_m\}$ inductively. As the base case, when $m = 0$, we define

$$\psi_{i,i}^{F_4} = \alpha_i^* \quad \text{for } i = 1, 2, 3, 4.$$

Explicitly,

$$\begin{aligned} \psi_{1,1}^{F_4} &= 2\partial_1, \\ \psi_{2,2}^{F_4} &= \partial_1 + \partial_2, \\ \psi_{3,3}^{F_4} &= 2\partial_1 + \partial_2 + \partial_3, \\ \psi_{4,4}^{F_4} &= 3\partial_1 + \partial_2 + \partial_3 + \partial_4. \end{aligned}$$

Proceeding inductively, for $m > 0$ and $i \in \Lambda_m$ we write

$$(5.38) \quad \psi_{i,i+m}^{F_4} = \sum_{j \in \Lambda_m} p_{ij}^{(m)} \alpha_{j,j+m} \psi_{j,j+m-1}^{F_4}$$

for some rational numbers $p_{ij}^{(m)}$. We determine the rational numbers $p_{ij}^{(m)}$ such that all of the derivations $\psi_{i,i+m}$ form uniform bases. As the proof of Proposition 5.5, we consider the maximal lower ideal $I^{(i,i+m)}$ containing the root $\alpha_{i,i+m}$ with respect to the inclusion. Since $\psi_{i,i+m}^{F_4}(\alpha) \in \mathcal{R}\alpha$ for any $\alpha \in I^{(i,i+m)}$, by (5.38) the following has to hold

$$\sum_{j \in \Lambda_m} p_{ij}^{(m)} \alpha_{j,j+m} \psi_{j,j+m-1}^{F_4}(\alpha) \equiv 0 \quad \text{mod } \alpha.$$

Now we know an explicit formula for $\psi_{j,j+m-1}^{F_4}$ ($j \in \Lambda_m$) by inductive step, so we obtain a linear equation in $p_{ij}^{(m)}$ ($j \in \Lambda_m$) for each $\alpha \in I^{(i,i+m)}$. We computed a solution of the system of the linear equations in $p_{ij}^{(m)}$ ($j \in \Lambda_m$), and the result is given as follows. (We also checked the solution by using Maple⁶.) For each m with $0 \leq m \leq 11 = \text{ht}(\Phi_{F_4}^+)$, the matrix $P_m^{F_4} = (p_{ij}^{(m)})_{i,j \in \Lambda_m}$ is given by

$$\begin{aligned} P_0^{F_4} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_1^{F_4} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix}, \quad P_2^{F_4} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad P_3^{F_4} = \begin{pmatrix} 1 & -\frac{1}{2} & -1 \\ 1 & 1 & -1 \\ \frac{1}{2} & \frac{1}{2} & 1 \end{pmatrix}, \\ P_4^{F_4} &= \begin{pmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix}, \quad P_5^{F_4} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -\frac{1}{2} & \frac{1}{2} & 1 \end{pmatrix}, \quad P_6^{F_4} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad P_7^{F_4} = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}, \\ P_m^{F_4} &= (1) \quad (8 \leq m \leq 11). \end{aligned}$$

Here, we arrange indices for rows and columns of the matrix $P_m^{F_4}$ as in increasing order. Note that

$$\begin{aligned} [\psi_{i,i}^{F_4}]_{i \in \Lambda_0} &= P_0^{F_4} [\alpha_i^*]_{i \in \Lambda_0} \\ [\psi_{i,i+m}^{F_4}]_{i \in \Lambda_m} &= P_m^{F_4} [\alpha_{i,i+m} \psi_{i,i+m-1}^{F_4}]_{i \in \Lambda_m} \quad \text{for } 1 \leq m \leq 11. \end{aligned}$$

One can check that $\det P_m^{F_4} \neq 0$, so we obtain the following theorem by Proposition 3.4 (see also Remark 3.5).

Theorem 5.13. *The derivations $\{\psi_{i,i+m}^{F_4} \mid 0 \leq m \leq 11, i \in \Lambda_m\}$ form uniform bases for the ideal arrangements of type F_4 .*

5.7. Type E_n ($n = 6, 7, 8$). We consider types E_n ($n = 6, 7, 8$) by case-by-case to compute the invertible matrices.


5.7.1. Type E_8 . Let \mathfrak{t}_{E_8} be the euclidean space $V = \mathbb{R}^8$ and we have

$$\mathcal{R}_{E_8} = \text{Sym}(\mathfrak{t}_{E_8}^*) = \mathbb{R}[x_1, \dots, x_8].$$

We set the exponents $e_1^{E_8}, \dots, e_8^{E_8}$ as

$$e_1^{E_8} = 19, \quad e_2^{E_8} = 29, \quad e_3^{E_8} = 23, \quad e_4^{E_8} = 13, \quad e_5^{E_8} = 11, \quad e_6^{E_8} = 7, \quad e_7^{E_8} = 1, \quad e_8^{E_8} = 17.$$

We take the simple roots $\alpha_1 = \frac{1}{2}(x_1 - x_2 - x_3 - x_4 - x_5 - x_6 - x_7 - x_8)$, $\alpha_2 = x_2 - x_3$, $\alpha_3 = x_3 - x_4$, $\alpha_4 = x_4 - x_5$, $\alpha_5 = x_5 - x_6$, $\alpha_6 = x_6 - x_7$, $\alpha_7 = x_7 - x_8$, $\alpha_8 = x_7 + x_8$ so that a labeling of the Dynkin diagram is as follows:



⁶The program is available at <https://researchmap.jp/ehrhart/Database/>.

Then, the positive roots are the following forms

$$(5.39) \quad x_i \pm x_j \quad (1 \leq i, j \leq 8),$$

$$(5.40) \quad \frac{1}{2}(x_1 \pm x_2 \pm x_3 \pm x_4 \pm x_5 \pm x_6 \pm x_7 \pm x_8).$$

We arrange all positive roots of E_8 as shown in Figure 14. For simplicity, we denote the positive root $x_i \pm x_j$ of the form (5.39) by $i \pm j$. Also, we denote by $\frac{1}{2}(i_1 i_2 \dots i_k)$ the positive root of the form (5.40) such that coefficients of x_i are positive for $i = i_1, i_2, \dots, i_k$. For example, the notation $\frac{1}{2}(12478)$ means the positive root $\frac{1}{2}(x_1 + x_2 - x_3 + x_4 - x_5 - x_6 + x_7 + x_8)$. Using the simple notations above, we arrange all positive roots of E_8 as shown in Figure 14.




FIGURE 14. The arrangement of all positive roots for type E_8 .

In Figure 14 the partial order \preceq on $\Phi_{E_8}^+$ is defined as follows. In each area separated by dotted lines we have the following relation

- (1) if a root α is left-adjacent to a root β , then $\alpha \prec \beta$,
- (2) if a root α is lower-adjacent to a root β , then $\alpha \prec \beta$.

For the blocks (\bullet) and $(\bar{\bullet})$ with same shape where the symbol \bullet means a, b, \dots, k , we have the following relation

- (3) if a root α is in the block (\bullet) and β is in the same place as α in the block $(\bar{\bullet})$, then $\alpha \prec \beta$.

For two positive roots α, β , we define $\alpha \preceq \beta$ if there exist positive roots $\gamma_0, \dots, \gamma_N$ such that $\alpha = \gamma_0 \prec \gamma_1 \prec \dots \prec \gamma_N = \beta$.

Now we fix a decomposition $\Phi_{E_8}^+ = \coprod_{i=1}^8 \Phi_i^+$ satisfying (3.1) and (3.2). Using the simple notations above again, we define the positive roots $\alpha_{i,j}$ as follows:

positive roots $\setminus j$	2	3	4	5	6	7	8	9	10	11	12
$\alpha_{1,j}$	$\frac{1}{2}(1)$	$\frac{1}{2}(178)$	$\frac{1}{2}(168)$	$\frac{1}{2}(158)$	$\frac{1}{2}(148)$	$\frac{1}{2}(138)$	$\frac{1}{2}(137)$	$\frac{1}{2}(136)$	$\frac{1}{2}(135)$	$\frac{1}{2}(125)$	$\frac{1}{2}(124)$
$\alpha_{2,j}$		$2-3$	$2-4$	$2-5$	$2-6$	$2-7$	$2-8$	$2+7$	$2+6$	$2+5$	$2+4$
$\alpha_{3,j}$			$3-4$	$3-5$	$3-6$	$3-7$	$3-8$	$3+7$	$3+6$	$3+5$	$3+4$
$\alpha_{4,j}$				$4-5$	$4-6$	$4-7$	$4-8$	$4+7$	$4+6$	$4+5$	$\frac{1}{2}(145)$
$\alpha_{5,j}$					$5-6$	$5-7$	$5-8$	$5+7$	$5+6$	$\frac{1}{2}(156)$	$\frac{1}{2}(15678)$
$\alpha_{6,j}$						$6-7$	$6-8$	$6+7$	$\frac{1}{2}(167)$	$\frac{1}{2}(157)$	$\frac{1}{2}(147)$
$\alpha_{7,j}$							$7-8$				
$\alpha_{8,j}$								$7+8$	$6+8$	$5+8$	$4+8$

	13	14	15	16	17	18	19	20	21
$\alpha_{1,j}$	$\frac{1}{2}(12478)$	$\frac{1}{2}(12468)$	$\frac{1}{2}(12458)$	$\frac{1}{2}(12457)$	$\frac{1}{2}(12456)$	$\frac{1}{2}(1245678)$	$\frac{1}{2}(1235678)$	$\frac{1}{2}(1234678)$	
$\alpha_{2,j}$	$2+3$	$\frac{1}{2}(123)$	$\frac{1}{2}(12378)$	$\frac{1}{2}(12368)$	$\frac{1}{2}(12358)$	$\frac{1}{2}(12348)$	$\frac{1}{2}(12347)$	$\frac{1}{2}(12346)$	$\frac{1}{2}(12345)$
$\alpha_{3,j}$	$\frac{1}{2}(134)$	$\frac{1}{2}(13478)$	$\frac{1}{2}(13468)$	$\frac{1}{2}(13458)$	$\frac{1}{2}(13457)$	$\frac{1}{2}(13456)$	$\frac{1}{2}(1345678)$	$1-2$	$1-3$
$\alpha_{4,j}$	$\frac{1}{2}(14578)$	$\frac{1}{2}(14568)$	$\frac{1}{2}(14567)$	$\frac{1}{2}(13567)$	$\frac{1}{2}(13467)$				
$\alpha_{5,j}$	$\frac{1}{2}(14678)$	$\frac{1}{2}(13678)$	$\frac{1}{2}(13578)$	$\frac{1}{2}(13568)$					
$\alpha_{6,j}$	$\frac{1}{2}(146)$								
$\alpha_{7,j}$									
$\alpha_{8,j}$	$3+8$	$2+8$	$\frac{1}{2}(128)$	$\frac{1}{2}(127)$	$\frac{1}{2}(126)$	$\frac{1}{2}(12678)$	$\frac{1}{2}(12578)$	$\frac{1}{2}(12568)$	$\frac{1}{2}(12567)$

	22	23	24	25	26	27	28	29	30	31
$\alpha_{1,j}$										
$\alpha_{2,j}$	$\frac{1}{2}(1234578)$	$\frac{1}{2}(1234568)$	$\frac{1}{2}(1234567)$	$1-8$	$1+7$	$1+6$	$1+5$	$1+4$	$1+3$	$1+2$
$\alpha_{3,j}$	$1-4$	$1-5$	$1-6$	$1-7$	$1+8$					
$\alpha_{4,j}$										
$\alpha_{5,j}$										
$\alpha_{6,j}$										
$\alpha_{7,j}$										
$\alpha_{8,j}$	$\frac{1}{2}(12467)$	$\frac{1}{2}(12367)$	$\frac{1}{2}(12357)$	$\frac{1}{2}(12356)$						

Note that $\text{ht}(\Phi_{E_8}^+) = 29$ and the set Λ_m in (3.6) is given by

$$(5.41) \quad \Lambda_m = \begin{cases} \{1, 2, 3, 4, 5, 6, 7, 8\} & \text{for } m = 0, 1, \\ \{1, 2, 3, 4, 5, 6, 8\} & \text{for } 2 \leq m \leq 7, \\ \{1, 2, 3, 4, 5, 8\} & \text{for } 8 \leq m \leq 11, \\ \{1, 2, 3, 4, 8\} & \text{for } 12 \leq m \leq 13, \\ \{1, 2, 3, 8\} & \text{for } 14 \leq m \leq 17, \\ \{1, 2, 3\} & \text{for } 18 \leq m \leq 19, \\ \{2, 3\} & \text{for } 20 \leq m \leq 23, \\ \{2\} & \text{for } 24 \leq m \leq 29. \end{cases}$$

For the decomposition $\Phi_{E_8}^+ = \coprod_{i=1}^8 \Phi_i^+$ above, a **Hessenberg function for type E_8** is defined to be a function $h^{E_8} : \{1, \dots, 8\} \rightarrow \{1, \dots, 31\}$ satisfying the following conditions

- (1) $i \leq h^{E_8}(i) \leq i + e_i$ for $i = 1, \dots, 8$,
- (2) if $h^{E_8}(1) \geq k$, then $h^{E_8}(2) \geq k$ for $k = 11, 12, 20$,
- (3) if $h^{E_8}(1) \geq k$, then $h^{E_8}(3) \geq k + 1$ for $k = 8, 9, 10, 12, \dots, 18$,
- (4) if $h^{E_8}(1) \geq k$, then $h^{E_8}(4) \geq k + 2$ for $k = 10$,

- (5) if $h^{E_8}(1) \geq k$, then $h^{E_8}(6) \geq k + 4$ for $k = 8, 9$,
- (6) if $h^{E_8}(1) \geq k$, then $h^{E_8}(8) \geq k + 6$ for $k = 3, \dots, 7, 11, 13, 14, 16, 19$,
- (7) if $h^{E_8}(2) \geq k$, then $h^{E_8}(1) \geq k - 2$ for $k = 14, \dots, 17, 22$,
- (8) if $h^{E_8}(2) \geq k$, then $h^{E_8}(3) \geq k$ for $k = 4, \dots, 12, 25, 26$,
- (9) if $h^{E_8}(2) \geq k$, then $h^{E_8}(8) \geq k + 5$ for $k = 9, 19, 20$,
- (10) if $h^{E_8}(3) \geq k$, then $h^{E_8}(1) \geq k - 3$ for $k = 13, 21, 22, 23$,
- (11) if $h^{E_8}(3) \geq k$, then $h^{E_8}(2) \geq k - 2$ for $k = 24, 25$,
- (12) if $h^{E_8}(3) \geq k$, then $h^{E_8}(4) \geq k$ for $k = 5, \dots, 11, 17$,
- (13) if $h^{E_8}(3) \geq k$, then $h^{E_8}(5) \geq k + 1$ for $k = 14, 15$,
- (14) if $h^{E_8}(3) \geq k$, then $h^{E_8}(8) \geq k + 4$ for $k = 9$,
- (15) if $h^{E_8}(4) \geq k$, then $h^{E_8}(3) \geq k - 2$ for $k = 17$,
- (16) if $h^{E_8}(4) \geq k$, then $h^{E_8}(5) \geq k$ for $k = 6, \dots, 10, 13, 16$,
- (17) if $h^{E_8}(4) \geq k$, then $h^{E_8}(6) \geq k + 1$ for $k = 12$,
- (18) if $h^{E_8}(4) \geq k$, then $h^{E_8}(8) \geq k + 3$ for $k = 9$,
- (19) if $h^{E_8}(5) \geq k$, then $h^{E_8}(1) \geq k - 5$ for $k = 14, 15$,
- (20) if $h^{E_8}(5) \geq k$, then $h^{E_8}(4) \geq k - 2$ for $k = 15, 16$,
- (21) if $h^{E_8}(5) \geq k$, then $h^{E_8}(6) \geq k$ for $k = 7, 8, 9, 11, 13$,
- (22) if $h^{E_8}(5) \geq k$, then $h^{E_8}(8) \geq k + 2$ for $k = 9$,
- (23) if $h^{E_8}(6) \geq k$, then $h^{E_8}(1) \geq k - 6$ for $k = 10, 11, 12$,
- (24) if $h^{E_8}(6) \geq k$, then $h^{E_8}(4) \geq k - 3$ for $k = 12, 13$,
- (25) if $h^{E_8}(6) \geq k$, then $h^{E_8}(5) \geq k - 2$ for $k = 11, 13$,
- (26) if $h^{E_8}(6) \geq k$, then $h^{E_8}(7) \geq k$ for $k = 8$,
- (27) if $h^{E_8}(6) \geq k$, then $h^{E_8}(8) \geq k + 1$ for $k = 9$,
- (28) if $h^{E_8}(8) \geq k$, then $h^{E_8}(1) \geq k - 8$ for $k = 15, 16, 17, 19, 22, 24, 25$,
- (29) if $h^{E_8}(8) \geq k$, then $h^{E_8}(2) \geq k - 7$ for $k = 14, 16, 17, 23, 24$,
- (30) if $h^{E_8}(8) \geq k$, then $h^{E_8}(3) \geq k - 6$ for $k = 13$,
- (31) if $h^{E_8}(8) \geq k$, then $h^{E_8}(4) \geq k - 5$ for $k = 12, 21, 22$,
- (32) if $h^{E_8}(8) \geq k$, then $h^{E_8}(5) \geq k - 4$ for $k = 11, 18, 19, 20$,
- (33) if $h^{E_8}(8) \geq k$, then $h^{E_8}(6) \geq k - 3$ for $k = 10$.

Under this decomposition $\Phi_{E_8}^+ = \coprod_{i=1}^8 \Phi_i^+$, one can see that the set of lower ideals $I \subset \Phi_{E_8}^+$ and the set of Hessenberg functions h for type E_8 are in one-to-one correspondence which sends I to h_I in (3.3).

By similar discussion on the case of type F_4 , we obtain uniform bases $\{\psi_{i,i+m}^{E_8} \in \text{Der } \mathcal{R} \mid 0 \leq m \leq 29, i \in \Lambda_m\}$ as follows. For each m with $0 \leq m \leq 29 = \text{ht}(\Phi_{E_8}^+)$, we define the matrix $P_m^{E_8} = (p_{ij}^{(m)})_{i,j \in \Lambda_m}$ by

$$P_0^{E_8} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_1^{E_8} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$\begin{aligned}
P_2^{E_8} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ -1 & 1 & 1 & 1 & 1 & -1 & -1 \\ -1 & 1 & 1 & 1 & 1 & 1 & -1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}, \quad P_3^{E_8} = \begin{pmatrix} 1 & -\frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} & -\frac{1}{4} & -\frac{1}{2} & -\frac{1}{4} \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & -1 & -\frac{1}{2} & \frac{1}{2} \\ 2 & 2 & 2 & 2 & 1 & -1 & 1 \\ 1 & 1 & 1 & 1 & \frac{1}{2} & 1 & \frac{1}{2} \\ 2 & -4 & -4 & -4 & -2 & -1 & 1 \end{pmatrix}, \\
P_4^{E_8} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & -\frac{1}{2} & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ -\frac{2}{3} & 1 & 1 & -\frac{1}{3} & 0 & \frac{1}{3} & -\frac{1}{6} \\ -2 & 3 & 3 & 1 & 0 & 1 & -\frac{1}{2} \\ -2 & 3 & 3 & 1 & 1 & 3 & -\frac{1}{2} \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 4 & 6 & 6 & 2 & 0 & -2 & 1 \end{pmatrix}, \quad P_5^{E_8} = \begin{pmatrix} 1 & -\frac{3}{8} & -\frac{3}{16} & -\frac{1}{8} & -\frac{1}{8} & -\frac{1}{4} & -\frac{1}{32} \\ \frac{2}{3} & 1 & -\frac{3}{4} & -\frac{1}{12} & -\frac{1}{12} & -\frac{1}{6} & \frac{1}{12} \\ \frac{4}{3} & 2 & 1 & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{3} & \frac{1}{6} \\ 2 & 3 & \frac{3}{2} & 1 & -\frac{3}{2} & -3 & \frac{1}{4} \\ 2 & 3 & \frac{3}{2} & 1 & 1 & -3 & \frac{1}{4} \\ 1 & \frac{3}{2} & \frac{3}{4} & \frac{1}{2} & \frac{1}{2} & 1 & \frac{1}{8} \\ 8 & -18 & -9 & -1 & -1 & -2 & 1 \end{pmatrix}, \\
P_6^{E_8} &= \begin{pmatrix} 1 & \frac{3}{4} & 0 & 0 & 0 & 0 & \frac{1}{16} \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ \frac{8}{3} & 4 & 1 & 0 & 0 & 0 & 1/6 \\ 16 & 18 & 3 & 1 & 0 & -2 & 1 \\ -8 & -6 & 0 & 0 & 1 & 2 & -\frac{1}{2} \\ -4 & -3 & 0 & 0 & 0 & 1 & -\frac{1}{4} \\ 0 & 12 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_7^{E_8} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \frac{1}{16} \\ 0 & 1 & 0 & 0 & 0 & 0 & \frac{1}{12} \\ \frac{8}{3} & 2 & 1 & 0 & 0 & 0 & \frac{1}{3} \\ 0 & 6 & 3 & 1 & 1 & 2 & \frac{1}{2} \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ -8 & -3 & -\frac{3}{2} & -\frac{1}{2} & \frac{1}{2} & 1 & -\frac{3}{4} \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \\
P_8^{E_8} &= \begin{pmatrix} 1 & 0 & 0 & \frac{1}{16} & \frac{1}{8} & \frac{1}{16} \\ 0 & 1 & 0 & 0 & 0 & \frac{1}{12} \\ \frac{4}{3} & 2 & 1 & \frac{1}{12} & \frac{1}{6} & \frac{1}{4} \\ -16 & 0 & 0 & 1 & -2 & -1 \\ -8 & 0 & 0 & \frac{1}{2} & 1 & -\frac{1}{2} \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_9^{E_8} = \begin{pmatrix} 1 & \frac{3}{2} & \frac{3}{4} & 0 & \frac{1}{8} & -\frac{1}{8} \\ -\frac{1}{3} & 1 & -\frac{1}{4} & 0 & -\frac{1}{24} & \frac{1}{24} \\ -\frac{2}{3} & 2 & 1 & 0 & -\frac{1}{12} & \frac{1}{12} \\ -\frac{16}{3} & -8 & -4 & 1 & \frac{4}{3} & -\frac{4}{3} \\ -4 & -6 & -3 & 0 & 1 & -1 \\ 4 & 6 & 3 & 0 & 1/2 & 1 \end{pmatrix}, \\
P_{10}^{E_8} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & \frac{1}{4} \\ -\frac{1}{3} & 1 & 0 & 0 & 0 & -\frac{1}{12} \\ \frac{2}{3} & 0 & 1 & 0 & 0 & \frac{1}{6} \\ -\frac{16}{3} & 0 & -8 & 1 & \frac{4}{3} & -\frac{8}{3} \\ -4 & 0 & -6 & 0 & 1 & -2 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_{11}^{E_8} = \begin{pmatrix} 1 & -3 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & \frac{1}{8} & \frac{1}{6} & -\frac{1}{6} \\ 0 & 0 & 0 & 1 & 0 & 0 \\ -8 & 24 & -6 & \frac{3}{4} & 1 & -1 \\ 0 & 0 & -6 & \frac{3}{4} & 1 & 1 \end{pmatrix},
\end{aligned}$$

$$\begin{aligned}
P_{12}^{E_8} &= \begin{pmatrix} 1 & -3 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ \frac{8}{3} & -8 & 1 & \frac{1}{4} & \frac{1}{6} \\ \frac{16}{3} & -16 & 0 & 1 & \frac{2}{3} \\ 8 & -24 & 0 & 0 & 1 \end{pmatrix}, \quad P_{13}^{E_8} = \begin{pmatrix} 1 & -3 & -\frac{3}{4} & -\frac{3}{16} & 0 \\ 0 & 1 & 0 & 0 & 0 \\ \frac{4}{3} & -4 & 1 & -\frac{1}{4} & 0 \\ \frac{16}{3} & -16 & 4 & 1 & \frac{4}{3} \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \\
P_{14}^{E_8} &= \begin{pmatrix} 1 & -6 & 0 & -\frac{1}{4} \\ \frac{1}{6} & 1 & 0 & \frac{1}{24} \\ \frac{2}{3} & -4 & 1 & -\frac{1}{6} \\ 4 & -24 & 0 & 1 \end{pmatrix}, \quad P_{15}^{E_8} = \begin{pmatrix} 1 & 0 & 0 & \frac{1}{4} \\ 0 & 1 & 0 & \frac{1}{24} \\ \frac{2}{3} & 0 & 1 & \frac{1}{6} \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_{16}^{E_8} = \begin{pmatrix} 1 & 6 & 0 & \frac{1}{4} \\ 0 & 1 & 0 & 0 \\ \frac{2}{3} & 4 & 1 & \frac{1}{6} \\ 0 & 24 & 0 & 1 \end{pmatrix}, \\
P_{17}^{E_8} &= \begin{pmatrix} 1 & 0 & \frac{3}{2} & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 4 & 24 & 6 & 1 \end{pmatrix}, \quad P_{18}^{E_8} = \begin{pmatrix} 1 & 0 & \frac{3}{2} \\ -\frac{1}{6} & 1 & -\frac{1}{4} \\ 0 & 0 & 1 \end{pmatrix}, \quad P_{19}^{E_8} = \begin{pmatrix} 1 & -6 & \frac{3}{2} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\
P_{20}^{E_8} &= P_{21}^{E_8} = \begin{pmatrix} 1 & \frac{1}{4} \\ 0 & 1 \end{pmatrix}, \quad P_{22}^{E_8} = P_{23}^{E_8} = \begin{pmatrix} 1 & 0 \\ 4 & 1 \end{pmatrix}, \quad P_m^{E_8} = (1) \quad (24 \leq m \leq 29).
\end{aligned}$$

Here, we think of indexes for rows and columns of the matrix $P_m^{E_8}$ as Λ_m given in (5.41), and we arrange them as in increasing order. We set

$$\begin{aligned}
[\psi_{i,i}^{E_8}]_{i \in \Lambda_0} &= P_0^{E_8} [\alpha_i^*]_{i \in \Lambda_0}, \\
[\psi_{i,i+m}^{E_8}]_{i \in \Lambda_m} &= P_m^{E_8} [\alpha_{i,i+m} \psi_{i,i+m-1}^{E_8}]_{i \in \Lambda_m} \quad \text{for } 1 \leq m \leq 29.
\end{aligned}$$

Note that $\psi_{i,i}^{E_8} = \alpha_i^*$ for $1 \leq i \leq 8$, namely

$$\begin{aligned}
\psi_{1,1}^{E_8} &= 2\partial_1, \\
\psi_{2,2}^{E_8} &= \partial_1 + \partial_2, \\
\psi_{3,3}^{E_8} &= 2\partial_1 + \partial_2 + \partial_3, \\
\psi_{4,4}^{E_8} &= 3\partial_1 + \partial_2 + \partial_3 + \partial_4, \\
\psi_{5,5}^{E_8} &= 4\partial_1 + \partial_2 + \partial_3 + \partial_4 + \partial_5, \\
\psi_{6,6}^{E_8} &= 5\partial_1 + \partial_2 + \partial_3 + \partial_4 + \partial_5 + \partial_6, \\
\psi_{7,7}^{E_8} &= \frac{1}{2}(5\partial_1 + \partial_2 + \partial_3 + \partial_4 + \partial_5 + \partial_6 + \partial_7 - \partial_8), \\
\psi_{8,8}^{E_8} &= \frac{1}{2}(7\partial_1 + \partial_2 + \partial_3 + \partial_4 + \partial_5 + \partial_6 + \partial_7 + \partial_8).
\end{aligned}$$

We can check by using Maple that $\{\psi_{i,h_I(i)}^{E_8} \mid 1 \leq i \leq 8\} \subset D(\mathcal{A}_I)$ for any lower ideal $I \subset \Phi_{E_8}^+$. From this together with $\det P_m^{E_8} \neq 0$ ($1 \leq m \leq 29$), we obtain the following theorem by Proposition 3.4 (see also Remark 3.5).

Theorem 5.14. *The derivations $\{\psi_{i,i+m}^{E_8} \mid 0 \leq m \leq 29, i \in \Lambda_m\}$ form uniform bases for the ideal arrangements of type E_8 .*

5.7.2. *Type E_7 .* Let \mathfrak{t}_{E_7} be the hyperplane in $\mathfrak{t}_{E_8} = \mathbb{R}^8$ defined by the linear function $\alpha_2^* = x_1 + x_2$:

$$\mathfrak{t}_{E_7} = \{(x_1, \dots, x_8) \in \mathbb{R}^8 \mid x_1 + x_2 = 0\}.$$

Then we have

$$\mathcal{R}_{E_7} = \text{Sym}(\mathfrak{t}_{E_7}^*) = \mathbb{R}[x_1, \dots, x_8]/(x_1 + x_2).$$

Taking the exponents $e_1^{E_7}, e_3^{E_7}, \dots, e_8^{E_7}$ as

$$e_1^{E_7} = 9, e_3^{E_7} = 17, e_4^{E_7} = 13, e_5^{E_7} = 11, e_6^{E_7} = 7, e_7^{E_7} = 1, e_8^{E_7} = 5,$$

we define the positive root $\alpha_{i,j}$ ($i = 1, 3, \dots, 8, i < j \leq i + e_i^{E_7}$) for E_7 by the $\alpha_{i,j}$ for E_8 . The poset structure of the positive roots for type E_7 can be obtained from Figure 14 by removing the first row in Figure 14.

Note that $\text{ht}(\Phi_{E_7}^+) = 17$ and the set Λ_m in (3.6) is given by

$$(5.42) \quad \Lambda_m = \begin{cases} \{1, 3, 4, 5, 6, 7, 8\} & \text{for } m = 0, 1, \\ \{1, 3, 4, 5, 6, 8\} & \text{for } 2 \leq m \leq 5, \\ \{1, 3, 4, 5, 8\} & \text{for } m = 6, 7, \\ \{1, 3, 4, 8\} & \text{for } m = 8, 9, \\ \{1, 3, 8\} & \text{for } m = 10, 11, \\ \{1, 3\} & \text{for } m = 12, 13, \\ \{3\} & \text{for } 14 \leq m \leq 17. \end{cases}$$

A Hessenberg function h^{E_8} for type E_8 such that $h^{E_8}(2) = 2$ and $h^{E_8}(i) \leq i + e_i^{E_7}$ for $i = 1, 3, \dots, 8$ is called a **Hessenberg function for type E_7** , denoted by h^{E_7} . We may omit the 2nd value $h^{E_7}(2)$, namely h^{E_7} is a function on $\{1, 3, \dots, 8\}$ given by $h^{E_7}(i) = h^{E_8}(i)$ for some h^{E_8} with $h^{E_8}(i) \leq i + e_i^{E_7}$ for $i = 1, 3, \dots, 8$. Under the decomposition $\Phi_{E_7}^+ = \Phi_1^+ \coprod (\coprod_{i=3}^8 \Phi_i^+)$ above, one can see that the set of lower ideals $I \subset \Phi_{E_7}^+$ and the set of Hessenberg functions h for type E_7 are in one-to-one correspondence which sends I to h_I in (3.3).

To explain uniform bases $\{\psi_{i,i+m}^{E_7} \in \text{Der } \mathcal{R} \mid 0 \leq m \leq 17, i \in \Lambda_m\}$, we first define the matrix $P_m^{E_7} = (p_{ij}^{(m)})_{i,j \in \Lambda_m}$ ($0 \leq m \leq 17 = \text{ht}(\Phi_{E_7}^+)$) by

$$P_0^{E_7} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_1^{E_7} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$\begin{aligned}
P_2^{E_7} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \\ -1 & 1 & 1 & 1 & -1 & -1 \\ -1 & 1 & 1 & 1 & 1 & -1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}, \quad P_3^{E_7} = \begin{pmatrix} 1 & -\frac{1}{2} & -\frac{1}{2} & -\frac{1}{4} & -\frac{1}{2} & -\frac{1}{4} \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & -1 & -\frac{1}{2} & \frac{1}{2} \\ 2 & 2 & 2 & 1 & -1 & 1 \\ 1 & 1 & 1 & \frac{1}{2} & 1 & \frac{1}{2} \\ 2 & -4 & -4 & -2 & -1 & 1 \end{pmatrix}, \\
P_4^{E_7} &= \begin{pmatrix} 1 & 0 & 0 & 0 & -\frac{1}{2} & 0 \\ -\frac{2}{3} & 1 & -\frac{1}{3} & 0 & \frac{1}{3} & -\frac{1}{6} \\ -2 & 3 & 1 & 0 & 1 & -\frac{1}{2} \\ -2 & 3 & 1 & 1 & 3 & -\frac{1}{2} \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 4 & 6 & 2 & 0 & -2 & 1 \end{pmatrix}, \quad P_5^{E_7} = \begin{pmatrix} 1 & -\frac{3}{8} & -\frac{1}{8} & -\frac{1}{8} & -\frac{1}{4} & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 3 & 1 & -1 & -2 & 0 \\ 0 & 3 & 1 & 1 & -2 & 0 \\ 0 & \frac{3}{2} & \frac{1}{2} & \frac{1}{2} & 1 & 0 \\ 8 & -9 & -1 & -1 & -2 & 1 \end{pmatrix}, \\
P_6^{E_7} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ \frac{4}{3} & 1 & 0 & 0 & 0 \\ 16 & 6 & 1 & 0 & -2 \\ -8 & 0 & 0 & 1 & 2 \\ -4 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_7^{E_7} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ \frac{4}{3} & 1 & 0 & 0 & 0 \\ 8 & 6 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ -8 & -3 & -\frac{1}{2} & \frac{1}{2} & 1 \end{pmatrix}, \\
P_8^{E_7} &= \begin{pmatrix} 1 & 0 & \frac{1}{8} & \frac{1}{8} \\ \frac{2}{3} & 1 & \frac{1}{12} & \frac{1}{12} \\ -8 & 0 & 1 & -1 \\ -8 & 0 & 1 & 1 \end{pmatrix}, \quad P_9^{E_7} = \begin{pmatrix} 1 & \frac{3}{2} & 0 & \frac{1}{8} \\ 0 & 1 & 0 & 0 \\ -8 & -12 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_{10}^{E_7} = \begin{pmatrix} 1 & 0 & 0 \\ -12 & 1 & 1 \\ -12 & 0 & 1 \end{pmatrix}, \\
P_{11}^{E_7} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -12 & 1 & 1 \end{pmatrix}, \quad P_{12}^{E_7} = \begin{pmatrix} 1 & \frac{1}{12} \\ 0 & 1 \end{pmatrix}, \quad P_{13}^{E_7} = \begin{pmatrix} 1 & 0 \\ 12 & 1 \end{pmatrix}, \quad P_m^{E_7} = (1) \quad (14 \leq m \leq 17).
\end{aligned}$$

Here, we think of indices for rows and columns of the matrix $P_m^{E_7}$ as Λ_m given in (5.42), and we arrange them as in increasing order. We set

$$\begin{aligned}
[\psi_{i,i}^{E_7}]_{i \in \Lambda_0} &= P_0^{E_7}[\alpha_i^*]_{i \in \Lambda_0}, \\
[\psi_{i,i+m}^{E_7}]_{i \in \Lambda_m} &= P_m^{E_7}[\alpha_{i,i+m} \psi_{i,i+m-1}^{E_7}]_{i \in \Lambda_m} \quad \text{for } 1 \leq m \leq 17.
\end{aligned}$$

Note that $\psi_{i,i}^{E_7} = \alpha_i^*$ for $i = 1, 3, \dots, 8$, namely

$$\begin{aligned}
\psi_{1,1}^{E_7} &= \psi_{1,1}^{E_8} - \psi_{2,2}^{E_8} = \partial_1 - \partial_2, \\
\psi_{3,3}^{E_7} &= \psi_{3,3}^{E_8} - \frac{3}{2}\psi_{2,2}^{E_8} = \frac{1}{2}\partial_1 - \frac{1}{2}\partial_2 + \partial_3, \\
\psi_{4,4}^{E_7} &= \psi_{4,4}^{E_8} - 2\psi_{2,2}^{E_8} = \partial_1 - \partial_2 + \partial_3 + \partial_4, \\
\psi_{5,5}^{E_7} &= \psi_{5,5}^{E_8} - \frac{5}{2}\psi_{2,2}^{E_8} = \frac{3}{2}\partial_1 - \frac{3}{2}\partial_2 + \partial_3 + \partial_4 + \partial_5,
\end{aligned}$$

$$\begin{aligned}
\psi_{6,6}^{E_7} &= \psi_{6,6}^{E_8} - 3\psi_{2,2}^{E_8} = 2\partial_1 - 2\partial_2 + \partial_3 + \partial_4 + \partial_5 + \partial_6, \\
\psi_{7,7}^{E_7} &= \psi_{7,7}^{E_8} - \frac{3}{2}\psi_{2,2}^{E_8} = \frac{1}{2}(2\partial_1 - 2\partial_2 + \partial_3 + \partial_4 + \partial_5 + \partial_6 + \partial_7 - \partial_8), \\
\psi_{8,8}^{E_7} &= \psi_{8,8}^{E_8} - 2\psi_{2,2}^{E_8} = \frac{1}{2}(3\partial_1 - 3\partial_2 + \partial_3 + \partial_4 + \partial_5 + \partial_6 + \partial_7 + \partial_8).
\end{aligned}$$

We can check by using Maple that $\{\psi_{i,h_I(i)}^{E_7} \mid i = 1, 3, \dots, 8\} \subset D(\mathcal{A}_I)$ for any lower ideal $I \subset \Phi_{E_7}^+$. From this together with $\det P_m^{E_7} \neq 0$ ($1 \leq m \leq 17$), we obtain the following theorem by Proposition 3.4 (see also Remark 3.5).

Theorem 5.15. *The derivations $\{\psi_{i,i+m}^{E_7} \mid 0 \leq m \leq 17, i \in \Lambda_m\}$ form uniform bases for the ideal arrangements of type E_7 .*

5.7.3. *Type E_6 .* Let \mathfrak{t}_{E_6} be the hyperplane in \mathfrak{t}_{E_7} defined by the linear function $\alpha_3^* = \frac{1}{2}x_1 - \frac{1}{2}x_2 + x_3$:

$$\mathfrak{t}_{E_6} := \{(x_1, \dots, x_8) \in \mathfrak{t}_{E_7} \mid \frac{1}{2}x_1 - \frac{1}{2}x_2 + x_3 = 0\}.$$

Here, we remark that $\alpha_1^*, \alpha_3^*, \dots, \alpha_8^*$ is the dual basis of the simple roots $\alpha_1, \alpha_3, \dots, \alpha_8$ for type E_7 . Then we have

$$\mathcal{R}_{E_6} = \text{Sym}(\mathfrak{t}_{E_6}^*) = \mathbb{R}[x_1, \dots, x_8] / (x_1 + x_2, \frac{1}{2}x_1 - \frac{1}{2}x_2 + x_3).$$

Taking the exponents $e_1^{E_6}, e_4^{E_6}, \dots, e_8^{E_6}$ as

$$e_1^{E_6} = 5, \quad e_4^{E_6} = 11, \quad e_5^{E_6} = 8, \quad e_6^{E_6} = 7, \quad e_7^{E_6} = 1, \quad e_8^{E_6} = 4,$$

we define the positive root $\alpha_{i,j}$ ($i = 1, 4, \dots, 8, i < j \leq i + e_i^{E_6}$) for E_6 by the $\alpha_{i,j}$ for E_7 . The poset structure of the positive roots for type E_7 can be obtained from Figure 14 by removing the first row and the second row in Figure 14.

Note that $\text{ht}(\Phi_{E_6}^+) = 11$ and the set Λ_m in (3.6) is given by

$$(5.43) \quad \Lambda_m = \begin{cases} \{1, 4, 5, 6, 7, 8\} & \text{for } m = 0, 1, \\ \{1, 4, 5, 6, 8\} & \text{for } 2 \leq m \leq 4, \\ \{1, 4, 5, 8\} & \text{for } m = 5, \\ \{1, 4, 8\} & \text{for } m = 6, 7, \\ \{1, 8\} & \text{for } m = 8, \\ \{1\} & \text{for } 9 \leq m \leq 11. \end{cases}$$

A Hessenberg function h^{E_7} for type E_7 such that $h^{E_7}(3) = 3$ and $h^{E_7}(i) \leq i + e_i^{E_6}$ for $i = 1, 4, \dots, 8$ is called a **Hessenberg function for type E_6** , denoted by h^{E_6} . We may omit the 3rd value $h^{E_6}(3)$, namely h^{E_6} is a function on $\{1, 4, \dots, 8\}$ given by $h^{E_6}(i) = h^{E_7}(i)$ for some h^{E_7} with $h^{E_7}(i) \leq i + e_i^{E_6}$ for $i = 1, 4, \dots, 8$. Under the decomposition $\Phi_{E_6}^+ = \Phi_1^+ \coprod (\coprod_{i=4}^8 \Phi_i^+)$ above, one can see that the set of lower ideals $I \subset \Phi_{E_6}^+$ and the set of Hessenberg functions h for type E_6 are in one-to-one correspondence which sends I to h_I in (3.3).

As well as type E_7 , to give uniform bases $\{\psi_{i,i+m}^{E_6} \in \text{Der } \mathcal{R} \mid 0 \leq m \leq 11, i \in \Lambda_m\}$, we first define the matrix $P_m^{E_6} = (p_{ij}^{(m)})_{i,j \in \Lambda_m}$ ($0 \leq m \leq 11 = \text{ht}(\Phi_{E_6}^+)$) by

$$\begin{aligned}
P_0^{E_6} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad P_1^{E_6} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \\
P_2^{E_6} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & 1 & 1 & -1 & -1 & -1 \\ -1 & 1 & 1 & 1 & -1 & -1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}, \quad P_3^{E_6} = \begin{pmatrix} 1 & -\frac{1}{2} & -\frac{1}{4} & -\frac{1}{2} & -\frac{1}{4} \\ 1 & 1 & -1 & -\frac{1}{2} & \frac{1}{2} \\ 2 & 2 & 1 & -1 & 1 \\ 1 & 1 & \frac{1}{2} & 1 & \frac{1}{2} \\ 2 & -4 & -2 & -1 & 1 \end{pmatrix}, \\
P_4^{E_6} &= \begin{pmatrix} 1 & 0 & 0 & -\frac{1}{2} & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 2 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 4 & 2 & 0 & -2 & 1 \end{pmatrix}, \quad P_5^{E_6} = \begin{pmatrix} 1 & -\frac{1}{4} & -\frac{1}{8} & -\frac{1}{4} \\ -2 & 1 & 0 & 0 \\ -4 & 2 & 1 & 0 \\ 0 & 1 & \frac{1}{2} & 1 \end{pmatrix}, \quad P_6^{E_6} = \begin{pmatrix} 1 & 0 & -\frac{1}{2} \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \\
P_7^{E_6} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 1 & 1 \end{pmatrix}, \quad P_8^{E_6} = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}, \quad P_m^{E_6} = (1) \quad (9 \leq m \leq 11).
\end{aligned}$$

Here, we think of indices for rows and columns of the matrix $P_m^{E_6}$ as Λ_m given in (5.43), and we arrange them as in increasing order. We set

$$\begin{aligned}
[\psi_{i,i}^{E_6}]_{i \in \Lambda_0} &= P_0^{E_6}[\alpha_i^*]_{i \in \Lambda_0}, \\
[\psi_{i,i+m}^{E_6}]_{i \in \Lambda_m} &= P_m^{E_6}[\alpha_{i,i+m} \psi_{i,i+m-1}^{E_6}]_{i \in \Lambda_m} \quad \text{for } 1 \leq m \leq 11.
\end{aligned}$$

Note that $\psi_{i,i}^{E_6} = \alpha_i^*$ for $i = 1, 4, \dots, 8$, namely

$$\begin{aligned}
\psi_{1,1}^{E_6} &= \psi_{1,1}^{E_7} - \frac{2}{3}\psi_{3,3}^{E_7} = \frac{2}{3}\partial_1 - \frac{2}{3}\partial_2 - \frac{2}{3}\partial_3, \\
\psi_{4,4}^{E_6} &= \psi_{4,4}^{E_7} - \frac{4}{3}\psi_{3,3}^{E_7} = \frac{1}{3}\partial_1 - \frac{1}{3}\partial_2 - \frac{1}{3}\partial_3 + \partial_4, \\
\psi_{5,5}^{E_6} &= \psi_{5,5}^{E_7} - \frac{5}{3}\psi_{3,3}^{E_7} = \frac{2}{3}\partial_1 - \frac{2}{3}\partial_2 - \frac{2}{3}\partial_3 + \partial_4 + \partial_5, \\
\psi_{6,6}^{E_6} &= \psi_{6,6}^{E_7} - 2\psi_{3,3}^{E_7} = \partial_1 - \partial_2 - \partial_3 + \partial_4 + \partial_5 + \partial_6, \\
\psi_{7,7}^{E_6} &= \psi_{7,7}^{E_7} - \psi_{3,3}^{E_7} = \frac{1}{2}(\partial_1 - \partial_2 - \partial_3 + \partial_4 + \partial_5 + \partial_6 + \partial_7 - \partial_8), \\
\psi_{8,8}^{E_6} &= \psi_{8,8}^{E_7} - \frac{4}{3}\psi_{3,3}^{E_7} = \frac{1}{2}\left(\frac{5}{3}\partial_1 - \frac{5}{3}\partial_2 - \frac{5}{3}\partial_3 + \partial_4 + \partial_5 + \partial_6 + \partial_7 + \partial_8\right).
\end{aligned}$$

We can check by using Maple that $\{\psi_{i,h_I(i)}^{E_6} \mid i = 1, 4, \dots, 8\} \subset D(\mathcal{A}_I)$ for any lower ideal $I \subset \Phi_{E_6}^+$. From this together with $\det P_m^{E_6} \neq 0$ ($1 \leq m \leq 11$), we obtain the following theorem by Proposition 3.4 (see also Remark 3.5).

Theorem 5.16. *The derivations $\{\psi_{i,i+m}^{E_6} \mid 0 \leq m \leq 11, i \in \Lambda_m\}$ form uniform bases for the ideal arrangements of type E_6 .*

6. UNIFORM BASES FOR THE IDEAL ARRANGEMENTS IN A ROOT SUBSYSTEM

In this section we prove that uniform bases for the ideal arrangements in a root subsystem of given a root system can be obtained from that of the given root system. From this point of view, one can see that uniform bases of types E_6 and E_7 can be obtained from that of type E_8 .

Recall that \mathfrak{t} is a real euclidean space of dimension n and $\Phi \subset \mathfrak{t}^*$ an irreducible root system. Under the isomorphism $\mathfrak{t} \cong \mathfrak{t}^*$ induced from the inner product $(\ , \)$ on \mathfrak{t} , the image of a root $\alpha \in \mathfrak{t}^*$ is denoted by $\hat{\alpha} \in \mathfrak{t}$. We also denote $\check{\beta} \in \mathfrak{t}^*$ by the image of $\beta \in \mathfrak{t}$, that is,

$$\mathfrak{t} \cong \mathfrak{t}^*; \hat{\alpha} \leftarrow \alpha, \beta \mapsto \check{\beta}.$$

Let $\alpha_1, \dots, \alpha_n$ be the simple roots and its dual basis is denoted by β_1, \dots, β_n in this section. Let S be a nonempty subset of $[n]$ and \mathfrak{t}' a subspace of \mathfrak{t} spanned by $\hat{\alpha}_i$ for $i \in S$. Note that \mathfrak{t}' is orthogonal to β_i for $i \notin S$:

$$\mathfrak{t}' = \{x \in \mathfrak{t} \mid (\beta_i, x) = 0 \text{ for all } i \notin S\}.$$

The inner product on \mathfrak{t} naturally induces that on \mathfrak{t}' . Then, the isomorphisms $\mathfrak{t} \cong \mathfrak{t}^*$ and $\mathfrak{t}' \cong (\mathfrak{t}')^*$ via the inner products on \mathfrak{t} and \mathfrak{t}' respectively make the following commutative diagram:

$$\begin{array}{ccc} \mathfrak{t} & \xrightarrow{\cong} & \mathfrak{t}^* \\ \uparrow & & \downarrow \\ \mathfrak{t}' & \xrightarrow{\cong} & (\mathfrak{t}')^* \end{array}$$

Note that $(\mathfrak{t}')^*$ is isomorphic to the quotient space $\mathfrak{t}^*/\text{span}\{\check{\beta}_i \mid i \notin S\}$. We denote by $\bar{\alpha}$ the image of $\alpha \in \mathfrak{t}^*$ under the surjective map $\mathfrak{t}^* \rightarrow (\mathfrak{t}')^*$. We define $\Phi' \subset (\mathfrak{t}')^*$ as the image of a set $\{\alpha \in \Phi \mid \alpha(\beta_i) = 0 \text{ for } i \notin S\}$ under the surjection $\mathfrak{t}^* \rightarrow (\mathfrak{t}')^*$. Then, Φ' is a root system on $(\mathfrak{t}')^*$ and we can take $\{\bar{\alpha}_i \mid i \in S\}$ as the simple roots of Φ' .

Let e_1, \dots, e_n (resp. e'_i ($i \in S$)) be the exponents of the Weyl group W (resp. W') where W and W' are the Weyl groups associated with (\mathfrak{t}, Φ) and (\mathfrak{t}', Φ') respectively. A decomposition $\Phi^+ = \coprod_{i=1}^n \Phi_i^+$ satisfying (3.1) and (3.2) induces the decomposition $\Phi'^+ = \coprod_{i \in S} \Phi'_i{}^+$ satisfying (3.1) and (3.2) where $\Phi'_i{}^+$ is defined as $\Phi'_i{}^+ = \{\bar{\alpha}_{i,i+1}, \dots, \bar{\alpha}_{i,i+e'_i}\}$ for $i \in S$. Let $\mathcal{R} = \text{Sym } \mathfrak{t}^*$ and $\mathcal{R}' = \text{Sym } (\mathfrak{t}')^* \cong \mathcal{R}/(\check{\beta}_i \mid i \notin S)$. Let $\{\psi_{i,j} \in \text{Der } \mathcal{R} \mid i \in [n] \text{ and } i \leq j \leq e_i\}$ be uniform bases for the ideal arrangements in (\mathfrak{t}, Φ) of the form in Theorem 4.1. We denote the invertible matrices associated with the uniform bases by P_m for $0 \leq m \leq \text{ht}(\Phi^+)$. Namely, we can write

$$\begin{aligned} \psi_{i,i} &= p_i \beta_i \quad \text{for } i \in [n], \\ \psi_{i,i+m} &= \sum_{j \in \Lambda_m} p_{ij} \alpha_{j,j+m} \psi_{j,j+m-1} \quad \text{for } 1 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m, \end{aligned}$$

where $P_0 = \text{diag}(p_1, \dots, p_n)$ and $P_m = (p_{ij})_{i,j \in \Lambda_m} \in \text{GL}(\Lambda_m; \mathbb{Q})$. Then, we define $\{\psi'_{i,j} \in \text{Der } \mathcal{R}' = \mathcal{R}' \otimes \mathfrak{t}' \mid i \in [n] \text{ and } i \leq j \leq e_i\}$ as follows:

$$\begin{aligned} \psi'_{i,i} &= p_i \gamma_i \quad \text{for } i \in [n], \\ \psi'_{i,i+m} &= \sum_{j \in \Lambda_m} p_{ij} \overline{\alpha_{j,j+m}} \psi'_{j,j+m-1} \quad \text{for } 1 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m, \end{aligned}$$

where $\{\gamma_i \in \mathfrak{t}' \mid i \in S\}$ is the dual basis of the simple roots $\{\overline{\alpha_i} \in (\mathfrak{t}')^* \mid i \in S\}$ for Φ' with the convention $\gamma_j = 0$ whenever $j \notin S$.

Lemma 6.1. *Let i and j be positive integers with $1 \leq i \leq n$ and $i \leq j \leq e_i$. Then,*

$$\psi'_{i,j}(\overline{\alpha}) = \overline{\psi_{i,j}(\alpha)}$$

for any $\overline{\alpha} \in \Phi'$.

Proof. Since we can take $\{\overline{\alpha_k} \mid k \in S\}$ as the simple roots of Φ' , it is enough to check that $\psi'_{i,j}(\overline{\alpha_k}) = \overline{\psi_{i,j}(\alpha_k)}$ for any $k \in S$. We prove this by induction on $m := j - i$. As the base case, when $m = 0$, we have

$$\psi'_{i,i}(\overline{\alpha_k}) = p_i \gamma_i(\overline{\alpha_k}) = p_i \delta_{ik} = \overline{\psi_{i,i}(\alpha_k)}.$$

Now we assume that $m > 0$ and the claim holds for $m - 1$. Then, for any $i \in [n]$, we have

$$\begin{aligned} \psi'_{i,i+m}(\overline{\alpha_k}) &= \sum_{j \in \Lambda_m} p_{ij} \overline{\alpha_{j,j+m}} \psi'_{j,j+m-1}(\overline{\alpha_k}) \\ &= \sum_{j \in \Lambda_m} p_{ij} \overline{\alpha_{j,j+m}} \overline{\psi_{j,j+m-1}(\alpha_k)} \quad (\text{by the inductive assumption}) \\ &= \overline{\psi_{i,i+m}(\alpha_k)}. \end{aligned}$$

This completes the proof. \square

Proposition 6.2. *A set of derivations $\{\psi'_{i,j} \in \text{Der } \mathcal{R}' \mid i \in S \text{ and } i \leq j \leq e'_i\}$ forms uniform bases for the ideal arrangements in (\mathfrak{t}', Φ') .*

Proof. Let I' be a lower ideal in Φ^+ and $h_{I'} : S \rightarrow \mathbb{Z}_{\geq 0}$ the associated Hessenberg function. We first show that $\psi'_{i,h_{I'}(i)} \in D(\mathcal{A}_{I'})$ for any $i \in S$. Namely, we prove that $\psi'_{i,h_{I'}(i)}(\overline{\alpha}) \in \mathcal{R}'\overline{\alpha}$ for any $\overline{\alpha} \in I'$. Since we have $\psi'_{i,h_{I'}(i)}(\overline{\alpha}) = \overline{\psi_{i,h_{I'}(i)}(\alpha)}$ by Lemma 6.1, it is enough to prove that $\psi_{i,h_{I'}(i)}(\alpha) \in \mathcal{R}\alpha$. Let I be a lower ideal in Φ^+ defined as

$$(6.1) \quad I = \{\alpha_{i,j} \mid i \in S \text{ and } \overline{\alpha_{i,j}} \in I'\}.$$

Note that the Hessenberg function h_I associated with I is given by

$$h_I(i) = \begin{cases} h_{I'}(i) & \text{if } i \in S, \\ i & \text{if } i \notin S. \end{cases}$$

Since $\{\psi_{i,j} \in \text{Der } \mathcal{R} \mid i \in [n] \text{ and } i \leq j \leq e_i\}$ forms uniform bases for the ideal arrangements in (\mathfrak{t}, Φ) , we have $\psi_{i,h_{I'}(i)}(\alpha) = \psi_{i,h_I(i)}(\alpha) \in \mathcal{R}\alpha$. Hence, we obtain that $\psi'_{i,h_{I'}(i)} \in D(\mathcal{A}_{I'})$ for any $i \in S$.

From Theorem 2.1 it suffices to show that $\det(\psi'_{i,h_{I'}(i)}(\overline{\alpha_j}))_{i,j \in S}$ is equal to $\prod_{\overline{\alpha} \in I'} \overline{\alpha}$ up to a non-zero scalar multiplication. Considering the lower ideal I in (6.1), from Theorem 2.1 we have

$$\det(\psi_{i,h_I(i)}(\alpha_j))_{i,j \in [n]} \doteq \prod_{\alpha \in I} \alpha$$

because $\{\psi_{i,j} \in \text{Der } \mathcal{R} \mid i \in [n] \text{ and } i \leq j \leq e_i\}$ forms uniform bases for the ideal arrangements in (\mathfrak{t}, Φ) . This implies that

$$(6.2) \quad \det(\overline{\psi_{i,h_I(i)}(\alpha_j)})_{i,j \in [n]} \doteq \left(\prod_{\alpha \in I} \overline{\alpha} \right) = \left(\prod_{\overline{\alpha} \in I'} \overline{\alpha} \right) \quad \text{in } \mathcal{R}',$$

where the second equality follows from the definition (6.1). Noting that $\psi_{i,h_I(i)}(\alpha_j) = \psi_{i,i}(\alpha_j) \doteq \delta_{ij}$ for $i \notin S$, we have

$$\det(\psi_{i,h_I(i)}(\alpha_j))_{i,j \in [n]} \doteq \det(\psi_{i,h_I(i)}(\alpha_j))_{i,j \in S} \doteq \det(\psi_{i,h_{I'}(i)}(\alpha_j))_{i,j \in S}.$$

From this together with (6.2) we obtain

$$\det(\overline{\psi_{i,h_{I'}(i)}(\alpha_j)})_{i,j \in S} \doteq \left(\prod_{\overline{\alpha} \in I'} \overline{\alpha} \right).$$

But the left hand side coincides with $\det(\psi'_{i,h_{I'}(i)}(\overline{\alpha_j}))_{i,j \in S}$ by Lemma 6.1, so this completes the proof. \square

We can apply Proposition 6.2 to type E . Let $\psi_{i,j}^E = \psi_{i,j}^{E_8}$ for $1 \leq i \leq 8$ and $i \leq j \leq e_i^{E_8}$. For each m with $0 \leq m \leq 29 = \text{ht}(\Phi_{E_8}^+)$, we denote by $P_m^E = (p_{ij})_{i,j \in \Lambda_m}$ the matrix $P_m^{E_8}$ in Section 5.7.1 for simplicity. We define $\psi_{i,j}'^E$ and $\psi_{i,j}''^E$ for $1 \leq i \leq 8$ and $i \leq j \leq e_i^{E_8}$ as follows:

$$\begin{aligned} \psi_{i,i}'^E &= \begin{cases} \psi_{i,i}^{E_7} & \text{if } i = 1, 3, 4, 5, 6, 7, 8, \\ 0 & \text{if } i = 2, \end{cases} \\ \psi_{i,i+m}'^E &= \sum_{j \in \Lambda_m} p_{ij} \overline{\alpha_{j,j+m}} \psi_{j,j+m-1}'^E \quad \text{for } 1 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m, \\ \psi_{i,i}''^E &= \begin{cases} \psi_{i,i}^{E_6} & \text{if } i = 1, 4, 5, 6, 7, 8, \\ 0 & \text{if } i = 2, 3, \end{cases} \\ \psi_{i,i+m}''^E &= \sum_{j \in \Lambda_m} p_{ij} \overline{\alpha_{j,j+m}} \psi_{j,j+m-1}''^E \quad \text{for } 1 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m. \end{aligned}$$

Corollary 6.3. *A set of derivations $\{\psi_{i,j}'^E \in \text{Der } \mathcal{R}_{E_7} \mid i = 1, 3, 4, 5, 6, 7, 8 \text{ and } i \leq j \leq e_i^{E_7}\}$ forms uniform bases for the ideal arrangements in type E_7 . A set of derivations $\{\psi_{i,j}''^E \in \text{Der } \mathcal{R}_{E_6} \mid i = 1, 4, 5, 6, 7, 8 \text{ and } i \leq j \leq e_i^{E_6}\}$ forms uniform bases for the ideal arrangements in type E_6 .*

7. THE COHOMOLOGY RINGS OF REGULAR NILPOTENT HESSENBERG VARIETIES

The logarithmic derivation modules $D(\mathcal{A}_I)$ for the lower ideals I have a surprising connection with the cohomology rings of regular nilpotent Hessenberg varieties by the work of [4]. Originally, explicit presentations of the cohomology rings of regular nilpotent Hessenberg varieties in type A_{n-1} is given by [1] using the localization technique. Motivated by this work, uniform bases for the lower ideals of types A_{n-1}, B_n, C_n, G_2 are constructed in [4]. In this section we first introduce the Hessenberg varieties. We then explain the work of [4] and an explicit presentation of the cohomology rings of regular nilpotent Hessenberg varieties in all Lie types. Throughout this paper, all cohomology rings will be taken with real coefficients.

Let G be a semisimple linear algebraic group of rank n . We fix a Borel subgroup B of G and a maximal torus T included in B . The Lie algebras of G and B are denoted by \mathfrak{g} and \mathfrak{b} , respectively. A **Hessenberg space** H is a \mathfrak{b} -submodule of \mathfrak{g} containing \mathfrak{b} . One can see that there is one-to-one correspondence between the set of lower ideals and the set of Hessenberg spaces which sends $I \subset \Phi^+$ by

$$H(I) := \mathfrak{b} \oplus \left(\bigoplus_{\alpha \in I} \mathfrak{g}_{-\alpha} \right),$$

where \mathfrak{g}_α is the root space for a root α . The **Hessenberg variety** $\text{Hess}(X, I)$ associated with an element $X \in \mathfrak{g}$ and a lower ideal $I \subset \Phi^+$ is defined as the following subvariety of the flag variety G/B :

$$\text{Hess}(X, I) := \{gB \in G/B \mid \text{Ad}(g^{-1})(X) \in H(I)\}.$$

In what follows, we concentrate on Hessenberg varieties $\text{Hess}(N, I)$ for a regular nilpotent element N and we call them **regular nilpotent Hessenberg varieties**. Here, we recall that an element $X \in \mathfrak{g}$ is nilpotent if $\text{ad}(X)$ is nilpotent, i.e., $\text{ad}(X)^k = 0$ for some $k > 0$. An element $X \in \mathfrak{g}$ is regular if its G -orbit of the adjoint action has the largest possible dimension.

Let \hat{T} be the character group of T . Since any $\alpha \in \hat{T}$ extends to a character of B , α defines a complex line bundle $L_\alpha := G \times_B \mathbb{C}_\alpha$ where \mathbb{C}_α is the one-dimensional B -module via α . We may regard \hat{T} as an additive group so that $\hat{T} \otimes_{\mathbb{Z}} \mathbb{R}$ is identified with the dual space \mathfrak{t}^* of the Lie algebra of the maximal compact torus. To each $\alpha \in \mathfrak{t}^*$ we can assign the Euler class $e(L_{-\alpha}) \in H^2(G/B)$. This assignment induces a ring homomorphism

$$(7.1) \quad \varphi: \mathcal{R} = \text{Sym}(\mathfrak{t}^*) \rightarrow H^*(G/B)$$

which doubles the grading on \mathcal{R} . According to Borel's theorem [5], the map φ is surjective and its kernel is the ideal generated by W -invariants in \mathcal{R} with zero constant term. Here, W is the Weyl group. Composing φ with the restriction map $H^*(G/B) \rightarrow H^*(\text{Hess}(N, I))$, we have a ring homomorphism

$$(7.2) \quad \varphi_I: \mathcal{R} \rightarrow H^*(\text{Hess}(N, I)).$$

The map φ_I is surjective from the result of [4] (Theorem 7.1 below). Moreover, its kernel can be described in terms of the logarithmic derivation module $D(\mathcal{A}_I)$. We can identify \mathfrak{t} and \mathfrak{t}^* via the Killing form which implies the isomorphism $\mathcal{R} \otimes \mathfrak{t} \cong \mathcal{R} \otimes \mathfrak{t}^*$. Composing the isomorphism with the multiplication map $\mathcal{R} \otimes \mathfrak{t}^* \rightarrow \mathcal{R}$, we obtain an \mathcal{R} -module map

$$q: \text{Der } \mathcal{R} = \mathcal{R} \otimes \mathfrak{t} \cong \mathcal{R} \otimes \mathfrak{t}^* \rightarrow \mathcal{R}.$$

We define an ideal $\mathfrak{a}(I)$ as the image of the logarithmic derivation module $D(\mathcal{A}_I)$ under the map q :

$$\mathfrak{a}(I) := q(D(\mathcal{A}_I)).$$

Theorem 7.1 ([4]). *The map φ_I in (7.2) is surjective and its kernel coincides with the ideal $\mathfrak{a}(I)$. In particular, φ_I induces the ring isomorphism*

$$H^*(\text{Hess}(N, I)) \cong \mathcal{R}/\mathfrak{a}(I).$$

From Theorem 7.1 together with explicit uniform bases for the lower ideals, we obtain an explicit presentation of the cohomology rings of regular nilpotent Hessenberg varieties in all Lie types. In fact, motivated by the explicit presentation given in [1, Theorem A] for type A_{n-1} , in [4] the explicit presentation for types A_{n-1}, B_n, C_n, G_2 are obtained from this perspective ([4, Corollary 10.4, Corollary 10.10, Corollary 10.15, Corollary 10.18]). We now state that an explicit presentation of the cohomology rings of regular nilpotent Hessenberg varieties in all Lie types which includes that of [1, 4] for types A_{n-1}, B_n, C_n, G_2 .

Let $\varpi_1, \dots, \varpi_n$ be the fundamental weights. In what follows, L means either of Lie types $A_n, B_n, C_n, D_n, E_6, E_7, E_8, F_4, G_2$. Let $\alpha_{i,j}^L$ be the positive root for type L defined in Section 5. Using the invertible matrices P_m^L for type L defined in Section 5, we define inductively polynomials $f_{i,i+m}^L \in \mathcal{R}$ for $0 \leq m \leq \text{ht}(\Phi^+)$ and $i \in \Lambda_m$ as follows:

$$\begin{aligned} [f_{i,i}^L]_{i \in \Lambda_0} &= \tilde{P}_0^L [\varpi_i]_{i \in \Lambda_0}, \\ [f_{i,i+m}^L]_{i \in \Lambda_m} &= P_m^L [\alpha_{i,i+m}^L \psi_{i,i+m-1}^L]_{i \in \Lambda_m} \quad \text{for } 1 \leq m \leq \text{ht}(\Phi^+), \end{aligned}$$

where \tilde{P}_0^L is the diagonal matrix with entries $\frac{2p_1}{\|\alpha_1\|^2}, \dots, \frac{2p_n}{\|\alpha_n\|^2}$ and p_1, \dots, p_n are the entries of the diagonal matrix P_0^L . Noting that $q(\alpha_i^*) = \frac{2}{\|\alpha_i\|^2} \varpi_i$, we have

$$(7.3) \quad f_{i,i+m}^L = q(\psi_{i,i+m}^L) \quad \text{for } 0 \leq m \leq \text{ht}(\Phi^+) \text{ and } i \in \Lambda_m.$$

Here, for only type $L = D_n$ we think of the equality (7.3) as $f_{i,i+m}^{D_n} = q(\tilde{\psi}_{i,i+m}^{D_n})$.

Recall that the set of lower ideals $I \subset \Phi_L^+$ and the set of Hessenberg functions h for type L defined in Section 5 are in one-to-one correspondence which sends I to h_I in (3.3). We denote a regular nilpotent Hessenberg variety $\text{Hess}(N, I)$ by $\text{Hess}(N, h)$ where h is corresponding to the lower ideal I , i.e. $h = h_I$. Since $\{\psi_{i,i+m}^L \mid 0 \leq m \leq \text{ht}(\Phi^+), i \in \Lambda_m\}$ forms uniform bases for the lower ideals of type L , we obtain the explicit presentation of the cohomology rings of regular nilpotent Hessenberg varieties in all Lie types from Theorem 7.1 and (7.3).

Corollary 7.2. *Let h be a Hessenberg function for type L and $\text{Hess}(N, h)$ the associated regular nilpotent Hessenberg variety for type L . Then, the following ring isomorphism holds*

$$H^*(\text{Hess}(N, h)) \cong \mathcal{R}/(f_{1,h(1)}^L, \dots, f_{n,h(n)}^L).$$

Remark 7.3. The regular nilpotent Hessenberg variety $\text{Hess}(N, h)$ for the special case when $h(i) = i + 1$ for $i \in [n]$ is called the **Peterson variety**, denoted by Pet . An explicit presentation of $H^*(Pet)$ is given by [8] as follows:

$$H^*(Pet) \cong \mathcal{R}/(\alpha_i \varpi_i \mid 1 \leq i \leq n).$$

One can see that Corollary 7.2 generalizes the explicit presentation above.

The polynomials $f_{i,j}^L$ have the explicit formula for types $L = A_{n-1}, B_n, C_n, G_2$ by [1, 4]. For type D_n we also have an explicit formula for $g_{i,j}^{D_n} := q(\psi_{i,j}^{D_n})$ because q sends ∂_i to x_i . That is, for $1 \leq i \leq n-1$ we have

$$\begin{aligned} g_{i,j}^{D_n} &= \sum_{k=1}^i (x_k - x_{i+1}) \cdots (x_k - x_j) x_k \quad \text{for } i \leq j \leq n-2 \ (i \neq n-1), \\ g_{i,n-1}^{D_n} &= \sum_{k=1}^i ((x_k - x_{i+1}) \cdots (x_k - x_{n-1})(x_k + x_n)) + (-1)^{n-i} n x_{i+1} \cdots x_n, \\ g_{i,n+j}^{D_n} &= \sum_{k=1}^i ((x_k - x_{i+1}) \cdots (x_k - x_n)(x_k + x_n) \cdots (x_k + x_{n-j})) \\ &\quad + (-1)^{n-i+1} n x_{i+1} \cdots x_{n-1-j} x_{n-j}^2 \cdots x_n^2 \quad \text{for } 0 \leq j \leq n-1-i. \end{aligned}$$

We also have

$$g_{n,2n-1-r}^{D_n} = \sum_{k=1}^r ((-1)^{n-r+1} (x_k - x_{r+1}) \cdots (x_k - x_{n-1})(x_k - x_n)) + n x_{r+1} \cdots x_n$$

for $0 \leq r \leq n-1$.

From Theorems 5.6 and 7.1 we obtain the following corollary.

Corollary 7.4. *Let h be a Hessenberg function for type D_n and $\text{Hess}(N, h)$ the associated regular nilpotent Hessenberg variety for type D_n . Then, the following ring isomorphism holds*

$$H^*(\text{Hess}(N, h)) \cong \mathbb{R}[x_1, \dots, x_n] / (g_{1,h(1)}^{D_n}, \dots, g_{n,h(n)}^{D_n}).$$

Also, we can describe systematically explicit presentations of the cohomology rings of the regular nilpotent Hessenberg varieties for type E . Using notations in Section 6, we define $g_{i,j}^E = q(\psi_{i,j}^E)$, $g'_{i,j}^E = q(\psi'_{i,j}^E)$, $g''_{i,j}^E = q(\psi''_{i,j}^E)$. Note that $g_{i,j}^E = f_{i,j}^{E_8}$. One can see inductively that

$$\begin{aligned} g'_{i,j}^E &\equiv g_{i,j}^E \pmod{g_{2,2}^E}, \\ g''_{i,j}^E &\equiv g_{i,j}^E \pmod{(g_{2,2}^E, g_{3,3}^E)}. \end{aligned}$$

This together with Corollary 6.3 implies the following corollary. Recall that Hessenberg functions h^{E_7} and h^{E_6} for types E_7 and E_6 respectively satisfy $h^{E_7}(2) = 2$ and $h^{E_6}(2) = 2, h^{E_6}(3) = 3$.

Corollary 7.5. *Let h be a Hessenberg function for type E_n ($n = 6, 7, 8$) and $\text{Hess}(N, h)$ the associated regular nilpotent Hessenberg variety for type E . Then, the following ring isomorphism holds*

$$H^*(\text{Hess}(N, h)) \cong \mathbb{R}[x_1, \dots, x_8] / (g_{1,h(1)}^E, \dots, g_{8,h(8)}^E).$$

REFERENCES

- [1] H. Abe, M. Harada, T. Horiguchi, and M. Masuda, *The cohomology rings of regular nilpotent Hessenberg varieties in Lie type A*, Int. Math. Res. Not. IMRN., **2019** (2019), 5316–5388.
- [2] H. Abe and T. Horiguchi, *A survey of recent developments on Hessenberg varieties*, arXiv:1904.11155.
- [3] T. Abe, M. Barakat, M. Cuntz, T. Hoge, and H. Terao, *The freeness of ideal subarrangements of Weyl arrangements*, J. European Math. Soc. **18** (2016), 1339–1348.
- [4] T. Abe, T. Horiguchi, M. Masuda, S. Murai, and T. Sato, *Hessenberg varieties and hyperplane arrangements*, to appear in J. Reine Angew. Math., DOI: 10.1515/crelle-2018-0039.
- [5] A. Borel, *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Ann. of Math. (2) **57** (1953), 115–207.
- [6] F. De Mari, C. Procesi, and M. A. Shayman, *Hessenberg varieties*, Trans. Amer. Math. Soc. **332** (1992), no. 2, 529–534.
- [7] F. De Mari and M. A. Shayman, *Generalized Eulerian numbers and the topology of the Hessenberg variety of a matrix*, Acta Appl. Math. **12** (1988), no. 3, 213–235.
- [8] M. Harada, T. Horiguchi, and M. Masuda, *The equivariant cohomology rings of Peterson varieties in all Lie types*, Canad. Math. Bull. **58** (2015), no. 1, 80–90.
- [9] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, 29. Cambridge University Press, Cambridge, 1990.
- [10] P. Orlik and H. Terao, *Arrangements of hyperplanes*, Grundlehren der Mathematischen Wissenschaften, **300**, Springer-Verlag, Berlin, 1992.
- [11] K. Saito, *Theory of logarithmic differential forms and logarithmic vector fields*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **27** (1980), 265–291.
- [12] H. Terao, *Arrangements of hyperplanes and their freeness I, II.*, J. Fac. Sci. Univ. Tokyo, **27**(2): 293–320, 1980.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, TOKYO UNIVERSITY OF SCIENCE, 2641 YAMAZAKI, NODA, CHIBA 278-8510, JAPAN
E-mail address: enokizono_makoto@ma.noda.tus.ac.jp

DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, 1-5, YAMADAOKA, SUITA, OSAKA, 565-0871, JAPAN
E-mail address: tatsuya.horiguchi0103@gmail.com

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KYOTO UNIVERSITY, KYOTO, 606-8522, JAPAN
E-mail address: tnagaoka@math.kyoto-u.ac.jp

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, KOMABA, MEGURO-KU, TOKYO 153-8914, JAPAN
E-mail address: akiyoshi@ms.u-tokyo.ac.jp