

TM₁ neutrino mixing with $\sin \theta_{13} = \frac{1}{\sqrt{3}} \sin \frac{\pi}{12}$

R. Krishnan^{a,1}

¹Saha Institute of Nuclear Physics, 1/AF Bidhannagar, Kolkata 700064, India,
<https://orcid.org/0000-0002-0707-3267>

Published: 21 April 2022

Abstract We construct a neutrino model using the flavour group $S_4 \times C_4 \times C_3 \times C_2$ under the type-1 seesaw mechanism. The vacuum alignments of the flavons in the model lead to TM₁ mixing with $\sin \theta_{13} = \frac{1}{\sqrt{3}} \sin \frac{\pi}{12}$. The mixing also exhibits μ - τ reflection symmetry. By fitting the eigenvalues of the effective seesaw mass matrix with the observed neutrino mass-squared differences, we predict the individual light neutrino masses. The vacuum alignment of the S_4 triplet appearing in the Majorana mass term plays a key role in obtaining the aforementioned TM₁ scenario. Since the symmetries of the flavour group are not sufficient to define this alignment, we apply the recently proposed framework of the auxiliary group in our model. Using this framework, the S_4 triplet is obtained by coupling together several irreducible multiplets that transform under an expanded flavour group consisting of the original flavour group as well as an auxiliary group. The vacuum alignment of each of these multiplets is uniquely defined in terms of its residual symmetries under the expanded flavour group. As a result, the S_4 triplet constructed from these multiplets also becomes uniquely defined.

Keywords Neutrino phenomenology · Discrete flavour symmetries

1 Introduction

The tribimaximal mixing (TBM) [1],

$$U_{\text{TBM}} = \begin{pmatrix} \frac{\sqrt{2}}{\sqrt{3}} & \frac{1}{\sqrt{3}} & 0 \\ \frac{-1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ \frac{-1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{-1}{\sqrt{2}} \end{pmatrix}, \quad (1)$$

has been the most widely studied mixing ansatz in literature. Though the discovery of nonzero reactor angle has ruled out this ansatz, model builders often use it as a starting point. One approach is to use TBM as the lowest order approximation and then obtain realistic mixing patterns through higher-order corrections. Another approach is to directly obtain realistic ansatzes that are modifications of TBM. Such mixing scenarios preserve certain symmetries of TBM while breaking certain others. A nomenclature for the mixing matrix that preserves the i^{th} column (row) of U_{TBM} was proposed as TM _{i} (TM ^{i}) [2, 3].

In this paper, we construct a model that results in TM₁ mixing [4]. TM₁ preserves the first column of U_{TBM} and mixes its second and third columns. Therefore, we have

$$U_{\text{TM}_1}(\theta, \zeta) = U_{\text{TBM}} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & \sin \theta e^{-i\zeta} \\ 0 & -\sin \theta e^{i\zeta} & \cos \theta \end{pmatrix} \\ = \begin{pmatrix} \frac{\sqrt{2}}{\sqrt{3}} & \frac{\cos \theta}{\sqrt{3}} & \frac{\sin \theta}{\sqrt{3}} e^{-i\zeta} \\ \frac{-1}{\sqrt{6}} \frac{\cos \theta}{\sqrt{3}} - \frac{\sin \theta}{\sqrt{2}} e^{i\zeta} \frac{\sin \theta}{\sqrt{3}} e^{-i\zeta} + \frac{\cos \theta}{\sqrt{2}} \\ \frac{-1}{\sqrt{6}} \frac{\cos \theta}{\sqrt{3}} + \frac{\sin \theta}{\sqrt{2}} e^{i\zeta} \frac{\sin \theta}{\sqrt{3}} e^{-i\zeta} - \frac{\cos \theta}{\sqrt{2}} \end{pmatrix}. \quad (2)$$

Several models have been proposed that generate TM₁ mixing. A model with two highly degenerate right-handed neutrinos resulting in resonant leptogenesis and TM₁ mixing is constructed in Ref. [4]. The discrete group S_4 is often used to implement TM₁ mixing [5, 6, 7, 8]. In Ref. [9], a modified Friedberg-Lee symmetry is utilised. A TM₁ scenario with one texture zero is discussed in Ref. [10]. TM₁ scenarios with μ - τ reflection symmetry and maximally broken CP are discussed in Refs. [6, 11]. Constrained sequential dominance (CSD) models often

^akrishnan.rama@saha.ac.in; krisphysics@gmail.com

predict TM_1 mixing along with the lightest neutrino mass being equal to zero. We have several such CSD models implemented using A_4 [12, 13, 14] and S_4 [15, 16] symmetries.

Using the parametrisation, Eq. (2), we obtain

$$\sin^2 \theta_{13} = \frac{\sin^2 \theta}{3}, \quad (3)$$

$$\sin^2 \theta_{12} = 1 - \frac{2}{3 - \sin^2 \theta}, \quad (4)$$

$$\sin^2 \theta_{23} = \frac{1}{2} \left(1 + \frac{\sqrt{6} \sin 2\theta \cos \zeta}{3 - \sin^2 \theta} \right), \quad (5)$$

$$J = \frac{\sin 2\theta \sin \zeta}{6\sqrt{6}}, \quad (6)$$

where

$$J = \text{Im}(U_{\mu 3} U_{e 3}^* U_{e 2} U_{\mu 2}^*) \\ = \frac{1}{8} \sin \delta \sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \cos \theta_{13} \quad (7)$$

is the Jarlkog's rephasing invariant [17, 18, 19, 20, 21].

Global fit [22] of neutrino oscillation data gives

$$\sin^2 \theta_{13} = 0.02241_{-0.00065}^{+0.00066}, \quad (8)$$

$$\sin^2 \theta_{12} = 0.310_{-0.012}^{+0.013}, \quad (9)$$

This leads to $|U_{12}|^2 = \sin^2 \theta_{12} \cos^2 \theta_{13} = 0.303_{-0.012}^{+0.012}$ and $|U_{12}|^2 + |U_{13}|^2 = \sin^2 \theta_{12} \cos^2 \theta_{13} + \sin^2 \theta_{13} = 0.325_{-0.012}^{+0.013}$. The TBM prediction of $|U_{12}|^2 = \frac{1}{3}$ is disfavoured while the TM_1 constraint $|U_{12}|^2 + |U_{13}|^2 = \frac{1}{3}$ is consistent with the experimental fit at 1σ level. Therefore, we may argue that the most promising among TM_i (TM^i) mixings is perhaps TM_1 .

Discrete groups have been used extensively to model the observed flavour symmetries [23, 24, 25, 26, 27, 28, 29, 30, 31, 32]. The three families of fermions are often assumed to transform as a triplet under a given discrete group. Flavour models may also include scalar fields called flavons which transform as various multiplets under the group. The Yukawa couplings and thus the fermion mass matrices in the Standard Model (SM) are obtained in terms of the Vacuum Expectation Values (VEVs) of these flavon fields. The VEVs emerge through Spontaneous Symmetry Breaking (SSB) of the flavon potentials. Therefore, the features of the mass matrices have their origin in the properties of the flavour group such as the types of its irreducible multiplets, the Clebsch-Gordan (C-G) coefficients appearing in the tensor products of these multiplets, the alignment of the flavon VEVs etc. These general principles underlie the model presented in our paper also.

The paper can be broadly divided into two parts. The first part consists of Sections 2-4 where a type-1 seesaw model is constructed using the discrete group

S_4 . In Section 2, we briefly mention the essential features of the S_4 group and its representations. In Section 3, we introduce the flavon content of the model which includes a triplet (χ) in the charged-lepton sector and a singlet (s) and a triplet (ϕ) in the Majorana neutrino sector. We construct the mass terms using these flavons, the SM fields and the right-handed neutrinos. We also construct the potential terms of the flavons and obtain their VEVs through the mechanism of SSB. The charged-lepton and the neutrino mass matrices are obtained in terms of these VEVs. Section 4 covers the phenomenology where we extract the PMNS mixing parameters as well as the light neutrino masses. We compare the model's predictions with the experimental data. The symmetries and the features of S_4 as well as the alignments of the flavon VEVs lead to the results predicted by the model. Notably, the result $\sin \theta_{13} = \frac{1}{\sqrt{3}} \sin \frac{\pi}{12}$ is obtained as the consequence of a specific VEV chosen for the triplet flavon, ϕ .

In the second part of the paper, consisting of Sections 5-7, we discuss the problem of obtaining the alignments of the VEVs based on discrete symmetries. In Section 5, we study all possible alignments of the triplets of S_4 that can be uniquely defined in terms of their residual symmetries alone. We find that the residual symmetries under S_4 are not sufficient to define the vacuum alignment of the triplet ϕ that we mentioned earlier. In this context, we briefly describe the recently proposed framework of the auxiliary group [33, 34] with which additional symmetries can be incorporated to obtain a richer choice of vacuum alignments. To apply this framework to our model, we construct a discrete group (which we call Y_{24}) with suitable symmetries in Section 6. We use this group in Section 7 to recast the model in the framework of the auxiliary group. Here, the triplet ϕ is replaced with an 'effective' triplet consisting of $\hat{\phi}$, $\tilde{\phi}$ and Δ which are 'elementary' flavons transforming as multiplets of the expanded flavour group in our framework. Their VEVs are fully defined in terms of their respective residual symmetries and thus the VEV of the effective triplet also becomes uniquely defined. We finally conclude in Section 8.


2 The Group S_4

The group S_4 has been used extensively in neutrino models [5, 7, 8, 15, 16, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45]. S_4 is the group of permutations of four objects. We may define it using the presentation,

$$\langle P, Q, R \mid P^2 = Q^3 = R^4 = PQR = I \rangle. \quad (10)$$

Therefore, S_4 is the von Dyck group with parameters (2,3,4). As can be inferred from its presentation, only

	()	(12)(34)	(12)	(1234)	(123)
1	1	1	1	1	1
1'	1	1	-1	-1	1
2	2	2	0	0	-1
3	3	-1	-1	1	0
3'	3	-1	1	-1	0

Table 1: The character table of S_4 .Fig. 1: The basis states and the symmetries of the representations **3** and **2** of S_4 .

two of its generators are independent, i.e.

$$P = QR, \quad Q = PR^3, \quad R = Q^2P. \quad (11)$$

The conjugacy classes and the irreducible representations of S_4 are listed in Table 1.

The defining triplet representation, **3**, can be generated using the matrices,

$$P = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad Q = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad R = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}. \quad (12)$$

This triplet representation denotes the rotational symmetries of a cube, Figure 1(a). In the figure, the unit vectors \hat{i} , \hat{j} and \hat{k} correspond to the basis states $(1, 0, 0)^T$, $(0, 1, 0)^T$ and $(0, 0, 1)^T$ respectively. P , Q and R are rotations (clockwise) by angles π , $\frac{2\pi}{3}$ and $\frac{\pi}{2}$ about the axes aligned along the directions $\hat{i} + \hat{k}$, $\hat{i} + \hat{j} + \hat{k}$ and \hat{i} respectively. S_4 has 24 elements which fall under five conjugacy classes, Table 1. The class (123) consists of the elements conjugate to Q . They represent rotations by angles $\frac{2\pi}{3}$ and $\frac{4\pi}{3}$ about the body diagonals of the cube. There are 8 elements in this conjugacy class. The elements in the class (12) are those that are conjugate to P . They represent π -rotations about the axes passing through the centres of the opposite edges of the

cube. We have 6 such rotations. The conjugacy class (1234) also has 6 elements. They are conjugate to R and they represent rotations by angles $\frac{\pi}{2}$ and $\frac{3\pi}{2}$ about the axes aligned along the basis states. On the other hand, π -rotations about these axes constitute the conjugacy class (12)(34). We have 3 such rotations and they are conjugate to R^2 .

In model building, a commonly adopted complex basis for **3** involves the generators S , T and U ,

$$S = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \bar{\omega} & 0 \\ 0 & 0 & \omega \end{pmatrix}, \quad (13)$$

$$U = - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix},$$

where $\omega = e^{i\frac{2\pi}{3}}$ and $\bar{\omega} = e^{-i\frac{2\pi}{3}}$ are the complex cube roots of unity. The two sets of bases, Eqs.(12, 13), are related by

$$S = U_\omega R^2 U_\omega^\dagger, \quad (14)$$

$$T = U_\omega Q U_\omega^\dagger, \quad (15)$$

$$U = U_\omega P R Q P U_\omega^\dagger = P R Q P, \quad (16)$$

where

$$U_\omega = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \bar{\omega} \\ 1 & \bar{\omega} & \omega \end{pmatrix} \quad (17)$$

is the 3×3 trimaximal matrix. Unlike P , Q and R , the generators S , T and U are independent. S_4 can not be generated from any two of them. In this paper, we adopt the basis, Eqs.(12), so as to provide a geometric interpretation to our analysis.

Besides the defining triplet **3**, S_4 has another triplet **3'**. **3'** differs from **3** by having a space inversion (multiplication by -1) in addition to the usual rotations for the elements of the conjugacy classes (12) and (1234). We have

$$P(\mathbf{3}') = -P, \quad Q(\mathbf{3}') = Q, \quad R(\mathbf{3}') = -R. \quad (18)$$

Like **3**, **3'** is also a faithful representation. The singlet **1'** involves multiplication with -1 corresponding to the space inversion in **3'**, i.e.

$$P(\mathbf{1}') = -1, \quad Q(\mathbf{1}') = 1, \quad R(\mathbf{1}') = -1. \quad (19)$$

The generators of the doublet, **2**, are

$$P(\mathbf{2}) = \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \quad Q(\mathbf{2}) = \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix}, \quad (20)$$

$$R(\mathbf{2}) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

	L	e_R	μ_R	τ_R	N	ρ	χ	s	ϕ	η
S_4	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{3}'$	$\mathbf{2}$
$C_4 \times C_3$	i	i	$i\omega$	$i\bar{\omega}$	i	1	ω	-1	-1	1
C_2	-1	1	1	-1	1	-1	1	1	1	1

Table 2: The fields in the model as the multiplets under $S_4 \times C_4 \times C_3 \times C_2$. The standard model Higgs (H) is assumed to remain invariant under the flavour group.

$\mathbf{2}$ forms the symmetry group of an equilateral triangle, Figure 1(b). $P(\mathbf{2})$ and $R(\mathbf{2})$ are the reflections about the directions $\frac{-1}{2}\hat{u} + \frac{\sqrt{3}}{2}\hat{v}$ and \hat{u} respectively where \hat{u} and \hat{v} are the basis states $(1, 0)^T$ and $(0, 1)^T$ respectively. $Q(\mathbf{2})$ is the rotation (clockwise) by an angle $\frac{2\pi}{3}$ in the \hat{u} - \hat{v} plane.

The tensor product expansion of two triplets ($\mathbf{3}'$'s) is given by

$$\mathbf{3}' \times \mathbf{3}' = \mathbf{1} + \mathbf{2} + \mathbf{3}' + \mathbf{3}. \quad (21)$$

With $a = (a_1, a_2, a_3)$ and $b = (b_1, b_2, b_3)$ transforming as $\mathbf{3}'$'s, we obtain

$$(ab)_1 = a_1b_1 + a_2b_2 + a_3b_3, \quad (22)$$

$$(ab)_2 = \left(2a_1b_1 - a_2b_2 - a_3b_3, \sqrt{3}(a_2b_2 - a_3b_3)\right)^T, \quad (23)$$

$$(ab)_{\mathbf{3}'} = (a_2b_3 + a_3b_2, a_3b_1 + a_1b_3, a_1b_2 + a_2b_1)^T, \quad (24)$$

$$(ab)_3 = (a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1)^T, \quad (25)$$

where $(ab)_1$, $(ab)_2$, $(ab)_{\mathbf{3}'}$ and $(ab)_3$ denote the irreducible parts in the tensor product of a and b that transform as $\mathbf{1}$, $\mathbf{2}$, $\mathbf{3}'$ and $\mathbf{3}$ respectively. The tensor product of two doublets, $p = (p_1, p_2)$ and $q = (q_1, q_2)$, leads to

$$\mathbf{2} \times \mathbf{2} = \mathbf{1} + \mathbf{1}' + \mathbf{2}, \quad (26)$$

with

$$(pq)_1 = p_1q_1 + p_2q_2, \quad (27)$$

$$(pq)_{\mathbf{1}'} = p_1q_2 - p_2q_1, \quad (28)$$

$$(pq)_2 = (-p_1q_1 + p_2q_2, p_1q_2 + p_2q_1)^T. \quad (29)$$

3 The model constructed using $S_4 \times C_4 \times C_3 \times C_2$

The field content of the model is given in Table 2. The three families of the left-handed weak-isospin lepton doublets and the three right-handed heavy neutrinos form the S_4 triplets, L and N , respectively. H is the

SM Higgs. The flavons, s , χ , ϕ , η and ρ are scalar fields and are gauge invariants. Using this field content and their symmetry properties, we obtain the following Lagrangian:

$$\begin{aligned} \mathcal{L} = & y_\tau \bar{L} \frac{\chi}{\Lambda} \tau_R H + y_\mu \bar{L} \frac{\rho \chi^*}{\Lambda^2} \mu_R H + y_e \bar{L} \frac{\rho(\chi^* \chi)_{\mathbf{3}'}}{\Lambda^3} e_R H \\ & + y_\nu \bar{L} N \frac{\rho}{\Lambda} \tilde{H} + y_s (\bar{N}^c N)_1 s + y_\phi (\bar{N}^c N)_{\mathbf{3}'} \phi, \end{aligned} \quad (30)$$

where Λ is the cut-off scale of the theory and y_x are the Yukawa-like coupling constants. Along with the symmetries given Table 2, we also impose CP symmetry. This implies that the coupling constants y_x are real. In Eq. (30), $(\cdot)_1$, $(\cdot)_3$ and $(\cdot)_{\mathbf{3}'}$ denote the tensor products that transform as $\mathbf{1}$, $\mathbf{3}$ and $\mathbf{3}'$ respectively under S_4 .

The flavon χ transforms as $\mathbf{3}'$ under S_4 and ω under C_3 . We have introduced the C_3 group so that χ , χ^* and $(\chi^* \chi)_{\mathbf{3}'}$ couple with τ_R , μ_R and e respectively along with the S_4 triplet L . They form the mass terms in the charged-lepton sector. Note that χ is the only flavon that is complex, and its complex conjugation corresponds to charge conjugation. The VEV of χ spontaneously breaks CP symmetry, and it is the only source of CP violation in the model. The flavon ρ and its C_2 charge aids in creating the observed charged-lepton mass hierarchy through the Froggatt-Nielsen mechanism.

The next term in the Lagrangian, $y_\nu \bar{L} N \frac{\rho}{\Lambda} \tilde{H}$, is the Dirac mass term for the neutrinos. The flavons, s and ϕ , transform as $\mathbf{1}$ and $\mathbf{3}'$ and couple with $(\bar{N}^c N)_1$ and $(\bar{N}^c N)_{\mathbf{3}'}$ respectively. They constitute the Majorana mass term for the neutrinos. We have introduced the C_4 group to prevent unwanted couplings. The flavour symmetry breaking scale is expected to be very high, typically the grand unification scale. Through the type-1 seesaw mechanism, the light-neutrino masses get suppressed by this scale.

The Higgs acquires the VEV,

$$\langle H \rangle = (0, v)^T, \quad (31)$$

through SSB. The VEV corresponds to a minimum point in the Higgs potential. The flavon fields also acquire VEVs through SSB,

$$\langle \rho \rangle = v_\rho, \quad (32)$$

$$\langle \chi \rangle = v_\chi (1, \omega, \bar{\omega})^T, \quad (33)$$

$$\langle s \rangle = v_s, \quad (34)$$

$$\langle \eta \rangle = v_\eta \left(-\frac{1}{2}, \frac{\sqrt{3}}{2} \right)^T, \quad (35)$$

$$\langle \phi \rangle = v_\phi \left(-\frac{1}{2\sqrt{2}}, -\sqrt{3}, \frac{1}{2\sqrt{2}} \right)^T. \quad (36)$$

In the following section, we construct the flavon potentials that lead to these VEVs. Note that the flavon η does not couple to any of the fermions in the Lagrangian at the lowest order and hence it does not contribute directly to the fermion mass matrices. We use it as a driving field [46] in the construction of the potential of the flavon ϕ .

3.1 The flavon potentials

The charged-lepton sector (χ)

For the singlet flavon ρ , we write the potential,

$$\mathcal{V}_\rho = k_\rho \rho^4 - 2k_\rho v_\rho^2 \rho^2, \quad (37)$$

where k_ρ is a real dimensionless parameter, and v_ρ is a real parameter of mass dimension one. For $k_\rho > 0$, this potential has two points of minima,

$$\rho = \pm v_\rho. \quad (38)$$

Through SSB, we obtain one of these minima as the VEV,

$$\langle \rho \rangle = v_\rho. \quad (39)$$

With $\chi = (\chi_1, \chi_2, \chi_3)$ and using Eqs. (22-24), we construct the following quadratic expressions:

$$(\chi^* \chi)_1 = \chi_1^* \chi_1 + \chi_2^* \chi_2 + \chi_3^* \chi_3, \quad (40)$$

$$(\chi \chi)_1 = \chi_1^2 + \chi_2^2 + \chi_3^2, \quad (41)$$

$$(\chi \chi)_2 = \left(2\chi_1^2 - \chi_2^2 - \chi_3^2, \sqrt{3}(\chi_2^2 - \chi_3^2) \right)^T, \quad (42)$$

$$(\chi \chi)_{3'} = 2(\chi_2 \chi_3, \chi_3 \chi_1, \chi_1 \chi_2)^T, \quad (43)$$

The antisymmetric expression $(\chi \chi)_3$ vanishes. At the quadratic order, the only invariant term is

$$\mathcal{T}(\chi^2) = (\chi^* \chi)_1. \quad (44)$$

At the cubic order, we have the invariant,

$$\mathcal{T}(\chi^3) = \text{Re}[(\chi \chi)_{3'}^T \chi] = 6 \text{Re}[\chi_1 \chi_2 \chi_3] \quad (45)$$

At the quartic order, we obtain the following invariants:

$$\mathcal{T}_1(\chi^4) = (\chi \chi)_1^* (\chi \chi)_1, \quad (46)$$

$$\mathcal{T}_2(\chi^4) = (\chi \chi)_2^\dagger (\chi \chi)_2, \quad (47)$$

$$\mathcal{T}_3(\chi^4) = (\chi \chi)_{3'}^\dagger (\chi \chi)_{3'}. \quad (48)$$

$\mathcal{T}(\chi^2)^2$ is related to the above invariants through the relation,

$$6(\mathcal{T}(\chi^2))^2 = 2\mathcal{T}_1(\chi^4) + \mathcal{T}_2(\chi^4) + 3\mathcal{T}_3(\chi^4). \quad (49)$$

Hence we have only three independent invariants at the quartic order. We use them to construct the flavon potential,

$$\begin{aligned} \mathcal{V}_\chi = & \kappa_0 v_\chi \mathcal{T}(\chi^3) + \kappa_1 \mathcal{T}_1(\chi^4) + \kappa_2 \mathcal{T}_2(\chi^4) + \kappa_3 \mathcal{T}_3(\chi^4) \\ & - (3\kappa_0 + 12\kappa_2 + 8\kappa_3) v_\chi^2 \mathcal{T}(\chi^2), \end{aligned} \quad (50)$$

where $\kappa_0, \kappa_1, \kappa_2, \kappa_3$ are dimensionless real parameters and v_χ is a real parameter of mass dimension one. By calculating the first derivative of this potential with respect to the components of χ , we can show that it has a set of extremum points,

$$\chi = g_i v_\chi (1, \omega, \bar{\omega})^T, \quad (51)$$

where g_i are the elements of the group $S_4 \times C_3$. Through SSB, the flavon acquires one among these extrema as its VEV¹,

$$\langle \chi \rangle = v_\chi (1, \omega, \bar{\omega})^T. \quad (52)$$

The neutrino sector (s, η, ϕ)

Construction of potential for the singlet s is similar to that of ρ , i.e.

$$\mathcal{V}_s = k_s s^4 - 2k_s v_s^2 s^2, \quad (53)$$

resulting in the VEV

$$\langle s \rangle = v_s. \quad (54)$$

We use Eqs. (27, 29) to obtain the tensor product of two flavon doublets, $\eta = (\eta_1, \eta_2)$, resulting in the quadratic invariant,

$$\mathcal{T}(\eta^2) = (\eta \eta)_1 = \eta_1^2 + \eta_2^2, \quad (55)$$

and the doublet,

$$(\eta \eta)_2 = (-\eta_1^2 + \eta_2^2, 2\eta_1 \eta_2)^T. \quad (56)$$

The antisymmetric expression $(\eta \eta)_{1'}$ vanishes. Using $(\eta \eta)_2$, we obtain the cubic invariant,

$$\mathcal{T}(\eta^3) = \eta^T (\eta \eta)_2 = -\eta_1^3 + 3\eta_1 \eta_2^2. \quad (57)$$

At the quartic order, we have the invariant

$$\mathcal{T}(\eta^4) = (\eta \eta)_2^T (\eta \eta)_2 = (\eta_1^2 + \eta_2^2)^2. \quad (58)$$

Note that $\mathcal{T}(\eta^4) = (\mathcal{T}(\eta^2))^2$.

¹For a large region of the parameter space $(\kappa_0, \kappa_1, \kappa_2, \kappa_3, v_\chi)$, the extrema correspond to the minima. For the stability of the VEV, we assume that the parameters fall in this region.

Using the flavon ϕ , we obtain the following quadratic expressions:

$$(\phi\phi)_1 = \phi_1^2 + \phi_2^2 + \phi_3^2, \quad (59)$$

$$(\phi\phi)_2 = \left(2\phi_1^2 - \phi_2^2 - \phi_3^2, \sqrt{3}(\phi_2^2 - \phi_3^2)\right)^T, \quad (60)$$

$$(\phi\phi)_{3'} = 2(\phi_2\phi_3, \phi_3\phi_1, \phi_1\phi_2)^T, \quad (61)$$

with

$$\mathcal{T}(\phi^2) = (\phi\phi)_1 \quad (62)$$

being the quadratic invariant. At the quartic order, we obtain the invariants,

$$\mathcal{T}_1(\phi^4) = (\phi\phi)_1^2, \quad (63)$$

$$\mathcal{T}_2(\phi^4) = (\phi\phi)_2^T (\phi\phi)_2, \quad (64)$$

$$\mathcal{T}_3(\phi^4) = (\phi\phi)_{3'}^T (\phi\phi)_{3'}. \quad (65)$$

These invariants are related by

$$4\mathcal{T}_1(\phi^4) = \mathcal{T}_2(\phi^4) + 3\mathcal{T}_3(\phi^4), \quad (66)$$

so only two of the quartic invariants are independent.

We can also couple η and ϕ to obtain the following invariants:

$$\mathcal{T}(\eta\phi^2) = \eta^T (\phi\phi)_2, \quad (67)$$

$$\mathcal{T}(\eta^2\phi^2) = (\eta\eta)_2^T (\phi\phi)_2. \quad (68)$$

As a result, we obtain a total of eight independent invariant terms involving η and ϕ : $\mathcal{T}(\eta^2)$, $\mathcal{T}(\eta^3)$, $\mathcal{T}(\eta^4)$, $\mathcal{T}(\phi^2)$, $\mathcal{T}_1(\phi^4)$, $\mathcal{T}_2(\phi^4)$, $\mathcal{T}(\eta\phi^2)$, $\mathcal{T}(\eta^2\phi^2)$.

Using these terms, we construct the potential,

$$\begin{aligned} \mathcal{V}_{\eta\phi} = & c_1\mathcal{T}(\eta^2) + c_2\mathcal{T}(\eta^3) + c_3\mathcal{T}(\eta^4) + c_4\mathcal{T}(\phi^2) \\ & + c_5\mathcal{T}_1(\phi^4) + c_6\mathcal{T}_2(\phi^4) + c_7\mathcal{T}(\eta\phi^2) + c_8\mathcal{T}(\eta^2\phi^2). \end{aligned} \quad (69)$$

These terms can be rearranged to obtain

$$\begin{aligned} \mathcal{V}_{\eta\phi} = & k_1(\mathcal{T}(\eta^2) - v_\eta^2)^2 + k_2(3\mathcal{T}(\phi^2) - (k_c v_\eta^2 + 4v_\phi^2))^2 \\ & + k_3(v_\eta\eta + (\eta\eta)_2)^T (v_\eta\eta + (\eta\eta)_2) \\ & + k_4(k_c v_\eta\eta - (\phi\phi)_2)^T (k_c v_\eta\eta - (\phi\phi)_2) \\ & + k_5(k_c(\eta\eta)_2 + (\phi\phi)_2)^T (k_c(\eta\eta)_2 + (\phi\phi)_2) \\ & - k_1 v_\eta^4 - k_2(k_c v_\eta^2 + 4v_\phi^2)^2, \end{aligned} \quad (70)$$

where v_η and v_ϕ are parameters of mass dimension one. The parameters k_1, \dots, k_5 and k_c are dimensionless;

k_1, \dots, k_5 are assumed to be positive. The eight parameters, $v_\eta, v_\phi, k_1, \dots, k_5, k_c$ are related to c_1, \dots, c_8 through the equations,

$$c_1 = (-2k_1 + k_3 + k_4 k_c^2) v_\eta^2, \quad c_5 = 9k_2, \quad (71)$$

$$c_2 = 2k_3 v_\eta, \quad c_6 = k_4 + k_5, \quad (72)$$

$$c_3 = k_1 + k_3 + k_5 k_c^2, \quad c_7 = -2k_4 k_c v_\eta, \quad (73)$$

$$c_4 = -6k_2(k_c v_\eta^2 + 4v_\phi^2), \quad c_8 = 2k_5 k_c. \quad (74)$$

The potential, Eq. (70), consists of five terms with the coefficients k_1 to k_5 . Each of them is positive semidefinite and hence they should all vanish when minimised. The constant factor, $-k_1 v_\eta^4 - k_2(k_c v_\eta^2 + 4v_\phi^2)^2$, does not play any role in the minimisation of the potential; it is added only for equating Eq. (69) with Eq. (70). The first term, $(\mathcal{T}(\eta^2) - v_\eta^2)^2$, vanishes when $|\eta|^2 = v_\eta^2$. Its minimum corresponds to a continuous set of points. The third term, $((\eta\eta)_2 + v_\eta\eta)^T ((\eta\eta)_2 + v_\eta\eta)$, breaks this continuous symmetry. This term vanishes for

$$\eta = g_i v_\eta \left(-\frac{1}{2}, \frac{\sqrt{3}}{2} \right)^T, \quad (75)$$

where g_i are the elements of the doublet representation of the group S_4 .

The second term, $(3\mathcal{T}(\phi^2) - (k_c v_\eta^2 + 4v_\phi^2))^2$, is invariant under the three-dimensional orthogonal transformation, $O(3)$, of the flavon ϕ , and this term vanishes when $3|\phi|^2 = k_c v_\eta^2 + 4v_\phi^2$. This minimum represents a continuous set of points corresponding to the $O(3)$ symmetry. The fourth and the fifth terms couple ϕ with η and they break this continuous symmetry to S_4 . These terms vanish when

$$\phi = g_i v_\phi \left(\frac{1}{\sqrt{2}} \sin \alpha, -2 \cos \alpha, -\frac{1}{\sqrt{2}} \sin \alpha \right)^T, \quad (76)$$

where

$$\sin \alpha = \frac{1}{3} \sqrt{8 - k_c \frac{v_\eta^2}{v_\phi^2}}, \quad \cos \alpha = \frac{1}{3} \sqrt{1 + k_c \frac{v_\eta^2}{v_\phi^2}} \quad (77)$$

and g_i are the elements of the triplet representation ($3'$) of S_4 . As a result, we obtain a discrete set of minima for the potential.

Through SSB, the flavons acquire one among the minima, Eqs. (75-76), as their VEVs,

$$\langle \eta \rangle = v_\eta \left(-\frac{1}{2}, \frac{\sqrt{3}}{2} \right)^T, \quad (78)$$

$$\langle \phi \rangle = v_\phi \left(\frac{1}{\sqrt{2}} \sin \alpha, -2 \cos \alpha, -\frac{1}{\sqrt{2}} \sin \alpha \right)^T. \quad (79)$$

The VEV assumed in our model, Eq. (36), corresponds to Eq. (79) with $\alpha = -\frac{\pi}{6}$. We may assign specific values to the parameters in the potential, v_ϕ , v_η , k_c , so that, using Eq. (77), we obtain $\alpha = -\frac{\pi}{6}$ and thus obtain the desired VEV. However, tuning the parameters in the potential to obtain a specific VEV is not justified if we expect that the underlying discrete symmetries naturally determine the alignment of the VEV. On the other hand, α being equal to $-\frac{\pi}{6}$ may point towards additional symmetries. This question is addressed in the second part of this paper where we use the framework of the auxiliary group to uniquely define the VEV in terms of these additional symmetries.

In the following section, we obtain the mass matrices in the charged-lepton and the neutrino sectors in terms of the VEVs, Eqs. (31-33, 36).

3.2 The charged-lepton mass matrix

The flavon χ couples in the charged-lepton mass term,

$$y_\tau \bar{L} \frac{\chi}{\Lambda} \tau_R H + y_\mu \bar{L} \frac{\rho \chi^*}{\Lambda^2} \mu_R H + y_e \bar{L} \frac{\rho(\chi^* \chi)_{\mathbf{3}'}}{\Lambda^3} e_R H. \quad (80)$$

Due to the C_3 assignments in Table 2, we can see that χ couples with τ_R , χ^* couples with μ_R and $(\chi^* \chi)_{\mathbf{3}'}$ couples with e_R at the lowest order. Using Eq. (25), we obtain

$$(\chi^* \chi)_{\mathbf{3}'} = (\chi_2^* \chi_3 + \chi_3^* \chi_2, \chi_3^* \chi_1 + \chi_1^* \chi_3, \chi_1^* \chi_2 + \chi_2^* \chi_1). \quad (81)$$

The VEV of χ , Eq. (33), and its conjugate,

$$\langle \chi \rangle = v_\chi(1, \omega, \bar{\omega}), \quad \langle \chi^* \rangle = v_\chi(1, \bar{\omega}, \omega), \quad (82)$$

couple with τ_R and μ_R respectively. Using Eqs. (81, 82), we obtain

$$\langle (\chi^* \chi)_{\mathbf{3}'} \rangle = -v_\chi^2(1, 1, 1), \quad (83)$$

which couples with e_R . Substituting the VEVs, Eqs. (82, 83) and the Higgs VEV, Eq. (31), in Eq. (80), we obtain the charged-lepton mass term after SSB,

$$\bar{l}_L M_l l_R, \quad (84)$$

where

$$l_L = (e_L, \mu_L, \tau_L)^T, \quad l_R = (e_R, \mu_R, \tau_R)^T, \quad (85)$$

and

$$M_l = v \frac{v_\chi}{\Lambda} \begin{pmatrix} -y_e \frac{v_\rho v_\chi}{\Lambda^2} & y_\mu \frac{v_\rho}{\Lambda} & y_\tau \\ -y_e \frac{v_\rho v_\chi}{\Lambda^2} & \bar{\omega} y_\mu \frac{v_\rho}{\Lambda} & \omega y_\tau \\ -y_e \frac{v_\rho v_\chi}{\Lambda^2} & \omega y_\mu \frac{v_\rho}{\Lambda} & \bar{\omega} y_\tau \end{pmatrix}. \quad (86)$$

is the charged-lepton mass matrix.

3.3 The neutrino mass matrices

The Dirac mass term for the neutrinos is $y_\nu \bar{L} N \frac{\rho}{\Lambda} \tilde{H}$. Substituting the VEVs of H and ρ in this term, we obtain

$$\bar{\nu}_L M_D N, \quad (87)$$

where

$$\nu_L = (\nu_e, \nu_\mu, \nu_\tau)^T, \quad N = (N_1, N_2, N_3)^T, \quad (88)$$

and

$$M_D = v \frac{v_\rho}{\Lambda} y_\nu I. \quad (89)$$

M_D is the Dirac mass matrix for the neutrinos, and is proportional to the identity, I , at the lowest order.

In the Majorana mass terms,

$$y_s (\bar{N}^c N)_1 s + y_\phi (\bar{N}^c N)_{\mathbf{3}'} \phi, \quad (90)$$

we substitute the VEVs, Eqs. (34, 36), to obtain

$$\bar{N}^c M_M N \quad (91)$$

where

$$M_M = v_s y_s I + v_\phi y_\phi \Theta \quad (92)$$

is the Majorana mass matrix with Θ being the off-diagonal matrix,

$$\Theta = \begin{pmatrix} 0 & \frac{1}{2\sqrt{2}} & -\sqrt{3} \\ \frac{1}{2\sqrt{2}} & 0 & -\frac{1}{2\sqrt{2}} \\ -\sqrt{3} & -\frac{1}{2\sqrt{2}} & 0 \end{pmatrix}. \quad (93)$$

The effective seesaw mass matrix is given by

$$M_{ss} = -M_D M_M^{-1} M_D^T. \quad (94)$$

Using Eqs. (89, 94), we obtain

$$M_{ss} = -v^2 \frac{v_\rho^2}{\Lambda^2} y_\nu^2 M_M^{-1}. \quad (95)$$

4 Phenomenology

We diagonalise the charged-lepton mass matrix, M_l , Eq. (86), using the 3×3 -trimaximal matrix, U_ω , Eq. (17),

$$U_\omega M_l \text{diag}(-1, 1, 1) = \text{diag}(m_e, m_\mu, m_\tau), \quad (96)$$

where $m_e = \sqrt{3} y_e v \frac{v_\rho v_\chi}{\Lambda^3}$, $m_\mu = \sqrt{3} y_\mu v \frac{v_\rho v_\chi}{\Lambda^3}$ and $m_\tau = \sqrt{3} y_\tau v \frac{v_\rho v_\chi}{\Lambda^3}$ are the charged-lepton masses. Let us assume that the expansion parameters $\frac{v_\chi}{\Lambda}$ and $\frac{v_\rho}{\Lambda}$ are approximately equal to 0.012. With $m_e \approx 0.5$ MeV, $m_\mu \approx 0.1$ GeV, $m_\tau \approx 1.8$ GeV and $v \approx 170$ GeV, we obtain

the Yukawa-like couplings to be of the order of one, $y_e \approx 1.0$, $y_\mu \approx 2.4$ and $y_\tau \approx 0.5$. We assume that the couplings in the neutrino sector y_ν , y_s and y_ϕ are also of the order of one.

To diagonalise the effective seesaw mass matrix, M_{ss} , we study the diagonalisation of Θ , Eq. (93). Applying the (13)-bimaximal matrix,

$$U_{\text{BM}} = \begin{pmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{-1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{pmatrix}, \quad (97)$$

on Θ , we obtain,

$$\begin{aligned} U_{\text{BM}}^T \Theta U_{\text{BM}} &= \begin{pmatrix} -\sqrt{3} & 0 & 0 \\ 0 & 0 & -\frac{1}{2} \\ 0 & -\frac{1}{2} & \sqrt{3} \end{pmatrix} \\ &= \frac{\sqrt{3}}{2} I + \begin{pmatrix} -\frac{3\sqrt{3}}{2} & 0 & 0 \\ 0 & -\cos \alpha & \sin \alpha \\ 0 & \sin \alpha & \cos \alpha \end{pmatrix}, \end{aligned} \quad (98)$$

where $\alpha = -\frac{\pi}{6}$. The above matrix is diagonalised as follows:

$$U_\theta^T U_{\text{BM}}^T \Theta U_{\text{BM}} U_\theta = \text{diag} \left(-\sqrt{3}, \frac{\sqrt{3}}{2} - 1, \frac{\sqrt{3}}{2} + 1 \right), \quad (99)$$

where

$$U_\theta = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & \sin \theta \\ 0 & -\sin \theta & \cos \theta \end{pmatrix} \quad \text{with} \quad \theta = -\frac{\pi}{12}. \quad (100)$$

With the help of the above result, we diagonalise the effective seesaw mass matrix, Eq. (95),

$$\begin{aligned} U_\theta^T U_{\text{BM}}^T M_{ss} U_{\text{BM}} U_\theta &= -v^2 \frac{v_\rho^2}{\Lambda^2} y_\nu^2 U_\theta^T U_{\text{BM}}^T M_M^{-1} U_{\text{BM}} U_\theta \\ &= \text{diag}(m_1, m_2, m_3), \end{aligned} \quad (101)$$

where m_1, m_2, m_3 are the light neutrino masses,

$$m_1 = \left(\sqrt{3} - r \right)^{-1} m, \quad (102)$$

$$m_2 = \left(-\frac{\sqrt{3}}{2} + 1 - r \right)^{-1} m, \quad (103)$$

$$m_3 = \left(-\frac{\sqrt{3}}{2} - 1 - r \right)^{-1} m, \quad (104)$$

with

$$r = \frac{v_s y_s}{v_\phi y_\phi}, \quad m = \frac{v^2 v_\rho^2 y_\nu^2}{v_\phi \Lambda^2 y_\phi}. \quad (105)$$

To obtain Eq. (101) from Eqs. (92, 95, 99), we have used the fact that a real symmetric matrix, as well as its inverse, are diagonalised by the same orthogonal matrix. The eigenvalues of the inverse matrix are simply the inverse of the eigenvalues of the original matrix.

The PMNS matrix is obtained by multiplying the diagonalising matrices of the charged-lepton mass matrix, Eq. (96), and the effective seesaw mass matrix, Eq. (101),

$$U = U_\omega U_{\text{BM}} U_\theta. \quad (106)$$

The tribimaximal matrix, Eq. (1), can be obtained in terms of U_ω and U_{BM} ,

$$U_\omega U_{\text{BM}} = \text{diag}(1, \omega, \bar{\omega}) U_{\text{TBM}} \text{diag}(1, 1, i). \quad (107)$$

Substituting Eq. (107) in Eq. (106), we obtain

$$U = \text{diag}(1, \omega, \bar{\omega}) U_{\text{TBM}} \text{diag}(1, 1, i) U_\theta \quad (108)$$

Comparing the above equation with Eq. (2), we get

$$U = \text{diag}(1, \omega, \bar{\omega}) U_{\text{TM}_1} \left(\frac{-\pi}{12}, \frac{\pi}{2} \right) \text{diag}(1, 1, i). \quad (109)$$

Note that a TM_1 mixing scheme with $\theta = \frac{\pi}{12}$ and $\zeta = \frac{\pi}{2}$ was proposed in Ref. [47]. The unobservable phases, $\text{diag}(1, \omega, \bar{\omega})$, can be ignored. The Majorana phases, $\text{diag}(1, 1, i)$, are potentially observable in the neutrinoless double-beta decay experiments.

Substituting the values of $\theta = \frac{-\pi}{12}$ and $\zeta = \frac{\pi}{2}$ in Eqs. (3, 4, 5, 6), we obtain

$$\sin^2 \theta_{13} = \frac{1}{3} \sin^2 \frac{-\pi}{12} = 0.0223, \quad (110)$$

$$\sin^2 \theta_{12} = 1 - \frac{2}{3 - \sin^2 \frac{-\pi}{12}} = 0.318, \quad (111)$$

$$\sin^2 \theta_{23} = \frac{1}{2}, \quad (112)$$

$$J = -\frac{1}{12\sqrt{6}}. \quad (113)$$

Comparing the values of J and the mixing angles with Eq. (7), we obtain

$$\delta_{\text{CP}} = -\frac{\pi}{2}. \quad (114)$$

The VEV of the flavon χ , Eq. (33), spontaneously breaks CP symmetry and produces the complex charged-lepton mass matrix, Eq. (86). Since the neutrino mass matrices are real, the charge-lepton sector becomes the sole source of CP violation in the model. U_ω , which diagonalises the charged-lepton mass matrix, results in the appearance of i in Eq. (108), which manifests as $\zeta = \frac{\pi}{2}$ in TM_1 mixing. This leads to $\delta_{\text{CP}} = -\frac{\pi}{2}$. From Eq. (2), it is clear that $\zeta = \pm \frac{\pi}{2}$ leads to the μ and τ rows of the mixing matrix being conjugate to each other. This




Fig. 2: The allowed ranges of the parameters r and m at the 3σ level. The red point corresponds to the best fit.




Fig. 3: Σm_i and $m_{\beta\beta}$ as predicted by the model

property, termed as the μ - τ reflection symmetry [48, 49, 50, 51, 52], results in maximal atmospheric mixing, i.e. $\sin^2 \theta_{23} = \frac{1}{2}$ as predicted by our model. The origin of μ - τ reflection symmetry becomes apparent when we express the neutrino effective seesaw mass matrix M_{ss} , Eq. (95), in the basis where the charged-lepton mass matrix is diagonal, i.e. $U_\omega^* M_{ss} U_\omega^\dagger$. This matrix remains invariant under the C_2 symmetry generated by complex conjugation and simultaneous exchange of μ - τ rows and columns. This C_2 symmetry manifests as the μ - τ reflection symmetry of the mixing matrix.

Our predictions, $\sin^2 \theta_{13} = 0.0223$ and $\sin^2 \theta_{12} = 0.318$, are within the 1σ ranges of the global fit, Eqs. (8, 9). The prediction $\sin^2 \theta_{23} = \frac{1}{2}$ is expected in any mixing scenario with μ - τ reflection symmetry. There are indications that the atmospheric mixing may not be maximal, which gives rise to the problem of the octant degeneracy. However, at the 3σ level, the allowed range is $0.428 < \sin^2 \theta_{23} < 0.624$. Therefore, $\sin^2 \theta_{23} = \frac{1}{2}$ is not ruled out. CP violation in the lepton sector is still an open problem and we do not have an accurate measurement of δ_{CP} . At the 3σ level, we have a large range, $135^\circ < \delta_{CP} < 366^\circ$, and the maximal value of $\delta_{CP} = -\frac{\pi}{2}$ lies within this range.

The neutrino masses, Eqs. (102, 103, 104) are given in terms of two free parameters, r and m . Using the

known mass-squared differences,

$$\Delta m_{21}^2 = 67.9 \rightarrow 80.1 \text{ meV}^2, \quad (115)$$

$$\Delta m_{31}^2 = 2432 \rightarrow 2618 \text{ meV}^2, \quad (116)$$

we can fit these parameters and predict the individual light neutrino masses. The allowed parameter space is shown in Figure 2. The resulting values of the individual neutrino masses are

$$m_1 = 5.1 \rightarrow 5.6 \text{ meV}, \quad (117)$$

$$m_2 = 8.5 \rightarrow 9.3 \text{ meV}, \quad (118)$$

$$m_3 = 49.4 \rightarrow 51.3 \text{ meV}, \quad (119)$$

From Figure 2, Eqs. (105) and given $\frac{v_\rho}{\Lambda} \approx 0.01$, we can estimate that the flavon VEVs v_ϕ and v_s are of the order of 10^{11} GeV.

The sum of the neutrino masses is constrained by cosmological observations. The upper bounds provided by these observations are a few hundreds of meV [53, 54], the most stringent bound being $\Sigma m_i < 78 \text{ meV}$ [53]. From Eqs. (117)-(119), we predict

$$\Sigma m_i = 63.1 \rightarrow 66.1 \text{ meV}. \quad (120)$$

This range is below the experimental bounds.

The neutrinoless double-beta decay experiments seek to determine whether the neutrinos are Majorana particles. If the neutrinoless double-beta decay is observed, the measurement of half-life leads to the determination of the effective mass,

$$m_{\beta\beta} = U_{e1}^2 m_1 + U_{e2}^2 m_2 + U_{e3}^2 m_3, \quad (121)$$

where U_{e1} , U_{e2} and U_{e3} are the elements of the first row of the PMNS matrix. Substituting the values of these elements from Eq. (109) in Eq. (121), we obtain

$$m_{\beta\beta} = \frac{2}{3} m_1 + \frac{1}{12} (2 + \sqrt{3}) m_2 + \frac{1}{12} (2 - \sqrt{3}) m_3. \quad (122)$$

Given our predicted masses, Eqs. (117, 118, 119), we obtain

$$m_{\beta\beta} = 7.2 \rightarrow 7.7 \text{ meV}. \quad (123)$$

This range is well below the upper bounds set by the recent $0\nu\beta\beta$ experiments [55, 56]. Figure 3 shows the ranges of Σm_i and $m_{\beta\beta}$ predicted by our model along with the best-fit point.




Fig. 4: Orbits under $\mathbf{3}$ of S_4

5 Uniquely Defining the VEVs

5.1 Orbits and stabilisers

In this section, we describe the concepts of orbits and stabilisers by using the action of the triplet representations of S_4 as examples. We utilise these concepts later in the paper. Various representations of a group act on the corresponding vector spaces. The triplet representations of S_4 act on the three-dimensional real space. Consider the set of all unit vectors forming a sphere in the 3-D real space. The action of an element of $\mathbf{3}$ of S_4 on any unit vector rotates it from one point on the sphere to another. Since $\mathbf{3}$ has 24 distinct elements, the group action on a given point leads to a set of 24 points on the sphere in the most general case, as shown in Figure 4(a). This set of points forms the orbit of the given point. The group action on any point within the orbit produces the same orbit; therefore, the orbit remains closed under the group action.

It need not be the case that all the orbits have their cardinality equal to the number of distinct elements in the representation. Group action on $x = \frac{1}{\sqrt{3}}(1, 1, 1)$ will produce the points,

$$\frac{1}{\sqrt{3}} \{(\pm 1, \pm 1, \pm 1), (\pm 1, \pm 1, \mp 1), (\pm 1, \mp 1, \pm 1), (\mp 1, \pm 1, \pm 1)\}, \quad (124)$$

which form the vertices of a cube, Figure 4(b). Rather than 24, here we have only 8 points in the orbit because each point on the cube remains invariant under the action of certain elements of the group. The set of such elements form a subgroup and it is termed as the stabiliser of the point. The stabiliser of $\frac{1}{\sqrt{3}}(1, 1, 1)$ is $\{Q, Q^2, I\}$ which forms a C_3 subgroup of S_4 . We have four such C_3 subgroups, corresponding to the four pairs of opposite vertices of the cube. The orbit-stabiliser theorem states that

$$|\text{Orb}(x)| = \frac{|G|}{|\text{Stab}(x)|}, \quad (125)$$

where $\text{Orb}(x)$ and $\text{Stab}(x)$ denote the orbit and the stabiliser respectively of a point x and $||$ denotes the cardinality. In the case of the cube, $|\text{Stab}(x)| = 3$; therefore, we have

$$|\text{Orb}(x)| = \frac{24}{3} = 8, \quad (126)$$

which is consistent with the number of vertices of the cube.

The group action on $x = (1, 0, 0)^T$ produces an octahedron,

$$\{(\pm 1, 0, 0), (0, \pm 1, 0), (0, 0, \pm 1)\}, \quad (127)$$

Figure 4(c). The stabiliser of $x = (1, 0, 0)^T$ is the C_4 subgroup, $\{R, R^2, R^3, I\}$. Using the orbit-stabiliser theorem, we obtain the cardinality of the orbit (the number of vertices of the octahedron) to be $\frac{24}{4} = 6$.

The group action on $x = \frac{1}{\sqrt{2}}(1, 0, 1)$ produces a cuboctahedron,

$$\frac{1}{\sqrt{2}} \{(\pm 1, 0, \pm 1), (\pm 1, \pm 1, 0), (0, \pm 1, \pm 1), (\pm 1, 0, \mp 1), (\pm 1, \mp 1, 0), (0, \pm 1, \mp 1)\}, \quad (128)$$

Figure 4(d). In this case, the stabiliser of x is the C_2 subgroup, $\{P, I\}$ and we obtain the number of vertices of the cuboctahedron to be $\frac{24}{2} = 12$.

The cube, Eq. (124), the octahedron, Eq. (127), and the cuboctahedron, Eq. (128), are the only ‘uniquely defined’ orbits that can be constructed in the case of the representation $\mathbf{3}$. They are unique in the sense that the orientations of their elements are fully defined by their respective residual symmetries (subgroups of S_4). For any other orbit of $\mathbf{3}$, the stabiliser of a point in the orbit will be the trivial group and the orbit will have the cardinality of 24. Such orbits do not form uniquely defined geometric objects like the cube.

Now let us analyse the representation $\mathbf{3}'$. We have $P(\mathbf{3}') = -P, Q(\mathbf{3}') = Q, R(\mathbf{3}') = -R$, Eqs. (18). This representation is the symmetry group of a tetrahedron




Fig. 5: Orbits under $\mathbf{3}'$ of S_4

which includes 12 proper rotations and 12 improper rotations. The action of $\mathbf{3}'$ on the point $\frac{1}{\sqrt{3}}(1, 1, 1)$ produces the orbit,

$$\frac{1}{\sqrt{3}} \{(1, 1, 1), (1, -1, -1), (-1, 1, -1), (-1, -1, 1)\}, \quad (129)$$

which forms the vertices of the tetrahedron, Figure 5(a). The point $\frac{1}{\sqrt{3}}(1, 1, 1)$ has the stabiliser $\{Q, Q^2, -QPQ^2R^2, -QPQ^2R^2Q, -QPQ^2R^2Q^2, I\}$ which forms the subgroup D_6 (the dihedral group with 6 elements). The cardinality of the orbit (the number of vertices of the tetrahedron) is $\frac{24}{6} = 4$. The action of $\mathbf{3}'$ on the point $\frac{1}{\sqrt{3}}(-1, -1, -1)$ produces the orbit,

$$\frac{1}{\sqrt{3}} \{(-1, -1, -1), (-1, 1, 1), (1, -1, 1), (1, 1, -1)\}, \quad (130)$$

forming a second tetrahedron which is the space inversion of the first².

The group action of $\mathbf{3}'$ on the point $(1, 0, 0)$ produces the orbit,

$$\{(\pm 1, 0, 0), (0, \pm 1, 0), (0, 0, \pm 1)\}, \quad (131)$$

²Combining the two tetrahedra, we obtain a cube.

which forms an octahedron, Figure 5(b). The stabiliser of the point $(1, 0, 0)^T$ is $\{R^2, -QPQ^2, -QPQ^2R^2, I\}$ which forms the subgroup $C_2 \times C_2$. The cardinality of the orbit is $\frac{24}{2 \times 2} = 6$. The tetrahedra, Eqs.(129,130) and the octahedron, Eq. (131) are the only unique orbits that can be constructed in the case of the representation $\mathbf{3}'$.

The action of $\mathbf{3}'$ on a random point produces an orbit with 24 elements. This situation is shown in Figure 5(c). Here we have a trivial stabiliser. For $\mathbf{3}'$, a point having a non-trivial stabiliser is not the sufficient condition for its orbit to be uniquely defined. Consider a point $\propto \left(\frac{1}{\sqrt{2}} \sin \alpha, -2 \cos \alpha, -\frac{1}{\sqrt{2}} \sin \alpha\right)$, Eq. (79). This point remains invariant under the action of $\{-P, I\}$ which is a non-trivial stabiliser. The orbit of this point has $\frac{24}{2} = 12$ elements and is shown in Figure 5(d). It is clear that this orbit is not unique, rather it depends on the arbitrary parameter α .

Let us study the VEVs used in our model in terms of their residual symmetries and check whether these VEVs can be uniquely defined. Consider the VEV of the doublet flavon, $\langle \eta \rangle \propto \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$, Eq. (35). $P(2)$, Eq. (20), generates the C_2 residual symmetry of $\langle \eta \rangle$,

$$P(2)\langle \eta \rangle = \langle \eta \rangle. \quad (132)$$

This residual symmetry uniquely defines $\langle \eta \rangle^3$.

The flavon χ transforms as $\mathbf{3}$ under S_4 . It is also assigned a C_3 symmetry (ω), Table 2, and hence it has complex degrees of freedom. Its VEV, $\langle \chi \rangle \propto (1, \omega, \bar{\omega})$, Eq. (33), remains invariant under the action of ωQ^2 ,

$$\omega Q^2 \langle \chi \rangle = \langle \chi \rangle. \quad (133)$$

The group element ωQ^2 generates the C_3 subgroup $\{\omega Q^2, \bar{\omega} Q, I\}$, which forms the residual symmetry of $\langle \chi \rangle$. This C_3 residual symmetry uniquely defines $\langle \chi \rangle^3$.

The flavon ϕ transforms as $\mathbf{3}'$ under S_4 . Its VEV $\langle \phi \rangle \propto \left(-\frac{1}{2\sqrt{2}}, -\sqrt{3}, \frac{1}{2\sqrt{2}}\right)$, Eq. (36), has the following C_2 residual symmetry:

$$P(3')\langle \phi \rangle = -P\langle \phi \rangle = \langle \phi \rangle. \quad (134)$$

This symmetry ensures that the third component of the VEV is negative of the first component. However, it does not determine the value of the second component in relation to the others. In fact, this value can not be determined by any of the symmetries of S_4 . Rather we obtained it in terms of α , Eq. (79), which is a continuous function of the parameters, Eq. (77), appearing in the flavon potential, Eq. (70). This potential results in a set of minima that forms the orbit shown in Figure 5(d). We tuned the parameters in the potential to

³up to a proportionality constant.

make α equal to $-\frac{\pi}{6}$. This tuning of the parameters changes the orbit (resizes the four small triangles in Figure 5(d)) as a continuous function and enables us to choose the VEV from among an infinite set of possible alignments. We argue that such a dependence of the VEV on the parameters of the potential goes against the spirit of using discrete symmetries in model building. We can resolve this problem by incorporating additional symmetries in the model. A way to achieve this is by utilising the recently proposed framework of the auxiliary group [33].

5.2 The Framework of the Auxiliary Group

Various discrete subgroups of $U(3)$ with an irreducible triplet representation (A_4 and the higher members of the $\Delta(3n^2)$ series, S_4 and the higher members of the $\Delta(6n^2)$ series, $\Sigma(72 \times 3)$, $PSL(2, 7)$, etc.) combined with Abelian discrete groups (C_2 , C_3 , etc.) have been studied in the literature as discrete flavour groups. In those studies, the subgroups of $U(3)$ with triplets are considered because the fermions exist in three families. However, it was shown in Refs. [57, 58, 59] that by going beyond the $U(3)$ -subgroup paradigm we can effectively avoid certain undesirable cross terms in the flavon potential. These papers used an enlarged flavor group constructed as a semidirect product in which the conventional flavor group (the direct product of a subgroup of $U(3)$ with Abelian groups) appears as the quotient. In Ref. [33, 34], a simpler construction, in which the semidirect product was replaced with a direct product, was studied. We named this construction the ‘framework of the auxiliary group’ and used the notation

$$G_f = G_r \times G_x. \quad (135)$$

where G_f is the enlarged flavor group, G_r is the conventional flavor group and G_x is the so-called auxiliary group. The auxiliary group (G_x) is defined as the group under which the fermions remain invariant. Only the flavons transform nontrivially under it. On the other hand, both the fermions and the flavons transform nontrivially under the ‘conventional’ flavour group (G_r). In the flavon sector, to obtain VEVs similar to Eq. (36), we incorporate symmetries in addition to those originating from G_r with the help of G_x . Since G_x does not act on the fermions, it need not be a subgroup of $U(3)$.

An effective multiplet of a given irreducible representation of G_r can be obtained by coupling together several elementary flavons transforming as irreducible multiplets under $G_r \times G_x$. The vacuum alignments of these elementary multiplets are constructed in such a way that they are invariants under the action of various subgroups of $G_r \times G_x$, i.e. these subgroups form

the residual symmetries of the VEVs of the respective multiplets. Moreover, these VEVs are uniquely defined by their residual symmetries. Even though the VEV of the effective multiplet of G_r may not have any residual symmetry under G_r , it will be uniquely defined in terms of the residual symmetries of its constituent elementary multiplets (transforming under $G_r \times G_x$). We will get a clearer understanding of this framework over the course of the following sections when we apply it to our model.

6 The Group Y_{24}

In this section, we construct a group that we call Y_{24} and briefly study its structure. Y_{24} is later used to recast our model in the framework of the auxiliary group.

Consider the group generated by

$$A = \begin{pmatrix} 0 & 0 & 0 & \bar{\tau} & 0 & 0 \\ 0 & 0 & 0 & 0 & i\bar{\omega} & 0 \\ 0 & 0 & 0 & 0 & 0 & -i\omega \\ \tau & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & \bar{\tau} & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \tau & 0 \\ 0 & 0 & 0 & 0 & 0 & -i\omega \\ 0 & 0 & 0 & i\bar{\omega} & 0 & 0 \end{pmatrix}, \quad (136)$$

where $\tau = e^{i\frac{\pi}{4}}$ and $\bar{\tau} = e^{-i\frac{\pi}{4}}$ are the complex eighth roots of unity. Using A and B , we obtain the group elements $D = (AB)^3$ and $E = (AB)^2$ which are given by

$$D = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}, \quad E = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}. \quad (137)$$

We also obtain the following diagonal group elements consisting of complex phases,

$$C_{\omega 1} = (DA)^8 = \text{diag}(1, 1, 1, 1, \omega, \bar{\omega}), \quad (138)$$

$$C_{\omega 2} = E(DA)^8 E^2 = \text{diag}(1, 1, 1, \bar{\omega}, 1, \omega), \quad (139)$$

$$C_{\omega 3} = DC_{\omega 1}D = \text{diag}(1, \omega, \bar{\omega}, 1, 1, 1), \quad (140)$$

$$C_{\omega 4} = DC_{\omega 2}D = \text{diag}(\bar{\omega}, 1, \omega, 1, 1, 1), \quad (141)$$

$$C_{\tau 1} = (DA)^9 = \text{diag}(\tau, 1, 1, \bar{\tau}, i, -i), \quad (142)$$

$$C_{\tau 2} = E(DA)^9 E^2 = \text{diag}(1, \tau, 1, -i, \bar{\tau}, i), \quad (143)$$

$$C_{\tau 3} = E^2(DA)^9 E = \text{diag}(1, 1, \tau, i, -i, \bar{\tau}). \quad (144)$$

We obtain the structure of the group generated by A and B with the help of the above-mentioned group elements, D , E and C_i . We have

$$D^2 = I, \quad E^3 = I, \quad DE = ED. \quad (145)$$

D and E form the cyclic groups C_2 and C_3 respectively and they commute with each other. Therefore, the two of them form the group C_6 . The diagonal group elements $C_{\omega 1}$, $C_{\omega 2}$, $C_{\omega 3}$ and $C_{\omega 4}$ generate four separate C_3 groups. Similarly, the elements $C_{\tau 1}$, $C_{\tau 2}$ and $C_{\tau 3}$ generate three separate C_8 groups. Multiplication of D and E with the diagonal group elements leads to

$$\begin{aligned} DC_{\omega 1} &= C_{\omega 3}D, & EC_{\omega 1} &= C_{\omega 2}E, \\ DC_{\omega 2} &= C_{\omega 4}D, & EC_{\omega 2} &= C_{\omega 1}^2 C_{\omega 2}^2 E, \\ DC_{\omega 3} &= C_{\omega 1}D, & EC_{\omega 3} &= C_{\omega 4}E, \\ DC_{\omega 4} &= C_{\omega 2}D, & EC_{\omega 4} &= C_{\omega 3}^2 C_{\omega 4}^2 E, \\ DC_{\tau 1} &= C_{\tau 1}^7 C_{\tau 2}^2 C_{\tau 3}^6 D, & EC_{\tau 1} &= C_{\tau 2}E, \\ DC_{\tau 2} &= C_{\tau 1}^6 C_{\tau 2}^7 C_{\tau 3}^2 D, & EC_{\tau 2} &= C_{\tau 3}E, \\ DC_{\tau 3} &= C_{\tau 1}^2 C_{\tau 2}^6 C_{\tau 3}^7 D, & EC_{\tau 3} &= C_{\tau 1}E. \end{aligned} \quad (146)$$

Also, the diagonal group elements, C_i , commute with each other,

$$C_i C_j = C_j C_i \quad (147)$$

Eqs. (145, 146, 147) prove that the group generated by D , E and C_i is finite. The generators A and B can be expressed in terms of D , E and C_i ,

$$A = C_{\omega 3}^2 C_{\tau 1}^{15} C_{\tau 2}^2 C_{\tau 3}^6 D, \quad B = C_{\omega 1} C_{\tau 1}^{15} E^2. \quad (148)$$

Since A and B can be expressed in terms of D , E and C_i and vice versa, the group generated by A and B is the same as the one generated by D , E and C_i . The diagonal elements $C_{\omega 1}$, $C_{\omega 2}$, $C_{\omega 3}$, $C_{\omega 4}$, $C_{\tau 1}$, $C_{\tau 2}$, $C_{\tau 3}$ form the group $C_3 \times C_3 \times C_3 \times C_3 \times C_8 \times C_8 \times C_8$ which is equivalent to $C_3 \times C_{24} \times C_{24} \times C_{24}$. As mentioned earlier, D and E form the group C_6 . Therefore, the group generated by D , E and C_i (or the one generated by A and B) is simply the semidirect product,

$$G(A, B) = (C_3 \times C_{24} \times C_{24} \times C_{24}) \rtimes C_6. \quad (149)$$

We enlarge this group by adding another generator,

$$C_1 = \text{diag}(1, 1, 1, 1, -1, -1). \quad (150)$$

Using C_1 , we obtain the group element,

$$C_2 = EC_1 E^2 = \text{diag}(1, 1, 1, -1, 1, -1). \quad (151)$$

The finiteness of the enlarged group is ensured by the following equations:

$$\begin{aligned} DC_1 &= C_1 C_{\tau 2}^4 C_{\tau 3}^4 D, & EC_1 &= C_2 E, \\ DC_2 &= C_2 C_{\tau 1}^4 C_{\tau 3}^4 D, & EC_2 &= C_1 C_2 E. \end{aligned} \quad (152)$$

C_1 and C_2 generate two C_2 groups. As a result, we obtain

$$G(A, B, C_1) = (C_2 \times C_2 \times C_3 \times C_{24} \times C_{24} \times C_{24}) \rtimes C_6. \quad (153)$$

For convenience, we name $G(A, B, C_1)$ as Y_{24} ,

$$Y_{24} = G(A, B, C_1) = (C_2 \times C_6 \times C_{24} \times C_{24} \times C_{24}) \rtimes C_6. \quad (154)$$

Any element of Y_{24} can be uniquely expressed as,

$$g_i = C_1^{i_1} C_2^{i_2} C_{\omega 1}^{j_1} C_{\omega 2}^{j_2} C_{\omega 3}^{j_3} C_{\omega 4}^{j_4} C_{\tau 1}^{k_1} C_{\tau 2}^{k_2} C_{\tau 3}^{k_3} D^{i_3} E^{j_5}, \quad (155)$$

where $i_x = 1, 2$, $j_x = 1, 2, 3$ and $k_x = 1, 2, \dots, 8$. This group has a total of $2 \times 6 \times 24^3 \times 6 = 995328$ elements. We used the computational package GAP [60] to obtain the group with A , B and C_1 as the generators. According to GAP, such a group has 995328 elements confirming our analysis.

We name the sextet representation of Y_{24} defined by the generators Eqs. (136, 150) as $\mathbf{6}$. Its tensor product with itself leads to the following expansion,

$$\mathbf{6} \times \mathbf{6} = \mathbf{6}_0 + \mathbf{6}_1 + \mathbf{6}_2 + \mathbf{3} + \mathbf{6}_1 + \mathbf{6}_2 + \mathbf{3}'. \quad (156)$$

The first four multiplets in the RHS of the above equation form the symmetric part and the next three form the antisymmetric part of the tensor product. Besides $\mathbf{6}$, we also utilise the representation $\mathbf{6}_1$ in this paper. $\mathbf{6}_1$ constructed from the symmetric part of Eq. (156) is given by

$$\begin{aligned} \mathbf{6}_1 \equiv & \\ & (\{a_2, b_3\}, \{a_3, b_1\}, \{a_1, b_2\}, \{a_5, b_6\}, \{a_6, b_4\}, \{a_4, b_5\})^T \end{aligned} \quad (157)$$

where a_i and b_i form sextets transforming under $\mathbf{6}$ and $\{a_i, b_j\} = a_i b_j + a_j b_i$. In the basis, Eq. (157), the generators of $\mathbf{6}_1$ are

$$\begin{aligned} A(\mathbf{6}_1) &= \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\tau\omega & 0 \\ 0 & 0 & 0 & 0 & 0 & \tau\bar{\omega} \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \tau & 0 & 0 & 0 & 0 \\ 0 & 0 & \tau & 0 & 0 & 0 \end{pmatrix}, \quad B(\mathbf{6}_1) = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \bar{\tau} & 0 & 0 & 0 \\ \bar{\tau} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -\bar{\tau}\bar{\omega} \\ 0 & 0 & 0 & \bar{\tau}\omega & 0 & 0 \end{pmatrix}, \\ C_1(\mathbf{6}_1) &= \text{diag}(1, 1, 1, 1, -1, -1). \end{aligned} \quad (158)$$

This completes our discussion of the structure of the group Y_{24} .

	L	e_R	μ_R	τ_R	N	ρ	χ	\acute{S}	\grave{S}	S	$\acute{\phi}$	$\grave{\phi}$	Δ
S_4	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{3}'$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{3}$	$\mathbf{3}$	$\mathbf{1}$
$C_4 \times C_3$	i	i	$i\omega$	$i\bar{\omega}$	i	1	ω	1	1	-1	1	1	-1
C_2	-1	1	1	-1	1	-1	1	1	1	1	1	1	1
Y_{24}	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{6}$	$\mathbf{6}$	$\bar{\mathbf{6}}_1$
C_2	1	1	1	1	1	1	1	1	1	1	$*$	$*$	$*$
C_2	1	1	1	1	1	1	1	-1	1	-1	-1	1	-1
C_2	1	1	1	1	1	1	1	1	-1	-1	1	-1	-1

Table 3: The fields in the recast model as the multiplets under $(S_4 \times C_4 \times C_3 \times C_2) \times (Y_{24} \rtimes C_2 \times C_2 \times C_2)$.

7 The Model recast using

$$(S_4 \times C_4 \times C_3 \times C_2) \times (Y_{24} \rtimes C_2 \times C_2 \times C_2)$$

The field content of the model recast in the framework of the auxiliary group is given in Table 3. The left-handed lepton field (L), the right-handed neutrino field (N) and the flavon field χ are triplets ($\mathbf{3}$) under S_4 and invariants under Y_{24} . The flavons $\acute{\phi}$ and $\grave{\phi}$ are not only triplets ($\mathbf{3}$) under S_4 , but also sextets ($\mathbf{6}$) under Y_{24} . Δ transforms as $\bar{\mathbf{6}}_1$ under Y_{24} . We impose the complex conjugation symmetry, $\acute{\phi} \rightarrow \acute{\phi}^*$, $\grave{\phi} \rightarrow \grave{\phi}^*$ and $\Delta \rightarrow \Delta^*$, represented as the C_2 in the 5th row in Table 3. Since Y_{24} is a complex group, combining it with conjugation leads to the semidirect product, $Y_{24} \rtimes C_2$. Note that this conjugation symmetry is different from CP symmetry which involves the transformation of not only $\acute{\phi}$, $\grave{\phi}$ and Δ but also all other complex scalars as well fermions in the model. Two more C_2 groups are introduced (the last two rows in the table), so that \acute{S} , \grave{S} and S (similarly $\acute{\phi}$, $\grave{\phi}$ and Δ) couple together in the Lagrangian. The fermions, L , e_R , μ_R , τ_R and N are invariant under $Y_{24} \rtimes C_2 \times C_2 \times C_2$, hence it forms the auxiliary group, G_x . The rest of the flavour group $(S_4 \times C_4 \times C_3 \times C_2)$, which also appears in Table 2, forms G_r .

Based on our field content and their symmetries, we construct the following Lagrangian:

$$\begin{aligned} \mathcal{L} = & y_\tau \bar{L} \frac{\chi}{\Lambda} \tau_R H + y_\mu \bar{L} \frac{\rho \chi^*}{\Lambda^2} \mu_R H + y_e \bar{L} \frac{\rho(\chi^* \chi)_{\mathbf{3}'}}{\Lambda^3} e_R H \\ & + y_\nu \bar{L} N \frac{\rho}{\Lambda} \tilde{H} + y_s (\bar{N}^c N)_1 \frac{1}{\Lambda^2} \acute{S} S \grave{S} \\ & + y_\phi (\bar{N}^c N)_{\mathbf{3}'} \frac{1}{\Lambda^2} \left[(\acute{\phi} \Delta \grave{\phi})_{\mathbf{3}'} + (\acute{\phi}^* \Delta^* \grave{\phi}^*)_{\mathbf{3}'} \right]. \end{aligned} \quad (159)$$

As was done in our original Lagrangian, Eq. (30), we impose CP symmetry in the above Lagrangian, leading to real coupling constants, y_x . In Eq. (159), the elementary multiplets $\acute{\phi}$, $\grave{\phi}$ and Δ couple together resulting in

the effective triplet ($\mathbf{3}'$) of S_4 ,

$$\frac{1}{\Lambda^2} \left[(\acute{\phi} \Delta \grave{\phi})_{\mathbf{3}'} + (\acute{\phi}^* \Delta^* \grave{\phi}^*)_{\mathbf{3}'} \right]. \quad (160)$$

Eq. (159) is the same as Eq. (30), except for the fact that the triplet ϕ is replaced with the effective triplet and the singlet s is replaced with the effective singlet $\frac{1}{\Lambda^2} \acute{S} S \grave{S}$ ⁴.

The various components of the flavons Δ , $\acute{\phi}$ and $\grave{\phi}$ can be expressed as Δ_α , $\acute{\phi}_{\beta m}$ and $\grave{\phi}_{\gamma n}$ where the Greek and the Latin indices correspond to Y_{24} and S_4 respectively. To explicitly calculate the expression of the effective triplet, Eq. (160), in terms of the components of the elementary flavons, we use the tensor product expansions given in Eqs. (24, 157). Thus we obtain

$$(\acute{\phi} \Delta \grave{\phi})_{\mathbf{3}'} = \sum C'_{lmn} C_{\alpha\beta\gamma} \acute{\phi}_{\beta m} \grave{\phi}_{\gamma n} \Delta_\alpha, \quad (161)$$

where the summation is over all the repeated indices and

$$\begin{aligned} C'_{lmn} &= 1 \quad \forall l \neq m \neq n \\ &= 0 \quad \text{otherwise} \end{aligned} \quad (162)$$

and

$$\begin{aligned} C_{lmn} &= 1 \quad \forall (l \bmod 3) \neq (m \bmod 3) \neq (n \bmod 3) \quad \& \\ &\quad (l-1) \bmod 3 = (m-1) \bmod 3 = (n-1) \bmod 3 \\ &= 0 \quad \text{otherwise} \end{aligned} \quad (163)$$

are the C-G coefficients corresponding to Eq. (24) and Eq. (157) respectively. In Eq. (163), ‘mod’ denotes the modulo operation and ‘div’ denotes the integer division. $(\acute{\phi}^* \Delta^* \grave{\phi}^*)_{\mathbf{3}'}$ is simply the complex conjugate of $(\acute{\phi} \Delta \grave{\phi})_{\mathbf{3}'}$.

The elementary flavons are assigned the following VEVs:

$$\langle \acute{\phi} \rangle = v_\phi \begin{pmatrix} 1 & 0 & 0 \\ 0 & \tau & 0 \\ 0 & 0 & 1 \\ -i\omega & 0 & 0 \\ 0 & \bar{\tau} & 0 \\ 0 & 0 & i\bar{\omega} \end{pmatrix}, \quad \langle \grave{\phi} \rangle = v_\phi \begin{pmatrix} i\omega & 0 & 0 \\ 0 & \tau & 0 \\ 0 & 0 & -i\bar{\omega} \\ -1 & 0 & 0 \\ 0 & \bar{\tau} & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad (164)$$

$$\langle \Delta \rangle = v_\Delta (1 \ 1 \ 1 \ 1 \ 1 \ 1), \quad (165)$$

where v_ϕ and v_Δ are real. Here, $\langle \acute{\phi} \rangle$ and $\langle \grave{\phi} \rangle$ are given in matrix forms with the rows (columns) representing Y_{24}

⁴The only purpose of introducing the singlets \acute{S} and \grave{S} in the Lagrangian is to ensure that the effective singlet term has the same dimensionality as the effective triplet term.

(S_4). Using Eqs. (161, 164, 165), we obtain the vacuum alignment of the effective triplet, Eq. (160),

$$\begin{aligned} \frac{1}{\Lambda^2} \left\langle \left(\dot{\phi} \Delta \dot{\phi} \right)_{\mathbf{3}'} + \left(\dot{\phi}^* \Delta^* \dot{\phi}^* \right)_{\mathbf{3}'} \right\rangle \\ = \frac{4v_{\dot{\phi}}^2 v_{\Delta}}{\Lambda^2} \left(-\frac{1}{2\sqrt{2}}, -\sqrt{3}, \frac{1}{2\sqrt{2}} \right). \end{aligned} \quad (166)$$

This alignment is proportional to the VEV of the triplet flavon that we originally proposed in Eq. (36). We also assign

$$\langle \dot{S} \rangle = \langle \dot{S} \rangle = \langle S \rangle = v_S \quad (167)$$

so that

$$\frac{1}{\Lambda^2} \langle \dot{S} S \dot{S} \rangle = \frac{v_S^3}{\Lambda^2} \quad (168)$$

which corresponds to the VEV of the singlet flavon, Eq. (34)⁵. We noted in Section 4 that the VEVs of the flavons ϕ and s are of the order of 10^{11} GeV. In the recast model, we have replaced these flavons with the corresponding effective multiplets. If we assume that the VEVs of all flavons, say v_x , in the recast model are at the same scale, we obtain $v_x \approx 10^{15}$ GeV and $\Lambda \approx 10^{17}$ GeV.

7.1 The Residual Symmetries of $\langle \dot{\phi} \rangle$, $\langle \dot{\phi} \rangle$ and $\langle \Delta \rangle$

In this section, we show that each of the VEVs, $\langle \dot{\phi} \rangle$, $\langle \dot{\phi} \rangle$, $\langle \Delta \rangle$, is an invariant under the action of certain elements of the flavour group. The VEV does not fully break the flavour group, rather it breaks the group into a specific subgroup that constitutes its residual symmetries. In this way, the alignment of each of these VEVs is uniquely defined.

Let us first study the symmetries of the alignment

$$\langle \Phi \rangle = |\langle \dot{\phi} \rangle| = |\langle \dot{\phi} \rangle| = v_{\Phi} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (169)$$

⁵We did not include the S_4 effective singlet, $\frac{1}{\Lambda^2} \left[\left(\dot{\phi} \Delta \dot{\phi} \right)_1 + \left(\dot{\phi}^* \Delta^* \dot{\phi}^* \right)_1 \right]$, and the doublet, $\frac{1}{\Lambda^2} \left[\left(\dot{\phi} \Delta \dot{\phi} \right)_2 + \left(\dot{\phi}^* \Delta^* \dot{\phi}^* \right)_2 \right]$, in the Lagrangian, Eq. (159), because they vanish for the given set of VEVs, Eqs. (164, 165).

where v_{Φ} is real. This alignment remains invariant under the following group actions:

$$\mathcal{O}_1 \langle \Phi \rangle = C_1 D C_1 D \langle \Phi \rangle Q R Q P Q = \langle \Phi \rangle, \quad (170)$$

$$\mathcal{O}_2 \langle \Phi \rangle = C_2 D C_2 D \langle \Phi \rangle R Q P Q^2 = \langle \Phi \rangle, \quad (171)$$

$$\mathcal{O}_3 \langle \Phi \rangle = D \langle \Phi \rangle I = \langle \Phi \rangle, \quad (172)$$

$$\mathcal{O}_4 \langle \Phi \rangle = E \langle \Phi \rangle Q = \langle \Phi \rangle, \quad (173)$$

$$\mathcal{O}_5 \langle \Phi \rangle = \langle \Phi \rangle^* = \langle \Phi \rangle, \quad (174)$$

where

$$C_1 D C_1 D = \text{diag}(1, -1, -1, 1, -1, -1), \quad (175)$$

$$Q R Q P Q = \text{diag}(1, -1, -1), \quad (176)$$

$$C_2 D C_2 D = \text{diag}(-1, 1, -1, -1, 1, -1), \quad (177)$$

$$R Q P Q^2 = \text{diag}(-1, 1, -1). \quad (178)$$

The group actions \mathcal{O}_1 and \mathcal{O}_2 , Eqs. (170, 171), multiply certain components of $\langle \Phi \rangle$ with -1 . Invariance under this action ensures that those components vanish, i.e., Eqs. (170, 171) ensure that all the components marked with zeros in Eq. (169) vanish. \mathcal{O}_3 exchanges the first three rows of $\langle \Phi \rangle$ with the last three. Invariance under \mathcal{O}_3 ensures that the upper three rows are equal to the lower three. This condition is satisfied by $\langle \Phi \rangle$. The group action \mathcal{O}_4 cycles the following sets of components: $\{(1,1), (2,2), (3,3)\}$, $\{(1,3), (2,1), (3,2)\}$, $\{(1,2), (2,3), (3,1)\}$, $\{(4,1), (5,2), (6,3)\}$, $\{(4,3), (5,1), (6,2)\}$, $\{(4,2), (5,3), (6,1)\}$. Invariance under this action ensures that the components within each set are equal. This condition is also satisfied by the VEV. Finally, the invariance under the action of \mathcal{O}_5 ensures that the VEV is real. The conditions, Eqs. (170, 171, 172, 173, 174) uniquely lead to the required vacuum alignment, Eq. (169) up to a real proportionality constant.

\mathcal{O}_1 , \mathcal{O}_2 , \mathcal{O}_3 and \mathcal{O}_5 generate various C_2 subgroups and \mathcal{O}_4 generates a C_3 subgroup of the flavour group. It can be shown that the group generated by \mathcal{O}_1 , \mathcal{O}_2 , \mathcal{O}_3 , \mathcal{O}_4 and \mathcal{O}_5 has 192 elements and has the structure $(C_2 \times C_2 \times C_2 \times C_2) \rtimes C_3 \times C_2 \times C_2$. This group represents the residual symmetries of $\langle \Phi \rangle$ and it uniquely defines $\langle \Phi \rangle$.

$\langle \dot{\phi} \rangle$ and $\langle \dot{\phi} \rangle$, Eqs. (164), can also be defined in terms of specific $(C_2 \times C_2 \times C_2 \times C_2) \rtimes C_3 \times C_2 \times C_2$ subgroups of the flavour group. Consider the group elements

$$\dot{g} = C_{\omega 2}^2 C_{\tau 2} = \text{diag}(1, \tau, 1, -i\omega, \bar{\tau}, i\bar{\omega}) \quad (179)$$

$$\dot{g} = C_2 D C_{\omega 2}^2 C_{\tau 2}^7 D = \text{diag}(i\omega, \tau, -i\bar{\omega}, -1, \bar{\tau}, -1) \quad (180)$$

We may obtain $\langle \dot{\phi} \rangle$ and $\langle \dot{\phi} \rangle$ by the action of \dot{g} and \dot{g} respectively on $\langle \Phi \rangle$,

$$\dot{g} \langle \Phi \rangle = \langle \dot{\phi} \rangle, \quad \dot{g} \langle \Phi \rangle = \langle \dot{\phi} \rangle. \quad (181)$$

We define a new set of group elements,

$$\hat{\mathcal{O}}_i = \hat{g}\mathcal{O}_i\hat{g}^{-1}, \quad \hat{\mathcal{O}}_i = \hat{g}\mathcal{O}_i\hat{g}^{-1}, \quad (182)$$

where $i = 1, \dots, 5$. These elements uniquely define $\langle \hat{\Phi} \rangle$ and $\langle \hat{\Phi} \rangle$ in terms of their residual symmetries,

$$\hat{\mathcal{O}}_i \langle \hat{\Phi} \rangle = \langle \hat{\Phi} \rangle, \quad \hat{\mathcal{O}}_i \langle \hat{\Phi} \rangle = \langle \hat{\Phi} \rangle. \quad (183)$$

$\langle \hat{\Phi} \rangle$ and $\langle \hat{\Phi} \rangle$, Eqs. (164), uniquely break the flavour group into separate $(C_2 \times C_2 \times C_2 \times C_2) \rtimes C_3 \times C_2 \times C_2$ groups generated by $\hat{\mathcal{O}}_i$ and $\hat{\mathcal{O}}_i$ respectively. $\langle \hat{\Phi} \rangle$, $\langle \hat{\Phi} \rangle$ and $\langle \hat{\Phi} \rangle$ belong to the same orbit and the $(C_2 \times C_2 \times C_2 \times C_2) \rtimes C_3 \times C_2 \times C_2$ groups generated by \mathcal{O}_i , $\hat{\mathcal{O}}_i$ and $\hat{\mathcal{O}}_i$ form their respective stabilisers.

The flavon Δ transforms as $\bar{\mathbf{6}}_1$. The generators of $\bar{\mathbf{6}}_1$ are given in Eqs. (158). Using these generators we can show that $D(\bar{\mathbf{6}}_1) = D(\mathbf{6}) = D$ and $E(\bar{\mathbf{6}}_1) = E(\mathbf{6}) = E$. The vacuum alignment, $\langle \Delta \rangle = v_\Delta(1, 1, 1, 1, 1, 1)$, Eq. (165), remains invariant under the following group actions:

$$D\langle \Delta \rangle = \langle \Delta \rangle, \quad E\langle \Delta \rangle = \langle \Delta \rangle, \quad \langle \Delta \rangle^* = \langle \Delta \rangle. \quad (184)$$

D interchanges the first three components of $\langle \Delta \rangle$ with the last three. Invariance under this action ensures that the 1st, the 2nd and the 3rd components are equal to the 4th, the 5th and the 6th components respectively. E cycles the first three components as well as the last three. Invariance under E ensures that the first three components are equal and also the last three components are equal. Invariance under both D and E implies that all the elements of the sextet are equal which is true for $\langle \Delta \rangle$. The third condition, $\langle \Delta \rangle^* = \langle \Delta \rangle$, implies that the VEV is real. Eqs. (184) uniquely results in the required VEV, Eq. (165). D and E generate C_2 and C_3 respectively and together they generate C_6 . We also have a C_2 symmetry representing the conjugation. Therefore, $\langle \Delta \rangle$ has the residual symmetry, $C_6 \times C_2$. This residual symmetry uniquely defines $\langle \Delta \rangle$.

Construction of the most general potential for the flavons presented in this paper would be quite tedious because of the very large number of degrees of freedom involved. We would also need driving fields to ensure that accidental Lie symmetries of the renormalizable potential are broken to obtain the required finite symmetries. Such details are beyond the scope of this paper. However, it is known [34, 61, 62] that for the potential constructed with an irreducible multiplet, every alignment that is fully determined by its residual symmetries is guaranteed to form a stationary point. In our model, VEVs of all irreducible multiplets are fully determined by their respective residual symmetries. Therefore, the generation of these VEVs is automatically ensured. Undesirable cross terms among the various multiplets can

be avoided by invoking mechanisms that are widely assumed in flavour models such as localisation in branes in an extradimensional framework.

The residual symmetries that define the VEVs of the individual elementary multiplets $\hat{\Phi}$, $\hat{\Phi}$ and Δ may not survive when the flavons are taken together to form the effective triplet. On the other hand, we utilise the fact that the flavon VEVs are oriented along different directions (with different residual symmetries) to obtain the required VEV of the effective triplet. Even though the effective triplet is an invariant in the space of the auxiliary group, it is constructed from the multiplets ($\hat{\Phi}$, $\hat{\Phi}$ and Δ) transforming non-trivially in that space. This implies that the components of the effective triplet hold information about how these multiplets are oriented relative to each other⁶. We may recall that the second half of this paper is simply the pursuit of finding an explanation for the angle $\alpha = -\frac{\pi}{6}$ proposed for the triplet VEV, Eq. (79). The relative orientations of the VEVs of $\hat{\Phi}$, $\hat{\Phi}$ and Δ in the higher dimensional flavour space (uniquely defined by their respective residual symmetries) contains precisely the information about this angle, which in turn gets transferred to the VEV of the effective triplet living in the lower dimensional space of the S_4 triplet representation.

The order of the symmetry group in the model is extremely large ($> 10^6$). This can be viewed upon as an unpleasant feature of the framework of the auxiliary group. In this context, we may note that Ref. [57] used a group constructed as a wreath product (a special case of the semidirect product) with order $> 10^4$ to decouple the flavon VEVs in the charged-lepton and neutrino sectors so that tribimaximal mixing (TBM) can be obtained. We may contrast this with the size of A_4 group (order 12), which is widely used for constructing TBM even though it results in dangerous cross terms that spoil the required VEVs. As stated earlier, model builders forbid these terms with the help of assumptions like extra dimensions. Having a larger flavour group, as in Ref. [57], can be considered as a necessary trade-off to avoid such assumptions. A simpler semidirect product, $Q_8 \rtimes A_4$, which is smaller than the wreath product, was later proposed [58, 59] to achieve decoupling and obtain TBM. In the current paper, the flavour group is given as the direct product, $G_f = G_r \times G_x$. We may be able to reduce the size of G_f by a couple of orders of magnitude if we replace the direct product with a suitable semidirect product. The presence of a large number of zeros in the VEVs of the elementary multiplets, Eqs. (164), is a hint towards this.

⁶As an analogy, consider the scalar product of two vectors. It is nothing but an invariant constructed from the vectors and it holds information about their relative orientation.

Unlike Refs.[57,58,59] where the purpose of using an enlarged group was to decouple VEVs, the current paper uses it to construct a novel VEV for the effective multiplet, which in turn generates a small reactor angle $\sin^2 \theta_{13} = \frac{1}{3} \sin^2 \frac{\pi}{12}$. We constructed the group Y_{24} (the main constituent of our auxiliary group) as a six-dimensional representation containing phases $e^{i\frac{\pi}{12}}$. Such constructions are rather unavoidable given our philosophy of modelling small mixing angles using finite groups. Consequently, we obtain very large groups. Nevertheless, we argue that rather than looking at flavor groups from the perspective of their size, it would be instructive to study the nature of their generators. Even though the group Y_{24} is very large, its generators are fairly simple, Eqs. (136,150). They correspond to certain permutations and diagonal multiplications with complex phases. If there is an underlying mechanism behind the origin of these generators, we may expect it to be simple.

8 Conclusion

TM_1 mixing preserves the first column of the tribimaximal mixing. TM_1 can be parametrised using an angle (θ) and a phase (ζ). In this parametrisation, the reactor mixing angle is given by $\sin^2 \theta_{13} = \frac{1}{3} \sin^2 \theta$. In this paper, we construct a type-1 seesaw model based on the flavour group, $S_4 \times C_4 \times C_3$, and obtain TM_1 mixing with $\theta = -\frac{\pi}{12}$ and $\zeta = \frac{\pi}{2}$ consistent with the current experimental data. The mixing obtained exhibits μ - τ reflection symmetry so that we have $\theta_{23} = \frac{\pi}{4}$. We also obtain maximal CP violation with $\delta = -\frac{\pi}{2}$. The light neutrino masses are found to be functions of two model parameters and by fitting them with the experimental values of the mass-squared-differences we predict the individual masses. They follow the normal ordering. We also predict the effective neutrino mass applicable to the neutrinoless double-beta decay.

To construct the fermion mass terms, we introduce the flavons χ , s , ϕ which transform as **3**, **1** and **3'** respectively under S_4 . We construct the flavon potentials which when minimised lead to their vacuum expectation values. The flavon χ couples in the charged-lepton sector and its vacuum alignment leads to the 3×3 -trimaximal contribution to the neutrino mixing matrix. The flavons s and ϕ couple in the Majorana sector. Their vacuum alignments constitute the Majorana mass matrix. The product of the 3×3 -trimaximal matrix and the diagonalising matrix from the Majorana sector generates the TM_1 mixing with $\theta = -\frac{\pi}{12}$ and $\zeta = \frac{\pi}{2}$. Obtaining the required vacuum alignment of ϕ entails tuning of the parameters in the flavon potential. This

tuning implies that the symmetries of S_4 are not sufficient to uniquely determine this vacuum alignment. Therefore, we recast the model by introducing additional symmetries using the framework of the auxiliary group.

In this framework, the flavour group is given as the direct product of a ‘conventional’ flavour group and an auxiliary group. Our original flavour group ($S_4 \times C_4 \times C_3 \times C_2$) takes the place of the conventional flavour group in the recast model. The auxiliary group is given by $Y_{24} \rtimes C_2 \times C_2 \times C_2$ where Y_{24} is a discrete group that we constructed with the required symmetries. We introduce the flavons $\tilde{\phi}$, $\tilde{\phi}$ and Δ which transform under both the conventional flavour group and the auxiliary group. They are coupled together to obtain $(\tilde{\phi}\Delta\tilde{\phi})_{\mathbf{3}'} + (\tilde{\phi}^*\Delta^*\tilde{\phi}^*)_{\mathbf{3}'}$ which is a triplet under S_4 and an invariant under the auxiliary group. This effective triplet transforms the same way as the triplet ϕ present in the original model. We assign specific VEVs to $\tilde{\phi}$, $\tilde{\phi}$ and Δ and identify the residual symmetries corresponding to each of them as a subgroup of the flavour group, i.e. a subgroup of $(S_4 \times C_4 \times C_3 \times C_2) \times (Y_{24} \rtimes (C_2 \times C_2 \times C_2))$. We find that these subgroups uniquely define the VEVs. Using these VEVs, we uniquely obtain the VEV of the effective triplet also, which is in agreement with the VEV of ϕ proposed in the original model.

Acknowledgements I would like to thank Paul Harrison and Bill Scott for the stimulating discussions. I acknowledge the help and support from Ambar Ghosal and Debasish Majumdar. I am grateful to Sujatha Ramakrishnan for helping me with constructing the figures.

References

1. P. F. Harrison, D. H. Perkins and W. G. Scott, *Tri-bimaximal mixing and the neutrino oscillation data*, *Phys. Lett. B* **530** (2002) 167 [[hep-ph/0202074](#)].
2. C. H. Albright and W. Rodejohann, *Comparing Trimaximal Mixing and Its Variants with Deviations from Tri-bimaximal Mixing*, *Eur. Phys. J. C* **62** (2009) 599 [[0812.0436](#)].
3. C. H. Albright, A. Dueck and W. Rodejohann, *Possible Alternatives to Tri-bimaximal Mixing*, *Eur. Phys. J. C* **70** (2010) 1099 [[1004.2798](#)].
4. Z.-z. Xing and S. Zhou, *Tri-bimaximal Neutrino Mixing and Flavor-dependent Resonant Leptogenesis*, *Phys. Lett. B* **653** (2007) 278 [[hep-ph/0607302](#)].
5. C. Luhn, *Trimaximal TM_1 neutrino mixing in S_4 with spontaneous CP violation*, *Nucl. Phys. B* **875** (2013) 80 [[1306.2358](#)].
6. C.-C. Li and G.-J. Ding, *Generalised CP and trimaximal TM_1 lepton mixing in S_4 family symmetry*, *Nucl. Phys. B* **881** (2014) 206 [[1312.4401](#)].
7. I. de Medeiros Varzielas and L. Lavoura, *Flavour models for TM_1 lepton mixing*, *J. Phys. G* **40** (2013) 085002 [[1212.3247](#)].

8. Y. Shimizu, K. Takagi and M. Tanimoto, *Towards the minimal seesaw model via CP violation of neutrinos*, *JHEP* **11** (2017) 201 [1709.02136].
9. Z.-h. Zhao, *Modified Friedberg-Lee symmetry for neutrino mixing*, *Phys. Rev. D* **92** (2015) 113001 [1509.06915].
10. R. R. Gautam, *Trimaximal mixing with a texture zero*, *Phys. Rev. D* **97** (2018) 055022 [1802.00425].
11. W. Rodejohann and X.-J. Xu, *Trimaximal μ - τ reflection symmetry*, *Phys. Rev. D* **96** (2017) 055039 [1705.02027].
12. S. Antusch, S. F. King, C. Luhn and M. Spinrath, *Trimaximal mixing with predicted θ_{13} from a new type of constrained sequential dominance*, *Nucl. Phys. B* **856** (2012) 328 [1108.4278].
13. S. F. King, *Minimal see-saw model predicting best fit lepton mixing angles*, *Phys. Lett. B* **724** (2013) 92 [1305.4846].
14. S. F. King, *Minimal predictive see-saw model with normal neutrino mass hierarchy*, *JHEP* **07** (2013) 137 [1304.6264].
15. S. F. King, *Littlest Seesaw*, *JHEP* **02** (2016) 085 [1512.07531].
16. S. F. King and C. Luhn, *Littlest Seesaw model from $S_4 \times U(1)$* , *JHEP* **09** (2016) 023 [1607.05276].
17. C. Jarlskog, *Commutator of the Quark Mass Matrices in the Standard Electroweak Model and a Measure of Maximal CP Violation*, *Phys. Rev. Lett.* **55** (1985) 1039.
18. C. Jarlskog, *A Basis Independent Formulation of the Connection Between Quark Mass Matrices, CP Violation and Experiment*, *Z. Phys. C* **29** (1985) 491.
19. C. Jarlskog, *JARLSKOG RESPONDS*, *Phys. Rev. Lett.* **57** (1986) 2875.
20. C. Jarlskog, *REPLY TO 'COMMENT ON JARLSKOG'S CONDITIONS FOR CP INVARIANCE.'*, *Phys. Rev. D* **39** (1989) 988.
21. C. Jarlskog, *Invariants of lepton mass matrices and CP and T violation in neutrino oscillations*, *Phys. Lett. B* **609** (2005) 323 [hep-ph/0412288].
22. I. Esteban, M. C. Gonzalez-Garcia, A. Hernandez-Cabezudo, M. Maltoni and T. Schwetz, *Global analysis of three-flavour neutrino oscillations: synergies and tensions in the determination of θ_{23} , δ_{CP} , and the mass ordering*, *JHEP* **01** (2019) 106 [1811.05487].
23. H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada and M. Tanimoto, *Non-Abelian Discrete Symmetries in Particle Physics*, *Prog. Theor. Phys. Suppl.* **183** (2010) 1 [1003.3552].
24. G. Altarelli and F. Feruglio, *Discrete Flavor Symmetries and Models of Neutrino Mixing*, *Rev. Mod. Phys.* **82** (2010) 2701 [1002.0211].
25. W. Grimus, *Theory of Neutrino Masses and Mixing*, *Phys. Part. Nucl.* **42** (2011) 566 [1101.0137].
26. D. Hernandez and A. Y. Smirnov, *Lepton mixing and discrete symmetries*, *Phys. Rev. D* **86** (2012) 053014 [1204.0445].
27. H. Ishimori, T. Kobayashi, H. Ohki, H. Okada, Y. Shimizu and M. Tanimoto, *An introduction to non-Abelian discrete symmetries for particle physicists*, vol. 858. 2012, 10.1007/978-3-642-30805-5.
28. S. F. King and C. Luhn, *Neutrino Mass and Mixing with Discrete Symmetry*, *Rept. Prog. Phys.* **76** (2013) 056201 [1301.1340].
29. S. F. King, A. Merle, S. Morisi, Y. Shimizu and M. Tanimoto, *Neutrino Mass and Mixing: from Theory to Experiment*, *New J. Phys.* **16** (2014) 045018 [1402.4271].
30. S. F. King, *Unified Models of Neutrinos, Flavour and CP Violation*, *Prog. Part. Nucl. Phys.* **94** (2017) 217 [1701.04413].
31. D. Meloni, *GUT and flavor models for neutrino masses and mixing*, *Front. in Phys.* **5** (2017) 43 [1709.02662].
32. S. T. Petcov, *Discrete Flavour Symmetries, Neutrino Mixing and Leptonic CP Violation*, *Eur. Phys. J. C* **78** (2018) 709 [1711.10806].
33. R. Krishnan, *Fully Constrained Mass Matrix: Can Symmetries alone determine the Flavon Vacuum Alignments?*, *Phys. Rev. D* **101** (2020) 075004 [1901.01205].
34. R. Krishnan, *Symmetries of stationary points of the G-invariant potential and the framework of the auxiliary group*, *Phys. Rev. D* **103** (2021) 051701 [2011.11653].
35. T. Brown, S. Pakvasa, H. Sugawara and Y. Yamanaka, *Neutrino Masses, Mixing and Oscillations in $S(4)$ Model of Permutation Symmetry*, *Phys. Rev. D* **30** (1984) 255.
36. D.-G. Lee and R. N. Mohapatra, *An $SO(10) \times S(4)$ scenario for naturally degenerate neutrinos*, *Phys. Lett. B* **329** (1994) 463 [hep-ph/9403201].
37. R. N. Mohapatra, M. K. Parida and G. Rajasekaran, *High scale mixing unification and large neutrino mixing angles*, *Phys. Rev. D* **69** (2004) 053007 [hep-ph/0301234].
38. E. Ma, *Neutrino mass matrix from $S(4)$ symmetry*, *Phys. Lett. B* **632** (2006) 352 [hep-ph/0508231].
39. C. Hagedorn, M. Lindner and R. N. Mohapatra, *$S(4)$ flavor symmetry and fermion masses: Towards a grand unified theory of flavor*, *JHEP* **06** (2006) 042 [hep-ph/0602244].
40. H. Zhang, *Flavor $S(4) \times Z(2)$ symmetry and neutrino mixing*, *Phys. Lett. B* **655** (2007) 132 [hep-ph/0612214].
41. F. Caravaglios and S. Morisi, *Gauge boson families in grand unified theories of fermion masses: $E_6^A \rtimes S_4$* , *Int. J. Mod. Phys. A* **22** (2007) 2469 [hep-ph/0611078].
42. Y. Koide, *$S(4)$ flavor symmetry embedded into $SU(3)$ and lepton masses and mixing*, *JHEP* **08** (2007) 086 [0705.2275].
43. R. Krishnan, P. F. Harrison and W. G. Scott, *Simplest Neutrino Mixing from S_4 Symmetry*, *JHEP* **04** (2013) 087 [1211.2000].
44. R. Krishnan, *A Model for Large Θ_{13} Constructed using the Eigenvectors of the S_4 Rotation Matrices*, *J. Phys. Conf. Ser.* **447** (2013) 012043 [1211.3364].
45. M. Chakraborty, R. Krishnan and A. Ghosal, *Predictive S_4 flavon model with TM_1 mixing and baryogenesis through leptogenesis*, *JHEP* **09** (2020) 025 [2003.00506].
46. G. Altarelli and F. Feruglio, *Tri-bimaximal neutrino mixing, $A(4)$ and the modular symmetry*, *Nucl. Phys. B* **741** (2006) 215 [hep-ph/0512103].
47. S. Zhou, *Lepton Flavor Mixing Pattern and Neutrino Mass Matrix after the Daya Bay Experiment*, **1205.0761**.
48. P. F. Harrison and W. G. Scott, *Symmetries and generalizations of tri - bimaximal neutrino mixing*, *Phys. Lett. B* **535** (2002) 163 [hep-ph/0203209].
49. P. F. Harrison and W. G. Scott, *μ - τ reflection symmetry in lepton mixing and neutrino oscillations*, *Phys. Lett. B* **547** (2002) 219 [hep-ph/0210197].
50. E. Ma, *The All purpose neutrino mass matrix*, *Phys. Rev. D* **66** (2002) 117301 [hep-ph/0207352].
51. K. S. Babu, E. Ma and J. W. F. Valle, *Underlying $A(4)$ symmetry for the neutrino mass matrix and the quark mixing matrix*, *Phys. Lett. B* **552** (2003) 207 [hep-ph/0206292].

-
52. W. Grimus and L. Lavoura, *A Nonstandard CP transformation leading to maximal atmospheric neutrino mixing*, *Phys. Lett. B* **579** (2004) 113 [[hep-ph/0305309](#)].
 53. S. Roy Choudhury and S. Choubey, *Updated Bounds on Sum of Neutrino Masses in Various Cosmological Scenarios*, *JCAP* **09** (2018) 017 [[1806.10832](#)].
 54. A. Loureiro et al., *On The Upper Bound of Neutrino Masses from Combined Cosmological Observations and Particle Physics Experiments*, *Phys. Rev. Lett.* **123** (2019) 081301 [[1811.02578](#)].
 55. EXO-200 collaboration, *Search for Neutrinoless Double- β Decay with the Complete EXO-200 Dataset*, *Phys. Rev. Lett.* **123** (2019) 161802 [[1906.02723](#)].
 56. CUPID collaboration, *Final result of CUPID-0 phase-I in the search for the ^{82}Se Neutrinoless Double- β Decay*, *Phys. Rev. Lett.* **123** (2019) 032501 [[1906.05001](#)].
 57. K. S. Babu and S. Gabriel, *Semidirect Product Groups, Vacuum Alignment and Tribimaximal Neutrino Mixing*, *Phys. Rev. D* **82** (2010) 073014 [[1006.0203](#)].
 58. M. Holthausen and M. A. Schmidt, *Natural Vacuum Alignment from Group Theory: The Minimal Case*, *JHEP* **01** (2012) 126 [[1111.1730](#)].
 59. M. Holthausen, M. Lindner and M. A. Schmidt, *Lepton flavor at the electroweak scale: A complete A_4 model*, *Phys. Rev. D* **87** (2013) 033006 [[1211.5143](#)].
 60. The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.11.1*, 2021.
 61. L. Michel and B. Zhilinskii, *Symmetry, invariants, topology. basic tools*, *Physics Reports* **341** (2001) 11.
 62. L. Michel, *Points critiques des fonctions invariantes sur une g-variete*, *C. R. Acad. Sci. Paris A* **272** (1971) 433.