BLOW UP FOR SMALL-AMPLITUDE SEMILINEAR WAVE EQUATIONS WITH MIXED NONLINEARITIES ON ASYMPTOTICALLY EUCLIDEAN MANIFOLDS

MENGYUN LIU AND CHENGBO WANG

ABSTRACT. In this work, we investigate the problem of finite time blow up as well as the upper bound estimates of lifespan for solutions to small-amplitude semilinear wave equations with mixed nonlinearities $c_1|u_t|^p+c_2|u|^q$, posed on asymptotically Euclidean manifolds, which is related to both the Strauss conjecture and Glassey conjecture. In some cases, we obtain existence results, where the lower bound of the lifespan agrees with the upper bound in order. In addition, our results apply for semilinear damped wave equations, when the coefficient of the dissipation term is integrable (without sign condition) and space-independent.

1. Introduction

Let $(\mathbb{R}^n, \mathfrak{g})$ be a asymptotically Euclidean (Riemannian) manifold, with $n \geq 2$. By asymptotically Euclidean, we mean that $(\mathbb{R}^n, \mathfrak{g})$ is certain perturbation of the Euclidean space $(\mathbb{R}^n, \mathfrak{g}_0)$. More precisely, we assume \mathfrak{g} can be decomposed as

$$\mathfrak{g} = \mathfrak{g}_1 + \mathfrak{g}_2 \;,$$

where \mathfrak{g}_1 is a spherical symmetric, long range perturbation of \mathfrak{g}_0 , and \mathfrak{g}_2 is a short range perturbation. By definition, there exists polar coordinates (r,ω) for $(\mathbb{R}^n,\mathfrak{g}_1)$, in which we can write

(1.2)
$$g_1 = K^2(r)dr^2 + r^2d\omega^2 ,$$

where $d\omega^2$ is the standard metric on the unit sphere \mathbb{S}^{n-1} , and

$$(1.3) |\partial_r^k (K-1)| \leq \langle r \rangle^{-k-\rho}, k = 0, 1, 2,$$

for some given positive constant ρ . Here, $\langle x \rangle = \sqrt{1+|x|^2}$, and we use $A \lesssim B$ to stand for $A \leq CB$ for some constant C > 0. Equipped with the coordinates $x = r\omega$, we have

$$\mathfrak{g} = g_{jk}(x)dx^{j}dx^{k} \equiv \sum_{j,k=1}^{n} g_{jk}(x)dx^{j}dx^{k} , \ \mathfrak{g}_{2} = g_{2,jk}(x)dx^{j}dx^{k} ,$$

where we have used the convention that Latin indices j, k range from 1 to n and the Einstein summation convention for repeated indices. Concerning \mathfrak{g}_2 , we assume

Date: September 19, 2021.

²⁰¹⁰ Mathematics Subject Classification. 58J45, 58J05, 35L71, 35B40, 35B33, 35B44, 35B09, 35L05

Key words and phrases. blow up, Glassey conjecture, Strauss conjecture, lifespan, critical exponent, asymptotically Euclidean manifolds.

The authors were supported by NSFC 11971428.

it satisfies

(1.4)
$$\nabla^{\beta} g_{2,jk} = \mathcal{O}(\langle r \rangle^{-\rho - 1 - |\beta|}), |\beta| \le 2.$$

By these assumptions, it is clear that there exists a constant $\delta_0 \in (0,1)$ such that

(1.5)
$$\delta_0 |\xi|^2 \le g^{jk}(x)\xi_j \xi_k \le \delta_0^{-1} |\xi|^2, \forall x, \xi \in \mathbb{R}^n, K \in (\delta_0, 1/\delta_0),$$

where $(g^{jk}(x))$ denotes the inverse of $(g_{jk}(x))$.

In this paper, we are interested in the investigation of the blow up part of the analogs of the Glassey conjecture and related problems on asymptotically Euclidean manifolds. More precisely, we will study the blow up of solutions for the following semilinear wave equations with small initial data, posed on asymptotically Euclidean manifolds (1.1)-(1.4)

(1.6)
$$\begin{cases} \partial_t^2 u - \Delta_{\mathfrak{g}} u = c_1 |u_t|^p + c_2 |u|^q \\ u(0, x) = \varepsilon u_0(x), u_t(0, x) = \varepsilon u_1(x) \end{cases},$$

where, $\Delta_{\mathfrak{g}} = \nabla^j \partial_j$ is the standard Laplace-Beltrami operator, p, q > 1, $c_1, c_2 \geq 0$ and $\varepsilon > 0$ is a small parameter. Concerning the initial data, we assume

(1.7)
$$0 \le u_0, u_1 \in C_0^{\infty}(\mathbb{R}^n), \quad \text{supp}(u_0, u_1) \subset \{x \in \mathbb{R}^n; |x| \le R_0\} ,$$

for some $R_0 > 0$.

Before stating our results, let us briefly review the history of this problem. When $c_1 = 0 < c_2, \mathfrak{g} = \mathfrak{g}_0$, the problem is related to the Strauss conjecture, where the critical power is given by the Strauss exponent $p_S(n)$, where p_S is the positive root of the equation:

$$(n-1)q^2 - (n+1)q - 2 = 0.$$

When there is no global solutions, and initial data are nontrivial and nonnegative, it has been proved that the upper bound of the lifespan is

(1.8)
$$T_{\varepsilon} \leq S_{\varepsilon} := \begin{cases} C_0 \varepsilon^{\frac{2q(q-1)}{(n-1)q^2 - (n+1)q - 2}} & 1 < q < p_S(n); \\ \exp(C_0 \varepsilon^{-q(q-1)}) & q = p_S(n). \end{cases}$$

In addition, when n = 2, 1 < q < 2 and $u_1 \neq 0$, the upper bound of the lifespan can be improved to

(1.9)
$$T_{\varepsilon} \leq S_{\varepsilon}^{1} = C_{0} \varepsilon^{-\frac{q-1}{3-q}}, \ n = 2, 1 < q < 2,$$

for some constant $C_0 > 0$. We refer [20] for discussion of the history and related results.

When $\mathfrak{g} = \mathfrak{g}_0$ and $c_1 \neq 0 = c_2$, the problem of determining the sharp range of powers of p > 1 for global existence versus blow up for arbitrary small initial data, is known as the Glassey conjecture, where the critical power p for (1.6) is conjectured to be

$$p_G(n) := 1 + \frac{2}{n-1}$$
.

In the important particular case of p=2, the problem with n=3 has been well known to be critical, which admits almost global solutions in general, with the estimates of the lifespan, denoted by T_{ε} , given by $\ln T_{\varepsilon} \sim \varepsilon^{-1}$, see John [13] and John-Klainerman [14]. In general, the global existence with $p > p_G(n)$ is known for n=2,3 through the works of Sideris [24], Hidano-Tsutaya [8] and Tzvetkov [27],

while nonexistence of global solutions for $1 and any <math>n \ge 2$ as well as the upper bound of lifespan,

$$(1.10) T_{\varepsilon} \leq G_{\varepsilon} := \begin{cases} C_0 \varepsilon^{-\frac{2(p-1)}{2-(n-1)(p-1)}}, \ 1$$

has also been well-known (at least for the case $u_1 \neq 0$), see Zhou [34] and references therein. The high dimensional existence part remains open, except for the radial case, see Hidano-Wang-Yokoyama [9]. The upper bound (1.10) is also known to be sharp in general, at least for radial data, see Hidano-Jiang-Wang-Lee [7] and references therein.

When $\mathfrak{g} = \mathfrak{g}_0$ and $c_1c_2 \neq 0$, the problem is related to both the Glassey conjecture and the Strauss conjecture. It turns out that a new critical curve occur in the expected region of global existence $p > p_G(n)$, and $q > p_S(n)$, that is

$$q > p_S, p > p_G, (q-1)((n-1)p-2) = 4$$
,

where the critical and super-critical case is known to admit global existence, at least for n=2,3, see Hidano-Wang-Yokoyama [11], while there is non-existence of global existence for the sub-critical case, as well as an upper bound of the lifespan

(1.11)
$$T_{\varepsilon}^{p,q} \le Z_{\varepsilon} := C_0 \varepsilon^{-\frac{2p(q-1)}{2q+2-(n-1)p(q-1)}}.$$

in the region

$$(1.12) (q-1)((n-1)p-2) < 4, p \le \frac{2n}{n-1}, q < \frac{2n}{n-2},$$

when $u_1 \neq 0$, see Han-Zhou [6]. Note that, in the recent work of Lai-Takamura [18], they remove the restriction for p and obtain the upper bound when $p > \frac{2n}{n-1}$ and $u_1 \neq 0$:

(1.13)
$$T_{\varepsilon}^{p,q} \le C_0 \varepsilon^{-\frac{(q-1)}{q+1-n(q-1)}}, p > \frac{2n}{n-1}, q < \frac{n+1}{n-1}.$$

In fact, in this case, since we have $q < \frac{n+1}{n-1} < p_S$, we could compare the upper bound with S_{ε} in (1.8) and find that (1.13) improves the upper bound exactly when n=2, 1 < q < 2, that is (1.9). Moreover, we find that the additional restriction for q in (1.12), that is, $q < \frac{2n}{n-2}$ is also not necessary, except $q < 1 + \frac{4}{n-3}$ implied by the first inequality, by recasting the proof.

Notice that the results (1.8), (1.9) and (1.10) apply also in the case of $c_1c_2 \neq 0$, by comparing S_{ε} , G_{ε} with Z_{ε} , when $u_1 \neq 0$, we have

(1.14)
$$T_{\varepsilon}^{p,q} \lesssim \begin{cases} S_{\varepsilon}, & q \leq ps, p \geq q \\ Z_{\varepsilon}, & p \leq q \leq 2p-1, (q-1)((n-1)p-2) < 4 \\ G_{\varepsilon}, & p \leq p_{G}, q \geq 2p-1 \\ S_{\varepsilon}^{1}, & q \in (1,2), p \geq 2\frac{q+1}{5-q}, n = 2 \end{cases}$$

See Figure 1 for the illustration of lifespan estimate of $T_{\varepsilon}^{p,q}$ in blows up region. Concerning the sharpness of the upper bound of the lifespan, at least for n=2,3,4, $q,p\geq 2$ and q>2/(n-1), see [11], Wang-Zhou [32] and Dai-Fang-Wang [4].

In recent years, there have been many works concerning the analogs of the problem and related variants on asymptotically flat manifolds, including asymptotically Euclidean manifolds, black hole space-time. In general, we expect that the same




FIGURE 1. Upper bound of the lifespan $T_{\varepsilon}^{p,q}$ when $u_1 \neq 0$

critical power extends to asymptotically flat manifolds, at least when the manifolds enjoy certain geometric structure such as non-trapping.

The existence part of the analogs of the Glassey conjecture for asymptotically Euclidean manifolds and certain small space time perturbation of the flat metric have been partly investigated by Sogge-Wang [25] ($n=3,\ p=p_G=2$) and the second author in [29, 30]. For exterior domain, see Zhou-Han [35] for blow up results and the second author [30] for existence with non-trapping obstacles. For the related works on black hole, Luk [21] obtained the small data global existence on Kerr spacetime, when the nonlinear term satisfies the null condition. Recently, Lai [16] studied the blow up of Glassey conjecture on Schwarzschild spacetime for 1 .

A closely related problem to (1.6) is the following

(1.15)
$$\begin{cases} \partial_t^2 u - \Delta_{\mathfrak{g}} u + b(t) u_t = c_1 |u_t|^p + c_2 |u|^q \\ u(0, x) = \varepsilon u_0(x), u_t(0, x) = \varepsilon u_1(x) \end{cases},$$

where $b \in L^1$. There have been some recent works concerning this problem, with typical damping coefficient $b(t) = \mu(1+t)^{-\beta}$. It turns out that the behavior of (1.15) depends on μ and β . When $\beta > 1$ and $\mathfrak{g} = \mathfrak{g}_0$, it is referred as "scattering" since the linear part of (1.15) behaves like linear wave equations. In this case, when $c_1 > c_2 = 0$, Lai-Takamura [17] proved blow-up results as well as estimates of the lifespan (1.10), when $\mathfrak{g} = \mathfrak{g}_0$, $1 , <math>0 \le b(t) \in L^1$ and $u_1 \ne 0$. Similarly, when $c_1c_2 \ne 0$, Lai-Takamura [18] obtained (1.11) for $\mathfrak{g} = \mathfrak{g}_0$, $u_1 \ne 0$ and typical damping $b(t) = \mu(1+t)^{-\beta}$, $\mu > 0$, $\beta > 1$.

On the other hand, the global existence for $p > p_G$ when n = 3 and $n \ge 4$ with $\mathfrak{g}_2 = 0$ was obtained by Bai-Liu [1] when $\beta > 1$ and $\mu \in \mathbb{R}$, posed on non-trapping asymptotically Euclidean manifolds (1.1)-(1.4). Notice that, the global existence part does not have the nonnegative assumption on b(t), so it is natural to infer that the nonnegative assumption might not be necessary for the blow up results as well.

In our recent work [20], where we consider the finite time blows up of solutions to (1.15) with $c_1 = 0 < c_2$, we have succeeded in proving blow up results without the sign conditions on b, by applying a variable transform. Let

(1.16)
$$m(t) = e^{\int_0^t b(\tau)d\tau}, \quad s = \int_0^t \frac{1}{m(\tau)} d\tau = h(t), t = \eta(s) ,$$

then (1.15) becomes

$$\partial_s^2 u - \tilde{m}^2(s) \Delta_{\mathfrak{g}} u = c_1 \tilde{m}^{2-p}(s) |u_s|^p + c_2 \tilde{m}^2(s) |u|^q, \quad \tilde{m}(s) = m(\eta(s)).$$

Traditionally, we still use t to denote time, and we are reduced to consider the following equation

$$(1.17) \qquad \partial_t^2 u - \tilde{m}^2(t) \Delta_{\mathfrak{g}} u = c_1 \tilde{m}^{2-p}(t) |u_t|^p + c_2 \tilde{m}^2(t) |u|^q, \quad \tilde{m}(t) = m(\eta(t)) .$$

Based on this transformation, as is typical for the test function method of proving blow up, one of the key points is to find appropriate test function adapted for the problem. For that purpose, we state the main assumption that we shall make.

Hypothesis: There exist $\lambda, c > 0$ and corresponding nonnegative nontrivial entire solution ϕ to $\Delta_{\mathfrak{g}} \phi = \lambda^2 \phi$ such that

(H)
$$0 < \phi(x) < c(\lambda r)^{-\frac{n-1}{2}} e^{\lambda \int_0^r K(\tau) d\tau}, \ \phi \neq 0.$$

Let us review some cases where the assumption (H) is valid. It is valid on Euclidean space, $\mathfrak{g} = \mathfrak{g}_0$, where ϕ with $\lambda > 0$ could be given by spherical average of $e^{\lambda x \cdot \omega}$,

$$\phi = \int_{\mathcal{S}^{n-1}} e^{\lambda x \cdot \omega} d\omega ,$$

see Yordanov-Zhang [33]. When \mathfrak{g}_3 is an exponential perturbation, that is, there exists $\alpha > 0$ so that

$$(1.18) |\nabla g_{3,jk}| + |g_{3,jk}| \lesssim e^{-\alpha \int_0^r K(\tau) d\tau},$$

the assumption (H) is recently verified for $\mathfrak{g} = \mathfrak{g}_1 + \mathfrak{g}_3$ by the authors [20] with uniform positive lower bound of ϕ , while the case $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_3$ was obtained by Wakasa and Yordanov in [28].

The main result of this paper then states that there are no global solutions to (1.6) or (1.15), for arbitrary small $\varepsilon > 0$, provided that $p \leq p_G$, or $q \leq p_S$, or

$$(1.19) (q-1)((n-1)p-2) < 4, p > 1, q > 1.$$

More precisely, we have

Theorem 1.1. Let $n \geq 2$, p,q > 1 and $b(t) \in L^1(\mathbb{R}_+)$. Assuming that (H) is valid. Consider (1.15) with $c_1, c_2 \geq 0$ and nontrivial initial data (1.7), posed on asymptotically Euclidean manifolds (1.1)-(1.4). Suppose it has a weak solution $u \in C^2([0,T_{\varepsilon}); \mathcal{D}'(\mathbb{R}^n))$ with $u_t \in L^p_{loc}([0,T_{\varepsilon}) \times \mathbb{R}^n)$, $|u|^q \in C([0,T_{\varepsilon}); \mathcal{D}'(\mathbb{R}^n))$ and

(1.20)
$$\operatorname{supp} u \subset \{(t, x); \int_0^r K(\tau) d\tau \le t + R_1\},$$

for some $R_1 \ge \int_0^{R_0} K(\tau) d\tau$. Then there exist constants $\varepsilon_0 > 0$ and $C_0 > 0$, such that for any $\varepsilon \in (0, \varepsilon_0)$, we have the following results on the upper bound of T_{ε} :

- (1) (1.10), if $c_1 > 0$, 1 ,
- (2) (1.11), if $c_1c_2 > 0$ and (1.19),
- (3) (1.8), if $c_2 > 0$, $1 < q < p_S^1$,
- (4) (1.9) if $c_2 > 0$, n = 2, 1 < q < 2 and $u_1 \neq 0$.

$$c_1 < \phi_{\lambda}(x) < c_1^{-1} \langle \lambda r \rangle^{-\frac{n-1}{2}} e^{\lambda \int_0^r K(\tau) d\tau}$$

for any $0 < \lambda \le \lambda_0$. See [20] for a proof.

¹When $q = p_S$, the situation is more delicate. It can be shown that we still have (1.8), provided that there exist $\lambda_0, c_1 > 0$ such that we have a class of solutions to $\Delta_{\mathfrak{g}} \phi_{\lambda} = \lambda^2 \phi_{\lambda}$ satisfying the uniform bound

Remark 1.1. As we have reviewed, in the previous works, the typical assumptions on the data are (1.7) and $u_1 \neq 0$ for (1.10) and (1.11). Here, we observe that we could actually relax the assumption to any nontrivial initial data with (1.7). It has the obvious benefit that we could then compare upper bounds between S_{ε} , G_{ε} and Z_{ε} .

As is clear, the upper bounds (1.8) and (1.9) could be easily adapted from the corresponding proof for the results for the case $c_1 = 0$, and we refer [20] for the proof.

For the strategy of proof of (1.10) and (1.11), we basically use the test function method, with the help of (H) and the transformation (1.16). We should note that, it seems that the test function $\tilde{\psi}(t,x) = e^{-\lambda \eta(t)}\phi$ used in [20] does not work equally well for the nonlinear term $|u_t|^p$, unless one assume $b(t) \geq 0$. To avoid this difficulty, we work directly on the precise solution of linear ODE

$$\psi_{tt} - \tilde{m}^2 \psi = 0 \ .$$

In this case, we do not have the explicit formula of the solution. Nevertheless, we can obtain the desired asymptotic behavior of the solution by applying the Levinson theorem and ODE arguments, which is still sufficient for our proof.

As we see, our result makes sense under the assumption that there exists a distribution solution that satisfies finite speed of propagation. Once we have the local well-posedness, we can remove this technical assumption. Before concluding the introduction, we discuss some situations where we do have local well-posedness.

At first, the local energy and KSS estimates are available for nontrapping asymptotically Euclidean manifolds, see Bony-Häfner [2] and Sogge-Wang [25]. With help of these estimates, we could prove not only local well-posedness, but also almost global existence, for the problem when n=3, $p=p_G=2$ with $c_2=0$. It has been obtained in [25], [29] for the case b=0. As the argument in [25], [29] could be easily adapted to the current setting, with additional absorption of the term $b(t) \in L^1(\mathbb{R}_+)$ by Gronwall's inequality, similar to that in [19], we omit the proof.

Theorem 1.2. Let n=3 and p=2. Consider (1.15) with $c_2=0$, $b(t) \in L^1(\mathbb{R}_+)$, posed on non-trapping asymptotically Euclidean manifolds (1.1)-(1.4). Assuming that (H) is valid. Then for any nontrivial initial data with (1.7), there is $\varepsilon_0 > 0$ so that for any $\varepsilon \in (0, \varepsilon_0)$, we have a unique local weak solution $(u, u_t) \in C([0, T]; H^3 \times H^2)$. Moreover, we have the following estimate for the lifespan T_{ε}

$$\exp(c_0/\varepsilon) \le T_{\varepsilon} \le \exp(C_0/\varepsilon),$$

for some $0 < c_0 < C_0$.

Recall that as a fundamental tool to prove local well-posedness, Strichartz estimates for wave operators with variable $C^{1,1}$ coefficients have been well-understood, see Smith [23], Tataru [26] and references therein. When $2 \le n \le 4$, it turns out that we could use Strichartz estimates to prove local well-posedness of (1.15) on general Riemannian manifolds, which apply for asymptotically Euclidean manifolds (1.1)-(1.4).

Theorem 1.3. Let $2 \le n \le 4$, $p > \max(1, (n-1)/2)$, $q > \max(1, (n-1)/2, n/2 - 1/(p-1))$. The problem (1.15) with $b(t) \in L^1(\mathbb{R}_+)$ and nontrivial data, posed on C^3 Riemannian $(\mathbb{R}^n, \mathfrak{g})$ with $\nabla^\beta g_{jk} \in L^\infty$, $|\beta| \le 3$, is locally well posed in $H^{s+1} \times H^s$




FIGURE 2. Local well posedness vs blow up, when n = 3, 4

for small enough $\varepsilon > 0$, where

$$(1.21) s \in \left(\max(\frac{n-1}{2}, \frac{n+1}{4}, \frac{n}{2} - \frac{1}{p-1}), \min(2, p, q)\right).$$

In addition, if the initial data satisfies (1.7), $\mathfrak{g} = \mathfrak{g}_1 + \mathfrak{g}_3$ with (1.2)-(1.3) and (1.18), we have upper bounds of the lifespan as in Theorem 1.1.

See Figure 2 for a comparison between region for local well posedness and blow up.

For high dimensional case, as $\min(p,q)$ is close to 1, it seems difficult to employ the approach of using Strichartz estimates obtain desired well-posed results. Instead, in the case of spherical symmetric, small metric perturbation, that is, $\mathfrak{g}_2 = 0$ and there exists $\theta > 0$, such that

(1.22)
$$\sum_{j \in \mathbb{Z}} 2^{jk} \|\partial_r^k (K-1)\|_{L_r^{\infty}(r \sim 2^j)} \le \theta, k = 0, 1, 2 ,$$

it turns out that the homogeneous local energy and KSS-type estimates are available (see Metcalfe-Sogge [22] and Hidano-Wang-Yokoyama [10, Lemma 2.3]). Thus, in spirit of [9] and [7], we could prove well-posed results for the problem with $c_2 = 0$ in the radial case, with sharp lower bound of the lifespan.

Theorem 1.4. Let $n \geq 3$, $1 and <math>\mathfrak{g} = \mathfrak{g}_1$ with (1.22). Consider (1.15) with $b(t) \in L^1(\mathbb{R}_+)$ and $c_2 = 0$. Then there exists $\theta_0 > 0$ such that the following statement holds for $\theta \leq \theta_0$: for any radial, nontrivial initial data satisfying (1.7), there is $\varepsilon_1 > 0$ so that for any $\varepsilon \in (0, \varepsilon_1)$, the problem (1.15) admits a unique local weak solution $(u, u_t) \in C([0, T]; H^2_{rad} \times H^1_{rad})$. Moreover, let T_{ε} be the lifespan of the local solution, we have

(1.23)
$$\begin{cases} c\varepsilon^{-\frac{2(p-1)}{2-(n-1)(p-1)}} \le T_{\varepsilon} \le C\varepsilon^{-\frac{2(p-1)}{2-(n-1)(p-1)}}, \ 1$$

for some 0 < c < C. Here, we use H^s_{rad} to denote the space of spherically symmetric functions in the usual Sobolev space H^s .

Finally, we give the local well posedness for $c_1c_2 \neq 0$ when $n \geq 5$ by exploiting local energy estimates.

Theorem 1.5. Let $n \geq 5$, $1 , <math>1 < q < 1 + \frac{4}{n-4}$. Assuming $\mathfrak{g} = \mathfrak{g}_1$ with (1.22). Consider (1.15) with $b(t) \in L^1(\mathbb{R}_+)$. Then there exists $\theta_1 > 0$ such that the following statement holds for $\theta \leq \theta_1$: for any nontrivial radial initial data




FIGURE 3. Local well posedness vs blow up, when $n \geq 5$

satisfying (1.7), there is $\varepsilon_1 > 0$ so that for any $\varepsilon \in (0, \varepsilon_1)$, the problem (1.15) admits a unique local weak solution $(u, u_t) \in C([0, T]; H^2_{rad} \times H^1_{rad})$. Moreover, we have upper bounds of the lifespan as in Theorem 1.1.

See Figure 3 for a comparison between region for local well posedness and blow up.

This paper is organized as follows. In the next section, we give the existence part of Theorems 1.2-1.5, i.e., local existence and uniqueness of weak solutions, which ensure the property of finite speed of propagation (1.20), instead of the assumption. In Section 3, we give the proof of Theorems 1.1-1.5 with $c_1 > 0$, that is, (1.10) for $1 . In the last Section 4, we present the proof of the blow up part, (1.11), for the case of mixed nonlinearities, with <math>c_1c_2 \ne 0$.

We close this section by listing some notations. For a norm X and a nonnegative integer m, we will use the shorthand notation

$$|\partial^{\leq m} u| = \sum_{|\mu| \leq m} |\partial^{\mu} u|, \quad \|\partial^{\leq m} u\|_X = \sum_{|\mu| \leq m} \|\partial^{\mu} u\|_X.$$

For $1 \leq r \leq \infty$ and $s \in \mathbb{R}$, we denote the norm

$$||u||_{\dot{l}_r^s(X)} = ||(\phi_j(x)u(t,x))||_{\dot{l}_r^s(X)} = ||(2^{js}||\phi_j(x)u(t,x)||_X)||_{\ell^r(j\in\mathbb{Z})},$$

for a partition of unity subordinate to the dyadic spatial annuli, $\sum_{j\in\mathbb{Z}}\phi_j^2(x)=1$, for any $x\neq 0$.

2. Local well-posedness

In this section, we present the proof of local well-posedness for equation (1.15). In the following, we recall the trace estimates we will use later.

Lemma 2.1 (Trace estimates). Let $n \ge 2$, 1/2 < s < n/2. Then

$$\|r^{n/2-s}u\|_{L^{\infty}_{r}L^{2}_{\omega}}\lesssim \|u\|_{\dot{H}^{s}}, \|r^{(n-1)/2}u\|_{L^{\infty}_{r}L^{2}_{\omega}}\lesssim \|u\|_{H^{s}},$$

in particular, for radial functions,

$$\|r^{n/2-s}\langle r\rangle^{s-1/2}u\|_{L^\infty_x}{\lesssim} \|u\|_{H^s_{rad}}\ .$$

See, e.g., Fang-Wang [5] or Wang-Yu [31].

2.1. Strichartz estimates.

Lemma 2.2 (Local homogeneous Strichartz estimates). Let $n \geq 2$, $b \in L^1$, $(\mathbb{R}^n, \mathfrak{g})$ be C^2 Riemannian metric with $\nabla^{\beta} g_{jk} \in L^{\infty}$, $|\beta| \leq 2$. Then for (s, q, r) admissible with $r < \infty$, that is

$$\frac{1}{q} \leq \min\left(\frac{1}{2}, \frac{n-1}{2}\left(\frac{1}{2} - \frac{1}{r}\right)\right), s = n\left(\frac{1}{2} - \frac{1}{r}\right) - \frac{1}{q} ,$$

we have the local homogeneous Strichartz estimates

(2.1)
$$\|\partial D^{-s}u\|_{L^{q}_{t\in[0,1]}L^{r}_{x}} + \|\partial u\|_{L^{\infty}_{t\in[0,1]}L^{2}} \lesssim \|\partial u(0)\|_{L^{2}} ,$$

for solutions to $(\partial_t^2 - \Delta_{\mathfrak{a}} + b(t)\partial_t)u = 0.$

See Smith [23] (n = 2, 3) and Tataru [26, Theorem 1.1] for the case b(t) = 0. The result when $b(t) \in L^1$ follows directly if we combine it with the Duhamel principle and the Gronwall inequality.

Corollary 2.3. Let $2 \le n \le 4$, $(\mathbb{R}^n, \mathfrak{g})$ be C^3 Riemannian metric with $\nabla^{\beta} g_{jk} \in L^{\infty}$, $|\beta| \le 3$, $\max(2, 4/(n-1)) < q < \infty$ and $\tau \in (\frac{n}{2} - \frac{1}{q}, 2]$. Then the solution of equation $(\partial_t^2 - \Delta_{\mathfrak{g}} + b(t)\partial_t)u = F$ satisfies

Proof. We need only to give the proof for F=0, by Duhamel's principle. By applying ∇^{α} , $|\alpha| \leq 2$, to $\partial_t^2 u - \Delta_{\mathfrak{g}} u + b(t) u_t = 0$, we get

$$(\partial_t^2 - \Delta_{\mathfrak{g}} + b(t)\partial_t)\nabla^\alpha u = [-\Delta_{\mathfrak{g}}, \nabla^\alpha]u = \mathcal{O}(\sum_{1 \le |\beta| \le 1 + |\alpha|} |\nabla^\beta u|) \ .$$

Then by (2.1) and Gronwall's inequality, we have

By interpolation, we know that

$$\|\partial D^{\tau-s}u\|_{L^q_{t\in[0,1]}L^r_x}+\|\partial u\|_{L^\infty_{t\in[0,1]}H^\tau}{\lesssim} \|\partial u(0)\|_{H^\tau}\ ,$$

for any $\tau \in [0,2]$. If $\tau - s > \frac{n}{r}$, that is, $\tau > \frac{n}{2} + s = \frac{n}{2} - \frac{1}{q}$, by the Sobolev inequality, we have

$$\|\partial u\|_{L^q_{t\in[0,1]}L^\infty_x} \leq \|\partial D^{\tau-s}u\|_{L^q_{t\in[0,1]}L^r_x} \lesssim \|\partial u(0)\|_{H^\tau} \ ,$$

which completes the proof.

Proposition 2.4. Let $2 \le n \le 4$, $p > \max(1, (n-1)/2)$, $q > \max(1, (n-1)/2, n/2-1/(p-1))$. The problem (1.15) with $b(t) \in L^1(\mathbb{R}_+)$, posed on Riemannian ($\mathbb{R}^n, \mathfrak{g}$) with $\nabla^{\beta}g_{jk} \in L^{\infty}$, $|\beta| \le 3$, is locally well posed in $H^{s+1} \times H^s$, where

(2.4)
$$s \in \left(\max(\frac{n-1}{2}, \frac{n+1}{4}, \frac{n}{2} - \frac{1}{p-1}), \min(2, p, q)\right),$$

for small enough $\varepsilon > 0$.

Proof. For fixed s satisfying (2.4), there is $r > \max(2, 4/(n-1), p-1)$ such that $s \in (n/2 - 1/r, 2)$ so that we could apply (2.2). Let

$$||u||_{X^s} = ||u||_{L^{\infty}_{t\in[0,1]}H^{s+1}} + ||u_t||_{L^{\infty}_{t\in[0,1]}H^s} + ||\partial u||_{L^r_t([0,1])L^{\infty}_x}.$$

For any $u \in X^s$, we define Πu to be the solution to the linear equation

(2.5)
$$\begin{cases} (\partial_t^2 - \Delta_{\mathfrak{g}} + b(t)\partial_t)\Pi u = F(u) := c_1|u_t|^p + c_2|u|^q, \\ \Pi u(0,x) = \varepsilon u_0, \partial_t \Pi u(0,x) = \varepsilon u_1. \end{cases}$$

In view of the Strichartz estimate (2.2), we have

$$\|\Pi u\|_{X^s} \lesssim \varepsilon(\|u_0\|_{H^{s+1}} + \|u_1\|_{H^s}) + \|F(u)\|_{L^1 H^s},$$

$$\|\Pi u - \Pi w\|_{L^{\infty} L^2} \lesssim \|F(u) - F(w)\|_{L^1 L^2}.$$

Then, when ε is small enough, by a standard iteration argument, the proof of local well posedness for $t \in [0,1]$ is reduced to the proof of the following two nonlinear estimates

$$(2.7) \|F(u) - F(w)\|_{L^{1}L^{2}} \lesssim \|(u, w)\|_{X^{s}}^{p-1} \|\partial_{t}u - \partial_{t}w\|_{L^{\infty}L^{2}} + \|(u, w)\|_{X^{s}}^{q-1} \|u - w\|_{L^{\infty}L^{2}}.$$

The second inequality (2.7) follows directly from the Sobolev embedding and Hölder's inequality, as s+1>n/2 and r>p-1. Concerning (2.6), since p,q>s and s+1>n/2, by the Moser type inequality, see, e.g., Hidano-Wang [12, Lemma 2.3], we have

$$||F(u)||_{H^s} \lesssim ||u_t||_{L^{\infty}}^{p-1} ||u_t||_{H^s} + ||u||_{L^{\infty}}^{q-1} ||u||_{H^s} \lesssim ||u_t||_{L^{\infty}}^{p-1} ||u_t||_{H^s} + ||u||_{H^{s+1}}^{q-1} ||u||_{H^s},$$
 and so, as $r > p-1$,

$$||F(u)||_{L^1H^s} \lesssim ||u_t||_{L^rL^\infty}^{p-1} ||u_t||_{L^\infty H^s} + ||u||_{L^\infty H^{s+1}}^{q-1} ||u||_{L^\infty H^s} \lesssim ||u||_{X^s}^p + ||u||_{X^s}^q$$
, which is (2.6). This completes the proof.

2.2. The Glassey conjecture for small, radial asymptotically Euclidean manifolds. As we mentioned before, when $\mathfrak{g} = \mathfrak{g}_0$ and $n \geq 2$, the local well-posedness and long time existence for (1.15) with $b = c_2 = 0$ has been studied in [9] and [7], by exploiting Morawetz type local energy estimates and its variants (see, e.g., [15], [22], [10]), together with the trace estimates. The similar approach could apply in the setting of small, radial asymptotically Euclidean manifolds, at least when $n \geq 3$.

In practice, the following estimates are useful in proving well-posedness for (1.15),

$$(2.8) \ T^{\frac{\mu-1}{2}} \| r^{-\frac{\mu}{2}} \tilde{\partial} u \|_{L^2_{T,x}} + (\ln(T+2))^{-\frac{1}{2}} \| r^{-\frac{\mu}{2}} \langle r \rangle^{-\frac{1-\mu}{2}} \tilde{\partial} u \|_{L^2_{T,x}} \lesssim \| u \|_{X_T} \ , \mu \in (0,1) \ ,$$

where, we denote L_T^q as $L_t^q([0,T])$, $\tilde{\partial}u = (\partial u, u/r)$ and

$$||u||_{X_T} := ||\tilde{\partial} u||_{\dot{\ell}_{\infty}^{-\frac{1}{2}} L_{t,x}^2 \cap L_t^{\infty} L_x^2([0,T] \times \mathbb{R}^n)}, ||F||_{N_T} := ||F||_{\dot{\ell}_1^{\frac{1}{2}} L_{t,x}^2 + L_t^1 L_x^2([0,T] \times \mathbb{R}^n)}.$$

See, e.g., [10], for a proof. Concerning X_T and N_T , we have the following version of the local energy estimates.

Lemma 2.5. Let $n \geq 3$, $b \in L^1$, $(\mathbb{R}^n, \mathfrak{g})$ with $\mathfrak{g} = \mathfrak{g}_1$ satisfying (1.22) and θ small enough. Then there exists a constant C > 0 such that

$$||u||_{X_T} \le C||\partial u(0)||_{L^2} + C||F||_{N_T} ,$$

for any solutions to $(\partial_t^2 - \Delta_{\mathfrak{g}} + b(t)\partial_t)u = F$ in $[0,T] \times \mathbb{R}^n$.

Proof. Recall that, when $\mathfrak{g} = \mathfrak{g}_1$ satisfies (1.22) for k = 0, 1 with θ small enough, the inequality (2.9) has essentially been proved in Metcalfe-Sogge [22] and Hidano-Wang-Yokoyama [10, Lemma 2.3] for b = 0, while the general case with $b \in L^1$ follows from the Gronwall inequality.

Since

$$(\partial_t^2 - \Delta_{\mathfrak{g}} + b(t)\partial_t)\nabla u = \nabla F + [\nabla, \Delta_{\mathfrak{g}}]u = \nabla F + \mathcal{O}(|(\nabla^2 g)(\nabla u)| + |(\nabla g)(\nabla^2 u)|)$$
, by (2.9), we have

$$\begin{split} \|\nabla u\|_{X_T} \lesssim &\|\partial u(0)\|_{\dot{H}^1} + \|\nabla F\|_{N_T} + \|\nabla^2 g \nabla u\|_{\dot{\ell}_1^{\frac{1}{2}}L_{t,x}^2} + \|\nabla g \nabla^2 u\|_{\dot{\ell}_1^{\frac{1}{2}}L_{t,x}^2} \\ \lesssim &\|\partial u(0)\|_{\dot{H}^1} + \|\nabla F\|_{N_T} + \|\tilde{\partial} \nabla g\|_{\dot{\ell}_1^{2}L_{\infty}^{\infty}} \|\nabla u\|_{X_T}. \end{split}$$

As
$$\|\tilde{\partial}\nabla g\|_{\dot{\ell}_1^2 L_{\infty}^{\infty}} \ll 1$$
 by assumption (1.22), we obtain (2.10).

Proposition 2.6. Let $n \geq 3$ and $1 . Consider (1.15) on asymtotically Euclidean manifolds <math>\mathfrak{g}$ with $\mathfrak{g} = \mathfrak{g}_1$ satisfying (1.22) and θ small enough. Then (1.15) with $c_2 = 0$ is locally well-posed in $H^2_{rad} \times H^1_{rad}$ for small enough ε . Moreover, there exists c > 0 such that, for any initial data $(u_0, u_1) \in H^2_{rad} \times H^1_{rad}$ with

$$||u_0||_{H^2} + ||u_1||_{H^1} \le 1,$$

there exists a unique solution $u \in C([0,T_{\varepsilon}];H^2_{rad}) \cap C^1([0,T_{\varepsilon}];H^1_{rad})$ with

(2.11)
$$T_{\varepsilon} \ge \begin{cases} c\varepsilon^{-\frac{2(p-1)}{2-(p-1)(n-1)}}, \ 1$$

Proof. We give the proof using the similar approach in Proposition 2.4. For the operator Π defined in (2.5), in view of the linear estimates in Lemma 2.5, we have

$$\|\nabla^{\leq 1}\Pi u\|_{X_T} \leq C\varepsilon + C\|\nabla^{\leq 1}F(u)\|_{N_T} ,$$

$$\|\Pi u - \Pi w\|_{X_T} \le C \|F(u) - F(w)\|_{N_T}$$
.

Then we need only to prove the following nonlinear estimates for radial functions u, w,

$$\|\nabla^{\leq 1}|u_t|^p\|_{N_T} \lesssim A(T) \|\nabla^{\leq 1}u\|_{X_T}^p \ ,$$

$$(2.13) ||u_t|^p - |w_t|^p||_{N_T} \lesssim A(T) ||\nabla^{\leq 1}(u, w)||_{X_T}^{p-1} ||u - w||_{X_T} ,$$

where

$$A(T) = \left\{ \begin{array}{ll} T^{1-\mu} & 1$$

Notice that the lower bound of T_{ε} is obtained from the requirement $A(T_{\varepsilon})\varepsilon^{p-1} \ll 1$, in the process of closing the iteration.

These nonlinear estimates are well-known, see, e.g., [9] and [7]. For reader's convenience, we give a proof of (2.12) here. By Lemma 2.1, we have

$$||r^{(n-1)/2}u_t||_{L^{\infty}} \lesssim ||r^{(n-2)/2}\langle r\rangle^{1/2}u_t||_{L^{\infty}} \lesssim ||u_t||_{H^1}$$
.

When $p \in (1, p_G)$, we have $\mu/(p-1) = (n-1)/2$ and so by (2.8), we see that $\|\nabla^{\leq 1}|u_t|^p\|_{N_T}$ is controlled by

$$T^{(1-\mu)/2} \| r^{\mu/2} \nabla^{\leq 1} |u_t|^p \|_{L^2_{T,x}} \lesssim T^{1-\mu} \| r^{\mu/(p-1)} u_t \|_{L^\infty}^{p-1} \| \nabla^{\leq 1} u \|_{X_T} \lesssim A(T) \| \nabla^{\leq 1} u \|_{X_T}^p .$$

When $p = p_G$, we have $\mu/(p-1) = (n-2)/2$ and then

$$\begin{split} \|\nabla^{\leq 1} |u_t|^p \|_{N_T} & \lesssim (\ln(2+T))^{1/2} \|r^{\mu/2} \langle r \rangle^{(1-\mu)/2} \nabla^{\leq 1} |u_t|^p \|_{L^2_{T,x}} \\ & \lesssim (\ln(2+T)) \|r^\mu \langle r \rangle^{1-\mu} |u_t|^{p-1} \|_{L^\infty_{T,x}} \|\nabla^{\leq 1} u\|_{X_T} \\ & \lesssim A(T) \|\nabla^{\leq 1} u\|_{X_T}^p \ . \end{split}$$

This completes the proof.

2.3. Local well-posedness for $c_1c_2 \neq 0$ in high dimension. When $n \geq 5$, it seems that the approach of using Strichartz estimate does not apply for small p, q. However, the local energy estimates in Lemma 2.5 works well for some powers. In summary, we can show (1.15) is locally well posed, when

(2.14)
$$1$$

Proposition 2.7. Let $n \geq 5$ and (p,q) in region (2.14) and $c_1c_2 \neq 0$. Consider (1.15) on asymtotically Euclidean manifolds \mathfrak{g} with $\mathfrak{g} = \mathfrak{g}_1$ satisfies (1.22) and θ small enough. Then (1.15) is locally well-posed in $H^2_{rad} \times H^1_{rad}$ for small enough ε .

Proof. The proof follows the similar way in Proposition 2.6, while for $p_G , we shall make a slightly modification of <math>\mu$, that is, we set $\mu = \frac{(n-2)(p-1)}{2} \in (0,1)$ and in this case, we see that

$$\begin{split} \|\nabla^{\leq 1}|u_t|^p\|_{N_T} \lesssim & T^{(1-\mu)/2} \|r^{\mu/2}\nabla^{\leq 1}|u_t|^p\|_{L^2_{T,x}} \\ \lesssim & T^{1-\mu} \|r^{(n-2)/2}u_t\|_{L^\infty}^{p-1} \|\nabla^{\leq 1}u\|_{X_T} \lesssim & T^{1-\mu} \|\nabla^{\leq 1}u\|_{X_T}^p . \end{split}$$

So we need only to show how to control the additional norm of $|u|^q$. With $\|\nabla^{\leq 1}u\|_{X_T}\lesssim \varepsilon$, we claim that

for some small T and ε .

On the one hand, by Hölder's inequality, we have

$$\|\nabla^{\leq 1}|u|^q\|_{N_T} \leq \|\nabla^{\leq 1}|u|^q\|_{L^1_TL^2_x} \lesssim T(\|u\|_{L^\infty_TL^{2q}}^q + \||u|^{q-1}\nabla u\|_{L^\infty_TL^2}) \ .$$

Notice that $\frac{n}{2} - \frac{1}{q-1}$, $\frac{n}{2} - \frac{n}{2q} \le 2$ when $1 < q \le 1 + \frac{2}{n-4}$, then by Sobolev embedding, we get

$$||u||_{L_x^{2q}} \lesssim ||u||_{\dot{H}^{\frac{n}{2} - \frac{n}{2q}}} \lesssim ||u||_{H^2},$$

$$\||u|^{q-1}\nabla u\|_{L^2_x} \lesssim \|\nabla u\|_{L^{\frac{2n}{n-2}}} \|u\|_{L^{n(q-1)}}^{q-1} \lesssim \|u\|_{\dot{H}^2} \|u\|_{\dot{H}^{\frac{n}{2}}}^{q-1} \lesssim \|u\|_{H^2}^q \ .$$

For the L^2 norm of u, by Newton-Leibniz formula, we have

$$||u||_{L^{\infty}_T L^2_x} \le T||u_t||_{L^{\infty}_T L^2_x} + ||u_0||_{L^2} \le T||u||_{X_T} + \varepsilon .$$

On the other hand, we also have

$$\|\nabla^{\leq 1}|u|^q\|_{N_T} \lesssim T^{\frac{1-\mu}{2}} \|r^{\frac{\mu}{2}}|u|^{q-1} \nabla^{\leq 1} u\|_{L^2_{T,x}}.$$

By Hölder's inequality, we get

$$\|r^{\frac{\mu}{2}}|u|^{q-1}\nabla^{\leq 1}u\|_{L^2_{T,x}}\lesssim \|r^{\frac{\mu+1}{q-1}}u\|_{L^\infty_{t,x}}^{q-1}\|r^{-1-\frac{\mu}{2}}\nabla^{\leq 1}u\|_{L^2_{T,x}}\;.$$

Note that $\frac{n}{2} - \frac{\mu+1}{q-1} \in (\frac{1}{2}, 2]$ if $1 + \frac{2}{n-1} < q < 1 + \frac{4}{n-4}$, then by applying trace estimates, we obtain that

$$||r^{\frac{\mu+1}{q-1}}u||_{L_x^{\infty}} \lesssim ||u||_{\dot{H}^{\frac{n}{2}-\frac{\mu+2}{q-1}}} \lesssim ||u||_{H^2}$$
.

Thus we have

$$\|\nabla^{\leq 1}|u|^q\|_{N_T} \lesssim T^{1-\mu}\|u\|_{L_T^\infty H^2}^{q-1} \varepsilon \lesssim \varepsilon^q \ll \varepsilon ,$$

if we take T small enough.

3. Finite time blow up of (1.15) with $c_1 > 0$

In this section we give the proof of Theorem 1.1, in the case $c_1 > 0$ with estimates (1.10), under the assumption there is a local solution $u \in C^2([0, T_{\varepsilon}); \mathcal{D}'(\mathbb{R}^n))$ with $u_t \in L^p_{loc,t,x}([0, T_{\varepsilon}) \times \mathbb{R}^n)$ and $|u|^q \in C([0, T_{\varepsilon}); \mathcal{D}'(\mathbb{R}^n))$ satisfies finite speed of propagation. And based on the local well-posedness in Section 2, we are ready to prove the finite time blows up in Theorem 1.2-Theorem 1.5 without these assumption.

By the transformation (1.16), for future reference, we record that

(3.1)
$$\eta'(t) = m(\eta(t)) = \tilde{m}(t), \ \tilde{m}'(t) = b(\eta(t))\tilde{m}^2(t), \ \eta(0) = 0, m(0) = \tilde{m}(0) = 1$$
.

3.1. Elliptic equation $\Delta_{\mathfrak{g}}\phi = \lambda^2\phi$. As we discussed in introduction, in our setting, (H) is assumed to be true for some fixed $\lambda > 0$ when $\mathfrak{g} = \mathfrak{g}_1 + \mathfrak{g}_2$ and we have (H) when $\mathfrak{g} = \mathfrak{g}_1 + \mathfrak{g}_3$. Then we will exploit it to construct the solution of linear wave equation

$$\partial_t^2 u - \tilde{m}^2 \Delta_{\mathfrak{g}} u = 0 .$$

3.2. **Test function** $\psi(t,x)$. As usual, we want to find a nonnegative solution $\psi(t,x)$ for (3.2) by methods of separation of variables. Let $\psi(t,x) = \varphi(t)\phi(x)$. Then we are reduced to consider the following ordinary differential equation:

(3.3)
$$\varphi'' - \lambda^2 \tilde{m}^2(t) \varphi = 0.$$

By applying Levinson theorem (see, e.g., [3, Chapter 3, Theorem 8.1]), there exist a T > 0 such that

$$\varphi \simeq e^{-\lambda \int_{t_0}^t \tilde{m}(\tau) d\tau} \simeq e^{-\lambda \eta(t)} , \forall t \geq T ,$$

(3.5)
$$\varphi' \simeq -\lambda k e^{-\lambda \int_{t_0}^t \tilde{m}(\tau) d\tau} \simeq -\lambda k e^{-\lambda \eta(t)}, \forall t \geq T,$$

where $k = \tilde{m}(\infty) = \exp(\int_0^\infty b(t)dt)$. In fact, let $Y(t) = (Y_1(t), Y_2(t))^T$ and $Y_1 = \varphi$, $Y_2 = \varphi'$, then we have

$$Y'=(A+V(t))Y, A=\left(\begin{array}{cc} 0 & 1 \\ \lambda^2 k^2 & 0 \end{array}\right), V(t)=\left(\begin{array}{cc} 0 & 0 \\ \lambda^2(\tilde{m}^2(t)-k^2) & 0 \end{array}\right) \; .$$

Noticing that $V' \in L^1$ and $\lim_{t\to\infty} V(t) = 0$, we could apply the Levinson theorem to the system. Then there exists $t_0 \in [0, \infty)$ so that we have a solution, which have the asymptotic form when $t \geq T$,

(3.6)
$$Y(t) = \begin{pmatrix} 1 + o(1) \\ -\lambda k + o(1) \end{pmatrix} e^{-\lambda \int_{t_0}^t \tilde{m}(\tau)d\tau},$$

which gives us (3.4)-(3.5).

Lemma 3.1. Let $\nu(t) = -\varphi'/\varphi$, then there exists $\delta_2 \in (0,1)$, such that for any $t \geq 0$, we have

$$(3.7) \lambda \delta_2 \le \nu(t) \le \lambda/\delta_2 .$$

Proof. By (3.6), we have

$$\nu(t) = -\frac{\varphi'}{\varphi} \to \lambda k, \ t \to \infty,$$

and there exists a N > T, such that

$$\nu(t) \in \left[\frac{\lambda k}{2}, 2\lambda k\right], \ t \ge N$$
.

For the remaining interval [0, N], with data $\varphi(N) > 0$, $\varphi'(N) < 0$ at t = N, we know from the equation (3.3) that

$$\varphi'(t) \le \varphi'(N) < 0, \ \varphi(t) \ge \varphi(N) > 0, \ \forall t \in [0, N]$$
.

Since $\nu \in C^1[0, N]$, it tells us that $\nu(t) > 0$ and so $\nu(t) \sim 1$ on the compact interval [0, N]. This completes the proof.

3.3. Finite speed of propagation. For linear wave equation $\partial_t^2 u - \Delta_{\mathfrak{g}_1} u + b(t) u_t = 0$ with the initial data (1.7) the support of solution u is

(3.8)
$$\sup u \subset \{(t,x); \int_0^{|x|} K(\tau)d\tau \le t + R_1\}, R_1 \ge \int_0^{R_0} K(\tau)d\tau .$$

As \mathfrak{g}_2 is short-range perturbation, which does not affect speed of propagation too much, we still have (3.8), with possibly bigger R_1 to (1.15). Thus with the change of variable $t \to \eta(t)$, the support of solution u of (1.17) is

(3.9)
$$\operatorname{supp} u \subset \{(t, x); \int_0^{|x|} K(\tau) d\tau \leq \eta(t) + R_1\} = D.$$

Recall that $m(t) \in [\delta_1, 1/\delta_1]$, by (3.1) we have

(3.10)
$$D \subset \{(t,x); |x| \le \frac{t}{\delta_0 \delta_1} + \frac{R_1}{\delta_0}\} = \tilde{D} .$$

Lemma 3.2. Let $q \ge 1$, then we have

$$\int_D \psi^q dv_{\mathfrak{g}} \lesssim (t+1)^{n-1-\frac{n-1}{2}q} .$$

Proof. We divide the region D into two disjoint parts: $D = D_1 \cup D_2$ where

$$D_1 = \{(t, x); \int_0^{|x|} K(\tau) d\tau \le \frac{\eta(t) + R_1}{2} \}.$$

For the region D_1 , by (\mathbb{H}), we have

$$\int_{D_1} \psi^q dv_{\mathfrak{g}} \lesssim e^{-\lambda \eta(t)q} \int_{D_1} (1+\lambda|x|)^{-\frac{n-1}{2}q} e^{q\lambda \int_0^{|x|} K(\tau) d\tau} dv_{\mathfrak{g}} \ .$$

Let $\tilde{r} = \int_0^{|x|} K(\tau) d\tau$, then $d\tilde{r} = K(r) dr$ and $\delta_0 r \leq \tilde{r} \leq r/\delta_0$ since $K \in [\delta_0, 1/\delta_0]$. Then we get

$$\begin{split} \int_{D_1} \psi^q dv_{\mathfrak{g}} \lesssim & e^{-\lambda \eta(t)q} \int_0^{\frac{\eta(t)+R_1}{2}} (1+\tilde{r})^{n-1-\frac{n-1}{2}q} e^{q\lambda \tilde{r}} d\tilde{r} \\ \lesssim & e^{-\lambda \eta(t)q} \int_0^{\frac{\eta(t)+R_1}{2}} e^{\frac{3}{2}q\lambda \tilde{r}} d\tilde{r} \\ \lesssim & e^{-\frac{\lambda}{4}\eta(t)} \lesssim & (1+t)^{n-1-\frac{n-1}{2}q}, \end{split}$$

where we have used the fact that e^{-t} decays faster than any polynomial. For the region D_2 , it is easy to see

$$\int_{D_2} \psi^q dv_{\mathfrak{g}} \lesssim e^{-\lambda \eta(t)q} \int_{\frac{\eta(t)+R_1}{2}}^{\eta(t)+R_1} e^{q\lambda \tilde{r}} (1+\tilde{r})^{n-1-\frac{n-1}{2}q} d\tilde{r}$$

$$\lesssim (1+\eta(t))^{n-1-\frac{n-1}{2}q} \int_{\frac{\eta(t)+R_1}{2}}^{\eta(t)+R_1} e^{q\lambda \tilde{r}-\lambda \eta(t)q} d\tilde{r}$$

$$\lesssim (1+t)^{n-1-\frac{n-1}{2}q} ,$$

which completes the proof.

3.4. **Proof of** (1.10). For simplicity, we first define some auxiliary functions

$$F(t) = \int_{\mathbb{R}^n} u_t(t, x) \psi(t, x) dv_{\mathfrak{g}} ,$$

$$G(t) = \int_{\mathbb{R}^n} u(t, x) \psi(t, x) dv_{\mathfrak{g}} ,$$

$$H(t) = c_1 \int_{\mathbb{R}^n} \tilde{m}^{2-p}(t) |u_t|^p \psi(t, x) dv_{\mathfrak{g}} ,$$

which are well-defind in the setting of Theorem 1.2-1.5. In the setting of Theorem 1.1, for fixed t, u, u_t , $|u_t|^p$, $|u|^q$ could be viewed as distribution with compact support, thanks to the finite speed of propagation. As a consequence, we see that $F, G \in C^1([0, T_{\varepsilon})$. Since $c_1, c_2 \geq 0$, both $c_1\tilde{m}^{2-p}(t)|u_t|^p$ and $c_2\tilde{m}^2(t)|u|^q$ are nonnegative distribution with compact support, which particularly gives us $H(t) \geq 0$, as $\psi \geq 0$.

Using $\psi, \psi_t \in C^{\infty}$ as the test functions, we know that $\langle u_t, \psi \rangle - \langle u, \psi_t \rangle \in C^1$ and we get from (1.17), $\psi \geq 0$ and $\partial_t^2 \psi = \Delta_{\mathfrak{g}} \psi$ that

$$\frac{d}{dt} \int (u_t \psi - u \psi_t) dv_{\mathfrak{g}} = \int (u_{tt} \psi - u \psi_{tt}) dv_{\mathfrak{g}}$$

$$= \int (u_{tt} \psi - \tilde{m}^2 u \Delta_{\mathfrak{g}} \psi) dv_{\mathfrak{g}} = \int (u_{tt} - \tilde{m}^2 \Delta_{\mathfrak{g}} u) \psi dv_{\mathfrak{g}}$$

$$\geq c_1 \int \tilde{m}^{2-p}(t) |u_t|^p \psi dv_{\mathfrak{g}} ,$$

and so

$$(3.11) (F + \nu G)' \ge H.$$

Then we get

(3.12)
$$F + \nu G \ge F(0) + \nu(0)G(0) + \int_0^t H(\tau)d\tau \ge 0,$$

thanks to the assumption (1.7) on the initial data. Notice that

$$(\nu G)' = -\frac{d}{dt} \int u\psi_t dv_{\mathfrak{g}} = -\left(\int u_t \psi_t dv_{\mathfrak{g}} + \int u\psi_{tt} dv_{\mathfrak{g}}\right)$$
$$= \nu \int u_t \psi dv_{\mathfrak{g}} - \lambda^2 \tilde{m}^2 \int u\psi dv_{\mathfrak{g}}$$
$$= \nu F - \lambda^2 \tilde{m}^2 G.$$

We get also from (3.11) that

$$(3.13) F' + \nu F - \lambda^2 \tilde{m}^2 G \ge H .$$

Based on (3.12) and (3.13), it is easy to see that

$$F(t) \ge 0$$
, $G(t) \ge 0$, $\forall t \ge 0$.

Actually, recall that $G' = \int u_t \psi + u \psi_t dv_{\mathfrak{g}} = F - \nu G$, we have

$$G' + 2\nu G = F + \nu G \ge 0$$

due to (3.12), which gives us $G \ge e^{-\int_0^t 2\nu d\tau} G(0) \ge 0$. Then, by (3.13), we see that $F' + \nu F \ge 0$, which yields $F \ge 0$.

Next, by multiplying $B_1 = \lambda \delta_1^2 \delta_2$ to (3.12) and adding it with (3.13), we obtain

$$F' + (\nu + B_1)F \ge (\lambda^2 \tilde{m}^2 - B_1 \nu)G + B_1 F(0) + B_1 \nu(0)G(0) + B_1 \int_0^t H d\tau + H$$
.

Note that $\nu \leq \lambda/\delta_2$, $\tilde{m} \geq \delta_1$ then $\lambda^2 \tilde{m}^2 - B_1 \nu \geq 0$ and

(3.14)
$$F' + (\frac{\lambda}{\delta_2} + B_1)F \ge B_1 F(0) + B_1 \nu(0) G(0) + B_1 \int_0^t H d\tau + H.$$

Let $B_2 = B_1/((\lambda/\delta_2) + B_1) < 1$, L(t) := F(t) - I(t) with increasing function

(3.15)
$$I(t) := B_2 \int_0^t H d\tau + B_2 F(0) + \frac{B_2}{2} \nu(0) G(0) ,$$

then $L' = F' - B_2 H$, $L(0) = (1 - B_2) F(0) - \frac{B_2}{2} \nu(0) G(0) \ge -\frac{B_2}{2} \nu(0) G(0)$ and

$$L' + \frac{B_1}{B_2}L = F' + \frac{B_1}{B_2}F - B_2H - B_1 \int_0^t Hd\tau - B_1F(0) - \frac{B_1}{2}\nu(0)G(0)$$
$$\geq (1 - B_2)H + \frac{B_1}{2}\nu(0)G(0) \geq \frac{B_1}{2}\nu(0)G(0) ,$$

thanks to (3.14). Thus we have

$$L(t) \ge e^{-\frac{B_1}{B_2}t} L(0) + e^{-\frac{B_1}{B_2}t} \frac{B_1}{2} \nu(0) G(0) \int_0^t e^{\frac{B_1}{B_2}\tau} d\tau$$

$$\ge \frac{B_2}{2} \nu(0) G(0) \left(1 - 2e^{-\frac{B_1}{B_2}t}\right) \ge 0,$$

when $t \geq t_0 = \frac{B_2}{B_1} \ln 2$, which yields

$$I(t) \le F(t) = \int u_t \psi dv_{\mathfrak{g}} , \ \forall t \ge t_0 .$$

By Hölder's inequality and Lemma 3.2 with q = 1, we get

$$I(t) \lesssim \left(\int |u_t|^p \psi dv_{\mathfrak{g}}\right)^{1/p} \left(\int_D \psi dv_{\mathfrak{g}}\right)^{1/p'} \lesssim H^{1/p} (t+1)^{\frac{n-1}{2p'}}, \ \forall t \geq t_0 \ ,$$

which gives us the following ordinary differential inequality for I(t)

(3.16)
$$I'(t) = B_2 H(t) \gtrsim \frac{I^p}{(1+t)^{\frac{n-1}{2}(p-1)}}, \ \forall t \ge t_0 \ .$$

On the other hand, as $I'(t) = B_2H(t) \ge 0$ and the initial data (u_0, u_1) are nontrivial with (1.7), we know that

(3.17)
$$I(t_0) \ge I(0) = B_2 \varepsilon \int u_1 \phi dv_{\mathfrak{g}} + \frac{B_2}{2} \nu(0) \varepsilon \int u_0 \phi dv_{\mathfrak{g}} \sim \varepsilon.$$

With help of (3.16) and (3.17), it is easy to conclude the proof of Theorem 1.1 for the case $c_1 > 0$ with estimates (1.10). Actually, when $\frac{n-1}{2}(p-1) = 1$, that is, $p = p_G$, we see that for some positive constants C, ε_0 , such that for any $\varepsilon \in (0, \varepsilon_0)$, we have

$$I(t) \ge \left(C\varepsilon^{1-p} - (p-1)\ln(1+t)\right)^{-1/(p-1)},$$

which will go to ∞ as $C\varepsilon^{1-p} - (p-1)\ln(1+t) \to 0$. Thus we have the existence time T_{ε} of I(t) satisfies

$$T_{\varepsilon} \leq \exp(\tilde{C}\varepsilon^{-(p-1)})$$
,

for some $\tilde{C} > 0$. Similarly, when $\frac{n-1}{2}(p-1) < 1$, that is, $p < p_G$, we get

$$I(t) \ge \left(C\varepsilon^{1-p} - \frac{p-1}{1-(n-1)(p-1)/2}(1+t)^{1-(n-1)(p-1)/2}\right)^{-1/(p-1)}$$
.

Hence the existence time T_{ε} satisfies

$$T_{\varepsilon} \le C_0 \varepsilon^{-\frac{2(p-1)}{2-(n-1)(p-1)}},$$

for some $C_0 > 0$ and small enough ε .

4. Finite time blow up of (1.15) with $c_1c_2 \neq 0$

In this section, we present a proof of (1.11), for (1.15) with $c_1c_2 \neq 0$. For that purpose, we introduce the function

$$F(t) = \int u(t, x) dv_{\mathfrak{g}} ,$$

which, as we shall see, will blow up in finite time.

In view of (1.17), we have

(4.1)
$$F'' = \int c_1 \tilde{m}^{2-p}(t) |u_t|^p + c_2 \tilde{m}^2(t) |u|^q dv_{\mathfrak{g}} \gtrsim \int |u|^q dv_{\mathfrak{g}}.$$

By Hölder's inequality, we have

$$|F| \le \int_{\tilde{D}} |u| dv_{\mathfrak{g}} \lesssim \left(\int_{\tilde{D}} |u|^q dv_{\mathfrak{g}}\right)^{1/q} (1+t)^{n/q'},$$

which, combined with (4.1), gives us

$$F'' \ge |F|^q (1+t)^{-n(q-1)}$$
.

Since $F'' \ge 0$, we have $F'(t) \ge F'(0) \ge 0$ and $F(t) \ge F(0) + tF'(0) \ge 0$. Then we get

$$(4.2) F'' \gtrsim |F|^q (1+t)^{-n(q-1)}, F(t) \geq F(0) + tF'(0) \geq 0, \ \forall t \geq 0.$$

To show blow up, we need to get better lower bound of F, for which purpose, we use the test function

$$\tilde{\psi}(t,x) = e^{-\lambda \eta(t)} \phi(x)$$

with ϕ satisfying (H), and introduce the auxiliary function

$$G(t) = \int u_t \tilde{\psi} dv_{\mathfrak{g}} .$$

Without loss of generality, we suppose $T_{\varepsilon} > 1$. Then we claim that

(4.3)
$$G(t) \ge C\varepsilon, \forall t \in [1, T_{\varepsilon}),$$

for some constant C > 0. With (4.3) in hand, we see that by Hölder's inequality

$$(4.4) C\varepsilon \leq G(t) \lesssim \left(\int_{D} |u_{t}|^{p} dv_{\mathfrak{g}}\right)^{1/p} \left(\int_{D} \tilde{\psi}^{p'} dv_{\mathfrak{g}}\right)^{1/p'},$$

Recall that by Lemma 3.2, we have

$$\int_D \tilde{\psi}^{p'} dv_{\mathfrak{g}} \lesssim (1+t)^{n-1-\frac{n-1}{2}p'},$$

which, recalling (4.1), yields

(4.5)
$$F'' \gtrsim \int |u_t|^p dv_{\mathfrak{g}} \gtrsim \varepsilon^p (1+t)^{-(n-1)(p-2)/2}, t \ge 1.$$

Notice that, if $(n-1)(p-2)/2 \le 1$, that is $p \le \frac{2n}{n-1}$ in the condition (1.12), then heuristically, we can integrate the above expression twice and obtain

(4.6)
$$F \gtrsim \varepsilon^p (1+t)^{2-(n-1)(p-2)/2}, t \ge 1,$$

which could be improved with the help of (4.2), if

$$\left(2 - \frac{(n-1)(p-2)}{2}\right)q - n(q-1) + 2 > 2 - \frac{(n-1)(p-2)}{2}$$
.

This is exactly the first condition of (1.12) on p,q. Then the classical Kato type lemma (see, e.g., [6, Lemma 2.1]) could be applied to prove blow up results and the desired upper bound of lifespan (1.11).

It remains to prove the claim (4.3). Let $H(t) = \int u(t,x)\phi(x)dv_{\mathfrak{q}}$, then

(4.7)
$$G(t) = e^{-\lambda \eta(t)} H'(t),$$

and H satisfies

$$H'' = \int u_{tt} \phi dv_{\mathfrak{g}} \ge \tilde{m}^2 \int \Delta_{\mathfrak{g}} u \phi dv_{\mathfrak{g}} \ge \tilde{m}^2 \lambda^2 H .$$

Let y be the solution of ordinary differential equation

$$y'' - \lambda^2 \tilde{m}^2 y = 0$$
, $y(0) = H(0)/\varepsilon = C_0$, $y'(0) = H'(0)/\varepsilon = C_1$,

for some $C_0, C_1 \ge 0$ with $C_0 + C_1 > 0$. We observe that $H' \ge \varepsilon y'$. For y, it is clear from continuity that we have y, y' > 0 for all t > 0. More precisely, we have $y(t) \geq C_0$, $y'' \geq \delta_1^2 \lambda^2 C_0$ for all $t \geq 0$, and so

$$(4.8) y' \ge C_1 + t\delta_1^2 \lambda^2 C_0 > 0 , \forall t > 0 .$$

Let $Y(t) = (Y_1(t), Y_2(t))^T$ and $Y_1 = y, Y_2 = y'$, then we have

$$Y' = (A + V(t))Y, A = \begin{pmatrix} 0 & 1 \\ \lambda^2 k^2 & 0 \end{pmatrix}, V(t) = \begin{pmatrix} 0 & 0 \\ \lambda^2 (\tilde{m}^2(t) - k^2) & 0 \end{pmatrix},$$

where $k = \tilde{m}(\infty) = \exp(\int_0^\infty b(t)dt)$. Noticing that $V' \in L^1$ and $\lim_{t \to \infty} V(t) = 0$, we could apply the Levinson theorem to the system. Then there exists $t_0 \in [0, \infty)$ so that we have two independent solutions, which have the asymptotic form as $t \to \infty$

$$Y_{\pm}(t) = \begin{pmatrix} 1 + o(1) \\ \pm \lambda k + o(1) \end{pmatrix} e^{\pm \lambda \int_{t_0}^t \tilde{m}(\tau)d\tau}$$

Thus, we have, for some d_1, d_2 ,

$$(4.9) y' = d_1(\lambda k + o(1))e^{\lambda \int_{t_0}^t \tilde{m}(\tau)d\tau} + d_2(-\lambda k + o(1))e^{-\lambda \int_{t_0}^t \tilde{m}(\tau)d\tau}$$

as $t \to \infty$. By (4.8), we know that $y' \ge C_1 + \delta_1^2 \lambda^2 C_0$ for all $t \ge 1$, it is clear that $d_1 > 0$. Then there exists some T > 1 such that

(4.10)
$$y' \simeq d_1 e^{\lambda \int_{t_0}^t \tilde{m}(\tau)d\tau} \simeq d_1 e^{\lambda \eta(t)} , \ \forall t \ge T .$$

So we have

$$y' \ge ce^{\lambda \eta(t)}, \forall t \ge 1,$$

for some C > 0.

In conclusion, we obtain

$$H' \ge \varepsilon y' \ge c\varepsilon e^{\lambda \eta(t)}, \forall t \ge 1$$
.

which completes the proof of (4.3).

References

- [1] Y. Bai and M. Liu, Global existence for semilinear damped wave equations in the scattering case, *Differential Integral Equations*, 32(3): 233–248, 2019.
- [2] J. Bony and D. Häfner, The semilinear wave equation on asymptotically Euclidean manifolds, Comm. Partial Differential Equations, 35(1): 23-67, 2010.
- [3] E. A. Coddington and N. Levinson, *Theory of ordinary differential equations*, McGraw-Hill Book Company, Inc., New York-Toronto-London, 1955.
- [4] W. Dai, D. Fang and C. Wang, Global existence and lifespan for semilinear wave equations with mixed nonlinear terms, J. Differential Equations, 267(5): 3328–3354, 2019.
- [5] D. Fang and C. Wang, Weighted Strichartz estimates with angular regularity and their applications, Forum Math. 23(1): 181–205, 2011.
- [6] W. Han and Y. Zhou, Blow up for some semilinear wave equations in multi-space dimensions, Comm. Partial Differential Equations, 39(4): 651–665, 2014.
- [7] K. Hidano, J. Jiang, S. Lee and C. Wang, Weighted fractional chain rule and nonlinear wave equations with minimal regularity, Rev. Mat. Iberoam. doi: 10.4171/rmi/1130 (to appear in print).
- [8] K. Hidano and K. Tsutaya, Global existence and asymptotic behavior of solutions for nonlinear wave equations, *Indiana Univ. Math. J.*, 44: 1273–1305, 1995.
- [9] K. Hidano, C. Wang and K. Yokoyama, The Glassey conjecture with radially symmetric data, J. Math. Pures Appl., 9: 518-541, 2012.
- [10] K. Hidano, C. Wang and K. Yokoyama, On almost global existence and local well posedness for some 3-D quasi-linear wave equations, Adv. Differential Equations, 17(3): 267–306, 2012.
- [11] K. Hidano, C. Wang and K. Yokoyama, Combined effects of two nonlinearities in lifespan of small solutions to semi-linear wave equations, Math. Ann., 366(1-2): 667–694, 2016.
- [12] K. Hidano and C. Wang, Fractional derivatives of composite functions and the Cauchy problem for the nonlinear half wave equation, Selecta Math. (N.S.), 25(1): 2–28, 2019.
- [13] F. John, Blow-up for quasilinear wave equations in three dimensions, Comm. Pure Appl. Math., 34: 29–51, 1981.
- [14] F. John and S. Klainerman, Almost global existence to nonlinear wave equations in three space dimensions, Comm. Pure Appl. Math., 37(4): 443–455, 1984.
- [15] M. Keel, H. Smith and C. Sogge, Almost global existence for some semilinear wave equations, J. Anal. Math., 87: 265–279, 2002.

- [16] N. Lai, Blow-up and lifespan estimate to a nonlinear wave equation in Schwarzschild spacetime, arXiv:1812.06542, 2018.
- [17] N. Lai and H. Takamura, Nonexistence of global solutions of nonlinear wave equations with weak time-dependent damping related to Glassey's conjecture, *Differential Integral Equa*tions, 32(1): 37–48, 2019.
- [18] N. Lai and H. Takamura, Nonexistence of global solutions of wave equations with weak time-dependent damping and combined nonlinearity, *Nonlinear Anal. Real World Appl.*, 45: 83–96, 2019.
- [19] M. Liu and C. Wang, Global existence for semilinear damped wave equations in relation with the strauss conjecture. Discrete Contin. Dyn. Syst. 40(2): 709–724, 2020.
- [20] M. Liu and C. Wang, The blow up of solutions to semilinear wave equations on asymptotically Euclidean manifolds, preprint.
- [21] J. Luk, The null condition and global existence for nonlinear wave equations on slowly rotating Kerr spacetimes, J. Eur. Math. Soc., 15(5): 1629–1700, 2013.
- [22] J. Metcalfe and C. Sogge, Long-time existence of quasilinear wave equations exterior to starshaped obstacles via energy methods, SIAM J. Math. Anal., 38(1): 188–209, 2006.
- [23] H. Smith, Wave equations with low regularity coefficients, Doc. Math., 2: 723-730, 1998.
- [24] T. C. Sideris, Global behavior of solutions to nonlinear wave equations in three dimentions, Comm. Partial Differential Equations, 8: 1291–1323, 1983.
- [25] C. Sogge and C. Wang, Concerning the wave equation on asymptotically Euclidean manifolds, J. Anal. Math., 112: 1–32, 2010.
- [26] D. Tataru, Strichartz estimates for second order hyperbolic operators with nonsmooth coefficients. III, J. Amer. Math. Soc., 15(2): 419–442, 2002.
- [27] N. Tzvetkov, Existence of global solutions to nonlinear massless Dirac system and wave equation with small data, *Tsukuba J. Math*, 22: 193–211, 1998.
- [28] K. Wakasa and B. Yordanov, Blow-up of solutions to critical semilinear wave equations with variable coefficients, J. Differential Equations, 266(9): 5360-5376, 2019.
- [29] C. Wang, The Glassey conjecture on asymptotically flat manifolds, Trans. Amer. Math. Soc., 367: 7429–7451, 2015.
- [30] C. Wang, The Glassey conjecture for nontrapping obstacles, J. Differential Equations, 259: 510–530, 2015.
- [31] C. Wang and X. Yu, Recent works on the Strauss conjecture, Contemp. Math., 581: 235–256, 2012.
- [32] C. Wang and H. Zhou, Almost global existence for semilinear wave equations with mixed nonlinearities in four space dimensions. J. Math. Appl. 459(1): 236–246, 2018.
- [33] B. Yordanov and Q. Zhang, Finite time blow up for critical wave equations in high dimensions, J. Funct. Anal., 231(2): 361–374, 2006.
- [34] Y. Zhou, Blow up of solutions to the Cauchy problem for nonlinear wave equations, Chinese Ann. Math. Ser. B, 22: 275–280, 2001.
- [35] Y. Zhou and W. Han, Blow-up of solutions to semilinear wave equations with variable coefficients and boundary, J. Math. Appl., 374(2): 585–601, 2011.

Department of Mathematics, Zhejiang Sci-Tech University, Hangzhou 310018, P. R. China

 $E ext{-}mail\ address: mengyunliu@zstu.edu.cn}$

School of Mathematical Sciences, Zhejiang University, Hangzhou 310027, P. R. China $E\text{-}mail\ address:\ wangcbo@zju.edu.cn$

URL: http://www.math.zju.edu.cn/wang