

# Another look at the Lady Tasting Tea and differences between permutation tests and randomization tests

Jesse Hemerik<sup>\*†</sup> and Jelle Goeman<sup>‡</sup>

February 6, 2022

## Summary

The statistical literature is known to be inconsistent in the use of the terms “permutation test” and “randomization test”. Several authors successfully argue that these terms should be used to refer to two distinct classes of tests and that there are major conceptual differences between these classes. The present paper explains an important difference in mathematical reasoning between these classes: a permutation test fundamentally requires that the set of permutations has a group structure, in the algebraic sense; the reasoning behind a randomization test is not based on such a group structure and it is possible to use an experimental design that does not correspond to a group. In particular, we can use a randomization scheme where the number of possible treatment patterns is larger than in standard experimental designs. This leads to exact  $p$ -values of improved resolution, providing increased power for very small significance levels, at the cost of decreased power for larger significance levels. We discuss applications in randomized trials and elsewhere. Further, we explain that Fisher’s famous Lady Tasting Tea experiment, which is commonly referred to as the first permutation test, is in fact a randomization test. This distinction is important to avoid confusion and invalid tests.

*keywords:* Permutation test; Group invariance test; Randomization test; Lady Tasting Tea;

## 1 Introduction

The statistical literature is very inconsistent in the use of the terms “permutation tests” and “randomization tests” (Onghena, 2018; Rosenberger et al., 2019). Both terms are often used to refer to tests that involve permutations. Sometimes these two terms are considered to refer to strictly distinct classes, sometimes to the same and sometimes to partly overlapping classes. The confusion surrounding differences between such tests is an important issue, because there are major differences between permutation tests and randomization tests in the sense of e.g. Onghena (2018), Kempthorne and Doerfler (1969) and Rosenberger et al. (2019), whose

---

<sup>\*</sup>Biometris, Wageningen University & Research

<sup>†</sup>Address for correspondence: Jesse Hemerik, Biometris, Wageningen University & Research, P.O. Box 16, 6700 AC Wageningen, The Netherlands. e-mail: jesse.hemerik@wur.nl

<sup>‡</sup>Biomedical Data Sciences, Leiden University Medical Center, Einthovenweg 20, 2333 ZC Leiden, The Netherlands

definitions we will follow here. Those authors use these terms to refer to strictly distinct classes of tests and discuss the terms in detail. Permutation tests are based on random sampling from populations and randomization tests are based on experimental randomization of treatments. With “randomization of treatments” we refer to the (physical) randomization that is part of the experimental design. With “permutation-based tests” we will refer to all tests involving permutations, including randomization tests involving permutations.

Another point of confusion has been the role of a *group structure*, in the algebraic sense, in permutation-based tests. Southworth et al. (2009), among others, explain that for permutation tests to have proven properties, it is important that the set of permutations used has such a group structure, as we discuss in Section 2. For example, the set of *balanced permutations*, which is a subset of a permutation group, does not have a group structure, and using it within a permutation test tends to lead to a very anti-conservative test. A balanced permutation, roughly speaking, is a permutation map that moves exactly 50 percent of the cases to the control group and 50 percent of the controls to the case group. Balanced permutations (not to be confused with stratified permutations) have been used in several publications (Fan et al., 2004; Jones-Rhoades et al., 2007) but Southworth et al. (2009) warn against their use.

A common reference in the permutation literature is the “Lady Tasting Tea” experiment, described in Fisher (1935a, Ch. II). This experiment is commonly referred to as the first published permutation test (Wald and Wolfowitz, 1944; Hoeffding, 1952; Anderson and Robinson, 2001; Lehmann and Romano, 2005; Langsrud, 2005; Mielke and Berry, 2007; Phipson and Smyth, 2010; Winkler et al., 2014). Indeed, like permutation tests, this test is based on permutations. We will see, however, that it is not a permutation test, but a randomization test, in the sense of Onghena (2018), Rosenberger et al. (2019) and Kempthorne and Doerfler (1969).

A main goal of the present paper is to explain the difference between two classes of tests involving permutations (or other transformations): tests that fundamentally rely on a group structure, and tests that do not, in an appropriate sense. To our knowledge this explicit distinction has not been made before. It is connected to the difference between permutation tests and randomization tests: the former fundamentally rely on a group structure and the latter do not – with the caveat that a randomization test should reflect the randomization scheme of the (physical) experiment. If the randomization scheme corresponds to a group, then the test also involves that group; otherwise the test does not involve a group. The fundamental point of this paper is as follows: the mathematical reasoning underlying a randomization test is not based on a group structure (even if a group happens to be used); the reasoning underlying a permutation test, on the other hand, is always based on a group structure and is completely different from the reasoning underlying randomization tests. In the existing literature, many randomization tests involve a group, but randomization tests that do not involve a group are also often considered (Onghena and Edgington, 1994, 2005; Rosenberger et al., 2019). The further contributions of this paper are as follows.

First of all, since permutation-based randomization tests do not require a group structure, it can be useful to consider a randomization scheme that does not correspond to a group. We introduce the idea of using an alternative randomization scheme to increase the number of possible treatment patterns. This increases the resolution of the  $p$ -value, thus improving power for very small significance levels  $\alpha$ , at the price of power loss for larger  $\alpha$ .

In addition, this paper provides the caveat that the Lady Tasting Tea experiment is rather different from permutation tests (in the sense of e.g. Onghena, 2018). Referring to

the Lady Tasting Tea experiment as an example of a permutation test, as is often done, can put readers on the wrong foot, since the reasoning underlying this experiment is not based on a group structure. Referring to the Lady Tasting Tea may have contributed to the confusion that has led researchers to design invalid permutation tests without a group structure (Southworth et al., 2009). The purpose of this paper is not to identify the first permutation test, which would not be straightforward (Berry et al., 2014).

This paper is built up as follows. In Section 2 we review existing results on permutation and group invariance tests, emphasizing the key role of the group structure of the permutations. In Section 3.1 we discuss the Lady Tasting Tea experiment, emphasizing why it does *not* require a group structure to control the type I error rate. In Section 3.2 we generalize the test of Section 3.1, providing a general randomization test and mentioning applications. In Section 3.3 we apply the general randomization test in a randomized trial setting, discussing how we can obtain higher-resolution  $p$ -values than with a canonical permutation-based test. The performance of our alternative test is illustrated with simulations in Section 4. We end with a discussion.

## 2 Permutation tests and group invariance tests

In the present section we explain the mathematical reasoning behind permutation tests, focusing on type I error control. As mentioned, the terms “permutation test” and “randomization test” have been used inconsistently in the literature. A typical example of a permutation test in the sense of Onghena (2018), Kempthorne and Doerfler (1969) and Rosenberger et al. (2019) is discussed in Fisher (1936, pp. 58-59). In this thought experiment, measurements of the statures of 100 Englishmen and 100 Frenchmen are considered. These observations are assumed to be randomly sampled from their respective populations. Such a model, where observations are randomly sampled from their populations, is typical for permutation tests in the sense of for example Kempthorne and Doerfler (1969), Onghena (2018) and Rosenberger et al. (2019). Note that in this example, there is no randomization of treatments as in, for example, clinical trials. In the example in Fisher (1936, pp. 58-59), to test whether “the two populations are homogeneous”, the difference between the two sample means is computed and this is repeated for each permutation of the 200 observations. The null hypothesis is rejected if the original difference is larger than most of the differences obtained after permutation. We will return to this example below.

Permutation tests are special cases of the general group invariance test. The definition of the group invariance test in, for example, Hoeffding (1952), Lehmann and Romano (2005) and Hemerik and Goeman (2018b) is rather general, so that many randomization tests also fall under it. The relationships between permutation tests, randomization tests and group invariance tests are illustrated in Figure 1. The principle underlying the group invariance test can also be used to prove properties of various permutation-based multiple testing methods (Westfall and Young, 1993; Tusher et al., 2001; Meinshausen and Bühlmann, 2005; Hemerik and Goeman, 2018a; Hemerik et al., 2019).

A general definition of a group invariance test is as follows. Generalizations of this framework, such as two-sided tests, are possible. Let  $X$  model random data with support  $\mathcal{X}$ . For example,  $X$  could be a random vector or matrix. Consider a set  $\mathcal{G}$  of permutation maps or other transformations  $g : \mathcal{X} \rightarrow \mathcal{X}$ . We will assume that  $\mathcal{G}$  is finite, although generalizations




Figure 1: A Venn diagram showing relationships between permutation tests ( $\mathbf{P}$ ), randomization tests ( $\mathbf{R}$ ) and group invariance tests ( $\mathbf{G}$ ). Permutation tests ( $\mathbf{P}$ ) are a subclass of group invariance tests ( $\mathbf{G}$ ) and are based on random sampling from populations. Randomization tests ( $\mathbf{R}$ ) are based on randomization of treatments. The randomization scheme sometimes corresponds to a group ( $\mathbf{R} \cap \mathbf{G}$ ) and sometimes does not ( $\mathbf{R} \setminus \mathbf{G}$ ). If the scheme corresponds to a group then this is often a set of permutations, but not always. There are also group invariance tests that fall outside all these categories ( $\mathbf{G} \setminus (\mathbf{P} \cup \mathbf{R})$ ). An example of a test from class  $\mathbf{P}$  is Fisher’s thought experiment with Englishmen and Frenchmen described in Section 2. An example from the class  $\mathbf{R} \cap \mathbf{G}$  is Fisher’s Lady Tasting Tea experiment. A different example, based on sign-flipping instead of permutation, is the test in Fisher (1935a, §21). An example from the class  $\mathbf{R} \setminus \mathbf{G}$  is in Section 3.3. An example from the class  $\mathbf{G} \setminus (\mathbf{P} \cup \mathbf{R})$  are certain tests based on rotations (Solari et al., 2014) or sign-flipping (Winkler et al., 2014).

are possible. The set  $\mathcal{G}$  is assumed to have a group structure with respect to the operation of composition of maps, which means that:  $\mathcal{G}$  contains the identity map  $x \mapsto x$ ; every element in  $\mathcal{G}$  has an inverse; and for all  $g, h \in \mathcal{G}$ ,  $g \circ h \in \mathcal{G}$  (Hoeffding, 1952). Further, we consider some test statistic  $T : \mathcal{X} \rightarrow \mathbb{R}$ . Consider a null hypothesis  $H_0$  that implies that the joint distribution of all test statistics  $T(g(X))$  with  $g \in \mathcal{G}$  is invariant under all transformations of  $X$  in  $\mathcal{G}$  (Hemerik and Goeman, 2018b). This holds in particular if the data  $X$  are themselves transformation-invariant, i.e., if

$$g(X) \stackrel{d}{=} X \quad (1)$$

for every  $g \in \mathcal{G}$ .

A typical example of such a setting is the thought experiment from Fisher (1936, pp. 58-59), mentioned above. Let  $X_1, \dots, X_{100}$  be the statures of the Englishmen and let  $X_{101}, \dots, X_{200}$  be the statures of the Frenchmen. The test statistic considered in Fisher (1936, pp. 58-59) is

$$T(X) = \left| \frac{1}{100} \sum_{i=1}^{100} X_i - \frac{1}{100} \sum_{i=101}^{200} X_i \right|. \quad (2)$$

The null hypothesis  $H_0$  is that  $X_1, \dots, X_{200}$  are i.i.d.. The null hypothesis is rejected if the original test statistic is larger than most of the statistics obtained after permutation. (In practice one would nowadays sample random permutations, to avoid having to use all possible permutations, see Hemerik and Goeman, 2018b.) The group  $\mathcal{G}$  that Fisher considers consists of all permutation maps  $g : \mathbb{R}^{200} \rightarrow \mathbb{R}^{200}$ . Here, every  $g \in \mathcal{G}$  is of the form

$$(x_1, \dots, x_{200}) \mapsto (x_{\pi_1}, \dots, x_{\pi_{200}}),$$

where  $(\pi_1, \dots, \pi_{200})$  is a permutation of  $(1, \dots, 200)$ . Note that  $X$  is then  $\mathcal{G}$ -invariant under  $H_0$ , i.e., (1) holds for every  $g \in \mathcal{G}$ .

As another example, consider random data  $X$  with support in  $\mathbb{R}^n$ , with independent entries that are symmetric about their means. Suppose that under  $H_0$ , the entries have mean 0. This may for example be the case if each entry of  $X$  is the difference of two paired observations. Then the distribution of  $X$  is invariant under all transformations in  $\mathcal{G}$  under  $H_0$  if we define  $\mathcal{G}$  to be the group of all sign-flipping maps of the form

$$(x_1, \dots, x_n) \mapsto (s_1 x_1, \dots, s_n x_n), \quad (3)$$

with  $(s_1, \dots, s_n) \in \{-1, 1\}^n$ . We may take  $T((x_1, \dots, x_n)) = \sum_{i=1}^n x_i$ . If the data generating mechanism involves randomization of treatments (rather than sampling from populations), such a test can be considered a randomization test. The test already appears in Fisher (1935a, §21), albeit without explicit proof. See also Basu (1980).

In both examples above, we can apply the general group invariance test to test  $H_0$ . This test already appears in the literature (Hoeffding, 1952; Lehmann and Romano, 2005; Hemerik and Goeman, 2018b), but for completeness we include the result and its proof.

We will write  $gX = g(X)$  for short. Let  $k = \lceil (1 - \alpha)|\mathcal{G}| \rceil$ , the smallest integer which is larger than or equal to  $(1 - \alpha)|\mathcal{G}|$ . Let  $T^{(1)}(X) \leq T^{(2)}(X) \leq \dots \leq T^{(k)}(X) \leq \dots \leq T^{(|\mathcal{G}|)}(X)$  be the sorted values  $T(gX)$  with  $g \in \mathcal{G}$ .

**Theorem 1.** *The size of the group invariance test is at most  $\alpha$ , i.e., under  $H_0$ ,  $\mathbb{P}\{T(X) > T^{(k)}(X)\} \leq \alpha$ .*

*Proof.* By the group structure,  $\mathcal{G}g = \mathcal{G}$  for all  $g \in \mathcal{G}$ . Hence  $T^{(k)}(gX) = T^{(k)}(X)$  for all  $g \in \mathcal{G}$ . Let  $G$  have the uniform distribution on  $\mathcal{G}$ . Then under  $H_0$ , the rejection probability is

$$\begin{aligned} \mathbb{P}\{T(X) > T^{(k)}(X)\} &= \\ \mathbb{P}\{T(GX) > T^{(k)}(GX)\} &= \\ \mathbb{P}\{T(GX) > T^{(k)}(X)\}. \end{aligned}$$

The first equality follows from the null hypothesis and the second equality holds since  $T^{(k)}(X) = T^{(k)}(GX)$ . Since  $G$  is uniform on  $\mathcal{G}$ , the above probability equals

$$\mathbb{E}\left[|\mathcal{G}|^{-1} \cdot \left|\{g \in \mathcal{G} : T(gX) > T^{(k)}(X)\}\right|\right] \leq \alpha,$$

as was to be shown.  $\square$

Under additional assumptions, the test is exact, i.e., the rejection probability is exactly  $\alpha$  under  $H_0$  (Hemerik and Goeman, 2018b). In the above proof we used the group structure, which guarantees the symmetry property  $\mathcal{G}g = \mathcal{G}$  for all  $g \in \mathcal{G}$ . A different proof, based on conditioning on the pooled sample, is also possible and also requires using this symmetry property (first proof of Theorem 1 in Hemerik and Goeman, 2018b).

Write  $\mathcal{GX} = \{gX : g \in \mathcal{G}\}$  and assume for convenience that  $gX$  and  $g'X$  are distinct with probability 1 if  $g, g' \in \mathcal{G}$  are distinct. This is usually the case if  $X$  is continuous. The permutation test is based on the fact that under  $H_0$ , for every permutation  $g \in \mathcal{G}$  the probability

$\mathbb{P}\{T(gX) > T^{(k)}(X)\}$  is the same. The reason is that under  $H_0$ , for every  $g \in \mathcal{G}$ , the joint distribution of  $(gX, \mathcal{G}X)$  is the same. This is because if  $g, g' \in \mathcal{G}$ , under  $H_0$  we have

$$(gX, \mathcal{G}X) = (gX, \mathcal{G}gX) \stackrel{d}{=} (X, \mathcal{G}X) \stackrel{d}{=} (g'X, \mathcal{G}g'X) = (g'X, \mathcal{G}X).$$

When  $\mathcal{G}g = \mathcal{G}$  does not hold for all  $g \in \mathcal{G}$ , then the above does not generally hold under  $H_0$ .

The group structure of  $\mathcal{G}$  implies that  $\mathcal{G}g = \mathcal{G}$  for all  $g \in \mathcal{G}$ . Under the mild condition that all  $g \in \mathcal{G}$  are surjective, the reverse implication also holds, i.e., if  $\mathcal{G}g = \mathcal{G}$  for all  $g \in \mathcal{G}$ , then  $\mathcal{G}$  is a group. For example, if  $\mathcal{G}g = \mathcal{G}$  for all  $g \in \mathcal{G}$ , there are  $h, g \in \mathcal{G}$  with  $hg = g$ . It follows that  $\mathcal{G}$  contains an identity element, and the other group properties also easily follow. We conclude that in the argument underlying the permutation test, the group structure is key.

In practice it is often computationally infeasible to use a test based on the full group  $\mathcal{G}$  of transformations. Researchers then usually resort to using a limited number of random transformations, uniformly sampled from the group  $\mathcal{G}$ . It is then still possible to obtain an exact test (see [Hemerik and Goeman, 2018b](#) and [Phipson and Smyth, 2010](#) for a detailed treatment).

### 3 The Lady Tasting Tea and randomization tests

Unlike a permutation test, a randomization test is based on data collected in an experiment involving randomization of treatments. The randomization scheme of the physical experiment does not necessarily correspond to a group, and if it does not, the statistical test does not involve a group either. Even if a group is used, the reasoning underlying a randomization test is not based on the group structure and is very different from the reasoning underlying permutation tests.

In this section, we first discuss the Lady Tasting Tea experiment, explaining that the reasoning underlying the test is not based on a group structure, because it is based on randomization. This experiment is a special case of a general randomization test that we discuss in Section 3.2. In Section 3.3 we apply this test to provide higher-resolution  $p$ -values in randomized trials.

#### 3.1 The Lady Tasting Tea experiment

In the Lady Tasting Tea experiment ([Fisher, 1935a](#), Ch. II), the null hypothesis is that a particular lady cannot distinguish between two types of cups of tea with milk: cups in which the tea was added first and cups in which the milk was added first. To test the null hypothesis, which we denote by  $H_0$ , the experimenter “mixes eight cups of tea, four in one way and four in the other,” and presents them “to the subject for judgment in a random order.” The experimental setup is made known to the lady. The lady then tastes from the cups and has to determine which four cups in the sequence of eight had milk added first. Fisher actually performed the experiment ([Box, 1978](#); [Berry et al., 2014](#)).

There are  $\binom{8}{4} = 70$  possible orders, with respect to the two types of cups. Suppose  $H_0$  is true. If the lady guesses every pattern with probability  $1/70$ , then the probability that she chooses the correct order is  $1/70$ . Even if she has an a priori preference for a certain order, the probability of guessing correct is  $1/70$ . Indeed, it is assumed that the researcher randomizes the true pattern, i.e., he chooses each pattern with equal probability. Thus, if

we reject  $H_0$  when the lady identifies all four “milk first” cups correctly, then the probability of a type I error is  $1/70$  (1.4%). The probability that she labels three of the “milk first” cups correctly is  $\binom{4}{3}\binom{4}{1}/70 = 16/70$  (22.9%) and the probability of two correct picks is  $36/70$  (51.4%). Thus, for example, when we reject  $H_0$  if at least three picks are correct, the level is  $16/70 + 1/70 = 17/70$  (24.3%). The test is equivalent to an instance of “Fisher’s exact test” (Yates, 1934; Fisher, 1935b; Berry et al., 2014) with pre-fixed marginal frequencies in the  $2 \times 2$  table. Fisher’s exact test, however, was not originally motivated from a permutation or randomization testing perspective.

Mathematically, we can describe the experiment as follows. Let  $\mathcal{W} \subset \{0, 1\}^8$  be the set of vectors containing four 0’s and four 1’s, so that the cardinality of  $\mathcal{W}$  is  $R := |\mathcal{W}| = 70$ . Let the decision of the lady be denoted by  $Y$  and let  $W$  denote the true order, i.e., the random decision by the experimenter. Here,  $Y$  and  $W$  are random variables taking values in  $\mathcal{W}$ . Note that  $W$  represents the ‘treatments’ given by the experimenter and  $Y$  represents the lady’s ‘responses’. The experimenter’s order  $W$  is assumed to be uniformly distributed on  $\mathcal{W}$ . The null hypothesis is

$$H_0 : Y \text{ is independent of } W.$$

Let  $\alpha \in (0, 1)$  be the desired type I error rate. If  $\alpha \in A = \{1/70, 17/70, 53/70, 69/70\}$ , then  $\alpha$  is called *attainable* in the Lady Tasting Tea experiment, meaning that we obtain a test of exactly level  $\alpha$  (Pesarin, 2015). If  $\alpha$  is not attainable, then we obtain a test with level strictly less than  $\alpha$ .

Let  $T : \mathcal{W} \times \mathcal{W} \rightarrow \mathbb{R}$  be a test statistic such that high values of  $T(w, y)$  indicate that the patterns  $w$  and  $y$  are similar, i.e., that there is evidence against  $H_0$ . Let

$$T^{(1)}(Y) \leq \dots \leq T^{(70)}(Y)$$

be the sorted statistics  $T(w, Y)$  with  $w \in \mathcal{W}$ . Whether the vector of sorted statistics  $(T^{(1)}, \dots, T^{(70)})$  actually depends on  $Y$  or not, depends on the definition of  $T$ ; in Fisher (1935a), the test statistic is

$$T(W, Y) = \sum_{i=1}^8 \mathbb{1}_{\{W_i=1\} \cap \{Y_i=1\}} = \sum_{i=1}^8 W_i Y_i \quad (4)$$

and it can be seen the sorted statistics do not depend on  $Y$ , since  $Y$  always has four entries that are 1. Let  $\lceil (1 - \alpha)R \rceil$  be the smallest integer that is at least  $(1 - \alpha)R$ . We have the following result (Fisher, 1935a).

**Theorem 2.** *The test that rejects  $H_0$  if and only if  $T(W, Y) > T^{(\lceil (1-\alpha)R \rceil)}(Y)$  has size at most  $\alpha$ .*

*Proof.* Assume  $H_0$  holds. Conditional on  $Y$ ,  $W$  is uniformly distributed on  $\mathcal{W}$  and  $T^{(\lceil (1-\alpha)R \rceil)}(Y)$  is known. Hence, conditional on  $Y$ , the rejection probability is

$$\begin{aligned} \mathbb{P}(W \in \{w \in \mathcal{W} : T(w, Y) > T^{(\lceil (1-\alpha)R \rceil)}(Y)\}) &= \\ \frac{1}{R} |\{w \in \mathcal{W} : T(w, Y) > T^{(\lceil (1-\alpha)R \rceil)}(Y)\}| &\leq \alpha. \end{aligned}$$

Thus, marginal over  $Y$ , the rejection probability is also at most  $\alpha$ .  $\square$



Observe that when we use the test statistic (4), then taking  $\alpha \in A$  indeed results in an exact test. This follows from the fact that

$$T^{(1)} < T^{(2)} = T^{(3)} = \dots = T^{(17)} < T^{(18)} = T^{(19)} = \dots = T^{(53)} < T^{(54)} = \dots = T^{(69)} < T^{(70)},$$

by the argument at the beginning of this section. If  $\alpha \in (0, 1) \setminus A$ , the level is strictly smaller than  $\alpha$ . If the experimenter does not choose randomly from all 70 possible patterns, but uses some smaller set of patterns for him and the lady to choose from, then there may not be any  $\alpha \in (0, 1)$  for which the test is exact, since the sorted test statistics may depend on  $Y$ . This is one of the reasons why using the full set of patterns, in combination with a suitable test statistic  $T$ , is useful. However, to prove Theorem 2, we did not need to use the group structure of the permutations. The reason is that in the Lady Tasting Tea experiment, under  $H_0$  the randomization  $W$  of the researcher is by design independent of the reference set  $\{(w, Y) : w \in \mathcal{W}\}$ . Further considerations follow below.

### 3.2 A general randomization test

Theorem 2 still applies if the researcher uses a set of permutations that does not correspond to a group. Suppose for example that the researcher omits one of the patterns, thus picking randomly from some set  $\mathcal{W}$  of 69 patterns, with or without the lady's knowledge. Denote the set that the lady chooses from by  $\mathcal{Y}$ . Then  $W$  and  $Y$  will still be independent and Theorem 2 still applies if we let  $R = |\mathcal{W}| = 69$  and let

$$T^{(1)}(W) \leq \dots \leq T^{(69)}(W)$$

be the sorted test statistics  $T(w, Y)$ ,  $w \in \mathcal{W}$ . Indeed, conditional on  $Y$ ,  $W$  will have a uniform distribution on  $\mathcal{W}$ . In fact, we have the following very general randomization test, of which the Lady Tasting Tea experiment is a special case. We refer to this result as a randomization test since in most applications of the theorem, the variable  $W$  will represent experimental randomization of treatments (Kempthorne and Doerfler, 1969; Ongheena, 2018). The idea of the theorem is certainly not new; it is at least implicitly present in earlier works (Morgan et al., 2012).

**Theorem 3** (General randomization test). *Let  $\mathcal{W}$  and  $\mathcal{Y}$  be nonempty sets, where  $\mathcal{W}$  is assumed to be finite. Write  $R = |\mathcal{W}|$ . Let  $Y$  be a variable taking values in  $\mathcal{Y}$  and assume  $W$  is uniformly distributed on  $\mathcal{W}$ . (Here  $Y$  and  $W$  are variables in a general sense, e.g. they may be random vectors.) Let  $T : \mathcal{W} \times \mathcal{Y} \rightarrow \mathbb{R}$  be some test statistic. Consider a null hypothesis  $H_0$  that implies that  $Y$  is independent of  $W$ . Let  $T^{(1)}(Y) \leq \dots \leq T^{(R)}(Y)$  be the sorted values  $T(w, Y)$  with  $w \in \mathcal{W}$ . Consider the test that rejects  $H_0$  if and only if  $T(W, Y) > T^{(\lceil (1-\alpha)R \rceil)}(Y)$ . Then the result of Theorem 2 still applies, i.e., the test has size at most  $\alpha$ .*

The proof is analogous to that of Theorem 2: under  $H_0$ , conditional on  $Y$ ,  $W$  is uniformly distributed on  $\mathcal{W}$  and  $T^{(\lceil (1-\alpha)R \rceil)}(Y)$  is known. Hence, conditional on  $Y$ , the rejection probability is

$$\mathbb{P}(W \in \{w \in \mathcal{W} : T(w, Y) > T^{(\lceil (1-\alpha)R \rceil)}(Y)\}) \leq \alpha$$

as before.



Under additional assumptions, the test is exact, i.e., the rejection probability is exactly  $\alpha$  under  $H_0$ . We assumed that  $\mathcal{W}$  is finite, but generalizations to infinite  $\mathcal{W}$  are possible, as well as generalizations to non-uniform  $W$ . We can also define a two-sided test. Moreover, as in Section 2, under straightforward additional assumptions, we can prove that the test of Theorem 3 is exact for certain  $\alpha$ , i.e., that the rejection probability is exactly  $\alpha$  under  $H_0$  (Hemerik and Goeman, 2018b).

Note that in Theorem 3,  $Y$  might be a constant, conditional on  $W$ . In fact, the randomization testing literature often views the outcomes as non-random, conditional on the treatments. This corresponds to the fact that randomization tests can be used without an assumption that the responses are randomly sampled from populations (Cox, 2009; Onghena, 2018; Rosenberger et al., 2019). We discuss this further in the context of randomized trials in Section 3.3.

The general randomization test of Theorem 3 has many applications. Examples are agricultural experiments and randomized clinical trials. Randomized trials will be discussed in Section 3.3. We mention a few other interesting applications here.

First of all, Theorem 3 has implications for the Lady Tasting Tea experiment. In Section 3.1, it is assumed that the lady knows beforehand that there are  $m$  cups of each type, where  $2m$  is the total number of cups she receives. If for some reason she does not know that, then she might label e.g.  $m+1$  of the  $2m$  items with the same label. In other words, she might pick a pattern from a set containing more patterns than the experimenter chooses from. Theorem 3 then says that the type I error probability will nevertheless be at most  $\alpha$  under  $H_0$ . Indeed, in Theorem 3,  $\mathcal{Y}$  is allowed to be any nonempty set, so in particular it can be larger than  $\mathcal{W}$ .

A further application of Theorem 3 are general sensory tests, of which the Lady Tasting Tea experiment is an example. It is interesting to note that in the literature on sensory tests, Fisher’s experiment has been regarded a “forerunner of modern sensory analysis” (Bi and Kuesten, 2015). For example, Harris and Kalmus (1949) perform a sensory experiment that is analogous to the Lady Tasting Tea experiment, as follows: “The glasses are arranged at random. The subject is told that four of them contain the substance and four contain water, and he is asked to taste them all and to separate them into the two groups of four.”

Another application of Theorem 3 are existing permutation-based randomization tests that are used to evaluate whether some classification algorithm has any predictive ability (both in-sample and out-of-sample). Such tests can be used to evaluate algorithms for, for example, text categorization, fraud detection, optical character recognition and medical diagnosis. Tests of this type are discussed in, for instance, Golland et al. (2005), Airola et al. (2010), Ojala and Garriga (2010), Schreiber and Krekelberg (2013) and Rosenblatt et al. (2019).

### 3.3 Randomization testing without a group structure: higher-resolution $p$ -values

In randomized trials, often we are interested in comparing two different treatments, for example a drug and a placebo. In such a setting, there is a treatment assignment randomized by the experimenter. In that case, we can use the randomization test of Theorem 3, as explained below. As discussed, we then do not require a group structure, to control the type I error rate. We now discuss such a setting in detail. The tests considered here will also be studied with simulations in Section 4.

Let  $n \geq 2$  be an integer, assumed even for convenience, and suppose we have  $n$  subjects,  $n/2$  of which receive one treatment and  $n/2$  of which receive the other treatment. Let  $W = (W_1, \dots, W_n)$  denote the treatments and  $Y = (Y_1, \dots, Y_n)$  the responses, taking values in  $\mathbb{R}^n$ . The treatment pattern  $W$  is uniformly sampled from a set  $\mathcal{W} \subseteq \{0, 1\}^n$ . The most common type of randomized trial is the *forced balance procedure*, where

$$\mathcal{W} = \{w \in \{0, 1\}^n : w \text{ contains } n/2 \text{ 1's}\} \quad (5)$$

(Rosenberger and Lachin, 2015; Lachin, 1988b; Braun and Feng, 2001). For each  $1 \leq i \leq n$ , the response  $Y_i \in \mathbb{R}$  is independent of all the other subjects' treatments and responses. We consider the null hypothesis  $H_0$  that  $Y$  is independent of  $W$ .

These assumptions are still rather general. It can be useful to consider a more specific randomization model as in Pitman (1937, §7), who assumes an additive treatment effect. An important property of randomization models, is that to test whether the treatment has an effect on our particular patients, we do not need to assume that they are random draws from populations. We could consider the patients as fixed and  $Y$  as constant, conditional on  $W$  (Pitman, 1937, §7). Indeed, "Any assumption that the units are, say, a random sample from a population of units [...] is additional to the specification" of the model (Cox, 2009). This property is discussed in detail in Onghena (2018) and Rosenberger et al. (2019).

We can invoke Theorem 3 to obtain a test that controls the type I error rate. We can also obtain an exact test, i.e., a test that rejects with probability exactly  $\alpha$  under  $H_0$ . Consider the test statistic  $T : \mathcal{W} \times \mathbb{R}^n \rightarrow \mathbb{R}$  defined as

$$T(W, Y) = \sum_{\{i: W_i=1\}} Y_i - \sum_{\{i: W_i=0\}} Y_i. \quad (6)$$

Recall that  $Y$  may be viewed as random or constant, conditional on  $W$ . In either case, assume that  $Y$  is such that (with probability 1), for all distinct  $w_1, w_2 \in \mathcal{W}$ ,  $T(w_1, Y) \neq T(w_2, Y)$ . This is satisfied in particular if  $Y_1, \dots, Y_n$  have continuous distributions. Write  $N = \binom{n}{n/2}$ . The test is exact if  $\alpha \in (0, 1)$  is a multiple of  $1/|\mathcal{W}|$ , where  $|\mathcal{W}|$  equals  $N$ . An exact  $p$ -value is

$$p(W, Y) = \frac{|\{w \in \mathcal{W} : T(w, Y) \geq T(W, Y)\}|}{|\mathcal{W}|}, \quad (7)$$

i.e., if  $\alpha \in (0, 1)$  is a multiple of  $1/|\mathcal{W}|$ , then  $\mathbb{P}(p \leq \alpha) = \alpha$  under  $H_0$ . A two-sided exact test can be obtained analogously.

Since Theorem 3 applies, the test essentially does not rely on a group structure. Hence, we may consider taking  $\mathcal{W}$  to be a set that does not correspond to a group. In a different context, this is also done in Onghena and Edgington (1994) and Onghena and Edgington (2005), where a set of permutations is used that is strictly smaller than the full set of permutations. This is done to avoid too repetitive treatment patterns such as ABBBBAAA. In our setting, if  $n = 8$ , instead of taking  $\mathcal{W}$  to be the set of all permutations of  $(0, 0, 0, 0, 1, 1, 1, 1)$  we could take  $\mathcal{W}$  to be a subset that does not correspond to a group, and still obtain an exact test (for certain  $\alpha$ ). As Onghena and Edgington (1994) illustrates, this may be useful in some settings. However, in a typical randomized trial there is no evident reason to only use a subset of the permutations, except to limit the number of permutations for computational reasons.

A more interesting alternative, when running the experiment, is to draw  $w$  from a set that is strictly larger than the set in (5), for example, from the set of all possible labelings,  $\{0, 1\}^n$

(a *Bernoulli trial*, [Imbens and Rubin, 2015](#)). Indeed, if the standard randomization scheme is used, i.e., the forced balance trial, then the smallest possible  $p$ -value that can be obtained is  $1/N$ , due to the discreteness of the  $p$ -value. If  $n = 8$  for example, then  $1/N = 1/70$ . This means that if the significance level is  $\alpha = 0.01$  for instance, we have a power of 0 to reject  $H_0$ . Such small  $\alpha$  are often used nowadays, for example due to multiple testing. The discreteness of the permutation  $p$ -value is a well-known downside of permutation-based tests ([Berger, 2000](#)). If we take  $\mathcal{W} = \{0, 1\}^n$ , however, then  $|\mathcal{W}| = 2^n$ , so that the smallest possible  $p$ -value is  $1/2^n = 1/256$ . If  $1/256 \leq \alpha < 1/70$ , this means a uniform improvement in power over the standard randomization test. Note that there only is a power improvement for very small  $\alpha$ ; for larger  $\alpha$  the Bernoulli trial has less power than the forced balance trial.

Under  $H_0$ , if  $\alpha$  is a multiple of  $1/2^n$ , the test with  $\mathcal{W} = \{0, 1\}^n$  rejects with probability exactly  $\alpha$ . Otherwise the test rejects with probability less than  $\alpha$  under  $H_0$ . For  $\mathcal{W} = \{0, 1\}^n$ , to our knowledge it is not known what the optimal choice of  $T$  is for testing an additive treatment effect. In Section 4 we will take

$$T(W, Y) = \sum_{\{i: W_i=1\}} (Y_i - \bar{Y}) - \sum_{\{i: W_i=0\}} (Y_i - \bar{Y}), \quad (8)$$

where  $\bar{Y} = n^{-1}(Y_1 + \dots + Y_n)$ . Using this test statistic ensures that under  $H_0$ , the expected value of  $T(W, Y)$  does not depend on the random labelling  $W$ . In Appendix A we show how the Bernoulli trial can be modified to enforce covariate balancing.

That it is possible to take  $\mathcal{W} = \{0, 1\}^n$  has been noted by several authors ([Pocock, 1979](#); [Kalish and Begg, 1985](#); [Lachin, 1988a](#); [Wei and Lachin, 1988](#); [Suresh, 2011](#); [Imbens and Rubin, 2015](#); [Rosenberger et al., 2019](#)). They do not recommend this approach, but merely mention it as a possibility, while focusing on more common randomization schemes. Their main argument against taking  $\mathcal{W} = \{0, 1\}^n$  is that it leads to less power than the usual approach of restricted randomization ([Pocock, 1979](#), p.188). This is true when  $\alpha$  is large enough and it is then better use the forced balance approach. When  $\alpha$  is rather small, however, that test has 0 power, while the Bernoulli trial may have substantial power. The idea that using  $\mathcal{W} = \{0, 1\}^n$  leads to higher-resolution  $p$ -values, has not been mentioned before to our knowledge. Nowadays, the use of large multiple testing corrections is more common than in the past, so higher-resolution, exact  $p$ -values can clearly be of interest.

Suppose we use  $\mathcal{W} = \{0, 1\}^n$ . Then, if we happen to draw  $W = (0, \dots, 0)$  or  $W = (1, \dots, 1)$ , the value of the statistic (8) is 0 and we can have no hope of rejecting  $H_0$  (if  $\alpha = 0.05$ ). Hence we might exclude  $(0, \dots, 0)$  and  $(1, \dots, 1)$ , and perhaps more elements, from  $\mathcal{W}$ . In any case, if  $\alpha < 1/N$ , it can be useful to consider a design with  $|\mathcal{W}|$  larger than  $N$ . Note that in practice, we should choose  $\mathcal{W}$  before administering the treatments. Once the treatments have been given, we cannot change our minds about  $\mathcal{W}$ . The randomization test that uses all patterns from  $\{0, 1\}^n$  except  $(0, \dots, 0)$  and  $(1, \dots, 1)$  is further studied with simulations in Section 4.

### 3.4 Randomization testing under a random sampling model

For completeness we note the following, but it can be skipped at a first read. Existing permutation tests based on random sampling from populations rely on a group structure. However, in some cases, we can use an alternative approach to avoid the requirement of a group structure also in this setting. This is a novel idea, to our knowledge (and arguably falls outside the categories of Figure 1). The approach is analogous to the test in Section 3.3.

Suppose that we are comparing two populations, for example, a population of cases and a population of controls, or Englishmen and Frenchmen. Let  $W$  have the uniform distribution on  $\mathcal{W} = \{0, 1\}^n$  or some subset thereof, as before. Then we could draw from the two populations as indicated by  $W$ , i.e., for every  $1 \leq i \leq n$ , the  $i$ -th individual is drawn from the first population if  $W_i = 0$  and from the second population if  $W_i = 1$ . For  $1 \leq i \leq n$ , let  $Y_i$  be the observation for the  $i$ -th individual, for example his or her stature. We can then perform a test exactly as in Section 3.3, using the test statistic (8) and the  $p$ -value (7).

If we take  $\mathcal{W}$  as in (5), then the test will be equivalent to a standard permutation test. For many other choices of  $\mathcal{W}$ , we obtain a novel type of test. If we take for instance  $\mathcal{W} = \{0, 1\}^n$ , then the number of observations drawn from each population will be random, with only the total number of observations being fixed at  $n$ . In many situations this would be impractical, for example because there is only a limited, fixed number of cases. We will not pursue such tests further here.

## 4 Empirical example

Here we illustrate the idea in Section 3.3 with a simple simulation study. We considered two randomization tests: a standard randomization test and an alternative test that provides higher-resolution  $p$ -values, as discussed in Section 3.3. The setting was as in the example in Section 3.3, with  $n = 8$ . Every  $Y_i$  was distributed as the absolute value of a  $N(0, 1)$  variable if  $W_i = 0$ ; if  $W_i = 1$  it had the same distribution, but with an increase in mean of 2. Under the null hypothesis, the distribution of  $Y_i$  does not depend on  $W_i$ . The first test considered was the standard randomization test. This test uses  $N = (n!)/((n/2)!(n/2)!) = 70$  permutations. The second test was based on all relabellings in  $\{0, 1\}^n$  excluding  $(0, \dots, 0)$  and  $(1, \dots, 1)$ . Thus, this test used  $2^n - 2 = 254$  relabellings. We used the test statistic (8). By Theorem 3, both tests control the type I error rate. Moreover, the first test is exact if  $\alpha \in (0, 1)$  is a multiple of  $1/70$ . The second test is exact if  $\alpha$  is multiple of  $1/254$ . It is important to note that the two tests are based on different randomization schemes, i.e., on different data gathering mechanisms. In practice the type of test should already be decided upon before running the physical experiment.

Table 1: Performance of an alternative to the standard randomization test. Test 1 is the standard test, based on a forced balance procedure. Test 2 is the alternative test, based on more relabellings.

		$\alpha$				
	test	$1/254 (\approx .0039)$	.005	.01	.02	.05
size	test 1	0	0	0	.0122	.0418
	test 2	.0034	.0034	.0076	.0190	.0464
power	test 1	0	0	0	.9011	.9725
	test 2	.5443	.5443	.7027	.8436	.9316

In Table 1, for different values of the significance level  $\alpha$ , the estimated level and power of the two tests are shown. Every estimate in the table is based on  $10^4$  repeated simulations. The regular randomization test had no power for  $\alpha < 1/70$ , due to the fact that only 70

relabellings are available with this approach. Test 2, which is based on 254 relabellings, however, did have substantial power for  $1/254 \leq \alpha < 1/70$ , as explained in Section 3.3. In the table, the estimated size for  $\alpha = 1/254$  is 0.0034, which is approximately the true size  $1/254 \approx 0.0039$ . Note that for  $\alpha = 0.005$ , the size and power are the same as for  $\alpha = 1/254$ . The reason is the discreteness of the  $p$ -value: 0.005 lies between  $1/254$  and  $2/254$ .

## 5 Discussion

In this paper, we have distinguished between two types of permutation-based tests: tests that fundamentally rely on a group structure and tests based on treatment randomization, which do not necessarily require a group structure. We have discussed that in settings where treatments are randomly assigned, it can be useful to consider a randomization scheme that does not correspond to a group. In particular, this allows obtaining higher-resolution exact  $p$ -values than are possible with standard randomization tests. This paper also provides the caveat that referring to the Lady Tasting Tea experiment as an example of a permutation test can be misleading, since the reasoning underlying this experiment is not based on a group structure.

The two types of tests between which we distinguish roughly correspond to respectively “permutation tests” and “randomization tests” in the sense of Onghena (2018) and Rosenberger et al. (2019). As we mentioned, the use of these terms has been rather inconsistent throughout the literature. For example, Edgington and Onghena (2007, p.1) write that “*randomization tests* are a subclass of statistical tests called *permutation tests*”, while Onghena (2018) proposes to use the terms for strictly distinct classes of tests. In any case, we propose to use the term “randomization test” only when there is some form of treatment randomization. This is in line with Kempthorne and Doerfler (1969), Edgington and Onghena (2007), Onghena (2018) and Rosenberger et al. (2019).

As mentioned in the introduction, it would not be straightforward to identify the first permutation test (Berry et al., 2014). In any case, it is clear that, once the concepts of randomization of treatments and random sampling from populations had been established in the 1920’s (Rubin, 1990; Fisher, 1925; Neyman and Pearson, 1928), the way was paved for the theoretical development of permutation-based tests. However, until the 1980’s, there was limited interest in permutation-based procedures, due to lack of access to fast computers. Nowadays, the opposite is true (Albajes-Eizaguirre et al., 2019; Hemerik et al., 2019; Rao et al., 2019) and this article discusses important differences between two types of tests involving permutations or other reassignments.

## A Enforced covariate balance in a Bernoulli trial

Consider the Bernoulli trial discussed in Section 3.3. Here we illustrate how we can use this type of approach while also enforcing covariate balance. We provide a simple example where there is one (binary) covariate, say sex, and we want the percentage of women to be the same in the two treatment groups. Generalizations can also be formulated.

Suppose that  $n > 0$  is divisible by 4 and that there are  $n/2$  men and  $n/2$  women. The goal is to achieve covariate balance, which means that we want the same fraction of women in both treatment groups. One way to proceed is as follows. Randomly and indepently allocate

treatments to the  $n/2$  women. There are  $2^{n/2}$  ways to do this. Let  $l$  be the number of times that the women received treatment I. Then, randomly allocate treatment I to  $l$  of the men. In this way, covariate balance is achieved:  $l$  women and  $l$  men have received treatment I and  $n/2 - l$  men and  $n/2 - l$  women have received treatment II. In each treatment group, the percentage of women is the same (although there is a small probability that one treatment group is empty).

We now compute the total number of possible treatment allocation patterns. Given  $l$ , there are

$$\binom{n/2}{l} \binom{n/2}{l}$$

possible patterns. Hence the total number of possible patterns is

$$\sum_{l=0}^{n/2} \binom{n/2}{l} \binom{n/2}{l}. \quad (9)$$

Correspondingly, the smallest possible  $p$ -value is the inverse of this number. Note that with the common design that enforces equally large treatment groups, the total number of possible treatment allocation patterns is

$$\binom{n/2}{n/4} \binom{n/2}{n/4}.$$

This is smaller than (9), which means that the smallest possible  $p$ -value is larger than for the Bernoulli trial.

## References

- Airola, A., Pahikkala, T., Boberg, J., and Salakoski, T. Applying permutation tests for assessing the statistical significance of wrapper based feature selection. In *2010 Ninth International Conference on Machine Learning and Applications*, pages 989–994. IEEE, 2010.
- Albajes-Eizaguirre, A., Solanes, A., Vieta, E., and Radua, J. Voxel-based meta-analysis via permutation of subject images (psi): Theory and implementation for sdm. *NeuroImage*, 186:174–184, 2019.
- Anderson, M. J. and Robinson, J. Permutation tests for linear models. *Australian & New Zealand Journal of Statistics*, 43(1):75–88, 2001.
- Basu, D. Randomization analysis of experimental data: the fisher randomization test. *Journal of the American Statistical Association*, 75, 1980.
- Berger, V. W. Pros and cons of permutation tests in clinical trials. *Statistics in medicine*, 19(10):1319–1328, 2000.
- Berry, K. J., Johnston, J. E., and Mielke Jr, P. W. A chronicle of permutation statistical methods. *Cham: Springer*, 2014.
- Bi, J. and Kuesten, C. Revisiting Fisher’s ‘Lady Tasting Tea’ from a perspective of sensory discrimination testing. *Food quality and preference*, 43:47–52, 2015.
- Box, J. R. A. Fisher: The life of a scientist, 1978.
- Braun, T. M. and Feng, Z. Optimal permutation tests for the analysis of group randomized trials. *Journal of the American Statistical Association*, 96(456):1424–1432, 2001.
- Cox, D. Randomization in the design of experiments. *International Statistical Review*, 77(3):415–429, 2009.

- Edgington, E. and Onghena, P. *Randomization tests*. Chapman and Hall/CRC, 2007.
- Fan, J., Tam, P., Woude, G. V., and Ren, Y. Normalization and analysis of cDNA microarrays using within-array replications applied to neuroblastoma cell response to a cytokine. *Proceedings of the National Academy of Sciences*, 101(5):1135–1140, 2004.
- Fisher, R. A. *Statistical methods for research workers*. Oliver & Boyd, 1925.
- Fisher, R. A. *The design of experiments*. Oliver and Boyd, 1935a.
- Fisher, R. A. The logic of inductive inference. *Journal of the royal statistical society*, 98(1):39–82, 1935b.
- Fisher, R. A. “The coefficient of racial likeness” and the future of craniometry. *Journal of the Anthropological Institute of Great Britain and Ireland*, 66:57–63, 1936.
- Golland, P., Liang, F., Mukherjee, S., and Panchenko, D. Permutation tests for classification. In *International Conference on Computational Learning Theory*, pages 501–515. Springer, 2005.
- Harris, H. and Kalmus, H. The measurement of taste sensitivity to phenylthiourea (P.T.C). *Annals of Eugenics*, 15(1):24–31, 1949.
- Hemerik, J. and Goeman, J. J. False discovery proportion estimation by permutations: confidence for significance analysis of microarrays. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 80(1):137–155, 2018a.
- Hemerik, J., Solari, A., and Goeman, J. Permutation-based simultaneous confidence bounds for the false discovery proportion. *Biometrika*, 106(3):635–649, 2019.
- Hemerik, J. and Goeman, J. J. Exact testing with random permutations. *TEST*, 27(4):811–825, 2018b.
- Hoeffding, W. The large-sample power of tests based on permutations of observations. *The Annals of Mathematical Statistics*, 23:169–192, 1952.
- Imbens, G. W. and Rubin, D. B. *Causal inference in statistics, social, and biomedical sciences*. Cambridge University Press, 2015.
- Jones-Rhoades, M. W., Borevitz, J. O., and Preuss, D. Genome-wide expression profiling of the arabidopsis female gametophyte identifies families of small, secreted proteins. *PLoS genetics*, 3(10):e171, 2007.
- Kalish, L. A. and Begg, C. B. Treatment allocation methods in clinical trials: a review. *Statistics in Medicine*, 4(2):129–144, 1985.
- Kempthorne, O. and Doerfler, T. The behaviour of some significance tests under experimental randomization. *Biometrika*, 56(2):231–248, 1969.
- Lachin, J. M. Properties of simple randomization in clinical trials. *Controlled clinical trials*, 9(4):312–326, 1988a.
- Lachin, J. M. Statistical properties of randomization in clinical trials. *Controlled Clinical Trials*, 9(4):289–311, 1988b.
- Langsrud, Ø. Rotation tests. *Statistics and computing*, 15(1):53–60, 2005.
- Lehmann, E. L. and Romano, J. P. *Testing statistical hypotheses*. Springer Science & Business Media, 2005.
- Meinshausen, N. and Bühlmann, P. Lower bounds for the number of false null hypotheses for multiple testing of associations under general dependence structures. *Biometrika*, 92(4):893–907, 2005.
- Mielke, P. W. and Berry, K. J. *Permutation methods: a distance function approach*. Springer Science & Business Media, 2007.
- Morgan, K. L., Rubin, D. B., et al. Rerandomization to improve covariate balance in experiments. *The Annals of Statistics*, 40(2):1263–1282, 2012.
- Neyman, J. and Pearson, E. S. On the use and interpretation of certain test criteria for purposes of statistical inference: Part I. *Biometrika*, 20(1/2):175–240, 1928.



- Ojala, M. and Garriga, G. C. Permutation tests for studying classifier performance. *Journal of Machine Learning Research*, 11(Jun):1833–1863, 2010.
- Onghena, P. Randomization tests or permutation tests? A historical and terminological clarification. *Randomization, masking, and allocation concealment*, pages 209–227, 2018.
- Onghena, P. and Edgington, E. S. Randomization tests for restricted alternating treatments designs. *Behaviour Research and Therapy*, 32(7):783–786, 1994.
- Onghena, P. and Edgington, E. S. Customization of pain treatments: Single-case design and analysis. *The Clinical journal of pain*, 21(1):56–68, 2005.
- Pesarin, F. Some elementary theory of permutation tests. *Communications in Statistics-Theory and Methods*, 44(22):4880–4892, 2015.
- Phipson, B. and Smyth, G. K. Permutation p-values should never be zero: calculating exact p-values when permutations are randomly drawn. *Statistical applications in genetics and molecular biology*, 9(1):39, 2010.
- Pitman, E. J. G. Significance tests which may be applied to samples from any populations. *Supplement to the Journal of the Royal Statistical Society*, 4(1):119–130, 1937.
- Pocock, S. J. Allocation of patients to treatment in clinical trials. *Biometrics*, pages 183–197, 1979.
- Rao, K., Drikvandi, R., and Saville, B. Permutation and bayesian tests for testing random effects in linear mixed-effects models. *Statistics in medicine*, 2019.
- Rosenberger, W. F. and Lachin, J. M. *Randomization in clinical trials: theory and practice*. John Wiley & Sons, 2015.
- Rosenberger, W. F., Uschner, D., and Wang, Y. Randomization: The forgotten component of the randomized clinical trial. *Statistics in medicine*, 38(1):1–12, 2019.
- Rosenblatt, J. D., Benjamini, Y., Gilron, R., Mukamel, R., and Goeman, J. J. Better-than-chance classification for signal detection. *Biostatistics*, kxz035, 2019.
- Rubin, D. B. Comment: Neyman (1923) and causal inference in experiments and observational studies. *Statistical Science*, 5(4):472–480, 1990.
- Schreiber, K. and Krekelberg, B. The statistical analysis of multi-voxel patterns in functional imaging. *PLoS One*, 8(7):e69328, 2013.
- Solari, A., Finos, L., and Goeman, J. J. Rotation-based multiple testing in the multivariate linear model. *Biometrics*, 70(4):954–961, 2014.
- Southworth, L. K., Kim, S. K., and Owen, A. B. Properties of balanced permutations. *Journal of Computational Biology*, 16(4):625–638, 2009.
- Suresh, K. An overview of randomization techniques: an unbiased assessment of outcome in clinical research. *Journal of human reproductive sciences*, 4(1):8, 2011.
- Tusher, V. G., Tibshirani, R., and Chu, G. Significance analysis of microarrays applied to the ionizing radiation response. *Proceedings of the National Academy of Sciences*, 98(9):5116–5121, 2001.
- Wald, A. and Wolfowitz, J. Statistical tests based on permutations of the observations. *The Annals of Mathematical Statistics*, 15(4):358–372, 1944.
- Wei, L. and Lachin, J. M. Properties of the urn randomization in clinical trials. *Controlled clinical trials*, 9(4):345–364, 1988.
- Westfall, P. H. and Young, S. S. *Resampling-based multiple testing: Examples and methods for p-value adjustment*, volume 279. John Wiley & Sons, 1993.
- Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M., and Nichols, T. E. Permutation inference for the general linear model. *Neuroimage*, 92:381–397, 2014.
- Yates, F. Contingency tables involving small numbers and the  $\chi^2$  test. *Supplement to the Journal of the Royal Statistical Society*, 1(2):217–235, 1934.