

Manifolds of triangulations, braid groups of manifolds, and the groups Γ_n^k

Denis A. Fedoseev, Vassily O. Manturov and Igor M. Nikonov

Abstract The spaces of triangulations of a given manifold have been widely studied. The celebrated theorem of Pachner [13] says that any two triangulations of a given manifold can be connected by a sequence of bistellar moves, or Pachner moves, see also [3, 12]. In the present paper we consider groups which naturally appear when considering the set of triangulations with fixed number of simplices of maximal dimension. There are three ways of introducing this groups: the geometrical one, which depends on the metric, the topological one, and the combinatorial one. The second one can be thought of as a “braid group” of the manifold and, by definition, is an invariant of the topological type of manifold; in a similar way, one can construct the smooth version. We construct a series of groups Γ_n^k corresponding to Pachner moves of $(k - 2)$ -dimensional manifolds and construct a canonical map from the braid group of any k -dimensional manifold to Γ_n^k thus getting topological/smooth invariants of these manifolds.

Introduction

Returning back to 1954, J.W. Milnor in his first paper on link groups [11] formulated the idea that manifolds which share lots of well known invariants (homotopy type, etc.) may have different *link groups*.

Denis A. Fedoseev
Moscow State University, Moscow Center for Fundamental and Applied Mathematics, e-mail: denfedex@yandex.ru

Vassily O. Manturov
Moscow Institute of Physics and Technology, Russia; North-Eastern University, China; Kazan Federal University, Russia, e-mail: vomanturov@yandex.ru

Igor M. Nikonov
Moscow State University, e-mail: nim@mail.ru

In 2015 the second named author [9] defined a series of groups G_n^k for natural numbers $n > k$ and formulated the following principle:

➤ **The main principle**

If dynamical systems describing the motion of n particles possess a nice codimension one property depending exactly on k particles, then these dynamical systems admit a topological invariant valued in G_n^k .

We shall define braids on a manifold as loops in some configuration space related to the manifold. Following G_n^k -ideology, we mark singular configurations, while moving along a loop in the configuration space, in order to get a word — an element of a “ G_n^k -like” group Γ_n^k . Singular configurations will arise here from the moments of transformation of triangulations spanned by the configuration points; and in this case the good property of codimension one is “*k points of the configuration lie on a sphere of dimension $k - 3$ and there are no points inside the sphere*”.

In the present paper, we shall consider an arbitrary closed d -manifold M^d and for n large enough we shall construct the braid group $B_n(M^d) = \pi_1(C_n(M^d))$ (see Definitions Definition 1, Definition 2, Definition 3), where $C_n(M^d)$ is the space of generic (in some sense) configurations of points in M^d which are dense enough to span a triangulation of M . This braid group will have a natural map to the group Γ_n^{d+2} . We shall consider point configurations for which the Delaunay triangulation exists, see [1].

The idea of this work comes from the second named author, and the third named author defined the constructions of Γ_n^k , $k \geq 5$, given in Sect. 2.

Acknowledgements The authors are very grateful to S. Kim for extremely useful discussions and comments.

The first named author was supported by the Russian Foundation for Basic Research (grant No. 19-01-00775-a, grant No. 20-51-53022). The second named author was supported by the Russian Foundation for Basic Research (grant No. 20-51-53022, grant No. 19-51-51004). The third named author was supported by the Russian Foundation for Basic Research (grant No. 18-01-00398-a, grant No. 19-51-51004). The work of V.O.M. was also funded by the development program of the Regional Scientific and Educational Mathematical Center of the Volga Federal District, agreement No. 075-02-2020.

1 The Manifold of Triangulations

In the present section we define three types of manifold of triangulations: a geometrical one, a topological one, and a combinatorial one. We begin with geometrical approach, considering a manifold with a Riemannian metric.

1.1 Geometrical manifold of triangulations

Fix a smooth manifold M of dimension d with a Riemannian metric g on it. Let $n \gg d$ be a large natural number. Consider the set of all Delaunay triangulations of M with n vertices x_1, \dots, x_n (if such triangulations exist). Such Delaunay triangulations are indexed by sets of vertices, hence, the set of such triangulation forms a subset of the configuration space, which we denote by $C(M, n)$. The above subset will be an open (not necessarily connected) manifold of dimension dn : if some n pairwise distinct points (x_1, \dots, x_n) form a Delaunay triangulation then the same is true for any collection of points (x'_1, \dots, x'_n) in the neighbourhood of (x_1, \dots, x_n) . We call a set of points x_1, \dots, x_n *admissible* if such a Delaunay triangulation exists.

Hence, we get a non-compact (open) manifold M_g^{dn} . We call two sets of points (x_1, \dots, x_n) and (x'_1, \dots, x'_n) *adjacent* if there is a path $(x_{1,t}, \dots, x_{n,t})$ for $t \in [0, 1]$ such that $(x_{1,0}, \dots, x_{n,0}) = (x_1, \dots, x_n)$ and $(x_{1,1}, \dots, x_{n,1}) = (x'_1, \dots, x'_n)$ and there exists exactly one $t_0 \in [0, 1]$ such that when passing through $t = t_0$ the Delaunay triangulation $(x_{1,t}, \dots, x_{n,t})$ undergoes a flip (also called *Pachner moves* or *bistellar moves*).

The flip (Pachner move) corresponds to a position when some $d + 2$ points $(x_{i_1,t_0}, \dots, x_{i_{d+2},t_0})$ belong to the same $(d - 1)$ -sphere S^{d-1} such that no other point x_j lies inside the ball B bounded by this sphere.

Every manifold of triangulations described above has a natural stratification. Namely, every point of M_g^{dn} is given by a collection (x_1, \dots, x_n) . Such a collection is *generic* if there is no sphere S^{d-1} such that exactly $d + 2$ points among x_k belong to this sphere without any points inside the sphere.

We say that a point (x_1, \dots, x_n) is of *codimension 1*, if there exists exactly one sphere with exactly $d + 2$ points on it and no points inside it; analogously, *codimension 2 strata* correspond to either one sphere with $d + 3$ points or two spheres containing $d + 2$ points each.

In codimension 2 this corresponds to either one point of valency five in Voronoï tiling or two points of valency four in Voronoï tiling.

Hence, we have constructed a stratified (open) manifold M_g^{dn} . We call it the *geometrical manifold of triangulations*.

Note that the manifold M_g^{dn} may be not connected, i.e. there can exist non-equivalent triangulations. On the other hand, if one considers the spines of some manifold, they all can be transformed into each other by Matveev–Piergallini moves [10].

Denote the connected components of M_g^{dn} by $(M_g^{dn})_1, \dots, (M_g^{dn})_p$.

Definition 1 The *geometrical n -strand braid groups* of the manifold M_g are the fundamental groups

$$B_g(M_g, n)_j = \pi_1((M_g^{dn})_j), \quad j = 1, \dots, p.$$

1.2 Topological manifold of triangulations

The definition of geometrical manifold of triangulations heavily depends on the metric of the manifold M . For example, if we take M to be the torus glued from the square 1×10 , the combinatorial structure of the manifold of triangulations will differ from the combinatorial structure for the case of the manifold of triangulations for the case of the torus glued from the square 1×1 . Its analogue which is independent on the metric is the *topological manifold of triangulations*. We shall construct the 2-frame of this manifold. The idea is to catch all simplicial decompositions which may eventually happen for the manifold of triangulations for whatever metrics, and glue them together to get an open (non-compact) manifold.

Consider a topological manifold M^d . We consider all Riemannian metrics g_α on this manifold. They give manifolds $M_{g_\alpha}^{dn}$ as described in Sect. 1.1, which are naturally stratified. By a *generalised cell* of such a stratification we mean a connected component of the set of generic points of $M_{g_\alpha}^{dn}$.

We say that two generalised cells C_1 and C_2 are *adjacent* if there exist two points, say, $x = (x_1, \dots, x_n)$ in C_1 and $x' = (x'_1, \dots, x'_n)$ in C_2 , and a path $x_t = (x_1(t), \dots, x_n(t))$, such that $x_i(0) = x_i$ and $x_i(1) = x'_i(i)$ such that all points on this path are generic except for exactly one point, say, corresponding to $t = t_0$, which belongs to the stratum of codimension 1.

We say that two generic strata of $M_{g_\alpha}^{dn}, M_{g_\beta}^{dn}$ are *equivalent* if there is a homeomorphism of $M_{g_\alpha}^{dn} \rightarrow M_{g_\beta}^{dn}$ taking one stratum to the other.

A generalised 0-cell of the manifold of triangulations is an equivalence class of generic strata.

Analogously, we define generalised 1-cells of the manifold of triangulations as equivalence classes of pairs of adjacent vertices for different metrics $M_{g_\alpha}^{dn}$.

In a similar manner, we define generalised 2-cells as equivalence classes of discs for metrics $M_{g_\alpha}^{dn}$ such that:

1. vertices of the disc are points in 0-strata;
2. edges of the disc connect vertices from adjacent 0-strata; each edge intersects codimension 1 set exactly in one point;
3. the cycle is spanned by a disc which intersects codimension 2 set exactly at one point;
4. equivalence is defined by homeomorphism taking disc to disc, edge to an equivalent edge and vertex to an equivalent vertex and respects the stratification.

Thus, we get the 2-frame of the manifold M_{top}^{dn} . This manifold might be disconnected.

Definition 2 The *topological n -strand braid groups* of the manifold M are the fundamental groups

$$B_t(M, n)_j = \pi_1((M_{top}^{dn})_j), \quad j = 1, \dots, q.$$




Fig. 1 Codimension 1 corresponds to a flip




Fig. 2 Codimension 2 corresponds to a pentagon

1.3 Combinatorial manifold of triangulations

In Sect. 1.1 and Sect. 1.2 we constructed manifolds of triangulations basing on the notion of *Delaunay triangulation*. We did not actually discuss the *existence* of such triangulation for a given manifold and a given set of points, requesting only that the number of points is sufficiently large and points are sufficiently dense. It turns out that this condition is not always sufficient, as shows in [1]. Even for a large number of points and a Riemannian manifold with metric arbitrary close to Euclidian one, there may not exist a Delaunay triangulation.

One may impose additional restrictions on the vertex set or on the manifold itself to overcome this difficulty, but to work with the most general situation we shall consider the third notion of manifolds of triangulations: the *combinatorial manifold of triangulations* which we denote by M_{comb}^{dn} . We construct the 2-frame of the manifold M_{comb}^{dn} which is needed to get the fundamental group.

First, we fix n points on the manifold M^d and consider triangulations of the manifold with vertices in those points. Now we do not restrict ourselves to Delaunay triangulations, but consider all triangulations of the manifold. Moreover, we work with *equivalence classes* of triangulations modulo the following relation: two triangulations T_1, T_2 with (fixed) vertices v_1, \dots, v_n are said to be *equivalent* if and only

if for each pair $(i, j) \in \bar{n}$ the vertices v_i, v_j are connected by an edge of the triangulation T_1 if and only if they are connected by an edge of the triangulation T_2 . From now on saying “triangulation” we mean such equivalence class of triangulations. Those triangulations are the vertices (0-cells) of the frame of the manifold M_{comb}^{dn} .

Two triangulations are connected by an edge (a 1-cell of the frame) if and only if they differ by a *flip* — a Pachner move.

Finally, the 2-cells of the frame are chosen to correspond to the relations of the groups Γ_n^k (see Sect. 2). There are two types of such relations. One relation corresponds to the far commutativity, i.e., for any two *independent* flips α, β (flips related to different edges) we define a quadrilateral consisting of subsequent flips corresponding to $\alpha, \beta, \alpha^{-1}, \beta^{-1}$. The other relations correspond to all possible simplicial polytopes with $d+3$ vertices inscribed in the unit sphere. With each polytope of such sort, we associate a relation of length $d+3$ as shown in Sect. 2.

Definition 3 The *combinatorial n -strand braid groups* of the manifold M are the fundamental groups

$$B_c(M, n)_j = \pi_1((M_{comb}^{dn})_j), \quad j = 1, \dots, q.$$

2 The groups Γ_n^k

In the present section we consider the case of the manifold $M^d = \mathbb{R}^d$ being a vector space. The case of Euclidian space gives groups which are important for the general case. In the present paper we deal mostly with the case of $d \geq 3$. The very interesting initial case $d = 2$ is studied in [6]. We start with the case $d = 3$.

2.1 The case $d = 3$

Consider a configuration of n points in general position in \mathbb{R}^3 . We can think of these points as lying in a fixed tetrahedron $ABCD$. The points induce a unique Delaunay triangulation of the tetrahedron: four points form a simplex of the triangulation if and only if there is no other points inside the sphere circumscribed over these points. The triangulation transforms when the points move in the space.

In order to avoid degenerate Delaunay triangulations we exclude configurations where four points lie on one circle (intersection of a plane and a sphere).

Transformations of the combinatorial structure of the Delaunay triangulation correspond to configurations of codimension 1 when five points lie on a sphere which does not contain any points inside. At this moment two simplices of the triangulation are replaced with three simplices as shown in Fig. 3 (or vice versa). This transformation is called a *2-3 Pachner move*.

To trace the evolution of triangulations that corresponds to a dynamics of the points we attribute a generator to each Pachner move. For the move that replace the simplices $iklm, jklm$ with the simplices $ijkl, ijk, ijl, ijm$ in Fig. 3 we use the generator $a_{ij,klm}$. Note that 1) we can split the indices into two subsets according to the combinatorics of the transformation; 2) the generator $a_{ij,klm}$ is not expected to be involutive because it changes the number of simplices of the triangulation.




Fig. 3 A Pachner move

The relations on generators $a_{ij,klm}$ correspond to configurations of codimension 2 which occurs when either 1) six points lie on the same sphere with empty interior, or 2) there are two spheres with five points on each of them, or 3) five points on one sphere compose a codimension 1 configuration.

The last case means that the convex hull of the five points has a quadrilateral face (Fig. 4). The vertices of this face lie on one circle so we exclude this configuration.

If there are two different spheres with five points on each of them, then there is no simplex inscribed into the both spheres (otherwise its vertices would belong to one plane which contains the intersection of the spheres). Hence, the simplices inscribed into the first spheres and the simplices inscribed into the second one have no common internal points. For each sphere we can suppose that the faces of the convex hull of the inscribed simplices has only triangular faces; in the other case, that would be a configuration of codimension greater than two. So the convex hulls can have at most one common face, so we can transform them independently. This gives us a commutation relation

$$a_{ij,klm}a_{i'j',k'l'm'} = a_{i'j',k'l'm'}a_{ij,klm}, \quad (1)$$

where

$$|\{i, j, k, l, m\} \cap \{i', j', k', l', m'\}| < 4, \quad (2)$$

$$|\{i, j\} \cap \{i', j', k', l', m'\}| < 2,$$

and

$$|\{i', j'\} \cap \{i, j, k, l, m\}| < 2. \quad (3)$$

Consider now the case of six points on one sphere. The convex hull of these points is a convex polyhedron. The polyhedron must have only triangular faces; otherwise,




Fig. 4 A quadrilateral pyramid

there is an additional linear condition (four points lie on one plane) which raises the codimension beyond 2. There are two such polyhedra, see Fig. 5.




Fig. 5 Convex polyhedra with 6 vertices

For the octahedron on the left we specify the geometrical configuration assuming that the orthogonal projection along the direction ij maps the points i and j near the projection of the edge kl and the line ln is higher than km if you look from i to j (Fig. 6). Note that we live in \mathbb{R}^3 which we may consider as oriented. In this case we have six triangulations:

1. $ijkl, ijkn, ijl m, i j m n$;
2. $iklm, ikmn, jkl m, j k m n$;
3. $ikln, ilmn, jkln, j l m n$;
4. $ijkl, i j k m, i j l m, i k m n, j k m n$;
5. $ijkl, i j k n, i j l n, i l m n, j l m n$;
6. $ikln, ilmn, jkl m, j k m n, k l m n$.

The first three have four simplices each, the last three have five simplices each. The Pachner moves between the triangulations are shown in Fig. 7. Thus, we have the relation

$$a_{km,ijn} a_{ij,klm}^{-1} a_{ln,ikm} a_{km,jln}^{-1} a_{ij,kln} a_{ln,ijm}^{-1} = 1. \quad (4)$$




Fig. 6 An orthogonal projection of the octahedron




Fig. 7 Triangulation graph for the octahedron

In the case of the shifted octahedron (Fig. 5, right) we assume the line ln lies higher than the line km when one looks from the vertex i to the vertex j (Fig. 8). Then we have six triangulations:

1. $ijkl, ijk, ijm$;
2. $ijkl, ijln, ilmn, jlm$;
3. ijk, ijm, ikl, jkl ;
4. $ijk, ikln, ilmn, jkl, jlm$;
5. ijk, ikl, ikm, jkl, jkm ;
6. $ijk, ikln, ilmn, jkl, jkm, klm$.




Fig. 8 An orthogonal projection of the shifted octahedron

The Pachner moves between the triangulations are shown in Fig. 9. This gives us the relation

$$a_{ln,ijm} a_{kn,ijl} a_{km,jln} = a_{km,ijl} a_{kn,ijm} a_{ln,ikm}. \quad (5)$$

Definition 4 The group Γ_n^5 is the group with generators

$$\{a_{ij,klm} \mid \{i, j, k, l, m\} \in \bar{n}, |\{i, j, k, l, m\}| = 5\}$$

and relations




Fig. 9 Triangulation graph for the shifted octahedron

1. $a_{ij,klm} = a_{ji,klm} = a_{ij,lkm} = a_{ij,kml}$,
2. $a_{ij,klm}a_{i'j',k'l'm'} = a_{i'j',k'l'm'}a_{ij,klm}$, for

$$|\{i, j\} \cap \{i', j', k', l', m'\}| < 2, \quad |\{k, l, m\} \cap \{i', j', k', l', m'\}| < 3$$

and

$$|\{i', j'\} \cap \{i, j, k, l, m\}| < 2, \quad |\{k', l', m'\} \cap \{i, j, k, l, m\}| < 3$$

3. $a_{km,ijn}a_{ij,klm}^{-1}a_{ln,ikm}a_{km,jln}^{-1}a_{ij,klm}a_{ln,ijm}^{-1} = 1$ for distinct i, j, k, l, m, n ,
4. $a_{ln,ijm}a_{kn,ijl}a_{km,jln} = a_{km,ijl}a_{kn,ijm}a_{ln,ikm}$, for distinct i, j, k, l, m, n .

The group Γ_n^5 can be used to construct invariants of braids and dynamic systems. Consider the configuration space $\tilde{C}_n(\mathbb{R}^3)$ which consists of nonplanar n -tuples (x_1, x_2, \dots, x_n) of points in \mathbb{R}^3 such that for all distinct i, j, k, l the points x_i, x_j, x_k, x_l do not lie on the same circle.

We construct a homomorphism from $\pi_1(\tilde{C}_n(\mathbb{R}^3))$ to Γ_n^5 . Let

$$\alpha = (x_1(t), \dots, x_n(t)), \quad t \in [0, 1],$$

be a loop in $\tilde{C}_n(\mathbb{R}^3)$. For any t the set $\mathbf{x}(t) = (x_1(t), \dots, x_n(t))$ determines a Delaunay triangulation $T(t)$ of the polytope $\text{conv } \mathbf{x}(t)$. If α is in general position there will be a finite number of moments $0 < t_1 < t_2 < \dots < t_N < 1$ when the combinatorial structure of $T(t)$ changes, and for each p the transformation of the triangulation at the moment t_p will be the Pachner move on simplices with vertices i_p, j_p, k_p, l_p, m_p . We assign to this move the generator $a_{i_p j_p, k_p l_p m_p}$ or $a_{i_p j_p, k_p l_p m_p}^{-1}$ and denote $\phi(\alpha) = \prod_{p=1}^N a_{i_p j_p, k_p l_p m_p} \in \Gamma_n^5$.

Theorem 1 *The homomorphism $\phi: \pi_1(\tilde{C}_n(\mathbb{R}^3)) \rightarrow \Gamma_n^5$ is well defined.*

Proof We need to show that the element $\phi(\alpha)$ does not depend on the choice of the representative α in a given homotopy class. Given a homotopy in general position $\alpha(\tau), \tau \in [0, 1]$, of loops in $\tilde{C}_n(\mathbb{R}^3)$, the transformations of the words $\phi(\alpha(\tau))$ correspond to point configurations of codimension 2, considered above the definition of Γ_n^5 , and thus are counted by the relations in the group Γ_n^5 . Therefore, the element $\phi(\alpha(\tau))$ of the group Γ_n^5 remains the same when τ changes. \square

2.2 General case

The braids in higher dimensional Euclidean spaces were defined by the second named author in [7]. For our needs we modify that definition slightly.

Consider the configuration space $\tilde{C}_n(\mathbb{R}^{k-2})$, $4 \leq k \leq n$, consisting of n -point configurations $\mathbf{x} = (x_1, x_2, \dots, x_n)$ in \mathbb{R}^{k-2} , such that $\dim \operatorname{conv} \mathbf{x} = k - 2$ and there are no $k - 1$ points which lie on one $(k - 4)$ -dimensional sphere (the intersection of a $(k - 3)$ -sphere and a hyperplane in \mathbb{R}^{k-2}).

A configuration $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \tilde{C}_n(\mathbb{R}^{k-2})$ determines a Delaunay triangulation of $\operatorname{conv} \mathbf{x}$ which is unique when \mathbf{x} is in general position. If the vertices $x_{i_1}, \dots, x_{i_{k-1}}$ span a simplex of the Delaunay triangulation then the interior of the circumscribed sphere over these points does not contain other points of the configuration. The inverse statement is true when \mathbf{x} is generic. The condition that no $k - 1$ points lie on one $(k - 4)$ -dimensional sphere ensures that there are no degenerate simplices in the Delaunay triangulation.

Let $\alpha = (x_1(t), \dots, x_n(t))$, $t \in [0, 1]$, be a loop in $\tilde{C}_n(\mathbb{R}^{k-2})$. For any t the configuration $\mathbf{x}(t) = (x_1(t), \dots, x_n(t))$ determines a Delaunay triangulation $T(t)$ of the polytope $\operatorname{conv} \mathbf{x}(t)$. If α is generic then there will be a finite number of moments $0 < t_1 < t_2 < \dots < t_L < 1$ when the combinatorial structure of $T(t)$ changes. We shall call such configurations *singular*.

For each singular configuration $\mathbf{x}(t_i)$, either one simplex degenerates and disappears on the boundary $\partial \operatorname{conv} \mathbf{x}(t)$ or the Delaunay triangulation is not unique. That means that there is a sphere in \mathbb{R}^{k-2} which contains k points of \mathbf{x} on it and no points of \mathbf{x} inside. Assuming \mathbf{x} is generic, the span of these k points is a simplicial polytope. Below we shall count only the latter type of singular configurations.

The simplicial polytopes in \mathbb{R}^{k-2} with k vertices are described in [4]. Each of them is the join $\Delta_P * \Delta_Q$ of simplices $\Delta_P = \operatorname{conv}(x_{p_1}, \dots, x_{p_l})$ of dimension $l - 1 \geq 1$ and $\Delta_Q = \operatorname{conv}(x_{q_1}, \dots, x_{q_{k-l}})$ of dimension $k - l - 1 \geq 1$ such that the intersection $\operatorname{relint}(x_{p_1}, \dots, x_{p_l}) \cap \operatorname{relint}(x_{q_1}, \dots, x_{q_{k-l}})$ consists of one point.

We recall that the *join* of two sets $X, Y \subset \mathbb{R}^m$ is defined as $X * Y = \{\lambda x + (1 - \lambda)y \mid x \in X, y \in Y, \lambda \in [0, 1]\}$, and the *relative interior* of a finite set $X \subset \mathbb{R}^m$ is defined as

$$\operatorname{relint} X = \left\{ \sum_{x \in X} \lambda_x x \mid \forall x \lambda_x > 0, \sum_{x \in X} \lambda_x = 1 \right\}.$$

The polytope $\Delta_P * \Delta_Q$ has two triangulations:

$$T_P = \{\mathbf{x}_{P \cup Q} \setminus \{x_p\}\}_{p \in P} = \{x_{p_1} \cdots x_{p_{l-1}} x_{p_{l+1}} \cdots x_{p_l} x_{q_1} \cdots x_{q_{k-l}}\}_{i=1}^l$$

and

$$T_Q = \{\mathbf{x}_{P \cup Q} \setminus \{x_q\}\}_{q \in Q} = \{x_{p_1} \cdots x_{p_l} x_{q_1} \cdots x_{q_{l-1}} x_{q_{l+1}} \cdots x_{q_{k-l}}\}_{i=1}^{k-l}.$$

Here $P = \{p_1, \dots, p_l\}$, $Q = \{q_1, \dots, q_{k-l}\}$ and $\mathbf{x}_J = \{x_j\}_{j \in J}$ for any $J \subset \{1, \dots, n\}$.

The condition $\text{relint}(\mathbf{x}_P) \cap \text{relint}(\mathbf{x}_Q) = \{z\}$ implies $P \cap Q = \emptyset$. Thus, when the configuration $\mathbf{x}(t)$ goes over a singular value $t_i, i = 1, \dots, L$, in the Delaunay triangulation simplices T_{P_i} are replaced by simplices T_{Q_i} for some subsets $P_i, Q_i \subset \{1, \dots, n\}$, $P_i \cap Q_i = \emptyset$, $|P_i|, |Q_i| \geq 2$, $|P_i \cup Q_i| = k$. This transformation is called a *Pachner move*. The pair of subsets (P_i, Q_i) we call the *type* of the Pachner move. We assign to the transformation the letter a_{P_i, Q_i} .

Hence, the loop α produces a word

$$\Phi(\alpha) = \prod_{i=1}^L a_{P_i, Q_i} \quad (6)$$

in the alphabet $\mathcal{A}_n^k = \{a_{P, Q} \mid P, Q \subset \{1, \dots, n\}, P \cap Q = \emptyset, |P \cup Q| = k, |P|, |Q| \geq 2\}$.

Now let us consider a generic homotopy $\alpha_s, s \in [0, 1]$ between two generic loops α_0 and α_1 . A loop $\alpha_s = \{\mathbf{x}(s, t)\}_{t \in [0, 1]}$ can contain a configuration of codimension 2. This means that for some t the configuration $\mathbf{x}(s, t)$ has two different k -tuples of points, such that each of them lies on a sphere whose interior contains no points of $\mathbf{x}(s, t)$. If these spheres do not coincide then their intersection contains at most $k - 2$ points (the intersection can not contain $k - 1$ points because $\mathbf{x}(s, t) \in \tilde{C}_n(\mathbb{R}^{k-2})$). Hence, the simplices involved in one Pachner move can not be involved in the other one, so the Pachner moves can be fulfilled in any order.

If the k -tuples of points lie on one sphere, there is a sphere with $k + 1$ points of $\mathbf{x}(s, t)$ on it and its interior contains no points of $\mathbf{x}(s, t)$. These $k + 1$ points span a simplicial polytope in \mathbb{R}^{k-2} . Such polytopes are described in [4]. The description uses the notion of *Gale transform*.

Let $X = \{x_1, \dots, x_n\}$ be a set of n points in \mathbb{R}^d such that $\dim \text{conv } X = d$. Then $n \geq d + 1$. Let $x_i = (x_{1i}, \dots, x_{di}) \in \mathbb{R}^d, i = 1, \dots, n$, be the coordinates of the points of X . The matrix

$$M = \begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ x_{d1} & x_{d2} & \cdots & x_{dn} \\ 1 & 1 & \cdots & 1 \end{pmatrix}$$

has rank $d + 1$. Then the dimension of the space $\ker M = \{b \in \mathbb{R}^n \mid Mb = 0\}$ of dependencies between columns of M is equal to $n - (d + 1)$. Take any basis $b_j = (b_{j1}, b_{j2}, \dots, b_{jn}), j = 1, \dots, n - d - 1$, of $\ker M$ and write it in matrix form

$$B = \begin{pmatrix} b_{11} & \cdots & b_{1n} \\ \cdots & \cdots & \cdots \\ b_{n-d-1,1} & \cdots & b_{n-d-1,n} \end{pmatrix}. \quad (7)$$

The columns of the matrix B form a set $Y = y_1, \dots, y_n, y_i = (b_{1i}, \dots, b_{n-d-1,i})$, in \mathbb{R}^{n-d-1} . The set Y is called a *Gale transform* of the point set X . Gale transforms which correspond to different bases of $\ker M$ are linearly equivalent. The vectors of the Gale transform Y may coincide.

Let $Y = y_1, \dots, y_n$ be a Gale transform of X . The set

$$\bar{Y} = \{\bar{y}_1, \dots, \bar{y}_n\}, \quad \bar{y}_i = \begin{cases} \frac{y_i}{\|y_i\|}, & y_i \neq 0, \\ 0, & y_i = 0 \end{cases}$$

is called a *Gale diagram* of the point set X . It is a subset of $S^{n-d-2} \cup \{0\}$.

Denote $N = \{1, \dots, n\}$. Two subsets $\bar{Y} = \{\bar{y}_1, \dots, \bar{y}_n\}$ and $\bar{Y}' = \{\bar{y}'_1, \dots, \bar{y}'_n\}$ in $S^{n-d-2} \cup \{0\}$ are called *equivalent* if there is a permutation σ of N such that for any $J \subset N$

$$0 \in \text{relint } \bar{Y}_J \iff 0 \in \text{relint } \bar{Y}'_{\sigma(J)}.$$

Here we denote $\bar{Y}_J = \{\bar{y}_i\}_{i \in J}$ and $\bar{Y}'_J = \{\bar{y}'_i\}_{i \in J}$.

The properties of Gale diagrams can be summarized as follows [4].

- Theorem 2** 1. Let X be a set of n points which are vertices of some polytope P in \mathbb{R}^d and \bar{Y} be its Gale diagram. Then the set of indices $J \subset N$ defines a face of P if and only if $0 \in \text{relint } \bar{Y}_{N \setminus J}$.
2. Let X and X' be sets of vertices of polytopes P and P' , $|X| = |X'|$, and \bar{Y} and \bar{Y}' be their Gale diagrams. Then P and P' are combinatorially equivalent if and only if \bar{Y} and \bar{Y}' are equivalent.
3. For any n -point set $\bar{Y} \in S^{n-d-2} \cup \{0\}$ such that \bar{Y} spans \mathbb{R}^{n-d-1} and 0 lies in the interior of $\text{conv } \bar{Y}$, there is an n -point set X in \mathbb{R}^d such that \bar{Y} is a Gale diagram of X .

The theorem implies (see [4]) that simplicial polytopes with $k+1$ vertices in \mathbb{R}^{k-2} are in a bijection with standard Gale diagrams in \mathbb{R}^2 (defined uniquely up to isometries of the plane).

A *standard Gale diagram* of order $l = k+1$ is a subset \bar{Y} , $|\bar{Y}| = l$, of the vertices set $\{e^{\frac{2\pi i p}{l}}\}_{p=0}^{2l-1}$ of the regular $2l$ -gon inscribed in the unit circle S^1 , such that:

1. any diameter of S^1 contains at most one point of \bar{Y} ;
2. for any diameter of S^1 , any open half-plane determined by it contains at least two points of \bar{Y} .

The first property means the corresponding polytope is simplicial, the second means any of the $k+1$ vertices of the polytope is a face.

The number c_l of standard Gale diagrams of order l is equal to

$$c_l = 2^{\lfloor \frac{l-3}{2} \rfloor} - \left\lfloor \frac{l+1}{4} \right\rfloor + \frac{1}{4l} \sum_{h: 2 \nmid h|l} \varphi(h) \cdot 2^{\frac{l}{h}},$$

where $l = 2^{a_0} \prod_{i=1}^t p_i^{a_i}$ is the prime decomposition of l , and φ is Euler's function. For small l the numbers are $c_5 = 1, c_6 = 2, c_7 = 5$, see Fig. 10.

Let us describe the triangulations of the simplicial polytopes with $k+1$ vertices in \mathbb{R}^{k-2} .

Let $X = \{x_1, \dots, x_n\}$ be a subset in \mathbb{R}^d , so $x_i = (x_{i1}, \dots, x_{id})$, $i = 1, \dots, n$. Let $P = \text{conv } X$ be the convex hull of X , assume that $\dim P = d$. A triangulation T of the polytope P with vertices in X is called *regular* if there is a height function $h: X \rightarrow \mathbb{R}$ such that T is the projection of the lower convex of the lifting $X^h = \{x_1^h, \dots, x_n^h\} \subset$




Fig. 10 Standard Gale diagrams of small order

\mathbb{R}^{d+1} , where $x_i^h = (x_i, h(x_i))$. This means that a set of indices $J \subset \{1, \dots, n\}$ determines a face of T if and only if there exists a linear functional ϕ on \mathbb{R}^{d+1} such that $\phi(0, \dots, 0, 1) > 0$ and $J = \{i \mid \phi(x_i^h) = \min_{x^h \in X^h} \phi(x^h)\}$. In case T is regular we write $T = T(X, h)$. Any generic height function induces a regular triangulation.

The Delaunay triangulation of X is regular with the height function $h: \mathbb{R}^d \rightarrow \mathbb{R}$, $h(z) = \|z\|^2 = \sum_{i=1}^d z_i^2$ if $z = (z_1, \dots, z_d) \in \mathbb{R}^d$.

A height function $h: X \rightarrow \mathbb{R}$ can be regarded as a vector $h = (h_1, \dots, h_n) \in \mathbb{R}^n$ where $h_i = h(x_i)$. Denote $\beta(h) = Bh \in \mathbb{R}^{n-d-1}$ where B is the matrix (7) used to define a Gale transform of X . Let $\bar{Y} = \{\bar{y}_1, \dots, \bar{y}_n\}$ be a Gale diagram of X . Convex cones generated by the subsets of \bar{Y} split the space \mathbb{R}^{n-d-1} into a union of conic cells. A relation between the triangulation $T(X, h)$ and the Gale diagram \bar{Y} can be described as follows.

- Theorem 3** 1. If $T(X, h)$ is a regular triangulation of X then $\beta(h)$ belongs to a conic cell of maximal dimension in the splitting of \mathbb{R}^{n-d-1} induced by \bar{Y} .
2. Let $J \subset N = \{1, \dots, n\}$. The set X_J spans a cells of the triangulation $T(X, h)$ if and only if $\beta(h) \in \text{concone}(\bar{Y}_{N \setminus J})$, where $\text{concone}(X_{N \setminus J})$ is the convex cone spanned by the set $\bar{Y}_{N \setminus J}$.

Let P be a simplicial polytopes with $l = k+1$ vertices in \mathbb{R}^{l-3} and $\bar{Y} = \{\bar{y}_1, \dots, \bar{y}_l\}$ be the corresponding standard Gale diagram. By Theorem 3 there are l different regular triangulation which correspond to open sectors between the rays spanned by the vectors of \bar{Y} . The graph whose vertices are combinatorial classes of triangulations of P and the edges are Pachner moves, is a cycle. Let us find which Pachner moves appear in this cycle.

We change the order of vertices of P (and hence, the order of the points of \bar{Y}) so that the points $\bar{y}_1, \dots, \bar{y}_l$ appear in this sequence when one goes counterclockwise on the unit circle. For each i denote $R_{\bar{Y}}(i)$ (respectively, $L_{\bar{Y}}(i)$) be the set of indices j of vectors \bar{y}_j that lie in the right (respectively, left) open half-plane incident to the oriented line spanned by the vector \bar{y}_i . Then the Pachner move which occurs when the vector $\beta(h)$ passes \bar{y}_i from right to left, will be marked with the letter

$a_{R_{\bar{Y}}(i), L_{\bar{Y}}(i)} \in \mathcal{A}_l^{l-1}$. The Pachner moves of the whole cycle of triangulation give the word $w_{\bar{Y}} = \prod_{i=1}^l a_{R(i), L(i)}$.

Thus, we have the relation $w_{\bar{Y}} = 1$ that we should impose in order to make the words like $\Phi(\alpha)$ independent on resolutions of configurations of codimension 2.

Example 1 Consider the standard Gale diagram of order 5, Fig. 11. Then we have $R(1) = \{4, 5\}$, $L(1) = \{2, 3\}$, $R(2) = \{1, 5\}$, $L(2) = \{3, 4\}$ etc. The corresponding word is equal to

$$w = a_{45,23}a_{15,34}a_{12,45}a_{23,15}a_{34,12}.$$




Fig. 11 Standard Gale diagram of order 5

We can give now the definition of Γ_n^k groups.

Definition 5 Let $4 \leq k \leq n$. The group Γ_n^k is the group with the generators

$$a_{P,Q}, \quad P, Q \subset \{1, \dots, n\}, P \cap Q = \emptyset, |P \cup Q| = k, |P|, |Q| \geq 2,$$

and the relations:

1. $a_{Q,P} = a_{P,Q}^{-1}$;
2. *far commutativity*: $a_{P,Q}a_{P',Q'} = a_{P',Q'}a_{P,Q}$ for each generators $a_{P,Q}$, $a_{P',Q'}$ such that

$$\begin{aligned} |P \cap (P' \cup Q')| &< |P|, & |Q \cap (P' \cup Q')| &< |Q|, \\ |P' \cap (P \cup Q)| &< |P'|, & |Q' \cap (P \cup Q)| &< |Q'|; \end{aligned}$$

3. $(k+1)$ -gon relations: for any standard Gale diagram \bar{Y} of order $k+1$ and any subset $M = \{m_1, \dots, m_{k+1}\} \subset \{1, \dots, n\}$

$$\prod_{i=1}^{k+1} a_{M_R(\bar{Y}, i), M_L(\bar{Y}, i)} = 1,$$

where $M_R(\bar{Y}, i) = \{m_j\}_{j \in R_{\bar{Y}}(i)}$, $M_L(\bar{Y}, i) = \{m_j\}_{j \in L_{\bar{Y}}(i)}$.

Example 2 Let us write the $(k+1)$ -gon relations in Γ_n^k for small k .

The group Γ_n^4 has one pentagon relation

$$a_{m_4 m_5, m_2 m_3} a_{m_1 m_5, m_3 m_4} a_{m_1 m_2, m_4 m_5} a_{m_2 m_3, m_1 m_5} a_{m_3 m_4, m_1 m_2} = 1.$$

Our definition of Γ_n^4 differs slightly from the definition of these groups in [6]. To obtain the groups defined in [6] we should add relations $a_{m_1 m_2, m_3 m_4}^2 = 1$ and

$$\begin{aligned} a_{m_1 m_2, m_3 m_4} a_{m_1 m_2, m_5 m_6} &= a_{m_1 m_2, m_5 m_6} a_{m_1 m_2, m_3 m_4}, \\ a_{m_1 m_2, m_3 m_4} a_{m_1 m_5, m_2 m_6} &= a_{m_1 m_5, m_2 m_6} a_{m_1 m_2, m_3 m_4}, \end{aligned}$$

where $|\{m_3, m_4\} \cap \{m_5, m_6\}| \leq 1$.

The group Γ_n^5 has two hexagon relations

$$\begin{aligned} a_{m_5 m_6, m_2 m_3 m_4} a_{m_1 m_5 m_6, m_3 m_4} a_{m_1 m_2, m_4 m_5 m_6} a_{m_1 m_2 m_3, m_5 m_6} a_{m_3 m_4, m_1 m_2 m_6} a_{m_3 m_4 m_5, m_1 m_2} &= 1, \\ a_{m_5 m_6, m_2 m_3 m_4} a_{m_1 m_5 m_6, m_3 m_4} a_{m_1 m_2 m_6, m_4 m_5} a_{m_1 m_2 m_3, m_5 m_6} a_{m_3 m_4, m_1 m_2 m_6} a_{m_4 m_5, m_1 m_2 m_3} &= 1 \end{aligned}$$

as we have seen in the previous subsection.

Theorem 4 *Formula (6) defines a correct homomorphism*

$$\Phi: \pi_1(\tilde{C}_n(\mathbb{R}^{k-2})) \rightarrow \Gamma_n^k.$$

Proof We need to show that the element $\Phi(\alpha)$ does not depend on the choice of the representative α in a given homotopy class. Given a generic homotopy $\alpha(\tau)$, $\tau \in [0, 1]$, of loops in $\tilde{C}_n(\mathbb{R}^{k-2})$, the transformations of the words $\Phi(\alpha(\tau))$ correspond to point configurations of codimension 2, considered above the definition of Γ_n^k , and thus are counted by the relations in the group Γ_n^k . Therefore, the element $\Phi(\alpha(\tau))$ of the group Γ_n^k remains the same when τ changes. \square

3 Main results

3.1 Γ_n^k -valued invariant of braids on manifolds

Theorem 5 *Let (M^d, g) be a manifold with a Riemannian metric and $B(M_g, n)_j$, $j = 1, \dots, p$ be its geometrical braid groups. Then for any j there is a well defined mapping $B(M_g, n)_j \rightarrow \Gamma_n^{d+2}$.*

Proof By Definition 1, the space of geometrical braids on the manifold M^d is the fundamental group of the manifold of triangulations: $B(M_g, n)_j = \pi_1((M_g^{dn})_j)$. Hence, we need to construct a mapping from homotopy classes of loops in $(M_g^{dn})_j$ into the group Γ_n^{d+2} .

Let us fix a number j and consider a loop $\alpha = (x_1(t), \dots, x_n(t))$, $t \in [0, 1]$ in $(M_g^{dn})_j$, where for each $i = 1, \dots, n$ the point $x_i(t) \in M^d$. For each t the collection of points $(x_1(t), \dots, x_n(t))$ defines a Delaunay triangulation $T(t)$ of the manifold

M. If the loop α is in general position, then there is a finite number of moments $0 < t_1 < \dots < t_N < 1$ when the combinatorial structure of the triangulation $T(t)$ changes. Each of those moments t_p corresponds to a codimension 1 configuration of points of the manifold M^d , and at each of them the triangulation undergoes some Pachner move, which transforms the triangulation simplices T_{P_p} into simplices T_{Q_p} , where $P_p, Q_p \subset \{1, \dots, n\}$, $P_p \cap Q_p = \emptyset$, $|P_p|, |Q_p| \geq 2$ and $|P_p \cup Q_p| = d + 2$. To each of those moments we attribute the element a_{P_p, Q_p} of the group Γ_n^{d+2} .

Thus a loop α produces a word $\Phi(\alpha) = \prod_{p=1}^N a_{P_p, Q_p}$.

Now we need to prove that this mapping $\Phi: \pi((M_g^{dn})_j) \rightarrow \Gamma_n^{d+2}$ is well defined: the element $\Phi(\alpha)$ does not depend on the choice of a representative in a given homotopy class.

Let $\alpha(\tau)$ be a generic homotopy, where $\tau \in [0, 1]$ and $\alpha(0) = \alpha$. With it we associate a family of words $\Phi(\alpha(\tau)) \in \Gamma_n^{d+2}$. Note that the word $\Phi(\alpha(\tau))$ changes whenever the loop passes through a point of codimension 2 in the manifold of triangulations M_g^{dn} . But such configurations of points in M^d exactly correspond to the relations of the group Γ_n^{d+2} . Therefore the element $\Phi(\alpha(\tau))$ of the group Γ_n^{d+2} remains the same when τ changes. \square

Quite analogously, we get the following

Theorem 6 *Let M^d be a smooth manifold of dimension d and $B(M, n)_j$, $j = 1, \dots, q$, be its topological braid groups. Then for any j there is a well defined mapping $B(M, n)_j \rightarrow \Gamma_n^{d+2}$.*

Proof By Definition 2, the topological braids group on the manifold M^d is the fundamental group of the manifold of triangulations: $B(M, n)_j = \pi_1((M_{top}^{dn})_j)$.

Consider a loop γ in the manifold M_{top}^{dn} and let γ be in general position. There is a finite number of intersections $\mathbf{x}_1, \dots, \mathbf{x}_k$ between the loop γ and the codimension 1 stratum of the manifold of triangulations, where the triangulation undergoes a flip. Consider an intersection point \mathbf{x}_i and its neighbourhood U . The intersection point \mathbf{x}_i is a set of points $x_{1,i}, \dots, x_{n,i} \in M$.

Consider two metrics g_α and g_β on the manifold M , yielding a structure of geometrical manifold of triangulations on the neighbourhood U . Let us denote the neighbourhood U equipped with the corresponding metric by U_{g_α} and U_{g_β} , respectively. As it was described in Sect. 1, equivalence of strata of the manifolds of triangulation $M_{g_\alpha}^{dn}$ and $M_{g_\beta}^{dn}$ is defined by an existence of a homeomorphism $h: M_{g_\alpha}^{dn} \rightarrow M_{g_\beta}^{dn}$ with certain properties.

The arc $l = \gamma \cap U \subset M_{top}^{dn}$ gives rise to an arc $l_1 \subset U_{g_\alpha}$ and an arc $l_2 \subset U_{g_\beta}$, and $l_2 = h(l_1)$ where h is the homeomorphism from the definition of cell equivalence. Likewise, the intersection point $X_i \in U$ gives a point $X'_i \in U_{g_\alpha}$ and a point $X''_i \in U_{g_\beta}$. Naturally, those points are the intersections between the codimension 1 stratum of the corresponding manifold and the arcs l_1, l_2 , respectively.

At the point \mathbf{x}_i a certain Pachner move occurs. The type of this move is independent of the choice of metric on the manifold. Therefore, the Pachner moves occurring in the points $\mathbf{x}'_i, \mathbf{x}''_i$ of the manifolds $U_{g_\alpha}, U_{g_\beta}$ are one and the same. Choosing the

numbering of the n points which form the vertices of the triangulations, we hence obtain the same generator of the group Γ_n^{d+2} corresponding to this intersection point. Therefore the letter which appears in the word $\Phi(\gamma)$ does not depend on the choice of the metric.

Therefore the mapping Φ is defined on the fundamental group of the topological manifold of triangulations M_{top}^{dn} and due to Theorem 5 it does not depend on the choice of a representative in the homotopy class of a loop. Hence, it is a well defined mapping from $B(M, n)_j$ to the group Γ_n^{d+2} . \square

Hence, we get invariants of smooth metrical manifolds and of topological manifolds: these invariants are the corresponding images of the fundamental group of the manifold of triangulations in the corresponding group Γ_n^k .

3.2 Triangulations of polyhedra and groups Γ_n^k

Triangulations of a given convex polyhedra are well studied. If we deal with stable triangulations, the set of those form a simplicial complex [3]. In the present section we show that they can be treated as elements of a certain group provided that some fixed triangulation plays the role of the unit element.

Let $\mathbf{x} = \{x_1, \dots, x_n\} \subset \mathbb{R}^d$, $n > d$, be the vertex set of a polytope W in general position, namely, any $d + 1$ points of \mathbf{x} do not belong to one hyperplane. Let \mathcal{T} be the set of regular triangulations of W (see page 13). We can consider \mathcal{T} as a graph whose vertices are regular triangulations and whose edges are Pachner moves. This graph is connected, see [10].

Choose a regular triangulation $T_0 \in \mathcal{T}$. For any other triangulation $T \in \mathcal{T}$ there is a path $\gamma = e_1 e_2 \dots e_l$ from T_0 to T where each edge e_i is a Pachner move of type (P_i, Q_i) , see page 12. We assign the word $\varphi(T) = \prod_{i=1}^l a_{P_i, Q_i} \in \Gamma_n^{d+2}$ to the triangulation T .

Let $\text{Cay}(\Gamma_n^{d+2})$ be the Cayley graph of the group Γ_n^{d+2} corresponding to the group presentation given in Definition 5.

Theorem 7 *The correspondence $T \mapsto \varphi(T)$ defines an embedding of the graph \mathcal{T} into the graph $\text{Cay}(\Gamma_n^{d+2})$.*

Proof We must show first that the map φ is well-defined, i.e. that the element $\varphi(T)$ does not depend on the choice of a path connecting T_0 with T .

It follows from the theory of I.M. Gelfand, M.M. Kapranov and A.V. Zelevinsky [3, Chapter 7] that the graph \mathcal{T} is the edge graph of some polytope $\Sigma(W)$ (called the *secondary polytope* of W), see Fig. 12. The faces of $\Sigma(W)$ correspond to triangulations which coincide everywhere except for divisions of two subpolytopes with $d + 2$ vertices or a subpolytope with $d + 3$ vertices in W . Hence, the faces give exactly the relations in Γ_n^{d+2} which appear in Definition 5. Thus, two different paths γ and γ' , connecting T_0 with some triangulation T , produce words $\varphi(\gamma)$ and $\varphi(\gamma')$ which differ by relations in Γ_n^{d+2} , so $\varphi(\gamma) = \varphi(\gamma') \in \Gamma_n^{d+2}$.




Fig. 12 A pentagon and its secondary polytope.

In order to proof that φ is injective we construct a left inverse map to it. A *formal l -dimensional simplex* is any subset $P \subset \{1, \dots, n\}$, $|P| = l + 1$. Let $C_l(n)$ be the linear space over \mathbb{Z}_2 whose basis consists of all the l -dimensional formal simplices. We can identify $C_l(n) \simeq \mathbb{Z}_2^{\binom{n}{l+1}}$ with the space of l -chains of the simplicial complex for a $(n - 1)$ -dimensional simplex whose vertices are marked with numbers $1, 2, \dots, n$. The differential $\partial: C_l(n) \rightarrow C_{l-1}(n)$ of this complex is defined by the formula $\partial(P) = \sum_{p \in P} P \setminus \{p\}$.

Any triangulation T of W defines a d -chain $c(T) = \sum_{P: x_P \in T} P$ such that $\partial c(T) = \partial W$ where ∂W is the formal sum of $(d - 1)$ -dimensional faces forming the boundary of the polytope W . The triangulation T can be restored from its chain $c(T)$ as the support of the chain. Thus, we can identify triangulations of W with a subset of $C_d(n)$.

If a triangulation T' differs from a triangulation T by applying a Pachner move of type (P, Q) then $c(T') - c(T) = \partial(P \cup Q)$. Define a homomorphism $\psi: \Gamma_n^{d+2} \rightarrow C_d(n)$ by the formula

$$\psi(a_{P,Q}) = \partial(P \cup Q) = \sum_{i \in P \cup Q} (P \cup Q) \setminus \{i\}.$$

The map ψ annihilates the relations of Γ_n^{d+2} because they come from cycles of triangulations of some polytope with $d + 3$ vertices (or two polytopes with $d + 2$ vertices). The image of a relation is equal to the difference of the chains for the final triangulation and the initial one. But these triangulations coincide so the image is zero. Thus, ψ defines a correct homomorphism from Γ_n^{d+2} to $C_d(n) \simeq \mathbb{Z}_2^{\binom{n}{d+1}}$.

For any regular triangulation T consider a path $\gamma = e_1 e_2 \dots e_l$ which connects T_0 with T in \mathcal{T} . Any edge e_i of γ corresponds to a Pachner move of type (P_i, Q_i) . Then

$$c(T) - c(T_0) = \sum_{i=1}^l \partial(P_i \cup Q_i) = \sum_{i=1}^l \psi(a_{P_i, Q_i}) = \psi(\varphi(T)).$$

Thus, the element $\varphi(T)$ determines the chain $c(T)$, hence, it uniquely determines the triangulation T . \square

Example 3 Let $W \subset \mathbb{R}^2$ be a convex n -gon. Its secondary polytope $\Sigma(W)$ is the associahedron (Stasheff polytope) K_{n-1} [14], see Fig. 13. Thus, we have the following statement:




Fig. 13 Associahedron K_5 .

Corollary 1 *The graph $\text{Cay}(\Gamma_n^4)$ contains a subgraph isomorphic to the edge graph of the associahedron K_{n-1} .*

4 Other developments: the groups $\tilde{\Gamma}_n^k$

In the present section we provide constructions resembling the Γ_n^k groups but different in some aspects, which prove useful in certain situations where the Γ_n^k groups are not sufficient. Geometrically speaking, here we consider oriented triangulations. Therefore, the indices of the generators of the groups are not independent and do not freely commute as was seen, for example, in the groups Γ_n^5 (see Definition 4, first relation).

To be precise, we introduce the following:

Definition 6 Let $5 \leq k \leq n$. The group $\tilde{\Gamma}_n^k$ is the group with the generators

$$\{a_{P,Q} \mid P, Q \text{ — cyclically ordered subsets of } \{1, \dots, n\}, P \cap Q = \emptyset, |P \cup Q| = k, |P|, |Q| \geq 2\},$$

and the following relations:

- 1,2,3. the relations 1, 2, 3 from Definition 5;
4. $a_{Q,P} = a_{Q',P'}$, where $Q = Q', P = P'$ as unordered sets, and as cyclically ordered sets Q differs from Q' by one transposition and P differs from P' by one transposition.

These groups have the same connection to geometry and dynamics as the groups Γ_n^k defined above. To illustrate that, consider the following dynamical system.




Fig. 14 A movement of a point around the configuration of four points on one circle

Example 4 Let us have a dynamical system describing a movement of a point around the configuration of four points on one circle, see Fig. 14. Such system may be presented as the word in the group $\tilde{\Gamma}_6^5$

$$w = a_{35,164} a_{46,253}^{-1} a_{46,135} a_{35,246}^{-1}.$$

We can show that $\phi(\alpha)$ is nontrivial in the abelianisation $(\tilde{\Gamma}_6^5)_{ab} = \tilde{\Gamma}_6^5 / [\tilde{\Gamma}_6^5, \tilde{\Gamma}_6^5]$ of the group $\tilde{\Gamma}_6^5$. Computer calculations show that the group $(\tilde{\Gamma}_6^5)_{ab}$ can be presented as the factor of a free commutative group with 120 generators modulo $2 \cdot 6! = 1440$ relations that span a space of rank 90 if we work over \mathbb{Z}_2 . Adding the element w to the relations increases the rank to 91, so the element w is nontrivial in $(\tilde{\Gamma}_6^5)_{ab}$, therefore it is nontrivial in $\tilde{\Gamma}_6^5$.


Hence, we have encountered a peculiar new effect in the behaviour of $\tilde{\Gamma}_n^5$ which is not the case of G_n^k . Certainly, the abelianisation of G_n^k is non-trivial and very easy to calculate since any generator enters each relation evenly many times. However, this happens not only for relations but for any words which come from braids. Thus, any abelianisations are trivial in interesting cases. For $\tilde{\Gamma}_n^k$, it is an interesting new phenomenon, and the invariants we have demonstrated so far are just the tip of the iceberg to be investigated further.



Note that both for the groups Γ_n^k and $\tilde{\Gamma}_n^k$ we have many invariants since the corank of those groups is big.


References


1. J. D. Boissonnat, R. Dyer, A. Ghosh, N. Martynchuk, “An obstruction to Delaunay triangulations in Riemannian manifolds”, *Discrete Comput. Geom.* (2018) 59:1, pp. 226–237.
2. V. M. Buchstaber, T. E. Panov, “Toric Topology”, Mathematical Surveys and Monographs, vol. 204, *Amer. Math. Soc., Providence, RI*, 2015.


3. I. M. Gelfand, M. M. Kapranov, A. V. Zelevinsky, “Discriminants, Resultants, and Multidimensional Determinants”, *Birkhäuser Boston*, 1994.
4. B. Grünbaum, “Convex polytopes”, 2nd ed., *Springer-Verlag*, 2003.
5. A. J. Kanel-Belov, “Sensationally non-planar interlocking structures”, Designing non-traditional materials based on geometrical principles (19-22 June 2005, Hanover, Germany), eds. Y. Estrin, Abstracts, Workshop in Hanover, 2005, 12
6. S. Kim, V. O. Manturov, “Artin’s braids, Braids for three space, and groups Γ_n^4 and G_n^k ”, arXiv:1902.11238
7. V. O. Manturov, “The Groups G_n^k and fundamental groups of configuration spaces”, *J. Knot Theory & Ramifications*, 2015, Vol. 26
8. V. O. Manturov, “The groups G_{k+1}^k and fundamental groups of configuration spaces”, *Proceedings of the third international congress on algebra and combinatorics (E. I. Zelmanov, K. P. Shum, P. S. Kolesnikov, Editors)*, World Scientific, Singapore (2019), pp. 310–323
9. V. O. Manturov, I. M. Nikonov, “On braids and groups G_n^k ”, *J. Knot Theory & Ramifications*, 2015, Vol. 24, No. 13.
10. S. V. Matveev, “Algorithmic topology and classification of 3-manifolds”, *Springer*, 2007.
11. J. Milnor, “Link groups”, *Annals of Mathematics*, 1954, 59(2), 177–195.
12. A. Nabutovsky, “Fundamental group and contractible closed geodesics”, *Comm. Pure Appl. Math.* 49(12) (1996), 1257–1270.
13. U. Pachner, “P.L. homeomorphic manifolds are equivalent by elementary shellings”, *European Journal of Combinatorics* (1991), 12 (2): 129–145.
14. J. D. Stasheff, “Homotopy associativity of H-spaces. I, II”, *Transactions of the American Mathematical Society* (1963), 108: 293–312.











y_4

y_2

y_5

y_1

y_3

