

Surveying the reach and maturity of machine learning and artificial intelligence in astronomy

Christopher J. Fluke^{1,2} and Colin Jacobs²

¹Advanced Visualisation Laboratory, DRICP, Swinburne University of Technology,
Hawthorn, Victoria, 3122, Australia, cfluke@swin.edu.au

²Centre for Astrophysics & Supercomputing, Swinburne University of Technology,
Hawthorn, Victoria, 3122, Australia

January 17, 2022

Abstract

Machine learning (automated processes that learn by example in order to classify, predict, discover or generate new data) and artificial intelligence (methods by which a computer makes decisions or discoveries that would usually require human intelligence) are now firmly established in astronomy. Every week, new applications of machine learning and artificial intelligence are added to a growing corpus of work. Random forests, support vector machines, and neural networks (artificial, deep, and convolutional) are now having a genuine impact for applications as diverse as discovering extrasolar planets, transient objects, quasars, and gravitationally-lensed systems, forecasting solar activity, and distinguishing between signals and instrumental effects in gravitational wave astronomy. This review surveys contemporary, published literature on machine learning and artificial intelligence in astronomy and astrophysics. Applications span seven main categories of activity: classification, regression, clustering, forecasting, generation, discovery, and the development of new scientific insight. These categories form the basis of a hierarchy of maturity, as the use of machine learning and artificial intelligence emerges, progresses or becomes established.

1 Introduction

Astronomy has a rich history of data gathering and record keeping [Brunner et al., 2002, Feigelson and Babu, 2012, Jäschek, 1968, 1978, Zhang and Zhao, 2015]. Data about and from celestial objects is collected using an assortment of telescopes, photon detectors and particle detectors. While all of the electromagnetic spectrum is of interest, the bulk of observational data comes from the visible/infrared (wavelengths from 400 nm to 1 mm) and radio (wavelengths from 1 cm to 1 km) portions of the spectrum. Much of this data is recorded in the form of two-dimensional pixel-based images and one-dimensional spectra. Secondary data products are derived from observational data, often as catalogues of individual source properties: position, size, mass, chemical composition, and so forth. Observational data can be recorded and analyzed at a single epoch, or the properties of astronomical sources – especially brightness and position – can be monitored over time.

Complementing observational data gathering are the dual fields of numerical simulation and astrophysical theory, although there is a great deal of overlap between the two. While some branches of theory do not in themselves produce large quantities of data, focusing instead on mathematical descriptions of

cosmic phenomena, computer simulations generate data that can be used to model, predict, and support analysis of the observational data.

Modern astronomical data is measured in Terabytes, Petabytes and, soon, Exabytes. When astronomy crossed the 100 Terabyte scale near the end of the 20th century [e.g., Brunner et al., 2002, Szalay and Gray, 2001] a new data-driven astronomy emerged: where data mining was mooted as the likely future [Bell et al., 2009, Ivezić et al., 2014, Szalay and Gray, 2006]. The expectation was an increased reliance on automated systems to locate, classify, and characterise objects. At the same time, new fundamental relationships between derived properties might be found, by allowing clever algorithms to search through complex, multi-dimensional data catalogues [e.g. Graham et al., 2013].

1.1 Data mining

The background and early history of data mining in astronomy is covered in some detail by Borne [2009], Ball and Brunner [2010], and the collection of articles in Way et al. [2012]. An early emphasis of data mining was to find new samples of rare sources, by applying workflows that gathered data from large, often online, repositories. Lépine and Shara [2005] applied a software blink comparator to 615,800 sub-fields downloaded from the Digitized Sky Survey, identifying and cataloging 61,977 stars with high proper motions ($\text{Dec } j \ 0^\circ$). Targeting the planned SkyMapper [Keller et al., 2007] Southern Sky Survey, Walsh et al. [2007] mined the Sloan Digitized Sky Survey (SDSS) Data Release 5 [DR5; Adelman-McCarthy et al., 2006] for stellar over-densities, uncovering a new Milky Way dwarf galaxy satellite (Boötes II). González-Solares et al. [2008] opened up their INT/WFC Photometric $\text{H}\alpha$ Survey of the Northern Galactic Plane (IPHAS) dataset through the AstroGrid Virtual Observatory Desktop,¹ with the goal of making their 200 million-object photometric catalog available for data mining. Discoveries resulting from exploration of the IPHAS initiative included new samples of young stars [Vink et al., 2008], planetary nebulae [Viironen et al., 2009], and galactic supernova remnants [Sabin et al., 2013]. Virtual Observatory infrastructure was also utilized by Chilingarian et al. [2009] in a workflow to identify a sample of compact elliptical galaxies. Candidates were selected by leveraging a combination of resources including imaging data from the VizieR Catalogue Service² at the Centre de Données Astronomiques de Strasbourg, the NASA/IPAC (National Aeronautics and Space Administration/Infrared Processing and Analysis Center) Extragalactic Database (NED)³, the Hubble Legacy Archive⁴, and photometric and spectroscopic results from SDSS Data Release 7 [DR7; Abazajian and et al., 2009]. SDSS catalogues also played a role in projects such as the identification of dwarf novae candidates [Wils et al., 2010], found by cross-matching DR7 with an astrometric catalogue from the Galaxy Evolution Explorer (GALEX) space mission [Martin et al., 2005]. The need for practical data mining infrastructure led to the development of tools such as DAMEWARE [Data Mining & Exploration Web Application and REsource; Brescia et al., 2014, 2016]. See also Ivezić et al. [2014] and the AstroML Python module⁵ for a selection of data mining implementations, with an emphasise on large-scale observational surveys.

1.2 The emergence of machine learning and artificial intelligence in astronomy

The value of automated data mining as an approach to knowledge discovery in astronomy has been firmly established across a broad range of sub-disciplines of astronomical interest. Within many fields of astronomy, though, the discussion of data mining is evolving rapidly to focus almost exclusively on *machine learning* (ML; automated processes that learn by example in order to classify, predict, discover or generate

¹<http://www.astrogrid.org>

²<http://vizier.u-strasbg.fr>

³<http://nedwww.ipac.caltech.edu>

⁴<http://hla.stsci.edu>

⁵<https://www.astroml.org>

new data) and, to a lesser extent, *artificial intelligence* (AI; methods by which a computer makes decisions or discoveries that would usually require human intelligence).

The developing use of ML and AI in astronomy has mirrored the broader use in computer science and the scientific community. Traditional statistical techniques found application first. Principal component analysis (PCA) was used, for instance: in the 1980s for morphological classification of spiral galaxies [Whitmore, 1984]; in the 1990s for quasar detection [Francis et al., 1992] and stellar spectral classification [Singh et al., 1998]; and in the 2000s for galaxy classification [Conselice, 2006] and quasar detection in the Sloan Digital Sky Survey [Yip et al., 2004]. PCA is now a standard technique, which continues to be used in hundreds of astronomy projects and papers per year.

By the early 1990s, astronomers begin to take advantage of more complex methods requiring labelled training sets. In the 1990s, decision Trees (DTs) began to be employed for tasks such as star-galaxy separation [Weir et al., 1995] and galaxy morphology classification [Kriessler et al., 1998, Owens et al., 1996]. By the 2000s, use of the technique proliferated and random forests (RFs) begin to dominate, with a key application being photometric redshift estimation [Carrasco Kind and Brunner, 2013]. Boosted decision tree techniques, such as AdaBoost, appeared in more recent years and continue to be used, including for assignment of photometric redshifts [Hoyle et al., 2015a] and for star-galaxy separation [Sevilla-Noarbe and Etayo-Sotos, 2015].

Support Vector Machines (SVMs) also found application in the 2000s and beyond, for instance in the detection of red variable stars [Woźniak et al., 2004], determination of photometric redshifts [Wadadekar, 2004], prediction of solar flares [Qahwaji and Colak, 2007], star-galaxy separation [Fadely et al., 2012], and noise analysis in gravitational wave detection [Biswas et al., 2013].

One of the dominant machine learning techniques, the artificial neural networks (ANN), appeared in the field at the end of the 1980s [Angel et al., 1990, Rosenthal, 1988] and by the 1990s was applied across a broad range of problems in astronomy. Early applications included star-galaxy separation [Odewahn et al., 1992], galaxy morphology classification [Lahav et al., 1996, Storrie-Lombardi et al., 1992], and object detection in the staple astronomical software Source Extractor [SExtractor; Bertin and Arnouts, 1996]. By the 2000s, ANNs were playing a key role in photometric redshift estimation [Collister and Lahav, 2004, Firth et al., 2003, Vanzella et al., 2004], galaxy classification [Ball et al., 2004] and the detection of gamma ray bursts [GRBs; Ball et al., 2004]. Paving the way for the “Deep Learning” era, the use of ANNs in astronomy has accelerated over the last decade, for instance in the analysis of asteroid composition [de León et al., 2010], pulsar detection [Eatough et al., 2010], and finding gravitationally lensed quasars [Agnello et al., 2015].

Two strongly linked occurrences have had a significant impact on the growth of adoption of ML and AI in astronomy. First was the appearance of graphics processing units (GPUs) as affordable, massively parallel computational accelerators, with applicability to a wide range of computationally-demanding problems [see, for example, Barsdell et al. [2010], Fluke et al. [2011] for adoption strategies in astronomy]. Secondly was the emergence of deep neural networks and convolutional neural networks. These approaches – extensions to the “vanilla” ANN – benefit from GPU acceleration to perform computationally arduous calculations in parallel in a reasonable time and at relatively low cost. For data-rich fields, such as astronomy, the predictive performance of these deep learning networks improves as more data is provided for training and tuning.

In the field of computer vision – the computational analysis of image data – the use of deep neural networks also accelerated after the spectacular demonstration by Krizhevsky et al. [2012] of the power of convolution neural networks applied to classifying images of millions of everyday objects. Astronomers were quick to take advantage of this revolution, with Dieleman et al. [2015] and Huertas-Company et al. [2015] achieving human-level performance on galaxy morphology classification and Hoyle [2016] demonstrating the possibility of estimating photometric redshifts directly from images using convolutional neural networks.

Recent applications in astronomy utilizing ML and AI include: the discovery of extrasolar planets [Pearson et al., 2018, Shallue and Vanderburg, 2018] and gravitationally-lensed systems [Jacobs et al., 2017, Lanusse et al., 2018, Pourrahmani et al., 2018]; discovery and classification of transient objects [Connor and van Leeuwen, 2018, Farah et al., 2018, Mahabal et al., 2019]; forecasting solar activity [Florios et al.,

2018, Inceoglu et al., 2018, Nishizuka et al., 2017]; assignment of photometric redshifts within large-scale galaxy surveys [Bilicki et al., 2018, Ruiz et al., 2018, Speagle and Eisenstein, 2017]; and the classification of gravitational wave signals and instrumental noise [George and Huerta, 2018b,a, Powell et al., 2017].

1.3 Scope and structure

This advanced review surveys progress in the wide-scale adoption of machine learning and artificial intelligence within astronomy, as evidenced by a collection of recently published works. Techniques are not explained and referenced in detail, except with respect to their particular adoption in astronomy. Most advances in astronomical and astrophysical knowledge have relied on a relatively small number of general methods (see Section 2).

The primary avenue for identifying relevant literature was NASA’s Astrophysics Data System⁶ [Chyla et al., 2015, Kurtz et al., 2000] – a knowledge discovery tool without equal. To gather an extensive and representative current sample of publications, abstracts of published, peer reviewed journal articles were probed for key words such as “machine learning”, “artificial intelligence”, “neural networks”, and “data mining”. Bayesian methods are intentionally omitted, as these align more naturally with traditional statistical methods [Heck et al., 1985]. Such an approach does miss important research results, so no claim is made as to the exhaustiveness or completeness of the review. Progress in the adoption of ML and AI in astronomy is occurring rapidly. However, through the qualitative examination of ~ 200 refereed publications from 2017 to February 2019, using an approach sharing elements with Grounded Theory,⁷ a broad collection of astronomy applications has been assessed such that common themes have emerged regarding the reach and maturity of ML and AI in astronomy.

Indicative examples are drawn from the recent published literature to highlight how ML and AI techniques are used across seven categories of activity (Section 2) and three phases of maturity (Section 3). It is important to remember that a successful use of a machine learning algorithm is more likely to be reported than one where a method failed to work. Counter examples or cautionary tales [e.g. Connor and van Leeuwen, 2018], where an algorithm may not have performed as well as hoped, are rare.

2 Machine learning and artificial intelligence in astronomy

Data-driven scientific discovery occurs through a combination of statistical methods, machine learning and artificial intelligence techniques, and the use of database systems.⁸ Scientific discovery requires techniques for identifying patterns within datasets (the original scope of *data mining*) as part of a multi-stage process for selecting, cleaning, processing, and transforming raw data into useful knowledge (i.e., *knowledge discovery in databases*, as described in Fayyad et al. [1996]).

As highlighted in Section 1, global astronomy data collections are approaching the Exabyte scale. A key motivation for many applications of ML and AI to astronomical data is the need to prepare for the data streams expected from near-term observatories and space missions. The Large Synoptic Survey Telescope [Ivezić and et al., 2019, LSST Science Collaboration and et al., 2009]; the Euclid satellite [Laureijs and et al., 2011, Amendola et al., 2013]; MeerKAT [Booth et al., 2009]; the Australian Square Kilometre Array Pathfinder [Johnston et al., 2007, 2008]; and the Square Kilometre Array [Dewdney et al., 2009], among others, will all generate datasets on scales (volumes and velocities) that vastly exceed the

⁶<http://ui.adsabs.harvard.edu>

⁷Grounded Theory is a qualitative analysis strategy, used, for example, in education research and social sciences, where a sequence of reader-assigned codes is applied to allow identification and tracking of themes within a sample of relevant literature. Codes do not have to be selected in advance, but are defined dynamically in multiple iterations through the literature.

⁸As proposed by Ball and Brunner [2010], a sufficiently flexible definition for a database in astronomy is “*any machine-readable astronomical data*”.

discovery capabilities of humans. In the interim, the Sloan Digital Sky Survey [SDSS; York and et al., 2000, Stoughton and et al., 2002, Abazajian and et al., 2009],

the Panoramaic Survey Telescope and Rapid Response System [Pan-STARRS; Kaiser, 2004], the Catalina Real-Time Transient Survey [CRTS; Drake et al., 2009, Mahabal et al., 2011] and the Zwicky Transient Facility [ZTF; Bellm et al., 2019], the Kilo Degree Survey [KiDS; de Jong et al., 2013] and the Fornax Deep Survey [Iodice et al., 2016], both using the the VLT Survey Telescope⁹, LOFAR [van Haarlem and et al., 2013], the Solar Dynamic Observatory [SDO; Lemen et al., 2012, Pesnell et al., 2012], the Kepler Planet-Detection Mission [Borucki and et al., 2010], and the GAIA space mission [Gaia Collaboration and et al., 2016b,a, 2018], are generating data with which ML and AI has enabled classification, regression, forecasting, and discovery, leading to new knowledge and new insights.

2.1 The nature of the data

The applicability and efficacy of any ML or AI technique depends on the nature of the data. Brunner et al. [2002] classified astronomical data into five domains, extended slightly here to allow a clearer connection to the specific applications of ML and AI within astronomy. In this section, references are given to recent works that apply ML and AI to each of the data types, rather than to the originator(s) of the data type.

Images are pictures of astronomical objects (usually as a pixel grid of numerical intensity values), such that the appearance informs a classification [e.g., Aniyan and Thorat, 2017, Xin et al., 2017, Domínguez Sánchez et al., 2018, Kuminski and Shamir, 2018, Ma et al., 2019] or provides insight about physical processes that are occurring [Müller et al., 2018]. For visible/infrared (IR) observations (i.e. optical astronomy), light passes through, and is focused by, a telescope’s optical system to be captured on a charge-coupled device (CCD). Various filters are used to select only specific regions of the visible/IR spectrum. For radio observations, it is common to refer to the frequency bandwidth over which flux is recorded from a particular location in the sky, with most radio images created from interferometers using the technique of aperture synthesis.

Spectroscopy refers to wavelength-dependent numerical intensity measurements over a finite range of wavelengths (or frequencies), but often with very high resolution. Spectroscopy provides information on the atomic and molecular composition, from which other physical properties (temperature, density, metallicity, etc.) can be inferred [e.g., Li et al., 2018, Márquez-Neila et al., 2018, Miettinen, 2018, Ucci et al., 2018].

A special case at the intersection of imaging and spectroscopy is the spectral cube [e.g., Araya et al., 2018, Bron et al., 2018]. This is a volumetric dataset comprising a sequence of images, each captured over a very narrow wavelength or frequency range. When looking for structures within a spectral cube, it is treated as an image. When extracting a spectrum at a fixed spatial location, it is treated as a spectroscopic data product. Spectroscopic data cubes, with their high dimensionality, may prove a challenge for established machine learning methods to handle. Convolutional neural networks have proved to work robustly on astronomical image data in several photometric bands, so there is no theoretical obstacle to extending this to thousands of spectroscopic frequencies. However, the practical challenges are yet to be fully explored.

Photometry is concerned with accurate measurements of the brightness (i.e. intensity, luminosity, flux) of an object recorded through a filter. It is a secondary, numerical data product derived from a calibrated image. Comparisons between photometric measurements through different filters are often used as an alternative to detailed spectroscopic observations, with specific application to determining the distance to a celestial source [e.g. Cavuoti et al. [2017a], Morrison et al. [2017], Beck et al. [2018], Bilicki et al. [2018]].

Images, spectroscopic, and photometric measurements can be made as a function of time. For optical astronomers, a *light curve* is time-based photometry, where the variation in intensity of a source over time helps with the identification and classification of a variety of variable star types [e.g. Cohen et al., 2017, Naul et al., 2018, Papageorgiou et al., 2018] or indicates the presence of otherwise unseen objects, such

⁹These last two survey projects provide data products for the SURvey Network for Deep Imaging Analysis & Learning (SUNDIAL) which is building inter-disciplinary teams of astronomers, computer scientists and industry partners. See <https://www.astro.rug.nl/sundial/>

as an extrasolar planet [e.g. Mislis et al., 2018, Pearson et al., 2018, Shallue and Vanderburg, 2018]. In this review, other time-based measurements in radio astronomy [e.g. pulsar and transient object searches - Connor and van Leeuwen [2018], Michilli et al. [2018], Pang et al. [2018], Tan et al. [2018], Farah et al. [2018]], and the emerging field of observational gravitational wave astronomy [Powell et al., 2017, Zevin et al., 2017, George and Huerta, 2018b,a], are categorized as *time series*.

The end product of many data gathering programs is a *catalogue*, which comprises one or more numerical or categorical data types. Some may be derived from the standard data gathering approaches introduced above, while others are calculated or otherwise derived – including through ML [e.g. Marchetti et al., 2017, Tachibana and Miller, 2018]. Within a catalogue, *astrometry* refers to the accurate measurement of the spatial locations of objects, on the celestial sphere or with respect to an alternative coordinate system [Castro-Ginard et al., 2018, Gao, 2018a,b]. Most objects are reported with at least one measurement of position. Some local objects, such as the Solar System’s planets and minor planets, and the growing number of stars within the reach of the GAIA space mission, move with respect to the coordinate system over time [Chen et al. [2018], Lin et al. [2018]]. A *morphological classification* places a particular type of object into an object-based category where a common physical process (or set of processes) is thought to drive the appearance of an object [e.g. Aniyan and Thorat, 2017, Domínguez Sánchez et al., 2018, Kuminski and Shamir, 2018, Ma et al., 2019].

Finally, although a single catch-all name does not do such a diverse field justice, *simulation* will be used to describe the data products from any numerical or computational method. For example, cosmological simulations [e.g. Agarwal et al., 2018, Hui et al., 2018, Lucie-Smith et al., 2018, Nadler et al., 2018, Rodríguez et al., 2018] follow the gravity-induced formation and growth of structures, requiring approximations to various physical mechanisms, a suitable choice of initial conditions, and a strategy for time-based evolution (down to some minimum level of accuracy).

2.2 From classification to insight

For the science of astronomy to progress through the use of ML and AI, it must be possible to demonstrate that the outcome is not merely an automated classification or a numerical prediction, but that astronomers are using the data mining phase to discover new objects or generate new insights into the underlying physical processes and relationships. In assessing the recently published literature, seven categories emerge pertaining to how ML and AI are used in astronomy (cf. Fayyad et al. [1996] and Zhang and Zhao [2015], who identify similar categories). The last two categories (discovery and insight) are those where a higher order scientific outcome arises.

1. **Classification:** Categories or labels are applied to objects or features. Based on a training set (labelled or unlabelled), the machine learning algorithm learns the characteristics that relate an instance to a category. When applied to a new instance, the algorithm assigns the most likely category label.
2. **Regression:** Assignment of a numerical value (or values) based on the characteristics that are learnt or otherwise predicted by the machine learning algorithm. As with classification, a training set may be used or the characteristics may be inferred from the dataset.
3. **Clustering:** These algorithms determine whether an object or a feature is part of (i.e. a member of) something. This might be a physical structure or association – as in the more familiar usage of the term in astronomy as applied to open, globular, or galactic clusters – or a region within an N -dimensional parameter space.
4. **Forecasting:** The purpose of the machine learning algorithm is to learn from previous events, and predict or forecast that a similar event is going to occur. There is an implicit time-dependence to the prediction.

5. **Generation and Reconstruction:** Missing information is created, expected to be consistent with the underlying truth. The cause of the missing information might be due to the presence of noise, processing artefacts, or additional astronomical phenomena, all of which conspire to obscure the required signal.
6. **Discovery:** New celestial objects, features or relationships are identified as a consequence of the application of a ML or AI method.
7. **Insight:** Moving beyond the discovery of celestial objects, new scientific knowledge is demonstrated as a consequence of applying machine learning or AI. This includes cases where insight is gained into the suitability of applying machine learning, choice of data set, hyperparameters, and comparisons with human-based classification.

Classification, regression and clustering processes are often presented as a comparison with a similar human-centred approach – but with a need to “scale-up” in terms of either the size of the dataset to be explored or “speed-up” the time taken to achieve the task. Classification and regression outcomes can either be the end-point of an investigation, or the input to a forecasting, generation, discovery or insight process. In Section 3, these categories will be used to make an assessment of the maturity of adoption of ML and AI in various sub-fields of astronomy.

A subset of discovery is the field of anomaly or outlier detection. Many of the most exciting discoveries to come are likely to lie among the “unknown unknowns” in new areas of parameter space captured within the Petascale and Exascale datasets of the future. New methods are being developed to find anomalous objects in astronomical datasets, such as the work by Baron and Poznanski [2016] using an unsupervised Random Forest to find outliers amongst SDSS galaxies. Promising avenues involve a combination of unsupervised learning methods, such as isolation forests [Liu et al., 2008]; dimensionality reduction, such as PCA, t-Distributed Stochastic Neighbor Embedding [t-SNE; van der Maaten and Hinton, 2008], e.g. Reis et al. [2018] and Nakoneczny et al. [2019], self-organizing maps [SOMs; Kohonen, 1990], e.g. Carrasco Kind and Brunner [2014] and Armstrong et al. [2016], or the latent space of a variational encoder [e.g. Yang and Li, 2015, Ma et al., 2019]. Novel visualization techniques are also contributing. For instance, Masters et al. [2015], using self-organizing maps to visualize the distribution of galaxies in photometric color space, were able to identify regions that were under-sampled spectroscopically and develop an optimal strategy for the Euclid mission’s photometric redshift calibration efforts.

Based on a qualitative examination of a sample of ~ 200 refereed publications from 2017 to February 2019, Table 1 summarises the mapping between the most common astronomical data types with the seven categories of ML/AI methods. Classification and regression algorithms are being applied to all of the data types. Although clustering activities span a number of data types, overall they were not common outside of studies of stellar clusters [Castro-Ginard et al., 2018, Gao, 2018a,b] or segmentation processes [Bron et al., 2018, Yang et al., 2018].

Forecasting outcomes were mostly confined to images [e.g. Mukkavilli et al. [2018] with Mars images; Liu et al. [2017], Nishizuka et al. [2017], Inceoglu et al. [2018], and Florios et al. [2018] using Solar magnetograms], photometric measurements [French and Zabludoff [2018] predicted likely tidal disruption events in post-starburst galaxies using a random forest algorithm] and catalogue data [forecasts of coronal mass ejections from the Sun based on ~ 180 similar events using a support vector machine (SVM) Liu et al. [2018]]. Generation methods, in particular generative adversarial networks, have been used with images from observations [Vavilova et al., 2018], and to simplify or remove the need for expensive numerical simulation [e.g. Diakogiannis et al., 2019, Rodríguez et al., 2018, Fussell and Moews, 2019].

With regards to the role of ML and AI in advancing knowledge in astronomy, there was clear evidence from the sample of recent publications that discovery tasks are being performed with all of the data types: images [Hartley et al., 2017, Ciuca and Hernández, 2017, Gomez Gonzalez et al., 2018, Pourrahmani et al., 2018, Lanusse et al., 2018, Wan et al., 2018, Morello et al., 2018, Jacobs et al., 2017]; spectroscopy [Bu et al., 2017, Li et al., 2018]; photometry [Timlin et al., 2018, Vida and Roettenbacher, 2018, Ostrovski et al.,

Table 1: From a qualitative examination of a sample of ~ 200 refereed publications from 2017 to February 2019, a mapping emerges between the nature of astronomical data and the way that machine learning and artificial intelligence is actively been pursued. The table presents a qualitative summary of the categories of ML/AI algorithms and the most common types of astronomical data in the sample of publications.

Nature/Type	Classification	Regression	Clustering	Forecasting	Generation	Discovery	Insight
Image	•	•	•	•	•	•	•
Spectroscopy	•	•	•			•	•
Photometry	•	•	•	•		•	•
Light curve	•	•				•	•
Time Series	•	•	•			•	•
Catalogue	•	•	•	•		•	•
Simulation	•	•			•	•	

2017]; light-curves [Armstrong et al., 2018, Peña et al., 2018, Cohen et al., 2017, Hedges et al., 2018, Heinze et al., 2018, van Roestel et al., 2018, Giles and Walkowicz, 2019]; time-series [Michilli et al., 2018, Tan et al., 2018, Pang et al., 2018, Connor and van Leeuwen, 2018, Farah et al., 2018, Morello et al., 2019]; catalogues [Nguyen et al., 2018, Yan et al., 2018, Lin et al., 2018, Marchetti et al., 2017]; and simulation: [Xu and Offner, 2017, Nadler et al., 2018, Kuntzer and Courbin, 2017].

For most data types, outcomes have progressed to the insight phase, for example: enhanced understanding of human biases in classification of Galaxy Zoo project images [Peng et al., 2018, Cabrera-Vives et al., 2018]; determination of the evolution of the effective radius and stellar mass of Kilo Degree Survey [KiDS; de Jong et al., 2013] galaxies based on photometric redshifts derived from ML [Roy et al., 2018]; and new relationships between physical and environmental properties of galaxies by applying an SVM to the results of a cosmological simulation [Hui et al., 2018].

2.3 Techniques

Machine learning algorithms are usually classified as being either supervised or unsupervised. Supervised methods rely on a pre-labeled dataset, which is used to help train and tune the algorithm. This learning allows for new instances to be assigned a label (classification) or numerical value (regression). Unsupervised methods allow the data to speak for itself, but do not necessarily make use of any existing knowledge. Although there is no shortage of data in astronomy, there is often a paucity of relevant pre-labeled data. For example, when the discovery of rare events is the target of an observational program, it is very difficult to train a network on sufficient examples and counter-examples. Moreover, astronomical discovery does rely on serendipity – anomalous cases that are potentially unlike anything that has previously been examined, and hence no exemplars exist [Norris, 2017].

The lion’s share of machine learning in astronomy is performed with five classes of algorithms: artificial neural networks; convolutional neural networks; decision trees; random forests; and support vector machines. These are primarily used as supervised learning algorithms. Since the Ball and Brunner [2010] review, convolutional neural networks are the only new method amongst these five to emerge and reach wide-spread usage in astronomy.

Artificial neural networks [ANNs; Rosenblatt, 1957, Fukushima, 1980] are the key technique behind the recent AI boom, but date back to the 1950s. They are designed by analogy to a biological neuron, with signals from multiple inputs weighted and added together; a biological neuron sends an electrical

signal if these weighted input signals cross a certain activation threshold. In the case of an artificial neuron (“perceptron”), the inputs and trainable weights are vectors of real valued numbers, and the output is a scalar value. A single artificial neuron can be employed as a classifier, however they are typically combined in ANNs where the outputs of an array (layer) of neurons form the inputs to a subsequent layer. At the output layer, the values of one or more neurons are interpreted according to the problem domain. Optimization of the weights using a labelled training set follows the gradient descent paradigm with the backpropagation algorithm [LeCun et al., 1989].

Support vector machines [SVMs; Cortes and Vapnik, 1995], similarly to ANNs, learn non-linear decision boundaries in spaces of arbitrary dimension, finding a hyperplane in a space of arbitrary dimensionality that distinctly separates the supplied data. The algorithm works by finding the hyperplane with the maximum separation between extreme examples (support vectors). A linear SVM (LSVM) finds the optimal hyperplane for the supplied input features, but using the so-called ‘kernel trick’ a SVM projects the data into higher dimensions where the data is linearly separable.

Decision trees [DTs; Quinlan, 1986] perform classification by recursive binary splitting of the data, learning through the training process a series of decisions based on features of the input data. The root node (the entire data space) is repeatedly split into two child nodes based on the most discriminative feature of the data, until at the leaf node a category is determined. **Random forests** [RFs; Ho, T.K., 1995, Breiman, 2001] are an extension of decision trees, improving accuracy by constructing an ensemble of decision trees, trained on subsets of the training data [bagging; Breiman, 1996] and/or feature set (feature randomness), and using the median or mode of the ensemble as the final output value. **AdaBoost** [Freund and Schapire, 1995] is another ensemble method that weights the contribution of each decision tree based on misclassifications; similarly, **gradient boosting** [Friedman, 2001] uses decision trees trained in sequence on the residual errors of other DTs.

A convolutional neural network (CNN) is an extension of the simple ANN but with many hidden layers (i.e. a deep neural network). CNNs are characterized by their use of convolutional layers, which are sensitive to specific features – usually within images – that may have undergone transformations through translation, rotation, or scaling. Working in conjunction with pooling layers, which reduce the spatial size of image features within the network, the final stage of a CNN is often a fully-connected ANN to generate a classification or numerical prediction.

CNNs have now been used in astronomy for a variety of image-based classification, regression and discovery activities. They appear in the literature as: *binary classifiers* [Gieseke et al. [2017], Jacobs et al. [2017] and Shallue and Vanderburg [2018]], where the training sets comprise two distinct categories representing “present” and “not present” examples; *morphological classifiers* [Aniyan and Thorat [2017], Domínguez Sánchez et al. [2018], González and Guzmán [2018], Huertas-Company et al. [2018] and Ma et al. [2019]], where there are multiple categories that have been determined previously, usually by human inspection, but also using other machine learning approaches [Kim and Brunner, 2017]; and for *detection*, with applications to discovery of exoplanet candidates in light-curves [Pearson et al., 2018], or real-time discovery of transient objects [Connor and van Leeuwen, 2018] and gravitational wave events [George and Huerta, 2018b,a].

Outside of this core group, are several unsupervised methods: k -nearest neighbours (k -NN), k -means clustering, and the DBSCAN method. The latter has been used as a discovery tool to improve efficiency at detecting exoplanet transits from light curves, based on the recovery of artificial transit signatures [Mislis et al., 2018], and in partnership with an ANN to identify open clusters [Castro-Ginard et al., 2018] in the GAIA DR2 [Gaia Collaboration and et al., 2018] – an example of a clustering process.

Other techniques that have been investigated, often in conjunction with one or more the above methods, include: AdaBoost [Xin et al., 2017, Bethapudi and Desai, 2018]; genetic algorithms [Sarro et al., 2018]; self-organizing maps [Armstrong et al., 2017, Süveges et al., 2017, Armstrong et al., 2018]; recurrent neural networks [Naul et al., 2018]; auto-encoders [Vincent et al., 2008, Sedaghat and Mahabal, 2018]; and transfer learning [Benavente et al., 2017]. Falling within the generation and reconstruction category (Section 2.2), generative adversarial networks (GANs) are likely to be the next most significant machine

Table 2: From a qualitative examination of a sample of ~ 200 refereed publications from 2017 to February 2019, a mapping emerges between the nature of astronomical data and the types of machine learning and artificial intelligence algorithms that are being applied. The table presents a summary of the types of astronomical data and the algorithms that appeared most regularly. The purpose of the table is to provide a convenient starting point for selecting an algorithm that has been used successfully for each data type.

Data/Method	ANN	CNN	GAN	SVM	DT	RF	DBSCAN	k-NN	k-M
Image	•	•	•	•	•	•		•	
Spectroscopy	•	•		•		•			•
Photometry	•				•	•	•		•
Light curve		•				•			
Time Series	•	•			•	•	•		
Catalogue	•			•	•	•	•	•	
Simulation	•	•	•	•		•			

ANN = Artificial Neural Network; CNN = Convolutional Neural Network; GAN = Generative Adversarial Network; SVM = Support Vector Machine; DT = Decision Tree; RF = Random Forest; DBSCAN = Density-based spatial clustering of applications with noise; k-NN = k-Nearest Neighbours; k-M = k-means clustering

learning approach for astronomy. Early applications of GANs include generating dark matter structures in cosmological simulations [Rodríguez et al., 2018, Diakogiannis et al., 2019], the creation of realistic images of galaxies as an input to weak gravitational lensing analysis [Fussell and Moews, 2019], and deblending overlaps between foreground and background galaxies in highly-crowded images [Reiman and Göhre, 2019].

Machine learning has also been used to identify the astrophysical features most significant for classification. For example, in the area of photometric redshift estimation, Polsterer et al. [2014] used GPUs to conduct an exhaustive feature search of over 341,000 feature combinations to identify the four most significant ones; and Hoyle et al. [2015b], who used Random Forests with AdaBoost to select the top photometric features to increase the performance of ANN-based redshift estimators. Frontera-Pons et al. [2017] used denoising autoencoders for unsupervised feature learning from galaxy spectral energy distributions (SEDs).

Table 2 summarises the relationships found between the main types of astronomical data and specific techniques in the sample of recently-published papers. It is expected that specific techniques have been applied to other data types, and it is important to remember that some fields will have trialled particular methods and moved on as new alternatives appear. The Table’s purpose is to emphasise areas of current activity and interest only, and thus provides a starting point for astronomers wishing to adopt a ML/AI approach by matching the data types to the methods.

ANNs [Ciuca and Hernández, 2017, Marchetti et al., 2017, Bethapudi and Desai, 2018, Bilicki et al., 2018, Fujimoto et al., 2018, Ho, 2019], random forest methods [Schindler et al., 2017, Goulding et al., 2018, Hedges et al., 2018, Reis et al., 2018, Pang et al., 2018, Tachibana and Miller, 2018, Nadler et al., 2018], and SVM algorithms [Hartley et al., 2017, Hui et al., 2018, Kong et al., 2018, Yan et al., 2018, Zhang et al., 2018] have been used extensively across most data types. CNNs are more suitable for image-style data (see above), although they have been used successfully with one-dimensional light curves [Shallue and Vanderburg, 2018, identification and ranking of transiting exoplanet candidates in Kepler light curves, including the discovery of two new exoplanets] and time series [George and Huerta, 2018b,a, identification of gravitational wave signatures within noisy time series data – a solution that scales better than template matching as the number of templates grows]. DBSCAN [Castro-Ginard et al., 2018] and k-NN [Smirnov and Markov, 2017] have been used to find structures in multi-dimensional catalogues. Given the prevalence of imaging data in astronomy, it is not surprising that images are being analyzed with the largest range of ML/AI methods.

There is still plenty of scope for studies that perform structured comparisons between multiple methods. This can occur more easily when reference datasets are made accessible to the community. For

example, the availability of the PHoto-z Accuracy Testing datasets [Hildebrandt et al., 2010] allowed Cavuoti et al. [2012] and Brescia et al. [2013] to establish the efficacy of a multi-layer perceptron (i.e. neural network) method coupled with the Quasi Newton Algorithm (MLPQNA) at assigning photometric redshifts for galaxies and quasars respectively. Training on the PHAT-1 spectroscopic sample, MLPQNA out-performed alternative statistical and neural network-based methods [e.g. ANNz; Collister and Lahav [2004]] with regards to bias¹⁰ for all objects, bright objects, and distant vs near objects in the PHAT-1 sample [Cavuoti et al., 2012]. Studies into the accuracy and validity of ML-based photometric redshifts continues today – see, for example, Almosallam et al. [2016], Cavuoti et al. [2017a] and Amaro et al. [2019].

Looking more broadly, these systematic comparisons tend to be occurring more often in solar astronomy than in other disciplines [e.g. Nishizuka et al. [2017], Florios et al. [2018] and Inceoglu et al. [2018]], although see Ksoll et al. [2018], Pashchenko et al. [2018], and Zhang et al. [2018] for examples pertaining to stellar and variable star classifications. While certain disciplines have adopted specific methods, experimentation with emerging techniques is on-going [e.g. probabilistic random forests and transfer learning Reis et al. [2019]].

3 Assessing the maturity of adoption

The seven categories introduced in Section 2.2 allow an assessment of the maturity of the use of machine learning and artificial intelligence within a sub-field of astronomy, as they represent a loose hierarchy of sophistication. The common starting point is to apply a machine learning technique to perform a classification, regression or clustering task. Once established as being comparable to, or exceeding, a more traditional approach, machine learning can be used to forecast likely future outcomes [e.g. solar flares [Nishizuka et al., 2017, Florios et al., 2018] or coronal mass ejections from the Sun [Inceoglu et al., 2018]] or make new discoveries [e.g. classification schemes for stellar types permitting the identification of new candidates of rare objects as in Bu et al. [2017], van Roestel et al. [2018], and Zhang et al. [2018]]. The most mature disciplines move beyond classification and discovery as ends in their own to that of gaining insight – where new physical knowledge is identified, often for the first time, because a machine learning approach was used.

The hierarchy of categories is used to assess the maturity of ML and AI within a set of sub-fields of astronomy as one of *emerging*, *progressing*, or *established*. In all cases, the reader should refer to the highlighted works in order to understand the scientific background, historical context for the establishment of a particular method, and the technical details of the data mining, machine learning or artificial intelligence approach that was applied.

3.1 Emerging

The emerging stage is applied to sub-fields of astronomy and astrophysics that are starting to investigate the use of ML and AI, often by tackling the “low-hanging fruit”. This includes a problem that requires a classification or regression approach, or through a comparison between machine learning and an alternative, established method. While some of the emerging disciplines show evidence of reaching the discovery and insight phases, the approaches are not as firmly established, or the size of the community is small. Emerging fields include:

- *Planetary studies.* ML-based identification and classification of clouds, dust storms and surface features on Mars [Gichu and Ogohara, 2019], with potential to forecast future dust storms [Mukkavilli et al., 2018], and the discovery of previously unknown impact craters [Xin et al., 2017] using the AdaBoost algorithm.

¹⁰the mean of $\Delta z \equiv (z_{\text{spec}} - z_{\text{phot}}) / (1 + z_{\text{spec}})$, where z_{spec} and z_{phot} are the known spectroscopic and predicted photometric redshifts respectively.

- *Non-stellar components of the Milky Way.* The primary components of the Milky Way are stars (see below), dust and gas, which can be concentrated in atomic and molecular clouds or more diffusely in the interstellar medium. Ucci et al. [2018] developed the **GAME** (GALaxy Machine learning for Emission lines) code to study physical properties of the interstellar medium. Segmentation and clustering algorithms are used to identify individual components of clouds of atomic and molecular gas within the Milky Way. Bron et al. [2018] used the Meanshift clustering algorithm to identify regions within molecular clouds that based on physical/chemical properties, instead of seeking purely spatial connections; Dénes et al. [2018] used ML to determine individual Gaussian components of the Riegel-Crutcher cloud, based on 21-cm neutral hydrogen (HI) observations of extra-galactic continuum sources behind the cloud complex; and SVM was used by Yan et al. [2018] to classify, and hence select, HII regions (gas clouds comprised mostly of singly-ionized hydrogen) from the Infrared Astronomical Satellite (IRAS) Point Source Catalogue. Time-consuming human classifications of Milky Way “bubbles”, caused by stellar feedback within molecular clouds, was enhanced through ML [Xu and Offner, 2017]. A random forest method was used by Chen et al. [2019] to determine the amount of dust reddening of more than 56 million stars, with application to future GAIA datasets. Using a GAN, Vavilova et al. [2018] demonstrated how to generate missing parts of the cosmological large-scale structure that are obscured by the Milky Way’s zone of avoidance.
- *Stellar clusters.* Application of the density-based DBSCAN algorithm to the Gaia Data Release 2 (DR2) has lead to the discovery of new open clusters [Castro-Ginard et al., 2018], with an ANN used to separate real clusters from spatial over-densities of stars. Multi-dimensional clustering processes have been used to determine the components of several open clusters, including M67 Gao [2018a] and NGC188 [Gao, 2018b].
- *Instrumentation.* Li and Yang [2018] presented an AI solution for identifying faults for a telescope drive system. Through an automated expert system, a series of self-healing decisions are made until an appropriate solution is found. The knowledge base is updated in real-time with human input for faults that have not previously been diagnosed or corrected.

Other emerging disciplines include: information retrieval systems, matching queries regarding specific instruments [Mukund et al., 2018]; identification of cosmic strings in all-sky maps [e.g. Ciuca and Hernández [2017] using a neural network; Vafaei Sadr et al. [2018]]; and the detection and classification of cosmic ray events [Krause et al., 2017, Zhao et al., 2018].

3.2 Progressing

Characteristics of disciplines identified as progressing in their use of ML and AI include a broader variety of techniques being applied, or a particular technique is used multiple times, or there is an immediate move to the forecasting, discovery or insight phases. Sub-fields at this stage of maturity include:

- *Solar System objects.* Due to their relative proximity to the Earth, the motion of Solar System objects is key to their discovery. ML and AI have enabled removal, or reduction, of false detections from the moving object detection pipeline in the Subaru/Hyper-Suprime-Cam Strategic Survey Program [Lin et al., 2018], with applications to Trans-Neptunian Objects [Chen et al., 2018]; and detection and classification of asteroids [Erasmus et al., 2017, 2018, Smirnov and Markov, 2017]. Duev et al. [2019] trained a CNN to discover fast-moving candidates from ZTF observations in order to more reliably identify potentially hazardous near-Earth objects.
- *Active galactic nuclei and quasars.* A common theme in this field is the need for classification and detection methods, including assigning morphological types to radio-detected active galactic nuclei with a CNN [Ma et al., 2019], identifying blazar candidates in the Fermi-LAT (3LAC) Clean Sample [Kang et al., 2019], detecting rare high-redshift, extremely luminous quasars [Schindler et al., 2017],

and discriminating populations of broad absorption line quasars (BALQs) from non-BALQs in SDSS data releases [Yong et al., 2018].

- *Cosmological simulations.* ML is providing new methods for examining the outputs of cosmological simulations, leading to new insights about the connections between physical properties of galaxies, dark matter halos and the cosmic environment. Examples include the use of an ANN to aid in determining the total mass of the Milky Way and the Andromeda Galaxy from the Small MultiDark simulation [McLeod et al., 2017], and both classification of sub-halos [Nadler et al., 2018] and assignment of galaxies to halos [Agarwal et al., 2018] in dark matter-only simulations.

3.3 Established

In the established sub-fields, the use of ML and AI have become essential, a substantial body of literature exists, and the focus is mostly on forecasting, discoveries, or insight. Here, there is no longer a need to evaluate a suitability of machine learning – its usage has become ingrained. Established sub-fields include:

- *Solar astronomy.* Machine learning has been used for classification of solar flares [e.g., Liu [2017], Liu et al. [2017], and Benvenuto et al. [2018] via a hybrid method using both supervised and unsupervised methods]; clustering [e.g. Yang et al. [2018] presented the simulated annealing genetic (SAG) AI method to distinguish between the umbra, penumbra and solar photosphere through a segmentation approach]; and forecasting of coronal mass ejections with a SVM [Liu et al., 2018], and SVM and multilayer perceptrons [Inceoglu et al., 2018]. A number of systematic comparison studies have been conducted in order to assess which ML methods perform best at forecasting solar events. Nishizuka et al. [2017] compared three ML methods to forecast solar activity from time-based features using data from the SDO and the Geostationary Operational Environmental Satellite (GEOS) system. They determined that k-NN performed more effectively than SVM or extremely randomized trees. For a different data set from the SDO, Florios et al. [2018] found that random forests provided greater prediction accuracy.
- *Extra-solar planets.* The Kepler Mission, amongst other search programs, has been a rich source of light curve data. ML and AI have enhanced and improved candidate selection [Armstrong et al., 2018], and classification of light-curves, by removing false positives [Armstrong et al., 2017], raising detection efficiency [Mislis et al., 2018] and accuracy [Pearson et al., 2018], and identifying anomalies [Giles and Walkowicz, 2019]. ML techniques have allowed discovery of fainter candidates than were accessible with existing methods using neural network and random forest algorithms [Gomez Gonzalez et al., 2018] and a CNN [Shallue and Vanderburg, 2018]. Insight into candidate and confirmed extrasolar planets is also being achieved with ML and AI, such as through the determination of a “habitability score” for extra-solar planets [Saha et al., 2018] and improved model-fitting of atmospheric composition [Márquez-Neila et al., 2018].
- *Stars and stellar products.* Two key activities in stellar astronomy are spectral classification [e.g. Wang et al. [2017]; Garcia-Dias et al. [2018] with k-means clustering; Kong et al. [2018]; classification of young stellar objects with eight different methods by Miettinen [2018]] and photometric classification [e.g., Ksoll et al. [2018]; Zhang et al. [2018] with SVM, RF and Fast Boxes]. Many new examples of specific stellar classes have been discovered, such as Wolf-Rayet stars [Morello et al., 2018], blue horizontal branch stars [Wan et al., 2018], hot sub dwarf stars [Bu et al., 2017], and rare hypervelocity stars [Marchetti et al., 2017]. ML/AI have also led to the discovery of unresolved binary stars in simulated catalogues using RF and ANN algorithms [Kuntzer and Courbin, 2017], and new pulsars, and fewer false positives, from the LOFAR Tied-Array All-Sky Survey [Michilli et al., 2018, Tan et al., 2018]. Fujimoto et al. [2018] used an ANN to gain new insights into neutron star equation of state from numerical simulations.

- *Variable stars.* Time-based photometric observations of variable stars provide extensive data sets of many millions of individual objects for mining. There has been highly productive use of ML and AI for classification [e.g. Benavente et al. [2017], Naul et al. [2018], and Papageorgiou et al. [2018]] and discovery activities [e.g. comparative study with a variety of techniques, including SVM, k-NN, neural networks and random forests by Pashchenko et al. [2018]; discovery of new EL CVn-type binaries from the Palomar Transient Factory [van Roestel et al., 2018] supported with data from other large-scale surveys]. Valenzuela and Pichara [2018] used an unsupervised method to overcome the limitations of available training samples, particularly when approaching new survey data, by identifying similarities between light curves rather than relying on pre-classified examples.
- *Transient object detection.* While not yet the preferred method for all observational programs, the applicability of ML and AI has been firmly established – particularly for the real-time detection of transient objects, which allows new classes of celestial objects to be discovered. The Catalina Real-Time Transient Survey [Drake et al., 2009] was utilized as a large-scale test-bed for the use of ML to aid the optical detection and monitoring of variable and transient objects [e.g. Mahabal et al., 2009, Djorgovski, 2014, Djorgovski et al., 2016]. These early successes paved the way for the Zwicky Transient Factory [Bellm et al., 2019]. The multi-filter optical survey with the ZTF was designed to provide a rich exploration of the transient sky, generating hundreds of thousands of real-time candidate alerts for each night of operation. Machine learning is fundamental to accelerating and enabling the data analysis and candidate identification (and rejection) workflows of the ZTF. ML is being used to perform time critical tasks such as morphological star/galaxy classification [Tachibana and Miller, 2018], binary real/bogus classification of candidates, and asteroid detection [Mahabal et al., 2019], and can play a role in the brokering of alerts with application to the LSST Alert Stream [Narayan et al., 2018]. In a radio-based transient object project, Farah et al. [2018] used a random forest as part of the UTMOST real-time detection pipeline, leading to the discovery of Fast Radio Burst FRB170827, however, Connor and van Leeuwen [2018] determined that CNNs were sub-optimal for some radio transient tasks, such as reducing the need for GPU-accelerated, brute-force dedispersion of time series signals.
- *Galaxies.* One of the major areas of ML application has been in the classification of galaxies from optical and radio imaging surveys. Recent examples include: neural network-based Faranoff-Riley classifications of radio galaxies [Aniyani and Thorat, 2017]; automated morphological annotation and assignment [Beck et al., 2018, Domínguez Sánchez et al., 2018, Kuminski and Shamir, 2018] and labelling [Hocking et al., 2018] of galaxy images, including detections from radio surveys [Lukic et al., 2018]. The reference point for many of the automated classifiers is the work done by human volunteers for projects like Galaxy Zoo [Lintott et al., 2008]. Cabrera-Vives et al. [2018] uncovered human biases that existed in morphological classification, which could be reduced through supervised ML. Other applications included predicting the HI content of galaxies based on optical observations [Rafieeantsoa et al., 2018], determining physical properties of galaxies from their emission-line spectra [Ucci et al., 2017], point source detection from radio interferometry surveys [Vafaei Sadr et al., 2019], and cross-identification of sources from the Radio Galaxy Zoo [Alger et al., 2018]. Training a CNN on mock images of rare “blue nugget” galaxies from cosmological simulations, such objects were successfully found in an observational sample from the CANDELS survey [Huertas-Company et al., 2018].
- *Distance measures.* Estimates of the distances to galaxies, quasars and other remote celestial objects has benefited greatly from the adoption of ML. Redshifts can be accurately inferred from photometric measurements of galaxies [e.g. Koo, 1985, 1999, Bolzonella et al., 2000], by training on samples where spectroscopic redshifts are also available. In general, it is more challenging to make the required spectroscopic measurements for large samples of galaxies [Ball and Brunner, 2010], whereas many surveys are able to provide a wealth of features for ML algorithms to learn from. Recent work has

included: comparisons between Gaussian Processes and other machine learning methods – including ANNz [Collister and Lahav, 2004] – from SDSS Data Release 12 Almosallam et al. [2016]; removal of anomalies from training data Hoyle et al. [2015]; application of deep neural networks Hoyle [2016] and SVM Jones and Singal [2017]; and the use of k-means clustering to identify features for input to photometric redshift estimation from SDSS datasets [Stensbo-Smidt et al., 2017]. Multiple machine learning methods have been utilized for determining photometric redshifts for the Dark Energy Survey Science Verification shear catalogue (DES SV) Bonnett et al. [2016]. See also Morrison et al. [2017], Cavuoti et al. [2017a], Leistedt and Hogg [2017] and the review by Salvato et al. [2018]. Cavuoti et al. [2017b] compared multiple machine learning methods with Bayesian and spectral energy template fitting, showing that ML achieved the best accuracy at prediction when there was appropriate coverage by spectroscopic templates. Beck et al. [2017] reported similar outcomes when comparing machine learning with template-fitting approaches, highlighting the “expected bad results” for machine learning methods when no suitable spectroscopic templates were available. Comparison between probability density functions obtained with ANNz2 [Sadeh et al., 2016] and METAPHOR (Machine-learning Estimation Tool for Accurate PHotometric Redshifts) in Amaro et al. [2019].

- *Gravitational lensing.* Concentrations of matter on galactic and cosmological scales bend and deflect the path of light rays from more distance sources. Lensing provides unique probes of dark matter distributions, tests of cosmological models, and magnified views of otherwise faint objects. However, finding lensed systems is observationally challenging. ML has helped in the discovery of previously unknown lensed quasars, e.g. Ostrovski et al. [2017, Gaussian mixture models] and Timlin et al. [2018, RFs/k-NN]. A major challenge for deep learning methods in lens finding is paucity of training data; this has been solved using simulated lenses at galaxy scale. Deep neural nets trained on simulations have resulted in lens discoveries in survey data including the Kilo-Degree Survey [de Jong et al., 2015] by Petrillo et al. [2017, 2019]; and the Dark Energy Survey [Dark Energy Survey Collaboration et al., 2016] by Jacobs et al. [2019b,a]. A strong lens finding challenge was recently conducted using simulated data [Metcalf et al., 2019], and deep learning-based methods outperformed all other methodologies including examination by human experts.
- *Gravitational wave astronomy* The recent detection of gravitational wave signals from coalescing black hole binaries [Abbott et al., 2018], and other related compact systems, has relied on real-time computation and analysis of streams of data from the Advanced Laser Interferometer Gravitational-Wave Observatory (LIGO) detectors [Harry and LIGO Scientific Collaboration, 2010]. By incorporating machine learning, Powell et al. [2017] improved performance in distinguishing between sources and noise signals, along with reducing the latency of the detection pipeline. Zevin et al. [2017] used crowd-sourced categorization of common “glitch” signals in order to train a ML system for real-time glitch classification. George and Huerta [2018b,a] developed Deep Filtering, which utilizes two CNNs for detecting signals (classification) and performing parameter estimation (regression) in real time. Testing first on mock data, they successfully recovered events from LIGO observations. Theoretical insight into binary black hole mergers has also been achieved through ML, training on outputs from numerical relativity simulations [Huerta et al., 2018].

4 Concluding remarks

Every week, new astronomical applications of machine learning and artificial intelligence are added to a growing corpus of work. Random forests, support vector machines, neural networks (artificial, deep, and convolutional), and generative adversarial networks are now having a genuine impact across all domains of astronomy. ML and AI simplify the processes of classification and regression, determination of clustering relationships, forecasting of time-based events, and the generation or reconstruction of missing information.

As methods become more sophisticated, the volume of training data grows, and classifications become more robust, ML and AI allow for new objects to be discovered, and for new scientific insight to be gathered.

The adoption of ML and AI is emerging in planetary studies, investigations of the non-stellar components of the Milky Way and of stellar clusters, and in real-time monitoring of instruments. Elsewhere, rapid progress is occurring in the use of ML and AI for classification and detection of Solar System objects, and the discovery of rare types of active galactic nuclei and quasars. ML, in particular through early experimentation with GANs, offer an intriguing alternative to generating and understanding complex structures in cosmological simulations. ML and AI are now firmly established in: solar astronomy (particularly forecasting of solar activity); the discovery of extra-solar planets and transient objects; and classification, discovery and gaining insights into the properties of all types of stars, variable stars, and stellar evolutionary products (neutron stars, pulsars, and black holes). ML and AI offer new ways to find and understand galaxies, gravitationally-lensed sources, and gravitational wave candidates.

As astronomy moves ever closer to the Exascale data era of the Square Kilometre Array, an increasing number of human-centred tasks and processes are being replaced by faster, automated processing. The adoption of ML and AI techniques is driving a fundamental change in the way future astronomers will approach the process of “discovery”. To date, in the vast majority of cases, discoveries have occurred when astronomers look directly at their data: qualitative inspection supported by quantitative analysis (e.g. model fitting, simulation, etc.). The volume (e.g. number of sources or quantity of data recorded per source) and the complexity (e.g. dimensionality) of data has not vastly exceeded available computing or visual resources. It has been possible to look at the majority of potential sources and false detections by eye, and to conduct visualization and analysis using a desktop-bound workspace. This is no longer the case. Continuous human monitoring of data streams from the SKA is likely to be a tedious task, and discoveries will be missed. Computers excel at such repetitive actions [see, for example, Yeakel et al. [2018], who investigated the role of AI as a means to reduce tedium and detect anomalies in spacecraft systems, through the Cassini-Huygens mission’s study of Saturn’s magnetosphere], and allow astronomers to focus their attention on interpreting and explaining new types of astronomical phenomena and their connection to fundamental physics.

As the use of deep neural networks increases in astronomy (and a great many other fields), the question arises: what is going on inside the networks? Where AI systems are making medical diagnoses and driving autonomous vehicles this may be an urgent question; but an understanding of the errors, biases and limitations is also of growing importance in a scientific context. In the computer vision realm several attempts have been made to develop techniques for visualizing and interpreting deep neural network outputs [see Montavon et al., 2018], for instance visualizing the feature detectors or building “saliency maps” of the important input pixels [Selvaraju et al., 2017]. However, the utility and adequacy of these methods in astronomy, where precisely quantified errors are often required, is far from apparent. Several advancements have been made, for example the use of Bayesian neural networks [Denker and Lecun, 1991] – where the outputs are probability distributions – in estimating gravitational lens model errors [Perreault Levasseur et al., 2017] or the uncertainties in neutron capture mass models [Utama and Piekarewicz, 2017]. Further work in understanding the internals of deep networks will be needed if the promise of deep neural networks for astronomical discovery is to be fully realized.

As progress in artificial intelligence and machine learning accelerates, particularly through advances in deep learning, the gap between human and automated pattern recognition capabilities is narrowing. However, it is still not always obvious why and how classifications or discoveries are made by ever more complex neural networks. There is still scope for more studies that consider the strengths and weaknesses of different ML and AI approaches when applied to a specific dataset – particularly as new, experimental techniques continue to appear. Learning which types of objects are harder to detect or classify also provides insight, along with highlighting potential biases in human input. As Ball and Brunner [2010] stated, and as still holds true, “there is no simple method to select the optimal algorithm to use”. For the time being, traditional statistical methods or visualization are still highly productive first steps, providing astronomers with a detailed understanding of their data. In the future, there is no doubt that the reach and maturity

of machine learning and artificial intelligence in astronomy will continue to grow.

acknowledgements

CJF and CJ thank WIREs for the invitation and opportunity to write this advanced review, and the two anonymous referees for their insightful comments. CJF thanks Wael Farah for helpful discussions. This research has made extensive use of NASA's Astrophysics Data System.

References

- Keivork N. Abazajian and et al. The Seventh Data Release of the Sloan Digital Sky Survey. *ApJSS*, 182: 543–558, Jun 2009. doi: 10.1088/0067-0049/182/2/543.
- T. M. C. Abbott, F. B. Abdalla, A. Alarcon, S. Allam, F. Andrade-Oliveira, J. Annis, S. Avila, M. Banerji, N. Banik, K. Bechtol, R. A. Bernstein, G. M. Bernstein, E. Bertin, D. Brooks, E. Buckley-Geer, D. L. Burke, H. Camacho, A. Carnero Rosell, M. Carrasco Kind, J. Carretero, F. J. Castander, R. Cawthon, K. C. Chan, M. Crocce, C. E. Cunha, C. B. D’Andrea, L. N. da Costa, C. Davis, J. De Vicente, D. L. DePoy, S. Desai, H. T. Diehl, P. Doel, A. Drlica-Wagner, T. F. Eifler, J. Elvin-Poole, J. Estrada, A. E. Evrard, B. Flaugher, P. Fosalba, J. Frieman, J. García-Bellido, E. Gaztanaga, D. W. Gerdes, T. Giannantonio, D. Gruen, R. A. Gruendl, J. Gschwend, G. Gutierrez, W. G. Hartley, D. Hollowood, K. Honscheid, B. Hoyle, B. Jain, D. J. James, T. Jeltema, M. D. Johnson, S. Kent, N. Kokron, E. Krause, K. Kuehn, S. Kuhlmann, N. Kuropatkin, F. Lacasa, O. Lahav, M. Lima, H. Lin, M. A. G. Maia, M. Manera, J. Marriner, J. L. Marshall, P. Martini, P. Melchior, F. Menanteau, C. J. Miller, R. Miquel, J. J. Mohr, E. Neilsen, W. J. Percival, A. A. Plazas, A. Porredon, A. K. Romer, A. Roodman, R. Rosenfeld, A. J. Ross, E. Rozo, E. S. Rykoff, M. Sako, E. Sanchez, B. Santiago, V. Scarpine, R. Schindler, M. Schubnell, S. Serrano, I. Sevilla-Noarbe, E. Sheldon, R. C. Smith, M. Smith, F. Sobreira, E. Suchyta, M. E. C. Swanson, G. Tarle, D. Thomas, M. A. Troxel, D. L. Tucker, V. Vikram, A. R. Walker, R. H. Wechsler, J. Weller, B. Yanny, and Y. Zhang. Dark Energy Survey Year 1 Results: Measurement of the Baryon Acoustic Oscillation scale in the distribution of galaxies to redshift 1. *MNRAS*, 483:4866–4883, Dec 2018. doi: 10.1093/mnras/sty3351.
- Jennifer K. Adelman-McCarthy, Marcel A. Agüeros, Sahar S. Allam, Kurt S. J. Anderson, Scott F. Anderson, James Annis, Neta A. Bahcall, Ivan K. Baldry, J. C. Barentine, Andreas Berlind, Mariangela Bernardi, Michael R. Blanton, William N. Boroski, Howard J. Brewington, Jarle Brinchmann, J. Brinkmann, Robert J. Brunner, Tamás Budavári, Larry N. Carey, Michael A. Carr, Francisco J. Castander, A. J. Connolly, István Csabai, Paul C. Czarapata, Julianne J. Dalcanton, Mamoru Doi, Feng Dong, Daniel J. Eisenstein, Michael L. Evans, Xiaohui Fan, Douglas P. Finkbeiner, Scott D. Friedman, Joshua A. Frieman, Masataka Fukugita, Bruce Gillespie, Karl Glazebrook, Jim Gray, Eva K. Grebel, James E. Gunn, Vijay K. Gurbani, Ernst de Haas, Patrick B. Hall, Frederick H. Harris, Michael Harvanek, Suzanne L. Hawley, Jeffrey Hayes, John S. Hendry, Gregory S. Hennessy, Robert B. Hindsley, Christopher M. Hirata, Craig J. Hogan, David W. Hogg, Donald J. Holmgren, Jon A. Holtzman, Shin-ichi Ichikawa, Željko Ivezić, Sebastian Jester, David E. Johnston, Anders M. Jorgensen, Mario Jurić, Stephen M. Kent, S. J. Kleinman, G. R. Knapp, Alexei Yu. Kniazev, Richard G. Kron, Jurek Krzesinski, Nikolay Kuropatkin, Donald Q. Lamb, Hubert Lampeitl, Brian C. Lee, R. French Leger, Huan Lin, Daniel C. Long, Jon Loveday, Robert H. Lupton, Bruce Margon, David Martínez-Delgado, Rachel Mandelbaum, Takahiko Matsubara, Peregrine M. McGehee, Timothy A. McKay, Avery Meiksin, Jeffrey A. Munn, Reiko Nakajima, Thomas Nash, Jr. Neilsen, Eric H., Heidi Jo Newberg, Peter R. Newman, Robert C. Nichol, Tom Nicinski, Maria Nieto-Santisteban, Atsuko Nitta, William O’Mullane, Sadanori Okamura, Russell Owen, Nikhil Padmanabhan, George Pauls, Jr. Peoples, John, Jeffrey R.

- Pier, Adrian C. Pope, Dimitri Pourbaix, Thomas R. Quinn, Gordon T. Richards, Michael W. Richmond, Constance M. Rockosi, David J. Schlegel, Donald P. Schneider, Joshua Schroeder, Ryan Scranton, Uroš Seljak, Erin Sheldon, Kazu Shimasaku, J. Allyn Smith, Vernesa Smolčić, Stephanie A. Snedden, Chris Stoughton, Michael A. Strauss, Mark SubbaRao, Alexander S. Szalay, István Szapudi, Paula Szkody, Max Tegmark, Aniruddha R. Thakar, Douglas L. Tucker, Alan Uomoto, Daniel E. Vanden Berk, Jan Vandenberg, Michael S. Vogeley, Wolfgang Voges, Nicole P. Vogt, Lucianne M. Walkowicz, David H. Weinberg, Andrew A. West, Simon D. M. White, Yongzhong Xu, Brian Yanny, D. R. Yocum, Donald G. York, Idit Zehavi, Stefano Zibetti, and Daniel B. Zucker. The Fourth Data Release of the Sloan Digital Sky Survey. *ApJSS*, 162(1):38–48, Jan 2006. doi: 10.1086/497917.
- Shankar Agarwal, Romeel Davé, and Bruce A. Bassett. Painting galaxies into dark matter haloes using machine learning. *MNRAS*, 478:3410–3422, Aug 2018. doi: 10.1093/mnras/sty1169.
- Adriano Agnello, Brandon C. Kelly, Tommaso Treu, and Philip J. Marshall. Data mining for gravitationally lensed quasars. *MNRAS*, 448(2):1446–1462, April 2015. ISSN 0035-8711. doi: 10.1093/mnras/stv037.
- M. J. Alger, J. K. Banfield, C. S. Ong, L. Rudnick, O. I. Wong, C. Wolf, H. Andernach, R. P. Norris, and S. S. Shabala. Radio Galaxy Zoo: machine learning for radio source host galaxy cross-identification. *MNRAS*, 478:5547–5563, Aug 2018. doi: 10.1093/mnras/sty1308.
- Ibrahim A. Almosallam, Matt J. Jarvis, and Stephen J. Roberts. GPZ: non-stationary sparse Gaussian processes for heteroscedastic uncertainty estimation in photometric redshifts. *MNRAS*, 462(1):726–739, Oct 2016. doi: 10.1093/mnras/stw1618.
- V. Amaro, S. Cavauoti, M. Brescia, C. Vellucci, G. Longo, M. Bilicki, J. T. A. de Jong, C. Tortora, M. Radovich, N. R. Napolitano, and H. Buddelmeijer. Statistical analysis of probability density functions for photometric redshifts through the KiDS-ESO-DR3 galaxies. *MNRAS*, 482(3):3116–3134, Jan 2019. doi: 10.1093/mnras/sty2922.
- Luca Amendola, Stephen Appleby, David Bacon, Tessa Baker, Marco Baldi, Nicola Bartolo, Alain Blanchard, Camille Bonvin, Stefano Borgani, Enzo Branchini, Clare Burrage, Stefano Camera, Carmelita Carbone, Luciano Casarini, Mark Cropper, Claudia de Rham, Cinzia Di Porto, Anne Ealet, Pedro G. Ferreira, Fabio Finelli, Juan García-Bellido, Tommaso Giannantonio, Luigi Guzzo, Alan Heavens, Lavinia Heisenberg, Catherine Heymans, Henk Hoekstra, Lukas Hollenstein, Rory Holmes, Ole Horst, Knud Jahnke, Thomas D. Kitching, Tomi Koivisto, Martin Kunz, Giuseppe La Vacca, Marisa March, Elisabetta Majerotto, Katarina Markovic, David Marsh, Federico Marulli, Richard Massey, Yannick Mellier, David F. Mota, Nelson J. Nunes, Will Percival, Valeria Pettorino, Cristiano Porciani, Claudia Quercellini, Justin Read, Massimiliano Rinaldi, Domenico Sapone, Roberto Scaramella, Constantinos Skordis, Fergus Simpson, Andy Taylor, Shaun Thomas, Roberto Trotta, Licia Verde, Filippo Vernizzi, Adrian Vollmer, Yun Wang, Jochen Weller, and Tom Zlosnik. Cosmology and Fundamental Physics with the Euclid Satellite. *Living Reviews in Relativity*, 16:6, Sep 2013. doi: 10.12942/lrr-2013-6.
- J. R. P. Angel, P. Wizinowich, M. Lloyd-Hart, and D. Sandler. Adaptive optics for array telescopes using neural-network techniques. *Nature*, 348(6298):221–224, November 1990. ISSN 1476-4687. doi: 10.1038/348221a0.
- A. K. Aniyani and K. Thorat. Classifying Radio Galaxies with the Convolutional Neural Network. *ApJSS*, 230:20, Jun 2017. doi: 10.3847/1538-4365/aa7333.
- M. Araya, M. Mendoza, M. Solar, D. Mardones, and A. Bayo. Unsupervised learning of structure in spectroscopic cubes. *A&C*, 24:25–35, Jul 2018. doi: 10.1016/j.ascom.2018.06.001.

- D. J. Armstrong, J. Kirk, K. W. F. Lam, J. McCormac, H. P. Osborn, J. Spake, S. Walker, D. J. A. Brown, M. H. Kristiansen, D. Pollacco, R. West, and P. J. Wheatley. K2 variable catalogue – II. Machine learning classification of variable stars and eclipsing binaries in K2 fields 0–4. *MNRAS*, 456(2): 2260–2272, February 2016. ISSN 0035-8711. doi: 10.1093/mnras/stv2836.
- D. J. Armstrong, D. Pollacco, and A. Santerne. Transit shapes and self-organizing maps as a tool for ranking planetary candidates: application to Kepler and K2. *MNRAS*, 465:2634–2642, Mar 2017. doi: 10.1093/mnras/stw2881.
- David J. Armstrong, Maximilian N. Günther, James McCormac, Alexis M. S. Smith, Daniel Bayliss, François Bouchy, Matthew R. Burleigh, Sarah Casewell, Philipp Eigmüller, Edward Gillen, Michael R. Goad, Simon T. Hodgkin, James S. Jenkins, Tom Loudon, Lionel Metrailler, Don Pollacco, Katja Poppenhaeger, Didier Queloz, Liam Raynard, Heike Rauer, Stéphane Udry, Simon R. Walker, Christopher A. Watson, Richard G. West, and Peter J. Wheatley. Automatic vetting of planet candidates from ground-based surveys: machine learning with NGTS. *MNRAS*, 478:4225–4237, Aug 2018. doi: 10.1093/mnras/sty1313.
- N. M. Ball, J. Loveday, M. Fukugita, O. Nakamura, S. Okamura, J. Brinkmann, and R. J. Brunner. Galaxy types in the Sloan Digital Sky Survey using supervised artificial neural networks. *MNRAS*, 348 (3):1038–1046, March 2004. ISSN 0035-8711. doi: 10.1111/j.1365-2966.2004.07429.x.
- Nicholas M. Ball and Robert J. Brunner. Data Mining and Machine Learning in Astronomy. *International Journal of Modern Physics D*, 19:1049–1106, January 2010. doi: 10.1142/S0218271810017160.
- Dalya Baron and Dovi Poznanski. The weirdest SDSS galaxies: Results from an outlier detection algorithm. *arXiv:1611.07526*, November 2016.
- B. R. Barsdell, D. G. Barnes, and C. J. Fluke. Analysing astronomy algorithms for graphics processing units and beyond. *MNRAS*, 408:1936–1944, Nov 2010. doi: 10.1111/j.1365-2966.2010.17257.x.
- Melanie R. Beck, Claudia Scarlata, Lucy F. Fortson, Chris J. Lintott, B. D. Simmons, Melanie A. Galloway, Kyle W. Willett, Hugh Dickinson, Karen L. Masters, Philip J. Marshall, and Darryl Wright. Integrating human and machine intelligence in galaxy morphology classification tasks. *MNRAS*, 476:5516–5534, Jun 2018. doi: 10.1093/mnras/sty503.
- R. Beck, C. A. Lin, E. E. O. Ishida, F. Gieseke, R. S. de Souza, M. V. Costa-Duarte, M. W. Hattab, and A. Krone-Martins. On the realistic validation of photometric redshifts. *MNRAS*, 468:4323–4339, Jul 2017. doi: 10.1093/mnras/stx687.
- Gordon Bell, Tony Hey, and Alex Szalay. Beyond the data deluge. *Science*, 323(5919):1297–1298, 2009. ISSN 0036-8075. doi: 10.1126/science.1170411.
- Eric C. Bellm, Shrinivas R. Kulkarni, Matthew J. Graham, Richard Dekany, Roger M. Smith, Reed Riddle, Frank J. Masci, George Helou, Thomas A. Prince, Scott M. Adams, C. Barbarino, Tom Barlow, James Bauer, Ron Beck, Justin Belicki, Rahul Biswas, Nadejda Blagorodnova, Dennis Bodewits, Bryce Bolin, Valery Brinnel, Tim Brooke, Brian Bue, Mattia Bulla, Rick Burruss, S. Bradley Cenko, Chan-Kao Chang, Andrew Connolly, Michael Coughlin, John Cromer, Virginia Cunningham, Kishalay De, Alex Delacroix, Vandana Desai, Dmitry A. Duev, Gwendolyn Eadie, Tony L. Farnham, Michael Feeney, Ulrich Feindt, David Flynn, Anna Franckowiak, S. Frederick, C. Fremling, Avishay Gal-Yam, Suvi Gezari, Matteo Giomi, Daniel A. Goldstein, V. Zach Golkhou, Ariel Goobar, Steven Groom, Eugene Hecopians, David Hale, John Henning, Anna Y. Q. Ho, David Hover, Justin Howell, Tiara Hung, Daniela Huppenkothen, David Imel, Wing-Huen Ip, Željko Ivezić, Edward Jackson, Lynne Jones, Mario Juric, Mansi M. Kasliwal, S. Kaspi, Stephen Kaye, Michael S. P. Kelley, Marek Kowalski, Emily Kramer, Thomas Kupfer, Walter

- Landry, Russ R. Laher, Chien-De Lee, Hsing Wen Lin, Zhong-Yi Lin, Ragnhild Lunnan, Matteo Giomi, Ashish Mahabal, Peter Mao, Adam A. Miller, Serge Monkenwitz, Patrick Murphy, Chow-Choong Ngeow, Jakob Nordin, Peter Nugent, Eran Ofek, Maria T. Patterson, Bryan Penprase, Michael Porter, Ludwig Rauch, Umaa Rebbapragada, Dan Reiley, Mickael Rigault, Hector Rodriguez, Jan van Roestel, Ben Rusholme, Jakob van Santen, S. Schulze, David L. Shupe, Leo P. Singer, Maayane T. Soumagnac, Robert Stein, Jason Surace, Jesper Sollerman, Paula Szkody, F. Taddia, Scott Terek, Angela Van Sistine, Sjoert van Velzen, W. Thomas Vestrand, Richard Walters, Charlotte Ward, Quan-Zhi Ye, Po-Chieh Yu, Lin Yan, and Jeffry Zolkower. The Zwicky Transient Facility: System Overview, Performance, and First Results. *PASP*, 131(995):018002, Jan 2019. doi: 10.1088/1538-3873/aaecbe.
- Patricio Benavente, Pavlos Protopapas, and Karim Pichara. Automatic Survey-invariant Classification of Variable Stars. *ApJ*, 845:147, Aug 2017. doi: 10.3847/1538-4357/aa7f2d.
- Federico Benvenuto, Michele Piana, Cristina Campi, and Anna Maria Massone. A Hybrid Supervised/Unsupervised Machine Learning Approach to Solar Flare Prediction. *ApJ*, 853:90, Jan 2018. doi: 10.3847/1538-4357/aaa23c.
- E. Bertin and S. Arnouts. SExtractor: Software for source extraction. *A&ASS*, 117:393, June 1996. doi: 10.1051/aas:1996164.
- S. Bethapudi and S. Desai. Separation of pulsar signals from noise using supervised machine learning algorithms. *A&C*, 23:15–26, Apr 2018. doi: 10.1016/j.ascom.2018.02.002.
- M. Bilicki, H. Hoekstra, M. J. I. Brown, V. Amaro, C. Blake, S. Caviuoti, J. T. A. de Jong, C. Georgiou, H. Hildebrandt, C. Wolf, A. Amon, M. Brescia, S. Brough, M. V. Costa-Duarte, T. Erben, K. Glazebrook, A. Grado, C. Heymans, T. Jarrett, S. Joudaki, K. Kuijken, G. Longo, N. Napolitano, D. Parkinson, C. Vellucci, G. A. Verdoes Kleijn, and L. Wang. Photometric redshifts for the Kilo-Degree Survey. Machine-learning analysis with artificial neural networks. *A&A*, 616:A69, August 2018. doi: 10.1051/0004-6361/201731942.
- Rahul Biswas, Lindy Blackburn, Junwei Cao, Reed Essick, Kari Alison Hodge, Erotokritos Katsavounidis, Kyungmin Kim, Young-Min Kim, Eric-Olivier Le Bigot, Chang-Hwan Lee, John J. Oh, Sang Hoon Oh, Edwin J. Son, Ye Tao, Ruslan Vaulin, and Xiaoge Wang. Application of machine learning algorithms to the study of noise artifacts in gravitational-wave data. *Phys. Rev. D*, 88(6):062003, September 2013. doi: 10.1103/PhysRevD.88.062003.
- M. Bolzonella, J. M. Miralles, and R. Pelló. Photometric redshifts based on standard SED fitting procedures. *A&A*, 363:476–492, Nov 2000.
- C. Bonnett, M. A. Troxel, W. Hartley, A. Amara, B. Leistedt, M. R. Becker, G. M. Bernstein, S. L. Bridle, C. Bruderer, M. T. Busha, M. Carrasco Kind, M. J. Childress, F. J. Castander, C. Chang, M. Crocce, T. M. Davis, T. F. Eifler, J. Frieman, C. Gangkofner, E. Gaztanaga, K. Glazebrook, D. Gruen, T. Kacprzak, A. King, J. Kwan, O. Lahav, G. Lewis, C. Lidman, H. Lin, N. MacCrann, R. Miquel, C. R. O’Neill, A. Palmese, H. V. Peiris, A. Refregier, E. Roza, E. S. Rykoff, I. Sadeh, C. Sánchez, E. Sheldon, S. Uddin, R. H. Wechsler, J. Zuntz, T. Abbott, F. B. Abdalla, S. Allam, R. Armstrong, M. Banerji, A. H. Bauer, A. Benoit-Lévy, E. Bertin, D. Brooks, E. Buckley-Geer, D. L. Burke, D. Capozzi, A. Carnero Rosell, J. Carretero, C. E. Cunha, C. B. D’Andrea, L. N. da Costa, D. L. DePoy, S. Desai, H. T. Diehl, J. P. Dietrich, P. Doel, A. Fausti Neto, E. Fernandez, B. Flaugher, P. Fosalba, D. W. Gerdes, R. A. Gruendl, K. Honscheid, B. Jain, D. J. James, M. Jarvis, A. G. Kim, K. Kuehn, N. Kuropatkin, T. S. Li, M. Lima, M. A. G. Maia, M. March, J. L. Marshall, P. Martini, P. Melchior, C. J. Miller, E. Neilsen, R. C. Nichol, B. Nord, R. Ogando, A. A. Plazas, K. Reil, A. K. Romer, A. Roodman, M. Sako, E. Sanchez, B. Santiago, R. C. Smith, M. Soares-Santos, F. Sobreira, E. Suchyta, M. E. C. Swanson,

- G. Tarle, J. Thaler, D. Thomas, V. Vikram, A. R. Walker, and Dark Energy Survey Collaboration. Redshift distributions of galaxies in the Dark Energy Survey Science Verification shear catalogue and implications for weak lensing. *PRD*, 94(4):042005, Aug 2016. doi: 10.1103/PhysRevD.94.042005.
- R. S. Booth, W. J. G. de Blok, J. L. Jonas, and B. Fanaroff. MeerKAT Key Project Science, Specifications, and Proposals. art. arXiv:0910.2935, Oct 2009.
- Kirk Borne. Scientific Data Mining in Astronomy. In H. Kargupta, J. Han, P.S. Yu, R. Motwani, and V. Kumar, editors, *Next Generation of Data Mining*, pages 91–114, 2009.
- William J. Borucki and et al. Kepler Planet-Detection Mission: Introduction and First Results. *Science*, 327:977, Feb 2010. doi: 10.1126/science.1185402.
- Leo Breiman. Bagging predictors. *Mach Learn*, 24(2):123–140, August 1996. ISSN 1573-0565. doi: 10.1007/BF00058655.
- Leo Breiman. Random Forests. *Machine Learning*, 45(1):5–32, October 2001. ISSN 1573-0565. doi: 10.1023/A:1010933404324.
- M. Brescia, S. Cavuoti, R. D’Abrusco, G. Longo, and A. Mercurio. Photometric Redshifts for Quasars in Multi-band Surveys. *ApJ*, 772(2):140, Aug 2013. doi: 10.1088/0004-637X/772/2/140.
- Massimo Brescia, Stefano Cavuoti, Giuseppe Longo, Alfonso Nocella, Mauro Garofalo, Francesco Manna, Francesco Esposito, Giovanni Albano, Marisa Guglielmo, Giovanni D’Angelo, Alessandro Di Guido, S. George Djorgovski, Ciro Donalek, Ashish A. Mahabal, Matthew J. Graham, Michelangelo Fiore, and Raffaele D’Abrusco. DAMEWARE: A Web Cyberinfrastructure for Astrophysical Data Mining. *PASP*, 126(942):783, Aug 2014. doi: 10.1086/677725.
- Massimo Brescia, Stefano Cavuoti, Francesco Esposito, Michelangelo Fiore, Mauro Garofalo, Marisa Guglielmo, Giuseppe Longo, Francesco Manna, Alfonso Nocella, and Civita Vellucci. DAMEWARE - Data Mining & Exploration Web Application Resource. art. arXiv:1603.00720, Mar 2016.
- Emeric Bron, Chloé Daudon, Jérôme Pety, François Levrier, Maryvonne Gerin, Pierre Gratier, Jan H. Orkisz, Viviana Guzman, Sébastien Bardeau, Javier R. Goicoechea, Harvey Liszt, Karin Öberg, Nicolas Peretto, Albrecht Sievers, and Pascal Tremblin. Clustering the Orion B giant molecular cloud based on its molecular emission. *A&A*, 610:A12, Feb 2018. doi: 10.1051/0004-6361/201731833.
- Robert J. Brunner, S. George Djorgovski, Thomas A. Prince, and Alex S. Szalay. *Massive Datasets in Astronomy*, pages 931–979. Springer US, Boston, MA, 2002. ISBN 978-1-4615-0005-6.
- Yude Bu, Zhenxin Lei, Gang Zhao, Jingde Bu, and Jingchang Pan. Searching for Hot Subdwarf Stars from the LAMOST Spectra. I. Method. *ApJSS*, 233:2, Nov 2017. doi: 10.3847/1538-4365/aa91cd.
- Guillermo Cabrera-Vives, Christopher J. Miller, and Jeff Schneider. Systematic Labeling Bias in Galaxy Morphologies. *AJ*, 156:284, Dec 2018. doi: 10.3847/1538-3881/aae9f4.
- Matias Carrasco Kind and Robert J. Brunner. TPZ: Photometric redshift PDFs and ancillary information by using prediction trees and random forests. *MNRAS*, 432(2):1483–1501, June 2013. ISSN 0035-8711. doi: 10.1093/mnras/stt574.
- Matias Carrasco Kind and Robert J. Brunner. SOMz: Photometric redshift PDFs with self-organizing maps and random atlas. *MNRAS*, 438(4):3409–3421, March 2014. ISSN 0035-8711. doi: 10.1093/mnras/stt2456.

- A. Castro-Ginard, C. Jordi, X. Luri, F. Julbe, M. Morvan, L. Balaguer-Núñez, and T. Cantat- Gaudin. A new method for unveiling open clusters in Gaia. New nearby open clusters confirmed by DR2. *A&A*, 618:A59, October 2018. doi: 10.1051/0004-6361/201833390.
- S. Cavuoti, M. Brescia, G. Longo, and A. Mercurio. Photometric redshifts with the quasi Newton algorithm (MLPQNA) Results in the PHAT1 contest. *A&A*, 546:A13, Oct 2012. doi: 10.1051/0004-6361/201219755.
- S. Cavuoti, V. Amaro, M. Brescia, C. Vellucci, C. Tortora, and G. Longo. METAPHOR: a machine-learning-based method for the probability density estimation of photometric redshifts. *MNRAS*, 465: 1959–1973, Feb 2017a. doi: 10.1093/mnras/stw2930.
- S. Cavuoti, C. Tortora, M. Brescia, G. Longo, M. Radovich, N. R. Napolitano, V. Amaro, C. Vellucci, F. La Barbera, F. Getman, and A. Grado. A cooperative approach among methods for photometric redshifts estimation: an application to KiDS data. *MNRAS*, 466:2039–2053, Apr 2017b. doi: 10.1093/mnras/stw3208.
- B. Q. Chen, Y. Huang, H. B. Yuan, C. Wang, D. W. Fan, M. S. Xiang, H. W. Zhang, Z. J. Tian, and X. W. Liu. Three-dimensional interstellar dust reddening maps of the Galactic plane. *MNRAS*, 483: 4277–4289, Mar 2019. doi: 10.1093/mnras/sty3341.
- Ying-Tung Chen, Hsing-Wen Lin, Mike Alexandersen, Matthew J. Lehner, Shiang-Yu Wang, Jen-Hung Wang, Fumi Yoshida, Yutaka Komiyama, and Satoshi Miyazaki. Searching for moving objects in HSC-SSP: Pipeline and preliminary results. *PASJ*, 70:S38, Jan 2018. doi: 10.1093/pasj/psx145.
- Igor Chilingarian, Véronique Cayatte, Yves Revaz, Serguei Dodonov, Daniel Durand, Florence Durret, Alberto Micol, and Eric Slezak. A Population of Compact Elliptical Galaxies Detected with the Virtual Observatory. *Science*, 326(5958):1379, Dec 2009. doi: 10.1126/science.1175930.
- R. Chyla, A. Accomazzi, A. Holachek, C. S. Grant, J. Elliott, E. A. Henneken, D. M. Thompson, M. J. Kurtz, S. S. Murray, and V. Sudilovsky. ADS 2.0: New Architecture, API and Services. In A. R. Taylor and E. Rosolowsky, editors, *Astronomical Data Analysis Software and Systems XXIV (ADASS XXIV)*, volume 495 of *Astronomical Society of the Pacific Conference Series*, page 401, Sep 2015.
- Razvan Ciuca and Oscar F. Hernández. A Bayesian framework for cosmic string searches in CMB maps. *Journal of Cosmology and Astro-Particle Physics*, 2017:028, Aug 2017. doi: 10.1088/1475-7516/2017/08/028.
- Judith G. Cohen, Branimir Sesar, Sophianna Bahnlzer, Kevin He, Shrinivas R. Kulkarni, Thomas A. Prince, Eric Bellm, and Russ R. Laher. The Outer Halo of the Milky Way as Probed by RR Lyr Variables from the Palomar Transient Facility. *ApJ*, 849:150, Nov 2017. doi: 10.3847/1538-4357/aa9120.
- Adrian A. Collister and Ofer Lahav. ANNz: Estimating Photometric Redshifts Using Artificial Neural Networks. *PASP*, 116(818):345–351, Apr 2004. doi: 10.1086/383254.
- Liam Connor and Joeri van Leeuwen. Applying Deep Learning to Fast Radio Burst Classification. *AJ*, 156:256, Dec 2018. doi: 10.3847/1538-3881/aae649.
- Christopher J. Conselice. The fundamental properties of galaxies and a new galaxy classification system. *MNRAS*, 373(4):1389–1408, December 2006. ISSN 0035-8711. doi: 10.1111/j.1365-2966.2006.11114.x.
- Corinna Cortes and Vladimir Vapnik. Support-vector networks. *Mach Learn*, 20(3):273–297, September 1995. ISSN 0885-6125, 1573-0565. doi: 10.1007/BF00994018.

- Dark Energy Survey Collaboration, T. Abbott, F. B. Abdalla, J. Aleksić, S. Allam, A. Amara, D. Bacon, E. Balbinot, M. Banerji, K. Bechtol, A. Benoit-Lévy, G. M. Bernstein, E. Bertin, J. Blazek, C. Bonnett, S. Bridle, D. Brooks, R. J. Brunner, E. Buckley-Geer, D. L. Burke, G. B. Caminha, D. Capozzi, J. Carlsen, A. Carnero-Rosell, M. Carollo, M. Carrasco-Kind, J. Carretero, F. J. Castander, L. Clerkin, T. Collett, C. Conselice, M. Crocce, C. E. Cunha, C. B. D’Andrea, L. N. da Costa, T. M. Davis, S. Desai, H. T. Diehl, J. P. Dietrich, S. Dodelson, P. Doel, A. Drlica-Wagner, J. Estrada, J. Etherington, A. E. Evrard, J. Fabbri, D. A. Finley, B. Flaugher, R. J. Foley, P. Fosalba, J. Frieman, J. García-Bellido, E. Gaztanaga, D. W. Gerdes, T. Giannantonio, D. A. Goldstein, D. Gruen, R. A. Gruendl, P. Guarnieri, G. Gutierrez, W. Hartley, K. Honscheid, B. Jain, D. J. James, T. Jeltema, S. Jouvel, R. Kessler, A. King, D. Kirk, R. Kron, K. Kuehn, N. Kuropatkin, O. Lahav, T. S. Li, M. Lima, H. Lin, M. A. G. Maia, M. Makler, M. Manera, C. Maraston, J. L. Marshall, P. Martini, R. G. McMahon, P. Melchior, A. Merson, C. J. Miller, R. Miquel, J. J. Mohr, X. Morice-Atkinson, K. Naidoo, E. Neilsen, R. C. Nichol, B. Nord, R. Ogando, F. Ostrovski, A. Palmese, A. Papadopoulos, H. V. Peiris, J. Peoples, W. J. Percival, A. A. Plazas, S. L. Reed, A. Refregier, A. K. Romer, A. Roodman, A. Ross, E. Roza, E. S. Rykoff, I. Sadeh, M. Sako, C. Sánchez, E. Sanchez, B. Santiago, V. Scarpine, M. Schubnell, I. Sevilla-Noarbe, E. Sheldon, M. Smith, R. C. Smith, M. Soares-Santos, F. Sobreira, M. Soumagnac, E. Suchyta, M. Sullivan, M. Swanson, G. Tarle, J. Thaler, D. Thomas, R. C. Thomas, D. Tucker, J. D. Vieira, V. Vikram, A. R. Walker, R. H. Wechsler, J. Weller, W. Wester, L. Whiteway, H. Wilcox, B. Yanny, Y. Zhang, and J. Zuntz. The Dark Energy Survey: More than dark energy - an overview. *MNRAS*, 460: 1270–1299, August 2016. ISSN 0035-8711. doi: 10.1093/mnras/stw641.
- Jelte T. A. de Jong, Gijs A. Verdoes Kleijn, Konrad H. Kuijken, and Edwin A. Valentijn. The Kilo-Degree Survey. *Experimental Astronomy*, 35(1-2):25–44, Jan 2013. doi: 10.1007/s10686-012-9306-1.
- Jelte T. A. de Jong, Gijs A. Verdoes Kleijn, Danny R. Boxhoorn, Hugo Buddelmeijer, Massimo Capaccioli, Fedor Getman, Aniello Grado, Ewout Helmich, Zhuoyi Huang, Nancy Irisarri, Konrad Kuijken, Francesco La Barbera, John P. McFarland, Nicola R. Napolitano, Mario Radovich, Gert Sikkema, Edwin A. Valentijn, Kor G. Begeman, Massimo Brescia, Stefano Cavuoti, Ami Choi, Oliver-Mark Cordes, Giovanni Covone, Massimo Dall’Ora, Hendrik Hildebrandt, Giuseppe Longo, Reiko Nakajima, Maurizio Paolillo, Emanuella Puddu, Agatino Rifatto, Crescenzo Tortora, Edo van Uitert, Axel Buddendiek, Joachim Harnois-Déraps, Thomas Erben, Martin B. Eriksen, Catherine Heymans, Henk Hoekstra, Benjamin Joachimi, Thomas D. Kitching, Dominik Klaes, Léon V. E. Koopmans, Fabian Köhlinger, Nivya Roy, Cristóbal Sifón, Peter Schneider, Will J. Sutherland, Massimo Viola, and Willem-Jan Vriend. The first and second data releases of the Kilo-Degree Survey. *A&A*, 582:A62, October 2015. doi: 10.1051/0004-6361/201526601.
- J. de León, J. Licandro, M. Serra-Ricart, N. Pinilla-Alonso, and H. Campins. Observations, compositional, and physical characterization of near-Earth and Mars-crosser asteroids from a spectroscopic survey. *A&A*, 517:A23, July 2010. ISSN 0004-6361, 1432-0746. doi: 10.1051/0004-6361/200913852.
- H. Dénes, N. M. McClure-Griffiths, J. M. Dickey, J. R. Dawson, and C. E. Murray. Calibrating the HISA temperature: Measuring the temperature of the Riegel-Crutcher cloud. *MNRAS*, 479:1465–1490, Sep 2018. doi: 10.1093/mnras/sty1384.
- John Denker and Yann Lecun. Transforming Neural-Net Output Levels to Probability Distributions. In *Advances in Neural Information Processing Systems 3*, pages 853–859. Morgan Kaufmann, 1991.
- P. E. Dewdney, P. J. Hall, R. T. Schilizzi, and T. J. L. W. Lazio. The Square Kilometre Array. *IEEE Proceedings*, 97:1482–1496, Aug 2009. doi: 10.1109/JPROC.2009.2021005.
- Foivos I. Diakogiannis, Geraint F. Lewis, Rodrigo A. Ibata, Magda Guglielmo, Mark I. Wilkinson, and Chris Power. Reliable mass calculation in spherical gravitating systems. *MNRAS*, 482:3356–3372, Jan 2019. doi: 10.1093/mnras/sty2931.

- Sander Dieleman, Kyle W. Willett, and Joni Dambre. Rotation-invariant convolutional neural networks for galaxy morphology prediction. *MNRAS*, 450:1441–1459, June 2015. ISSN 0035-8711. doi: 10.1093/mnras/stv632.
- G. Djorgovski. Astrophysics in the Era of Massive Time-Domain Surveys. In P. R. Wozniak, M. J. Graham, A. A. Mahabal, and R. Seaman, editors, *The Third Hot-wiring the Transient Universe Workshop*, pages 215–215, Jan 2014.
- S.G. Djorgovski, M.J. Graham, C. Donalek, A.A. Mahabal, A.J. Drake, M. Turmon, and T. Fuchs. Real-time data mining of massive data streams from synoptic sky surveys. *Future Gener. Comput. Syst.*, 59 (C):95–104, June 2016. ISSN 0167-739X. doi: 10.1016/j.future.2015.10.013.
- H. Domínguez Sánchez, M. Huertas-Company, M. Bernardi, D. Tuccillo, and J. L. Fischer. Improving galaxy morphologies for SDSS with Deep Learning. *MNRAS*, 476:3661–3676, May 2018. doi: 10.1093/mnras/sty338.
- A. J. Drake, S. G. Djorgovski, A. Mahabal, E. Beshore, S. Larson, M. J. Graham, R. Williams, E. Christensen, M. Catelan, A. Boattini, A. Gibbs, R. Hill, and R. Kowalski. First Results from the Catalina Real-Time Transient Survey. *ApJ*, 696(1):870–884, May 2009. doi: 10.1088/0004-637X/696/1/870.
- Dmitry A. Duev, Ashish Mahabal, Quanzhi Ye, Kushal Tirumala, Justin Belicki, Richard Dekany, Sara Frederick, Matthew J. Graham, Russ R. Laher, Frank J. Masci, Thomas A. Prince, Reed Riddle, Philippe Rosnet, and Maayane T. Soumagnac. DeepStreaks: identifying fast-moving objects in the Zwicky Transient Facility data with deep learning. *MNRAS*, 486(3):4158–4165, Jul 2019. doi: 10.1093/mnras/stz1096.
- R. P. Eatough, N. Molkenhuth, M. Kramer, A. Noutsos, M. J. Keith, B. W. Stappers, and A. G. Lyne. Selection of radio pulsar candidates using artificial neural networks. *MNRAS*, 407(4):2443–2450, October 2010. ISSN 0035-8711. doi: 10.1111/j.1365-2966.2010.17082.x.
- N. Erasmus, M. Mommert, D. E. Trilling, A. A. Sickafoose, C. van Gend, and J. L. Hora. Characterization of Near-Earth Asteroids Using KMTNET-SAAO. *AJ*, 154:162, Oct 2017. doi: 10.3847/1538-3881/aa88be.
- N. Erasmus, A. McNeill, M. Mommert, D. E. Trilling, A. A. Sickafoose, and C. van Gend. Taxonomy and Light-curve Data of 1000 Serendipitously Observed Main-belt Asteroids. *ApJSS*, 237:19, Jul 2018. doi: 10.3847/1538-4365/aac38f.
- Ross Fadely, David W. Hogg, and Beth Willman. STAR-GALAXY CLASSIFICATION IN MULTI-BAND OPTICAL IMAGING. *ApJ*, 760(1):15, October 2012. ISSN 0004-637X. doi: 10.1088/0004-637X/760/1/15.
- W. Farah, C. Flynn, M. Bailes, A. Jameson, K. W. Bannister, E. D. Barr, T. Bateman, S. Bhandari, M. Caleb, D. Campbell-Wilson, S. W. Chang, A. Deller, A. J. Green, R. Hunstead, F. Jankowski, E. Keane, J. P. Macquart, A. Möller, C. A. Onken, S. Osłowski, A. Parthasarathy, K. Plant, V. Ravi, R. M. Shannon, B. E. Tucker, V. Venkatraman Krishnan, and C. Wolf. FRB microstructure revealed by the real-time detection of FRB170827. *MNRAS*, 478:1209–1217, July 2018. doi: 10.1093/mnras/sty1122.
- U. Fayyad, G. Piatetsky-Shapiro, and P. Smyth. From data mining to knowledge discovery in databases. *AI Magazine*, 17(3):37–53, 1996.
- E.D. Feigelson and G.J. Babu. Big data in astronomy. *Significance*, 9(4):22–25, 2012. doi: 10.1111/j.1740-9713.2012.00587.x.
- Andrew E. Firth, Ofer Lahav, and Rachel S. Somerville. Estimating photometric redshifts with artificial neural networks. *MNRAS*, 339(4):1195–1202, March 2003. ISSN 0035-8711. doi: 10.1046/j.1365-8711.2003.06271.x.

- Kostas Florios, Ioannis Kontogiannis, Sung-Hong Park, Jordan A. Guerra, Federico Benvenuto, D. Shaun Bloomfield, and Manolis K. Georgoulis. Forecasting Solar Flares Using Magnetogram-based Predictors and Machine Learning. *SoPh*, 293:28, Feb 2018. doi: 10.1007/s11207-018-1250-4.
- Christopher J. Fluke, David G. Barnes, Benjamin R. Barsdell, and Amr H. Hassan. Astrophysical Supercomputing with GPUs: Critical Decisions for Early Adopters. *PASA*, 28:15–27, Jan 2011. doi: 10.1071/AS10019.
- Paul J. Francis, Paul C. Hewett, Craig B. Foltz, and Frederic H. Chaffee. An Objective Classification Scheme for QSO Spectra. *ApJ*, 398:476, October 1992. ISSN 0004-637X. doi: 10.1086/171870.
- K. Decker French and Ann I. Zabludoff. Identifying Tidal Disruption Events via Prior Photometric Selection of Their Preferred Hosts. *ApJ*, 868:99, Dec 2018. doi: 10.3847/1538-4357/aaea64.
- Yoav Freund and Robert E. Schapire. A decision-theoretic generalization of on-line learning and an application to boosting. In Paul Vitányi, editor, *Computational Learning Theory*, Lecture Notes in Computer Science, pages 23–37. Springer Berlin Heidelberg, 1995. ISBN 978-3-540-49195-8.
- Jerome H. Friedman. Greedy Function Approximation: A Gradient Boosting Machine. *The Annals of Statistics*, 29(5):1189–1232, 2001. ISSN 0090-5364.
- J. Frontera-Pons, F. Sureau, J. Bobin, and E. Le Floch. Unsupervised feature-learning for galaxy SEDs with denoising autoencoders. *A&A*, 603:A60, Jul 2017. doi: 10.1051/0004-6361/201630240.
- Yuki Fujimoto, Kenji Fukushima, and Koichi Murase. Methodology study of machine learning for the neutron star equation of state. *PhysRevD*, 98:023019, Jul 2018. doi: 10.1103/PhysRevD.98.023019.
- Kunihiko Fukushima. Neocognitron: A self-organizing neural network model for a mechanism of pattern recognition unaffected by shift in position. *Biol. Cybernetics*, 36(4):193–202, April 1980. ISSN 0340-1200, 1432-0770. doi: 10.1007/BF00344251.
- Levi Fussell and Ben Moews. Forging new worlds: high-resolution synthetic galaxies with chained generative adversarial networks. *MNRAS*, 485:3203–3214, Mar 2019. doi: 10.1093/mnras/stz602.
- Gaia Collaboration and et al. The Gaia mission. *A&A*, 595:A1, Nov 2016a. doi: 10.1051/0004-6361/201629272.
- Gaia Collaboration and et al. Gaia Data Release 1. Summary of the astrometric, photometric, and survey properties. *A&A*, 595:A2, Nov 2016b. doi: 10.1051/0004-6361/201629512.
- Gaia Collaboration and et al. Gaia Data Release 2. Summary of the contents and survey properties. *A&A*, 616:A1, Aug 2018. doi: 10.1051/0004-6361/201833051.
- Xin-Hua Gao. Memberships, distance and proper-motion of the open cluster NGC 188 based on a machine learning method. *Ap&SS*, 363:232, Nov 2018b. doi: 10.1007/s10509-018-3453-4.
- Xinhua Gao. A Machine-learning-based Investigation of the Open Cluster M67. *ApJ*, 869:9, Dec 2018a. doi: 10.3847/1538-4357/aae8dd.
- Rafael Garcia-Dias, Carlos Allende Prieto, Jorge Sánchez Almeida, and Ignacio Ordovás-Pascual. Machine learning in APOGEE. Unsupervised spectral classification with K-means. *A&A*, 612:A98, May 2018. doi: 10.1051/0004-6361/201732134.
- Daniel George and E. A. Huerta. Deep neural networks to enable real-time multimessenger astrophysics. *PRD*, 97:044039, February 2018a. doi: 10.1103/PhysRevD.97.044039.

- Daniel George and E. A. Huerta. Deep Learning for real-time gravitational wave detection and parameter estimation: Results with Advanced LIGO data. *Physics Letters B*, 778:64–70, March 2018b. doi: 10.1016/j.physletb.2017.12.053.
- Ryusei Gichu and Kazunori Ogohara. Segmentation of dust storm areas on Mars images using principal component analysis and neural network. *Progress in Earth and Planetary Science*, 6:19, Feb 2019. doi: 10.1186/s40645-019-0266-1.
- Fabian Gieseke, Steven Bloemen, Cas van den Bogaard, Tom Heskes, Jonas Kindler, Richard A. Scalzo, Valério A. R. M. Ribeiro, Jan van Roestel, Paul J. Groot, Fang Yuan, Anais Möller, and Brad E. Tucker. Convolutional neural networks for transient candidate vetting in large-scale surveys. *MNRAS*, 472:3101–3114, Dec 2017. doi: 10.1093/mnras/stx2161.
- Daniel Giles and Lucianne Walkowicz. Systematic serendipity: a test of unsupervised machine learning as a method for anomaly detection. *MNRAS*, 484:834–849, Mar 2019. doi: 10.1093/mnras/sty3461.
- C. A. Gomez Gonzalez, O. Absil, and M. Van Droogenbroeck. Supervised detection of exoplanets in high-contrast imaging sequences. *A&A*, 613:A71, Jun 2018. doi: 10.1051/0004-6361/201731961.
- J. A. González and F. S. Guzmán. Characterizing the velocity of a wandering black hole and properties of the surrounding medium using convolutional neural networks. *PhysRevD*, 97:063001, Mar 2018. doi: 10.1103/PhysRevD.97.063001.
- E. A. González-Solares, N. A. Walton, R. Greimel, J. E. Drew, M. J. Irwin, S. E. Sale, K. Andrews, A. Aungwerojwit, M. J. Barlow, E. van den Besselaar, R. L. M. Corradi, B. T. Gänsicke, P. J. Groot, A. S. Hales, E. C. Hopewell, Haili Hu, J. Irwin, C. Knigge, E. Lagadec, P. Leisy, J. R. Lewis, A. Mampaso, M. Matsuura, B. Moont, L. Morales-Rueda, R. A. H. Morris, T. Naylor, Q. A. Parker, P. Prema, S. Pyrzas, G. T. Rixon, P. Rodríguez-Gil, G. Roelofs, L. Sabin, I. Skillen, J. Suso, R. Tata, K. Viironen, J. S. Vink, A. Witham, N. J. Wright, A. A. Zijlstra, A. Zurita, J. Drake, J. Fabregat, D. J. Lennon, P. W. Lucas, E. L. Martín, S. Phillipps, D. Steeghs, and Y. C. Unruh. Initial data release from the INT Photometric H α Survey of the Northern Galactic Plane (IPHAS). *MNRAS*, 388(1):89–104, Jul 2008. doi: 10.1111/j.1365-2966.2008.13399.x.
- Andy D. Goulding, Jenny E. Greene, Rachel Bezanson, Johnny Greco, Sean Johnson, Alexie Leauthaud, Yoshiaki Matsuoka, Elinor Medezinski, and Adrian M. Price-Whelan. Galaxy interactions trigger rapid black hole growth: An unprecedented view from the Hyper Suprime-Cam survey. *PASJ*, 70:S37, January 2018. doi: 10.1093/pasj/psx135.
- Matthew J. Graham, S. G. Djorgovski, Ashish A. Mahabal, Ciro Donalek, and Andrew J. Drake. Machine-assisted discovery of relationships in astronomy. *MNRAS*, 431(3):2371–2384, May 2013. doi: 10.1093/mnras/stt329.
- Gregory M. Harry and LIGO Scientific Collaboration. Advanced LIGO: the next generation of gravitational wave detectors. *Classical and Quantum Gravity*, 27(8):084006, Apr 2010. doi: 10.1088/0264-9381/27/8/084006.
- P. Hartley, R. Flamary, N. Jackson, A. S. Tagore, and R. B. Metcalf. Support vector machine classification of strong gravitational lenses. *MNRAS*, 471:3378–3397, Nov 2017. doi: 10.1093/mnras/stx1733.
- A. Heck, F. Murtagh, and D. Ponz. The Increasing Importance of Statistical Methods in Astronomy. *The Messenger*, 41:22–25, September 1985.

- Christina Hedges, Simon Hodgkin, and Grant Kennedy. Discovery of new dipper stars with K2: a window into the inner disc region of T Tauri stars. *MNRAS*, 476:2968–2998, May 2018. doi: 10.1093/mnras/sty328.
- A. N. Heinze, J. L. Tonry, L. Denneau, H. Flewelling, B. Stalder, A. Rest, K. W. Smith, S. J. Smartt, and H. Weiland. A First Catalog of Variable Stars Measured by the Asteroid Terrestrial-impact Last Alert System (ATLAS). *AJ*, 156:241, November 2018. doi: 10.3847/1538-3881/aae47f.
- H. Hildebrandt, S. Arnouts, P. Capak, L. A. Moustakas, C. Wolf, F. B. Abdalla, R. J. Assef, M. Banerji, N. Benítez, G. B. Brammer, T. Budavári, S. Carliles, D. Coe, T. Dahlen, R. Feldmann, D. Gerdes, B. Gillis, O. Ilbert, R. Kotulla, O. Lahav, I. H. Li, J. M. Miralles, N. Purger, S. Schmidt, and J. Singal. PHAT: PHoto-z Accuracy Testing. *A&A*, 523:A31, Nov 2010. doi: 10.1051/0004-6361/201014885.
- I. Ting Ho. A Machine Learning Artificial Neural Network Calibration of the Strong-Line Oxygen Abundance. *MNRAS*, 485:3569–3579, Mar 2019. doi: 10.1093/mnras/stz649.
- Ho, T.K. Random decision forests. In *Proceedings of 3rd International Conference on Document Analysis and Recognition*, volume 1, pages 278–282, August 1995. doi: 10.1109/ICDAR.1995.598994.
- Alex Hocking, James E. Geach, Yi Sun, and Neil Davey. An automatic taxonomy of galaxy morphology using unsupervised machine learning. *MNRAS*, 473:1108–1129, Jan 2018. doi: 10.1093/mnras/stx2351.
- B. Hoyle. Measuring photometric redshifts using galaxy images and Deep Neural Networks. *A&C*, 16: 34–40, Jul 2016. doi: 10.1016/j.ascom.2016.03.006.
- B. Hoyle. Measuring photometric redshifts using galaxy images and Deep Neural Networks. *A&C*, 16: 34–40, July 2016. ISSN 2213-1337. doi: 10.1016/j.ascom.2016.03.006.
- Ben Hoyle, Markus Michael Rau, Kerstin Paech, Christopher Bonnett, Stella Seitz, and Jochen Weller. Anomaly detection for machine learning redshifts applied to SDSS galaxies. *MNRAS*, 452(4):4183–4194, Oct 2015. doi: 10.1093/mnras/stv1551.
- Ben Hoyle, Markus Michael Rau, Roman Zitlau, Stella Seitz, and Jochen Weller. Feature importance for machine learning redshifts applied to SDSS galaxies. *MNRAS*, 449(2):1275–1283, May 2015a. ISSN 0035-8711. doi: 10.1093/mnras/stv373.
- Ben Hoyle, Markus Michael Rau, Roman Zitlau, Stella Seitz, and Jochen Weller. Feature importance for machine learning redshifts applied to SDSS galaxies. *MNRAS*, 449(2):1275–1283, May 2015b. ISSN 0035-8711. doi: 10.1093/mnras/stv373.
- E. A. Huerta, C. J. Moore, Prayush Kumar, Daniel George, Alvin J. K. Chua, Roland Haas, Erik Wessel, Daniel Johnson, Derek Glennon, Adam Rebei, A. Miguel Holgado, Jonathan R. Gair, and Harald P. Pfeiffer. Eccentric, nonspinning, inspiral, Gaussian-process merger approximant for the detection and characterization of eccentric binary black hole mergers. *PhysRevD*, 97:024031, Jan 2018. doi: 10.1103/PhysRevD.97.024031.
- M. Huertas-Company, R. Gravet, G. Cabrera-Vives, P. G. Pérez-González, J. S. Kartaltepe, G. Barro, M. Bernardi, S. Mei, F. Shankar, P. Dimauro, E. F. Bell, D. Kocevski, D. C. Koo, S. M. Faber, and D. H. McIntosh. A CATALOG OF VISUAL-LIKE MORPHOLOGIES IN THE 5 CANDELS FIELDS USING DEEP LEARNING. *ApJS*, 221(1):8, October 2015. ISSN 0067-0049. doi: 10.1088/0067-0049/221/1/8.
- M. Huertas-Company, J. R. Primack, A. Dekel, D. C. Koo, S. Lapiner, D. Ceverino, R. C. Simons, G. F. Snyder, M. Bernardi, Z. Chen, H. Domínguez-Sánchez, C. T. Lee, B. Margalef-Bentabol, and D. Tuccillo. Deep Learning Identifies High-z Galaxies in a Central Blue Nugget Phase in a Characteristic Mass Range. *ApJ*, 858:114, May 2018. doi: 10.3847/1538-4357/aabfed.

- Jianan Hui, Miguel Aragon, Xinping Cui, and James M. Flegal. A machine learning approach to galaxy-LSS classification - I. Imprints on halo merger trees. *MNRAS*, 475:4494–4503, Apr 2018. doi: 10.1093/mnras/stx3235.
- Fadil Inceoglu, Jacob H. Jeppesen, Peter Kongstad, Néstor J. Hernández Marcano, Rune H. Jacobsen, and Christoffer Karoff. Using Machine Learning Methods to Forecast if Solar Flares Will Be Associated with CMEs and SEPs. *ApJ*, 861:128, Jul 2018. doi: 10.3847/1538-4357/aac81e.
- E. Iodice, M. Capaccioli, A. Grado, L. Limatola, M. Spavone, N. R. Napolitano, M. Paolillo, R. F. Peletier, M. Cantiello, T. Lisker, C. Wittmann, A. Venhola, M. Hilker, R. D’Abrusco, V. Pota, and P. Schipani. The Fornax Deep Survey with VST. I. The Extended and Diffuse Stellar Halo of NGC 1399 out to 192 kpc. *ApJ*, 820(1):42, Mar 2016. doi: 10.3847/0004-637X/820/1/42.
- Željko Ivezić and et al. LSST: From Science Drivers to Reference Design and Anticipated Data Products. *ApJ*, 873:111, Mar 2019. doi: 10.3847/1538-4357/ab042c.
- Željko Ivezić, Andrew J. Connolly, Jacob T. VanderPlas, and Alexander Gray. *Statistics, Data Mining, and Machine Learning in Astronomy*. 2014.
- C. Jacobs, K. Glazebrook, T. Collett, A. More, and C. McCarthy. Finding strong lenses in CFHTLS using convolutional neural networks. *MNRAS*, 471:167–181, Oct 2017. doi: 10.1093/mnras/stx1492.
- C. Jacobs, T. Collett, K. Glazebrook, E. Buckley-Geer, H. T. Diehl, H. Lin, C. McCarthy, A. K. Qin, C. Odden, M. Caso Escudero, P. Dial, V. J. Yung, S. Gaitsch, A. Pellico, K. A. Lindgren, T. M. C. Abbott, J. Annis, S. Avila, D. Brooks, D. L. Burke, A. Carnero Rosell, M. Carrasco Kind, J. Carretero, L. N. da Costa, J. De Vicente, P. Fosalba, J. Frieman, J. García-Bellido, E. Gaztanaga, D. A. Goldstein, D. Gruen, R. A. Gruendl, J. Gschwend, D. L. Hollowood, K. Honscheid, B. Hoyle, D. J. James, E. Krause, N. Kuropatkin, O. Lahav, M. Lima, M. A. G. Maia, J. L. Marshall, R. Miquel, A. A. Plazas, A. Roodman, E. Sanchez, V. Scarpine, S. Serrano, I. Sevilla-Noarbe, M. Smith, F. Sobreira, E. Suchyta, M. E. C. Swanson, G. Tarle, V. Vikram, A. R. Walker, and Y. Zhang and. An Extended Catalog of Galaxy–Galaxy Strong Gravitational Lenses Discovered in DES Using Convolutional Neural Networks. *ApJS*, 243(1): 17, July 2019a. ISSN 0067-0049. doi: 10.3847/1538-4365/ab26b6.
- C. Jacobs, T. Collett, K. Glazebrook, C. McCarthy, A. K. Qin, T. M. C. Abbott, F. B. Abdalla, J. Annis, S. Avila, K. Bechtol, E. Bertin, D. Brooks, E. Buckley-Geer, D. L. Burke, A. Carnero Rosell, M. Carrasco Kind, J. Carretero, L. N. da Costa, C. Davis, J. De Vicente, S. Desai, H. T. Diehl, P. Doel, T. F. Eifler, B. Flaugher, J. Frieman, J. García-Bellido, E. Gaztanaga, D. W. Gerdes, D. A. Goldstein, D. Gruen, R. A. Gruendl, J. Gschwend, G. Gutierrez, W. G. Hartley, D. L. Hollowood, K. Honscheid, B. Hoyle, D. J. James, K. Kuehn, N. Kuropatkin, O. Lahav, T. S. Li, M. Lima, H. Lin, M. a. G. Maia, P. Martini, C. J. Miller, R. Miquel, B. Nord, A. A. Plazas, E. Sanchez, V. Scarpine, M. Schubnell, S. Serrano, I. Sevilla-Noarbe, M. Smith, M. Soares-Santos, F. Sobreira, E. Suchyta, M. E. C. Swanson, G. Tarle, V. Vikram, A. R. Walker, Y. Zhang, and J. Zuntz. Finding high-redshift strong lenses in DES using convolutional neural networks. *MNRAS*, 484(4):5330–5349, April 2019b. ISSN 0035-8711. doi: 10.1093/mnras/stz272.
- C. Jaschek. Data Growth in Astronomy. *QJRAS*, 19:269, September 1978.
- Carlos Jaschek. Information problems in astrophysics. *PASP*, 80:654, dec 1968. doi: 10.1086/128707.
- S. Johnston, M. Bailes, N. Bartel, C. Baugh, M. Bietenholz, C. Blake, R. Braun, J. Brown, S. Chatterjee, J. Darling, A. Deller, R. Dodson, P. G. Edwards, R. Ekers, S. Ellingsen, I. Feain, B. M. Gaensler, M. Haverkorn, G. Hobbs, A. Hopkins, C. Jackson, C. James, G. Joncas, V. Kaspi, V. Kilborn, B. Koribalski, R. Kothes, T. L. Landecker, E. Lenc, J. Lovell, J. P. Macquart, R. Manchester, D. Matthews, N. M.

- McClure-Griffiths, R. Norris, U. L. Pen, C. Phillips, C. Power, R. Protheroe, E. Sadler, B. Schmidt, I. Stairs, L. Staveley-Smith, J. Stil, R. Taylor, S. Tingay, A. Tzioumis, M. Walker, J. Wall, and M. Wolleben. Science with the Australian Square Kilometre Array Pathfinder. *PASA*, 24:174–188, Dec 2007. doi: 10.1071/AS07033.
- S. Johnston, R. Taylor, M. Bailes, N. Bartel, C. Baugh, M. Bietenholz, C. Blake, R. Braun, J. Brown, S. Chatterjee, J. Darling, A. Deller, R. Dodson, P. Edwards, R. Ekers, S. Ellingsen, I. Feain, B. Gaensler, M. Haverkorn, G. Hobbs, A. Hopkins, C. Jackson, C. James, G. Joncas, V. Kaspi, V. Kilborn, B. Koribalski, R. Kothes, T. Landecker, E. Lenc, J. Lovell, J. P. Macquart, R. Manchester, D. Matthews, N. McClure-Griffiths, R. Norris, U. L. Pen, C. Phillips, C. Power, R. Protheroe, E. Sadler, B. Schmidt, I. Stairs, L. Staveley-Smith, J. Stil, S. Tingay, A. Tzioumis, M. Walker, J. Wall, and M. Wolleben. Science with ASKAP. The Australian square-kilometre-array pathfinder. *ExA*, 22:151–273, Dec 2008. doi: 10.1007/s10686-008-9124-7.
- E. Jones and J. Singal. Analysis of a custom support vector machine for photometric redshift estimation and the inclusion of galaxy shape information. *A&A*, 600:A113, Apr 2017. doi: 10.1051/0004-6361/201629558.
- Nicholas Kaiser. Pan-STARRS: a wide-field optical survey telescope array. In Jr. Oschmann, Jacobus M., editor, *Ground-based Telescopes*, volume 5489 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, pages 11–22, Oct 2004. doi: 10.1117/12.552472.
- Shi-Ju Kang, Jun-Hui Fan, Weiming Mao, Qingwen Wu, Jianchao Feng, and Yue Yin. Evaluating the Optical Classification of Fermi BCUs Using Machine Learning. *ApJ*, 872:189, Feb 2019. doi: 10.3847/1538-4357/ab0383.
- S. C. Keller, B. P. Schmidt, M. S. Bessell, P. G. Conroy, P. Francis, A. Granlund, E. Kowald, A. P. Oates, T. Martin-Jones, T. Preston, P. Tisserand, A. Vaccarella, and M. F. Waterson. The SkyMapper Telescope and The Southern Sky Survey. *PASA*, 24(1):1–12, May 2007. doi: 10.1071/AS07001.
- Edward J. Kim and Robert J. Brunner. Star-galaxy classification using deep convolutional neural networks. *MNRAS*, 464:4463–4475, Feb 2017. doi: 10.1093/mnras/stw2672.
- T. Kohonen. The self-organizing map. *Proceedings of the IEEE*, 78(9):1464–1480, September 1990. ISSN 0018-9219. doi: 10.1109/5.58325.
- Xiao Kong, A. Li Luo, Xiang-Ru Li, You-Fen Wang, Yin-Bi Li, and Jing-Kun Zhao. Spectral Feature Extraction for DB White Dwarfs Through Machine Learning Applied to New Discoveries in the Sdss DR12 and DR14. *PASP*, 130:084203, Aug 2018. doi: 10.1088/1538-3873/aac7a8.
- D. C. Koo. Optical multicolors : a poor person’s Z machine for galaxies. *AJ*, 90:418–440, Mar 1985. doi: 10.1086/113748.
- D. C. Koo. Overview - Photometric Redshifts: A Perspective from an Old-Timer[!] on their Past, Present, and Potential. In Ray Weymann, Lisa Storrie-Lombardi, Marcin Sawicki, and Robert Brunner, editors, *Photometric Redshifts and the Detection of High Redshift Galaxies*, volume 191 of *Astronomical Society of the Pacific Conference Series*, page 3, Jan 1999.
- Maria Krause, Elisa Pueschel, and Gernot Maier. Improved γ /hadron separation for the detection of faint γ -ray sources using boosted decision trees. *Astroparticle Physics*, 89:1–9, Mar 2017. doi: 10.1016/j.astropartphys.2017.01.004.

- J. R. Kriessler, E. H. Han, S. C. Odewahn, and T. C. Beers. Automated Morphological Classification of Galaxies and the Morphology-Density Relation. In *American Astronomical Society Meeting Abstracts*, volume 193, page 38.20, Dec 1998.
- Alex Krizhevsky, Ilya Sutskever, and Geoffrey E. Hinton. ImageNet Classification with Deep Convolutional Neural Networks. In F. Pereira, C. J. C. Burges, L. Bottou, and K. Q. Weinberger, editors, *Advances in Neural Information Processing Systems 25*, pages 1097–1105. Curran Associates, Inc., 2012.
- Victor F. Ksoll, Dimitrios A. Gouliermis, Ralf S. Klessen, Eva K. Grebel, Elena Sabbi, Jay Anderson, Daniel J. Lennon, Michele Cignoni, Guido de Marchi, Linda J. Smith, Monica Tosi, and Roeland P. van der Marel. Hubble Tarantula Treasury Project - VI. Identification of pre-main-sequence stars using machine-learning techniques. *MNRAS*, 479:2389–2414, Sep 2018. doi: 10.1093/mnras/sty1317.
- E. Kuminski and L. Shamir. A hybrid approach to machine learning annotation of large galaxy image databases. *A&C*, 25:257–269, Oct 2018. doi: 10.1016/j.ascom.2018.10.008.
- T. Kuntzer and F. Courbin. Detecting unresolved binary stars in Euclid VIS images. *A&A*, 606:A119, Oct 2017. doi: 10.1051/0004-6361/201730792.
- Michael J. Kurtz, Guenther Eichhorn, Alberto Accomazzi, Carolyn S. Grant, Stephen S. Murray, and Joyce M. Watson. The NASA Astrophysics Data System: Overview. *A&ASS*, 143:41–59, Apr 2000. doi: 10.1051/aas:2000170.
- O. Lahav, A. Nairn, L. Sodré, and M. C. Storrie-Lombardi. Neural computation as a tool for galaxy classification: Methods and examples. *MNRAS*, 283(1):207–221, October 1996. ISSN 0035-8711. doi: 10.1093/mnras/283.1.207.
- François Lanusse, Quanbin Ma, Nan Li, Thomas E. Collett, Chun-Liang Li, Siamak Ravanbakhsh, Rachel Mandelbaum, and Barnabás Póczos. CMU DeepLens: deep learning for automatic image-based galaxy-galaxy strong lens finding. *MNRAS*, 473:3895–3906, January 2018. doi: 10.1093/mnras/stx1665.
- R. Laureijs and et al. Euclid Definition Study Report. art. arXiv:1110.3193, Oct 2011.
- Y. LeCun, B. Boser, J. S. Denker, D. Henderson, R. E. Howard, W. Hubbard, and L. D. Jackel. Back-propagation Applied to Handwritten Zip Code Recognition. *Neural Comput.*, 1(4):541–551, December 1989. ISSN 0899-7667. doi: 10.1162/neco.1989.1.4.541.
- Boris Leistedt and David W. Hogg. Data-driven, Interpretable Photometric Redshifts Trained on Heterogeneous and Unrepresentative Data. *ApJ*, 838:5, Mar 2017. doi: 10.3847/1538-4357/aa6332.
- James R. Lemen, Alan M. Title, David J. Akin, Paul F. Boerner, Catherine Chou, Jerry F. Drake, Dexter W. Duncan, Christopher G. Edwards, Frank M. Friedlaender, Gary F. Heyman, Neal E. Hurlburt, Noah L. Katz, Gary D. Kushner, Michael Levay, Russell W. Lindgren, Dnyanesh P. Mathur, Edward L. McFeaters, Sarah Mitchell, Roger A. Rehse, Carolus J. Schrijver, Larry A. Springer, Robert A. Stern, Theodore D. Tarbell, Jean-Pierre Wuelser, C. Jacob Wolfson, Carl Yanari, Jay A. Bookbinder, Peter N. Cheimets, David Caldwell, Edward E. Deluca, Richard Gates, Leon Golub, Sang Park, William A. Podgorski, Rock I. Bush, Philip H. Scherrer, Mark A. Gummin, Peter Smith, Gary Auken, Paul Jerram, Peter Pool, Regina Soufli, David L. Windt, Sarah Beardsley, Matthew Clapp, James Lang, and Nicholas Waltham. The Atmospheric Imaging Assembly (AIA) on the Solar Dynamics Observatory (SDO). *Sol Phys*, 275:17–40, Jan 2012. doi: 10.1007/s11207-011-9776-8.
- Sébastien Lépine and Michael M. Shara. A Catalog of Northern Stars with Annual Proper Motions Larger than 0.15” (LSPM-NORTH Catalog). *AJ*, 129(3):1483–1522, Mar 2005. doi: 10.1086/427854.

- Yin-Bi Li, A. Li Luo, Chang-De Du, Fang Zuo, Meng-Xin Wang, Gang Zhao, Bi-Wei Jiang, Hua-Wei Zhang, Chao Liu, Li Qin, Rui Wang, Bing Du, Yan-Xin Guo, Bo Wang, Zhan-Wen Han, Mao-Sheng Xiang, Yang Huang, Bing-Qiu Chen, Jian-Jun Chen, Xiao Kong, Wen Hou, Yi-Han Song, You-Fen Wang, Ke-Fei Wu, Jian-Nan Zhang, Yong Zhang, Yue-Fei Wang, Zi-Huang Cao, Yong-Hui Hou, and Yong-Heng Zhao. Carbon Stars Identified from LAMOST DR4 Using Machine Learning. *ApJSS*, 234: 31, Feb 2018. doi: 10.3847/1538-4365/aaa415.
- Yun Li and Shihai Yang. Research on the fault diagnosis and self-healing technology of unattended Antarctic telescope. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, volume 10700, page 107004W, Jul 2018. doi: 10.1117/12.2311946.
- Hsing-Wen Lin, Ying-Tung Chen, Jen-Hung Wang, Shiang-Yu Wang, Fumi Yoshida, Wing-Huen Ip, Satoshi Miyazaki, and Tsuyoshi Terai. Machine-learning-based real-bogus system for the HSC-SSP moving object detection pipeline. *PASJ*, 70:S39, Jan 2018. doi: 10.1093/pasj/psx082.
- Chris J. Lintott, Kevin Schawinski, Anže Slosar, Kate Land, Steven Bamford, Daniel Thomas, M. Jordan Raddick, Robert C. Nichol, Alex Szalay, Dan Andreescu, Phil Murray, and Jan Vandenberg. Galaxy Zoo: morphologies derived from visual inspection of galaxies from the Sloan Digital Sky Survey. *MNRAS*, 389(3):1179–1189, Sep 2008. doi: 10.1111/j.1365-2966.2008.13689.x.
- Chang Liu, Na Deng, Jason T. L. Wang, and Haimin Wang. Predicting Solar Flares Using SDO/HMI Vector Magnetic Data Products and the Random Forest Algorithm. *ApJ*, 843:104, Jul 2017. doi: 10.3847/1538-4357/aa789b.
- F. T. Liu, K. M. Ting, and Z. Zhou. Isolation Forest. In *2008 Eighth IEEE International Conference on Data Mining*, pages 413–422, December 2008. doi: 10.1109/ICDM.2008.17.
- Jiajia Liu, Yudong Ye, Chenglong Shen, Yuming Wang, and Robert Erdélyi. A New Tool for CME Arrival Time Prediction using Machine Learning Algorithms: CAT-PUMA. *ApJ*, 855:109, Mar 2018. doi: 10.3847/1538-4357/aaae69.
- Junyu Liu. Artificial neural network in cosmic landscape. *Journal of High Energy Physics*, 2017:149, Dec 2017. doi: 10.1007/JHEP12(2017)149.
- LSST Science Collaboration and et al. LSST Science Book, Version 2.0. art. arXiv:0912.0201, Dec 2009.
- Luisa Lucie-Smith, Hiranya V. Peiris, Andrew Pontzen, and Michelle Lochner. Machine learning cosmological structure formation. *MNRAS*, 479:3405–3414, Sep 2018. doi: 10.1093/mnras/sty1719.
- V. Lukic, M. Brüggen, J. K. Banfield, O. I. Wong, L. Rudnick, R. P. Norris, and B. Simmons. Radio Galaxy Zoo: compact and extended radio source classification with deep learning. *MNRAS*, 476:246–260, May 2018. doi: 10.1093/mnras/sty163.
- Zhixian Ma, Haiguang Xu, Jie Zhu, Dan Hu, Weitian Li, Chenxi Shan, Zhenghao Zhu, Liyi Gu, Jinjin Li, Chengze Liu, and Xiangping Wu. A Machine Learning Based Morphological Classification of 14,245 Radio AGNs Selected from the Best-Heckman Sample. *ApJSS*, 240:34, Feb 2019. doi: 10.3847/1538-4365/aaf9a2.
- Zhixian Ma, Haiguang Xu, Jie Zhu, Dan Hu, Weitian Li, Chenxi Shan, Zhenghao Zhu, Liyi Gu, Jinjin Li, Chengze Liu, and Xiangping Wu. A Machine Learning Based Morphological Classification of 14,245 Radio AGNs Selected from the Best-Heckman Sample. *ApJS*, 240(2):34, February 2019. ISSN 0067-0049. doi: 10.3847/1538-4365/aaf9a2.

- A. A. Mahabal, S. G. Djorgovski, A. J. Drake, C. Donalek, M. J. Graham, R. D. Williams, Y. Chen, B. Moghaddam, M. Turmon, E. Beshore, and S. Larson. Discovery, classification, and scientific exploration of transient events from the Catalina Real-time Transient Survey. *Bulletin of the Astronomical Society of India*, 39(3):387–408, Sep 2011.
- Ashish Mahabal, Umaa Rebbapragada, Richard Walters, Frank J. Masci, Nadejda Blagorodnova, Jan van Roestel, Quan-Zhi Ye, Rahul Biswas, Kevin Burdge, Chan-Kao Chang, Dmitry A. Duev, V. Zach Golkhou, Adam A. Miller, Jakob Nordin, Charlotte Ward, Scott Adams, Eric C. Bellm, Doug Branton, Brian Bue, Chris Cannella, Andrew Connolly, Richard Dekany, Ulrich Feindt, Tiara Hung, Lucy Fortson, Sara Frederick, C. Fremling, Suvi Gezari, Matthew Graham, Steven Groom, Mansi M. Kasliwal, Shrinivas Kulkarni, Thomas Kupfer, Hsing Wen Lin, Chris Lintott, Ragnhild Lunnan, John Parejko, Thomas A. Prince, Reed Riddle, Ben Rusholme, Nicholas Saunders, Nima Sedaghat, David L. Shupe, Leo P. Singer, Maayane T. Soumagnac, Paula Szkody, Yutaro Tachibana, Kushal Tirumala, Sjoert van Velzen, and Darryl Wright. Machine Learning for the Zwicky Transient Facility. *PASP*, 131:038002, Mar 2019. doi: 10.1088/1538-3873/aaf3fa.
- Ashish A. Mahabal, S. G. Djorgovski, A. J. Drake, B. Hensley, C. Donalek, M. J. Graham, R. D. Williams, E. Glikman, C. Baltay, D. Rabinowitz, and PQ Survey Team. Towards the Automated Classification of Variable Objects and Transients. *Bulletin of the American Astronomical Society*, 42:427.06, Jan 2009.
- T. Marchetti, E. M. Rossi, G. Kordopatis, A. G. A. Brown, A. Rimoldi, E. Starkenburg, K. Youakim, and R. Ashley. An artificial neural network to discover hypervelocity stars: candidates in Gaia DR1/TGAS. *MNRAS*, 470:1388–1403, Sep 2017. doi: 10.1093/mnras/stx1304.
- Pablo Márquez-Neila, Chloe Fisher, Raphael Sznitman, and Kevin Heng. Supervised machine learning for analysing spectra of exoplanetary atmospheres. *Nature Astronomy*, 2:719–724, Jun 2018. doi: 10.1038/s41550-018-0504-2.
- D. Christopher Martin, James Fanson, David Schiminovich, Patrick Morrissey, Peter G. Friedman, Tom A. Barlow, Tim Conrow, Robert Grange, Patrick N. Jelinsky, Bruno Milliard, Oswald H. W. Siegmund, Luciana Bianchi, Yong-Ik Byun, Jose Donas, Karl Forster, Timothy M. Heckman, Young-Wook Lee, Barry F. Madore, Roger F. Malina, Susan G. Neff, R. Michael Rich, Todd Small, Frank Surber, Alex S. Szalay, Barry Welsh, and Ted K. Wyder. The Galaxy Evolution Explorer: A Space Ultraviolet Survey Mission. *ApJL*, 619(1):L1–L6, Jan 2005. doi: 10.1086/426387.
- Daniel Masters, Peter Capak, Daniel Stern, Olivier Ilbert, Mara Salvato, Samuel Schmidt, Giuseppe Longo, Jason Rhodes, Stephane Paltani, Bahram Mobasher, Henk Hoekstra, Hendrik Hildebrandt, Jean Coupon, Charles Steinhardt, Josh Speagle, Andreas Faisst, Adam Kalinich, Mark Brodwin, Massimo Brescia, and Stefano Cavuoti. MAPPING THE GALAXY COLOR–REDSHIFT RELATION: OPTIMAL PHOTOMETRIC REDSHIFT CALIBRATION STRATEGIES FOR COSMOLOGY SURVEYS. *ApJ*, 813(1):53, October 2015. ISSN 0004-637X. doi: 10.1088/0004-637X/813/1/53.
- M. McLeod, N. Libeskind, O. Lahav, and Y. Hoffman. Estimating the mass of the Local Group using machine learning applied to numerical simulations. *Journal of Cosmology and Astro-Particle Physics*, 2017:034, Dec 2017. doi: 10.1088/1475-7516/2017/12/034.
- R. B. Metcalf, M. Meneghetti, C. Avestruz, F. Bellagamba, C. R. Bom, E. Bertin, R. Cabanac, F. Courbin, A. Davies, E. Decencière, R. Flamary, R. Gavazzi, M. Geiger, P. Hartley, M. Huertas-Company, N. Jackson, C. Jacobs, E. Jullo, J.-P. Kneib, L. V. E. Koopmans, F. Lanusse, C.-L. Li, Q. Ma, M. Makler, N. Li, M. Lightman, C. E. Petrillo, S. Serjeant, C. Schäfer, A. Sonnenfeld, A. Tagore, C. Tortora, D. Tuccillo, M. B. Valentín, S. Velasco-Forero, G. A. Verdoes Kleijn, and G. Vernardos. The strong gravitational lens finding challenge. *A&A*, 625:A119, May 2019. ISSN 0004-6361, 1432-0746. doi: 10.1051/0004-6361/201832797.

- D. Michilli, J. W. T. Hessels, R. J. Lyon, C. M. Tan, C. Bassa, S. Cooper, V. I. Kondratiev, S. Sanidas, B. W. Stappers, and J. van Leeuwen. Single-pulse classifier for the LOFAR Tied-Array All-sky Survey. *MNRAS*, 480:3457–3467, Nov 2018. doi: 10.1093/mnras/sty2072.
- O. Miettinen. Protostellar classification using supervised machine learning algorithms. *ApES*, 363:197, Sep 2018. doi: 10.1007/s10509-018-3418-7.
- D. Mislis, S. Pyrzas, and K. A. Alsubai. TSARDI: a Machine Learning data rejection algorithm for transiting exoplanet light curves. *MNRAS*, 481:1624–1630, Dec 2018. doi: 10.1093/mnras/sty2361.
- Grégoire Montavon, Wojciech Samek, and Klaus-Robert Müller. Methods for interpreting and understanding deep neural networks. *Digital Signal Processing*, 73:1–15, February 2018. ISSN 1051-2004. doi: 10.1016/j.dsp.2017.10.011.
- Giuseppe Morello, P. W. Morris, S. D. Van Dyk, A. P. Marston, and J. C. Mauerhan. Applications of machine-learning algorithms for infrared colour selection of Galactic Wolf-Rayet stars. *MNRAS*, 473: 2565–2574, Jan 2018. doi: 10.1093/mnras/stx2474.
- V. Morello, E. D. Barr, S. Cooper, M. Bailes, S. Bates, N. D. R. Bhat, M. Burgay, S. Burke-Spolaor, A. D. Cameron, D. J. Champion, R. P. Eatough, C. M. L. Flynn, A. Jameson, S. Johnston, M. J. Keith, E. F. Keane, M. Kramer, L. Levin, C. Ng, E. Petroff, A. Possenti, B. W. Stappers, W. van Straten, and C. Tiburzi. The High Time Resolution Universe survey - XIV. Discovery of 23 pulsars through GPU-accelerated reprocessing. *MNRAS*, 483:3673–3685, Mar 2019. doi: 10.1093/mnras/sty3328.
- C. B. Morrison, H. Hildebrandt, S. J. Schmidt, I. K. Baldry, M. Bilicki, A. Choi, T. Erben, and P. Schneider. the-wizz: clustering redshift estimation for everyone. *MNRAS*, 467:3576–3589, May 2017. doi: 10.1093/mnras/stx342.
- S. K. Mukkavilli, D. Meger, and G. Dudek. EnviRoNet - Planetary Science Applications. In *AGU Fall Meeting Abstracts*, volume 2018, pages P43J–3875, Dec 2018.
- Nikhil Mukund, Saurabh Thakur, Sheelu Abraham, A. K. Aniyar, Sanjit Mitra, Ninan Sajeeth Philip, Kaustubh Vaghmare, and D. P. Acharjya. An Information Retrieval and Recommendation System for Astronomical Observatories. *ApJSS*, 235:22, Mar 2018. doi: 10.3847/1538-4365/aaadb2.
- Ancla Müller, Moritz Hackstein, Maksim Greiner, Philipp Frank, Dominik J. Bomans, Ralf-Jürgen Dettmar, and Torsten Enßlin. Sharpening up Galactic all-sky maps with complementary data. A machine learning approach. *A&A*, 620:A64, Nov 2018. doi: 10.1051/0004-6361/201833604.
- Ethan O. Nadler, Yao-Yuan Mao, Risa H. Wechsler, Shea Garrison-Kimmel, and Andrew Wetzel. Modeling the Impact of Baryons on Subhalo Populations with Machine Learning. *ApJ*, 859:129, June 2018. doi: 10.3847/1538-4357/aac266.
- S. Nakoneczny, M. Bilicki, A. Solarz, A. Pollo, N. Maddox, C. Spiniello, M. Brescia, and N. R. Napolitano. Catalog of quasars from the Kilo-Degree Survey Data Release 3. *A&A*, 624:A13, April 2019. ISSN 0004-6361, 1432-0746. doi: 10.1051/0004-6361/201834794.
- Gautham Narayan, Tayeb Zaidi, Monika D. Soraisam, Zhe Wang, Michelle Lochner, Thomas Matheson, Abhijit Saha, Shuo Yang, Zhenge Zhao, John Kececioglu, Carlos Scheidegger, Richard T. Snodgrass, Tim Axelrod, Tim Jenness, Robert S. Maier, Stephen T. Ridgway, Robert L. Seaman, Eric Michael Evans, Navdeep Singh, Clark Taylor, Jackson Toeniskoetter, Eric Welch, Songzhe Zhu, and ANTARES Collaboration. Machine-learning-based Brokers for Real-time Classification of the LSST Alert Stream. *ApJSS*, 236:9, May 2018. doi: 10.3847/1538-4365/aab781.

- Brett Naul, Joshua S. Bloom, Fernando Pérez, and Stéfan van der Walt. A recurrent neural network for classification of unevenly sampled variable stars. *Nature Astronomy*, 2:151–155, Nov 2018. doi: 10.1038/s41550-017-0321-z.
- T. Nguyen, V. Pankratius, L. Eckman, and S. Seager. Computer-aided discovery of debris disk candidates: A case study using the Wide-Field Infrared Survey Explorer (WISE) catalog. *A&C*, 23:72–82, Apr 2018. doi: 10.1016/j.ascom.2018.02.004.
- N. Nishizuka, K. Sugiura, Y. Kubo, M. Den, S. Watari, and M. Ishii. Solar Flare Prediction Model with Three Machine-learning Algorithms using Ultraviolet Brightening and Vector Magnetograms. *ApJ*, 835: 156, Feb 2017. doi: 10.3847/1538-4357/835/2/156.
- Ray P. Norris. Discovering the Unexpected in Astronomical Survey Data. *PASA*, 34:e007, Jan 2017. doi: 10.1017/pasa.2016.63.
- S. C. Odewahn, E. B. Stockwell, R. L. Pennington, R. M. Humphreys, and W. A. Zumach. Automated Star/Galaxy Discrimination With Neural Networks. *The Astronomical Journal*, 103:318, January 1992. ISSN 0004-6256. doi: 10.1086/116063.
- Fernanda Ostrovski, Richard G. McMahon, Andrew J. Connolly, Cameron A. Lemon, Matthew W. Auger, Manda Banerji, Johnathan M. Hung, Sergey E. Koposov, Christopher E. Lidman, Sophie L. Reed, Sahar Allam, Aurélien Benoit-Lévy, Emmanuel Bertin, David Brooks, Elizabeth Buckley-Geer, Aurelio Carnero Rosell, Matias Carrasco Kind, Jorge Carretero, Carlos E. Cunha, Luiz N. da Costa, Shantanu Desai, H. Thomas Diehl, Jörg P. Dietrich, August E. Evrard, David A. Finley, Brenna Flaugher, Pablo Fosalba, Josh Frieman, David W. Gerdes, Daniel A. Goldstein, Daniel Gruen, Robert A. Gruendl, Gaston Gutierrez, Klaus Honscheid, David J. James, Kyler Kuehn, Nikolay Kuropatkin, Marcos Lima, Huan Lin, Marcio A. G. Maia, Jennifer L. Marshall, Paul Martini, Peter Melchior, Ramon Miquel, Ricardo Ogando, Andrés Plazas Malagón, Kevin Reil, Kathy Romer, Eusebio Sanchez, Basilio Santiago, Vic Scarpine, Ignacio Sevilla-Noarbe, Marcelle Soares-Santos, Flavia Sobreira, Eric Suchyta, Gregory Tarle, Daniel Thomas, Douglas L. Tucker, and Alistair R. Walker. VDES J2325-5229 a $z = 2.7$ gravitationally lensed quasar discovered using morphology-independent supervised machine learning. *MNRAS*, 465: 4325–4334, Mar 2017. doi: 10.1093/mnras/stw2958.
- E. A. Owens, R. E. Griffiths, and K. U. Ratnatunga. Using oblique decision trees for the morphological classification of galaxies. *MNRAS*, 281(1):153–157, July 1996. ISSN 0035-8711. doi: 10.1093/mnras/281.1.153.
- Di Pang, Katerina Goseva-Popstojanova, Thomas Devine, and Maura McLaughlin. A novel single-pulse search approach to detection of dispersed radio pulses using clustering and supervised machine learning. *MNRAS*, 480:3302–3323, Nov 2018. doi: 10.1093/mnras/sty1992.
- Athanasios Papageorgiou, Márcio Catelan, Panagiota-Eleftheria Christopoulou, Andrew J. Drake, and S. G. Djorgovski. An Updated Catalog of 4680 Northern Eclipsing Binaries with Algol-type Light-curve Morphology in the Catalina Sky Surveys. *ApJSS*, 238:4, Sep 2018. doi: 10.3847/1538-4365/aad8a9.
- Ilya N. Pashchenko, Kirill V. Sokolovsky, and Panagiotis Gavras. Machine learning search for variable stars. *MNRAS*, 475:2326–2343, April 2018. doi: 10.1093/mnras/stx3222.
- J. Peña, C. Fuentes, F. Förster, J. C. Maureira, J. San Martín, J. Littín, P. Huijse, G. Cabrera-Vives, P. A. Estévez, L. Galbany, S. González-Gaitán, J. Martínez, Th. de Jaeger, and M. Hamuy. Asteroids in the High Cadence Transient Survey. *AJ*, 155:135, Mar 2018. doi: 10.3847/1538-3881/aaaaed.
- Kyle A. Pearson, Leon Palafox, and Caitlin A. Griffith. Searching for exoplanets using artificial intelligence. *MNRAS*, 474:478–491, February 2018. doi: 10.1093/mnras/stx2761.

- Tianrui Rae Peng, John Edward English, Pedro Silva, Darren R. Davis, and Wayne B. Hayes. SpArcFiRe: morphological selection effects due to reduced visibility of tightly winding arms in distant spiral galaxies. *MNRAS*, 479:5532–5543, Oct 2018. doi: 10.1093/mnras/sty546.
- Laurence Perreault Levasseur, Yashar D. Hezaveh, and Risa H. Wechsler. Uncertainties in Parameters Estimated with Neural Networks: Application to Strong Gravitational Lensing. *ApJ*, 850(1):L7, November 2017. doi: 10.3847/2041-8213/aa9704.
- W. Dean Pesnell, B. J. Thompson, and P. C. Chamberlin. The Solar Dynamics Observatory (SDO). *Sol Phys*, 275:3–15, Jan 2012. doi: 10.1007/s11207-011-9841-3.
- C. E. Petrillo, C. Tortora, S. Chatterjee, G. Vernardos, L. V. E. Koopmans, G. Verdoes Kleijn, N. R. Napolitano, G. Covone, P. Schneider, A. Grado, and J. McFarland. Finding strong gravitational lenses in the Kilo Degree Survey with Convolutional Neural Networks. *MNRAS*, 472(1):1129–1150, November 2017. ISSN 0035-8711. doi: 10.1093/mnras/stx2052.
- C. E. Petrillo, C. Tortora, G. Vernardos, L. V. E. Koopmans, G. Verdoes Kleijn, M. Bilicki, N. R. Napolitano, S. Chatterjee, G. Covone, A. Dvornik, T. Erben, F. Getman, B. Giblin, C. Heymans, J. T. A. de Jong, K. Kuijken, P. Schneider, H. Shan, C. Spiniello, and A. H. Wright. LinKS: Discovering galaxy-scale strong lenses in the Kilo-Degree Survey using Convolutional Neural Networks. *MNRAS*, 484:3879, January 2019.
- K. L. Polsterer, F. Gieseke, C. Igel, and T. Goto. Improving the Performance of Photometric Regression Models via Massive Parallel Feature Selection. *Astronomical Society of the Pacific Conference Series*, 485:425, 2014.
- Milad Pourrahmani, Hooshang Nayyeri, and Asantha Cooray. LensFlow: A Convolutional Neural Network in Search of Strong Gravitational Lenses. *ApJ*, 856:68, Mar 2018. doi: 10.3847/1538-4357/aaae6a.
- Jade Powell, Alejandro Torres-Forné, Ryan Lynch, Daniele Trifirò, Elena Cuoco, Marco Cavaglià, Ik Siong Heng, and José A. Font. Classification methods for noise transients in advanced gravitational-wave detectors II: performance tests on Advanced LIGO data. *Classical and Quantum Gravity*, 34:034002, Feb 2017. doi: 10.1088/1361-6382/34/3/034002.
- R. Qahwaji and T. Colak. Automatic Short-Term Solar Flare Prediction Using Machine Learning and Sunspot Associations. *Sol Phys*, 241(1):195–211, March 2007. ISSN 1573-093X. doi: 10.1007/s11207-006-0272-5. URL <https://doi.org/10.1007/s11207-006-0272-5>.
- J. R. Quinlan. Induction of decision trees. *Mach Learn*, 1(1):81–106, March 1986. ISSN 1573-0565. doi: 10.1007/BF00116251.
- Mika Rafieferantsoa, Sambatra Andrianomena, and Romeel Davé. Predicting the neutral hydrogen content of galaxies from optical data using machine learning. *MNRAS*, 479:4509–4525, Oct 2018. doi: 10.1093/mnras/sty1777.
- David M. Reiman and Brett E. Göhre. Deblending galaxy superpositions with branched generative adversarial networks. *MNRAS*, 485:2617–2627, Feb 2019. doi: 10.1093/mnras/stz575.
- Itamar Reis, Dovi Poznanski, Dalya Baron, Gail Zasowski, and Sahar Shahaf. Detecting outliers and learning complex structures with large spectroscopic surveys - a case study with APOGEE stars. *MNRAS*, 476:2117–2136, May 2018. doi: 10.1093/mnras/sty348.
- Itamar Reis, Dovi Poznanski, Dalya Baron, Gail Zasowski, and Sahar Shahaf. Detecting outliers and learning complex structures with large spectroscopic surveys – a case study with APOGEE stars. *MNRAS*, 476(2):2117–2136, May 2018. ISSN 0035-8711. doi: 10.1093/mnras/sty348.

- Itamar Reis, Dalya Baron, and Sahar Shahaf. Probabilistic Random Forest: A Machine Learning Algorithm for Noisy Data Sets. *AJ*, 157:16, Jan 2019. doi: 10.3847/1538-3881/aaf101.
- Andres C. Rodríguez, Tomasz Kacprzak, Aurelien Lucchi, Adam Amara, Raphaël Sgier, Janis Fluri, Thomas Hofmann, and Alexandre Réfrégier. Fast cosmic web simulations with generative adversarial networks. *Computational Astrophysics and Cosmology*, 5:4, Nov 2018. doi: 10.1186/s40668-018-0026-4.
- Frank Rosenblatt. The Perceptron-a perceiving and recognizing automaton. *Cornell Aeronautical Lab*, 1957.
- D. A. Rosenthal. Applying artificial intelligence to astronomical databases - a survey of applicable technology. *European Southern Observatory Conference and Workshop Proceedings*, 28:245, 1988.
- N. Roy, N. R. Napolitano, F. La Barbera, C. Tortora, F. Getman, M. Radovich, M. Capaccioli, M. Brescia, S. Cavuoti, G. Longo, M. A. Raj, E. Puddu, G. Covone, V. Amaro, C. Vellucci, A. Grado, K. Kuijken, G. Verdoes Kleijn, and E. Valentijn. Evolution of galaxy size-stellar mass relation from the Kilo-Degree Survey. *MNRAS*, 480:1057–1080, Oct 2018. doi: 10.1093/mnras/sty1917.
- A. Ruiz, A. Corral, G. Mountrichas, and I. Georgantopoulos. XMMPZCAT: A catalogue of photometric redshifts for X-ray sources. *A&A*, 618:A52, Oct 2018. doi: 10.1051/0004-6361/201833117.
- L. Sabin, Q. A. Parker, M. E. Contreras, L. Olguín, D. J. Frew, M. Stupar, R. Vázquez, N. J. Wright, R. L. M. Corradi, and R. A. H. Morris. New Galactic supernova remnants discovered with IPHAS. *MNRAS*, 431(1):279–291, May 2013. doi: 10.1093/mnras/stt160.
- I. Sadeh, F. B. Abdalla, and O. Lahav. ANNz2: Photometric Redshift and Probability Distribution Function Estimation using Machine Learning. *PASP*, 128(968):104502, Oct 2016. doi: 10.1088/1538-3873/128/968/104502.
- S. Saha, S. Basak, M. Safonova, K. Bora, S. Agrawal, P. Sarkar, and J. Murthy. Theoretical validation of potential habitability via analytical and boosted tree methods: An optimistic study on recently discovered exoplanets. *A&C*, 23:141–150, Apr 2018. doi: 10.1016/j.ascom.2018.03.003.
- Mara Salvato, Olivier Ilbert, and Ben Hoyle. The many flavours of photometric redshifts. *Nature Astronomy*, 3:212–222, Jun 2018. doi: 10.1038/s41550-018-0478-0.
- L. M. Sarro, J. Ordieres-Meré, A. Bello-García, A. González-Marcos, and E. Solano. Estimates of the atmospheric parameters of M-type stars: a machine-learning perspective. *MNRAS*, 476:1120–1139, May 2018. doi: 10.1093/mnras/sty165.
- Jan-Torge Schindler, Xiaohui Fan, Ian D. McGreer, Qian Yang, Jin Wu, Linhua Jiang, and Richard Green. The Extremely Luminous Quasar Survey in the SDSS Footprint. I. Infrared-based Candidate Selection. *ApJ*, 851:13, Dec 2017. doi: 10.3847/1538-4357/aa9929.
- Nima Sedaghat and Ashish Mahabal. Effective image differencing with convolutional neural networks for real-time transient hunting. *MNRAS*, 476:5365–5376, June 2018. doi: 10.1093/mnras/sty613.
- Ramprasaath R. Selvaraju, Michael Cogswell, Abhishek Das, Ramakrishna Vedantam, Devi Parikh, and Dhruv Batra. Grad-CAM: Visual Explanations From Deep Networks via Gradient-Based Localization. In *Proceedings of the IEEE International Conference on Computer Vision*, volume 2017-October, 2017.
- I. Sevilla-Noarbe and P. Etayo-Sotos. Effect of training characteristics on object classification: An application using Boosted Decision Trees. *A&C*, 11:64–72, June 2015. ISSN 2213-1337. doi: 10.1016/j.ascom.2015.03.010.

- Christopher J. Shallue and Andrew Vanderburg. Identifying Exoplanets with Deep Learning: A Five-planet Resonant Chain around Kepler-80 and an Eighth Planet around Kepler-90. *AJ*, 155:94, Feb 2018. doi: 10.3847/1538-3881/aa9e09.
- Harinder P. Singh, Ravi K. Gulati, and Ranjan Gupta. Stellar spectral classification using principal component analysis and artificial neural networks. *MNRAS*, 295(2):312–318, April 1998. ISSN 0035-8711. doi: 10.1046/j.1365-8711.1998.01255.x.
- Evgeny A. Smirnov and Alexey B. Markov. Identification of asteroids trapped inside three-body mean motion resonances: a machine-learning approach. *MNRAS*, 469:2024–2031, Aug 2017. doi: 10.1093/mnras/stx999.
- Joshua S. Speagle and Daniel J. Eisenstein. Deriving photometric redshifts using fuzzy archetypes and self-organizing maps - II. Implementation. *MNRAS*, 469:1205–1224, Jul 2017. doi: 10.1093/mnras/stx510.
- Kristoffer Stensbo-Smidt, Fabian Gieseke, Christian Igel, Andrew Zirm, and Kim Steenstrup Pedersen. Sacrificing information for the greater good: how to select photometric bands for optimal accuracy. *MNRAS*, 464(3):2577–2596, Jan 2017. doi: 10.1093/mnras/stw2476.
- M. C. Storrie-Lombardi, O. Lahav, L. Sodré, and L. J. Storrie-Lombardi. Morphological Classification of galaxies by Artificial Neural Networks. *MNRAS*, 259(1):8P–12P, November 1992. ISSN 0035-8711. doi: 10.1093/mnras/259.1.8P.
- Chris Stoughton and et al. Sloan Digital Sky Survey: Early Data Release. *AJ*, 123:485–548, Jan 2002. doi: 10.1086/324741.
- M. Süveges, F. Barblan, I. Lecoœur-Taïbi, A. Prša, B. Holl, L. Eyer, A. Kochoska, N. Mowlavi, and L. Rimoldini. Gaia eclipsing binary and multiple systems. Supervised classification and self-organizing maps. *A&A*, 603:A117, Jul 2017. doi: 10.1051/0004-6361/201629710.
- Alexander Szalay and Jim Gray. The World-Wide Telescope. *Science*, 293:2037–2040, Sep 2001. doi: 10.1126/science.293.5537.2037.
- Alexander Szalay and Jim Gray. 2020 computing: Science in an exponential world. *Nature*, 440(7083):413–414, 03 2006.
- Yutaro Tachibana and A. A. Miller. A Morphological Classification Model to Identify Unresolved PanSTARRS1 Sources: Application in the ZTF Real-time Pipeline. *PASP*, 130:128001, Dec 2018. doi: 10.1088/1538-3873/aae3d9.
- C. M. Tan, R. J. Lyon, B. W. Stappers, S. Cooper, J. W. T. Hessels, V. I. Kondratiev, D. Michilli, and S. Sanidas. Ensemble candidate classification for the LOTAAS pulsar survey. *MNRAS*, 474:4571–4583, Mar 2018. doi: 10.1093/mnras/stx3047.
- John D. Timlin, Nicholas P. Ross, Gordon T. Richards, Adam D. Myers, Andrew Pellegrino, Franz E. Bauer, Mark Lacy, Donald P. Schneider, Edward J. Wollack, and Nadia L. Zakamska. The Clustering of High-redshift ($2.9 \leq z \leq 5.1$) Quasars in SDSS Stripe 82. *ApJ*, 859:20, May 2018. doi: 10.3847/1538-4357/aab9ac.
- G. Ucci, A. Ferrara, S. Gallerani, and A. Pallottini. Inferring physical properties of galaxies from their emission-line spectra. *MNRAS*, 465:1144–1156, Feb 2017. doi: 10.1093/mnras/stw2836.
- G. Ucci, A. Ferrara, A. Pallottini, and S. Gallerani. GAME: GALaxy Machine learning for Emission lines. *MNRAS*, 477:1484–1494, Jun 2018. doi: 10.1093/mnras/sty804.

- R. Utama and J. Piekarewicz. Refining mass formulas for astrophysical applications: A Bayesian neural network approach. *Phys. Rev. C*, 96(4):044308, October 2017. doi: 10.1103/PhysRevC.96.044308.
- A. Vafaei Sadr, M. Farhang, S. M. S. Movahed, B. Bassett, and M. Kunz. Cosmic string detection with tree-based machine learning. *MNRAS*, 478:1132–1140, Jul 2018. doi: 10.1093/mnras/sty1055.
- A. Vafaei Sadr, Etienne E. Vos, Bruce A. Bassett, Zafirah Hosenie, N. Oozeer, and Michelle Lochner. DEEPSOURCE: point source detection using deep learning. *MNRAS*, 484:2793–2806, Apr 2019. doi: 10.1093/mnras/stz131.
- Lucas Valenzuela and Karim Pichara. Unsupervised classification of variable stars. *MNRAS*, 474:3259–3272, Mar 2018. doi: 10.1093/mnras/stx2913.
- Laurens van der Maaten and Geoffrey Hinton. Visualizing data using t-SNE. *Journal of machine learning research*, 9(Nov):2579–2605, 2008.
- M. P. van Haarlem and et al. LOFAR: The LOw-Frequency ARray. *A&A*, 556:A2, Aug 2013. doi: 10.1051/0004-6361/201220873.
- J. van Roestel, T. Kupfer, R. Ruiz-Carmona, P. J. Groot, T. A. Prince, K. Burdge, R. Laher, D. L. Shupe, and E. Bellm. Discovery of 36 eclipsing EL CVn binaries found by the Palomar Transient Factory. *MNRAS*, 475:2560–2590, Apr 2018. doi: 10.1093/mnras/stx3291.
- E. Vanzella, S. Cristiani, A. Fontana, M. Nonino, S. Arnouts, E. Giallongo, A. Grazian, G. Fasano, P. Popesso, P. Saracco, and S. Zaggia. Photometric redshifts with the Multilayer Perceptron Neural Network: Application to the HDF-S and SDSS. *A&A*, 423(2):761–776, August 2004. ISSN 0004-6361, 1432-0746. doi: 10.1051/0004-6361:20040176.
- I. B. Vavilova, A. A. Elyiv, and M. Yu. Vasylenko. Behind the Zone of Avoidance of the Milky Way: what can we Restore by Direct and Indirect Methods? *Russian Radio Physics and Radio Astronomy*, 23: 244–257, Dec 2018. doi: 10.15407/rpra23.04.244.
- Krisztián Vida and Rachael M. Roettenbacher. Finding flares in Kepler data using machine-learning tools. *A&A*, 616:A163, Sep 2018. doi: 10.1051/0004-6361/201833194.
- K. Viironen, A. Mampaso, R. L. M. Corradi, M. Rodríguez, R. Greimel, L. Sabin, S. E. Sale, Y. Unruh, G. Delgado-Inglada, J. E. Drew, C. Giammanco, P. Groot, Q. A. Parker, J. Sokoloski, and A. Zijlstra. New young planetary nebulae in IPHAS. *A&A*, 502(1):113–129, Jul 2009. doi: 10.1051/0004-6361/200811575.
- Pascal Vincent, Hugo Larochelle, Yoshua Bengio, and Pierre-Antoine Manzagol. Extracting and composing robust features with denoising autoencoders. In *Proceedings of the 25th International Conference on Machine Learning*, ICML ’08, pages 1096–1103, New York, NY, USA, 2008. ACM. ISBN 978-1-60558-205-4. doi: 10.1145/1390156.1390294.
- Jorick S. Vink, Janet E. Drew, Danny Steeghs, Nick J. Wright, Eduardo L. Martin, Boris T. Gänsicke, Robert Greimel, and Jeremy Drake. IPHAS discoveries of young stars towards Cyg OB2 and its southern periphery. *MNRAS*, 387(1):308–318, Jun 2008. doi: 10.1111/j.1365-2966.2008.13220.x.
- Yogesh Wadadekar. Estimating Photometric Redshifts Using Support Vector Machines. *PASP*, 117(827): 79, December 2004. ISSN 1538-3873. doi: 10.1086/427710.
- S. M. Walsh, H. Jerjen, and B. Wiallman. A Pair of Boötes: A New Milky Way Satellite. *ApJL*, 662(2): L83–L86, Jun 2007. doi: 10.1086/519684.

- Zhen Wan, Prajwal R. Kafle, Geraint F. Lewis, Dougal Mackey, Sanjib Sharma, and Rodrigo A. Ibata. Galactic cartography with SkyMapper - I. Population substructure and the stellar number density of the inner halo. *MNRAS*, 480:1218–1228, Oct 2018. doi: 10.1093/mnras/sty1880.
- Ke Wang, Ping Guo, and A. Li Luo. A new automated spectral feature extraction method and its application in spectral classification and defective spectra recovery. *MNRAS*, 465:4311–4324, Mar 2017. doi: 10.1093/mnras/stw2894.
- Michael J. Way, Jeffrey D. Scargle, Kamal M. Ali, and Ashok N. Srivastava. *Advances in Machine Learning and Data Mining for Astronomy*. 2012.
- Nicholas Weir, Usama M. Fayyad, S. G. Djorgovski, and Joseph Roden. The SKICAT System for Processing and Analyzing Digital Imaging Sky Surveys. *PASP*, 107:1243, December 1995. doi: 10.1086/133683. URL <https://ui.adsabs.harvard.edu/abs/1995PASP..107.1243W/abstract>.
- B. C. Whitmore. An objective classification system for spiral galaxies. I. The two dominant dimensions. *ApJ*, 278:61, March 1984. ISSN 0004-637X. doi: 10.1086/161768.
- Patrick Wils, Boris T. Gänsicke, Andrew J. Drake, and John Southworth. Data mining for dwarf novae in SDSS, GALEX and astrometric catalogues. *MNRAS*, 402(1):436–446, Feb 2010. doi: 10.1111/j.1365-2966.2009.15894.x.
- P. R. Woźniak, S. J. Williams, W. T. Vestrand, and V. Gupta. Identifying Red Variables in the Northern Sky Variability Survey. *AJ*, 128(6):2965–2976, Dec 2004. doi: 10.1086/425526.
- Xin Xin, Kaichang Di, Yexin Wang, Wenhui Wan, and Zongyu Yue. Automated detection of new impact sites on Martian surface from HiRISE images. *Advances in Space Research*, 60:1557–1569, Oct 2017. doi: 10.1016/j.asr.2017.06.044.
- Duo Xu and Stella S. R. Offner. Assessing the Performance of a Machine Learning Algorithm in Identifying Bubbles in Dust Emission. *ApJ*, 851:149, Dec 2017. doi: 10.3847/1538-4357/aa9a42.
- Qing-Zeng Yan, Ye Xu, A. J. Walsh, J. P. Macquart, G. C. MacLeod, Bo Zhang, P. J. Hancock, Xi Chen, and Zheng-Hong Tang. Improved selection criteria for H II regions, based on IRAS sources. *MNRAS*, 476:3981–3990, May 2018. doi: 10.1093/mnras/sty518.
- Tan Yang and Xiangru Li. An autoencoder of stellar spectra and its application in automatically estimating atmospheric parameters. *MNRAS*, 452(1):158–168, September 2015. ISSN 0035-8711. doi: 10.1093/mnras/stv1210.
- Yunfei Yang, Hongjuan Yang, Xianyong Bai, Huituan Zhou, Song Feng, and Bo Liang. Automatic Detection of Sunspots on Full-disk Solar Images using the Simulated Annealing Genetic Method. *PASP*, 130:104503, Oct 2018. doi: 10.1088/1538-3873/aadbfa.
- K. Yeakel, J. D. Vande-griff, D. G. Mitchell, D. C. Hamilton, C. M. Jackman, P. A. Delamere, and E. Rousos. Automatic Detection of Magnetospheric Regions around Saturn using Cassini Data. In *AGU Fall Meeting Abstracts*, volume 2018, pages P41D–3769, Dec 2018.
- C. W. Yip, A. J. Connolly, D. E. Vanden Berk, Z. Ma, J. A. Frieman, M. SubbaRao, A. S. Szalay, G. T. Richards, P. B. Hall, D. P. Schneider, A. M. Hopkins, J. Trump, and J. Brinkmann. Spectral Classification of Quasars in the Sloan Digital Sky Survey: Eigenspectra, Redshift, and Luminosity Effects. *AJ*, 128(6):2603, December 2004. ISSN 1538-3881. doi: 10.1086/425626.

- Suk Yee Yong, Anthea L. King, Rachel L. Webster, Nicholas F. Bate, Matthew J. O’Dowd, and Kathleen Labrie. Using the Properties of Broad Absorption Line Quasars to Illuminate Quasar Structure. *MNRAS*, 479:4153–4171, Sep 2018. doi: 10.1093/mnras/sty1540.
- Donald G. York and et al. The Sloan Digital Sky Survey: Technical Summary. *AJ*, 120:1579–1587, Sep 2000. doi: 10.1086/301513.
- M. Zevin, S. Coughlin, S. Bahaadini, E. Besler, N. Rohani, S. Allen, M. Cabero, K. Crowston, A. K. Katsaggelos, S. L. Larson, T. K. Lee, C. Lintott, T. B. Littenberg, A. Lundgren, C. Østerlund, J. R. Smith, L. Trouille, and V. Kalogera. Gravity Spy: integrating advanced LIGO detector characterization, machine learning, and citizen science. *Classical and Quantum Gravity*, 34:064003, Mar 2017. doi: 10.1088/1361-6382/aa5cea.
- Jingyi Zhang, Yanxia Zhang, and Yongheng Zhao. Imbalanced Learning for RR Lyrae Stars Based on SDSS and GALEX Databases. *AJ*, 155:108, Mar 2018. doi: 10.3847/1538-3881/aaa5b1.
- Yanxia Zhang and Yongheng Zhao. Astronomy in the Big Data Era. *Data Science Journal*, 14:11, May 2015. doi: 10.5334/dsj-2015-011.
- Hao Zhao, Wen-Xi Peng, Huan-Yu Wang, Rui Qiao, Dong-Ya Guo, Hong Xiao, and Zhao-Min Wang. A machine learning method to separate cosmic ray electrons from protons from 10 to 100 GeV using DAMPE data. *Research in Astronomy and Astrophysics*, 18:071, Jun 2018. doi: 10.1088/1674-4527/18/6/71.