FROM BRACES TO HECKE ALGEBRAS & QUANTUM GROUPS

ANASTASIA DOIKOU AND AGATA SMOKTUNOWICZ

ABSTRACT. We examine links between the theory of braces and set theoretical solutions of the Yang-Baxter equation, and fundamental concepts from the theory of quantum integrable systems. More precisely, we make connections with Hecke algebras and we identify new quantum groups associated to set-theoretic solutions coming from braces. We also construct a novel class of quantum discrete integrable systems and we derive symmetries for the corresponding periodic transfer matrices.

1. INTRODUCTION

The Yang-Baxter equation is a fundamental equation in the theory of quantum integrable models and solvable statistical systems, as well as in the formulation of quantum groups [18, 37, 15]. It was introduced in [53] as a main tool for the investigation of many particle systems with δ -type interactions, and in [4] for the study of a two-dimensional solvable statistical model. Since Drinfield [14] suggested a theory of set-theoretic solutions to the Yang-Baxter equation be developed, set-theoretic solutions have been extensively investigated using braided groups, and more recently by applying the theory of braces and skew-braces.

Set theoretical solutions and Yang-Baxter maps have been also extensively studied in the context of classical fully discrete integrable systems linked to the notion of Darboux-Bäcklund transformation within the Lax pair formulation [1, 52, 44]. In classical integrable systems usually a Poisson structure exists associated to a classical *r*-matrix, which is a solution of the classical Yang-Baxter equation [19]. Also, relevent recent results on Yang-Baxter maps, when the quantum group symmetry is an priori requirement can be found in [5].

It is worth noting that [28] provides one of the first instances of classification of set-theoretical solutions of Yang-Baxter equation. Various connections between the set theoretical Yang-Baxter equation and geometric crystals [17, 6], or soliton cellular automatons [50, 27] have been also demonstrated.

The theory of braces was established around 2005, when Wolfgang Rump developed a structure called a brace to describe all finite involutive set-theoretic

Date: December 17, 2019.

solutions of the Yang-Baxter equation. Rump showed that every brace yields a solution to the Yang-Baxter equation, and every non-degenerate, involutive settheoretic solution of the Yang-Baxter equation can be obtained from a brace, a structure which generalises nilpotent rings. Subsequently skew-braces were developed by Guareni and Vendramin to describe non-involutive solutions [26]. In this paper we will follow this direction.

The theory of braces and skew braces has connections with numerous research areas, for example with group theory (Garside groups, regular subgroups, factorised groups– see for example [3, 30, 31, 49]), algebraic number theory, Hopf-Galois extensions [2, 47], non-commutative ring theory [46, 40, 41], Knot theory [39, 42], Hopf algebras, quantum groups [16], universal algebra, groupoids [29], semi-braces [8], trusses [7] and Yang-Baxter maps. Moreover, skew braces are related to non-commutative physics, Yetter-Drinfield modules and Nichols algebras.

Because set-theoretic solutions coming from braces are involutive, it is possible to *Baxterise* them [38] and obtain solutions to the parameter dependent Yang-Baxter equation, which appear in quantum integrable systems.

The aim of this paper is to investigate connections between the theory of braces and selected topics from the theory of quantum integrable systems. More precisely:

- (1) We derive new quantum groups associated to braces.
- (2) We construct a novel class of quantum discrete integrable systems.
- (3) We identify symmetries of the periodic transfer matrices of the novel integrable systems.

Note that in [16] Etingof, Schedler and Soloviev constructed quantum groups associated to set-theoretic solutions, however we use a different construction coming from parameter dependent solutions of the Yang-Baxter equation and our quantum groups differ from these in [16].

Rump showed that every nilpotent ring is a brace, therefore readers who are not familiar with the theory of braces may replace use of the word brace in this paper with the words nilpotent ring. Readers interested in learning more about the theory of braces are referred to [45, 46, 9, 24, 48].

Structure of the paper. This paper is divided into four sections:

- (1) Section 1 contains the introduction of the paper.
- (2) Section 2 shows how to construct *R*-matrices associated to non-degenerate, involutive, set-theoretic solutions of the Yang-Baxter equation in preparation for Section 3.
- (3) Section 3 contains information on how to go about connecting the theory of quantum integrable systems with R-matrices constructed from braces as per Section 2. We construct the new quantum algebra associated to

braces. We also construct various realizations of the relevant quantum algebras using classical results from the theory of braces, along with posing some more open questions. Note that the Yangian is a special case within the larger class of quantum algebras emerging from braces.

The section consists of three subsections:

- 3.1: The Yang-Baxter equation & A-type Hecke algebra.
- 3.2: Quantum algebras from braces.
- 3.3: Representations of quantum algebras.

At the end of each section there are relevant questions and lines of enquiry for further research.

(4) Section 4, offers information on the construction of a new class of integrable quantum spin chain-like systems associated to braces. Spin chainlike systems are typically constructed by means of tensor realizations of the underlying quantum algebra, by introducing the so called transfer matrix. We also employ the theory of braces to construct symmetries of the corresponding periodic transfer matrices.

The section consists of two subsections:

- 4.1: Quantum spin chains from braces: novel class of integrable systems.
- 4.2: New symmetries of the periodic transfer matrix.

2. Basic information about braces & set-theoretic solutions

For a set-theoretic solution of the Braid equation, we will use notation (X, \check{r}) , instead of the usual notation (X, r), to be consistent with notations used in quantum integrable systems.

Let n be a natural number, and let $e_{i,j}$ denote the $n \times n$ matrix whose all entries are 0 except for the i, j-th entry, which is 1.

Let $X = \{x_1, \ldots, x_N\}$ be a set and $\check{r} : X \times X \to X \times X$. Denote

$$\check{r}(x,y) = (\sigma_x(y), \tau_y(x)) = (^x y, x^y).$$

We say that \check{r} is non-degenerate if σ_x and τ_y are bijective functions. Suppose that (X, \check{r}) is an involutive, non-degenerate set-theoretic solution of the Braid equation:

$$(\check{r} \times Id_X)(Id_X \times \check{r})(\check{r} \times Id_X) = (Id_X \times \check{r})(\check{r} \times Id_X)(Id_X \times \check{r}).$$

With a slight abuse of notation, let \check{r} also denote the *R*-matrix associated to the linearisation of \check{r} on $V = \mathbb{C}X$ (see [48] for more details).

This matrix is also called a *check-matrix*. Then the check-matrix related to (X, \check{r}) is $\check{r} = \check{r}_{i,j;k,l}$, where $\check{r}_{i,j;k,l} = 1$ if and only if $\check{r}(i,j) = (k,l)$, and is zero otherwise. Notice that the matrix $\check{r} : V \otimes V \to V \otimes V$ satisfies the (constant)

Braid equation:

$$(\check{r} \otimes Id_V)(Id_V \otimes \check{r})(\check{r} \otimes Id_V) = (Id_V \otimes \check{r})(\check{r} \times Id_V)(Id_V \otimes \check{r}).$$

Notice that $\check{r}^2 = I_{V \otimes V}$ the identity matrix, because \check{r} is involutive.

Let $r = \tau \check{r}$ be the corresponding solution of the quantum Yang-Baxter equation (where $\tau(x, y) = (y, x)$) and let r denote the matrix associated to the linearisation of r on $V = \mathbb{C}X$. Notice that the matrix r satisfies the constant Yang-Baxter equation.

Let $e_{x,y}$ be the matrix with x, y entry equal to 1 and all the other entries 0.

Lemma 2.1. Let notation be as above. Then the matrix \check{r} has the form:

(2.1)
$$\check{r} = \sum_{x,y \in X} e_{x,\sigma_x(y)} \otimes e_{y,\tau_y(x)}$$

The matrix r has the form:

$$r = \mathcal{P} \cdot \check{r} = (\sum_{x,y \in X} e_{y,x} \otimes e_{x,y}) (\sum_{x,y \in X} e_{x,\sigma_x(y)} \otimes e_{y,\tau_y(x)}),$$

consequently

(2.2)
$$r = \sum_{x,y \in X} e_{y,\sigma_x(y)} \otimes e_{x,\tau_y(x)}$$

Moreover because \check{r} is involutive we get $\check{r}(\sigma_x(y), \tau_y(x)) = (x, y)$, therefore

$$r = \sum_{x,y \in X} e_{\tau_y(x),x} \otimes e_{\sigma_x(y),y}$$

Proof. We use a direct calculation using the way in which the *R*-matrix associated to a set-theoretic solution (X, \check{r}) is built (for more details, see Definition 2.3 in [48]). It is worth noticing that, in some books on quantum groups, the obtained checkmatrix is also transposed. However, for involutive solutions, which we consider in this paper, the obtained R-matrix is symmetric, so it is the same.

In [45, 46] Rump showed that every solution (X, r) can be in a good way embedded in a brace.

Definition 2.2 (Proposition 4, [46]). A left brace is an abelian group (A; +) together with a multiplication \cdot such that the circle operation $a \circ b = a \cdot b + a + b$ makes A into a group, and $a \cdot (b + c) = a \cdot b + a \cdot c$.

In many papers, the following equivalent definition from [10] is used:

Definition 2.3 ([10]). A left brace is a set G together with binary operations + and \circ such that (G, +) is an abelian group, (G, \circ) is a group, and $a \circ (b + c) + a = a \circ b + a \circ c$ for all $a, b, c \in G$.

The additive identity of a brace A will be denoted by 0 and the multiplicative identity by 1. In every brace 0 = 1. The same notation will be used for skew braces (in every skew brace 0 = 1).

Some authors use the notation \cdot instead of \circ and * instead of \cdot (see for example [10, 24, 23]).

In [16], Etingof, Schedler and Soloviev introduced the retract relation for any solution (X, r). Denote $X = \{x_1, \ldots, x_N\}$ and $r(x, y) = (\sigma_x(y), \tau_y(x))$. Recall that the retract relation \sim on X is defined by $x_i \sim x_j$ if $\sigma_i = \sigma_j$. The induced solution $Ret(X, r) = (X/\sim, r^{\sim})$ is called the *retraction* of X. A solution (X, r) is called a *multi-permutation solution of level* m if m is the smallest non-negative integer such that after m retractions we obtain the solution with one element.

Throughout this paper we will use the following result, which is implicit in [45, 46] and explicit in Theorem 4.4 of [10].

Theorem 2.4. (Rump's theorem, [45, 46, 10]). It is known that for an involutive, non degenerate solution of the braid equation there is always an underlying brace $(B, \circ, +)$, such that the maps σ_x and τ_y come from this brace, and X is a subset in this brace such that $\check{r}(X, X) \subseteq (X, X)$ and $\check{r}(x, y) = (\sigma_x(y), \tau_y(x))$, where $\sigma_x(y) = x \circ y - x, \tau_y(x) = t \circ x - t$, where t is the inverse of $\sigma_x(y)$ in the circle group (B, \circ) . Moreover, we can assume that every element from B belongs to the additive group (X, +) generated by elements of X. In addition every solution of this type is a non-degenerate, involutive set-theoretic solution of the braid equation.

We will call the brace B an underlying brace of the solution (X, \check{r}) , or a brace associated to the solution (X, \check{r}) . We will also say that the solution (X, \check{r}) is associated to brace B. Notice that this is also related to the formula of settheoretic solutions associated to the braided group (see [16] and [24]).

The following remark was also discovered by Rump.

Remark 2.5. Let $(N, +, \cdot)$ be an associative ring which is a nilpotent ring. For $a, b \in N$ define

$$a \circ b = a \cdot b + a + b,$$

then $(N, +, \circ)$ is a brace.

Definition 2.6. Let X, Y be sets and $\check{r} : X \times X \to X \times X$, $\check{r}' : Y \times Y \to Y \times Y$ be functions. Let (X,\check{r}) and (Y,\check{r}') be set-theoretic solutions of the Braid equation, and left $f : X \to Y$ be a function onto X such that $\check{r}'(f(x), f(y)) = (f \times f)(\check{r}(x, y))$, for all $x, y \in X$. Then f is called a homomorphism of solutions. If f is one -to - one then f is called an isomorphism of solutions.

Lemma 2.7. Notice that if a solution (X, r) comes from brace B, and J is an ideal in B and $X_J = \{x + J : j \in J\}$ is a subset of the factor brace B/J then the

map $f: X \to X_J$ given by f(x) = x + J is a homomorphism of solutions (X, \check{r}) and (X_J, \check{r}_J) , where (X_J, \check{r}_J) is the solution associated to brace B/J on the set $X_J = X + J$.

Proof. It follows immediately from the properties of an ideal in a brace (ideals in braces were defined in [46]. See also [10]). \Box

3. Hecke Algebras & Quantum groups from braces

3.1. The Yang-Baxter equation & Hecke algebras. In this section we explore various connections between braces, representations of the A-type Hecke algebras, and quantum algebras. In particular, after showing some fundamental properties for the brace R-matrices and making the direct connection with A-type Hecke algebras, we derive new quantum algebras coming from braces. The Yangian $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$ turns out to be a special case within this larger class of quantum algebras.

Before we start our investigation on the aforementioned connections let us first derive some preliminary results, that will be essential especially when proving the integrability of open spin-chain like systems, this issue however will be discussed in a forthcoming work [13]. Recall the Yang-Baxter equation in the braid form $(\delta = \lambda_1 - \lambda_2)$:

(3.1)
$$\mathring{R}_{12}(\delta) \mathring{R}_{23}(\lambda_1) \mathring{R}_{12}(\lambda_2) = \mathring{R}_{23}(\lambda_2) \mathring{R}_{12}(\lambda_1) \mathring{R}_{23}(\delta)$$

We focus here on brace type solutions of (3.1), i.e.

$$\check{R}(\lambda) = \lambda \check{r} + \mathbb{I}$$

where $\mathbb{I} = I_X \otimes I_X$ and I_X is the identity matrix of dimension equal to the cardinality of the set X. Also, we recall the notation introduced in Lemma 2.1 for the matrix \check{r} (2.1). Let also, $R = \mathcal{P}\check{R}$, then

$$(3.3) R(\lambda) = \lambda r + \mathcal{P}$$

where r is defined in (2.2), and R is a solution of the Yang-Baxter equation in the familiar form:

(3.4)
$$R_{12}(\delta) R_{13}(\lambda_1) R_{23}(\lambda_2) = R_{23}(\lambda_2) R_{13}(\lambda_1) R_{12}(\delta)$$

Remark 3.1. It would be useful for the following Proposition to introduce the notion of partial transposition. Let $A \in End(\mathbb{C}^{N} \otimes \mathbb{C}^{N})$ expressed as: $A = \sum_{i,j=1}^{N} A_{ij,kl} e_{i,j} \otimes e_{k,l}$. We define the partial transposition as follows (in the index notation):

(3.5)
$$A_{12}^{t_1} = \sum_{i,j=1}^{\mathcal{N}} A_{ij,kl} \ e_{i,j}^t \otimes e_{k,l}, \quad A_{12}^{t_2} = \sum_{i,j=1}^{\mathcal{N}} A_{ij,kl} \ e_{i,j} \otimes e_{k,l}^t$$

where $e_{i,j}^t = e_{j,i}$.

Proposition 3.2. The brace *R*-matrix satisfies the following fundamental properties:

(3.6)
$$R_{12}(\lambda) R_{21}(-\lambda) = (-\lambda^2 + 1)\mathbb{I}, \ Unitarity$$

(3.7)
$$R_{12}^{t_1}(\lambda) \ R_{12}^{t_2}(-\lambda - \mathcal{N}) = \lambda(-\lambda - \mathcal{N})\mathbb{I}, \ Crossing-unitarity$$

(3.8) $R_{12}^{t_1 t_2}(\lambda) = R_{21}(\lambda),$

where $t_{1,2}$ denotes transposition on the fist, second space respectively.

Proof. Recall $R_{21} = \mathcal{P}R_{12}\mathcal{P}$, the proof of unitarity is straightforward due to $\check{r}^2 = \mathcal{P}^2 = \mathbb{I}$. To prove crossing-unitarity (3.7) it suffices to show the following identities:

(3.9)
$$(\mathcal{P}_{12}^{t_1})^2 = \mathcal{N}\mathcal{P}_{12}^{t_1}, \quad r_{12}^{t_1}\mathcal{P}_{12}^{t_1} = \mathcal{P}_{12}^{t_1}r_{12}^{t_2} = \mathcal{P}_{12}^{t_1}, \quad r_{12}^{t_1}r_{12}^{t_2} = \mathbb{I}.$$

The above can be easily shown, from the definitions of $\mathcal{P} = \sum_{x,y} e_{x,y} \otimes e_{y,x}$ and $r = \mathcal{P}\check{r}$ (2.2). Given (3.9) the crossing-unitarity immediately follows. The last property (3.8) immediately follows from the definitions of R_{12} , R_{21} and the brace representation.

We can now state the obvious connection of the brace representation with the *A*-type Hecke algebra.

Definition 3.3. The A-type Hecke algebra $\mathcal{H}_N(q)$ is defined by the generators g_l , $l \in \{1, 2, ..., N-1\}$ and the exchange relations:

$$(3.10) g_l g_{l+1} g_l = g_{l+1} g_l g_{l+1},$$

(3.11)
$$\left[g_l, g_m\right] = 0, \ |l-m| > 1$$

(3.12)
$$(g_l - q)(g_l + q^{-1}) = 0.$$

Remark 3.4. The brace solution \check{r} (2.1) is a representation of the A-type Hecke algebra for q = 1.

Indeed, \check{r} satisfies the braid relation and $\check{r}^2 = \mathbb{I}$, which can be easily shown by using the involution. We can then define $g_1 = \check{r} \otimes I^{\otimes N}$ and $g_i = I^{\otimes i-1} \otimes \check{r} \otimes I^{\otimes N-i-1}$. Let us also show below the braid relation:

$$(3.13) \qquad (\check{r} \otimes I_X) \ (I_X \otimes \check{r}) \ (\check{r} \otimes I_X) = (I_X \otimes \check{r}) \ (\check{r} \otimes I_X) \ (\check{r} \otimes I_X).$$

Starting from the LHS of the equation above we conclude:

(3.14)
$$\sum e_{x,\sigma_{\bar{x}}(\bar{y})} \otimes e_{y,\tau_{\bar{y}}(\bar{x})} \otimes e_{\hat{y},\tau_{\bar{y}}(\hat{x})}$$

provided that $\hat{x} = \tau_y(x)$, $\bar{x} = \sigma_x(y)$, $\bar{y} = \sigma_{\hat{x}}(\hat{y})$. Similarly the RHS leads to

(3.15)
$$\sum e_{x,\sigma_x(y)} \otimes e_{\hat{x},\sigma_{\bar{x}}(\bar{y})} \otimes e_{\hat{y},\tau_{\bar{y}}(\bar{x})}.$$

provided that $y = \sigma_{\hat{x}}(\hat{y}), \ \bar{x} = \tau_y(x), \ \bar{y} = \tau_{\hat{y}}(\hat{x}).$

Comparison between the LHS and RHS expressions corresponds to the set-theoretic version and follows from the second part of Rump's theorem (Theorem 2.4).

3.2. Quantum algebras from braces. Given a solution of the Yang-Baxter equation, the quantum algebra is defined via the fundamental relation [18] (we have multiplied the familiar RTT relation by the permutation operator):

(3.16)
$$\check{R}_{12}(\lambda_1 - \lambda_2) L_1(\lambda_1) L_2(\lambda_2) = L_1(\lambda_2) L_2(\lambda_1) \check{R}_{12}(\lambda_1 - \lambda_2).$$

 $R(\lambda) \in \operatorname{End}(\mathbb{C}^{\mathcal{N}} \otimes \mathbb{C}^{\mathcal{N}}), \ L(\lambda) \in \operatorname{End}(\mathbb{C}^{\mathcal{N}}) \otimes \mathfrak{A},$ where \mathfrak{A} is the quantum algebra defined by (3.16).

Proposition 3.5. The quantum algebra associated to the brace solution

$$\check{r} = \sum_{x,y \in X} e_{x,\sigma_x(y)} \otimes e_{y,\tau_y(x)}$$

is defined by generators $L_{zw}^{(m)}, \ z, w \in X$, and defining relations

$$(3.17) L_{zw}^{(n)} L_{\hat{z}\hat{w}}^{(m)} - L_{zw}^{(m)} L_{\hat{z}\hat{w}}^{(n)} = L_{z\sigma_w(\hat{w})}^{(m)} L_{\hat{z}\tau_{\hat{w}}(w)}^{(n+1)} - L_{z\sigma_w(\hat{w})}^{(m+1)} L_{\hat{z}\tau_{\hat{w}}(w)}^{(n)} - L_{\sigma_z(\hat{z})w}^{(n+1)} L_{\tau_{\hat{z}}(z)\hat{w}}^{(m)} + L_{\sigma_z(\hat{z})w}^{(n)} L_{\tau_{\hat{z}}(z)\hat{w}}^{(m+1)}.$$

Proof. Using the index notation we define:

(3.18)
$$L_1(\lambda) = \sum_{z,w \in X} e_{z,w} \otimes I_X \otimes L_{z,w}(\lambda)$$

(3.19)
$$L_2(\lambda) = \sum_{z,w \in X} I_X \otimes e_{z,w} \otimes L_{z,w}(\lambda)$$

(3.20)
$$\check{R}_{12} = \check{R} \otimes \mathrm{id}_{\mathfrak{A}}$$

where recall I_X is the identity matrix of dimension equal to the cardinality of the set X, and $id_{\mathfrak{A}}$ is the identity element of the algebra \mathfrak{A} . $L_{z,w}(\lambda)$ are elements of the affine algebra \mathfrak{A} and \check{R} is given in (3.2), (2.1). The exchange relations among the various generators of the affine algebra are derived below via (3.16).

Let us express L as a formal power series expansion $L(\lambda) = \sum_{n=0}^{\infty} \frac{L^{(n)}}{\lambda^n}$. Substituting expressions (3.2), and the λ^{-1} expansion in (3.16) we obtain the defining relations of the quantum algebra associated to a brace *R*-matrix (we focus on terms $\lambda_1^{-n} \lambda_2^{-m}$):

(3.21)
$$\check{r}_{12}L_1^{(n+1)}L_2^{(m)} - \check{r}_{12}L_1^{(n)}L_2^{(m+1)} + L_1^{(n)}L_2^{(m)} = L_1^{(m)}L_2^{(n+1)}\check{r}_{12} - L_1^{(m+1)}L_2^{(n)}\check{r}_{12} + L_1^{(m)}L_2^{(n)}.$$

This immediately leads to the quantum algebra relations (3.17), after recalling similarly to (3.18)-(3.20):

$$L_{1}^{(k)} = \sum_{i,j \in X} e_{i,j} \otimes I_{X} \otimes L_{i,j}^{(k)}, \quad L_{2}^{(k)} = \sum_{i,j \in X} I_{X} \otimes e_{i,j} \otimes L_{i,j}^{(k)},$$

and $\check{r}_{12} = \check{r} \otimes \mathrm{id}_{\mathfrak{A}}$. $L_{i,j}^{(k)}$ are the generators of the associated quantum algebra.

By substituting the above expressions for $L_1^{(k)}$ and $L_2^{(k)}$ and $\check{r} = \sum_{i,j \in X} e_{i,i,j} \otimes e_{j,i,j}$ in (3.21) and computing both sides we obtain:

$$\sum_{x,j,y,i\in X} e_{x,j} \otimes e_{y,i} \otimes Q_{x,j,y,i}^{(m,n)} = \sum_{x,j,y,i\in X} e_{x,j} \otimes e_{y,i} \otimes P_{x,j,y,i}^{(m,n)}$$

where

$$Q_{x,j,y,i}^{(m,n)} = L_{xy,j}^{(n+1)} L_{x^y,i}^{(m)} - L_{xy,j}^{(n)} L_{x^{y},i}^{(m+1)} + L_{x,j}^{(n)} L_{y,i}^{(m)}$$

$$P_{x,j,y,i}^{(m,n)} = L_{x,ji}^{(m)} L_{y,ji}^{(n+1)} - L_{x,ii}^{(m+1)} L_{y,ji}^{(n)} + L_{x,j}^{(m)} L_{y,ii}^{(n)}$$

Notice that $Q_{x,j,y,l}^{(m,n)} - P_{x,j,y,l}^{(m,n)}$ are the defining relations in our quantum algebra.

Definition 3.6. Let (X, \check{r}) be a set-theoretic solution of the Yang-Baxter equation, with $\check{r}(x,y) = ({}^{x}y, {}^{x}y)$. Recall that $P = \sum_{i,j \in X} e_{i,j} \otimes e_{j,i}$. The quantum algebra associated to the brace R matrix $R(u) = P + uP\check{r}$ is defined by generators $L_{zw}^{(m)}$, $z, w \in X$, m = 0, 1, 2, ... and defining relations

$$\begin{split} L^{(n+1)}_{xy,j}L^{(m)}_{x^{y},i} - L^{(n)}_{xy,j}L^{(m+1)}_{x^{y},i} + L^{(n)}_{x,j}L^{(m)}_{y,i} = \\ L^{(m)}_{x,ji}L^{(n+1)}_{y,j^{i}} - L^{(m+1)}_{x,ji}L^{(n)}_{y,j^{i}} + L^{(m)}_{x,j}L^{(n)}_{y,i} \end{split}$$

for $x, j, y, i \in X$. This algebra will be denoted as $\mathfrak{A}(X, \check{r})$

This is the same algebra as in Proposition 3.5.

The quantum algebra is a Hopf algebra also equipped with a co-product [18, 15]

$$(3.22) (I \otimes \Delta)L(\lambda) = L_{13}(\lambda)L_{12}(\lambda)$$

as well as an anti-pode $(I \otimes s)L(\lambda) = L^{-1}(\lambda)$ and a co-unit $(I \otimes \epsilon)L(\lambda) = \mathbb{I}$.

Special case: The Yangian $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$. In this part we recall some basic notions about $\mathfrak{gl}_{\mathcal{N}}$ and the Yangian $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$. $\mathfrak{gl}_{\mathcal{N}}$ is the algebra (over field \mathbb{C}) with generators denoted as $\mathfrak{L}_{i,j}$ for $i, j \in \{1, \ldots, \mathcal{N}\}$, and satisfying the following defining relations:

(3.23)
$$\left[\mathfrak{L}_{i,j}, \ \mathfrak{L}_{k,l}\right] = \mathfrak{L}_{k,j}\delta_{i,l} - \mathfrak{L}_{i,l}\delta_{k,j},$$

where $[a, b] = a \cdot b - b \cdot a$ as usual.

Let A be an associative algebra over the field \mathbb{C} . The matrices $e_{i,j} \in M_N(A)$ are the generators of \mathfrak{gl}_N in the fundamental representation. For $y \in \mathfrak{gl}_{\mathcal{N}}$ we define for $Y \in \mathfrak{gl}_{\mathcal{N}}$ (we also recall the usual notation $\Delta^{(2)}(Y) = \Delta(Y)$)

(3.24)
$$\Delta^{(N)}(\mathbf{Y}) = \sum_{n=1}^{N} Y_n$$
$$= \sum_{n=1}^{N} \mathrm{id} \otimes \ldots \otimes \underbrace{\mathbf{Y}}_{n^{th}} \underbrace{\mathbf{Y}}_{\text{position}} \otimes \ldots \otimes \mathrm{id},$$

the element Y_n , in the standard *index notation* above, appears in the n^{th} position of the N co-product. The elements $\Delta^{(N)}(e_{i,j})$ are tensor representations of \mathfrak{gl}_N .

We recall two well known results about Yangians. For more information on Yangians we refer the interested reader to [15, 43]. The Yangian for $\mathfrak{gl}_{\mathcal{N}}$ will be denoted as $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$. We recall the defining relations of $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$ in the following Corollary 3.7.

Corollary 3.7. In the special case $\check{r} = \mathcal{P}$ the $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$ algebra is recovered as the corresponding quantum algebra.

Proof. The quantum algebra is constructed using the above fundamental relation. Let us consider the special case where $\check{r} = \mathcal{P}$, which corresponds to the Yangian $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$. Let us express L as a formal power series expansion $L(\lambda) = \sum_{n=0}^{\infty} \frac{L^{(n)}}{\lambda^n}$. Then the fundamental relation leads to:

(3.25)
$$\left[L_1^{(n+1)}, L_2^{(m)}\right] - \left[L_1^{(n)}, L_2^{(m+1)}\right] = \mathcal{P}_{12}\left(L_1^{(m)}L_2^{(n)} - L_1^{(n)}L_2^{(m)}\right).$$

Recalling the definitions (3.20) and $\mathcal{P}_{12} = \sum e_{ij} \otimes e_{ji} \otimes id$, we conclude (see also [15])

(3.26)
$$\begin{bmatrix} L_{i,j}^{(n+1)}, \ L_{k,l}^{(m)} \end{bmatrix} - \begin{bmatrix} L_{i,j}^{(n)}, \ L_{k,l}^{(m+1)} \end{bmatrix} = L_{k,j}^{(m)} L_{i,l}^{(n)} - L_{k,j}^{(n)} L_{i,l}^{(m)}$$

The latter relations are, up to a minus sign, the defining relations of the $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$ algebra.

Corollary 3.8. In the case of the Yangian the finitely dimensional subalgebra $\mathfrak{gl}_{\mathcal{N}}$ emerges from (3.25), realized by the elements of $L^{(1)}$.

Proof. Let us first consider terms proportional to $\lambda_1 \lambda_2^{-m}$ and λ_2^{-m} from the relations of the quantum algebra (3.21) for any \hat{r} :

(3.27)
$$\check{r}_{12}L_1^{(0)}L_2^{(m)} = L_1^{(m)}L_2^{(0)}\check{r}_{12},$$

(3.28)
$$\check{r}_{12}L_1^{(1)}L_2^{(m)} - L_1^{(m)}L_2^{(1)}\check{r}_{12} = L_1^{(m)}L_2^{(0)} - L_1^{(0)}L_2^{(m)}.$$

When considering the Yangian ($\check{r} = \mathcal{P}$), and focusing on the case where m = 1, and also consider $L^{(0)} = I_X \otimes id$ (in this case (3.27) is automatically satisfied, since $\check{r} = \mathcal{P}$), we deduce from (3.28) that the elements $L_{i,j}^{(1)}$ satisfy the defining relations of $\mathfrak{gl}_{\mathcal{N}}$ (3.7). Note that our choise here $L^{(0)} = I_X \otimes \operatorname{id}$ is compatible with the fact that $L(\lambda) = \lambda I_X \otimes \operatorname{id} + \mathfrak{P}$, where $\mathfrak{P} = \sum_{i,j=1}^{\mathcal{N}} e_{i,j} \otimes \mathfrak{P}_{i,j}$ and $\mathfrak{P}_{i,j}$ are the generators of $\mathfrak{gl}_{\mathcal{N}}$, provides a realization of $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$, [18, 43] (see also next section, comments in the proof of Corollary 4.2 on tensor representations of the quantum algebra in the special case of Yangian).

After the brief "interlude" regarding the Yangian case we return to quantum algebras associated to general brace solutions.

Proposition 3.9. Let X, Y be sets and $\check{r} : X \times X \to X \times X$, $\check{r}' : Y \times Y \to Y \times Y$ be functions. Let (X, \check{r}) and (Y, \check{r}') be set-theoretic solutions of the Braid equation. Let $f : X \to Y$ be a homomorphism of solutions (X, \check{r}) and (Y, \check{r}') , and f is onto Y. Then the map

$$L_{x,y}^{(k)} \to L_{f(x),f(y)}^{(k)}$$

is a homomorphism of quantum algebras $\mathfrak{A}(X,\check{r})$ and $\mathfrak{A}(Y,\check{r}')$.

Proof. We can verify that this function maps the defining relations of the quantum algebra $\mathfrak{A}(X, \check{r})$ onto defining relations of $\mathfrak{A}(Y, \check{r}')$.

Proposition 3.10. Let B be a brace, X be a subset of B and (X, \check{r}) be an involutive solution of the Braid relation obtained from this brace as in Lemma 1. Let J be an ideal of brace B. Let (X_J, \check{r}_J) be the solution associated to brace B/J on the subset $X_J = X + J$ of the factor brace B/J. Then the quantum group $\mathfrak{A}(X_J, \check{r}_J)$ is a representation of the quantum group $\mathfrak{A}(X, \check{r})$.

We call a solution (Y, \check{r}) of the Braid equation trivial if and only if $\check{r}(x, y) = (y, x)$ for all $x, y \in Y$. In this case we may denote \check{r} as τ .

Proposition 3.11. A set-theoretic solution (X, \check{r}) which can be homomorphically mapped onto a trivial solution (Y, τ) will have $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$ as its quantum group, where \mathcal{N} is the cardinality of Y.

Proof. The trivial solution (Y, τ) of cardinality \mathcal{N} has $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$ as its quantum group by Corollary 3.7. By Proposition 3.10, a solution (X, \check{r}) which can be homomorphically mapped onto a trivial solution (Y, τ) will have $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$ as its quantum group.

Recall that if (X, \check{r}) is a non-degenerate, involutive set-theoretic solution of the Braid equation, and $Y \subseteq X$, $e \in Y$, then Y is an orbit of e if for $x \in X, y \in Y$ we have $\sigma_x(y) \in Y$ and $\tau_x(y) \in Y$ and Y is the smallest set with this property.

Corollary 3.12. Let (X,\check{r}) is an involutive, non-degenerate solution with \mathcal{N} orbits, then $\mathcal{Y}(\mathfrak{gl}_{\mathcal{N}})$ is a representation of the quantum group $\mathfrak{A}(X,\check{r})$.

Proof. We can map (X, \check{r}) onto a trivial solution by mapping each element on its orbit. It is easy to check that this map is a homomorphism of set-theoretic solutions. The result now follows from Proposition 3.11.

Some basic information about orbits and examples of orbits can be found in [48], page 90 (just above section 2.2). Solutions which have only one orbit are called indecomposable solutions. Indecomposable involutive non-degenerate set-theoretic solutions of the Yang-Baxter equation can be constructed using one-generator braces [48].

Let $m \in \mathbb{Z}^+$ and $X_m = \{1, 2, ..., m\}$, define also $\check{r}_m(i, j) = (j + 1, i - 1)$ where addition and subtraction are taken modulo m. This is a special type of Lyubashenko solution [14]. We define \mathfrak{A}_m to be the quantum group $\mathfrak{A}(X_m, \check{r}_m)$. The following shows that Lyubashenko's solutions are useful for constructing representations of quantum groups constructed from braces.

Proposition 3.13. Let (X, \check{r}) be a finite, indecomposable, involutive and nondegenerate set theoretic solution of a finite multipermutation level. Suppose that $\sigma_x(z) \neq \sigma_y(z)$ for some $x, y, z \in X$. Then the quantum group $\mathfrak{A}(X, \check{r})$ associated to the solution (X, \check{r}) can be mapped onto the quantum algebra \mathfrak{A}_m for some m > 1.

Proof. Let $(X_{ret}, \check{r}_{ret})$ be the retraction of (X', \check{r}) . Notice that the retraction of (X, \check{r}) has more than one element, since $\sigma_x(z) \neq \sigma_y(z)$ for some $x, y, z \in X$. Notice that $(X_{ret}, \check{r}_{ret})$ satisfies the assumptions of Proposition 7.1 from [48], hence it can be mapped onto solution (X_m, \check{r}_m) for some m > 1.

Proposition 3.13 suggests the following question:

Question 1. Investigate representations of the quantum group \mathfrak{A}_m associated to the Lyubashenko solutions.

3.3. Representations of quantum groups. Let (X, \check{r}) be a set theoretic solution of the braid equation and let $A_{(X,\check{r})}$ be arbitrary associative algebra generated by elements from the set X and satisfying relations xy = uv whenever $\check{r}(x, y) = (u, v)$.

We show several representations of our quantum group using algebra $A_{(X,\check{r})}$ give representations of the quantum group $\mathfrak{A}(X,\check{r})$ associated to solution (X,\check{r}) .

Proposition 3.14. Let (X, \check{r}) be a set theoretic solution of the braid equation and let $A_{(X,\check{r})}$ be an arbitrary associative algebra generated by elements from the set X and satisfying relations xy = uv whenever $\check{r}(x, y) = (u, v)$. The following holds:

(1) $A_{(X,\check{r})}$ is a representation of the quantum group $\mathfrak{A}(X,\check{r})$ when we map $L_{x,y}^{(n)}$ to $x \in A_{(X,\check{r})}$ for every n.

- (2) $A_{(X,\check{r})} \otimes A_{(X,\check{r})}$ is a representation of the quantum group $\mathfrak{A}(X,\check{r})$ when we map $L_{x,y}^{(n)}$ to $x \otimes y \in A_{(X,\check{r})} \otimes A_{(X,\check{r})}$ for every n.
- (3) Let R be an arbitrary commutative associative algebra over the field \mathbb{C} generated by elements c_1, c_2, \ldots Then $A_{(X,\check{r})} \otimes R$ is a representation of the quantum group $\mathfrak{A}(X,\check{r})$ when we map $L_{x,y}^{(n)}$ to $x \otimes c_n \in A_{(X,\check{r})} \otimes R$ for every n.
- (4) In the above point 3 if $0 = c_2 = c_3 = \dots$ then we obtain a representation satisfying $L(\lambda) = L_0 + \lambda L_1$ (where L_1, L_0 are independent of λ).

Proof. The proof is by verifying that the quantum algebra relations will go to zero after applying the above homomorphic images of algebras (related to the above representations). \Box

Examples of algebras $A_{(X,\check{r})}$. Let (X,\check{r}) be an indecomposable, involutive settheoretic solution of the Yang-Baxter equation. Various types of algebras belonging to the class $A_{(X,\check{\tau})}$ were investigated extensively by several authors, and a lot is known about them [33, 34]. Quantum binomial algebras were introduced and investigated by Gateva-Ivanova in [20], [22] and [21]. The monomial algebras of I type were investigated for involutive solutions in [25], [32], [35], [36], and recently for both involutive and non-involutive solutions in [23] and [30]. The structure algebras of set-theoretic solutions. The algebra generated by the set X and with defining relations xy = uv if $\check{r}(x,y) = (u,v)$ was investigated in [30], where in section 5 they study prime ideals in such algebras and hence representations of such algebras which are prime (they showed that under mild assumptions they correspond to prime ideals of a group algebra associated to the same set-theoretic solution). For involutive solutions such algebras were previously investigated in [25, 32, 35, 36]. Let G be a permutation group of a finite, non-degenerate, involutive set-theoretic solution (X,\check{r}) of the Braid equation. Then $A_{(X,\check{r})}$ can be taken to be the group algebra $\mathbb{C}[G]$. Such algebras were investigated in [16].

Research directions and open questions

Question 1. Let (X, \check{r}) be an involutive, non-degenerate set-theoretic solution. Does the quantum algebra $\mathfrak{A}_{(X,\check{r})}$ have a finite Gröbner basis?

Question 2. What can be said about the quantum group $\mathfrak{A}_{(X,\tilde{r})}$ associated to a solution (X,\tilde{r}) of a finite multi-permutation level? Or of an indecomposable solution? Does $\mathfrak{A}_{(X,\tilde{r})}$ have any representations of small dimensions?

4. Novel class of quantum integrable systems & associated symmetries

4.1. Quantum spin chains from braces: novel class of integrable systems.

We introduce in this subsection physical spin-chain like systems, with periodic boundary conditions, associated to braces, and we investigate the corresponding symmetries. Our main objectives in this and the subsequent subsection is the derivation of tensor representations of the underlying quantum algebra, and the identification of potential symmetries of the transfer matrix of spin chain-like models. The transfer matrix, which will be defined below, is the generating function of a family of mutually commuting quantities, which gaurantee in principle the quantum integrability of the spin-chain system.

Indeed, let us introduce the monodromy matrix, which is a tensor representation of the quantum group (3.16), [18]

(4.1)
$$T_0(\lambda) = R_{0N}(\lambda) \dots R_{02}(\lambda) R_{01}(\lambda)$$

 $T \in \operatorname{End}(\mathbb{C}^{\mathcal{N}} \otimes (\mathbb{C}^{\mathcal{N}})^{\otimes N})$, and recall $R = \mathcal{P}\check{R}$, and in the case of brace solutions is given by (3.3), (2.2). We define also the transfer matrix $\mathfrak{t}(\lambda) = tr_0(T_0(\lambda)) \in$ $\operatorname{End}((\mathbb{C}^{\mathcal{N}})^{\otimes N})$. Given that T satisfies (3.16), we conclude that the transfer matrix provides mutually commuting quantities: $(\mathfrak{t}(\lambda) = \lambda^N \sum_k \frac{\mathfrak{t}^{(k)}}{\lambda^k})$

(4.2)
$$\left[\mathfrak{t}(\lambda), \ \mathfrak{t}(\mu)\right] = 0 \ \Rightarrow \ \left[\mathfrak{t}^{(k)}, \ \mathfrak{t}^{(l)}\right] = 0.$$

We present below one of our main Propositions, which will have important implications when studying the symmetries of the related periodic transfer matrices, as will become clear in the next subsection. This result becomes even more prominent when integrable boundary conditions are implemented to integrable systems, especially those coming from braces [13]. The following Proposition 4.1 and Lemma 4.2 are quite general and hold for any $R(\lambda) = \lambda \mathcal{P}\check{r} + \mathcal{P}$, where \check{r} provides a representation of the A-type Hecke algebra $\mathcal{H}_N(q=1)$, i.e. satisfies the braid relation and $\check{r}^2 = \mathbb{I}$, and \mathcal{P} is the permutation operator.

Proposition 4.1. Consider the λ -series expansion of the monodromy matrix: $T(\lambda) = \lambda^N \sum_{k=0}^{N} \frac{T^{(k)}}{\lambda^k}$ for any $R(\lambda) = \lambda \mathcal{P}\check{r} + \mathcal{P}$, where \check{r} provides a representation of the A-type Hecke algebra $\mathcal{H}_N(q = 1)$. Let also $H^{(k)} = \mathfrak{t}^{(k)}(\mathfrak{t}^{(N)})^{-1}$, $k = 0, \ldots, N-1$ and $H^{(N)} = \mathfrak{t}^{(N)}$, where $\mathfrak{t}^{(k)} = Tr_0(T_0^{(k)})$. Then the commuting quantities, $H^{(k)}$ for $k = 1, \ldots, N-1$, are expressed exclusively in terms of the elements \check{r}_{nn+1} , $n = 1, \ldots, N-1$, and \check{r}_{N1} .

Proof. Let us introduce some useful notation. We define, for $0 \le m < n \le N + 1$: $\mathbb{P}_{n-1;m+1} = \mathcal{P}_{0n-1} \dots \mathcal{P}_{0m+1}, \ n > m+2, \quad \mathbb{P}_{n-1;n} = \mathrm{id}, \quad \mathbb{P}_{n;n} = \mathcal{P}_{0n}$ and for $1 \le n \le N$: $\Pi = \mathcal{P}_{12} \ \mathcal{P}_{23} \dots \mathcal{P}_{N-1N}, \qquad \check{\mathfrak{R}}_{n;m} = \check{r}_{n-1n} \ \check{r}_{n-2n-1} \dots \check{r}_{mm+1}, \qquad n > m+1, \\ \check{\mathfrak{R}}_{n;n} = \mathrm{id}, \ \check{\mathfrak{R}}_{n+1;n} = \check{r}_{nn+1}. \text{ Note that } \log(\Pi) \text{ is the momentum operator of the system. Let us also define the ordered product:}$

$$\prod_{1 \le j \le k}^{\leftarrow} \check{r}_{n_j n_{j+1}} = \check{r}_{n_1 n_1 + 1} \check{r}_{n_2 n_2 + 1} \dots \check{r}_{n_k n_k + 1} : \quad n_1 > n_2 > \dots n_k.$$

We compute all the members of the expansion of the monodromy $T^{(k)}$, using the notation introduced above and the definition (4.1):

$$\begin{split} T_{0}^{(N)} &= \mathbb{P}_{N;1} = \mathcal{P}_{01}\Pi, \\ T_{0}^{(N-1)} &= \sum_{n=1}^{N} \mathbb{P}_{N;n+1} r_{0n} \mathbb{P}_{n-1;1} = \Big(\sum_{n=1}^{N-1} \check{r}_{nn+1} + \check{r}_{N0}\Big) \mathcal{P}_{01}\Pi, \dots \\ T_{0}^{(N-k)} &= \sum_{1 \le n_k < \dots < n_1 \le N} \prod_{j=1}^{k} \mathbb{P}_{n_{j-1}-1;n_j+1} r_{0n_j} \mathbb{P}_{n_k-1;1} = \\ \Big(\sum_{1 \le n_k < \dots < n_1 < N} \prod_{1 \le j \le k}^{\leftarrow} \check{r}_{n_j n_{j+1}} + \sum_{1 \le n_k < \dots < n_2 < N} \prod_{2 \le j \le k}^{\leftarrow} \check{r}_{N0} \check{r}_{n_j n_{j+1}}\Big) \mathcal{P}_{01}\Pi, \dots \\ T_{0}^{(1)} &= \Big(\sum_{n=1}^{N-1} \check{r}_{N0} \check{\Re}_{N;n+1} \check{\Re}_{n;1} + \check{\Re}_{N;1}\Big) \mathcal{P}_{01}\Pi \\ T_{0}^{(0)} &= \mathbb{R}_{N;1} = \check{r}_{N0} \check{\Re}_{N;1} \mathcal{P}_{01}\Pi. \end{split}$$

Recall that we can express the monodromy matrix and consequently all $T^{(k)}$ in a block form, i.e. $T^{(k)} = \sum_{x,y \in X} e_{x,y} \otimes T^{(k)}_{x,y}$, thus $\mathfrak{t}^{(k)} = tr_0(T^{(k)}_0) = \sum_{x \in X} T^{(k)}_{x,x}$. $\mathfrak{t}^{(k)}$ commute among each other, and hence any combination of them also provides a family of mutually commuting quantities. For instance, we consider the following convenient combination: $H^{(k)} = \mathfrak{t}^{(k)}(\mathfrak{t}^{(N)})^{-1}$, $k = 1, \ldots, N-1$ and $H^{(N)} = \mathfrak{t}^{(N)} =$ Π , then (periodicity is naturally imposed after taking the trace, $N + 1 \equiv 1$):

$$\begin{split} H^{(N-1)} &= \sum_{n=1}^{N} \check{r}_{nn+1}, \\ H^{(N-2)} &= \sum_{1 \le m < n \le N} \check{r}_{nn+1} \check{r}_{mm+1} + \sum_{n=1}^{N-2} \check{r}_{nn+1} \check{r}_{N1} + \check{r}_{N1} \check{r}_{N-1N}, \ \dots \\ H^{(N-k)} &= \sum_{1 \le n_k < \dots < n_1 < N} \prod_{1 \le j \le k}^{\leftarrow} \check{r}_{n_j n_{j+1}} + \sum_{1 \le n_k < \dots < n_2 < N-1} \prod_{2 \le j \le k}^{\leftarrow} \check{r}_{n_j n_{j+1}} \check{r}_{N1} \\ &+ \sum_{1 \le n_k < \dots < n_2 = N-1} \prod_{l+1 \le j \le k}^{\leftarrow} \check{r}_{n_j n_{j+1}} \check{r}_{N1} \prod_{2 \le j \le l}^{\leftarrow} \check{r}_{n_j n_{j+1}} \Big|_{c_j = 0, \ c_l > 0}, \ \dots \\ H^{(1)} &= \sum_{n=1}^{N-1} \check{\Re}_{n;1} \check{r}_{N1} \check{\Re}_{N;n+1} + \check{\Re}_{N;1}, \end{split}$$

where we define $c_j = n_j - n_{j+1} - 1$, $2 \le j < l$ and $c_l = n_l - n_{l+1} - 1$. Indeed, the Hamiltonians $H^{(k)}$ for $k = 1, \ldots, N-1$ are expressed solely in terms of the elements \check{r}_{nn+1} and \check{r}_{N1} and this essentially concludes our proof. Let us also for the sake of completeness report $H^{(0)}$, which takes the simple form $H^{(0)} = tr_0 (\check{r}_{N0} \check{\mathfrak{R}}_{N;1} \mathcal{P}_{01})$, but as opposed to the rest of the commuting Hamiltonians $H^{(k)}$, $k = 1, \ldots, N-1$, it can not be expressed only in terms of \check{r}_{nn+1} and \check{r}_{N1} .

It will be also instructive for our purposes here, related especially with Proposition 4.11 (presented later in the text), to compute explicitly $T^{(0)}$ and $\mathfrak{t}^{(0)}$. Recalling expression (4.3) and the form of the brace solution (2.1) we have:

(4.3)
$$T^{(0)} = \sum_{x_1, \dots, x_N, y_1, \dots, y_N \in X} e_{y_N, \sigma_{x_1}(y_1)} \otimes e_{x_1}, \tau_{y_1(x_1)} \otimes \dots \otimes e_{x_N}, \tau_{y_N(x_N)}$$

and by taking the trace (periodic boundary conditions: $N + 1 \equiv 1$)

(4.4)
$$\mathfrak{t}^{(0)} = \sum_{x_1, \dots, x_N, y_1, \dots, y_N \in X} e_{x_1}, \tau_{y_1(x_1)} \otimes \dots \otimes e_{x_N}, \tau_{y_N(x_N)},$$

where both expressions above are subject to the constraints: $y_n = \sigma_{x_{n+1}}(y_{n+1})$.

We show below an interesting property regarding the element $\hat{\mathfrak{R}}_{N-1;1}$ introduced in the proof of the latter Proposition (see also relevant findings in connection to Murphy elements in Hecke algebras in [11]).

Lemma 4.2. The action of $\check{\mathfrak{R}}_{N;1} = \check{r}_{N-1N} \check{r}_{N-2N-1} \dots \check{r}_{12}$ on the elements of the A-type Hecke algebra is given by

$$\begin{aligned} \dot{\mathfrak{R}}_{N;1} \ \check{r}_{nn+1} &= \check{r}_{n-1n} \ \dot{\mathfrak{R}}_{N;1}, \quad \forall n \in \{2, \dots, N-2\} \\ \check{\mathfrak{R}}_{N:1} \ \check{r}_{12} &= \check{\mathfrak{R}}_{N:2}, \quad \check{r}_{N-1N} \ \dot{\mathfrak{R}}_{N:1} &= \check{\mathfrak{R}}_{N-1:1} \end{aligned}$$

Proof. The poof is straightforward via the use of the braid relation $\check{r}_{nn+1}\check{r}_{n+1n+2}\check{r}_{nn+1} = \check{r}_{n+1n+2}\check{r}_{nn+1}\check{r}_{n+1n+2}, \,\check{r}^2 = I \otimes I$, and the form of $\check{\Re}_{N;1}$. \Box

We showed that the Hamiltonians $H^{(k)}$, k = 1, ..., N - 1 introduced in Proposition 4.1 are expressed exclusively in terms of the A-type Hecke elements and the periodic element \check{r}_{N1} . This fact will be exploited later when investigating the symmetries of the conserved quantities for various brace solutions. In the special case where $\check{r} = \mathcal{P}$, i.e. the Yangian (see also Corollary below), the Hamiltonian is $\mathfrak{gl}_{\mathcal{N}}$ symmetric (see also (3.24)). However, if we focus on the more general brace solution we conclude that there is no non-abelian algebra as symmetry of the Hamiltonian or in general of the transfer matrix, with the exception of certain special cases that will be examined later in the text. Note that in [13] the existence of a non-abelian algebra that is also a symmetry of the open boundary Hamiltonian is shown (see also [12] and references therein for relevant findings). The notion of the so-called Murphy elements associated to Hecke algebras, emerging from open boundary transfer matrices [11], is also discussed in [13] for *R*-matrices that come from braces. Moreover, for a special class of set theoretical solutions and for a special choice of boundary conditions it is shown [13], that not only the corresponding Hamiltonian is $\mathfrak{gl}_{\mathcal{N}}$ symmetric, but also the boundary transfer matrix.

Let us recall in the next corollary the known result about the existence of a non-abelian algebra that is a symmetry of the transfer matrix in the Yangian case.

Corollary 4.3. In the case of Yangian ($\check{r} = \mathcal{P}$), $\mathfrak{gl}_{\mathcal{N}}$ is a symmetry of the periodic transfer matrix.

Proof. Indeed, if $r = I \otimes I$, then $T^{(0)} = I^{\otimes (N+1)}$, and from (3.28) it follows that

(4.5)
$$\left[T_1^{(1)}, \ T_2^{(m)}\right] = T_2^{(m)} \mathcal{P}_{12} - \mathcal{P}_{12} T_2^{(m)}.$$

By taking the trace over the second space we conclude

(4.6)
$$\left[T_1^{(1)}, \mathfrak{t}^{(m)}\right] = 0 \Rightarrow \left[T_{i,j}^{(1)}, \mathfrak{t}(\lambda)\right] = 0$$

 $T_{i,j}^{(1)} \in \operatorname{End}((\mathbb{C}^{\mathcal{N}})^{\otimes N})$ are the entries of the *T* matrix, and are tensor representations of the $\mathfrak{gl}_{\mathcal{N}}$ algebra (3.7),

(4.7)
$$T_{i,j}^{(1)} = \Delta^{(N)}(e_{i,j}), \quad i, \ j \in \{1, \ 2, \dots, N\}$$

where the co-product is defined in (3.24), i.e. the transfer matrix enjoys the $\mathfrak{gl}_{\mathcal{N}}$ symmetry.

4.2. New symmetries of the periodic transfer matrix. The ultimate goal in the context of quantum integrable systems, or any quantum system for that matter, is the identification of the eigenvalues and eigenvectors of the corresponding Hamiltonian. In the frame of quantum integrable systems more specifically there exists a set of mutually commuting "Hamiltonians", guaranteed by the existence of a quantum R-matrix that satisfies the Yang-Baxter equation. As already discussed in the previous subsection this set of mutually commuting objects is generated by the transfer matrix. Thus the derivation of the eigenvalues and eigenstates of the transfer matrix is the significant problem within quantum intgrability. This is in general an intricate task and the typical methodology used is the Bethe ansatz formulation, or suitable generalizations, depending on the problem at hand. A detailed study of this problem for transfer matrices associated to brace solutions will be presented in future investigations.

Here we are focusing primarily on the investigation of possible existing new symmetries of the periodic transfer matrix, as any information regarding the symmetries of the transfer matrix provides for instance valuable insight on the multiplicities occurring in the spectrum. It will be transparent in what follows that the study of the symmetries of the *R*-matrices is a first step towards formulating the symmetries of the transfer matrix. A detailed analysis on more generic symmetry algebras and boundary conditions is presented in [13].

Research Directions and Open Questions.

Question 1. Let (X, \check{r}) be a set theoretical solution of the Yang-Baxter equation. The next natural step is to investigate the existence of non-abelian algebras that are symmetries of the general open boundary transfer matrix, i.e. generalize Corollary 4.3 for any brace solution. This is a fundamental problem and is investigated in [13].

We will provide below some new examples of symmetries of transfer matrices. We will use the following known fact:

Lemma 4.4. Let R be a solution of the Yang-Baxter equation, and B be a $\mathcal{N} \times \mathcal{N}$ matrix such that

(4.8)
$$(B \otimes B)R(\lambda) = R(\lambda)(B \otimes B),$$

then

(4.9)
$$(B \otimes B^{\otimes N})T(\lambda) = T(\lambda)(B \otimes B^{\otimes N}),$$

where $T(\lambda)$ is the monodromy matrix.

Proof. To prove this it is convenient to employ the index notation. Indeed, in the index notation expression (4.8) is translated into: $\forall n \in \{1, \ldots, N\}$

$$B_0 B_n R_{0n}(\lambda) = R_{0n}(\lambda) B_0 B_n \Rightarrow$$

$$B_0 B_1 \dots B_N R_{0N}(\lambda) \dots R_{01}(\lambda) = R_{0N}(\lambda) \dots R_{01}(\lambda) B_0 B_1 \dots B_N.$$

The latter expression is apparently equivalent to (4.9).

We will also use the following obvious fact:

Lemma 4.5. If B is an $\mathcal{N} \times \mathcal{N}$ matrix and $\mathcal{P} = \sum_{1 \leq i, j \leq \mathcal{N}} e_{i,j} \otimes e_{j,i}$ then

$$(B \otimes B)\mathcal{P} = \mathcal{P}(B \otimes B).$$

Proposition 4.6. Let (X,\check{r}) be a set-theoretic solution of the braid equation and let $f: X \to X$ be an isomorphism of solutions, so $f(\sigma_x(y)) = \sigma_{f(x)}(f(y))$ and

 $f(\tau_y(x)) = \tau_{f(y)}(f(x))$. Denote $M = \sum_{x \in X} \alpha_x e_{x,f(x)}$ such that $a_x \in \mathbb{C}, \ \alpha_x \neq 0, \ \forall x \in X \text{ and } \alpha_x \alpha_y = \alpha_{\sigma_x(y)} \alpha_{\tau_y(x)}$, then

(4.10)
$$\left[M^{\otimes N}, \ \mathfrak{t}(\lambda)\right] = 0,$$

where $\mathfrak{t}(\lambda)$ is the transfer matrix for $R(\lambda) = \mathcal{P} + \lambda \mathcal{P}\check{r}$.

Proof. By means of Lemmata 4.4 and 4.5 it suffices to show that

$$(4.11) (M \otimes M)r = r(M \otimes M).$$

where recall $r = \mathcal{P}\check{r}$. Indeed, then by direct computation the LHS of (4.11) reduces to:

(4.12)
$$(M \otimes M)r = \sum_{x,y \in X} \alpha_x \alpha_y \ e_{y,f(\sigma_x(y))} \otimes e_{x,f(\tau_y(x))}$$

whereas the RHS gives:

(4.13)
$$r(M \otimes M) = \sum_{x,y \in X} \alpha_{\sigma_x(y)} \alpha_{\tau_y(x)} \ e_{y,f(\sigma_x(y))} \otimes e_{x,f(\tau_y(x))}$$

Comparing (4.12), (4.13) we arrive at (4.11), provided that $\alpha_x \alpha_y = \alpha_{\sigma_x(y)} \alpha_{\tau_y(x)}$.

Having shown (4.11) it then immediately follows from Lemmata 4.4 and 4.5 that

(4.14)
$$(M \otimes M^{\otimes N})T(\lambda) = T(\lambda)(M \otimes M^{\otimes N}).$$

From the latter equation we focus on each element of the matrices (LHS vs RHS in (4.14)) on the auxiliary space (recall the notation $T = \sum_{x,y} e_{x,y} \otimes T_{x,y}$):

(4.15)
$$\begin{aligned} \alpha_x M^{\otimes N} T_{f(x), f(y)} &= \alpha_y T_{x, y} M^{\otimes N} \Rightarrow \\ M^{\otimes N} T_{f(x), f(x)} &= T_{x, x} M^{\otimes N} \Rightarrow \left[M^{\otimes N}, \ \mathfrak{t}(\lambda) \right] = 0. \end{aligned}$$

We can explicitly express $M^{\otimes N}$ as

(4.16)
$$M^{\otimes N} = \sum_{x_1, x_2, \dots, x_N \in X} \prod_{k=1}^N \alpha_{x_k} e_{x_1, f(x_1)} \otimes e_{x_2, f(x_2)} \dots \otimes e_{x_N, f(x_N)}.$$

Notice also, that due to the symmetry of the *R*-matrix (4.11), one easily shows that if *T* satisfies the RTT relation then so *MT* does. If *M* is non-singular then $M^{\otimes N}$ is a similarity transformation that leaves the transfer matrix invariant, and naturally provides information on the multiplicities of the spectrum.

A special case of the proposition above is the obvious choice: $f(x) = x, \forall x \in X$. Also, we obtain the following as immediate corollaries. **Corollary 4.7.** Let (X,\check{r}) be a finite, non-degenerate involutive set-theoretic solution of the Yang-Baxter equation, and let Q_1, \ldots, Q_k be all the orbits of X. Let $\alpha_1, \ldots, \alpha_k \in \mathbb{C}$. Let $M = \sum_{j=1}^k \alpha_j M_j$, where $M_j = \sum_{i \in Q_j} e_{i,i}$, then

(4.17)
$$\left[M^{\otimes N}, \ \mathfrak{t}(\lambda)\right] = 0,$$

where $\mathfrak{t}(\lambda)$ is the transfer matrix for $R(\lambda) = \mathcal{P} + \lambda \mathcal{P}\check{r}$.

Proof. If all $\alpha_x \neq 0$ then the result follows from Corollary 4.6. Observe that

$$M^{\otimes N} = \sum_{i_1,\dots,i_n \in \mathbb{N}} M_{i_1,\dots,i_n} \alpha_1^{i_1} \cdots \alpha_N^{i_N}.$$

We know that for all non-zero choices of α_x the matrix $M^{\otimes N}$ commutes with $\mathfrak{t}(\lambda)$. By a "Vandermonde matrix argument" each matrix $M_{i_1,\ldots,i_n\in\mathbb{N}}$ commutes with $\mathfrak{t}(\lambda)$, which concludes the proof.

Corollary 4.8. Let (X,\check{r}) be a finite, non-degenerate involutive set-theoretic solution of the Yang-Baxter equation, and let G(X,r) be its structure group (i.e. the group generated by elements from X and their inverses subject to relations xy = uv when r(x, y) = (u, v)). Let $\alpha : G(X, r) \to \mathbb{C}^*$ be a group homomorphism. Let $M = \sum_{x \in X} \alpha_x e_{x,x}$, then $M^{\otimes N}$ commutes with the transfer matrix.

Similarly, as in Proposition 4.6 we use Lemmata 4.4 and 4.5 in the following proofs.

Proposition 4.9. Let (X, \check{r}) be a finite, non degenerate involutive set-theoretic solution of the braid equation. Let $x_1, \ldots, x_\alpha \in X$ for some $\alpha \in \{1, \ldots, \mathcal{N}\}$. Assume that $\check{r}(x_i, y) = (y, x_i)$ for every $y \in X$. Then $\forall i, j \in \{1, \ldots, \alpha\}$:

(4.18)
$$\left[\Delta^{(N)}(e_{x_i,x_j}), \ \mathfrak{t}(\lambda)\right] = 0,$$

where $\mathfrak{t}(\lambda)$ is the transfer matrix for $R(\lambda) = \mathcal{P} + \lambda \mathcal{P}\check{r}$.

Proof. We first recall that $\Delta^{(N)}(e_{x_i,x_j})$ is defined in (3.24). We will show that $\forall i, j \in \{1, \ldots, \alpha\},\$

(4.19)
$$\Delta(e_{x_i,x_j})r = r\Delta(e_{x_i,x_j}).$$

Indeed, by direct computation: $\Delta(e_{x_i,x_j})r = \Delta(e_{x_i,x_j}) = r\Delta(e_{x_i,x_j})$, i.e. (4.19). It then immediately follows from (4.19) that

(4.20)
$$\Delta^{(N+1)}(e_{x_i,x_j})T(\lambda) = T(\lambda)\Delta^{(N+1)}(e_{x_i,x_j}).$$

Recalling on the definition of the co-product (3.24), and the notation $T(\lambda) = \sum_{x,y \in X} e_{x,y} \otimes T_{x,y}(\lambda)$ we conclude that expression (4.20) leads to

(4.21)
$$\sum_{w} e_{x_{i},w} \otimes T_{x_{j},w} + \sum_{z,w} e_{z,w} \otimes \Delta^{(N)}(e_{x_{i},x_{j}})T_{z,w}(\lambda) = \sum_{w} e_{z,x_{j}} \otimes T_{z,x_{i}} + \sum_{z,w} e_{z,w} \otimes T_{z,w}(\lambda)\Delta^{(N)}(e_{x_{i},x_{j}})$$

We focus on the diagonal entries of the latter expression and we obtain:

$$\left[\Delta^{(N)}(e_{x_i,x_j}), \ T_{x_i,x_i}(\lambda)\right] = -T_{x_j,x_i}(\lambda) + \delta_{ij}T_{x_j,x_i}(\lambda)$$
$$\left[\Delta^{(N)}(e_{x_i,x_j}), \ T_{x_j,x_j}(\lambda)\right] = T_{x_j,x_i}(\lambda) - \delta_{ij}T_{x_j,x_i}(\lambda)$$
$$\left[\Delta^{(N)}(e_{x_i,x_j}), \ T_{z,z}(\lambda)\right] = 0, \quad z \neq x_i, \ x_j.$$

Summing up all the terms above we arrive at (4.18), $\left[\sum_{x \in X} T_{x,x}(\lambda), \Delta^{(N)}(e_{x_i,x_j})\right] = 0, \quad \forall i, j \in \{1, \ldots, \alpha\}$, i.e. we conclude that the transfer matrix is \mathfrak{gl}_{α} symmetric.

Lemma 4.10. Let $r = \mathcal{P}\check{r}$ and

(4.23)
$$\Delta(e_{x_i,x_j})r = r\Delta(e_{x_i,x_j})$$

for some $x_i, x_j \in X$, then

$$(4.24) e_{x_i,x_j} \otimes e_{x_i,x_j} r = r e_{x_i,x_j} \otimes e_{x_i,x_j}$$

and also, $\forall n \in \{1, \ldots, N\}$:

(4.25)
$$\left[\sum_{m_1 < m_2 \dots < m_n = 1}^N (e_{x_i, x_j})_{m_1} \dots (e_{x_i, x_j})_{m_n}, \ \mathfrak{t}(\lambda)\right] = 0,$$

where we use the standard index notation,

(4.26)
$$(e_{x_i,x_j})_m = I \otimes \dots I \otimes \underbrace{e_{x_i,x_j}}_{m^{th}} \bigotimes I \dots \otimes I,$$

where I appears N-1 times.

Proof. The proof is straightforward: from the definition of the co-product Δ , the fact that $e_{x,y}^2 = e_{x,y}\delta_{xy}$ and also $(\Delta(e_{x_i,x_j}))^2 r = r(\Delta(e_{x_i,x_j}))^2$, we arrive at (4.24).

Similarly as in the proof of Proposition 4.9 we know that if $\Delta(e_{x_i,x_j})r = r\Delta(e_{x_i,x_j})$, then $\left[\Delta^{(N)}(e_{x_i,x_j}), t(\lambda)\right] = 0$, and $\left[\left(\Delta^{(N)}(e_{x_i,x_j})\right)^n, t(\lambda)\right] = 0$, $n \in \{1, \ldots, N\}$, which together with $e_{x,y}^2 = e_{x,y}\delta_{xy}$ lead to (4.25).

We obtain in what follows a general class of symmetries, associated to solutions endowed with some extra special properties. Recall that solutions (X, r) such that r(x, x) = (x, x) for every $x \in X$ are called square free, and they were introduced by Gateva-Ivanova. There was a famous conjecture by Gateva-Ivanova as to whether or not these solutions need to have a finite multi-permutation level. Vendramin subsequently showed that this was not necessary [51]. Later Cedó, Jespers and Okniński investigated these solutions using wreath products of groups and gave many interesting examples of square-free solutions.

Proposition 4.11. Let (X, \check{r}) be a finite non degenerate involutive set-theoretic solution of the braid equation. Let $x_1, \ldots, x_\alpha \in X$ for some $\alpha \in \{1, \ldots, N\}$ be such that $\check{r}(x_j, x_j) = (x_j, x_j)$. Let $\mathfrak{t}(\lambda) = \lambda^N \sum_{k=0}^N \mathfrak{t}^{(k)} \lambda^{-k}$ be the transfer matrix for $R(u) = P + uP\check{r}$. Then $\forall i, j = 1, \ldots, \alpha$ and $k = 1, \ldots, N$

$$\left[e_{x_i,x_j}^{\otimes N}, \ \mathfrak{t}^{(k)}\right] = 0.$$

Proof. We first show by direct computation that:

$$e_{x_i,x_j} \otimes e_{x_i,x_j}\check{r} = \check{r}e_{x_i,x_j} \otimes e_{x_i,x_j} \Rightarrow$$

(4.27)
$$e_{x_i,x_i}^{\otimes N}\check{r}_{nn+1} = \check{r}_{nn+1}e_{x_i,x_i}^{\otimes N}, \quad n = 1, \dots, N-1,$$

(4.28)
$$e_{x_i,x_j}^{\otimes N}\check{r}_{1N} = \check{r}_{1N}e_{x_i,x_j}^{\otimes N}$$

By multiplying equation (4.28) with \mathcal{P}_{1N} we conclude that: $\left[e_{x_i,x_j}^{\otimes N}, \check{r}_{N1}\right] = 0.$

Then recall from Proposition 4.1 that $H^{(k)} = \mathfrak{t}^{(k)}(\mathfrak{t}^{(N)})^{-1}$ for k = 1, 2, ..., N-1are expressed exclusively in terms of \check{r}_{nn+1} , n = 1, ..., N-1 and \check{r}_{N1} . Recall also from the proof of Proposition 4.1 that $H^{(N)} = \mathfrak{t}^{(N)} = \mathcal{P}_{12}\mathcal{P}_{23}\ldots\mathcal{P}_{N-1N}$, which immediately leads to (from the definition of \mathcal{P}): $\begin{bmatrix} e_{x_i,x_j}^{\otimes N}, H^{(N)} \end{bmatrix} = 0$, which in turn together with expressions (4.27), (4.28) and Proposition 4.1 lead to: $\begin{bmatrix} e_{x_i,x_j}^{\otimes N}, H^{(k)} \end{bmatrix} = 0$, $k = 1, \ldots, N$. And due to $H^{(k)} = \mathfrak{t}^{(k)}(\mathfrak{t}^{(N)})^{-1}$ for $k = 1, \ldots, N-1$ and $H^{(N)} = \mathfrak{t}^{(N)}$, we conclude that $\begin{bmatrix} e_{x_i,x_j}^{\otimes N}, \mathfrak{t}^{(k)} \end{bmatrix} = 0$ for $k = 1, \ldots, N$ and $i, j = 1, \ldots, \alpha$.

Recall that by Remark 2.5 every nilpotent ring is a brace when we define $a \circ b = ab + a + b$ (we call this brace the corresponding brace).

We restrict our attention in what follows in the case where N is odd. Let $(B, +, \circ)$ be a brace. We say that $a \in B$ is central if $a \circ b = b \circ a$ for every $b \in B$.

Proposition 4.12. Let $(B, +, \cdot)$ be a nilpotent ring and $(B, +, \circ)$ be the corresponding brace. Let a be a central element in B, and a + a = 0, $a \circ a = 0$. Let $X \subseteq B$ and let (X, \check{r}) be a finite non degenerate involutive set-theoretic solution of the braid equation constructed from B as in Theorem 2.4. Let $x, y \in X$ and $x = \sigma_b(a) = ba + a$, $y = \sigma_c(a) = ca + a$ for some $b, c \in B$. Then

(4.29)
$$\left[e_{x,y}^{\otimes N}, \ \mathfrak{t}(\lambda)\right] = 0.$$

Proof. First recall that $\mathfrak{t}(\lambda) = \lambda^N \sum_{k=0}^N \frac{\mathfrak{t}^{(k)}}{\lambda^k}$, recall also that we can express the monodromy matrix and consequently all $T^{(k)}$ in a block form, i.e. $T^{(k)} = \sum_{z,w \in X} e_{z,w} \otimes T^{(k)}_{z,w}$, thus $\mathfrak{t}^{(k)} = tr_0(T^{(k)}_0) = \sum_{z \in X} T^{(k)}_{z,z}$. Via Proposition 4.11 it suffices to show that $e^{\otimes N}_{x,y}$ commutes with $\mathfrak{t}^{(0)}$.

Denote

$$W_{p_1,\ldots,p_N} = Q_{p_1,p_2} \otimes Q_{p_2,p_3} \otimes \cdots \otimes Q_{p_{N-1},p_N} \otimes Q_{p_N,p_1},$$

where $Q_{i,p} = \sum_{j \in W_{i,p}} e_{j,j^i}$ and $W_{i,p} = \{j : j^i = p\}$. Recall also the form of $\mathfrak{t}^{(0)}$ from the proof of Proposition 4.1, expressed as

$$\mathfrak{t}^{(0)} = \sum_{p_1,\dots,p_N \leq \mathcal{N}} W_{p_1,\dots,p_N} = Q_{p_1,p_2} \otimes Q_{p_2,p_3} \otimes \dots \otimes Q_{p_{N-1},p_N} \otimes Q_{p_N,p_1}$$

where \mathcal{N} is the cardinality of X. We will show that

$$e_{x,y}^{\otimes N}W_{p_1,\ldots,p_N} = W_{p_1,\ldots,p_N}e_{x,y}^{\otimes N},$$

for all $p_1, \ldots, p_N \leq \mathcal{N}$.

Part 1. We will first calculate $e_{x,y}^{\otimes N} W_{p_1,\dots,p_N}$.

Notice that $e_{x,y}^{\otimes N}W_{p_1,\ldots,p_N} = e_{x,y}Q_{p_1,p_2} \otimes e_{x,y}Q_{p_2,p_3} \otimes \cdots \otimes e_{x,y}Q_{p_N,p_1}$. If it is non zero then $e_{x,y}Q_{p_i,p_{i+1}} \neq 0$ for every *i*.

Notice that $e_{x,y}Q_{p_i,p_{i+1}} = \sum_{j \in W_{p_i,p_{i+1}}} e_{x,y}e_{j,j^{p_i}}$. If $e_{x,y}e_{j,j^{p_i}} \neq 0$ then j = y, and $j^{p_i} = y^{p_i}$, hence

$$e_{x,y}Q_{p_i,p_{i+1}} = e_{x,y^{p_i}}.$$

Notice that $y = j \in W_{p_i,p_{i+1}}$, hence ${}^{y}p_i = p_{i+1}$. Similarly ${}^{y}p_{i+1} = p_{i+2}$, this implies ${}^{y \circ y}p_i = p_{i+2}$. Observe that $y \circ y = 0$, and so $p_i = p_{i+2}$ for every *i* (where $p_{i+N} = p_i$). Since *N* is odd it follows that $p_1 = p_2 = \dots p_N$ and ${}^{y}p_i = p_i \ \forall i$.

Now ${}^{y}p_{i} = p_{i+1}$ implies ${}^{y}p_{i} = p_{i}$, hence ${}^{a}p_{i} = p_{i}$. It follows that $y^{p_{i}} = y$ (since a is central in B). Therefore, $e_{x,y}^{\otimes N}W_{p_{1},\ldots,p_{N}} = e_{x,y}^{\otimes N}$ provided that $p_{1} = \ldots = p_{N}$ and ${}^{a}p_{1} = p_{1}$, and otherwise it is zero.

Part 2. We will now calculate $W_{p_1,\ldots,p_N} e_{x,y}^{\otimes N}$.

Notice that if $W_{p_1,\ldots,p_N}e_{x,y}^{\otimes N} \neq 0$ then $Q_{p_i,p_{i+1}}e_{x,y} \neq 0$, for every *i*. Observe that $Q_{p_i,p_{i+1}}e_{x,y} = \sum_{j \in W_{p_i,p_{i+1}}} e_{j,j^{p_i}}e_{x,y}$. If $e_{j,j^{p_i}}e_{x,y} \neq 0$ then $j^{p_i} = x$. Hence $j = x^{q_i}$ where q_i is the inverse of p_i in the group (B, \circ) . It follows that $j = q_i \cdot a + a$. Therefore, for given p_i element j_i is uniquely determined. Consequently, $Q_{p_i,p_{i+1}}e_{x,y} = e_{j_i,y}$ and $j_i = j \in W_{p_i,p_{i+1}}$. This implies $j_i p_i = p_{i+1}$, similarly $j_{i+1}p_{i+1} = p_{i+2}$, etc. It follows that $jp_i = p_i$ where $j = j_{i+N-1} \circ \cdots \circ j_{i+1} \circ j_i$. Observe that j = a + qa for some $q \in B$ by assumptions on x. Previously it was shown that $j^i p_i = p_{i+1} = p_i$, hence $xp_i = p_i$. Therefore $ap_i = p_i$ for $i = 1, \ldots, N$. This implies $x^{p_i} = x$. Observe that $j_i = x^{q_i}$ notice that $q_i = p_i \circ \cdots \circ p_i$ and since $x^{p_i} = x$ it follows that $j_i = x$. Therefore, $W_{p_1,\ldots,p_N}e_{x,y}^{\otimes N} = e_{x,y}^{\otimes N}$ provided that $p_1 = \ldots = p_N$ and ${}^ap_1 = p_1$, and otherwise it is zero. Consequently, $e_{x,y}^{\otimes N}W_{p_1,\ldots,p_N} = W_{p_1,\ldots,p_N}e_{x,y}^{\otimes N}$.

Acknowledgments. We are indebted to Robert Weston for illuminating discussions. AS thanks Robert Weston for explaining to her several topics from quantum integrable systems. We also thank Vincent Caudrelier, Ferran Cedó, Eric Jespers, Jan Okniński and Michael West for useful suggestions on the manuscript. AD acknowledges support from the EPSRC research grant EP/R009465/1, AS acknowledges support from the EPSRC programme grant EP/R034826/1.

References

- V.E. Adler, A.I. Bobenko and Yu.B. Suris, Classification of integrable equations on quadgraphs. The consistency approach, Commun. Math. Phys. 233 (2003) 513.
- [2] D. Bachiller, Counterexample to a conjecture about braces, J. Algebra, 453 (2016) 160–176.
- [3] D. Bachiller, F. Cedó, E. Jespers and J. Okniński, Iterated matched products of finite braces and simplicity; new solutions of the Yang-Baxter equation, Trans. Amer. Math. Soc. 370 (2018), 4881–4907.
- [4] R.J. Baxter, Exactly solved models in statistical mechanics, (Academic Press, 1982).
- [5] V.V. Bazhanov and S.M. Sergeev, YangBaxter maps, discrete integrable equations and quantum groups, Nucl. Phys. B926 (2018) 509-543.
- [6] A. Berenstein and D. Kazhdan, *Geometric and unipotent crystals*, GAFA special volume (2000) 188.
- [7] T. Brzezinski, Trusses: Between braces and ring, Trans. Amer. Math. Soc. 372 (2019), 4149–4176.
- [8] F. Catino, I. Colazzo and P. Stefanelli, Semi-braces and the Yang-Baxter equation, J. Algebra, 483, (2017).
- [9] F. Cedó, Left braces: solutions of the Yang-Baxter equation, Adv. Group Theory Appl., Vol. 5 (2018), 33–90.
- [10] F. Cedó, E. Jespers, and J. Okninski, Braces and the Yang-Baxter equation. Comm. Math. Phys., 327(1) (2014) 101116.
- [11] A. Doikou, Murphy elements from the double-row transfer matrix, J. Stat. Mech. (2009) L03003.
- [12] A. Doikou and R.I. Nepomechie, Bulk and Boundary S Matrices for the SU(N) Chain, Nucl. Phys. B521 (1998) 547-572.
- [13] A. Doikou and A. Smoktunowicz, Set-theoretical Yang-Baxter and reflection equations & quantum group symmetries, in preparation (2019).
- [14] V.G. Drinfeld, On some unsolved problems in quantum group t (ed.) Quantum groups, in: Lecture Notes in Math., vol. 1510, Springer-Verlag, Berlin, 1992, pp. 1-8.heory, in P.P. Kulish.
- [15] V.G. Drinfeld, Hopf algebras and the quantum YangBaxter equation, Soviet. Math. Dokl. 32 (1985) 254;

A new realization of Yangians and quantized affine algebras, Soviet. Math. Dokl. 36 (1988) 212.

[16] P. Etingof, T. Schedler and A. Soloviev, Set-theoretical solutions to the quantum YangBaxter equation, Duke Math. J. 100 (1999) 169–209.

- [17] P. Etingof, Geometric crystals and set-theoretical solutions to the quantum Yang-Baxter equations, Commun.algebra 31 (2003) 1961.
- [18] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, Quantization of Lie groups and Lie algebras, Leningrad Math. J. 1 (1990) 193.
- [19] L.D. Faddeev and L.A. Takhtajan, Hamiltonian Methods in the Theory of Solitons, Springer-Verlag (1987).
- [20] T. Gateva-Ivanova, Quadratic algebras, Yang-Baxter equation, and Artin- Schelter regularity, Adv. in Math. 230 (2012) 2152–2175.
- [21] T. Gateva-Ivanova, Skew polynomial rings with binomial relations, J. Algebra 185 (1996) 710753.
- [22] T. Gateva-Ivanova, Binomial skew polynomial rings, Artin-Schelter regularity, and binomial solutions of the Yang-Baxter equation, Serdica Mathematical Journal (2004) Volume: 30, Issue: 2-3, page 431–470.
- [23] T. Gateva-Ivanova, A combinatorial approach to noninvolutive set-theoretic solutions of the Yang-Baxter equation, preprint (2018), arXiv:1808.03938.
- [24] T. Gateva-Ivanova, Set-theoretic solutions of the YangBaxter equation, braces and symmetric groups, Adv. Math., 388(7) (2018) 649–701.
- [25] T. Gateva–Ivanova and M. Van den Bergh, Semigroups of I-type, J. Algebra 206 (1997) 97–112.
- [26] L. Guarnieri and L. Vendramin, Skew braces and the YangBaxter equation, Math. Comp., 86(307), (2017) 25192534.
- [27] G. Hatayama, A. Kuniba and T. Takagi, Soliton cellular automata associated with crystal bases, Nucl. Phys. B577 (2000) 619.
- [28] J. Hietarinta, Permutation-type solutions to the Yang-Baxter and other nsimplex equations, J. Phys. A30 (1997) 4757-4771.
- [29] P. Jedlicka, A. Pilitowska and A. Zamojska-Dzienio, The retraction relation for biracks, J. Pure Appl. Algebra 223 (2019) 3594–3610.
- [30] E. Jespers, E. Kubat and A. Van Antwerpen, The structure monoid and algebra of a nondegenerate set-theoretic solution of the Yang-Baxter equation, Trans. Amer. Math. Soc. 372 (2019), no. 10, 7191–7223.
- [31] E. Jespers, E. Kubat, A. Van Antwerpen and L. Vendramin, Factorizations of skew braces, Math. Ann. 375 (2019), no. 3-4, 1649–1663.
- [32] E. Jespers and J. Okniński, Monoids and groups of I-type, Algebr. Represent. Theory 8 (2005) 709–729.
- [33] E. Jespers and J. Okniński, Binomial semigroups, Journal of Algebra, 202 (1998), 250–275.
- [34] E. Jespers and J. Okniński, Noetherian semigroup algebras, Series: Algebra and Applications, Vol. 7, 2007.
- [35] E. Jespers, J. Okniński, and M. Van Campenhout, Finitely generated algebras defined by homogeneous quadratic monomial relations and their underlying monoids, J. Algebra 440 (2015) 72–99.
- [36] E. Jespers and M. Van Campenhout, Finitely generated algebras defined by homogeneous quadratic monomial relations and their underlying monoids II, J. Algebra 492 (2017) 524– 546.
- [37] M. Jimbo, A q-difference analogue of U(g) and the Yang-Baxter equation, Lett. Math. Phys. 10 (1985) 63.
- [38] M. Jimbo, Quantum R-matrix for the generalized Toda system, Comm. Math. Phys.102 (1986) 537547.

- [39] L.H. Kauffman, Virtual knot theory, European Journal of Combinatorics 20 (1999) 663-691.
- [40] A. Konovalov, A. Smoktunowicz and L. Vendramin, On skew braces and their ideals, Experimental Mathematics, April 2018, DOI: 10.1080/10586458.2018.1492476.
- [41] I. Lau, Left Brace With The Operation * Associative Is A Two-sided Brace, to appear in J. Algebra and its Applications.
- [42] V. Lebed and L. Vendramin, On structure groups of set-theoretical solutions to the Yang-Baxter equation, Proc. Edinburgh Math. Soc. (2019) online first, http://doi.org/10.1017/S0013091518000548.
- [43] A. Molev, M. Nazarov and G. Olshanski, Yangians and classical Lie algebras, Russ. Math. Surveys 51 (1996) 205.
- [44] V.G. Papageorgiou, Yu.B. Suris, A.G. Tongas and A.P. Veselov, On quadrirational Yang-Baxter Maps, SIGMA 6 (2010) 033.
- [45] W. Rump, A decomposition theorem for square-free unitary solutions of the quantum Yang-Baxter equation. A dv. Math., 193(1) (2005) 4055.
- [46] W. Rump, Braces, radical rings, and the quantum Yang-Baxter equation, J. Algebra, 307(1) (2007) 153170.
- [47] A. Smoktunowicz and L. Vendramin, On Skew Braces (with an appendix by N. Byott and L. Vendramin), Journal of Combinatorial Algebra Volume 2, Issue 1, (2018) 47-86.
- [48] A. Smoktunowicz and A. Smoktunowicz, Set-theoretic solutions of the YangBaxter equation and new classes of R-matrices, Linear Algebra and its Applications, Volume 546, 1 June 2018, pages 86–114.
- [49] Y.P. Sysak, The adjoint group of radical rings and related questions, in: Ischia Group Theory 2010, (proceedings of the conference: Ischia, Naples, Italy, 14-17 April 2010), pp. 344365, World Scientifc, Singapore 2011.
- [50] D. Takahashi and J. Satsuma, A soliton cellular automaton, J. Phys. Soc. Japan 59 (1990) 3514.
- [51] L. Vendramin, Extensions of set-theoretic solutions of the Yang-Baxter equation and a conjecture of Gateva-Ivanova. Journal of Pure and Applied Algebra, Volume 220, Issue 5, (2016) 1681–2076.
- [52] A.P. Veselov, Yang-Baxter maps and integrable dynamics, Phys. Lett. A314 (2003) 214.
- [53] C.N. Yang, Some exact results for the many-body problem in one dimension with repulsive delta-function interaction, Rev. Lett. 19 (1967) 1312.

(A. Doikou) DEPT OF MATHEMATICS, HERIOT-WATT UNIVERSITY, EDINBURGH EH14 4AS, AND MAXWELL INSTITUTE FOR MATHEMATICAL SCIENCES, EDINBURGH *E-mail address*: A.Doikou@hw.ac.uk

(A. Smoktunowicz) School of Mathematics, The University of Edinburgh, James Clerk Maxwell Building, The Kings Buildings, Mayfield Road EH9 3JZ, and Maxwell Intitute for Mathematical Sciences, Edinburgh

E-mail address: A.Smoktunowicz@ed.ac.uk