

A homotopy BV algebra for Yang-Mills and color-kinematics

Michael Reiterer

The Hebrew University of Jerusalem, Israel

E-mail: michael.reiterer@protonmail.com

ABSTRACT: Yang-Mills gauge theory on Minkowski space supports a Batalin-Vilkovisky-infinity algebra structure, all whose operations are local. To make this work, the axioms for a BV-infinity algebra are deformed by a quadratic element, here the Minkowski wave operator. This homotopy structure implies BCJ/color-kinematics duality; a cobar construction yields a strict algebraic structure whose Feynman expansion for Yang-Mills tree amplitudes complies with the duality. It comes with a ‘syntactic kinematic algebra’.

Contents

1	Introduction	1
2	Application to BCJ	6
2.1	BCJ relations	7
2.2	BCJ/color-kinematics duality	9
2.3	Syntactic kinematic algebra	13
3	Definition of \mathfrak{a} and existence of h	14
4	Definition of BV_∞^Q-algebras	16
4.1	Recap of ordinary G_∞ - and BV_∞ -algebras	16
4.2	BV_∞^Q -algebras, with axioms deformed by a quadratic element Q	19
4.3	The auxiliary operator Γ	22
4.4	A special class	22
5	A vanishing theorem	25
6	BV_∞^\square-algebra for Yang-Mills	30
6.1	Proof of Theorem 3	30
6.2	Strict structure via a cobar construction	34
7	Acknowledgements	39
A	Addendum to Section 4.2	39
B	Some remarks about the BV formalism	41
C	Code	42

1 Introduction

We construct a Batalin-Vilkovisky-infinity algebra [23, 24] for Yang-Mills gauge theory on Minkowski spacetime. This implies the Bern-Carrasco-Johansson relations [1, 4] for tree amplitudes. Furthermore, a cobar construction yields a quasi-isomorphic strict algebra whose Feynman expansion complies with BCJ/color-kinematics duality [1–3]¹. Similar structures have been considered before in gauge theory, see Zeitlin’s [10–12]², but not a structure suitable for BCJ, for which one needs to deform the BV-infinity axioms.

¹This posits that YM tree amplitudes may be written as a sum of cubic trees where kinematic numerators satisfy ‘identities in one-to-one correspondence to the Jacobi identities obeyed by color factors’ [2].

²Zeitlin considers, among others, homotopy BV/Gerstenhaber algebras and generalized Yang-Mills equations associated to certain vertex operator algebras, via a quasi-classical limit of Lian-Zuckerman algebras.

One can formulate Yang-Mills in terms of a differential graded commutative algebra. To define this dgca [10, 13, 14], let $\Omega = \Omega(\mathbb{R}^4)$ be the smooth complex-valued differential forms on Minkowski space; d_{dR} the de Rham differential; $\Omega^2 = \Omega_+^2 \oplus \Omega_-^2$ the invariant decomposition³, and $\pi_+ : \Omega^2 \rightarrow \Omega_+^2$ the associated projection. Consider

$$\begin{array}{ccccccc}
 \Omega^0 & \xrightarrow{d_{\text{dR}}} & \Omega^1 & \xrightarrow{\pi_+ d_{\text{dR}}} & \Omega_+^2 & & \\
 & & & \searrow \mathbb{1} & & & \\
 & & \Omega_+^2 & \xrightarrow{d_{\text{dR}}} & \Omega^3 & \xrightarrow{d_{\text{dR}}} & \Omega^4
 \end{array} \tag{1.1}$$

Let $\mathfrak{a} = \mathfrak{a}^0 \oplus \mathfrak{a}^1 \oplus \mathfrak{a}^2 \oplus \mathfrak{a}^3$ be the four columns in (1.1) respectively, and $d : \mathfrak{a}^i \rightarrow \mathfrak{a}^{i+1}$ the differential given by all arrows in (1.1). The gca product $\mathfrak{a}^i \otimes \mathfrak{a}^j \rightarrow \mathfrak{a}^{i+j}$ is given by wedging forms: the product of an element in the upper row and one in the upper (lower) row is placed into the upper (lower) row; the product of two elements in the lower row is zero; see Section 3. Then \mathfrak{a} is a dgca: d is a differential; the gca product is associative and graded commutative; d is a derivation⁴. Given a finite-dimensional ‘color’ Lie algebra \mathfrak{c} , the classical Yang-Mills equation of motion is the Maurer-Cartan equation [19] in the differential graded Lie algebra $\mathfrak{a} \otimes \mathfrak{c}$ ⁵. The focus in this paper is on \mathfrak{a} which is the ‘kinematic’ part.

Denote by $h : \mathfrak{a}^i \rightarrow \mathfrak{a}^{i-1}$ any map⁶ of the form

$$\begin{array}{ccccccc}
 \Omega^0 & \xleftarrow{\quad} & \Omega^1 & \xleftarrow{\quad} & \Omega_+^2 & & \\
 & \nwarrow & & \nwarrow & & \nwarrow & \\
 & & \Omega_+^2 & \xleftarrow{\quad} & \Omega^3 & \xleftarrow{\quad} & \Omega^4
 \end{array} \tag{1.2}$$

subject to the following requirements: all arrows are invariant under translations on \mathbb{R}^4 ; the four horizontal arrows are homogeneous first order partial differential operators; the three diagonal arrows are $C^\infty(\mathbb{R}^4)$ -linear meaning they are without derivatives; and

$$\begin{aligned}
 h^2 &= 0 \\
 dh + hd &= \square \mathbb{1}
 \end{aligned} \tag{1.3}$$

where $\square = \eta^{\mu\nu} \partial_\mu \partial_\nu$ is the wave operator for the Minkowski metric η on \mathbb{R}^4 . Such an h exists but it is not unique; see Section 3. It need not be Lorentz invariant.

By definition, the failure of h to be second order on \mathfrak{a} is measured by^{7,8}

$$\begin{aligned}
 S(x, y, z) &= h(xyz) - h(xy)z - (-1)^x xh(yz) - (-1)^{(x+1)y} yh(xz) \\
 &\quad + h(x)yz + (-1)^x xh(y)z + (-1)^{x+y} xyh(z)
 \end{aligned} \tag{1.4}$$

³Decomposition into eigenspaces of the Hodge star operator for the Minkowski metric.

⁴Note that such a dgca \mathfrak{a} can be defined on any 4-manifold with a metric or conformal metric.

⁵The unknown in the Maurer-Cartan equation is an element of $\mathfrak{a}^1 \otimes \mathfrak{c} = (\Omega^1 \otimes \mathfrak{c}) \oplus (\Omega_+^2 \otimes \mathfrak{c})$, consisting of a \mathfrak{c} -valued 1-form, the vector potential; and the Ω_+^2 -half of a \mathfrak{c} -valued 2-form, the field strength.

⁶All maps are understood to be \mathbb{C} -linear.

⁷The order is in the graded sense, as in [30, Section 1], not the order as a partial differential operator.

⁸The $-h(1)xyz$ term is absent because $h(1) = 0$, since \mathfrak{a} is zero in degree -1 .

a graded symmetric map $S : \mathfrak{a}^{\otimes 3} \rightarrow \mathfrak{a}$. Here and in similar instances below, x, y, z, \dots are homogeneous elements of \mathfrak{a} , and juxtaposition denotes the gca product in \mathfrak{a} .

Theorem 1 (Second order up to homotopy). *Every such h is second order on \mathfrak{a} up to homotopy. Namely, there exists a map $\theta_3 : \mathfrak{a}^i \otimes \mathfrak{a}^j \otimes \mathfrak{a}^k \rightarrow \mathfrak{a}^{i+j+k-2}$ such that*

$$S(x, y, z) = d\theta_3(x, y, z) - \theta_3(dx, y, z) - (-1)^x \theta_3(x, dy, z) - (-1)^{x+y} \theta_3(x, y, dz) \quad (1.5)$$

The homotopy θ_3 is graded symmetric; it is $C^\infty(\mathbb{R}^4)$ -linear in each argument; it is invariant under translations on \mathbb{R}^4 ; and it satisfies the identities⁹

$$\begin{aligned} x\theta_3(y, u, v) - (-1)^{xy} y\theta_3(x, u, v) + \theta_3(x, yu, v) - (-1)^{xy} \theta_3(y, xu, v) &= 0 \\ \theta_3(xy, u, v) - \theta_3(x, yu, v) + \theta_3(x, y, uv) - (-1)^{x(y+u+v)} \theta_3(y, u, vx) &= 0 \end{aligned} \quad (1.6)$$

Corollary 2. *The BCJ relations in the form given in [4] hold.*

Theorem 1 suggests that the dgca operations together with h and θ_3 are but a fragment of a more comprehensive algebraic structure that subsumes (1.3), (1.5), (1.6) into its axioms. Specifically, \mathbf{BV}_∞ - and \mathbf{G}_∞ -algebras [23, 24] are used in deformation theory [21–23, 26], and Zeitlin [10–12] considered them in the context of generalized Yang-Mills. The \mathbf{BV}_∞ axioms require that $dh + hd$ be zero, which here fails due to \square in (1.3). This prompts one to define a deformed notion of a \mathbf{BV}_∞ -algebra, see Section 4.2, to be called a $\mathbf{BV}_\infty^\square$ -algebra¹⁰.

Theorem 3 (Main). *For every such h , there exists a $\mathbf{BV}_\infty^\square$ -algebra structure on \mathfrak{a} :*

- Its \mathbf{C}_∞ part is the dgca structure described above.
- Its unary degree -1 operator is h .
- All operations are invariant under translations and local¹¹.

In the notation of [24], all structure maps except $m_{1,\dots,1}^0$, $m_{1,\dots,1,2}^0$, m_1^1 vanish. The differential is m_1^0 ; the gca product is m_2^0 ; the homotopy θ_3 is $m_{1,2}^0$; and h is m_1^1 ; see (4.12).

The proof in Section 6.1 gives a definite choice and construction of all operations of this $\mathbf{BV}_\infty^\square$ -algebra; natural conditions based on the grading and the structure of \mathbb{R}^4 , cf. (6.4), fix all operations uniquely, given h ¹². This structure is a superstructure of the Yang-Mills dgca that also encompasses the ‘propagator numerator’ h .

⁹The first identity in (1.6) implies the second using graded symmetry (6.3).

¹⁰Given a graded vector space V that is also a module over a cocommutative Hopf algebra H , and a central quadratic element $Q \in H$, the dgLa controlling \mathbf{BV}_∞ -algebras is restricted to H -equivariant elements, then deformed by Q to a curved dgLa that, by definition, controls \mathbf{BV}_∞^Q -algebras. Every \mathbf{BV}_∞^Q -algebra has a differential and contains a \mathbf{C}_∞ -algebra, but not necessarily an \mathbf{L}_∞ -algebra. This notion is not specific to Yang-Mills. For Yang-Mills on Minkowski spacetime, set $V = \mathfrak{a}[2]$ and $H = \mathbb{C}[\partial_0, \dots, \partial_3]$ and $Q = \square$.

¹¹All operations belong to the classes of local operators in Definition 16; they are sums of translation-invariant $C^\infty(\mathbb{R}^4)$ -multilinear operations composed with at most one partial derivative.

¹²The construction does not reveal whether all but finitely many operations are identically zero. Computer calculations, using the code in Appendix C, show that there are nonzero operations of arity at least 6.

This has an application to BCJ/color-kinematics duality [1–3]. Let $\mathfrak{a}_w \hookrightarrow \mathfrak{a}$ be the subspace of finite linear combinations of plane waves¹³ with momenta in \mathbb{C}^4 . Then the L_∞ minimal model Feynman expansion for $\mathfrak{a}_w \otimes \mathfrak{c}$ using h as the propagator numerator gives the Yang-Mills amplitudes as a sum of cubic trees [14], but this does not yet comply with the duality. However, the upgrade to a BV_∞^\square -algebra structure in Theorem 3 does naturally lead to a Feynman expansion that complies with the duality¹⁴.

Namely, once Yang-Mills is cast as a BV_∞^\square -algebra, a generic cobar construction yields a quasi-isomorphic strict structure¹⁵, where the duality is more immediate. This construction in Section 6.2 is an adaptation of a standard one [24, 25, 33]. The next theorem states this result only as applied to \mathfrak{a} specifically. Denote by $H = \mathbb{C}[\partial_0, \dots, \partial_3]$ the Hopf algebra of constant coefficient differential operators on \mathbb{R}^4 , and note that \mathfrak{a} is an H -module.

Theorem 4 (Strict structure via a cobar construction). *Let A be the free Gerstenhaber algebra of the cofree Gerstenhaber coalgebra of \mathfrak{a} ¹⁶. Then a cobar construction applied to the BV_∞^\square -algebra \mathfrak{a} yields H -equivariant $d_A \in \text{End}^1(A)$ and $h_A \in \text{End}^{-1}(A)$ such that:*

- (A, d_A) is a dgca; (A, h_A) is a BV algebra; so h_A is genuinely second order; and

$$d_A^2 = 0 \quad d_A h_A + h_A d_A = \square \mathbb{1} \quad h_A^2 = 0 \quad (1.7)$$

where \square is in the sense of the induced H -module structure on A .

- The complexes (\mathfrak{a}, d) and (A, d_A) are quasi-isomorphic. More concretely, the canonical degree zero map $I : \mathfrak{a} \hookrightarrow A$ is part of an H -equivariant contraction¹⁷:

$$\begin{array}{ccc} \mathfrak{a} & \xrightleftharpoons[I]{I} & A \end{array} \quad \begin{array}{c} \hookrightarrow \\ \hookleftarrow \end{array} Z \quad (1.8)$$

The maps I and P are actually chain maps in two ways:

$$d_A I = I d \quad d P = P d_A \quad h_A I = I h \quad h P = P h_A \quad (1.9)$$

- Denoting by $\Theta : A \otimes A \rightarrow A$ and $\theta_2 : \mathfrak{a} \otimes \mathfrak{a} \rightarrow \mathfrak{a}$ the respective gca products:

$$P \circ \Theta \circ I^{\otimes 2} = \theta_2 \quad (1.10)$$

The cobar construction is functorial. Applied to $\mathfrak{a}_w \hookrightarrow \mathfrak{a}$ it yields a structure $A_w \hookrightarrow A$; both \mathfrak{a}_w and A_w are \mathbb{C}^4 -graded by momentum $k \simeq \partial$; H -equivariance implies momentum conservation. Abusing notation, denote by I also its restriction to A_w etc.

¹³By definition, an element $y \in \mathfrak{a}$ is a plane wave with momentum $k \in \mathbb{C}^4$ if and only if $\tau_a y = e^{c \cdot k_\mu a^\mu} y$ for all $a \in \mathbb{R}^4$, where τ_a denotes translation by a . The constant $c \neq 0$ is a matter of convention.

¹⁴With ‘local kinematic numerators’.

¹⁵Strict means only unary and binary operations; \mathfrak{a} is not strict as a BV_∞^\square -algebra because $\theta_3 \neq 0$.

¹⁶See Lemma 17 for details, including degree shifts; set $V = \mathfrak{a}[2]$ there, equivalently $V[-2] = \mathfrak{a}$.

¹⁷See Lemma 17 for details, including the equations defining a contraction.

Corollary 5 (BCJ duality). *Use \rightharpoonup to indicate partial maps defined for a dense subset of momenta. Consider the differential graded Lie algebra $A_w \otimes \mathfrak{c}$, with homology $H(\mathfrak{a}_w \otimes \mathfrak{c})$. The standard L_∞ minimal model Feynman expansion gives for every $n \geq 2$ a map*

$$M_n : H^1(\mathfrak{a}_w \otimes \mathfrak{c})^{\otimes n} \rightharpoonup H^2(\mathfrak{a}_w \otimes \mathfrak{c}) \quad (1.11)$$

as a sum over all non-planar cubic trees with one output and inputs $1 \dots n$ using:

tree element	operation	
vertex	gca product in A_w , Lie bracket in \mathfrak{c}	$(A_w^1 \otimes \mathfrak{c})^{\otimes 2} \rightarrow A_w^2 \otimes \mathfrak{c}$
internal line	$h_A/k^2 \otimes \mathbb{1}$	$A_w^2 \otimes \mathfrak{c} \rightharpoonup A_w^1 \otimes \mathfrak{c}$
input	$i \otimes \mathbb{1}$	$H^1(\mathfrak{a}_w \otimes \mathfrak{c}) \rightharpoonup A_w^1 \otimes \mathfrak{c}$
the output	$pP \otimes \mathbb{1}$	$A_w^2 \otimes \mathfrak{c} \rightharpoonup H^2(\mathfrak{a}_w \otimes \mathfrak{c})$

where $k^2 \simeq \square$ is the momentum squared; $i : H(\mathfrak{a}_w) \not\rightarrow \mathfrak{a}_w : p$ are any momentum-conserving quasi-isomorphisms that are nonzero only along $k^2 = 0$, undefined at $k = 0$. This expansion complies with BCJ/color-kinematics duality¹⁸; and the M_n are the Yang-Mills amplitudes.

The definition of the duality and the proof are in Section 2.2.

It has been suggested that a rationale for the duality would be provided by a hypothetical ‘kinematic Lie algebra’, see [6–9] for results in this direction. Theorem 4 provides such a Lie algebra, namely the Lie bracket

$$[-, -] : A_w^1 \otimes A_w^1 \rightarrow A_w^1 \quad (1.12)$$

Indeed, the Yang-Mills amplitudes are also obtained using these Feynman rules:

tree element	operation	
vertex	(1.12) in A_w^1 , Lie bracket in \mathfrak{c}	$(A_w^1 \otimes \mathfrak{c})^{\otimes 2} \rightarrow A_w^1 \otimes \mathfrak{c}$
internal line	$1/k^2 \otimes \mathbb{1}$	$A_w^1 \otimes \mathfrak{c} \rightharpoonup A_w^1 \otimes \mathfrak{c}$
input	$i \otimes \mathbb{1}$	$H^1(\mathfrak{a}_w \otimes \mathfrak{c}) \rightharpoonup A_w^1 \otimes \mathfrak{c}$
the output	$pP \otimes \mathbb{1}$	$A_w^1 \otimes \mathfrak{c} \rightharpoonup H^1(\mathfrak{a}_w \otimes \mathfrak{c})$

There is a paradox here. The bracket (1.12) in the free Gerstenhaber algebra is a syntactic construction, given by sticking together words modulo some relations, see (4.1). How can this give the Yang-Mills amplitudes? The resolution lies in P which translates the syntactic expression in A_w^1 back to \mathfrak{a}_w^1 , in a nontrivial way. The maps $H^1(\mathfrak{a}_w \otimes \mathfrak{c})^{\otimes n} \rightharpoonup H^1(\mathfrak{a}_w \otimes \mathfrak{c})$ so obtained are equivalent to the M_n , as discussed in Section 2.3. The Jacobi identity for the ‘syntactic kinematic algebra’ (1.12) implies BCJ duality.

The perspective on BCJ duality provided by Theorems 3 and 4 is that once \mathfrak{a} is up-graded to a BV_∞^\square -algebra, then the duality arises by rectifying this homotopy structure.

Beware that Theorem 4 does not actually say that A is a rectification of \mathfrak{a} in the sense of an ∞ -quasi-isomorphism, though this is very likely true, but (1.8), (1.9), (1.10) are partial

¹⁸With ‘local kinematic numerators’.

statements that suffice to show that one gets the Yang-Mills amplitudes, using the recursive characterization and tools from [14], see Section 2.2.

Some questions:

- Instead of the \mathbf{C}_∞ minimal model for \mathfrak{a}_w , one could consider its $\mathbf{BV}_\infty^\square$ minimal model. Assuming this can be defined, is it an interesting object?¹⁹
- A direct approach to the ‘double copy’ could be to consider $\mathfrak{a} \otimes \mathfrak{a}$ understood as a suitably defined tensor product of $\mathbf{BV}_\infty^\square$ -algebras.^{20,21}
- Are there invariant bilinear pairings that give a ‘cyclic’ $\mathbf{BV}_\infty^\square$ -algebra, for loops?
- What are other applications and ramifications of this $\mathbf{BV}_\infty^\square$ -algebra structure?

2 Application to BCJ

This section is intended as motivation for readers familiar with BCJ [1–3]. It includes proof sketches of Corollaries 2 and 5. This section uses results shown in detail later.

Plane waves and propagators. This is the only section where we work with plane waves. Assume an h as in Section 1 is fixed.

- Let $\mathfrak{a}_w \hookrightarrow \mathfrak{a}$ be the subalgebra given by finite linear combinations of plane waves with complex momentum denoted $k \in \mathbb{C}^4$. This defines a \mathbb{C}^4 -grading on \mathfrak{a}_w , conserved by all operations. Then h is, relative to standard trivializations, a matrix with entries that are affine linear functions of $k \simeq \partial$. Abbreviate $k^2 = \eta^{\mu\nu} k_\mu k_\nu \simeq \square$. Set

$$h_\sharp = h/k^2 \tag{2.1}$$

This ‘propagator’ is ill-defined when acting on plane waves with $k^2 = 0$; so we write $h_\sharp : \mathfrak{a}_w \rightharpoonup \mathfrak{a}_w$ to indicate that this is a partial map. The arrow \rightharpoonup is contagious, but all partial maps we encounter are defined on dense subsets of momenta. In every occurrence of h_\sharp it is implicit that one must have $k^2 \neq 0$. With this proviso,

$$dh_\sharp + h_\sharp d = \mathbb{1} \qquad h_\sharp^2 = h_\sharp h = h h_\sharp = 0 \tag{2.2}$$

by (1.3). The homology²² $H(\mathfrak{a}_w)$ is a direct sum of the fibers of a non-trivial vector bundle over the complex cone $k^2 = 0$ in \mathbb{C}^4 , assuming $k \neq 0$, see [14] and cf. Lemma 11. The homology at $k = 0$ is never used.

¹⁹At the very least, this conjectural $\mathbf{BV}_\infty^\square$ minimal model ought to subsume the BCJ relations into its axioms – rather than implying them by an ad-hoc calculation as in Section 2.1 – analogously to how the \mathbf{C}_∞ minimal model subsumes the Kleiss-Kuijf relations into its axioms; see [20] for the \mathbf{C}_∞ axioms.

²⁰Direct approach means as opposed to going through BCJ duality. The (completed) tensor product $\mathfrak{a} \otimes \mathfrak{a}$ naively gives objects on $\mathbb{R}^4 \oplus \mathbb{R}^4 \ni x \oplus y$, but can be restricted to things depending only on $x + y$, or analogously, products of pairs of plane waves with equal momentum.

²¹See also [12, Section 3] for considerations in this ballpark.

²²If h_\sharp was a true element of $\text{End}^{-1}(\mathfrak{a}_w)$ then (2.2) would erroneously imply that $H(\mathfrak{a}_w)$ vanishes.

- Analogous notation is used for the cobar construction $A_w \hookrightarrow A$ of $\mathfrak{a}_w \hookrightarrow \mathfrak{a}$; it is spanned by Gerstenhaber words of co-Gerstenhaber words (cf. [26]) in plane waves in \mathfrak{a}_w . The momentum $k \in \mathbb{C}^4$ of such a word-of-words is the total momentum of all plane waves in it; this defines a \mathbb{C}^4 -grading on A_w . Define $h_{A\sharp} : A_w \rightarrow A_w$ by

$$h_{A\sharp} = h_A/k^2 \quad (2.3)$$

It follows from (1.7) that

$$d_A h_{A\sharp} + h_{A\sharp} d_A = \mathbb{1} \quad h_{A\sharp}^2 = h_{A\sharp} h_A = h_A h_{A\sharp} = 0 \quad (2.4)$$

Feynman expansion. Given a contraction, aka deformation retract, from \mathfrak{a}_w (or A_w) to its homology $H(\mathfrak{a}_w) \simeq H(A_w)$, there are standard formulas for...

- ...the \mathbf{C}_∞ minimal model of the dgca \mathfrak{a}_w (or A_w) using planar cubic trees, defining minimal model operations $H(\mathfrak{a}_w)^{\otimes n} \rightarrow H(\mathfrak{a}_w)$, see [20] or [33, Section 13.1.9].
- ...the \mathbf{L}_∞ minimal model of the dgLa $\mathfrak{a}_w \otimes \mathfrak{c}$ (or $A_w \otimes \mathfrak{c}$) using cubic trees, defining minimal model operations $H(\mathfrak{a}_w \otimes \mathfrak{c})^{\otimes n} \rightarrow H(\mathfrak{a}_w \otimes \mathfrak{c})$, see [33, Section 10.3.4] or [14].

Let us call them Feynman-Kontsevich-Soibelman trees. These trees have one output and $n \geq 2$ inputs labeled $1 \dots n$. The rules for \mathbf{C}_∞ trees are summarized in the next table:

tree element	operation, using \mathfrak{a}_w		operation, using A_w	
vertex	gca product	$\mathfrak{a}_w^i \otimes \mathfrak{a}_w^j \rightarrow \mathfrak{a}_w^{i+j}$	gca product	$A_w^i \otimes A_w^j \rightarrow A_w^{i+j}$
internal line	h_\sharp	$\mathfrak{a}_w^i \rightarrow \mathfrak{a}_w^{i-1}$	$h_{A\sharp}$	$A_w^i \rightarrow A_w^{i-1}$
input	i	$H^i(\mathfrak{a}_w) \rightarrow \mathfrak{a}_w^i$	$\textcolor{blue}{I}i$	$H^i(\mathfrak{a}_w) \rightarrow A_w^i$
the output	p	$\mathfrak{a}_w^i \rightarrow H^i(\mathfrak{a}_w)$	$p\textcolor{blue}{P}$	$A_w^i \rightarrow H^i(\mathfrak{a}_w)$

and they involve signs. For \mathbf{L}_∞ one must suitably tensor with \mathfrak{c} , the Lie bracket $\mathfrak{c} \otimes \mathfrak{c} \rightarrow \mathfrak{c}$, and $\mathbb{1} : \mathfrak{c} \rightarrow \mathfrak{c}$. Canonically, $H(\mathfrak{a}_w \otimes \mathfrak{c}) \simeq H(\mathfrak{a}_w) \otimes \mathfrak{c}$. The quasi-isomorphisms

$$i : H(\mathfrak{a}_w) \not\rightarrow \mathfrak{a}_w : p \quad (2.5)$$

can be and are chosen to conserve the \mathbb{C}^4 -grading; they are nonzero only along $k^2 = 0$; undefined only at $k = 0$; otherwise arbitrary. By $\textcolor{blue}{H}$ -equivariance, $\textcolor{blue}{I} : \mathfrak{a}_w \not\rightarrow A_w : p\textcolor{blue}{P}$ in Theorem 4 conserve the \mathbb{C}^4 -grading. The minimal model operations do not depend on the contraction – this includes i and p and the propagator – for a proof in the \mathbf{L}_∞ case see [14].

Working with the operations on \mathfrak{a}_w , the Yang-Mills amplitudes correspond to the \mathbf{C}_∞ operation $H^1(\mathfrak{a}_w)^{\otimes n} \rightarrow H^2(\mathfrak{a}_w)$, or, including the color Lie algebra \mathfrak{c} , to the \mathbf{L}_∞ operation $H^1(\mathfrak{a}_w \otimes \mathfrak{c})^{\otimes n} \rightarrow H^2(\mathfrak{a}_w \otimes \mathfrak{c})$ [14]. Using A_w gives the same result, see Section 2.2.

2.1 BCJ relations

The following lemma is a direct consequence of the definition of S in (1.4), together with the dgca axioms and (1.3). One does not need Theorem 1 to prove this.

Lemma 6. *The map S has degree -1 , is graded symmetric²³, and*

$$dS(x, y, z) = -S(dx, y, z) - (-1)^x S(x, dy, z) - (-1)^{x+y} S(x, y, dz) \quad (2.6)$$

and

$$\begin{aligned} (-1)^x x S(y, u, v) - (-1)^{xy+y} y S(x, u, v) + S(x, yu, v) - (-1)^{xy} S(y, xu, v) &= 0 \\ S(xy, u, v) - S(x, yu, v) + S(x, y, uv) - (-1)^{x(y+u+v)} S(y, u, vx) &= 0 \end{aligned} \quad (2.7)$$

Proof. For (2.6) also use $\square(f_1 f_2 f_3) - \square(f_1 f_2) f_3 - \square(f_2 f_3) f_1 - \square(f_3 f_1) f_2 + \square(f_1) f_2 f_3 + \square(f_2) f_3 f_1 + \square(f_3) f_1 f_2 = 0$ for all $f_1, f_2, f_3 \in C^\infty(\mathbb{R}^4)$. \square

In this section we show the BCJ relations given by Tye and Zhang [4, equation (4.32)], that is, Corollary 2. For every $n \geq 3$, the argument is in two steps:

- *Step 1.* First we define a map $S_n : \mathfrak{a}_w^{\otimes n} \rightarrow \mathfrak{a}_w$, which is a certain signed sum of cubic trees with n inputs and one output, where each of the $n-1$ vertices is a gca product; and internal/input/output lines are decorated by a total of $n-3$ propagators $h_\#$ and a single $h = k^2 h_\#$. Lemma 6 is then used to show that S_n is a chain map:

$$dS_n(x_1, \dots, x_n) = \sum_{i=1}^n (-1)^{n+x_1+\dots+x_{i-1}} S_n(x_1, \dots, dx_i, \dots, x_n)$$

In particular, S_n induces a map on homology, $H(\mathfrak{a}_w)^{\otimes n} \rightarrow H(\mathfrak{a}_w)$.

- *Step 2.* Using Theorem 1 we then show that the induced $H(\mathfrak{a}_w)^{\otimes n} \rightarrow H(\mathfrak{a}_w)$ is zero. Namely, θ_3 is used to construct a $T_n : \mathfrak{a}_w^{\otimes n} \rightarrow \mathfrak{a}_w$ such that

$$S_n(x_1, \dots, x_n) = dT_n(x_1, \dots, x_n) + \sum_{i=1}^n (-1)^{n+x_1+\dots+x_{i-1}} T_n(x_1, \dots, dx_i, \dots, x_n)$$

The vanishing of the map $H^1(\mathfrak{a}_w)^{\otimes n} \rightarrow H^2(\mathfrak{a}_w)$ induced by S_n is [4, equation (4.32)].

It is useful to first discuss $n=3$, then $n=4$, then sketch the general n case.

- *Case $n=3$.* Set $S_3 = S$. Step 1 follows from (2.6). Step 2 follows from (1.5) using $T_3 = \theta_3$. This case goes back to an observation of Zhu [5].
- *Case $n=4$.* Set $E(x, y) = h_\#(xy)$ and set

$$\begin{aligned} S_4(x, y, u, v) &= S(E(x, y), u, v) - (-1)^x S(x, E(y, u), v) \\ &\quad + (-1)^{x+y} S(x, y, E(u, v)) - (-1)^{y+u+x(y+u+v)} S(y, u, E(v, x)) \end{aligned}$$

Note $dE(x, y) = xy - E(dx, y) - (-1)^x E(x, dy)$ by (2.2), with an ‘error term’ xy . Step 1 follows from this, from (2.6), and from the second of (2.7) to cancel ‘error terms’. Step 2 follows from (1.5) and the second of (1.6) using the witness

$$\begin{aligned} T_4(x, y, u, v) &= \theta_3(E(x, y), u, v) - (-1)^x \theta_3(x, E(y, u), v) \\ &\quad + (-1)^{x+y} \theta_3(x, y, E(u, v)) - (-1)^{y+u+x(y+u+v)} \theta_3(y, u, E(v, x)) \end{aligned}$$

²³Explicitly, $S(x, y, z) = (-1)^{xy} S(y, x, z) = (-1)^{yz} S(x, z, y)$.

- *General $n \geq 3$, sketch.* Set $E(x_1) = x_1$. For $n \geq 2$ let $E(x_1, \dots, x_n)$ be the sum, with suitable signs, of all planar cubic trees with n inputs and one output²⁴, where each vertex is a gca product; each internal line and the output line is decorated by h_\sharp ; the input lines are not decorated. Schematically set

$$S_n(x_1, \dots, x_n) = \sum \pm S(E(x_i, x_{i+1}, \dots, x_{j-1}), E(x_j, x_{j+1}, \dots, x_{k-1}), E(x_k, x_{k+1}, \dots, x_{i-1})) \quad (2.8)$$

where the subscripts are in $\mathbb{Z}/n\mathbb{Z}$ and the sum is over all decompositions of $\mathbb{Z}/n\mathbb{Z}$ into three nonempty contiguous subsets²⁵. Step 1 and Step 2 follow like before²⁶.

The translation to [4] uses the following remarks:

- Take the definition of S_n , (2.8), and substitute the definitions of S , (1.4), and of E . One obtains cubic trees as described in Step 1. Since $hh_\sharp = h_\sharp h = 0$, (2.2), a tree is zero unless all $n - 3$ propagators h_\sharp and the single h appear on *distinct* lines.
- There are some trees where h decorates an input or output line, but they do not contribute to $H^1(\mathfrak{a}_w)^{\otimes n} \rightarrow H^2(\mathfrak{a}_w)$ by (3.4); these degrees give the YM amplitudes. So only trees where all h_\sharp and h appear on *distinct internal* lines contribute to this.
- Replace the single h by $h = k^2 h_\sharp$, and regard k^2 (the square of the momentum flowing through an internal line) as a weight multiplying a cubic tree that uses h_\sharp on all internal lines, that is, a standard \mathbf{C}_∞ Feynman-Kontsevich-Soibelman tree.

The discussion in this section is ad-hoc. Presumably, if one can define the $\mathbf{BV}_\infty^\square$ minimal model of \mathfrak{a}_w , the BCJ relations will be subsumed into the $\mathbf{BV}_\infty^\square$ axioms.

2.2 BCJ/color-kinematics duality

Lemma 7. *If a functional $\ell \in A^*$ satisfies $\ell h_A = 0$, then for all $x, y, z \in \ker h_A$:*

$$\ell(h_A(xy)z) + (-1)^{x(y+z)}\ell(h_A(yz)x) + (-1)^{z(x+y)}\ell(h_A(zx)y) = 0 \quad (2.9)$$

where juxtaposition denotes the gca product in A . Analogous for $A_w \hookrightarrow A$.

Proof. Since (A, h_A) is a BV algebra by Theorem 4 (cf. Lemma 17), h_A is genuinely second order, so the right hand side of (1.4), with h_A in the role of h and all free variables in A , vanishes. Using graded symmetry of the gca product in A , the claim follows. \square

To prove Corollary 5, consider for every $n \geq 2$ the n -ary operation of the \mathbf{L}_∞ minimal model, but specialized to the degrees of interest, see (1.11). Concretely, M_n is a sum of Feynman-Kontsevich-Soibelman trees with propagator (2.3):

$$M_n = \sum_{T \in \mathcal{T}_n} M_{n,T'} \otimes c_{T'} \quad (2.10)$$

²⁴There are C_{n-1} such trees, with C the Catalan numbers.

²⁵The number of ways of decomposing $\mathbb{Z}/n\mathbb{Z}$ into m nonempty contiguous subsets is $\binom{n}{m}$.

²⁶In this entire section, only the second identity in (2.7) and (1.6) respectively is used; there should be an analogous discussion using the first in (2.7) and (1.6) respectively.




Figure 1: The graph \mathcal{T}_4 . Each vertex is a cubic tree with one output and four inputs $1, \dots, 4$. There are many 3-cliques; an example is given by the three black vertices. In \mathcal{T}_n , every vertex belongs to exactly $n - 2$ many 3-cliques, and there are no larger cliques.

Here \mathcal{T}_n is the set of all (non-planar) cubic trees with one output and n inputs labeled $1 \dots n$, so that $|\mathcal{T}_n| = (2n - 3)!!$, and T' denotes any planar embedding of T . Here $c_{T'} : \mathfrak{c}^{\otimes n} \rightarrow \mathfrak{c}$ nests $n - 1$ instances of the Lie bracket as dictated by the planar embedding T' , and

$$M_{n,T'} : H^1(\mathfrak{a}_w)^{\otimes n} \rightharpoonup H^2(\mathfrak{a}_w)$$

is defined by decorating T' as follows:

tree element	operation
vertex	gca product $A_w^1 \otimes A_w^1 \rightarrow A_w^2$
internal line	$h_{A\#} = h_A/k^2$ $A_w^2 \rightharpoonup A_w^1$
input	Ii $H^1(\mathfrak{a}_w) \rightharpoonup A_w^1$
the output	pP $A_w^2 \rightharpoonup H^2(\mathfrak{a}_w)$

e.g. $M_{3,(1(23))}(x, y, z) = pP((Iix)h_{A\#}((Iiy)(Iiz)))$. So (2.10) does not depend on the choice of the T' since the Lie bracket on \mathfrak{c} and the product $A_w^1 \otimes A_w^1 \rightarrow A_w^2$ are antisymmetric.

One can view \mathcal{T}_n as an undirected graph where an edge corresponds to²⁷ ‘one associativity move’, see Figure 1. By definition, (2.10) satisfies BCJ/color-kinematics duality [1–3] if and only if for every 3-clique²⁸ $U, V, W \in \mathcal{T}_n$ one has

$$k_U^2 M_{n,U'} + k_V^2 M_{n,V'} + k_W^2 M_{n,W'} = 0 \quad (2.11)$$

where k_U is the momentum flowing through the distinguished line in U ²⁹, and the planar embeddings U', V', W' are chosen compatibly in the sense that the four subtrees connecting to the distinguished lines, which appear jointly in U and V and W , are embedded in the same way as part of U' and V' and W' , and the relative planar embedding of the four subtrees is cyclic, in particular so that $c_{U'} + c_{V'} + c_{W'} = 0$ by the Jacobi identity in \mathfrak{c} .

²⁷Cf. the associahedron, aka Stasheff polytope, which is for planar cubic trees.

²⁸Given a graph, an n -clique is a subset of n vertices that are all pairwise connected by an edge.

²⁹The distinguished line in U is the one not present in V and W .

BCJ duality (2.11) for the expansion (2.10) follows from (2.9), by letting x, y, z be the outputs of three of the four subtrees, and letting ℓ be the input to the fourth. The degrees work out to be $x, y, z \in A_w^1$ and $\ell \in (A_w^2)^*$ so that the signs in (2.9) disappear.

- Proof of $x \in \ker h_A$: If x is not an overall input, then $x = h_A \xi$ for some $\xi \in A_w^2$, so $h_A x = (h_A)^2 \xi = 0$ by (1.7). If x is an overall input, then $x = I i \xi$ for some $\xi \in H^1(\mathfrak{a}_w)$, so $h_A x = h_A I i \xi = I h i \xi = 0$ by (1.9) and (3.4). Analogously $y, z \in \ker h_A$.
- Proof of $\ell h_A = 0$: If ℓ is not the overall output, then $\ell = \nu h_A$ for some $\nu \in (A_w^1)^*$, so $\ell h_A = \nu (h_A)^2 = 0$ by (1.7). If ℓ is the overall output, then $\ell = \nu p P$ for some $\nu \in H^2(\mathfrak{a}_w)^*$, so $\ell h_A = \nu p P h_A = \nu p h P = 0$ by (1.9) and (3.4).

To prove Corollary 5, we must also check that M_n are the ordinary YM amplitudes³⁰. We exploit the fact that the YM amplitudes can be recursively characterized in terms of the ‘factorization of their residues’. This is known from Britto-Cachazo-Feng-Witten recursion. We use the characterization of A. Nützi and the author in [14] that is simpler in that it does not involve any BCFW shift nor limiting conditions ‘at infinity’ under such shifts³¹.

Let k_1, \dots, k_n be the input momenta viewed as standard coordinates on $(\mathbb{C}^4)^n \simeq \mathbb{C}^{4n}$, and abbreviate $k_{n+1} = k_1 + \dots + k_n$ which is the output momentum. The argument that the M_n are the ordinary Yang-Mills amplitudes goes as follows (sketch):

- Recall (1.11); recall that M_n is undefined if (2.3) is undefined meaning $k_J^2 = 0$ for some $J \subseteq \{1, \dots, n\}$ with $1 < |J| < n$, where $k_J = \sum_{j \in J} k_j$; recall that the one-particle homology is a sheaf over the cone in \mathbb{C}^4 minus the origin, see [14] and cf. Lemma 11. So, geometrically, M_n is a section of an algebraic sheaf along $V_n - P_n$ where

$$V_n = \{k_1^2 = \dots = k_{n+1}^2 = 0\} \quad P_n = \bigcup_{j=1 \dots n+1} \{k_j = 0\} \cup \bigcup_{1 < |J| < n} \{k_J^2 = 0\}$$

Here P_n is a closed subset of the variety V_n , of codimension one if $n > 2$.

- M_n is homogeneous jointly in k_1, \dots, k_n with the natural degree of homogeneity, because all constructions in this paper, including A_w , scale naturally³². Thus M_n is, equivalently, a section of a sheaf on the projective variety associated to $V_n - P_n$.
- M_2 coincides with the map $H^1(\mathfrak{a}_w \otimes \mathfrak{c})^{\otimes 2} \rightarrow H^2(\mathfrak{a}_w \otimes \mathfrak{c})$ that is induced by the ga product $\mathfrak{a}_w^1 \otimes \mathfrak{a}_w^1 \rightarrow \mathfrak{a}_w^2$ times the Lie bracket $\mathfrak{c} \otimes \mathfrak{c} \rightarrow \mathfrak{c}$, because of (1.10), hence it is Lorentz invariant and nonzero on both irreducible components of $V_2 - P_2$.

Hereafter, $n > 2$. Then V_n is irreducible [14]. Decompose $\bigcup_{1 < |J| < n} \{k_J^2 = 0\} \subseteq V_n$ into its irreducible components denoted $V(\mathfrak{p}) \subseteq V_n$, where \mathfrak{p} is the corresponding prime divisor in the coordinate ring of V_n . A classification and properties of the \mathfrak{p} is in [14]; beware that one $V(\mathfrak{p})$ can be a component of several $\{k_J^2 = 0\}$; this happens but only for $n = 3$.

³⁰E.g., why does the Feynman expansion for $\mathfrak{a}_w \otimes \mathfrak{c}$, which fails BCJ duality, give the same result?

³¹These points ‘at infinity’ are irrelevant for the recursive characterization, by a Hartogs extension theorem, because they are contained in (projectively) a codimension two subset.

³²Since h scales naturally by the requirements in Section 1, all operations do.

- M_n has only simple poles along each $V(\mathfrak{p})$. More precisely, M_n extends, as a section of a sheaf that allows only these specific simple poles [14], to the bigger open set $V_n - Z_n$, where the bad set $Z_n = \bigcup_{\mathfrak{p} \neq \mathfrak{q}} V(\mathfrak{p}) \cap V(\mathfrak{q})$ has codimension two.
- The recursive characterization requires showing that $\text{Res}_{\mathfrak{p}} M_n$, the residue along $V(\mathfrak{p})$, equals some bilinear expression in M_2, \dots, M_{n-1} for every \mathfrak{p} , see [14]³³. This is not immediate when using the propagator (2.3). But one can use gauge independence in [14] to replace the relevant instances of (2.3) by another propagator – a composition of an optimal homotopy as in [14] with the contraction (1.8) – suitable to evaluate $\text{Res}_{\mathfrak{p}} M_n$ and to check that it is of the required form. Here are more details:
 - Let $\mathfrak{a}_w = \bigoplus_{k \in \mathbb{C}^4} \mathfrak{a}_{w,k}$ and $A_w = \bigoplus_{k \in \mathbb{C}^4} A_{w,k}$ be the \mathbb{C}^4 -gradings by momentum. Then $\mathfrak{a}_{w,k}$ is finite-dimensional, but $A_{w,k}$ is infinite-dimensional since it is the span of all words-of-words in plane waves in \mathfrak{a}_w with total momentum k .
 - By H -equivariance, $d(\mathfrak{a}_{w,k}) \subseteq \mathfrak{a}_{w,k}$ and $d_A(A_{w,k}) \subseteq A_{w,k}$ and the binary operations on \mathfrak{a}_w and A_w conserve the \mathbb{C}^4 -grading. Hence the gauge independence theorem as stated in [14] applies not only to $\mathfrak{a}_w \otimes \mathfrak{c}$ but also to $A_w \otimes \mathfrak{c}$.
 - Suppose we want to evaluate $\text{Res}_{\mathfrak{p}} M_n$ at a point

$$(k_{1\star}, \dots, k_{n\star}) \in V(\mathfrak{p}) - Z_n \quad (2.12)$$

For every J with $1 < |J| < n$ such that $V(\mathfrak{p}) \subseteq \{k_J^2 = 0\}$ (if $n > 3$ then J is unique) set $q_{\star} = k_{J\star} = \sum_{j \in J} k_{j\star}$, so $q_{\star} \in \mathbb{C}^4 - 0$ and $q_{\star}^2 = 0$. Denote by

$$i_{\star} : H(\mathfrak{a}_{w,q_{\star}}) \xrightarrow{\varphi} \mathfrak{a}_{w,q_{\star}} : p_{\star} \quad I_{\star} : \mathfrak{a}_{w,q_{\star}} \xrightarrow{\varphi} A_{w,q_{\star}} : P_{\star}$$

the restrictions of (2.5) respectively (1.8) to q_{\star} ; and by $\chi_{\star} : H(\mathfrak{a}_{w,q_{\star}}) \rightarrow H(\mathfrak{a}_{w,q_{\star}})$ the canonical degree -1 map whose only component is (3.5).

- An optimal homotopy [14] for \mathfrak{a}_w based at q_{\star} is a degree -1 map $h^{\text{opt}} : \mathfrak{a}_w \rightarrow \mathfrak{a}_w$ with these properties: it is given by a matrix whose entries are rational functions of k so that $k^2 h^{\text{opt}}$ is regular at q_{\star} ; it satisfies $dh^{\text{opt}} + h^{\text{opt}}d = \mathbb{1}$ and $(h^{\text{opt}})^2 = 0$ and $\lim_{k \rightarrow q_{\star}} k^2 h^{\text{opt}} = i_{\star} \chi_{\star} p_{\star}$. This is constructed in [14] by nesting two instances of the homological perturbation lemma. Set³⁴

$$h_A^{\text{opt}} = Z + I h^{\text{opt}} P : A_w \rightarrow A_w$$

It has these properties: $k^2 h_A^{\text{opt}}$ is regular at q_{\star} and (to check this one also uses the equations satisfied by the contraction (1.8), spelled out in Lemma 17)³⁵:

$$d_A h_A^{\text{opt}} + h_A^{\text{opt}} d_A = \mathbb{1} \quad (h_A^{\text{opt}})^2 = 0 \quad \lim_{k \rightarrow q_{\star}} k^2 h_A^{\text{opt}} = I_{\star} i_{\star} \chi_{\star} p_{\star} P_{\star} \quad (2.13)$$

³³This is known as ‘factorization of residues’.

³⁴The maps I, P, Z from (1.8) are unconditionally, as opposed to partially, defined.

³⁵A precise definition of the limit is omitted. One possibility is this: Take a word-of-words of $m \geq 1$ plane waves viewed as depending parametrically on the m momenta; then replace them by m sequences of momenta so that the m sequences converge separately and so that the total momentum converges to q_{\star} . Evaluating $k^2 h_A^{\text{opt}}$ along such a sequence gives the claimed result.

- For all $T \in \mathcal{T}_n$, consider the line – if there is one – with momentum k_J flowing through, and replace the corresponding $h_{A\sharp}$ by h_A^{opt} in the definition of $M_{n,T'}$. The first two in (2.13) mean that h_A^{opt} is an admissible propagator, hence this replacement does not change M_n , by gauge independence [14]. The last in (2.13) implies that $\text{Res}_p M_n$ is, at the point (2.12), of the required form.

Hence the M_n are the Yang-Mills amplitudes, by the recursive characterization in [14].

2.3 Syntactic kinematic algebra

A slightly different Feynman expansion, using (1.12) at each vertex, also produces the Yang-Mills amplitudes M_n . Namely, for every $n \geq 2$ set

$$\widetilde{M}_n = \sum_{T \in \mathcal{T}_n} \widetilde{M}_{n,T'} \otimes c_{T'} \quad : \quad H^1(\mathfrak{a}_w \otimes \mathfrak{c})^{\otimes n} \rightarrow H^1(\mathfrak{a}_w \otimes \mathfrak{c})$$

where $\widetilde{M}_{n,T'} : H^1(\mathfrak{a}_w)^{\otimes n} \rightarrow H^1(\mathfrak{a}_w)$ is defined by decorating T' as follows:

tree element	operation	
vertex	gLa product	$A_w^1 \otimes A_w^1 \rightarrow A_w^1$
internal line	$1/k^2$	$A_w^1 \rightarrow A_w^1$
input	Ii	$H^1(\mathfrak{a}_w) \rightarrow A_w^1$
the output	pP	$A_w^1 \rightarrow H^1(\mathfrak{a}_w)$

e.g. $\widetilde{M}_{3,(1(23))}(x, y, z) = pP[Iix, (1/k^2)[Iiy, Iiz]]$. Note that A_w^1 is an ordinary Lie algebra.

Lemma 8 (Equivalence). *For every $n \geq 2$,*

$$\widetilde{M}_n = (-1)^{n-1}(\chi \otimes \mathbb{1})M_n$$

where $\chi : H^2(\mathfrak{a}_w) \rightarrow H^1(\mathfrak{a}_w)$ is the canonical isomorphism, see (3.5).

Proof. Recall that (A_w, h_A) is a BV algebra, see (6.11). Hence for all $x, y \in A_w^1$:

$$[x, y] = -h_A(xy) + h_A(x)y - xh_A(y) \quad (2.14)$$

Rewrite each vertex in $\widetilde{M}_{n,T'}$ using (2.14) and multiply out, which produces 3^{n-1} terms. All terms but one vanish because $h_A Ii = Ihi = 0$ on $H^1(\mathfrak{a}_w)$ by (1.9) and (3.4); and because $h_A^2 = 0$ by (1.7). The only term that survives is the one that uses $(x, y) \mapsto -h_A(xy)$ at all vertices. Note that $pPh_A = phP$ by (1.9). Summing over all trees $T \in \mathcal{T}_n$ gives

$$\widetilde{M}_n = (-1)^{n-1}(ph \otimes \mathbb{1})M'_n \quad (2.15)$$

where $M'_n : H^1(\mathfrak{a}_w \otimes \mathfrak{c})^{\otimes n} \rightarrow \mathfrak{a}_w^2 \otimes \mathfrak{c}$ is defined like M_n but without the final p in all trees; in particular $(p \otimes \mathbb{1})M'_n = M_n$. But the image of M'_n is in the kernel of the differential, meaning $(d \otimes \mathbb{1})M'_n = 0$ ³⁶; and p applied to a plane wave with momentum k is nonzero only if $k^2 = 0$; therefore (3.5) implies that one can replace ph by χp in (2.15). \square

³⁶First use $dP = Pd_A$, see (1.9). Then this is a property of the construction of M'_n , not of contributions from individual trees but only of the sum over all trees, see the cancellation lemma in [14].

3 Definition of \mathfrak{a} and existence of h

Definition of \mathfrak{a} . Let Λ be the exterior \mathbb{C} -algebra generated by e^0, e^1, e^2, e^3 in degree one. The de Rham dgca on \mathbb{R}^4 is $\Omega = C^\infty(\mathbb{R}^4) \otimes \Lambda$ with differential $d_{\text{dR}} = \partial_\mu \otimes e^\mu$ where by a small abuse of notation, e^μ is also used as a multiplication operator from the left. Let $\mathbb{C} \oplus \mathbb{C}\epsilon$ be the dgca where ϵ is an element of degree -1 ; the product is given by $\epsilon^2 = 0$; and the differential is given by $1 \mapsto 0$ and $\epsilon \mapsto 1$. Consider the tensor product of dgca

$$\mathfrak{a}_\# = \Omega \otimes (\mathbb{C} \oplus \mathbb{C}\epsilon) \simeq \Omega \oplus \Omega\epsilon$$

This means that differential and product on $\mathfrak{a}_\#$ are respectively given by

$$\begin{aligned} d_\# : \quad x \oplus u\epsilon &\mapsto (d_{\text{dR}}x + (-1)^u u) \oplus (d_{\text{dR}}u)\epsilon \\ (x \oplus u\epsilon, y \oplus v\epsilon) &\mapsto xy \oplus (xv + (-1)^y uy)\epsilon \end{aligned}$$

for all $x, y, u, v \in \Omega$. In the diagram for $\mathfrak{a}_\#$ below, Ω occupies the upper row, $\Omega\epsilon$ occupies the lower row with ϵ suppressed. With this understanding, the differential $d_\#$ is as follows, where the six columns correspond to degrees $-1 \dots 4$ of $\mathfrak{a}_\#$ respectively:

$$\begin{array}{ccccccccc} & & \Omega^0 & \xrightarrow{d_{\text{dR}}} & \Omega^1 & \xrightarrow{d_{\text{dR}}} & \Omega^2 & \xrightarrow{d_{\text{dR}}} & \Omega^3 & \xrightarrow{d_{\text{dR}}} & \Omega^4 \\ & \nearrow \mathbb{1} & & \searrow -\mathbb{1} & & \nearrow \mathbb{1} & & \searrow -\mathbb{1} & & \nearrow \mathbb{1} & \\ \Omega^0 & \xrightarrow{d_{\text{dR}}} & \Omega^1 & \xrightarrow{d_{\text{dR}}} & \Omega^2 & \xrightarrow{d_{\text{dR}}} & \Omega^3 & \xrightarrow{d_{\text{dR}}} & \Omega^4 & & \end{array}$$

Set $\Omega_\pm^2 = C^\infty(\mathbb{R}^4) \otimes \Lambda_\pm^2$ where, by definition,

$$\Lambda_-^2 = \mathbb{C}(e^0 e^1 - i e^2 e^3) \oplus \mathbb{C}(e^0 e^2 - i e^3 e^1) \oplus \mathbb{C}(e^0 e^3 - i e^1 e^2) \quad (3.1)$$

$$\Lambda_+^2 = \mathbb{C}(e^0 e^1 + i e^2 e^3) \oplus \mathbb{C}(e^0 e^2 + i e^3 e^1) \oplus \mathbb{C}(e^0 e^3 + i e^1 e^2) \quad (3.2)$$

Using in particular $\Omega_-^2 \Omega_+^2 = 0$ one finds, see [10, 13, 14] and Appendix B for more details:

Lemma 9 (Definition of \mathfrak{a}). *The following is a dgca subquotient of $\mathfrak{a}_\#$:*

$$\mathfrak{a} = \frac{\Omega \oplus (\Omega_+^2 \oplus \Omega^3 \oplus \Omega^4)\epsilon}{(\Omega_-^2 \oplus \Omega^3 \oplus \Omega^4) \oplus 0\epsilon} \quad (3.3)$$

This means that the numerator is a sub-dgca of $\mathfrak{a}_\#$, and the denominator is a dgca ideal in the numerator. As a module, $\mathfrak{a} \simeq (\Omega^0 \oplus \Omega^1 \oplus \Omega_+^2) \oplus (\Omega_+^2 \oplus \Omega^3 \oplus \Omega^4)\epsilon$; as a complex, \mathfrak{a} coincides with (1.1); and as a dgca, \mathfrak{a} has the structure announced in Section 1.

Existence of h . Note that $\Lambda_-^2 = \{x \in \Lambda^2 \mid \star x = -ix\}$ and $\Lambda_+^2 = \{x \in \Lambda^2 \mid \star x = ix\}$, provided one picks the orientation appropriately, where $\star : \Lambda^i \rightarrow \Lambda^{4-i}$ is the Hodge star operator for the bilinear form $\eta = \text{diag}(-1, 1, 1, 1)$ on $\mathbb{C}e^0 \oplus \dots \oplus \mathbb{C}e^3$. Note that $\star^2 = 1$ on odd degree elements, $\star^2 = -1$ on even degree elements. Also write $\star : \Omega^i \rightarrow \Omega^{4-i}$. Denote by $\delta_{\text{dR}} = \star d_{\text{dR}} \star : \Omega^i \rightarrow \Omega^{i-1}$ the ‘adjoint’ of d_{dR} . Let $i_\mu : \Lambda^i \rightarrow \Lambda^{i-1}$ be the interior

multiplications using a dual basis of the e^μ , so $e^\mu e^\nu + e^\nu e^\mu = 0$ and $e^\mu i_\nu + i_\nu e^\mu = \delta_\nu^\mu$ and $i_\mu i_\nu + i_\nu i_\mu = 0$. Then equivalently

$$\delta_{\text{dR}} = -\eta^{\mu\nu} \partial_\mu \otimes i_\nu$$

and therefore $d_{\text{dR}} \delta_{\text{dR}} + \delta_{\text{dR}} d_{\text{dR}} = -\square$ where $\square = \eta^{\mu\nu} \partial_\mu \partial_\nu$. Define

$$h_\# = -\delta_{\text{dR}} \otimes \mathbb{1} \quad : \quad \mathfrak{a}_\#^i \rightarrow \mathfrak{a}_\#^{i-1}$$

Then $h_\#^2 = 0$ and $d_\# h_\# + h_\# d_\# = \square$. The map $h_\#$ is explicitly given as follows:

$$\Omega^0 \xleftarrow{-\delta_{\text{dR}}} \Omega^1 \xleftarrow{-\delta_{\text{dR}}} \Omega^2 \xleftarrow{-\delta_{\text{dR}}} \Omega^3 \xleftarrow{-\delta_{\text{dR}}} \Omega^4$$

$$\Omega^0 \xleftarrow{-\delta_{\text{dR}}} \Omega^1 \xleftarrow{-\delta_{\text{dR}}} \Omega^2 \xleftarrow{-\delta_{\text{dR}}} \Omega^3 \xleftarrow{-\delta_{\text{dR}}} \Omega^4$$

While $h_\#$ does not descend to the subquotient \mathfrak{a} , a modification of it does:

Lemma 10 (Existence of h). *Given \mathbb{C} -linear maps $z : \Lambda^2 \rightarrow \Lambda^0$ and $w : \Lambda^4 \rightarrow \Lambda^2$ that satisfy³⁷ $z\star = iz$ and $\star w = iw$, define $X : \mathfrak{a}_\#^i \rightarrow \mathfrak{a}_\#^{i-2}$ by*

$$\begin{array}{ccccccc} & & z & & i\star & & \\ & \swarrow & & \searrow & & \swarrow & \\ \Omega^0 & \xleftarrow{\quad} & \Omega^1 & \xleftarrow{\quad} & \Omega^2 & \xleftarrow{\quad} & \Omega^3 & \xleftarrow{\quad} & \Omega^4 \\ & \nwarrow & & \swarrow & & \nwarrow & \\ & & -i\star & & w & & \\ & \swarrow & & \searrow & & \swarrow & \\ \Omega^0 & \xleftarrow{\quad} & \Omega^1 & \xleftarrow{\quad} & \Omega^2 & \xleftarrow{\quad} & \Omega^3 & \xleftarrow{\quad} & \Omega^4 \end{array}$$

Then $h_\# + [d_\#, X] : \mathfrak{a}_\#^i \rightarrow \mathfrak{a}_\#^{i-1}$ descends to the subquotient \mathfrak{a} , meaning it leaves numerator and denominator in (3.3) invariant. The induced map $h : \mathfrak{a}^i \rightarrow \mathfrak{a}^{i-1}$ satisfies $dh + hd = \square$. If in addition³⁸ $zw = 2i\star$ then $h^2 = 0$ and h satisfies all requirements in Section 1.

Proof. Recall here that $\Omega\epsilon$ are things in the lower row in the diagrams. Descent amounts to checking that $h_\# + [d_\#, X]$ maps $\Omega_+^2\epsilon \rightarrow \Omega^0 \oplus 0\epsilon$, $\Omega^3\epsilon \rightarrow \Omega^1 \oplus \Omega_+^2\epsilon$, $\Omega_-^2 \rightarrow 0 \oplus 0\epsilon$, and $\Omega^3 \rightarrow \Omega_-^2 \oplus 0\epsilon$. For $h^2 = 0$ it suffices, since h^2 has degree -2 , to check that h^2 annihilates Ω_+^2 , $\Omega^3\epsilon$, $\Omega^4\epsilon$. Both these things are by direct calculation. \square

One can actually check that all h that satisfy the requirements of Section 1 are as in Lemma 10. Nevertheless, in this paper the specific form of h in Lemma 10 is never used.

Lemma 11 (Plane wave homology). *Let $\mathfrak{a}_{\mathbf{w},k} \subseteq \mathfrak{a}$ be the subspace of plane waves with momentum $k \in \mathbb{C}^4 - 0$. The homology of $d|_{\mathfrak{a}_{\mathbf{w},k}}$ is as follows. If $k^2 \neq 0$ then $H(\mathfrak{a}_{\mathbf{w},k}) = 0$. If $k^2 = 0$ then $H^0(\mathfrak{a}_{\mathbf{w},k}) = H^3(\mathfrak{a}_{\mathbf{w},k}) = 0$ and $H^1(\mathfrak{a}_{\mathbf{w},k}) \simeq H^2(\mathfrak{a}_{\mathbf{w},k}) \simeq \mathbb{C}^2$ and*

$$\begin{aligned} \ker(d : \mathfrak{a}_{\mathbf{w},k}^1 \rightarrow \mathfrak{a}_{\mathbf{w},k}^2) &\subseteq \ker(h : \mathfrak{a}_{\mathbf{w},k}^1 \rightarrow \mathfrak{a}_{\mathbf{w},k}^0) \\ \text{image}(h : \mathfrak{a}_{\mathbf{w},k}^3 \rightarrow \mathfrak{a}_{\mathbf{w},k}^2) &\subseteq \text{image}(d : \mathfrak{a}_{\mathbf{w},k}^1 \rightarrow \mathfrak{a}_{\mathbf{w},k}^2) \end{aligned} \tag{3.4}$$

³⁷These conditions are equivalent to $\Lambda_-^2 \subseteq \ker z$ and $\text{image } w \subseteq \Lambda_+^2$ respectively.

³⁸Clearly such z and w exist. Necessarily $z, w \neq 0$, so this h is not Lorentz invariant.

Further, for $k^2 = 0$, the map $h : \mathfrak{a}_{w,k}^2 \rightarrow \mathfrak{a}_{w,k}^1$ induces an isomorphism

$$H^2(\mathfrak{a}_{w,k}) \rightarrow H^1(\mathfrak{a}_{w,k}) \quad (3.5)$$

and this isomorphism is canonical, independent of h .

Proof. The homology is calculated in [14]. Let $k^2 = 0$. Then (i) $dh + hd = 0$ on $\mathfrak{a}_{w,k}$; (ii) $\ker(d : \mathfrak{a}_{w,k}^0 \rightarrow \mathfrak{a}_{w,k}^1) = 0$ because $H^0(\mathfrak{a}_{w,k}) = 0$ and $\mathfrak{a}_{w,k}^{-1} = 0$; (iii) $\mathfrak{a}_{w,k}^3 = \text{image}(d : \mathfrak{a}_{w,k}^2 \rightarrow \mathfrak{a}_{w,k}^3)$ because $H^3(\mathfrak{a}_{w,k}) = 0$ and $\mathfrak{a}_{w,k}^4 = 0$. They imply (3.4): if $x \in \mathfrak{a}_{w,k}^1$ and $dx = 0$ then $0 = hdx = -dhx$ by (i) hence $hx = 0$ by (ii); if $x \in \mathfrak{a}_{w,k}^3$ then $x = dy$ for $y \in \mathfrak{a}_{w,k}^2$ by (iii) hence $hx = hdy = -dhy$ by (i) so hx is in the image of d . By (i), h induces a map (3.5); that this is an isomorphism follows by expanding d and h to first order around the given momentum k and using (1.3), details omitted. The proof does not use $h^2 = 0$. \square

4 Definition of BV_∞^Q -algebras

Section 4.1 contains an account of G_∞ - and BV_∞ -algebras following [23], but restricted to the special case in [23, Remark 2.8], [24, Proposition 24]. In Section 4.2 this definition is deformed using a quadratic element in a Hopf algebra. Section 4.4 is about a special case where all but certain operations are zero, to be applied to Yang-Mills in Section 6.1.

Remark 1 (Dualization). Following [23], the account below avoids the cofree Gerstenhaber coalgebra. This requires the following clarification. Given a graded vector space V , the algebraic structures we define are collections of maps $V^* \rightarrow (V^*)^{\otimes n}$ subject to certain quadratic axioms; here V^* is the dual. In the unlikely case that $\dim_{\mathbb{C}} V < \infty$ this is, by taking duals, equivalent to collections of maps $V^{\otimes n} \rightarrow V$ subject to the corresponding dual axioms. Whether V is finite-dimensional or not, we agree that the proper way to interpret the definitions is that *one must dualize the axioms, rather than the maps, to get axioms for the collection of maps $V^{\otimes n} \rightarrow V$* . It is still sometimes convenient to say that we dualize the maps, but this is always meant formally and with the reservations in this remark.

4.1 Recap of ordinary G_∞ - and BV_∞ -algebras

Graded vector space. The ground field is \mathbb{C} . The constructions in this section require a graded vector space $V = \bigoplus_{i \in \mathbb{Z}} V^i$ as input data. Associated objects such as $V \otimes V$ or the dual $V^* = \text{Hom}_{\mathbb{C}}(V, \mathbb{C})$ carry an induced grading; a \mathbb{C} -linear map from one graded vector space to another has degree i iff it raises the degree by i , thus $(V^*)^i \simeq (V^{-i})^*$.

Gerstenhaber algebra. Briefly, a Gerstenhaber algebra is a graded vector space A that has two compatible structures:

- A graded commutative associative algebra or *gca* structure $A^i \otimes A^j \rightarrow A^{i+j}$.
- A degree -1 graded Lie algebra or *gLa* structure $[-, -] : A^i \otimes A^j \rightarrow A^{i+j-1}$.

The gca multiplication is denoted by juxtaposition. In full detail, the gca axioms, the gLa axioms and the compatibility axiom say that the following must be zero for all $a, b, c \in A$:

$$ab - (-1)^{ab}ba \quad (4.1a)$$

$$(ab)c - a(bc) \quad (4.1b)$$

$$[a, b] + (-1)^{(a+1)(b+1)}[b, a] \quad (4.1c)$$

$$[a, [b, c]] + (-1)^{(a+1)(b+c)}[b, [c, a]] + (-1)^{(c+1)(a+b)}[c, [a, b]] \quad (4.1d)$$

$$[ab, c] - a[b, c] - (-1)^{ab}b[a, c] \quad (4.1e)$$

A Gerstenhaber algebra may or may not have a unit.

Equivalently, a Gerstenhaber algebra is an algebra of the homology of the little disks operad. The little disks operad is a sequence of topological spaces indexed by an integer n , closely related to configuration spaces of n points in \mathbb{R}^2 , together with an operad structure, whose singular homology neatly captures the Gerstenhaber algebra axioms, see [27].

*Free Gerstenhaber algebra*³⁹. Let V be a graded vector space. The free Gerstenhaber algebra of V^* *without* unit will be denoted GV^* , with a degree zero map $V^* \hookrightarrow GV^*$. The free Gerstenhaber algebra GV^* is given by the following construction (sketch):

- Terms: Linear combinations of ‘Gerstenhaber words’ that in this context are build from elements of V^* combined using two formal operations: the gca product to-be denoted by juxtaposition; the gLa product to-be denoted by $[-, -]$. The products have no properties yet, except that all \mathbb{C} -linearities are already understood.
- Relations: Quotient by the relations in the Gerstenhaber algebra axioms (4.1).

The empty term is not allowed (no unit). An example of a term is $[v_1[v_2, v_3], v_4]v_5$ with $v_i \in V^*$; its degree is $\sum_i |v_i| - 2$, and it is equal to⁴⁰

$$v_1[[v_2, v_3], v_4]v_5 + (-1)^{v_1(v_2+v_3-1)}[v_2, v_3][v_1, v_4]v_5$$

in GV^* . Like in this example, one can always move all gca products to the outside.

Producing a \mathbb{C} -basis for GV^* is an interesting topic. This acquires a geometric flavor using the isomorphic⁴¹ $GV^* = \oplus_{n \geq 0} H(D_n) \otimes_{\mathbb{C}[S_n]} (V^*)^{\otimes n}$, with D_n the space of n little

³⁹The free Gerstenhaber algebra of V^* comes in for the following reason. In general, see [25], to every quadratic operad \mathbf{P} one can associate its Koszul dual cooperad \mathbf{P}^\vee [16]. Some quadratic operads are Koszul [16], an adjective. For conceptual reasons [25], if \mathbf{P} is quadratic and Koszul, a \mathbf{P}_∞ -algebra on V is defined to be a codifferential of degree 1 on the cofree \mathbf{P}^\vee -coalgebra cogenerated by V . The Gerstenhaber operad \mathbf{G} (axioms (4.1)) is quadratic and Koszul [17] (like Ass, Lie, Com). It is Koszul self-dual (like Ass, unlike Lie, Com), so, modulo degree shifts, a \mathbf{G}^\vee -coalgebra is the same thing as a Gerstenhaber coalgebra (operations and axioms linearly dual to (4.1)). Hence a \mathbf{G}_∞ -algebra on V is a codifferential of degree 1 on the cofree Gerstenhaber coalgebra cogenerated by V ; this is formally the same thing as a differential of degree 1 on the free Gerstenhaber algebra generated by V^* , cf. [16, Proposition 4.2.14] and [23] and Remark 1.

⁴⁰Additional brackets are omitted, using the associativity axiom in (4.1).

⁴¹This way of associating to a vector space a free algebra is known as a ‘Schur functor’, see e.g. [33]. The action of the permutation group S_n is the obvious one but involves degree-dependent signs.

disks, S_n the symmetric group. The size of GV^* is therefore related to the size of the singular homology $H(D_n)$ which may be explicitly calculated, together with the cohomology ring, using the de Rham pairing, that is, integration of forms [27].

The n -degree and the decreasing filtration. The n -degree is given by the word length on GV^* , so $n|_{V^*} = 1$ and $n(ab) = n([a, b]) = n(a) + n(b)$ for all n -homogeneous $a, b \in GV^*$. Abbreviate $n(a) = n_a$. The n -degree is different from the primary grading on GV^* coming from the grading on V . Define a decreasing filtration by letting $\mathcal{F}_n = \mathcal{F}_n GV^*$ be the \mathbb{C} -span of terms of n -degree $\geq n$. Then $\mathcal{F}_1 = GV^*$ and $\mathcal{F}_n \mathcal{F}_m \subseteq \mathcal{F}_{n+m}$ and $[\mathcal{F}_n, \mathcal{F}_m] \subseteq \mathcal{F}_{n+m}$; and $\bigcap_{n \geq 1} \mathcal{F}_n = 0$. An endomorphism is in $\mathcal{F}_n \text{End}_{\mathbb{C}}(GV^*)$ iff it maps $\mathcal{F}_m \rightarrow \mathcal{F}_{m+n}$ for all m .

Remark 2 (Completion). The completion relative to the filtration $\mathcal{F}_{\bullet} GV^*$ allows countably infinite linear combinations analogous to formal power series. In the remainder of this section, one should replace GV^* by this completion, however we will not introduce separate notation for this. The completion is also filtered.

Derivations. For every Gerstenhaber algebra A , a homogeneous element⁴² $\delta \in \text{End}_{\mathbb{C}}(A)$ is called a Gerstenhaber derivation, or simply derivation, if

$$\begin{aligned}\delta(ab) &= \delta(a)b + (-1)^{\delta_a} a\delta(b) \\ \delta([a, b]) &= [\delta(a), b] + (-1)^{\delta(a-1)} [a, \delta(b)]\end{aligned}$$

The space of all Gerstenhaber derivations is denoted $\text{Der}(A)$ and it is a gLa using the graded commutator, $[\delta_1, \delta_2] = \delta_1 \delta_2 - (-1)^{\delta_1 \delta_2} \delta_2 \delta_1$ for all $\delta_1, \delta_2 \in \text{Der}(A)$, so it is a sub-gLa of $\text{End}_{\mathbb{C}}(A)$; one checks that $[\delta_1, \delta_2]$ is itself a Gerstenhaber derivation.

Definition of \mathbf{G}_{∞} -algebra. For every graded vector space V define [23]:

$$\mathbf{G}_{\infty}\text{-algebra structure on } V \quad \Longleftrightarrow \quad \delta \in \text{Der}^1(GV^*) \text{ with } [\delta, \delta] = 0$$

A few comments are in order:

- δ being of degree one, the Maurer-Cartan equation $[\delta, \delta] = 0$ is equivalent to $\delta^2 = 0$. Therefore, a \mathbf{G}_{∞} structure is a derivation δ of degree one that is also a differential.
- There are analogous definitions of \mathbf{A}_{∞} , \mathbf{C}_{∞} , \mathbf{L}_{∞} using respectively the free graded associative or tensor algebra; the free gLa; the free gca, without unit. Note [23] that GV^* contains a copy of the free gLa as a quotient gLa and a copy of the free dgca as a quotient gca, and every Gerstenhaber derivation descends to these quotients. Therefore every \mathbf{G}_{∞} structure contains both a \mathbf{C}_{∞} and an \mathbf{L}_{∞} structure.
- δ is determined by its restriction to V^* and this data is unconstrained⁴³,

$$\text{Der}(GV^*) \simeq \text{Hom}_{\mathbb{C}}(V^*, GV^*) \tag{4.2}$$

By decomposing GV^* one obtains a decomposition of δ . Recall Remarks 1, 2 here.

⁴²Here and below, $\text{End}_{\mathbb{C}}$ are simply the \mathbb{C} -linear maps, not endomorphisms of algebras.

⁴³By an ‘infinitesimal’ version of the universal property of GV^* .

A \mathbf{G}_∞ -algebra with $\delta(V^*) \subseteq V^*V^* + [V^*, V^*]$ is an ordinary Gerstenhaber algebra⁴⁴.

The canonical operator α . The free Gerstenhaber algebra has an interesting canonical structure [23]. Namely, define $\alpha \in \text{End}_{\mathbb{C}}^{-1}(GV^*)$ by requiring⁴⁵

$$\begin{aligned}\alpha|_{V^*} &= 0 \\ \alpha(ab) &= (-1)^a[a, b] + \alpha(a)b + (-1)^a a\alpha(b) \\ \alpha([a, b]) &= [\alpha(a), b] + (-1)^{a-1}[a, \alpha(b)]\end{aligned}\tag{4.3}$$

for all $a, b \in GV^*$. This recursively fixes how α acts on all terms, and one checks that it is well-defined meaning compatible with the relations (4.1)⁴⁶. Further:

- α has degree -1 because $|[a, b]| = |ab| - 1$.
- $\alpha^2 = 0$ ⁴⁷ and therefore⁴⁸ $d_\alpha = [\alpha, -] \in \text{End}_{\mathbb{C}}^{-1}(\text{End}_{\mathbb{C}}(GV^*))$ is a differential, $d_\alpha^2 = 0$, and it satisfies the Leibniz rule for the gLa structure on $\text{End}_{\mathbb{C}}(GV^*)$ ⁴⁹.
- If $\delta \in \text{Der}(GV^*)$ then $d_\alpha(\delta) \in \text{Der}(GV^*)$ ⁵⁰. So, $d_\alpha \in \text{End}_{\mathbb{C}}^{-1}(\text{Der}(GV^*))$.

Hence $\text{Der}(GV^*)$ is a differential graded Lie algebra with a differential d_α of degree -1 .

Definition of \mathbf{BV}_∞ -algebra. With [23, Remark 2.8], [24, Proposition 24] define:

$$\mathbf{BV}_\infty\text{-algebra structure on } V \iff \begin{cases} \delta^i \in \text{Der}^i(GV^*) \text{ for } i = 1, -1, -3, -5, \dots \text{ with} \\ d_\alpha \delta + \frac{1}{2}[\delta, \delta] = 0 \\ \text{where } \delta = \delta^1 + \delta^{-1} + \delta^{-3} + \dots \end{cases}$$

For degree reasons it is necessary that $[\delta^1, \delta^1] = 0$, hence every \mathbf{BV}_∞ structure contains a \mathbf{G}_∞ structure. The next equation is $d_\alpha \delta^1 + [\delta^1, \delta^{-1}] = 0$, and so forth.

4.2 \mathbf{BV}_∞^Q -algebras, with axioms deformed by a quadratic element Q

Here we modify the definition of a \mathbf{BV}_∞ -algebra in the case when V is also a module over a cocommutative Hopf algebra H , and a central ‘quadratic’ element in H is fixed⁵¹.

⁴⁴Namely, after dualization, it is a Gerstenhaber algebra on $V[-2]$ where $V[-2]^i = V^{i-2}$. The part mapping into V^*V^* encodes the gLa product, the part mapping into $[V^*, V^*]$ encodes the gca product.

⁴⁵This operator is denoted Δ_{Lie} in [23]. In this paper, Δ is reserved for a coproduct in Section 4.2.

⁴⁶Direct calculation. See also [23].

⁴⁷Check $\alpha^2|_{V^*} = 0$ and $\alpha^2(ab) = \alpha^2(a)b + a\alpha^2(b)$ and $\alpha^2([a, b]) = [\alpha^2(a), b] + [a, \alpha^2(b)]$.

⁴⁸Explicitly, set $d_\alpha(e) = \alpha e - (-1)^e e\alpha$ for all $e \in \text{End}_{\mathbb{C}}(GV^*)$.

⁴⁹Explicitly, the Leibniz rule is $d_\alpha([e_1, e_2]) = [d_\alpha(e_1), e_2] + (-1)^{e_1}[e_1, d_\alpha(e_2)]$ for all $e_1, e_2 \in \text{End}_{\mathbb{C}}(GV^*)$. It is trivially true since d_α is given by a graded commutator.

⁵⁰By direct calculation, one finds $d_\alpha(\delta)|_{V^*} = 0$ and $d_\alpha(\delta)(ab) = (d_\alpha(\delta)(a))b + (-1)^{(\delta-1)a}a(d_\alpha(\delta)(b))$ and $d_\alpha(\delta)([a, b]) = [d_\alpha(\delta)(a), b] + (-1)^{(\delta-1)(a-1)}[a, d_\alpha(\delta)(b)]$, for all $a, b \in GV^*$.

⁵¹Operads \mathbf{P} in the category of H -modules, where H is a graded cocommutative Hopf algebra, have been discussed in the literature [28, 29]. One can then form a semidirect product of operads $\mathbf{P} \rtimes H$, see [28]. The action of H on \mathbf{P} can encode interesting information, for instance the semidirect product of the operad of Gerstenhaber algebras with the Hopf algebra of dual numbers, acting in a certain way, is isomorphic to the operad of BV algebras [29]. It is not excluded that some of this theory can be applied here.

Hopf module and quadratic element. In this section, V is both a graded vector space and an H -module. Here H is a cocommutative Hopf algebra with coproduct denoted Δ . The Hopf algebra is not graded and leaves the grading on V invariant. Further, the constructions in this section require the choice of an element $Q \in H$ with these properties:

- Q satisfies the 7-term identity

$$\Delta^2 Q - (\mathbb{1} + \sigma + \sigma^2)(1 \otimes \Delta Q) + (\mathbb{1} + \sigma + \sigma^2)(1 \otimes 1 \otimes Q) = 0 \quad (4.4)$$

where $1 \in H$ is the unit and where $\sigma : H^{\otimes 3} \rightarrow H^{\otimes 3}$, $h_1 \otimes h_2 \otimes h_3 \mapsto h_2 \otimes h_3 \otimes h_1$.

- Q is in the kernel of the counit $H \rightarrow \mathbb{C}$ of the Hopf algebra⁵².
- Q is in the center of the Hopf algebra.

The motivating example is when H is a polynomial algebra⁵³ and Q is a polynomial of degree at most two, without constant term. Another example, not in general commutative but cocommutative, is the universal enveloping algebra $H = U\mathfrak{g}$ ⁵⁴ of a Lie algebra \mathfrak{g} , and Q an element of degree at most two, without constant term, in the center of $U\mathfrak{g}$.

Hopf module structure on GV^ .* Since V is an H -module, so is $(V^*)^{\otimes n}$ ⁵⁵. By cocommutativity of H , on $(V^*)^{\otimes n}$ the S_n -module and H -module structures are compatible⁵⁶. Hence GV^* is an H -module. The following maps are H -equivariant: the inclusion $V^* \hookrightarrow GV^*$; the gca and gLa products as maps $GV^* \otimes GV^* \rightarrow GV^*$; the map $\alpha : GV^* \rightarrow GV^*$.

As an example, if $h \in H$ and $v_i \in V^*$ then

$$h([v_1[v_2, v_3], v_4]v_5) = [(h_1 v_1)[(h_2 v_2), (h_3 v_3)], (h_4 v_4)](h_5 v_5)$$

where $\Delta^4 h = h_1 \otimes \cdots \otimes h_5$ is Sweedler notation for the iterated coproduct $\Delta^4 : H \rightarrow H^{\otimes 5}$.

Equivariant derivations. Denote by $\text{Der}(GV^*)^H \subseteq \text{Der}(GV^*)$ the sub-graded Lie algebra of H -equivariant elements. Note that d_α maps this sub-gLa to itself.

The operator β . Here the fixed element $Q \in H$ is used to make a definition inspired by and superficially similar to (4.3). Namely define $\beta \in \text{End}_{\mathbb{C}}^1(GV^*)$ by

$$\begin{aligned} \beta|_{V^*} &= 0 \\ \beta(ab) &= \beta(a)b + (-1)^a a\beta(b) \\ \beta([a, b]) &= (-1)^a (Q(ab) - (Qa)b - a(Qb)) + [\beta(a), b] + (-1)^{a-1} [a, \beta(b)] \end{aligned} \quad (4.5)$$

for all $a, b \in GV^*$. This recursively fixes how β acts on all terms, and one checks that it is well-defined meaning compatible with the relations (4.1); this calculation requires the 7-term identity (4.4) and is in Appendix A. Further:

⁵²Actually this follows from (4.4) by applying $(\mathbb{1} \otimes \epsilon)(\mathbb{1} \otimes \mathbb{1} \otimes \epsilon)$ where $\epsilon : H \rightarrow \mathbb{C}$ is the counit, and using the Hopf algebra axioms. The counit plays the role of the trivial representation of a Hopf algebra.

⁵³On $\mathbb{C}[x_1, \dots, x_n]$ the coalgebra structure is determined by $\Delta x_i = x_i \otimes 1 + 1 \otimes x_i$.

⁵⁴On $U\mathfrak{g}$ the coalgebra structure is determined by $\Delta g = g \otimes 1 + 1 \otimes g$ for all $g \in \mathfrak{g}$.

⁵⁵For $n = 0$ the module structure is the counit, but we do not need $n = 0$ here.

⁵⁶That is, one has a bimodule.

- β has degree +1 because $|ab| = |[a, b]| + 1$.
- $\beta : GV^* \rightarrow GV^*$ is H -equivariant, by the assumption that Q is in the center.
- $\beta^2 = 0$ ⁵⁷ and therefore $d_\beta = [\beta, -] \in \text{End}_{\mathbb{C}}^1(\text{End}_{\mathbb{C}}(GV^*))$ is a differential, $d_\beta^2 = 0$, and it satisfies the Leibniz rule for the gLa structure on $\text{End}_{\mathbb{C}}(GV^*)$.
- If $\delta \in \text{Der}(GV^*)^H$ then $d_\beta(\delta) \in \text{Der}(GV^*)^H$. So, $d_\beta \in \text{End}_{\mathbb{C}}^1(\text{Der}(GV^*)^H)$.

The last claim, which is only for H -equivariant δ , is also proved in Appendix A.

A curvature operator. The sum $d_\alpha + d_\beta$ is not a differential because $d_\alpha d_\beta + d_\beta d_\alpha \neq 0$. Its restriction to H -equivariant derivations is explicitly given by

$$d_\alpha d_\beta + d_\beta d_\alpha = -[K, -] \quad \text{on } \text{Der}(GV^*)^H \quad (4.6)$$

where, by definition, $K \in \text{Der}^0(GV^*)^H$ is given by

$$\begin{aligned} K|_{V^*} &= Q\mathbb{1} \\ K(ab) &= K(a)b + aK(b) \\ K([a, b]) &= [K(a), b] + [a, K(b)] \end{aligned}$$

Note that K is H -equivariant since Q is in the center. One finds $d_\alpha(K) = d_\beta(K) = 0$. See Appendix A for proofs. In summary, on $\text{Der}(GV^*)^H$, the map $d_\alpha + d_\beta$ is not a differential, but together with $-K$ one has a curved differential $\mathbb{Z}/2\mathbb{Z}$ -graded Lie algebra. The Maurer-Cartan equation in this curved dgLa motivates the following definition.

Definition of BV_∞^Q -algebra. Set

$$BV_\infty^Q\text{-algebra structure on } V \iff \begin{cases} \delta^i \in \text{Der}^i(GV^*)^H \text{ for } i = 1, -1, -3, \dots \text{ with} \\ -K + (d_\alpha + d_\beta)\delta + \frac{1}{2}[\delta, \delta] = 0 \\ \text{where } \delta = \delta^1 + \delta^{-1} + \delta^{-3} + \dots \end{cases} \quad (4.7)$$

The first few degree levels of these equations are⁵⁸

$$\begin{aligned} \frac{1}{2}[\delta^1, \delta^1] &= -d_\beta \delta^1 \\ d_\alpha \delta^1 + [\delta^1, \delta^{-1}] &= K - d_\beta \delta^{-1} \\ d_\alpha \delta^{-1} + [\delta^1, \delta^{-3}] + \frac{1}{2}[\delta^{-1}, \delta^{-1}] &= -d_\beta \delta^{-3} \\ &\vdots \end{aligned} \quad (4.8)$$

The terms on the right correspond to the deformation by Q . Already the first equation is deformed, so that δ^1 is not necessarily a G_∞ -algebra⁵⁹. But by the definition of d_β , note that δ^1 must still contain a C_∞ -algebra, including a differential.

⁵⁷Check $\beta^2|_{V^*} = 0$ and $\beta^2(ab) = \beta^2(a)b + a\beta^2(b)$ and $\beta^2([a, b]) = [\beta^2(a), b] + [a, \beta^2(b)]$, which implies $\beta^2 = 0$. The 7-term identity (4.4) is not needed for this calculation.

⁵⁸In the special case $\delta^{-3} = \delta^{-5} = \dots = 0$, only the three displayed equations are non-vacuous.

⁵⁹Since it only involves δ^1 , we could say that it defines a G_∞^Q -algebra.

4.3 The auxiliary operator Γ

Here a $\Gamma \in \text{End}_{\mathbb{C}}^1(\text{Der}(GV^*))$ is defined such that $\Gamma^2 = 0$ and

$$d_{\alpha}\Gamma + \Gamma d_{\alpha} = \mathbb{1}_{>0} \quad (4.9)$$

where, for every derivation δ , set $\mathbb{1}_{>0}(\delta) = 0$ if $n_{\delta} = 0$ and $\mathbb{1}_{>0}(\delta) = \delta$ if $n_{\delta} > 0$. The n -degree was defined in Section 4.1; a derivation δ has n -degree equal to n_{δ} iff it increases the n -degree by n_{δ} . Further, Γ maps H -equivariant derivations to H -equivariant derivations.

Construction of Γ . Define $\gamma \in \text{End}_{\mathbb{C}}^1(GV^*)$ by

$$\begin{aligned} \gamma|_{V^*} &= 0 \\ \gamma(ab) &= \gamma(a)b + (-1)^a a\gamma(b) \\ \gamma([a, b]) &= (-1)^a n_a n_b ab + [\gamma(a), b] + (-1)^{a-1} [a, \gamma(b)] \end{aligned}$$

for all n -homogeneous $a, b \in GV^*$. It is well-defined on GV^* . Both α and γ preserve the n -degree. By direct calculation, $\gamma^2 = 0$, and $\kappa = \alpha\gamma + \gamma\alpha \in \text{End}_{\mathbb{C}}^0(GV^*)$ is given by⁶⁰

$$\kappa(a) = \frac{1}{2} n_a (n_a - 1) a \quad (4.10)$$

Now, if $\delta \in \text{Der}(GV^*)$ then $[\gamma, \delta]$ need not be a derivation. Hence, for every derivation δ of n -degree equal to n_{δ} , let $\Gamma(\delta) \in \text{Der}(GV^*)$ be the unique derivation with

$$\Gamma(\delta)|_{V^*} = \begin{cases} 0 & \text{if } n_{\delta} = 0 \\ \frac{2}{(n_{\delta}+1)n_{\delta}} [\gamma, \delta]|_{V^*} & \text{if } n_{\delta} > 0 \end{cases}$$

Then $\Gamma^2 = 0$ by $\gamma^2 = 0$. To see (4.9), suppose $n_{\delta} > 0$; the case $n_{\delta} = 0$ is trivial. Then $\delta(V^*)$ has n -degree equal to $n_{\delta} + 1$. Set $\delta' = (d_{\alpha}\Gamma + \Gamma d_{\alpha})\delta$ which is a derivation, $n_{\delta'} = n_{\delta}$, and $\delta'|_{V^*} = \frac{2}{(n+1)n} (\kappa\delta)|_{V^*}$, and therefore (4.10) implies $\delta' = \delta$ as required.

Another property of Γ . If $\delta \in \text{Der}(GV^*)$ and $n_{\delta} = 0$ then $[\delta, -]$ commutes with Γ up to sign. More explicitly, using the definition of Γ , this means that

$$\delta(V^*) \subseteq V^* \quad \implies \quad \Gamma[\delta, -] = (-1)^{\delta} [\delta, \Gamma-] \quad (4.11)$$

4.4 A special class

The special case. Throughout this section,

$$\delta = \sum_{k \geq 1} \mu_k + \sum_{k \geq 2} \nu_k$$

where $\mu_k, \nu_k \in \text{Der}^{\text{odd}}(GV^*)^H$ satisfy

$$\begin{aligned} \mu_k(V^*) &\subseteq (V^*)^{\otimes k} \\ \nu_k(V^*) &\subseteq [V^*, V^*](V^*)^{\otimes(k-2)} \end{aligned} \quad (4.12)$$

⁶⁰Check $\kappa|_{V^*} = 0$ and $\kappa(ab) = n_a n_b ab + \kappa(a)b + a\kappa(b)$ and $\kappa([a, b]) = n_a n_b [a, b] + [\kappa(a), b] + [a, \kappa(b)]$.

It is always understood that $\mu_k = 0$ for $k \leq 0$ respectively $\nu_k = 0$ for $k \leq 1$. Further assume that almost all operations have degree one: $\mu_k, \nu_k \in \text{Der}^1$ for all $k \geq 2$ and

$$\mu_1 = d + h$$

where $d \in \text{Der}^1$ and $h \in \text{Der}^{-1}$. (In the notation of [24], this means that only the structure maps $m_{1,\dots,1}^0$, $m_{1,\dots,1,2}^0$ and m_1^1 can be nonzero.) The notation d, h is merely suggestive for the synonymous operations in the Yang-Mills dgca \mathfrak{a} ; beware that in this section d and h are not primarily maps $V^* \rightarrow V^*$ but as derivations that map $V^* \rightarrow V^*$. Using (4.12),

$$d_\alpha \mu_1 = 0 \quad d_\alpha \nu_2 = 0 \quad d_\beta \mu_k = 0 \quad \Gamma \mu_k = 0 \quad (4.13)$$

Decomposition of the axioms. Define $A_k, B_k, C_k \in \text{Der}^{\text{even}}(GV^*)^H$ by

$$\begin{aligned} A_k &= -K\delta_{k1} + d_\beta \nu_k + \frac{1}{2} \sum_{m+n-1=k} [\mu_m, \mu_n] \\ B_k &= +d_\alpha \mu_k + \sum_{m+n-1=k} [\mu_m, \nu_n] \\ C_k &= +d_\alpha \nu_k + \frac{1}{2} \sum_{m+n-1=k} [\nu_m, \nu_n] \end{aligned}$$

with δ_{k1} the Kronecker delta. The sums are finite, and note that A_k, B_k, C_k are zero except when $k \geq 1, k \geq 2, k \geq 3$ respectively using (4.13). By (4.13),

$$-K + (d_\alpha + d_\beta)\delta + \frac{1}{2}[\delta, \delta] = \sum_{k \geq 1} A_k + \sum_{k \geq 2} B_k + \sum_{k \geq 3} C_k$$

Hence if all A_k, B_k, C_k vanish then δ is a BV_∞^Q -algebra structure, see (4.7). The converse is also true, because A_k, B_k, C_k map V^* to linearly independent sectors of GV^* :

$$\begin{aligned} A_k(V^*) &\subseteq (V^*)^{\otimes k} \\ B_k(V^*) &\subseteq [V^*, V^*](V^*)^{\otimes(k-2)} \\ C_k(V^*) &\subseteq [[V^*, V^*], V^*](V^*)^{\otimes(k-3)} + \underbrace{[V^*, V^*][V^*, V^*](V^*)^{\otimes(k-4)}}_{\text{only for } k \geq 4} \end{aligned}$$

The goal of this section is to spell out the linear dependencies among the A_k, B_k, C_k and to reduce the axioms to a form that one can check for Yang-Mills, in Section 6.1.

Linear dependencies. One finds

$$\Gamma A_k = 0 \quad (4.14)$$

$$d_\beta A_k = 0 \quad (4.15)$$

$$d_\beta B_k = \sum_{m+n-1=k} [A_m, \mu_n] \quad (4.16)$$

$$d_\alpha A_k + d_\beta C_k = \sum_{m+n-1=k} [A_m, \nu_n] + [B_m, \mu_n] \quad (4.17)$$

$$d_\alpha B_k = \sum_{m+n-1=k} [B_m, \nu_n] + [C_m, \mu_n] \quad (4.18)$$

$$d_\alpha C_k = \sum_{m+n-1=k} [C_m, \nu_n] \quad (4.19)$$

To see this, use (4.6); (4.13); $d_\alpha(K) = d_\beta(K) = 0$; $d_\alpha^2 = d_\beta^2 = 0$; the Leibniz rule for d_α and d_β ; and the Jacobi identity by which $[y, [y, y]] = 0$ for all $y \in \text{Der}^{\text{odd}}(GV^*)$.

Lemma 12 (Reduced axioms). δ is a BV_∞^Q -algebra iff

$$A_1 = 0 \quad \Gamma B_k = 0 \text{ for all } k \geq 2 \quad \Gamma C_k = 0 \text{ for all } k \geq 3 \quad (4.20)$$

More precisely, for every integer K , if (4.20) hold for $k < K$ then $A_k = B_k = C_k = 0$ for $k < K$, and if in addition $\Gamma C_K = 0$ then $C_K = 0$.

Proof. We prove \Leftarrow by induction on k . At each step we show that A_k, B_k, C_k vanish. Case $k = 1$: Vacuous since $B_1 = C_1 = 0$. Case $k = 2$: Use (4.18) to see that $d_\alpha B_2 = 0$ (note that $\nu_1 = 0$ and $C_2 = 0$) and by assumption $\Gamma B_2 = 0$, which, since B_2 has n -degree $+1$ and using (4.9), implies $B_2 = 0$. Now (4.17) implies $d_\alpha A_2 = 0$ which with (4.14) and (4.9) implies $A_2 = 0$. Case $k = 3$: $d_\alpha C_3 = 0$ by (4.19) which with $\Gamma C_3 = 0$ and (4.9) implies $C_3 = 0$. Now (4.18) implies $d_\alpha B_3 = 0$ which with $\Gamma B_3 = 0$ implies $B_3 = 0$. Now $d_\alpha A_3 = 0$ by (4.17), and (4.14), imply $A_3 = 0$. For $k \geq 4$, the argument is just like for $k = 3$. \square

Lemma 13 (A special case of the special case). Restrict to the case when δ is determined by $\mu_1 = d + h$ as before and by a sequence $\theta_k \in \text{Der}^2(GV^*)^H$, $k \geq 2$, with

$$\theta_k(V^*) \subseteq (V^*)^{\otimes k} \quad (4.21)$$

by setting

$$\begin{aligned} \nu_k &= d_\alpha \theta_k \\ \mu_k &= -\Gamma[h, \nu_k] \end{aligned} \quad (4.22)$$

for all $k \geq 2$; they have degree one and are consistent with (4.12). Then:

- Part A: δ is a BV_∞^Q -algebra iff
 - $[h, h] = 0$ and $[d, h] = K$ and $[d, d] = 0$.
 - $b_k = 0$ where, by definition, $b_k = \Gamma[d, \nu_k] + \Gamma \sum_{m+n-1=k, m \neq 1} [\mu_m, \nu_n]$, for $k \geq 2$.
 - $c_k = 0$ where, by definition, $c_k = \frac{1}{2} \Gamma \sum_{m+n-1=k} [\nu_m, \nu_n]$, for $k \geq 3$.
- Part B: Assuming only the first bullet in Part A, the following implications hold:
 - If $b_K = c_K = 0$ for all $K < k$ then $[d, c_k] = 0$.
 - If $b_K = c_K = 0$ for all $K < k$ and if $c_k = 0$ then $[d, b_k] = 0$.

Proof. In Part A, the first bullet means $A_1 = 0$, the three equations corresponding to degrees $-2, 0, 2$ respectively. Let $B_k = B_k^0 \oplus B_k^2$ and $C_k = C_k^0 \oplus C_k^2$ be the decompositions by degree. Then $\Gamma B_k^0 = \Gamma C_k^0 = 0$ by construction of μ_k and ν_k , in fact, using (4.9):

$$\begin{aligned} \Gamma B_k^0 &= \Gamma(d_\alpha \mu_k + [h, \nu_k]) = \Gamma(-d_\alpha \Gamma + \mathbb{1}_{>0})[h, \nu_k] = \Gamma^2 d_\alpha [h, \nu_k] = 0 \\ \Gamma C_k^0 &= \Gamma d_\alpha \nu_k = \Gamma d_\alpha^2 \theta_k = 0 \end{aligned}$$

Since $b_k = \Gamma B_k^2$, $c_k = \Gamma C_k^2$, Part A follows from Lemma 12. Part B: By Lemma 12, $B_K = C_K = 0$ for $K < k$, and if $c_k = 0$ then also $C_k = 0$. The degree 3 part of (4.18) (recall $\nu_1 = 0$) implies $[C_k^2, d] = 0$, so, by (4.11), $[d, c_k] = [d, \Gamma C_k^2] = -\Gamma[d, C_k^2] = 0$. If $c_k = 0$ then $C_k = 0$, so the degree 3 part of (4.17) implies $[B_k^2, d] = 0$, so $[d, b_k] = [d, \Gamma B_k^2] = -\Gamma[d, B_k^2] = 0$. \square

Note that $b_k, c_k \in \text{Der}^3(GV^*)^H$ satisfy

$$\begin{aligned} b_k(V^*) &\subseteq (V^*)^{\otimes k} \\ c_k(V^*) &\subseteq [V^*, V^*](V^*)^{\otimes(k-2)} \end{aligned} \quad (4.23)$$

Note that (4.21) implies $\Gamma\theta_k = 0$ and therefore $\Gamma\nu_k = \theta_k$ by (4.9). Hence (4.11) implies $\Gamma[d, \nu_k] = -[d, \Gamma\nu_k] = -[d, \theta_k]$. Therefore $b_k = -[d, \theta_k] + q_k$ where, by definition,

$$q_k = \Gamma \sum_{m+n-1=k, m \neq 1} [\mu_m, \nu_n] \quad (4.24)$$

Part B and $[d, d] = 0$ give the implication

$$\text{If } b_K = c_K = 0 \text{ for all } K < k \text{ and if } c_k = 0 \text{ then } [d, q_k] = 0. \quad (4.25)$$

5 A vanishing theorem

This section is stand-alone and uses its own notation. The goal is a homology vanishing theorem, Theorem 15, which is used as a lemma in Section 6.1.

This commutative diagram contains the objects used in this section, with $n \geq 1$:

$$\begin{array}{ccccccc} \textcolor{red}{U}_n & \hookrightarrow & \textcolor{blue}{T}_n & \twoheadrightarrow & \textcolor{orange}{\Lambda}_n & \twoheadrightarrow & A_n \\ f = f_n \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \textcolor{red}{U} & \hookrightarrow & \textcolor{blue}{T} & \twoheadrightarrow & \textcolor{orange}{\Lambda} & \twoheadrightarrow & A \end{array} \quad (5.1)$$

Here $\textcolor{red}{U}_n = \oplus_{\mu=0}^3 \oplus_{j=1}^n \mathbb{C}e_j^\mu$ is the $4n$ -dimensional vector space with basis elements e_j^μ ; $\textcolor{blue}{T}_n$ is the tensor algebra generated by $\textcolor{red}{U}_n$, with unit, graded so that the e_j^μ are in degree 1; $\textcolor{orange}{\Lambda}_n = \textcolor{blue}{T}_n/(v^2 \text{ with } v \in \textcolor{red}{U}_n)$ is the exterior algebra; and A_n is the quotient algebra

$$A_n = \textcolor{orange}{\Lambda}_n / (e_j^0 e_j^1 - ie_j^2 e_j^3, \quad e_j^0 e_j^2 - ie_j^3 e_j^1, \quad e_j^0 e_j^3 - ie_j^1 e_j^2 \text{ with } j = 1 \dots n)$$

All these algebras are graded. Set $\textcolor{red}{U} = \textcolor{red}{U}_1$ and for its basis we always use the shorthand $e^\mu = e_1^\mu$. Set $\textcolor{blue}{T} = \textcolor{blue}{T}_1$, $\textcolor{orange}{\Lambda} = \textcolor{orange}{\Lambda}_1$, $A = A_1$. Let $f = f_n$ be the \mathbb{C} -linear map

$$f : \textcolor{red}{U}_n \rightarrow \textcolor{red}{U}, \quad e_j^\mu \mapsto e^\mu \quad (5.2)$$

The other vertical arrows in (5.1) are algebra maps induced by f . The Hilbert series⁶¹ are:

$$\textcolor{blue}{T}_n : (1 - 4nt)^{-1} \quad \textcolor{orange}{\Lambda}_n : (1 + t)^{4n} = (1 + 4t + 6t^2 + 4t^3 + t^4)^n \quad A_n : (1 + 4t + 3t^2)^n$$

Remark 3 (Algebra isomorphisms). For $n \geq 2$ one has⁶² $\textcolor{blue}{T}_n \not\simeq \textcolor{blue}{T}^{\otimes n}$, in fact $\textcolor{blue}{T}_n$ is much bigger. By contrast $\textcolor{orange}{\Lambda}_n \simeq \textcolor{orange}{\Lambda}^{\otimes n}$, and this algebra isomorphism can be written down explicitly using Koszul signs in a standard way. It induces an isomorphism $A_n \simeq A^{\otimes n}$.

⁶¹Formal power series $\sum_{k \geq 0} d_k t^k$ where d_k is the \mathbb{C} -dimension of the subspace of degree k .

⁶²If A, B are unital graded associative \mathbb{C} -algebras, commutative or not, then the tensor product of unital associative \mathbb{C} -algebras is the vector space $A \otimes_{\mathbb{C}} B$ with product $(a \otimes b)(a' \otimes b') = (-1)^{ba'}(aa') \otimes (bb')$.

Given a graded left A -module M and a graded left A_n -module M_n , define

$$X^{k,\ell} = S^k U_n^* \otimes \text{Hom}_{\mathbb{C}}^\ell(M_n, M) \quad (5.3)$$

where $S^k U_n^*$ are the homogeneous polynomials of degree k on U_n ; and Hom^ℓ are the maps of degree ℓ . For $\phi \in X^{k,\ell}$, denote its evaluation at $v \in U_n$ by $\phi_v \in \text{Hom}_{\mathbb{C}}^\ell(M_n, M)$. Define

$$d^{k,\ell} : X^{k,\ell} \rightarrow X^{k+1,\ell+1}, \quad \phi \mapsto \psi \quad \text{where} \quad \psi_v(m) = f(v)\phi_v(m) - (-1)^k \phi_v(vm)$$

for all $m \in M_n$; here vm is A_n -module multiplication where $v \in U_n$ also stands for the corresponding element of A_n via (5.1); similarly $f(v)\phi_v(m)$ is A -module multiplication. This is well-defined: $v \mapsto \psi_v$ is a polynomial homogeneous of degree $k+1$; and $\psi_v \in \text{Hom}^{\ell+1}$.

Lemma 14. *For every integer ℓ , the following sequence is a complex:*

$$C_\ell : 0 \rightarrow X^{0,\ell} \xrightarrow{d^{0,\ell}} X^{1,\ell+1} \xrightarrow{d^{1,\ell+1}} X^{2,\ell+2} \xrightarrow{d^{2,\ell+2}} \dots$$

Its homologies will be denoted $H^0(C_\ell)$, $H^1(C_\ell)$, and so forth.

Proof. This calculation uses $f(v)^2 = 0 \in A$ and $v^2 = 0 \in A_n$:

$$\begin{aligned} (d^{k+1,\ell+1} d^{k,\ell} \phi)_v(m) &= f(v)(d^{k,\ell} \phi)_v(m) - (-1)^{k+1} (d^{k,\ell} \phi)_v(vm) \\ &= f(v)(f(v)\phi_v(m) - (-1)^k \phi_v(vm)) - (-1)^{k+1} (f(v)\phi_v(vm) - (-1)^k \phi_v(v^2 m)) = 0 \end{aligned}$$

□

Theorem 15 (A vanishing theorem for H^0 and H^1). *In both Case 1 and Case 2 below one has: If $\ell < 0$ then $H^0(C_\ell) = 0$; and if $\ell + 1 < 0$ then $H^1(C_\ell) = 0$.*

- Case 1: $M = Ag$ and $M_n = A_n g$ where g is a generator of degree 0⁶³.
- Case 2: $M = (Ag_1 \oplus Ag_2 \oplus Ag_3)/S$ and $M_n = (A_n g_1 \oplus A_n g_2 \oplus A_n g_3)/S_n$ where g_1, g_2, g_3 are generators of degree 0; S is the graded left A -submodule generated by

$$\begin{array}{cccc} e^1 g_1 - e^2 g_2 & e^1 g_1 - e^3 g_3 & e^1 g_2 + e^2 g_1 & e^2 g_3 + e^3 g_2 \\ e^3 g_1 + e^1 g_3 & e^0 g_1 + i e^2 g_3 & e^0 g_2 + i e^3 g_1 & e^0 g_3 + i e^1 g_2 \end{array} \quad (5.4)$$

and S_n is the graded left A_n -submodule generated by the same eight relations but with each occurrence of e^μ in (5.4) replaced by e_1^μ for $\mu = 0, 1, 2, 3$.

Remark 4. In Case 2, M is isomorphic as an Λ -module to the Λ -submodule of Λ generated by $e^0 e^1 + i e^2 e^3$, $e^0 e^2 + i e^3 e^1$, $e^0 e^3 + i e^1 e^2$ in the role of g_1, g_2, g_3 respectively, which one can check is also an A -module. The gradings match after a degree shift by 2.

⁶³So this is the case of two free modules of rank one.

The Hilbert series are as follows:

$$\begin{array}{ll} \text{Case 1:} & M : 1 + 4t + 3t^2 \quad M_n : (1 + 4t + 3t^2)^n \\ \text{Case 2:} & M : 3 + 4t + t^2 \quad M_n : (3 + 4t + t^2)(1 + 4t + 3t^2)^{n-1} \end{array}$$

As modules, M and M_n are finitely generated by degree zero elements.

Remark 5 (The multilinear perspective). This is a continuation of Remark 3. Denoting by \otimes the \mathbb{C} -tensor product as before, there are canonical isomorphisms

$$\begin{array}{ll} \text{Case 1:} & M \simeq E^0 \quad M_n \simeq M^{\otimes n} \simeq (E^0)^{\otimes n} \\ \text{Case 2:} & M \simeq E^1 \quad M_n \simeq M \otimes (Ag)^{\otimes(n-1)} \simeq E^1 \otimes (E^0)^{\otimes(n-1)} \end{array}$$

where $E^r = E^{r,0} \oplus E^{r,1} \oplus E^{r,2}$ with $r = 0, 1$ are the graded modules given, in the notation of Section 3, by $E^{0,0} = \Lambda^0$, $E^{0,1} = \Lambda^1$, $E^{0,2} = \Lambda_+^2$ and $E^{1,0} = \Lambda_+^2$, $E^{1,1} = \Lambda^3$, $E^{1,2} = \Lambda^4$. So elements of $\text{Hom}_{\mathbb{C}}(M_n, M)$ may equivalently be viewed as multilinear maps with n slots, and elements of $X^{k,\ell}$ as multilinear maps depending polynomially on a parameter in U_n . One can translate the definition of C_ℓ and its differential to this perspective.

Remark 6. This homology could also be computed using the BGG correspondence between graded Λ_n -modules and linear SU_n^* -complexes; see especially [15, Theorem 7.8]⁶⁴. To apply this here, one would endow $\text{Hom}_{\mathbb{C}}(M_n, M)$ with a Λ_n -module structure⁶⁵.

Proof. For $H^0(C_\ell) = 0$, we must show that if $\phi \in X^{0,\ell}$ and $d^{0,\ell}\phi = 0$ then $\phi = 0$. Since $v \mapsto \phi_v$ is a constant map, write $\phi_v = \phi$. Then $d\phi = 0$ is equivalent to $\phi(vm) = f(v)\phi(m)$ for all $v \in U_n$, $m \in M_n$. So to prove $\phi = 0$ it suffices to prove that ϕ annihilates the module generators of M_n , that is, to prove that $\phi(g)$ respectively $\phi(g_1), \phi(g_2), \phi(g_3)$ are zero. This holds because these elements of M have degree $\ell < 0$, but M is zero in negative degree.

For $H^1(C_\ell) = 0$, given $\psi \in X^{1,\ell+1}$ with $d^{1,\ell+1}\psi = 0$, we must show there exists⁶⁶ a $\phi \in X^{0,\ell}$ with $d^{0,\ell}\phi = -\psi$; the minus sign is for convenience. Here $d\psi = 0$ is equivalent to

$$f(v)\psi_v(m) + \psi_v(vm) = 0 \tag{5.5}$$

⁶⁴Summary: Let U be a vector space, $\dim U < \infty$; let $\Lambda = \Lambda U$ and $S = SU^*$; let $\mathfrak{m} \subseteq \Lambda$ be the unique maximal ideal; let P be a graded Λ -module, $\dim P < \infty$; let P^* be the dual vector space which is canonically a Λ -module; let $F \rightarrow P^* \rightarrow 0$ be a free resolution of P^* as a Λ -module. A canonical differential on $S \otimes_{\mathbb{C}} P$ is given by acting from the left with $\mathbb{1} \in \text{End}(U) \simeq U^* \otimes U$. Then there are canonical isomorphisms

$$H(F/\mathfrak{m}F) \simeq \text{Tor}^\Lambda(P^*, \Lambda/\mathfrak{m}) \simeq H(S \otimes_{\mathbb{C}} P)^*$$

of bigraded vector spaces; the bigradings are linear combinations of the homological and module degrees detailed in [15, Theorem 7.8]. This is based on the two ways of computing Tor : the right \simeq comes from using the minimal free resolution of Λ/\mathfrak{m} in the proof of [15, Theorem 7.8]; the left \simeq is from the definition of Tor using a free resolution of its left argument. If F is minimal then $H(F/\mathfrak{m}F) \simeq F/\mathfrak{m}F$. So the graded Betti numbers of P^* , which can be read off from any minimal free resolution, determine the dimensions of the individual parts of the bigraded vector space $H(S \otimes_{\mathbb{C}} P)$. No attempt was made to apply this.

⁶⁵Define a Λ_n -module structure on $\text{Hom}_{\mathbb{C}}(M_n, M)$ by $(v \cdot \phi)(m) = f(v)\phi(m) - (-1)^\ell \phi(vm)$. Then define $D^{k,\ell} : X^{k,\ell} \rightarrow X^{k+1,\ell+1}$, $\phi \mapsto \psi$ by $\psi_v(m) = f(v)\phi_v(m) - (-1)^\ell \phi_v(vm)$ or, using the module structure, $\psi_v = v \cdot \phi_v$. The $D^{k,\ell}$ are equivalent to the $d^{k,\ell}$. Define involutions $i^{k,\ell} \in \text{End}(X^{k,\ell})$ by $(i^{k,\ell}\phi)_v(m) = \phi_v(\sigma_{k+\ell}m)$, where $\sigma_p \in \text{End}(M_n)$ is given by $\sigma_p(m) = (-1)^{pm}m$, then $i^{k+1,\ell+1} \circ d^{k,\ell} \circ i^{k,\ell} = D^{k,\ell}$.

⁶⁶By $H^0(C_\ell) = 0$ for $\ell < 0$, hence for $\ell + 1 < 0$: If ϕ exists then it is unique.

for all $v \in \mathcal{U}_n$ and $m \in M_n$. Since $v \mapsto \psi_v$ is a linear map, $d\psi = 0$ is equivalent to

$$f(v)\psi_w(m) + f(w)\psi_v(m) + \psi_v(wm) + \psi_w(vm) = 0 \quad (5.6)$$

for all $v, w \in \mathcal{U}_n$ and $m \in M_n$. In the following, G stands for g in Case 1 respectively for either of g_1, g_2, g_3 in Case 2. Note that $\psi_v(G) = 0$ for all v because it has degree $\ell + 1 < 0$. We now define ϕ in several steps. First define a map ϕ^T on free T_n -modules:

$$\begin{aligned} \text{Case 1:} \quad & \phi^T \in \text{Hom}_{\mathbb{C}}^{\ell}(T_n g, M) \\ \text{Case 2:} \quad & \phi^T \in \text{Hom}_{\mathbb{C}}^{\ell}(T_n g_1 \oplus T_n g_2 \oplus T_n g_3, M) \end{aligned}$$

where, recursively in the degree $N \geq 0$ of the input,

$$\begin{aligned} \phi^T(v_N \cdots v_1 G) &= \psi_{v_N}(v_{N-1} \cdots v_1 G) + f(v_N)\phi^T(v_{N-1} \cdots v_1 G) \\ \phi^T(G) &= 0 \end{aligned} \quad (5.7)$$

for all $v_1, \dots, v_N \in \mathcal{U}_n$ and $G = g$ or $G = g_1, g_2, g_3$. Note that ϕ^T is well-defined and has degree ℓ as claimed. The recursion (5.7) means (here the final term is actually zero):

$$\begin{aligned} \phi^T(v_N \cdots v_1 G) &= \psi_{v_N}(v_{N-1} \cdots v_1 G) + f(v_N)\psi_{v_{N-1}}(v_{N-2} \cdots v_1 G) + \\ &+ \dots + f(v_N) \cdots f(v_3)\psi_{v_2}(v_1 G) + f(v_N) \cdots f(v_2)\psi_{v_1}(G) \end{aligned} \quad (5.8)$$

Let $I \subseteq T_n$ be the two-sided ideal generated by all v^2 with $v \in \mathcal{U}_n$. Recall $T_n/I = \Lambda_n$. Claim: $\phi^T(IG) = 0$. It suffices to check that $\phi^T(v_N \cdots v_1 G) = 0$ when $v_{p+1} = v_p = v$ for p with $0 \leq p \leq N-1$; here $N \geq 2$. Since $\psi_w(IG) = 0$ for all $w \in \mathcal{U}_n$ because $IG = 0 \subseteq M_n$, and since $f(v)^2 = 0 \in A$, it suffices by (5.8) to check that $\psi_v(v-) + f(v)\psi_v(-) = 0$, which follows from (5.5). Therefore $\phi^T(IG) = 0$, so ϕ^T descends to a map

$$\begin{aligned} \text{Case 1:} \quad & \phi^{\Lambda} \in \text{Hom}_{\mathbb{C}}^{\ell}(\Lambda_n g, M) \\ \text{Case 2:} \quad & \phi^{\Lambda} \in \text{Hom}_{\mathbb{C}}^{\ell}(\Lambda_n g_1 \oplus \Lambda_n g_2 \oplus \Lambda_n g_3, M) \end{aligned}$$

Denote by $J \subseteq \Lambda_n$ the left ideal generated by all

$$E_j^1 = e_j^0 e_j^1 - i e_j^2 e_j^3 \quad E_j^2 = e_j^0 e_j^2 - i e_j^3 e_j^1 \quad E_j^3 = e_j^0 e_j^3 - i e_j^1 e_j^2$$

with $j = 1 \dots n$. Since Λ_n is graded commutative, J is a two-sided ideal, and $A_n = \Lambda_n/J$. Claim: $\phi^{\Lambda}(JG) = 0$. To see this, it suffices to check that $\phi^{\Lambda}(v_N \cdots v_3 E_j^1 G) = 0$; here $N \geq 2$. (E_j^2, E_j^3 are analogous.) Since $\psi_w(JG) = 0$ for all $w \in \mathcal{U}_n$ because $JG = 0 \subseteq M_n$, and since $\psi_w(G) = 0$, it suffices by (5.8) to check that⁶⁷

$$\psi_{e_j^0}(e_j^1 G) = i\psi_{e_j^2}(e_j^3 G) \quad \psi_{e_j^0}(e_j^2 G) = i\psi_{e_j^3}(e_j^1 G) \quad \psi_{e_j^0}(e_j^3 G) = i\psi_{e_j^1}(e_j^2 G) \quad (5.9)$$

If $\ell + 2 < 0$ then all terms in (5.9) are separately zero; the proof of (5.9) in the remaining case $\ell + 2 = 0$ is given after this proof. Therefore $\phi^{\Lambda}(JG) = 0$, so ϕ^{Λ} descends to a map

$$\begin{aligned} \text{Case 1:} \quad & \phi^A \in \text{Hom}_{\mathbb{C}}^{\ell}(A_n g, M) \\ \text{Case 2:} \quad & \phi^A \in \text{Hom}_{\mathbb{C}}^{\ell}(A_n g_1 \oplus A_n g_2 \oplus A_n g_3, M) \end{aligned}$$

⁶⁷In each of these three claimed equations, G stands for the same element on both sides.

In Case 1 we have $M_n = A_n g$, so set $\phi = \phi^{A68}$. In Case 2 recall the eight generators of the graded left A_n -submodule $S_n \subseteq A_n g_1 \oplus A_n g_2 \oplus A_n g_3$. Denote them by

$$P_1 = e_1^1 g_1 - e_1^2 g_2 \quad \dots \quad P_8 = e_1^0 g_3 + i e_1^1 g_2$$

Claim: $\phi^A(S_n) = 0$. It suffices to check that $\phi^A(v_N \cdots v_2 P_1) = 0$; here $N \geq 1$. (P_2, \dots, P_8 are analogous.) Since $\psi_w(S_n) = 0$ for all $w \in U_n$ because $S_n = 0 \subseteq M_n$, it suffices by (5.8) to check $\psi_{e_1^1}(g_1) - \psi_{e_1^2}(g_2) = 0$, which is true because the ψ_w annihilate all generators for degree reasons. Hence $\phi^A(S_n) = 0$ and ϕ^A descends to a map

$$\text{Case 2:} \quad \phi \in \text{Hom}_{\mathbb{C}}^{\ell}(M_n, M) \simeq X^{0, \ell}$$

We are ready to check that $d^{0, \ell} \phi = -\psi$ which is equivalent to $\phi(vm) = \psi_v(m) + f(v)\phi(m)$ for all $v \in U_n$ and $m \in M_n$. This follows from the recursion (5.7) which is satisfied by ϕ^T by definition, and which is therefore also satisfied by ϕ . \square

Proof (of equation (5.9)). The case of interest is $\ell + 2 = 0$, but only assume $\ell + 1 < 0$ for the time being. Two observations:

- $\psi_v(wG) = -\psi_w(vG)$ for all $v, w \in U_n$, by (5.6).
- $\psi_v(e_j^0 e_j^1 G) = i \psi_v(e_j^2 e_j^3 G)$ for all $v \in U_n$, because $e_j^0 e_j^1 = i e_j^2 e_j^3$ in A_n .

Analogous identities are obtained by cyclically permuting the indices 1, 2, 3.

For fixed j and G abbreviate the following elements of M (recall (5.2)):

$$r_{\mu\nu\rho} = \psi_{e_j^\mu}(e_j^\nu e_j^\rho G) \quad s_{\mu\nu\rho} = e_j^\mu \psi_{e_j^\nu}(e_j^\rho G)$$

where $\mu, \nu, \rho = 0, 1, 2, 3$. By the two observations above and by (5.6):

$$\begin{aligned} r_{\mu\nu\rho} &= -r_{\mu\rho\nu} & r_{\mu 01} &= i r_{\mu 23} \\ s_{\mu\nu\rho} + s_{\nu\mu\rho} + r_{\mu\nu\rho} + r_{\nu\mu\rho} &= 0 & r_{\mu 02} &= i r_{\mu 31} \\ s_{\mu\nu\rho} &= -s_{\mu\rho\nu} & r_{\mu 03} &= i r_{\mu 12} \end{aligned}$$

Eliminating all $r_{\mu\nu\rho}$ from this finite linear system, one finds $(s_{001} - i s_{023}) + i(s_{302} - i s_{331}) = 0$ and $(s_{301} - i s_{323}) + i(s_{002} - i s_{031}) = 0$ and $s_{101} - i s_{123} = s_{202} - i s_{231}$ and analogous identities obtained by cyclically permuting 1, 2, 3. It is convenient to write this in matrix form, $WZ = 0$. Here W is an 8×3 matrix with entries in A of degree one, viewed as a degree one map $M^3 \rightarrow M^8$, and $Z \in M^3$ is a vector:

$$W = \begin{pmatrix} e^0 & i e^3 & 0 \\ e^3 & i e^0 & 0 \\ 0 & e^0 & i e^1 \\ 0 & e^1 & i e^0 \\ i e^2 & 0 & e^0 \\ i e^0 & 0 & e^2 \\ e^1 & -e^2 & 0 \\ 0 & e^2 & -e^3 \end{pmatrix} \quad Z = \begin{pmatrix} \psi_{e_j^0}(e_j^1 G) - i \psi_{e_j^2}(e_j^3 G) \\ \psi_{e_j^0}(e_j^2 G) - i \psi_{e_j^3}(e_j^1 G) \\ \psi_{e_j^0}(e_j^3 G) - i \psi_{e_j^1}(e_j^2 G) \end{pmatrix}$$

⁶⁸This does not conclude Case 1, one must still prove $d\phi = -\psi$.

If $\ell + 2 < 0$ then we have $Z = 0$ for degree reasons. If $\ell + 2 = 0$ then $Z \in (\mathbb{C}g)^3$ in Case 1; $Z \in (\mathbb{C}g_1 \oplus \mathbb{C}g_2 \oplus \mathbb{C}g_3)^3$ in Case 2. But one checks that W is injective as a map⁶⁹

$$\begin{aligned} \text{Case 1:} & \quad (\mathbb{C}g)^3 \rightarrow M^8 \\ \text{Case 2:} & \quad (\mathbb{C}g_1 \oplus \mathbb{C}g_2 \oplus \mathbb{C}g_3)^3 \rightarrow M^8 \end{aligned}$$

Hence $WZ = 0$ implies $Z = 0$ which is (5.9). \square

6 BV_∞^\square -algebra for Yang-Mills

6.1 Proof of Theorem 3

Here we construct the BV_∞^\square -algebra claimed in Theorem 3. (Then, Theorem 1 follows by writing out two axioms, namely, in the notation of Lemma 13: the 3-ary identity $b_3 = 0$ to get (1.5); the 4-ary identity $c_4 = 0$ to get (1.6).) The conventions in Remarks 1 and 2 are in force. This proof is organized around Lemma 13. Use the following data:

data in Section 4	Yang-Mills in this section
V	$\mathfrak{a}[2]$, see Section 3
H	$\mathbb{C}[\partial_0, \dots, \partial_3]$
Q	$\square = \eta^{\mu\nu} \partial_\mu \partial_\nu$

So $\mathfrak{a} = V[-2]$ ⁷⁰, that is $V^i = \mathfrak{a}^{i+2}$, therefore $V = V^{-2} \oplus \dots \oplus V^1$. The coproduct of H is given by $\partial_\mu \mapsto \partial_\mu \otimes \mathbb{1} + \mathbb{1} \otimes \partial_\mu$. The H -module structure on \mathfrak{a} is given by differentiation, the one on $\mathfrak{a}^{\otimes n}$ is induced by the coproduct. Note that $Q = \square$ satisfies (4.4).

The first three operations. Recall that, by assumption of Theorem 3, an h satisfying the requirements in Section 1 is given, in particular (1.3) holds.

- The differential in (1.1) maps $d : \mathfrak{a}^i \rightarrow \mathfrak{a}^{i+1}$, equivalently $V^i \rightarrow V^{i+1}$, so its dual maps $(V^*)^i \rightarrow (V^*)^{i+1}$. By (4.2) it defines a derivation also denoted $d \in \text{Der}^1(GV^*)^H$. Analogously, $h : \mathfrak{a}^i \rightarrow \mathfrak{a}^{i-1}$ in (1.3) gives $h \in \text{Der}^{-1}(GV^*)^H$. Set $\mu_1 = d + h$.
- The gca product maps $\mathfrak{a}^i \otimes \mathfrak{a}^j \rightarrow \mathfrak{a}^{i+j}$, or $V^i \otimes V^j \rightarrow V^{i+j+2}$, so formally its dual maps $(V^*)^i \rightarrow \sum_{i_1+i_2=i+2} (V^*)^{i_1} \otimes (V^*)^{i_2}$. It is graded commutative, so defines a derivation $\theta_2 \in \text{Der}^2(GV^*)^H$ by (4.2) consistent with (4.21). Set $\nu_2 = d_\alpha \theta_2 \in \text{Der}^1(GV^*)^H$.

Some assumptions in Lemma 13 follow: $[d, d] = [h, h] = 0$ and $[d, h] = K$ using (1.3); and $b_2 = c_3 = 0$ because $[d, \nu_2] = [\nu_2, \nu_2] = 0$ by the dgca Leibniz rule and associativity.

Definition 16 (Local operators). *Denote by \mathcal{C}_n the space of all maps $\mathfrak{a}^{\otimes n} \rightarrow \mathfrak{a}$ that are $C^\infty(\mathbb{R}^4)$ -linear in all n arguments and invariant under translations on \mathbb{R}^4 . Denote*

$$\mathcal{D}_n = \left\{ \sum_{j=1}^n c_j^\mu (\mathbb{1}^{\otimes(j-1)} \otimes \partial_\mu \otimes \mathbb{1}^{\otimes(n-j)}) : \mathfrak{a}^{\otimes n} \rightarrow \mathfrak{a} \mid c_j^\mu \in \mathcal{C}_n \right\} \quad (6.1)$$

⁶⁹The dimensions are as follows. Since W has degree one, let us only take into account the degree one subspace of M^8 which is $\simeq (\mathbb{C}^4)^8 \simeq \mathbb{C}^{32}$ in both cases. With this understanding, the injectivity claim is for \mathbb{C} -linear maps of dimensions $\mathbb{C}^3 \rightarrow \mathbb{C}^{32}$ respectively $\mathbb{C}^9 \rightarrow \mathbb{C}^{32}$.

⁷⁰For a graded vector space V , one denotes by $V[k]^i = V^{i+k}$ the shifted space.

a class of translation invariant, homogeneous first order partial differential operators. Note that \mathcal{C}_n and \mathcal{D}_n are finite-dimensional. All elements of \mathcal{C}_n and \mathcal{D}_n are H -equivariant.

Row decomposition. Denote by $\mathfrak{a}^i = \oplus_{r=0,1} \mathfrak{a}^{r,i}$ the decomposition of the complex (1.1) into the upper row, $r = 0$, and the lower row, $r = 1$. For instance, $\mathfrak{a}^{1,2} = \Omega^3$. This leads to decompositions $d = d^0 + d^1$ and $h = h^0 + h^1$ where

$$\begin{aligned} d^s &: \mathfrak{a}^{r,i} \rightarrow \mathfrak{a}^{r+s-1,i+1} \\ h^s &: \mathfrak{a}^{r,i} \rightarrow \mathfrak{a}^{r+s-1,i-1} \end{aligned}$$

for $s = 0, 1$. Here $d^0, h^0 \in \mathcal{C}_1$ and $d^1, h^1 \in \mathcal{D}_1$. In particular, d^1 maps the upper (lower) row of \mathfrak{a} to the upper (lower) row. In the notation of Section 3,

$$d^1 = \partial_\mu \otimes e^\mu : \mathfrak{a}^{r,i} \rightarrow \mathfrak{a}^{r,i+1} \quad (6.2)$$

here viewed as an element $d^1 \in \text{Der}^1(GV^*)^H$ with $[d^1, d^1] = 0$.

The notation θ_n . Below we inductively construct operations θ_n with $n \geq 2$. By abuse of notation, the operation θ_n is simultaneously (the correspondence is via (4.2)):

- A graded symmetric and H -equivariant map $\theta_n : \mathfrak{a}^{\otimes n} \rightarrow \mathfrak{a}$ of degree $4 - 2n$. Equivalently, a map $V^{\otimes n} \rightarrow V$ or formally $V^* \rightarrow (V^*)^{\otimes n}$ of degree 2.
- A derivation $\theta_n \in \text{Der}^2(GV^*)^H$, as in Lemma 13.

Graded symmetry means

$$\theta_n(\dots, x, y, \dots) = (-1)^{xy} \theta_n(\dots, y, x, \dots) \quad (6.3)$$

The inductive construction will be such that

$$\theta_n : \mathfrak{a}^{r_1, i_1} \otimes \dots \otimes \mathfrak{a}^{r_n, i_n} \rightarrow \mathfrak{a}^{r_1 + \dots + r_n, i_1 + \dots + i_n + 4 - 2n} \quad \text{and} \quad \theta_n \in \mathcal{C}_n \quad (6.4)$$

The operation θ_2 defined above is the gca product in \mathfrak{a} , and is consistent with (6.4). The operation θ_3 constructed below is the one that Theorem 1 refers to.

Outline of the induction. Below we inductively construct $\theta_3, \theta_4, \dots$ so that (in the notation of Lemma 13) $b_3, c_4, b_4, c_5, b_5, \dots$ vanish and so that (6.3), (6.4) hold. More precisely, using Part B of Lemma 13, one proceeds as follows, recalling $b_n = -[d, \theta_n] + q_n$: Use $[d, q_3] = 0$ to construct θ_3 so that $[d, \theta_3] = q_3$; use $[d, c_4] = 0$ to show that $c_4 = 0$; use $[d, q_4] = 0$ to construct θ_4 so that $[d, \theta_4] = q_4$; use $[d, c_5] = 0$ to show that $c_5 = 0$; etc. At each step, the construction of θ_n uses the homology vanishing theorem, Theorem 15.

Properties of q_n and c_n known inductively. The q_n and c_n are defined once $\theta_2, \dots, \theta_{n-1}$ are defined. Similar to the notation θ_n , here too we abuse notation, using (4.2) and (4.23): b_n and q_n are formally also graded symmetric maps $V^* \rightarrow (V^*)^{\otimes n}$ of degree 3; and c_n is also a map $V^* \rightarrow (V^*)^{\otimes n}$ of degree 4 that has some symmetry in the first two respectively

the remaining $n - 2$ factors. Suppose we are at a stage of the induction where $\theta_2, \dots, \theta_{n-1}$ have been constructed and satisfy (6.4), then one can check⁷¹ that the following is true:

$$c_n : \quad \mathfrak{a}^{r_1, i_1} \otimes \dots \otimes \mathfrak{a}^{r_n, i_n} \rightarrow \mathfrak{a}^{r_1 + \dots + r_n, i_1 + \dots + i_n + 6 - 2n} \quad (6.5a)$$

$$q_n^s : \quad \mathfrak{a}^{r_1, i_1} \otimes \dots \otimes \mathfrak{a}^{r_n, i_n} \rightarrow \mathfrak{a}^{r_1 + \dots + r_n + s - 1, i_1 + \dots + i_n + 5 - 2n} \quad (6.5b)$$

where $q_n = q_n^0 + q_n^1$ with

$$c_n, q_n^0 \in \mathcal{C}_n \quad q_n^1 \in \mathcal{D}_n \quad (6.6)$$

Induction step to prove $c_n = 0$, for $n \geq 4$. This comes after $\theta_2, \dots, \theta_{n-1}$ have been constructed and $b_2, c_3, b_3, \dots, c_{n-1}, b_{n-1}$ are known to vanish. (For $b_n = 0$, see below.)

- By Part B of Lemma 13 we have $[d, c_n] = 0$. This has the form

$$dc_n(x_1, \dots, x_n) = \sum_{i=1}^n \pm c_n(x_1, \dots, x_{i-1}, dx_i, x_{i+1}, \dots, x_n)$$

for all $x_1, \dots, x_n \in \mathfrak{a}$. The signs, which may depend on degrees, are irrelevant. By (6.6), $[d, c_n]$ is a first order partial differential operator. The homogeneous first order part must separately vanish, $[d^1, c_n] = 0$. Using (6.2) and $c_n \in \mathcal{C}_n$ this implies

$$e^\mu c_n(x_1, \dots, x_n) = \pm c_n(x_1, \dots, x_{i-1}, e^\mu x_i, x_{i+1}, \dots, x_n) \quad (6.7)$$

separately for all $\mu = 0, \dots, 3$ and $i = 1 \dots n$.

- Denote by $c_n^{r_1 \dots r_n}$ the map (6.5a), meaning the restriction of c_n where the i -th argument is in row r_i of \mathfrak{a} . The identities (6.7) hold separately for each $c_n^{r_1 \dots r_n}$ because e^μ maps the upper (lower) row of \mathfrak{a} to the upper (lower) row of \mathfrak{a} . Hence $c_n^{r_1 \dots r_n}$ vanishes iff its restriction to the generators $\oplus_{r=0,1} \mathfrak{a}^{r,r}$, meaning to the leftmost term in each row in (1.1), vanishes. By (6.5a), this restriction maps

$$\mathfrak{a}^{r_1, r_1} \otimes \dots \otimes \mathfrak{a}^{r_n, r_n} \rightarrow \mathfrak{a}^{r_1 + \dots + r_n, r_1 + \dots + r_n + 6 - 2n}$$

Since $6 - 2n < 0$, the space on the right is zero, so $c_n^{r_1 \dots r_n} = 0$, so $c_n = 0$.

Induction step to construct θ_n and to prove $b_n = 0$, for $n \geq 3$. This is after $\theta_2, \dots, \theta_{n-1}$ have been constructed and $b_2, c_3, b_3, \dots, c_{n-1}, b_{n-1}, c_n$ are known to vanish.

- By Part B of Lemma 13 we have $[d, q_n] = 0$. This means that

$$dq_n(x_1, \dots, x_n) + \sum_{i=1}^n (-1)^{x_1 + \dots + x_{i-1}} q_n(x_1, \dots, x_{i-1}, dx_i, x_{i+1}, \dots, x_n) = 0$$

Recall that $q_n : \mathfrak{a}^{\otimes n} \rightarrow \mathfrak{a}$ is a graded symmetric map of degree $5 - 2n$. By (6.6), $[d, q_n]$ is a second order partial differential operator. The homogeneous second order part must separately vanish, $[d^1, q_n^1] = 0$. Since $q_n^1 \in \mathcal{D}_n$ one has

$$q_n^1 = \sum_{j=1}^n u_j^\mu (\mathbb{1}^{\otimes(j-1)} \otimes \partial_\mu \otimes \mathbb{1}^{\otimes(n-j)})$$

⁷¹Using (4.22), note that q_n is a sum of composition of three operations, h and θ_m and θ_{n+1-m} with $1 < m < n$; and c_n is a sum of composition of two operations, θ_m and θ_{n+1-m} with $1 < m < n$.

for $u_j^\mu \in \mathcal{C}_n$. The $u_j^\mu : \mathfrak{a}^{\otimes n} \rightarrow \mathfrak{a}$ have degree $5 - 2n$ but may not be graded symmetric. Now $[d^1, q_n^1] = 0$ implies, separately for all $\mu, \nu = 0, \dots, 3$ and $i, j = 1, \dots, n$:

$$\begin{aligned} & e^\mu u_j^\nu(x_1, \dots, x_n) + e^\nu u_i^\mu(x_1, \dots, x_n) \\ & + (-1)^{x_1 + \dots + x_{i-1}} u_j^\nu(x_1, \dots, x_{i-1}, e^\mu x_i, x_{i+1}, \dots, x_n) \\ & + (-1)^{x_1 + \dots + x_{j-1}} u_i^\mu(x_1, \dots, x_{j-1}, e^\nu x_j, x_{j+1}, \dots, x_n) = 0 \end{aligned} \quad (6.8)$$

- Denote by $q_n^{1;r_1 \dots r_n}$ the map (6.5b), meaning the restriction of q_n^1 where the i -th argument is in row r_i of \mathfrak{a} ; let $u_j^{\mu;r_1 \dots r_n}$ be the corresponding piece of u_j^μ ; let $u^{r_1 \dots r_n}$ be the collection of all $4n$ maps ($u_j^{\mu;r_1 \dots r_n}$). Explicitly,

$$q_n^{1;r_1 \dots r_n}, u_j^{\mu;r_1 \dots r_n} : \mathfrak{a}^{r_1, i_1} \otimes \dots \otimes \mathfrak{a}^{r_n, i_n} \rightarrow \mathfrak{a}^{r_1 + \dots + r_n, i_1 + \dots + i_n + 5 - 2n}$$

The (6.8) hold separately for each collection $u^{r_1 \dots r_n}$. Distinguish Case 1 where $r_i = 0$ for all i ; Case 2 where $r_j = 1$ for one j and $r_i = 0$ for $i \neq j$; otherwise $u^{r_1 \dots r_n} = 0$ for degree reasons. The plan is to apply the respective Cases 1 and 2 of Theorem 15. To do this, use $u_j^{\mu;r_1 \dots r_n} \in \mathcal{C}_n$ to encode the collection $u^{r_1 \dots r_n}$ in a linear map

$$\begin{aligned} \psi^{r_1 \dots r_n} : \mathcal{U}_n = \oplus_{\mu, i} \mathbb{C} e_j^\mu & \rightarrow \text{Hom}_{\mathbb{C}}^{5-2n}(E^{r_1} \otimes \dots \otimes E^{r_n}, E^{r_1 + \dots + r_n}) \\ e_j^\mu & \mapsto u_j^{\mu;r_1 \dots r_n} \end{aligned} \quad (6.9)$$

using $\mathfrak{a}^{r, i} = C^\infty(\mathbb{R}^4) \otimes E^{r, i+r}$, where the vector spaces E^r are defined in Remark 5. So $\psi^{r_1 \dots r_n} \in X^{1, \ell+1} = X^{1, 5-2n}$ (see (5.3)) via the isomorphisms in Remarks 3 and 5. The identities (6.8) translate to (5.6) so that $d^{1, \ell+1} \psi^{r_1 \dots r_n} = 0$ with $\ell+1 = 5-2n < 0$.

- Hence $H^1(C_\ell) = H^1(C_{4-2n}) = 0$ in Theorem 15 implies that $\psi^{r_1 \dots r_n} = d^{0, \ell} \phi^{r_1 \dots r_n}$ for some $\phi^{r_1 \dots r_n} \in X^{0, \ell}$. By $H^0(C_\ell) = 0$ in Theorem 15 and $\ell = 4 - 2n < 0$, these $\phi^{r_1 \dots r_n}$ are unique. Translating the $\phi^{r_1 \dots r_n}$ using Remarks 3 and 5, there are unique

$$\theta_n^{r_1 \dots r_n} : \mathfrak{a}^{r_1, i_1} \otimes \dots \otimes \mathfrak{a}^{r_n, i_n} \rightarrow \mathfrak{a}^{r_1 + \dots + r_n, i_1 + \dots + i_n + 4 - 2n}$$

in \mathcal{C}_n with $[d^1, \theta_n^{r_1 \dots r_n}] = q_n^{1;r_1 \dots r_n}$. Denoting by $\theta_n : \mathfrak{a}^{\otimes n} \rightarrow \mathfrak{a}$ the direct sum of all these components, by construction we have $[d^1, \theta_n] = q_n^1$ which explicitly means

$$\begin{aligned} & q_n^1(x_1, \dots, x_n) \\ & = d^1 \theta_n(x_1, \dots, x_n) - \sum_{i=1}^n (-1)^{x_1 + \dots + x_{i-1}} \theta_n(x_1, \dots, x_{i-1}, d^1 x_i, x_{i+1}, \dots, x_n) \end{aligned}$$

Since q_n^1 is graded symmetric, and since θ_n is unique as stated, it follows that θ_n is also graded symmetric, as required. It is also H -equivariant.

- Recall that $b_n = -[d, \theta_n] + q_n$ is a graded symmetric map $\mathfrak{a}^{\otimes n} \rightarrow \mathfrak{a}$ of degree $5 - 2n$. We must show $b_n = 0$. By construction of θ_n , we have $b_n \in \mathcal{C}_n$, so its pieces are

$$b_n^{r_1 \dots r_n} : \mathfrak{a}^{r_1, i_1} \otimes \dots \otimes \mathfrak{a}^{r_n, i_n} \rightarrow \mathfrak{a}^{r_1 + \dots + r_n - 1, i_1 + \dots + i_n - 1 + (6 - 2n)} \quad (6.10)$$

Part B of Lemma 13 implies $[d, b_n] = 0$, and $b_n \in \mathcal{C}_n$ implies $[d^0, b_n] = [d^1, b_n] = 0$. But $[d^1, b_n] = 0$ splits into independent identities $[d^1, b_n^{r_1 \dots r_n}] = 0$. Hence $b_n^{r_1 \dots r_n}$

vanishes iff its restriction to $\oplus_{r=0,1} \mathfrak{a}^{r,r}$ vanishes⁷². If $n > 3$ then $6 - 2n < 0$, so this vanishes for degree reasons by (6.10), so $b_n = 0$. Hereafter, $n = 3$. We show that $[d^1, b_3^{r_1 r_2 r_3}] = 0$ implies $b_3^{r_1 r_2 r_3} = 0$ separately for every triple $r_1 r_2 r_3$. Recall

$$b_3^{r_1 r_2 r_3} : \mathfrak{a}^{r_1, i_1} \otimes \mathfrak{a}^{r_2, i_2} \otimes \mathfrak{a}^{r_3, i_3} \rightarrow \mathfrak{a}^{r_1 + r_2 + r_3 - 1, i_1 + i_2 + i_3 - 1}$$

This can only be nonzero if exactly one or two of the r_i are equal to 1. By graded symmetry, this leaves $u = b_3^{100}$ and $v = b_3^{110}$. By $[d^1, b_3] = 0$ they are determined by their restriction to $\oplus_{r=0,1} \mathfrak{a}^{r,r}$. So $u, v \in \mathcal{C}_3$ are determined by two linear maps,

$$\begin{aligned} U &: \Lambda_+^2 \otimes S^2 \Lambda^0 \rightarrow \Lambda^0 && \text{for } u \text{ restricted to generators} \\ V &: \Lambda^2 \Lambda_+^2 \otimes \Lambda^0 \rightarrow \Lambda_+^2 && \text{for } v \text{ restricted to generators} \end{aligned}$$

Below we use the following bases: $1 \in \mathbb{C} \simeq \Lambda^0$; and $g_1, g_2, g_3 \in \Lambda_+^2$ denote respectively the three elements in (3.2). We also use $u(e^\mu x, y, z) = -e^\mu u(x, y, z)$ and $v(e^\mu x, y, z) = -e^\mu v(x, y, z)$ which follow from $[d^1, b_3] = 0$. Consider now the two cases.

- Set $U_i = u(g_i, 1, 1) \in \mathbb{C}$. Note $e^1 g_1 = e^2 g_2 \in \Lambda^3$, so $u(e^1 g_1, 1, 1) = u(e^2 g_2, 1, 1)$, so $e^1 u(g_1, 1, 1) = e^2 u(g_2, 1, 1)$, so $e^1 U_1 = e^2 U_2 \in \Lambda^1$. But $e^1, e^2 \in \Lambda^1$ are linearly independent, so $U_1 = U_2 = 0$, analogously $U_3 = 0$, so $U = 0$.
- Set $V(g_j, g_k, 1) = \sum_{i=1}^3 V_{jk}^i g_i$, $V_{jk}^i = -V_{kj}^i \in \mathbb{C}$. Note that $g_i g_j = \delta_{ij} g_1 g_1 \in \Lambda^4$. So if $i \neq j$ then $v(g_i g_j, g_k, 1) = 0$, so $g_i v(g_j, g_k, 1) = 0$, so $V_{jk}^i = 0$, all when $i \neq j$. This and antisymmetry in the two lower indices implies $V = 0$.

This concludes the proof of Theorem 3.

6.2 Strict structure via a cobar construction

This section is a continuation of Section 4.2, and it is not specific to Yang-Mills. It contains an explicit cobar construction that associates to a BV_∞^Q -algebra a quasi-isomorphic strict BV_∞^Q -algebra, called a dgBVa^Q below. The main task, relative to standard cobar constructions, is to accommodate the Q -dependent terms in (4.7).

The standard cobar construction [25, 33] associates to a dg coalgebra C without counit⁷³ a dg algebra A without unit⁷⁴. Here A is an appropriate free algebra generated by C . Schematically, if u_C and b_C are the co-unary and co-binary co-operations on C respectively, u_A and b_A the unary and binary operations on A , then:

- u_A is given by an explicit formula in terms of u_C and b_C .
- b_A is the native algebra structure on A .

⁷²By an argument like in the induction step for c_n .

⁷³Or a conilpotent coaugmented dg coalgebra [33]. Then the cokernel of the coaugmentation yields the version without counit. In any event, the cobar construction involves removing the counit.

⁷⁴Or an augmented dg algebra [33]. Then the kernel of the augmentation yields the version without unit.

So all operations on C are packed into the unary one on A . The cobar and bar constructions form an adjunction; they work for various (co)algebraic structures; and they can be used to rectify ∞ -algebraic structures to ∞ -quasi-isomorphic strict structures [24, 25, 33]. The result below is not a full rectification result, because we only establish the quasi-isomorphism for the lowest operations, which suffices for our application⁷⁵.

Notation for the cobar construction. Let V be a graded vector space and H -module, and an element Q is fixed, just as in Section 4.2. Define:

- C is the cofree Gerstenhaber coalgebra without counit cogenerated by V . It is spanned by co-Gerstenhaber words (cf. [26]) in V . It is formally the linear dual of GV^* .
- A is the free Gerstenhaber algebra without unit generated by $C[-2]$ ⁷⁶, not completed. It is spanned by Gerstenhaber words in C . So it is spanned by words-of-words in V .

Both are H -modules and, canonically, $V[-2] \hookrightarrow C[-2] \hookrightarrow A$. Note that C is precisely the object that we avoided working with in Section 4, see Remark 1, but here we must make the switch. The gLa of H -equivariant Gerstenhaber coderivations is denoted $\text{Coder}(C)^H$, its elements are formal duals of the elements of $\text{Der}(GV^*)^H$.

Lemma 17 (Cobar construction). *Suppose $\delta \in \text{Coder}(C)^H$ is a BV_∞^Q -algebra structure on V , see (4.7), with the following additional assumptions⁷⁷:*

- $\delta = \delta^1 + \delta^{-1}$ where $\delta^{\pm 1} \in \text{Coder}^{\pm 1}(C)^H$, so δ only has degree 1 and -1 pieces⁷⁸.
- The only nonzero component of δ^{-1} is its $V \rightarrow V$ component⁷⁹.

Then a cobar construction⁸⁰ yields H -equivariant $d_A \in \text{End}^1(A)$, $h_A \in \text{End}^{-1}(A)$ such that:

- *Strict structure:* A is a strict BV_∞^Q -algebra, or dgBVa^Q , meaning:
 - A is a Gerstenhaber algebra, see the five axioms (4.1).
 - d_A is a derivation for the gca product and h_A is a BV operator, so for all $a, b \in A$:

$$\begin{aligned} d_A(ab) &= d_A(a)b + (-1)^a ad_A(b) \\ [a, b] &= (-1)^a h_A(ab) - (-1)^a h_A(a)b - ah_A(b) \end{aligned} \tag{6.11}$$

- $d_A^2 = 0$ and $d_A h_A + h_A d_A = Q\mathbb{1}$ and $h_A^2 = 0$.

So (A, d_A) is a dgca and (A, h_A) is a BV algebra, in particular h_A is second order.

⁷⁵It is possible that the extension to an ∞ -quasi-isomorphism is immediate, once things are set up, also because there is an explicit candidate for the ∞ -quasi-isomorphism, cf. [25, Proposition 3].

⁷⁶For a graded vector space C , one denotes by $C[k]^i = C^{i+k}$ the shifted space.

⁷⁷These assumptions hold for the algebra in Theorem 3.

⁷⁸The notation does not distinguish $\delta^{\pm 1} \in \text{Coder}^{\pm 1}(C)^H$ and its formal dual $\delta^{\pm 1} \in \text{Der}^{\pm 1}(GV^*)^H$.

⁷⁹The $V \rightarrow V$ component is the formal dual of the $V^* \rightarrow V^*$ component.

⁸⁰The cobar construction is applied to (C, δ) here interpreted as a co-dgBVa^Q . See the proof.

- *Quasi-isomorphism:* Denote by $I : V[-2] \hookrightarrow A$ the canonical degree zero map, and abbreviate the differentials⁸¹ $d = \delta^1|_V \in \text{End}^1(V)$ and $h = \delta^{-1}|_V \in \text{End}^{-1}(V)$. There exist H -equivariant maps P of degree zero⁸², Z of degree -1 , as in

$$V[-2] \begin{array}{c} \xrightarrow{I} \\ \xleftarrow{P} \end{array} A \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} Z \quad (6.12)$$

such that $d_A I = Id$ and $h_A I = Ih$ and $Pd_A = dP$ and $Ph_A = hP$ and $PI = \mathbb{1}$ and $IP = \mathbb{1} - d_A Z - Z d_A$ and $ZI = PZ = Z^2 = 0$. That is:

- I and P are chain maps using the differentials d and d_A respectively, and they are quasi-isomorphisms as witnessed by the contraction (6.12).
- I and P are chain maps using the differentials h and h_A respectively.

Furthermore, denoting by $\Theta : A \otimes A \rightarrow A$ the gca product on A , by $\theta : V \otimes V \rightarrow V$ the binary product of δ^1 as a \mathbf{C}_∞ -structure on $V[-2]$ ⁸³, one has

$$\theta = P \circ \Theta \circ I^{\otimes 2} \quad (6.13)$$

The construction of this dgBVa^Q , and of the contraction (6.12), is functorial.

Proof. Denote by $\Delta^0, \Delta^{-1} : C \rightarrow C \otimes C$ the native coproducts on C of degree 0 and -1 respectively, satisfying axioms dual to (4.1)⁸⁴. Abusing notation, denote by

$$\begin{array}{lll} \alpha \in \text{End}^{-1}(C) & \beta \in \text{End}^1(C) & K \in \text{Coder}^0(C) \\ d_\alpha \in \text{End}^{-1}(\text{Coder}(C)^H) & d_\beta \in \text{End}^1(\text{Coder}(C)^H) & \end{array}$$

the duals of the objects in Section 4 of the same name. They satisfy analogous identities, e.g. $\alpha^2 = \beta^2 = 0$. By (4.8), and this is our implicit definition of a co-dgBVa^Q on C :

$$\begin{aligned} \frac{1}{2}[\delta^1, \delta^1] &= -d_\beta \delta^1 \\ d_\alpha \delta^1 + [\delta^1, \delta^{-1}] &= K - d_\beta \delta^{-1} \\ d_\alpha \delta^{-1} + \frac{1}{2}[\delta^{-1}, \delta^{-1}] &= 0 \end{aligned} \quad (6.14)$$

The assumption that δ^{-1} only has a $V \rightarrow V$ component is only used much later in this proof. Denote the native products on A by juxtaposition and $[-, -]$ like in Section 4.1; later in this proof we also denote them by $\nabla^0, \nabla^{-1} : A \otimes A \rightarrow A$. All are H -equivariant. Recall that, ignoring degrees, $V \hookrightarrow C \hookrightarrow A$. First define $d_A|_C$ and $h_A|_C$ by

$$\begin{aligned} d_A(c) &= \beta(c) + \delta^1(c) + (-1)^{c_1} c_1 c_2 + (-1)^{z_1-1} [z_1, z_2] \\ h_A(c) &= \alpha(c) + \delta^{-1}(c) \end{aligned} \quad (6.15)$$

⁸¹That d and h are differentials follows from (6.14) below (which follows from (4.8)) using $\alpha|_V = \beta|_V = 0$.

⁸²This P is far from being the canonical map $A \rightarrow V[-2]$.

⁸³The dual of the $V^* \rightarrow [V^*, V^*]$ component of δ^1 . This need not be an associative product.

⁸⁴They are formal duals of the gca and gLa products on GV^* respectively.

for all $c \in C$ where, using Sweedler notation, $\Delta^{-1}c = c_1 \otimes c_2$ and $\Delta^0 c = z_1 \otimes z_2$. This is consistent with d_A and h_A having degrees 1 and -1 respectively, thanks to the -2 degree shift in the definition of A . Extend d_A and h_A uniquely by

$$\begin{aligned} d_A(ab) &= d_A(a)b + (-1)^a a d_A(b) \\ d_A([a, b]) &= (-1)^a (Q(ab) - (Qa)b - a(Qb)) + [d_A a, b] + (-1)^{a-1} [a, d_A b] \\ h_A(ab) &= (-1)^a [a, b] + h_A(a)b + (-1)^a a h_A(b) \\ h_A([a, b]) &= [h_A a, b] + (-1)^{a-1} [a, h_A b] \end{aligned} \tag{6.16}$$

for all $a, b \in A$. These extensions exist⁸⁵ and they are H -equivariant. To check that one has a dgBVa^Q as claimed in Lemma 17, only $d_A^2 = h_A^2 = 0$ and $[d_A, h_A] = Q\mathbb{1}$ remain todo. Using (6.16) one checks $d_A^2, h_A^2 \in \text{Der}(A)$, so they vanish if their restriction to C vanishes; and $[d_A, h_A]$ fails to be a derivation precisely in such a way⁸⁶ that it suffices to show that $[d_A, h_A]|_C = Q\mathbb{1}$. Lengthy calculations using (6.15) and (6.16) yield

$$\begin{aligned} h_A^2|_C &= d_\alpha \delta^{-1} + \tfrac{1}{2} [\delta^{-1}, \delta^{-1}] \\ [d_A, h_A]|_C &= d_\alpha \delta^1 + d_\beta \delta^{-1} + [\delta^1, \delta^{-1}] + \alpha\beta + \beta\alpha \\ d_A^2|_C &= d_\beta \delta^1 + \tfrac{1}{2} [\delta^1, \delta^1] \end{aligned}$$

where the first also uses $\alpha^2 = 0$; the second also uses $\delta^{-1} \in \text{Coder}^{-1}(C)^H$ and the dual of (4.3); the third also uses $\beta^2 = 0$ and $\delta^1 \in \text{Coder}^1(C)^H$ and the duals of (4.1) and (4.5). Now (6.14) imply $d_A^2|_C = h_A^2|_C = 0$ and $[d_A, h_A]|_C = K + \alpha\beta + \beta\alpha = Q\mathbb{1}$, where the very last equality uses (A.2). This concludes the construction of the dgBVa^Q.

To discuss (6.12), first note that $\Delta^{-1}|_V = \Delta^0|_V = 0$, because C is without counit, and $\beta|_V = \alpha|_V = 0$, so (6.15) gives $d_A|_V = \delta^1|_V$ and $h_A|_V = \delta^{-1}|_V$, so $I : V[-2] \hookrightarrow A$ so $d_A I = I d$ and $h_A I = I h$ as claimed. To continue, note the equivalent definitions

$$\begin{aligned} d_A &= \beta_A + \mathcal{D}(\beta) + \mathcal{D}(\delta^1) + \mathcal{D}(\pm \nabla^0 \circ \Delta^{-1}) + \mathcal{D}(\pm \nabla^{-1} \circ \Delta^0) \\ h_A &= \alpha_A + \mathcal{D}(\alpha) + \mathcal{D}(\delta^{-1}) \end{aligned} \tag{6.17}$$

where $\alpha_A \in \text{End}^{-1}(A)$, $\beta_A \in \text{End}^1(A)$ are defined like (4.3) and (4.5) respectively; where $\mathcal{D} : \text{Hom}(C, A) \rightarrow \text{Der}(A)$ extends maps to derivations; $\nabla^0, \nabla^{-1} : A \otimes A \rightarrow A$ are the two products⁸⁷; the \pm are detailed in (6.15); and $C \hookrightarrow A$ is implicit in a few places.

We also introduce auxiliary gradings. For every Gerstenhaber word, denote by n_{gca} and n_{gLa} the number of gca and gLa products that it uses; this yields a well-defined \mathbb{N}^2 -grading⁸⁸ on every free Gerstenhaber algebra. Dualization and sign reversal yields an \mathbb{N}^2 -grading on every cofree Gerstenhaber coalgebra⁸⁹. Note that A is spanned by Gerstenhaber words of co-Gerstenhaber words in V . To every such word-of-words $w \in A$ we associate: $(n_{\text{gca}}^A, n_{\text{gLa}}^A)$ is the \mathbb{N}^2 -degree of w as a Gerstenhaber word, and $(n_{\text{gca}}^C, n_{\text{gLa}}^C)$ is the sum of the \mathbb{N}^2 -degrees

⁸⁵ Just like α and β in Section 4 and Appendix A. This uses the 7-term identity (4.4).

⁸⁶ Analogous to the last two equations in (A.1).

⁸⁷ In particular, $\nabla^0 = \Theta$ are two notations for the same object.

⁸⁸ $\mathbb{N} = \{0, 1, 2, 3, \dots\}$.

⁸⁹ Dualization produces non-positive degrees, the sign reversal is to get an \mathbb{N}^2 -grading.

	n_{gca}^A	n_{gLa}^A	n_{gca}^C	n_{gLa}^C	p	q	r
β_A	1	-1	0	0	1	-1	0
$\mathcal{D}(\beta)$	0	0	-1	1	1	-1	0
$\mathcal{D}(\delta^1)$	0	0	≤ 0	≤ 0	≤ 0	≤ 0	≤ 0
$\mathcal{D}(\pm \nabla^0 \circ \Delta^{-1})$	1	0	0	-1	0	0	0
$\mathcal{D}(\pm \nabla^{-1} \circ \Delta^0)$	0	1	-1	0	0	0	0
α_A	-1	1	0	0	-1	1	0
$\mathcal{D}(\alpha)$	0	0	1	-1	-1	1	0
$\mathcal{D}(\delta^{-1})$	0	0	≤ 0	≤ 0	≤ 0	≤ 0	≤ 0

Table 1: Degrees of the summands in (6.17). They do not increase r . The entries for $\mathcal{D}(\delta^{-1})$ are actually all zero if one uses the assumption that δ^{-1} has only a $V \rightarrow V$ component.

of the co-Gerstenhaber words that constitute w . This yields a well-defined \mathbb{N}^4 -grading on A . Also introduce the following coarser gradings:

$$p = n_{\text{gLa}}^C + n_{\text{gca}}^A \quad q = n_{\text{gca}}^C + n_{\text{gLa}}^A \quad r = p + q$$

giving decompositions $A = \oplus_{p,q} A_{p,q} = \oplus_r A_r$. The summands in (6.17) do not increase r , as shown in Table 1. Now consider the components that neither depend on $\delta^{\pm 1}$ nor on the H -module structure or Q , only on V itself. Namely, set

$$u = \mathcal{D}(\pm \nabla^0 \circ \Delta^{-1}) + \mathcal{D}(\pm \nabla^{-1} \circ \Delta^0) \quad v = \alpha_A + \mathcal{D}(\alpha)$$

A byproduct of the above proof that we have a dgBVa Q is $u^2 = v^2 = uv + vu = 0$. Note that $u(A_r) \subseteq A_r$ and $v(A_r) \subseteq A_r$, and decomposing $A_r = \oplus_{p+q=r} A_{p,q}$ we get:

$$\begin{array}{ccccccc} \begin{array}{c} u \\ \curvearrowright \\ A_{r,0} \end{array} & \xrightarrow{v} & \begin{array}{c} u \\ \curvearrowright \\ A_{r-1,1} \end{array} & \xrightarrow{v} & \cdots & \xrightarrow{v} & \begin{array}{c} u \\ \curvearrowright \\ A_{1,r-1} \end{array} \xrightarrow{v} \begin{array}{c} u \\ \curvearrowright \\ A_{0,r} \end{array} \end{array} \quad (6.18)$$

To every graded vector space V one can functorially associate (6.18); it is for every r a universal construction coming with the Gerstenhaber operad that involves suitable tensoring with $V^{\otimes(r+1)}$. The homology of $u|_{A_r}$ vanishes if $r > 0$ ⁹⁰. The homology of $v|_{A_r}$ vanishes if $r > 0$ ⁹¹. This implies that there exists a $g \in \text{End}^{-1}(A)$ that satisfies⁹²

$$g(A_{p,q}) \subseteq A_{p,q} \quad g|_{A_0} = 0 \quad g^2 = 0 \quad ug + gu = \mathbb{1}_{>0} \quad vg + gv = 0 \quad (6.19)$$

where, by definition, $\mathbb{1}_{>0}|_{A_r} = \mathbb{1}$ if $r > 0$ and $\mathbb{1}_{>0}|_{A_0} = 0$. This g can be chosen functorially in V , specifically so that g is H -equivariant and commutes with every derivation on A

⁹⁰Because the Gerstenhaber operad is Koszul [17, 18, 25], and see [16, Theorem 4.2.5].

⁹¹Let $\gamma \in \text{End}^1(C)$, $\kappa \in \text{End}^0(C)$ be dual to the objects in Section 4.3; let $\gamma_A \in \text{End}^1(A)$, $\kappa_A \in \text{End}^0(A)$ be like in Section 4.3. Set $w = \gamma_A + \mathcal{D}(\gamma) \in \text{End}^1(A)$. Then $w(A_r) \subseteq A_r$, $w^2 = 0$ and $wv + vw = \kappa_A + \mathcal{D}(\kappa)$. The last operator commutes with v and w and is invertible on A_r when $r > 0$, using (4.10).

⁹²For $r > 0$, this would follow from $H(u|_{A_r}) = 0$, if it was not for the requirement $vg + gv = 0$. In (6.18), choose g from left to right, at each step consistent with $vg + gv = 0$, which works because $H(v|_{A_r}) = 0$.

that extends a coderivation on C that extends an endomorphism on V . In particular, for $d = \delta^1|_V$ and $h = \delta^{-1}|_V$, here viewed as elements of $\text{Coder}(C)$,

$$\mathcal{D}(d)g + g\mathcal{D}(d) = 0 \quad \mathcal{D}(h)g + g\mathcal{D}(h) = 0 \quad (6.20)$$

By (6.17), (6.19), (6.20) and Table 1, the term $\mathbb{1}_{>0} - d_A g - g d_A$ decreases $(r, q) \in \mathbb{N}^2$ ordered lexicographically⁹³, so its restriction to $A_0 \oplus \dots \oplus A_r$ is nilpotent. Define⁹⁴

$$\mathbb{Z} = \sum_{n \geq 0} g(\mathbb{1}_{>0} - d_A g - g d_A)^n \quad (6.21)$$

which, when applied to any given element of A , has finitely many nonzero summands. Set $\pi = \mathbb{1} - d_A \mathbb{Z} - \mathbb{Z} d_A$. On A/A_0 we have $\mathbb{1}_{>0} = \mathbb{1}$ and, considering (6.21) on this quotient, a standard calculation using $g^2 = 0$ implies $\pi = 0$ on A/A_0 . So $\text{image } \pi \subseteq A_0$. But $g|_{A_0} = 0$, hence $\mathbb{Z}|_{A_0} = 0$, hence $\pi|_{A_0} = \mathbb{1}$. Therefore, $\pi^2 = \pi$ with $\text{image } \pi = A_0$. Continuing, $g^2 = 0$ and $\mathbb{1}_{>0}g = g\mathbb{1}_{>0}$ imply $\mathbb{Z}g = 0$, hence $\mathbb{Z}^2 = 0$. This implies $\pi\mathbb{Z} = \mathbb{Z}\pi$. Since $\mathbb{Z}|_{A_0} = 0$ we have $\mathbb{Z}\pi = 0$, hence $\pi\mathbb{Z} = 0$. Note $A_0 \simeq V$, and $\text{image } \pi = A_0 = \text{image } I$. Let P be the composition of π with the canonical $A \rightarrow V[-2]$, so $PI = \mathbb{1}$ and $\ker P = \ker \pi$.

To prove (6.13), using $\text{image}(\Theta \circ I^{\otimes 2}) \subseteq A_0 \oplus A_{1,0}$ it suffices to consider the subcomplex $(A_0 \oplus A_{1,0}, d_A)$. The diagonal blocks $A_0 \rightarrow A_0$ and $A_{1,0} \rightarrow A_{1,0}$ are described above, the block $A_{1,0} \rightarrow A_0$ only uses θ , a piece of δ^1 . The construction of \mathbb{Z} implies (6.13).

Now recall the assumption that the only nonzero component of δ^{-1} is its $V \rightarrow V$ component, $\delta^{-1} = h$. Then $h_A(A_r) \subseteq A_r$ hence $\mathbb{1}_{>0}h_A = h_A\mathbb{1}_{>0}$. Using (6.19) and (6.20) we have $gh_A + h_Ag = 0$. Using $d_A h_A + h_A d_A = Q\mathbb{1}$ and $Qg = gQ$, by H -equivariance of g , it follows that h_A commutes with $\mathbb{1}_{>0} - d_A g - g d_A$. Hence $\mathbb{Z}h_A + h_A\mathbb{Z} = 0$. Now, using $Q\mathbb{Z} = \mathbb{Z}Q$, by H -equivariance of \mathbb{Z} , one gets $\pi h_A = h_A\pi$. Hence $Ph_A = hP$. \square

7 Acknowledgements

M.R. is grateful to have received funding from ERC through grant agreement No. 669655.

A Addendum to Section 4.2

Here are the details for several claims in Section 4.2.

To see that β is well-defined, recall the definition of GV^* via terms and relations. Set $C(a, b) = ab - (-1)^{ab}ba$, $A(a, b, c) = (ab)c - a(bc)$ and $X(a, b) = [a, b] + (-1)^{(a+1)(b+1)}[b, a]$ where here a, b, c are terms before imposing any relations. By direct calculation,

$$\begin{aligned} \beta(C(a, b)) &= C(\beta(a), b) + (-1)^a C(a, \beta(b)) \\ \beta(A(a, b, c)) &= A(\beta(a), b, c) + (-1)^a A(a, \beta(b), c) + (-1)^{a+b} A(a, b, \beta(c)) \\ \beta(X(a, b)) &= X(\beta(a), b) + (-1)^{a-1} X(a, \beta(b)) + (-1)^a (QC(a, b) - C(Qa, b) - C(a, Qb)) \end{aligned}$$

⁹³So it either decreases r , or it preserves r and decreases q .

⁹⁴This can be interpreted as a formula that implements back-substitution to ‘invert’ triangular matrices; or as a Neumann series; or as an application of the homological perturbation lemma.

and note that⁹⁵ $QC(a, b) = C(Q_1a, Q_2b)$ using Sweedler notation $\Delta Q = Q_1 \otimes Q_2$. Given this and given how β acts on products, see (4.5), one concludes that β preserves the ‘ideal’ generated by C, A, X . From here on we work modulo these relations. Set

$$\begin{aligned} P(a, b, c) &= [ab, c] - a[b, c] - (-1)^{ab}b[a, c] \\ J(a, b, c) &= [a, [b, c]] + (-1)^{(a+1)(b+c)}[b, [c, a]] + (-1)^{(c+1)(a+b)}[c, [a, b]] \end{aligned}$$

One finds

$$\begin{aligned} \beta(P(a, b, c)) &= P(\beta(a), b, c) + (-1)^a P(a, \beta(b), c) + (-1)^{a+b-1} P(a, b, \beta(c)) \\ &\quad + (-1)^{a+b} (Q(abc) - (Q(ab))c - aQ(bc) - (-1)^{ab}bQ(ac) + (Qa)bc + a(Qb)c + ab(Qc)) \end{aligned}$$

and note that the last line vanishes by the 7-term identity (4.4). One concludes that β also preserves the ‘ideal’ generated by P . We work modulo P from here on. Similarly,

$$\begin{aligned} \beta(J(a, b, c)) &= J(\beta(a), b, c) + (-1)^{a-1} J(a, \beta(b), c) + (-1)^{a+b-2} J(a, b, \beta(c)) \\ &\quad + (-1)^a (Q(a[b, c]) - (Qa)[b, c] - aQ[b, c]) \\ &\quad - (-1)^{a+b} [a, Q(bc) - (Qb)c - b(Qc)] \\ &\quad + (-1)^{a(b+c)+c} (Q(b[c, a]) - (Qb)[c, a] - bQ[c, a]) \\ &\quad - (-1)^{a(b+c)} [b, Q(ca) - (Qc)a - c(Qa)] \\ &\quad + (-1)^{c(a+b)+a+b+c} (Q(c[a, b]) - (Qc)[a, b] - cQ[a, b]) \\ &\quad - (-1)^{c(a+b)+b+c} [c, Q(ab) - (Qa)b - a(Qb)] \end{aligned}$$

To see that the terms involving Q cancel, replace $[c, Q(ab)] = [c, (Q_1a)(Q_2b)]$ and so forth, then rewrite using $P = 0$ and use (4.4). So, β is well-defined on GV^* .

Suppose now $\delta \in \text{Der}(GV^*)^H$. We show $d_\beta(\delta) \in \text{Der}(GV^*)^H$. By direct calculation,

$$d_\beta(\delta)(ab) = (d_\beta(\delta)(a))b + (-1)^{(\delta+1)a} a d_\beta(\delta)(b)$$

for all $a, b \in GV^*$ without using H -equivariance of δ . By direct calculation,

$$\begin{aligned} d_\beta(\delta)([a, b]) &= [d_\beta(\delta)(a), b] + (-1)^{(\delta+1)(a-1)} [a, d_\beta(\delta)(b)] \\ &\quad + (-1)^{\delta+a} (Q(\delta(a)b) - (Q\delta(a))b - \delta(a)(Qb) \\ &\quad + (-1)^{\delta a} Q(a\delta(b)) - (-1)^{\delta a} (Qa)\delta(b) - (-1)^{\delta a} a(Q\delta(b)) \\ &\quad - \delta(Q(ab)) + \delta((Qa)b) + \delta(a(Qb))) \end{aligned}$$

To see that the Q terms cancel, replace $Q(\delta(a)b) = (Q_1\delta(a))(Q_2b) = (\delta(Q_1a))(Q_2b)$ using H -equivariance of δ ; similarly $\delta(Q(ab)) = \delta((Q_1a)(Q_2b))$; and so on. Since δ is a derivation, the Q terms cancel. Here (4.4) is not used.

To check (4.6), set $N = \alpha\beta + \beta\alpha \in \text{End}^0(GV^*)$ and check by direct calculation that

$$\begin{aligned} N|_{V^*} &= 0 \\ N(ab) &= Q(ab) - (Qa)b - a(Qb) + N(a)b + aN(b) \\ N([a, b]) &= Q[a, b] - [Qa, b] - [a, Qb] + [N(a), b] + [a, N(b)] \end{aligned} \tag{A.1}$$

⁹⁵Using the H -module structure on the space of terms, and cocommutativity.

for all $a, b \in GV^*$; this uses the H -equivariance of α and β but not (4.4). Hence

$$N + K = Q\mathbb{1} \quad (\text{A.2})$$

Clearly $d_\alpha d_\beta + d_\beta d_\alpha = [N, -]$. Hence $[N, -] = -[K, -]$ when restricted to $\text{Der}(GV^*)^H$ since $[Q\mathbb{1}, -]$ annihilates H -equivariant derivations. This gives (4.6). To see $d_\alpha(K) = 0$, note that $d_\alpha(Q\mathbb{1}) = 0$ since α is H -equivariant, and $d_\alpha(N) = \alpha(\alpha\beta + \beta\alpha) - (\alpha\beta + \beta\alpha)\alpha = 0$ because $\alpha^2 = 0$. The proof that $d_\beta(K) = 0$ goes the same way.

B Some remarks about the BV formalism

The BV_∞^\square algebra in Theorem 3 is not directly related, to the author's knowledge, to the algebraic structures in the BV formalism for the quantization of gauge theories. Nevertheless, for comparison, here are some remarks about the BV formalism [31, 32]. Costello [13] uses the BV formalism to introduce the dgca denoted \mathfrak{a} in this paper.

1. Let M be a finite-dimensional manifold. The gca of polyvector fields $A = \Gamma(\wedge(TM))$ with the Schouten bracket $\{-, -\}$ is a Gerstenhaber algebra⁹⁶.
2. Suppose M has a volume form, μ . It induces an isomorphism between A and the de Rham complex of M . Define the degree -1 map $\Delta_\mu = \mu^{-1} \circ d_{\text{dR}} \circ \mu : A \rightarrow A$. Then Δ_μ is second order on A ; evidently $\Delta_\mu^2 = 0$; and the associated Gerstenhaber bracket is the Schouten bracket [31, Theorem 2.8]. This turns A into a BV algebra.
3. The supergeometry perspective is that A is the ‘algebra of functions’ on the graded supermanifold $T^*[-1]M$, the odd cotangent bundle of M ; and the Schouten bracket is an odd Poisson, in fact odd symplectic, structure [31, Section 2.3].
4. For every $S \in A^{\text{even}}$, the quantum master equation $\Delta_\mu S + \frac{1}{2}\{S, S\} = 0$ is formally equivalent to $\Delta_\mu e^S = 0$, and formally implies that the integration of e^S over a Lagrangian submanifold $L \subseteq T^*[-1]M$, a small perturbation of the canonical Lagrangian submanifold $M \subseteq T^*[-1]M$, is independent of L , see [32].
5. If M is a finite-dimensional vector space, then μ is canonical up to normalization. There is an analogous story where A is replaced by formal power series on the graded vector space $T^*[-1]M = M \oplus M^*[-1]$; the supergeometry perspective is that this is the ‘algebra of functions’ on a formal neighborhood of the origin $0 \in T^*[-1]M$.
6. Applying this to gauge theory involves generalizations: M is an infinite-dimensional vector space of ‘fields’, say the space of sections of a vector bundle on a 4-manifold; M is itself graded, say due to the addition of BRST ghosts. Passing from M to $T^*[-1]M$ is referred to as adding antifields. In [13, Chapter 6.2] the BV formalism is used to introduce fields for first order Yang-Mills and, based on this, to define the dgca \mathfrak{a} .

⁹⁶The differential on A is zero: By the Hochschild-Kostant-Rosenberg theorem, the graded vector space A is the homology of a subcomplex of the complex of Hochschild cochains for the algebra $C^\infty(M)$.

C Code

This Wolfram Mathematica code is not part of the logic of this paper. It can be used to construct θ_n for small arity n and check all axioms up to that arity (cf. Section 6.1).

The code uses ordered bases, so `e0i123` corresponds to $e^0e^1 + ie^2e^3 \in \Lambda_+^2$ in row 0 in (1.1), `s0i123` corresponds to $e^0e^1 + ie^2e^3 \in \Lambda_+^2$ in row 1. The degree `deg` is that in $V^{-2} \oplus \dots \oplus V^1$; `rdeg` is the row degree. The gca product and differential are given by pre-calculated arrays. For instance $e^0e^1 + ie^2e^3$ in row 0 times $e^0e^1 + ie^2e^3$ in row 1 yields $2ie^0e^1e^2e^3$ in row 1, corresponding to the entry `{16,6,9}->2*I` below. Further, `k[0],k[1],k[2],k[3]` are symbols for the four partial derivatives, or equivalently, for the four components of the momentum.

```
(* ordered bases and degrees *)
bas={one, e0,e1,e2,e3, e0i123,e0i23i1,e03i12,
      s0i123,s02i31,s03i12, s123,s023,s031,s012, s0123};
dual=Map[ToExpression["dual"<ToString[#]]&,bas];
MapThread[(deg[#1]=deg[#5]=#3;deg[#2]=-#3;rdeg[#1]=rdeg[#5]=#4;pos[#2]=#5;ndeg[#2]=1;)&,
  {bas,dual,{-2,-1,-1,-1,-1,0,0,0,-1,-1,-1,0,0,0,0,1},
    {0,0,0,0,0,0,0,0,1,1,1,1,1,1,1,1},Range[1,16]}];

(* pre-calculated gca product and differential, and h *)
prod=SparseArray[{{1,1,1}->1,{2,1,2}->1,{2,2,1}->1,{3,1,3}->1,{3,3,1}->1,{4,1,4}->1,{4,4,1}->1,{5,1,5}->1,
  {5,5,1}->1,{6,1,6}->1,{6,2,3}->1/2,{6,3,2}->-1/2,{6,6,1}->1,{6,4,5}->-I/2,{6,5,4}->I/2,{7,1,7}->1,
  {7,2,4}->1/2,{7,4,2}->-1/2,{7,7,1}->1,{7,3,5}->I/2,{7,5,3}->-I/2,{8,1,8}->1,{8,2,5}->1/2,{8,5,2}->-1/2,
  {8,8,1}->1,{8,3,4}->-I/2,{8,4,3}->I/2,{9,1,9}->1,{9,9,1}->1,{10,1,10}->1,{10,10,1}->1,{11,1,11}->1,
  {11,11,1}->1,{12,1,12}->1,{12,3,9}->I,{12,4,10}->I,{12,5,11}->I,{12,9,3}->-I,{12,10,4}->-I,{12,11,5}->-I,
  {12,12,1}->1,{13,1,13}->1,{13,2,9}->I,{13,4,11}->-1,{13,5,10}->1,{13,9,2}->-I,{13,10,5}->-1,{13,11,4}->1,
  {13,13,1}->1,{14,1,14}->1,{14,2,10}->I,{14,3,11}->1,{14,5,9}->-1,{14,9,5}->1,{14,10,2}->-I,{14,11,3}->-1,
  {14,14,1}->1,{15,1,15}->1,{15,2,11}->I,{15,3,10}->-1,{15,4,9}->1,{15,9,4}->-1,{15,10,3}->1,{15,11,2}->-I,
  {15,15,1}->1,{16,1,16}->1,{16,2,12}->1,{16,3,13}->-1,{16,4,14}->-1,{16,5,15}->-1,{16,6,9}->2*I,
  {16,7,10}->2*I,{16,8,11}->2*I,{16,9,6}->2*I,{16,10,7}->2*I,{16,11,8}->2*I,{16,12,2}->1,{16,13,3}->-1,
  {16,14,4}->-1,{16,15,5}->-1,{16,16,1}->1,{16,16,16}}];
dmat=SparseArray[{{2,1}->k[0],{3,1}->k[1],{4,1}->k[2],{5,1}->k[3],{6,9}->1,{6,4}->I*k[3]/2,
  {6,5}->-I*k[2]/2,{6,2}->-k[1]/2,{6,3}->k[0]/2,{7,10}->1,{7,3}->-I*k[3]/2,{7,2}->-k[2]/2,{7,5}->I*k[1]/2,
  {7,4}->k[0]/2,{8,11}->1,{8,2}->-k[3]/2,{8,3}->I*k[2]/2,{8,4}->-I*k[1]/2,{8,5}->k[0]/2,{12,11}->I*k[3],
  {12,10}->I*k[2],{12,9}->I*k[1],{13,10}->k[3],{13,11}->-k[2],{13,9}->I*k[0],{14,9}->-k[3],{14,11}->k[1],
  {14,10}->I*k[0],{15,9}->k[2],{15,10}->-k[1],{15,11}->I*k[0],{16,15}->-k[3],{16,14}->-k[2],{16,13}->-k[1],
  {16,12}->k[0]},{16,16}}];
nvar:=With[{x=Unique[]},x/:HoldPattern[NumericQ[x]]=True;x];
hmat=Outer[If[deg[#1]==deg[#2]-1,Switch[rdeg[#1]-rdeg[#2],0,
  Sum[nvar*k[i],{i,0,3}]],-1,nvar,_,0]&,bas,bas];
```

The ad-hoc implementation of a free Gerstenhaber algebra with product `mul`, bracket `lie` below is not guaranteed to reduce all expressions to a normal form⁹⁷, but suffices here.

```
(* degrees and linearity *)
setdeg[s_,offset_:0]:=( s[_?NumericQ]=0;
  s[mul[a_,b_]]:=s[a]+s[b];
  s[lie[a_,b_]]:=s[a]+s[b]+offset;
  s[(?NumericQ)k[_]|Power[k[_],_]]*a_:=s[a];);
lin[s_]:=( s[a___,(b_:1)*c_Plus,d___]:=Map[s[a,b*#,d]&,c];
  s[a___,n_*NumericQ*b_,c___]:=n*s[a,b,c];
  s[___,0,___]=0;);
setdeg[deg,-1]; setdeg[ndeg]; setdeg[rdeg]; lin[mul]; lin[lie];
set[pairs_]:=Scan[(Evaluate[First[#]]&Last[#])&,pairs];

(* ad-hoc implementation of free Gerstenhaber algebra *)
mul[a_]:=a; mul[a___,n_*NumericQ,b___]:=n*mul[a,b]; lie[_?NumericQ]=lie[_?NumericQ,_]=0;
HoldPattern[mul[a___,mul[b___],c___]]:=mul[a,b,c];
mul[a___,b_,c_,d___]/;Not[OrderedQ[{b,c}]]:=(-1)^(deg[b]*deg[c])*mul[a,c,b,d];
lie[a_,b_]/;Not[OrderedQ[{a,b}]]:=(-1)^(deg[a]-1)*(deg[b]-1)*lie[b,a];
lie[mul[a_,b_],c_]:=mul[a,lie[mul[b],c]]+(-1)^(deg[a]*deg[mul[b]])*mul[b,lie[a,c]];
lie[a_,mul[b_,c_]]:=mul[lie[a,b],c]+(-1)^(deg[a]-1)*deg[b]*mul[b,lie[a,mul[c]]];
```

⁹⁷Unlike, say, Gröbner basis reduction for the quotient of a polynomial algebra by an ideal.

```

mul/:Power[k[i_],j_:1]*mul[a_,b_]:=k[i]^(j-1)*(mul[k[i]*a,b]+mul[a,k[i]*mul[b]])//Expand;
lie/:Power[k[i_],j_:1]*lie[a_,b_]:=k[i]^(j-1)*(lie[k[i]*a,b]+lie[a,k[i]*b])//Expand;
Scan[{lie[#,#]=0;mul[_,_,#,#]=0}&,&,
  Flatten[Map[{1,k[0],k[1],k[2],k[3]}*#&,Select[dual,OddQ[deg[#]]&]]];
J[a_,b_,c_]:= (lie[lie[a,b],c]+(-1)^(deg[a]-1)*(deg[b]+deg[c]))*lie[lie[b,c],a]
+(-1)^(deg[c]-1)*(deg[a]+deg[b]))*lie[lie[c,a],b];
Outer[set[First[Solve[Thread[J[##]==0]]]&,dual,dual,dual]; (* Jacobi identities *)

```

Below is an implementation of Gerstenhaber derivations; the operations α , β , γ and d_α , d_β , Γ and the curvature K ; the operations ν_k , μ_k in terms of θ_k where at the moment θ_k is given by an ansatz with undetermined coefficients; and the axioms A_k , B_k , C_k .

```

(* Gerstenhaber derivation associated to f in the default case g=h=0 *)
op[ddeg_,dndeg_,f_,g_:(0&),h_:(0&)]:=Module[{s},
  lin[s]; s[_?NumericQ]=0; Scan[{s[#]:=s[#]=Expand[f[#]]&},dual];
  s[mul[a_,b_]]:=(-1)^deg[a]*g[a,mul[b]]+mul[s[a],b]+(-1)^(ddeg*deg[a])*mul[a,s[mul[b]]];
  s[lie[a_,b_]]:=(-1)^deg[a]*h[a,b]+lie[s[a],b]+(-1)^(ddeg*(deg[a]-1))*lie[a,s[b]];
  s/:deg[s]=ddeg; s/:ndeg[s]=dndeg;
  s[Power[k[i_],j_:1]*a_]:=k[i]^j*s[a];
  s/:(n?NumericQ)*s:=op[ddeg,dndeg,n*f[#]&,n*g[#]&,n*h[#]&];
  s];
comm[x_,y_]:=op[deg[x]+deg[y],ndeg[x]+dndeg[y],x[y[#]]-(-1)^(deg[x]*deg[y])*y[x[#]]&];

(* alpha, beta, gamma, curvature, etc *)
Q=-k[0]^2+k[1]^2+k[2]^2+k[3]^2;
alpha=op[-1,0,0&,lie,0&];
beta=op[1,0,0&,Q*mul[#1,#2]-mul[Q*#1,#2]-mul[#1,Q*#2]&];
gamma=op[1,0,0&,Q*ndeg[#1]*ndeg[#2]*mul[#1,#2]&];
da=comm[alpha,#]&; db=comm[beta,#]&; K=op[0,0,Q*#&];
GAMMA[x_]:=With[{n=ndeg[x]},op[deg[x]+1,n,If[n==0,0,2/((n+1)*n)]*gamma[x[#]]&];

(* d, h and gca product *)
d=op[1,0,0,mat[[pos[#]]].dual&]; h=op[-1,0,hmat[[pos[#]]].dual&];
theta[2]=op[2,1,Flatten[prod[[pos[#]]]].Flatten[Outer[mul,dual,dual]]&];

(* ansatz for theta_n, n>=3: graded symmetric, deg=2 and rdeg=0 *)
incr[n_]:=incr[n]=Select[Select[Tuples[Range[1,16],n],LessEqual@@#&],
  DuplicateFreeQ[Cases[#,Alternatives@@Select[Range[1,16],OddQ[deg[#]]&]]]&];
incr2[n_]:=incr2[n]=Table[Select[incr[n],And[Total[Map[deg,#]]+2==deg[i],
  Total[Map[rdeg,#]]==rdeg[i]]&],{i,1,16}];
theta[n_]:=theta[n]=op[2,n-1,Total[Map[nvar*(mul@@dual[[#]])&,incr2[n][[pos[#]]]]&];
nu[k_]:=nu[k]=da[theta[k]]; mu[k_]:=mu[k]=-GAMMA[comm[h,nu[k]]];

(* auxiliary derivations, and axioms A,B,C decomposed by degree *)
dd=comm[d,d]; dh=comm[d,h]; hh=comm[h,h];
dn[k_]:=dn[k]=comm[d,nu[k]]; dm[k_]:=dm[k]=comm[d,mu[k]];
hn[k_]:=hn[k]=comm[h,nu[k]]; hm[k_]:=hm[k]=comm[h,mu[k]];
mm[m_,n_]:=mm[m,n]=comm[mu[m],mu[n]]; mn[m_,n_]:=mn[m,n]=comm[mu[m],nu[n]];
nn[m_,n_]:=nn[m,n]=comm[nu[m],nu[n]];
dam[k_]:=dam[k]=da[mu[k]]; dan[k_]:=dan[k]=da[nu[k]]; dbn[k_]:=dbn[k]=db[nu[k]];
Unprotect[C];
A[1,2]=dd/2; A[1,0]=op[0,0,-K[#]+dh[#]&]; A[1,-2]=hh/2;
A[k_,2]:=A[k,2]=op[2,k-1,dbn[k][#]+dm[k][#]+1/2*Sum[mm[k+1-n,n][#],{n,2,k-1}&];
A[k_,0]:=A[k,0]=op[0,k-1,hm[k][#]&];
B[k_,2]:=B[k,2]=op[2,k-1,dn[k][#]+Sum[mn[k+1-n,n][#],{n,2,k-1}&];
B[k_,0]:=B[k,0]=op[0,k-1,dam[k][#]+hm[k][#]&];
C[k_,2]:=C[k,2]=op[2,k-1,1/2*Sum[nn[k+1-n,n][#],{n,2,k-1}&];
C[k_,0]:=C[k,0]=op[0,k-1,dan[k][#]&];
q[k_]:=q[k]=GAMMA[op[2,k-1,Sum[mn[k+1-n,n][#],{n,2,k-1}&]]&];
dt[k_]:=dt[k]=comm[d,theta[k]];
b[k_]:=b[k]=op[3,k-1,-dt[k][#]+q[k][#]&];

```

Here an h is fixed such that $A_1 = 0$, note that this h is not unique; then the coefficients of θ_n are computed for small n ; finally all the axioms up to that arity are checked.

```

(* fix h, not unique *)
coeffs[expr_]:=Map[Last,Flatten[CoefficientRules[expr,Join[Table[k[i],{i,0,3}],dual]]];
set[First[Solve[Thread[coeffs[Map[A[1,0],dual]]]==0]]];
set[With[{X=coeffs[Map[A[1,-2],dual]]},First[FindInstance[Thread[X==0],Variables[X]]]];

```

```

(* compute theta_n for n=3...nmax *)
nmax=3;
sol[expr_] := Module[{x}, CoefficientList[expr/.MapIndexed[(#1->x^First[#2])&,
DeleteDuplicates[Extract[expr, Position[expr, _mul]]], x]
//DeleteDuplicates//First[Solve[Thread[##==0]]]&];
Do[Scan[set[sol[b[n][#]]]&, dual], {n, 3, nmax}];

(* check all axioms up to arity nmax *)
Table[Map[A[1, i], dual], {i, {-2, 0, 2}}//Flatten//DeleteDuplicates
Table[Map[A[n, i], dual], {n, 2, nmax}, {i, {0, 2}}//Flatten//DeleteDuplicates
Table[Map[B[n, i], dual], {n, 2, nmax}, {i, {0, 2}}//Flatten//DeleteDuplicates
Table[Map[C[n, i], dual], {n, 3, nmax}, {i, {0, 2}}//Flatten//DeleteDuplicates

```

References

- [1] Z. Bern, J.J.M. Carrasco and H. Johansson, *Phys. Rev. D* **78**, 085011 (2008)
New Relations for Gauge-Theory Amplitudes
- [2] Z. Bern, T. Dennen, Y. Huang and M. Kiermaier, *Phys. Rev. D* **82** 065003 (2010)
Gravity as the Square of Gauge Theory
- [3] Z. Bern, J.J. Carrasco, M. Chiodaroli, H. Johansson and R. Roiban, *arXiv:1909.01358* (2019)
The Duality Between Color and Kinematics and its Applications
- [4] H. Tye and Y. Zhang, *JHEP* **1006** (2010) 071
Dual Identities inside the Gluon and the Graviton Scattering Amplitudes
- [5] D. Zhu, *Phys. Rev. D* **22**, 2266 (1980)
Zeros in scattering amplitudes and the structure of non-Abelian gauge theories
- [6] R. Monteiro and D. O’Connell, *JHEP* **1107** (2011) 007
The Kinematic Algebra From the Self-Dual Sector
- [7] C. Cheung and C.H. Shen, *Phys. Rev. Lett.* **118**, 121601 (2017)
Symmetry and Action for Flavor-Kinematics Duality
- [8] C.-H. Fu and K. Krasnov, *JHEP* **1701** (2017) 075
Colour-kinematics duality and the Drinfeld double of the Lie algebra of diffeomorphisms
- [9] G. Chen, H. Johansson, F. Teng and T. Wang, *arXiv:1906.10683* (2019)
On the kinematic algebra for BCJ numerators beyond the MHV sector
- [10] A.M. Zeitlin, *JHEP* **1003:056** (2010)
Conformal Field Theory and Algebraic Structure of Gauge Theory
- [11] A.M. Zeitlin, *Comm. Math. Physics* **303.2**, 331-359 (2011)
Quasiclassical Lian-Zuckerman Homotopy Algebras, Courant Algebroids and Gauge Theory
- [12] A.M. Zeitlin, *J. Physics, Conf. Ser.* **597**, 012074 (2015)
Sigma-models and Homotopy Algebras
- [13] K. Costello, *AMS* **170** (2011)
Renormalization and effective field theory
- [14] A. Nützi and M. Reiterer, *arXiv:1812.06454* (2018)
Scattering amplitudes in YM and GR as minimal model brackets and their recursive characterization
- [15] D. Eisenbud, *Graduate Texts in Math.*, Springer (2005)
The Geometry of Syzygies

- [16] V. Ginzburg and M. Kapranov, *Duke Math. J.* 76, 1, 203-272 (1994)
Koszul duality for Operads
- [17] E. Getzler and J.D.S. Jones, *arXiv:hep-th/9403055* (1994)
Operads, homotopy algebra and iterated integrals for double loop spaces
- [18] M. Markl, *Ann. Inst. Fourier (Grenoble)* 46, 2, 307-323 (1996)
Distributive laws and Koszulness
- [19] M. Manetti, *arXiv:math/0507284* (2005)
Deformation theory via differential graded Lie algebras
- [20] X.Z. Cheng and E. Getzler, *J. Pure Appl. Algebra* 212, 11, 2535-2542 (2008)
Transferring homotopy commutative algebraic structures
- [21] D.E. Tamarkin, *arXiv:math/9803025* (1998)
Another proof of M. Kontsevich formality theorem
- [22] D.E. Tamarkin, *Lett. Math. Phys.* 66:65-72 (2003)
Formality of chain operad of little discs
- [23] D. Tamarkin and B.L. Tsygan, *Methods Funct. Anal. Topology*, 6(2), 85-100 (2000)
Noncommutative differential calculus, homotopy BV algebras and formality conjectures
- [24] I. Galvez-Carrillo, A. Tonks and B. Vallette, *J. Noncommut. Geom.* 6, 539-602 (2012)
Homotopy Batalin-Vilkovisky algebras
- [25] V. Dolgushev, D. Tamarkin and B.L. Tsygan, *J. Noncommut. Geom.* 1, 1-25 (2007)
The homotopy Gerstenhaber algebra of Hochschild cochains of a regular algebra is formal
- [26] T. Willwacher, *Duke Math. J.* 165, 10, 1815-1964 (2016)
The Homotopy Braces Formality Morphism
- [27] D. Sinha, *arXiv:math/0610236* (2006)
The homology of the little disks operad
- [28] P. Salvatore and N. Wahl, *Quart. J. Math.* 54, 213-231 (2003)
Framed discs operads and Batalin-Vilkovisky algebras
- [29] O. Bellier, *Algebr. Geom. Topol.* 14, 1-35 (2014)
Koszul duality theory for operads over Hopf algebras
- [30] J.L. Koszul, *Astérisque*, S131, 257-271 (1985)
Crochet de Schouten-Nijenhuis et cohomologie
- [31] Y. Kosmann-Schwarzbach and J. Monterde, *Ann. Inst. Fourier*, 52, 2, 419-456 (2002)
Divergence operators and odd Poisson brackets
- [32] A. Schwartz, *Commun. Math. Phys.* 155, 249-260 (1993)
Geometry of Batalin-Vilkovisky quantization
- [33] J.L. Loday and B. Vallette, *Grundlehren*, 346, Springer (2012)
Algebraic operads