

QED Treatment of Linear Elastic Waves in Asymmetric Environments

Maysam Yousefian* and Mehrdad Farhoudi†

Department of Physics, Shahid Beheshti University G.C., Evin, Tehran 19839, Iran

(Dated: January 16, 2020)

Regarding some logical gaps in the existing asymmetric elastic theories, by generalizing and modifying the related Cosserat theory (CT) for an elastic environment, we have shown that its elastic wave equations are analogous to quantum electrodynamics (QED) ones. Thus on macro scales, elastic waves can possess the QED treatment. This analogy provides a new paradigm of fermions and bosons. Also, from experimental point of view, contrary to the Cosserat and discrete models, this amended CT has been indicated to be consistent with the corresponding empirical observations.

PACS numbers: 87.10.Pq; 62.20.D-; 45.70.-n; 12.20.-m; 31.30.Jv

Keywords: Linear Elastic Waves; Asymmetric Stress Tensor; Quantum Electrodynamics; Cosserat Theory

I. INTRODUCTION

In the classical elastic theory, the stress tensor has normally been considered to be symmetric. In this scope, the similarity between the elastic and electromagnetic waves is incomplete. For example, in Ref. [1–3], an analogy between the electromagnetic field and a fluid, which includes a large number of vortex filaments, has been investigated. In their work, the pressure of fluid, the fluid velocity and the density of the turbulence energy respectively are as the electrostatic potential, the magnetic vector potential and the electromotive force, but charge has no independent nature. Also in Ref. [4], it has been claimed that the electromagnetic equations fit the structure of a linear elastic continuum which is hard to compression though liable to shear deformation. Again in this work, charge still has no independent nature.

On the other hand, in the CT [5–7], by generalizing the elastic theory and assuming the presence of microstructures in an elastic environment, the stress tensor has been employed asymmetrically. However, still there are some logical gaps in these theories. For instance, any microstructure can have a vibration degree of freedom, which has not been talked about in these theories. Also, there is no mention of the gauge property of the displacement wave for the rotational wave of the microstructures. Whereas, the displacement wave can play a role in rotational wave transmission. Hence, in this work, in order to achieve a complete analogy between a linear asymmetric elastic medium and electrodynamics, we first generalize the CT by adding vibration degrees of freedom of microstructures. Then, using the gauge property of the displacement wave, we modify the CT and proceed in a way that charge and electromagnetic field to have independent nature.

II. THE COSSERAT THEORY IN MINKOWSKI METRIC

In the absence of any external force and torque, the corresponding dynamical equations for any homogeneous and isotropic linear asymmetric elastic environment, in the Minkowski metric, can be written [8] as¹

$$\rho u^{i,0} + \sigma^{ji}{}_{,j} = 0, \quad (1)$$

$$\mathfrak{S} \varphi^{i,0} + \epsilon^{ijk} \sigma_{jk} + \mathbf{m}^{ji}{}_{,j} = 0, \quad (2)$$

where ρ is constant average mass density of environment, \mathfrak{S} is the density of moment of inertia, σ^{ij} is the stress tensor, \mathbf{m}^{ij} is the couple-stress tensor, and u^i and φ^i respectively are the displacement vector and the rotation pseudovector of any point of environment. As the displacement and rotation satisfy wave equations and are defined over the entire elastic environment, we also refer to these as waves or fields. The Latin lowercase letters run from one to three, and the unit of speed has been selected to be one.

As Eqs. (1) and (2) contain four unknown parameters, in order to proceed, we appeal to definitions of the deformation tensors²

$$\Upsilon_{ij} \equiv u_{j,i} - \epsilon_{ijk} \varphi^k \quad \text{and} \quad \chi_{ij} \equiv \varphi_{j,i}. \quad (3)$$

In addition, when an environment behaves linearly, it has been shown [8] that one obtains

$$\sigma^{ij} = (\alpha + \mu) \Upsilon^{ij} + (\mu - \alpha) \Upsilon^{ji} + \lambda \delta^{ij} \Upsilon^k{}_k, \quad (4)$$

$$\mathbf{m}^{ij} = (\gamma + \varepsilon) \chi^{ij} + (\gamma - \varepsilon) \chi^{ji} + \beta \delta^{ij} \chi^k{}_k. \quad (5)$$

¹ In Ref. [8], these equations have been written in the Euclidean three-dimensional space, however, we have rewritten those based on the Minkowski metric with signature +2.

² Υ_{ij} has also been called strain tensor.

* M.Yousefian@sbu.ac.ir

† m-farhoudi@sbu.ac.ir

The six parameters α , μ , γ , ε , λ and β are characteristic coefficients of an elastic medium. These coefficients are measured in an adiabatic mode and are constants in a homogeneous environment, in a way that, the first four ones have positive values and the last two ones are required to satisfy the inequalities $3\lambda + 2\mu > 0$ and $3\beta + 2\gamma > 0$ [8]. Accordingly, substituting definitions (4) and (5) with relation (3) into Eqs. (1) and (2), these equations read³

$$\rho u^{i,0}_0 + (\alpha + \mu)u^{i,j}_j + (\mu - \alpha + \lambda)u^{j,i}_j + 2\alpha\epsilon^{ijk}\varphi_{j,k} = 0, \quad (6)$$

$$\mathfrak{S}\varphi^{i,0}_0 + (\gamma + \varepsilon)\varphi^{i,j}_j + (\gamma - \varepsilon + \beta)\varphi^{j,i}_j - 4\alpha\varphi^i - 2\alpha\epsilon^{ijk}u_{j,k} = 0. \quad (7)$$

III. FOUR-DIMENSIONAL REPRESENTATION

At this stage, we furthermore intend to extend the above obtained equations in a four-dimensional space-time.

For this purpose, first to introduce a zeroth component for the displacement, we utilize a scalar field in analogy with the issue of longitudinal waves. In there, a scalar field (as a kinetic potential) has been used to explain some parameters⁴ including the condensation,⁵ which is defined as minus the time derivative of it multiplied by the inverse square of the longitudinal component of the wave speed, v_L , [9]. Also, in the linear acoustics, it has been shown [10] that the condensation is equal to minus the divergence of the displacement in the Euclidean space. Thus, for such a scalar field/kinetic potential, say ϕ , one obtains

$$\phi_{,0} - v_L^2 u^i_{,i} = 0. \quad (8)$$

Incidentally, the longitudinal component of the wave speed is related to the elastic environment characteristic coefficients (the Lamé constants) as $v_L^2 \equiv (\lambda + 2\mu)/\rho$ [11]. With these respects, we rewrite relation (8) as

$$u^{\hat{\alpha}}_{,\hat{\alpha}} = 0 \quad (9)$$

with metric $g^{\hat{\alpha}\hat{\beta}} \equiv (-v_L^{-2}, 1, 1, 1)$ for hat-letters, where the kinetic potential has been assumed as zeroth component of the displacement, $\phi \equiv u_0$, and the Greek lowercase letters run from zero to three.

IV. MULTI-METRIC INSTEAD OF MULTI-SPEED

For simplicity of the appearances, and also for clarity, of the resulted equations, to raise down-indices of any tensor, we manifestly define five different diagonal metrics specified with normal-letters, hat-letters, bar-letters, check-letters and tilde-letters through the work. These metrics only differ with each other in their 00 components, which are related to different speeds (like $g^{\hat{\alpha}\hat{\beta}}$), rather than as in the usual natural units, assuming those to be one. In this respect, we also assume that the down-indices of any tensor (obviously, except the corresponding metrics themselves) being the same with regard to all of these metrics, say for example $T_\alpha = T_{\hat{\alpha}} = T_{\bar{\alpha}} = T_{\check{\alpha}} = T_{\tilde{\alpha}}$. This restriction makes only changes in the up-zeroth components being different, e.g. $T^{\hat{0}} = v_L^{-2}T^0$. However, for symmetry of the appearances of the relations, wherein the summation rule is used, we specify a down-index with the same notation as its corresponding up-one, e.g. as mentioned in relation (9).

With known definitions

$$\varphi^{ij} \equiv \epsilon^{ijk}\varphi_k \quad \text{and} \quad F^{\alpha\beta} \equiv u^{\alpha,\beta} - u^{\beta,\alpha}, \quad (10)$$

while using relation (9), we can rewrite Eq. (6) as

$$(\alpha + \mu)F^{i\hat{\alpha}}_{,\hat{\alpha}} - 2\alpha\varphi^{ij}_{,j} = 0 \quad (11)$$

with metric $g^{\bar{\alpha}\bar{\beta}} \equiv (-v_T^{-2}, 1, 1, 1)$ for bar-letters, where $v_T^2 \equiv (\alpha + \mu)/\rho$ is the transverse component of the wave speed of divergence-free displacement waves.

V. GENERALIZATION OF CT VIA NEW DEGREES OF FREEDOM

We secondly need to introduce another three components for the antisymmetric rotation wave φ^{ij} , namely φ^{0i} . For this purpose, as the moment of inertia density is proportional to the mass density of environment, thus varying it will cause that to change. The simplest way to explain variation of moment of inertia density is to assume the microstructures of medium as two same balls, each with mass m , connected by a massless spring, i.e. a 3-dimension isotropic harmonic oscillator where its symmetry group is isomorphic to $SO(4)$. To show this, let us write the energy of these mentioned two balls as the energy of another two balls, each also with mass m in a 4-dimension Euclidean space, connected by a massless rod with length $2R$, which are rotating around their center of mass. That is, a 4-dimension linear rigid rotor with

³ In Ref. [8], the term $4\alpha\varphi^i$ in Eq. (7) is missing, and it has been stated that total initial conditions have been marked with a symbol therein. However, an initial condition is obviously something different from a term in field equations, besides the mentioned term is a field, which its existence changes the nature of rotation wave to a massive wave (as it will be described below). Also, field equations have to be the same independently of any initial conditions, hence for simplicity, we consider initial conditions to be zero.

⁴ In linear acoustics, a real potential has been presented in a way that the turbulence of pressure and mass density, and speed can be provided in terms of it [9].

⁵ The ratio of difference between the instantaneous and the equilibrium mass densities to the equilibrium mass density of a medium, at a point, is called condensation [10].

total energy E and angular momentum $l_{(4)}$ of each ball, where $l_{(4)}^2 \equiv l^{AB}l_{AB}/2$ and the Latin uppercase letters run from one to four. Then, one can easily show that

$$\frac{l_{(4)}^2 + l^2}{2mR^2} = \frac{mr^2}{2} + \frac{Er^2}{R^2} + \frac{l^2}{2mr^2}, \quad (12)$$

where l and r respectively are the 3-dimension angular momentum of each ball (with $l^2 = l^{ij}l_{ij}/2$) and the distance between two balls in 3-dimension. To prove relation (12), note that $x^A x_A = R^2$ and $\dot{x}^A \dot{x}_A = 2E/m$ and $l^{AB} \equiv mx^A \dot{x}^B - mx^B \dot{x}^A$. This relation indicates that the symmetry group of the isotropic harmonic oscillator in 3-dimension is isomorphic to $SO(4)$.

In this way, we assume the microstructures of medium as 4-dimension linear rigid rotor with 4-dimension angular rotation $\varphi^{\alpha\beta}$, which φ^{0i} 's (as vibration and relative velocity of parts of each microstructure) are related to φ^{4i} 's (as rotation in fourth dimension) via the Wick rotation. On the other hand, by selecting an appropriate gauge, one can specify the type of environment and properties of waves within it. Hence, we choose such a gauge that causes the displacement and rotation wave equations of motion to be covariant, and in general, the corresponding waves velocities, masses and coupling (to the displacement field) of φ^{0i} being different from those of φ^{ij} . In this respect, we assume gauge

$$\varphi^{\tilde{\gamma}\tilde{\alpha}}_{,\tilde{\gamma}\tilde{\beta}} - \varphi^{\tilde{\gamma}\tilde{\beta}}_{,\tilde{\gamma}\tilde{\alpha}} = 0 \quad (13)$$

with metric $g^{\tilde{\alpha}\tilde{\beta}} = (-v_{\tilde{V}}^{-2}, 1, 1, 1)$ for check-letters, where $v_{\tilde{V}}^2 \equiv (\gamma + \varepsilon)/\mathfrak{S}$ is the wave speed of vibration φ^{0i} waves. Then, we plausibly generalize Eq. (7) as

$$(2\gamma + \beta) \left(\varphi^{\tilde{\alpha}\tilde{\beta}}_{,\tilde{\gamma}\tilde{\gamma}} + \varphi^{\tilde{\gamma}\tilde{\alpha}}_{,\tilde{\gamma}\tilde{\beta}} - \varphi^{\tilde{\gamma}\tilde{\beta}}_{,\tilde{\gamma}\tilde{\alpha}} \right) - 4\alpha\varphi^{\tilde{\alpha}\tilde{\beta}} - 2\alpha F^{\tilde{\alpha}\tilde{\beta}} = 0 \quad (14)$$

with metric $g^{\tilde{\alpha}\tilde{\beta}} = (-v_{\tilde{R}}^{-2}, 1, 1, 1)$ for tilde-letters, where $v_{\tilde{R}}^2 \equiv (2\gamma + \beta)/\mathfrak{S}$ is the wave speed of φ^{ij} rotation waves. In Eq. (14), the part

$$(2\gamma + \beta) \left(\varphi^{\tilde{i}\tilde{j}}_{,\tilde{\gamma}\tilde{\gamma}} + \varphi^{\tilde{\gamma}\tilde{i}}_{,\tilde{\gamma}\tilde{j}} - \varphi^{\tilde{\gamma}\tilde{j}}_{,\tilde{\gamma}\tilde{i}} \right) - 4\alpha\varphi^{\tilde{i}\tilde{j}} - 2\alpha F^{\tilde{i}\tilde{j}} = 0, \quad (15)$$

using gauge (13), is just Eq. (7), and the extra part

$$(2\gamma + \beta) \left(\varphi^{\tilde{0}\tilde{i}}_{,\tilde{j}\tilde{j}} + \varphi^{\tilde{j}\tilde{0}}_{,\tilde{j}\tilde{i}} - \varphi^{\tilde{j}\tilde{i}}_{,\tilde{j}\tilde{0}} \right) - 4\alpha\varphi^{\tilde{0}\tilde{i}} - 2\alpha F^{\tilde{0}\tilde{i}} = 0, \quad (16)$$

again due to gauge (13), is the wave equation of vibration φ^{0i} , i.e.

$$(\gamma + \varepsilon) \varphi^{\tilde{0}\tilde{i}}_{,\tilde{\gamma}\tilde{\gamma}} - 4\alpha\varphi^{\tilde{0}\tilde{i}} - 2\alpha F^{\tilde{0}\tilde{i}} = 0. \quad (17)$$

Eq. (14) is invariant under a gauge transformation with

respect to an arbitrary gauge field, say ζ_{α} , as

$$\begin{aligned} \varphi^{\tilde{\alpha}\tilde{\beta}} &\rightarrow \varphi^{\tilde{\alpha}\tilde{\beta}} - \left(\zeta^{\tilde{\alpha},\tilde{\beta}} - \zeta^{\tilde{\beta},\tilde{\alpha}} \right) \\ F^{\tilde{i}\tilde{j}} &\rightarrow F^{\tilde{i}\tilde{j}} + 2 \left(\zeta^{\tilde{i},\tilde{j}} - \zeta^{\tilde{j},\tilde{i}} \right) \\ F^{\tilde{i}\tilde{0}} &\rightarrow F^{\tilde{i}\tilde{0}} + 2 \frac{v_{\tilde{T}}^2}{v_{\tilde{R}}^2} \left(\zeta^{\tilde{i},\tilde{0}} - \zeta^{\tilde{0},\tilde{i}} \right). \end{aligned} \quad (18)$$

This gauge gives us a cue to manage Eq. (11) being also invariant under a gauge transformation with respect to another arbitrary gauge field, say ξ_{α} , as

$$\begin{aligned} \varphi^{\tilde{\alpha}\tilde{\beta}} &\rightarrow \varphi^{\tilde{\alpha}\tilde{\beta}} + \xi^{\tilde{\alpha},\tilde{\beta}} - \xi^{\tilde{\beta},\tilde{\alpha}} \\ F^{\tilde{i}\tilde{j}} &\rightarrow F^{\tilde{i}\tilde{j}} + \frac{2\alpha}{\alpha + \mu} \left(\xi^{\tilde{i},\tilde{j}} - \xi^{\tilde{j},\tilde{i}} \right) \\ F^{\tilde{i}\tilde{0}} &\rightarrow F^{\tilde{i}\tilde{0}} + \frac{2\alpha v_{\tilde{T}}^2}{(\alpha + \mu)v_{\tilde{R}}^2} \left(\xi^{\tilde{i},\tilde{0}} - \xi^{\tilde{0},\tilde{i}} \right), \end{aligned} \quad (19)$$

provided that we generalize Eq. (11) in a more general form

$$(\alpha + \mu) F^{i\tilde{\alpha}}_{,\tilde{\alpha}} - 2\alpha\varphi^{i\tilde{\alpha}}_{,\tilde{\alpha}} = 0, \quad (20)$$

however with $\varphi^{0i} = 0$ in order to reproduce Eq. (11). The generalization of adding such a degree of freedom (i.e., φ^{0i}) is physically meaningful. The reason is that, F^{0i} is composed of the velocity of microstructures and gradient of the kinetic potential (that, in turn, is related to change of the density of microstructures). On the other hand, the vibration φ^{0i} has also two features, namely the relative velocity of parts of each microstructure and alteration in the volume (and hence the density) of each one. Thus, F^{0i} and φ^{0i} have a similar role, which is missing in Eq. (11).

Now, performing the divergence of Eq. (20) yields

$$(\alpha + \mu) F^{i\tilde{0}}_{,\tilde{0}i} - 2\alpha\varphi^{i\tilde{0}}_{,\tilde{0}i} = 0. \quad (21)$$

Integrating Eq. (21) over time while considering the zeroth component of the displacement as a wave, it gives⁶

$$(\alpha + \mu) F^{\tilde{0}\tilde{\alpha}}_{,\tilde{\alpha}} - 2\alpha\varphi^{\tilde{0}\tilde{\alpha}}_{,\tilde{\alpha}} = 0. \quad (22)$$

Combining Eqs. (20) and (22), we finally end up with a generalized form of Eq. (11) as

$$(\alpha + \mu) F^{\tilde{\beta}\tilde{\alpha}}_{,\tilde{\alpha}} - 2\alpha\varphi^{\tilde{\beta}\tilde{\alpha}}_{,\tilde{\alpha}} = 0. \quad (23)$$

It would be instructive to respectively rewrite gauge conditions (9) and (13), and Eqs. (14) and (23) in the exterior algebra notations,⁷ namely

$$\delta\hat{\mathbf{u}} = 0 \quad \text{and} \quad \check{d}\check{\check{\varphi}} = 0, \quad (24)$$

⁶ Eq. (22), without the second interaction term while using relation (9), explains the kinetic potential as a wave [9].

⁷ See, e.g., Ref. [12].

$$(2\gamma + \beta)\tilde{\delta}\tilde{d}\tilde{\varphi} - 4\alpha\tilde{\varphi} - 2\alpha\tilde{d}\tilde{\mathbf{u}} = 0, \quad (25)$$

$$(\alpha + \mu)\tilde{\delta}\tilde{d}\tilde{\mathbf{u}} - 2\alpha\tilde{\delta}\tilde{\varphi} = 0, \quad (26)$$

where \mathbf{u} and φ are one-form and two-form fields, respectively.

VI. MODIFICATION OF GENERALIZED CT VIA GAUGE PROPERTY OF DISPLACEMENT WAVE

As the motion of an environment can make the motion of a wave within it ineffective, it is noteworthy to mention that when the displacement wave moves a microstructure of medium, then the microstructure movement can make the movement of rotation wave (the gradient or the momentum of rotation wave) ineffective. This matter means that the displacement field is capable to play the role of the momentum of rotation wave, similar to the photon field that acts as the role of the momentum or a gauge for the fermion field. However, there is no trace of such a feature in Eqs. (25) and (26). Hence, we still need to modify the generalized CT further in order to present such a gauge characteristic. For this purpose, let us first check whether the existed generalized CT without the interacting terms between the displacement wave and rotation wave (i.e., without the last terms in Eqs. (25) and (26)), is related to theory of Yang-Mills or not. If the result being positive, then we will need only to modify the interaction terms.

The non-interactive part of Eq. (26) with its gauge condition is well-known to be in accord with boson fields. For the other equation, it is also known [13] that a spinor field, say ψ , can be defined by a two-form field as

$$\psi \equiv \varphi_{\alpha\beta}\sigma^{\alpha\beta}v, \quad (27)$$

where $\sigma^{\alpha\beta} = \frac{i}{2}[\gamma^\alpha, \gamma^\beta]$, γ^α 's are the Dirac matrices and v is an arbitrary constant nonzero fiducial spinor for which $\gamma_5 v \neq iv$. Multiply the transpose conjugate of v to a Klein-Gordon equation for $\tilde{\psi} \equiv \varphi_{\tilde{\alpha}\tilde{\beta}}\sigma^{\tilde{\alpha}\tilde{\beta}}v$, i.e.

$$v^\dagger \left(i\tilde{\not{D}} + m \right) \left(i\tilde{\not{D}} - m \right) \tilde{\psi} = 0. \quad (28)$$

Then, substituting for $\tilde{\psi}$ while using [13]

$$8vv^\dagger = 2v^\dagger\gamma_\alpha v\gamma^\alpha + \omega_{\alpha\beta}\sigma^{\alpha\beta}, \quad (29)$$

in the exterior algebra, it yields

$$\tilde{\omega} \wedge * \left(\tilde{\delta}\tilde{d}\tilde{\varphi} - m^2\tilde{\varphi} \right) = 0, \quad (30)$$

with $\tilde{\omega}$ as an arbitrary two-form defined as $\omega^{\alpha\beta} \equiv v^\dagger\sigma^{\alpha\beta}v$ that, in turn while assuming $m \equiv \sqrt{4\alpha}/(2\gamma + \beta)$, gives the non-interactive part of Eq. (25). On the other hand, if the Dirac equation for a massive fermion, i.e.

$$\left(i\tilde{\not{D}} - m \right) \tilde{\psi} = 0, \quad (31)$$

is satisfied, then its corresponding Klein-Gordon equation and, in turn, Eq. (30) will be held.⁸ To check the corresponding gauge condition, i.e. the second one in gauges (24), we use the trivial condition

$$v^\dagger \partial_{\tilde{\alpha}} \partial_{\tilde{\beta}} \sigma^{\tilde{\alpha}\tilde{\beta}} \tilde{\psi} = 0, \quad (32)$$

which, in turn by relation (29), gives

$$\tilde{\omega} \wedge * \left(\tilde{\delta}\tilde{d}\tilde{\varphi} \right) = 0, \quad (33)$$

where, as $\tilde{\omega}$ is arbitrary, yields the desired result. In addition, the rotation field has two states of right and left, and analogously, the spin 1/2 of fermions with its gyromagnetic ratio causes the effective amount of spin, in reaction to a magnetic field, being ± 1 . Therefore, the non-interacting QED equations result the non-interacting generalized CT.

With respect to the gauge characteristic mentioned before, regarding the necessity of modifying the generalized CT, if we choose Eq. (31) to be modified as

$$\left(i\tilde{\not{D}} - e\tilde{\not{A}} - m \right) \tilde{\psi} = 0, \quad (34)$$

then consequently, Eq. (25) will be amended as

$$(2\gamma + \beta)\tilde{\Delta}\tilde{D}\tilde{\varphi} - 4\alpha\tilde{\varphi} = 0, \quad (35)$$

where e is a constant parameter depended on the characteristics of the elastic wave of environment, $D \equiv d + ie\mathbf{u}$ and $\Delta \equiv *D*$. Eq. (34) is invariant under $U(1)$ gauge transformation, which indicates that the displacement wave plays the role of momentum for the rotation wave. Moreover, we need to modify Eq. (23) in such a way that also remains invariant under $U(1)$. Accordingly, we choose

$$2(\alpha + \mu)F^{\tilde{\alpha}\tilde{\beta}}{}_{,\tilde{\beta}} + e\tilde{\psi}\gamma^{\tilde{\alpha}}\tilde{\psi} = 0. \quad (36)$$

With the above choices, Eqs. (34) and (36) can be gained from the variation of Lagrangian

$$L = \tilde{\psi} \left(i\tilde{\not{D}} - e\tilde{\not{A}} - m \right) \tilde{\psi} - \frac{1}{2}(\alpha + \mu)F^{\tilde{\alpha}\tilde{\beta}}F_{\tilde{\alpha}\tilde{\beta}}, \quad (37)$$

with respect to $\tilde{\psi}$ and $u_{\tilde{\alpha}}$, where $\tilde{\psi}$ is the adjoint of $\tilde{\psi}$.

VII. QED TREATMENT

Now, by employing the exchange of variables as

$$\sqrt{2(\alpha + \mu)}u^\alpha \longrightarrow u^\alpha \quad \text{and} \quad e/\sqrt{2(\alpha + \mu)} \longrightarrow e, \quad (38)$$

⁸ The comparison of fermions and rotation waves has also been considered in Refs. [14–16]. Besides, the anticommutation nature of spinor field (27) has been described in Ref. [17], which guarantees the characteristics of the Fermi-Dirac statistics of rotational waves.

Lagrangian (37) can be rewritten as

$$L = \bar{\psi} \left(i\tilde{\not{\partial}} - e\tilde{\not{\eta}} - m \right) \tilde{\psi} - \frac{1}{4} F^{\alpha\beta} F_{\alpha\beta}. \quad (39)$$

This Lagrangian is equivalent to the QED Lagrangian for photon with zero rest-mass and fermion with mass m and negative charge e [18], except that the metrics of different parts of it are different. However, for a specific elastic environment via the thermodynamic characteristics, namely the one with properties

$$\alpha = \mu + \lambda, \quad \varepsilon = \gamma + \beta \quad \text{and} \quad \frac{\alpha + \mu}{\rho} = \frac{\gamma + \varepsilon}{\mathfrak{S}}, \quad (40)$$

this equivalence is exact. In Lagrangian (39), the displacement vector u^α is analogous with the electromagnetic potential four-vector in electrodynamics, and the tensor $\varphi^{\alpha\beta}$ is comparable to the fermion spin tensor in QED. In particular, components φ^{0i} are analogous with the electric dipole ones and components φ^{ij} are comparable to the magnetic dipole ones.

VIII. EMPIRICAL ASSESSMENT

Let us justify our modifications. In this regard, from experimental point of view, a report has been presented [19] on the empirical assessment of the CT. Therein, a comparison has been made between the test results with predictions of the CT and also a discrete lattice model [20]. At last, it has been shown that the CT is at least inadequate. For our purpose, we have compared the experimental results of [19] with our amended CT.

In Fig. 3 of Ref. [19] (repeated here as Fig. 1 with some subregions marked on it), two regions have been specified. The first region is the low-frequency region (0 – 80 kHz) or the L-TR region. In this region, in two subregions of less than 20 kHz and 20 – 50 kHz (the subregions 1 and 2 in Fig. 1, respectively), the intensity of wave is maximal, and the arrival-time is almost constant with respect to the variations of frequency (as shown by the arrows in the subregions 1 and 2). Thus, the environment for the displacement wave is transparent and the displacement wave has constant speed and is massless. According to their figure, the subregion of less than 20 kHz is related to the longitudinal wave and the subregion 20 – 50 kHz corresponds to the shear wave that (due to the intensity of wave in these two subregions) is linearly coupled to the longitudinal wave. In the subregion 50 – 80 kHz (the subregion 3 in Fig. 1), in addition to the decrement of the wave intensity, the wave arrival-time increases with the increment of the wave frequency (as shown by the arrows in the subregion 3) and thus, the wave speed decreases. This situation is similar to when light travels in an environment in which its permittivity and permeability, due to the induction of electric and magnetic dipoles, increase. In analogous, this situation means that in this subregion, components φ^{0i} and φ^{ij}

get excited and the wave u^α gradually decays into the wave $\varphi^{\alpha\beta}$. Such a decay is visible in Fig. 2b of Ref. [19] at the end of this region. Also, this decay is similar to decay of high-energy cosmic gamma-rays in collision with low-energy background cosmic lights and their pair production. In Fig. 9 of Ref. [21], the spectral energy distribution of the Crab Nebula from soft to very high-energy γ -rays has been depicted, which is similar to Fig. 2b of Ref. [19] that shows transmittance measured in the granular crystal. This point indicates similarity between γ -rays treatment and our amended CT.




Figure 1. (colored online) This figure (the Fig 3 of Ref. [19] with some marks on it) shows the arrival-time of received signals from a shear transducer (left) and a longitudinal transducer (right) in terms of frequency and intensity after transmission through a crystal, that we have identified thereon the variations of the wave velocity based on the frequency variation.

The second region in Fig. 3 of Ref. [19] is the high-frequency region (80 – 150 kHz, the subregion 4 in Fig. 1) or the RT-region. In this region, waves show different behavior from the first region. That is, while the intensity of waves are very low, the wave arrival-time decreases with the increment of frequency (as shown by the arrows in the left-hand side of Fig. 1) and thus, the wave speed increases. This issue means waves in this region are massive. Hence, the hypothesis of the decay of massless waves u^α into massive waves $\varphi^{\alpha\beta}$ is confirmed, although due to the high mass of the vibrational waves φ^{i0} , there is no trace of this wave in the subregion 4 in the left-hand side of Fig. 1. Also, the comparison of difference in the intensity of wave, in the subregions 1 and 4 in Fig. 1, indicates that the wave intensity of $\varphi^{\alpha\beta}$ is approximately the square of the wave intensity of u^α . This result displays that, contrary to the Cosserat and discrete models, $\varphi^{\alpha\beta}$ wave is non-linearly coupled to u^α wave, which is consistent with our modification of interaction parts of the generalized CT.

Another point, which is not a new one, is how to apply a continuous theory to a discrete environment. In this respect, when one quantizes a theory, one needs an energy scale to get a renormalized theory. Such an energy scale should be proportional to the size of grains in a granular

system. If the frequency of wave in a granular medium is close to such an energy scale, it will be equivalent to when the wavelength approaches to the size of system grains. In this situation, loop-corrections should be considered and the coupling constant of U(1) gauge theory running and being a function of energy of system [18]. Such a situation is equivalent with an effective higher-gradient theory with increasing energy levels as it had been suggested in Ref. [19] that the CT should be combined with higher-gradient theories like those in Ref. [22], however therein, such terms have been manually added.

IX. CONCLUSIONS

In conclusion, we have noticed some logical gaps in the Cosserat [5] and discrete models [20], such as disregarding the possibility of degree of vibrational freedom of microstructures and also the gauge property of u^α wave for $\varphi^{\alpha\beta}$ wave. Hence, we first have generalized the CT in a four-dimensional form by adding degrees of freedom of vibration of microstructures. Then, via a gauge property of the displacement wave relative to rotation wave, we have logically amended the existing interaction terms

instead of somehow modifying those manually. Thus, we have established a fruitful analogy between different subfields of physics and have shown that in asymmetric elastic environments, the elastic wave equations are analogous to the QED ones. Hence on macro scales, elastic waves can possess the QED treatment.⁹ This analogy provides a new paradigm of fermions and bosons. Moreover, from the experimental point of view, due to the analogy between wave intensity of $\varphi^{\alpha\beta}$ and u^α , we have indicated that, contrary to the Cosserat and discrete models, our amended CT are consistent with the corresponding empirical observations. In this study, the elastic environment resembles an ether itself in an etheral model¹⁰ and the elastic waves resemble electromagnetic waves in the ether. Indeed, in another work [25], we have proposed an etheral model based on a third kind of quantization approach [26], in which electromagnetic waves analogize elastic waves in an elastic environment.

ACKNOWLEDGEMENTS

We thank the Research Council of Shahid Beheshti University.

-
- [1] W. Thomson, "On the propagation of laminar motion through a turbulently moving inviscid liquid", *Philos. Magaz. Ser. 5* (1887), 342.
- [2] V.P. Dmitriyev, "The elastic model of physical vacuum", *Mech. Solids (N.Y.)* **26** (1992), 60.
- [3] V.P. Dmitriyev, "Towards an exact mechanical analogy of particles and fields", *Nuov. Cim.* **111A** (1998), 501.
- [4] V.P. Dmitriyev, "Elasticity and electromagnetism (the Coulomb gauge)", *Electromag. Phenomena* **2** (2001), 474.
- [5] E. Cosserat and F. Cosserat, "*Théorie des Corps Déformables* (Theory of Deformable Bodies)", (Herman, Paris, 1909).
- [6] E.B. Wilson, "An advance in theoretical mechanics", *Bull. Am. Math. Soc.* **19** (1913), 242.
- [7] R.A. Toupin, "Theories of elasticity with couple-stress", *Arch. Rational Mech. Anal.* **17** (1964), 85.
- [8] W. Nowacki, "The linear theory of micropolar elasticity", In: "*Micropolar Elasticity*", Symposium organized by the department of mechanics of solids, June 1972, Edited by: W. Nowacki and W. Olszak, (Springer, Vienna, 1974), pp. 1-43.
- [9] D. Stanzial, D. Bonsi and G. Schiffrer, "Four-dimensional treatment of linear acoustic fields and radiation pressure", *Act. Acust. Un. Acustica* **88** (2002), 213.
- [10] L.E. Kinsler, A.R. Frey, A.B. Coppens and J.V. Sanders, "*Fundamentals of Acoustics*", (Wiley, New York, 1999).
- [11] J.G. Harris, "*Linear Elastic Waves*", (Cambridge University Press, Cambridge, 2004).
- [12] N. Straumann, "*General Relativity with Applications to Astrophysics*", (Springer, Berlin, 2004).
- [13] J.R. Klauder, "Linear representation of spinor fields by antisymmetric tensors", *J. Math. Phys.* **5** (1964), 1204.
- [14] J. Burnett and D. Vassiliev "Modeling the electron with Cosserat elasticity", *Mathematika* **58** (2012), 349.
- [15] C.T. Chan, Z.H. Hang and X. Huang, "Dirac dispersion in two-dimensional photonic crystals", *Adv. OptoElectron* **2012** (2012), 313984.
- [16] R.A. Close, "Exact description of rotational waves in an elastic solid", *Adv. Appl. Clifford Algeb.* **21** (2011), 273.
- [17] P.A. Deymier, K. Runge, N. Swintek and K. Muralidharan, "Torsional topology and fermion-like behavior of elastic waves in phononic structures", *Comptes Rendus Mécanique* **343** (2015), 700.
- [18] M.E. Peskin and D. Schroeder, "*An Introduction to Quantum Field Theory*", (Westview, New York, 1995).
- [19] A. Merkel, V. Tournat and V. Gusev, "Experimental evidence of rotational elastic waves in granular phononic crystals", *Phys. Rev. Lett.* **107** (2011), 225502.
- [20] A. Merkel, V. Tournat and V. Gusev, "Dispersion of elastic waves in three-dimensional noncohesive granular phononic crystals: Properties of rotational modes", *Phys. Rev. E* **82** (2010), 031305.
- [21] A.A. Abdo, *et al.*, "Fermi large area telescope observations of the Crab pulsar and nebula", *Astrophys. J.* **708** (2010), 1254.
- [22] A.S.J. Suiker, R. de Borst and C.S. Chang, "Micro-mechanical modeling of granular material", *Acta Mech.* **149** (2001), 161.

⁹ Also, in Ref. [23], we have shown a kind of symmetry that statistically implies uniformity of physics in large and small scales.

¹⁰ For a review on the ether, see, e.g., Ref. [24] and references therein.

- [23] M. Yousefian and M. Farhoudi, “Towards real amplituhedron via one-dimensional theory”, *arXiv:1910.05548*.
- [24] M. Farhoudi and M. Yousefian, “Ether and relativity”, *Int. J. Theor. Phys.* **55** (2016), 2436.
- [25] M. Yousefian and M. Farhoudi, “Fine structure constant variation and ultra high energy cosmic rays via a novel ethereal model”, work in progress.
- [26] M. Yousefian and M. Farhoudi, “Dark matter and hierarchy problems via vacuum quantization”, submitted to journal.