

GENERALIZED LOCALIZATION FOR SPHERICAL PARTIAL SUMS OF THE MULTIPLE FOURIER SERIES AND INTEGRALS

RAVSHAN ASHUROV

ABSTRACT. It is well known, that Luzin's conjecture has a positive solution for one dimensional trigonometric Fourier series and it is still open for the spherical partial sums $S_\lambda f(x)$, $f \in L_2(\mathbb{T}^N)$, of multiple Fourier series, while it has the solution for square and rectangular partial sums. Historically progress with solving Luzin's conjecture has been made by considering easier problems. One of such easier problems for $S_\lambda f(x)$ was suggested by V. A. Il'in in 1968 and this problem is called the generalized localization principle. In this paper we first give a short survey on convergence almost-everywhere of Fourier series and on generalized localization of Fourier integrals, then present a positive solution for the generalized localization problem for $S_\lambda f(x)$.

AMS 2000 Mathematics Subject Classifications : Primary 42B05; Secondary 42B99.
Key words: Multiple Fourier series and integrals, spherical partial sums, convergence almost-everywhere, generalized localization.

1. INTRODUCTION

Let $\{f_n\}$, $n \in \mathbb{Z}^N$, be the Fourier coefficients of a function $f \in L_2(\mathbb{T}^N)$, $N \geq 2$, i.e.

$$f_n = (2\pi)^{-N} \int_{\mathbb{T}^N} f(y) e^{-iny} dy,$$

where \mathbb{T}^N is N -dimensional torus: $\mathbb{T}^N = (\pi, \pi]^N$. Consider the spherical partial sums of the multiple Fourier series:

$$(1.1) \quad S_\lambda f(x) = \sum_{|n|^2 < \lambda} f_n e^{inx},$$

where $nx = n_1x_1 + n_2x_2 + \dots + n_Nx_N$ and $|n| = \sqrt{n_1^2 + n_2^2 + \dots + n_N^2}$.

The aim of this paper is to investigate convergence almost-everywhere (a.e.) of these partial sums. One of the first questions which arises in the study of a.e. convergence of the sums (1.1) is the question of the validity of the Luzin conjecture: is it true that the spherical sums (1.1) of N -fold Fourier series of an arbitrary function $f \in L_2(\mathbb{T}^N)$ converge a.e. on \mathbb{T}^N ? In other words, does Carleson's theorem extend to spherical partial sums (1.1)? The answer to this question is unknown so far (see [1] and [2]). What is known is only that Hunt's theorem does not extend to partial sums (1.1) [3]. Namely, for each $p \in [1, 2)$ there exists a function $f \in L_p(\mathbb{T}^N)$ such that on a set of positive measure

$$\overline{\lim}_{\lambda \rightarrow \infty} |S_\lambda f(x)| = +\infty.$$

As the authors of paper [3] showed, this result is a consequence of Fefferman's celebrated theorem to the effect that partial sums (1.1) does not converge in $L_p(\mathbb{T}^N)$, $1 \leq p < 2$.

Historically progress with solving the Luzin conjecture has been made by considering easier problems. One of such easier problems is the generalized principle of localization.

Il'in [4] was the first to introduce the concept of generalized localization principle for an arbitrary eigenfunction expansions. We present the corresponding definition for Fourier series; for Fourier integrals the definition is similar.

Definition 1.1. We say that for $S_\lambda f$ the generalized localization principle in function class $L_p(\mathbb{T}^N)$, $p \geq 1$, holds true, if for any function $f \in L_p(\mathbb{T}^N)$, with $f = 0$ on a open set $\Omega \subset \mathbb{T}^N$ the equality

$$\lim_{\lambda \rightarrow \infty} S_\lambda f(x) = 0$$

is valid for a.e. on Ω .

Recall that the validity of the classical Riemann localization principle means that the convergence at a point x_0 of the Fourier series of a function f depends only on the behavior of f in a small neighborhood of that point. More exactly, if $f = 0$ in an open set $\Omega \subset \mathbb{T}^N$ then the Fourier series of f converges to 0 at every point of Ω . For generalized localization one requires that the convergence to 0 occurs a.e. on Ω .

Why one should consider the generalized localization instead of the classical one in L_p -classes? According to the classical Riemann theorem, in the one-dimensional case, the localization principle is valid for any integrable function. The situation changes in the case of functions of two or more variables. In this case, there are examples of functions with high smoothness for which the classical principle of localization is not valid (see [1]). Therefore, the principle of generalized localization, introduced by Il'in, may become a replacement for the classical principle of localization.

Recently the problem of generalized localization was completely solved in $L_p(\mathbb{T}^N)$ classes [5]. In the present paper we first give a short survey on convergence a.e. of Fourier series, then indicate a sketch of the proof from [5].

2. CONVERGENCE ALMOST-EVERYWHERE OF FOURIER SERIES

1. One dimensional Fourier series. A problem of expanding of a function $f(t) \in L_1(\mathbb{T})$, $\mathbb{T} = (-\pi, \pi]$, in terms of trigonometric series

$$\sum a_k \cos kt + b_k \sin kt,$$

arises in mathematical physics and investigation of the convergence problems of such series is very important. Here a_k and b_k are as usual Fourier coefficients of function f .

Partial sums are defined as follows:

$$S_m(t) = \sum_{k=0}^m a_k \cos kt + b_k \sin kt.$$

If we define a new coefficients as $c_k = \frac{1}{2}(a_k - ib_k)$, $c_{-k} = \overline{c_k}$, then partial sums can be rewritten as

$$(2.1) \quad S_m(t) = \sum_{|k| \leq m} c_k e^{ikt}.$$

These partial sums can easily be generalized in multidimensional case.

In 1915 N.N.Luzin made the conjecture that the Fourier series of any function in $L_2(\mathbb{T})$ converges a.e.. In other words, if

$$\sum |c_k|^2 < \infty$$

then the series

$$\sum c_k e^{ikt}$$

converges for almost every $t \in \mathbb{T} = (-\pi, \pi]$.

More than 50 years the Luzin's conjecture attracted the attention of the specialists inducing a large number of investigations. In 1922 A.N.Kolmogorov constructed a remarkable example of a function $f \in L_1(\mathbb{T})$, whose Fourier series diverges a.e. and even at every point of \mathbb{T} (see [1] and [2]). After this result some specialists have doubted the validity of the the Luzin's conjecture, since Kolmogorov's counterexample in L_1 was unbounded in any interval, but it was thought to be only a matter of time before a continuous counterexample was found.

This long story has its end only in 1966 when Carleson proved that Luzin's problem has a positive solution. Now this is a fundamental result in mathematical analysis. It may be worth mentioning that even Carleson started (see [6]) by trying to find a continuous counterexample to the Luzin's conjecture making use of the Blaschke products and Zygmund was very positive

about that. But he realized eventually that his approach could not work. Only after this he tried instead to prove Luzin's conjecture.

Soon after two years (in 1968) R.A. Hunt, improving Carleson's method, showed that the Fourier series of any function $f \in L_p(\mathbb{T})$, $p > 1$, converges a.e. on \mathbb{T} . Thus the problem of convergence a.e. of Fourier series is completely solved in classes $L_p(\mathbb{T})$, $p \geq 1$: if $p > 1$ then we have the convergence and if $p = 1$, then the convergence a.e. fails.

2. Partial sums in multidimensional case. The multidimensional Fourier series has the form

$$(2.2) \quad \sum_{n \in \mathbb{Z}^N} f_n e^{inx}.$$

The definition (2.1) of partial sums generalizes to the multidimensional case in three obvious forms, which are often found in the literature.

Partial sums of series (2.2)

$$M_k f(x) = \sum_{|n_1| \leq k} \cdots \sum_{|n_N| \leq k} f_n e^{inx}$$

is called **square partial sums**. We say that the above series converges over squares if $\lim_{k \rightarrow \infty} M_k f(x)$ exists.

Second partial sums have the form

$$R_{k, \dots, l} f(x) = \sum_{|n_1| \leq k} \cdots \sum_{|n_N| \leq l} f_n e^{inx},$$

which is called **rectangular partial sums**. We say that series (2.2) converges over rectangles if

$$\lim_{\min\{k, \dots, l\} \rightarrow \infty} R_{k, \dots, l} f(x)$$

exists.

Finally, **spherical partial sums** of series (2.2) are defined as (see (1.1))

$$S_\lambda f(x) = \sum_{|n|^2 < \lambda} f_n e^{inx}.$$

If $\lim_{\lambda \rightarrow \infty} S_\lambda f(x)$ exists, then we say that series (2.2) converges over spheres.

All these partial sums are a natural generalization of one dimensional case (see (2.1)). But from spectral theory point of view, the natural generalization is the spherical partial sums. Note the eigenfunctions of the periodical Laplace operator are the functions $u_n(x) = (2\pi)^{-\frac{N}{2}} e^{inx}$, while the eigenvalues equal $\lambda_n = |n|^2$. Thus eigenfunction expansions of function f have the form (1.1).

3. Convergence almost-everywhere. In 1970 N. R. Tevzadze [7] has extended Carleson's theorem to double Fourier series with square partial sums. The idea of his proof is very simple. Indeed, let $f \in L_2(\mathbb{T}^2)$ be expanded in a double Fourier series whose quadratic partial sums equal

$$M_k f(x_1, x_2) = \sum_{|n_1| \leq k} \sum_{|n_2| \leq k} f_{n_1, n_2} e^{i(n_1 x_1 + n_2 x_2)}.$$

One can rewrite this sum as a sum of two "one dimensional" partial sums

$$(2.3) \quad M_k f(x_1, x_2) = \sum_{|n_1| \leq k} A_{n_1}(x_2) e^{in_1 x_1} + \sum_{|n_2| \leq k} B_{n_2}(x_1) e^{in_2 x_2},$$

where the coefficients have the form

$$A_{n_1}(x_2) = \sum_{|n_2| \leq |n_1|} f_{n_1, n_2} e^{in_2 x_2}, \quad B_{n_2}(x_1) = \sum_{|n_1| \leq |n_2|} f_{n_1, n_2} e^{in_1 x_1}.$$

It is not hard to prove, using Levi's theorem, that the series

$$\sum_{n_1 = -\infty}^{\infty} A_{n_1}^2(x_2)$$

converge for almost every $x_2 \in \mathbb{T}$. This means, that for almost every $x_2 \in \mathbb{T}$ we can consider the first of the sums in the right hand side of (2.3) as the partial sums of a Fourier series of a L_2 -

function of x_1 . Therefore by Carleson's theorem, this sum must converge for almost all $x_1 \in \mathbb{T}$. By the same way one can prove that the second sum in (2.3) for almost every x_1 converges in x_2 a.e. on \mathbb{T} , i.e.

$$\lim_{k \rightarrow \infty} M_k f(x_1, x_2) = f(x_1, x_2)$$

a.e. on \mathbb{T}^2 . Note that the convergence to f exactly follows from the fact that M_k converges to f in $L_2(\mathbb{T}^2)$.

In an analogous way one can prove, that

$$(2.4) \quad \lim_{k \rightarrow \infty} \widetilde{M}_k f(x_1, x_2) = f(x_1, x_2),$$

for almost all $(x_1, x_2) \in \mathbb{T}^2$, where

$$\widetilde{M}_k f(x_1, x_2) = \sum_{|n_1| \leq m_1(k)} \sum_{|n_2| \leq m_2(k)} f_{n_1, n_2} e^{i(n_1 x_1 + n_2 x_2)}$$

and $m_1(k)$ and $m_2(k)$ are arbitrary decreasing sequences of integers.

It should be noted that the proof presented in the above paper easily can be extended to N -dimensional case.

An analogue of the Hunt's theorem was obtained by P. Sjölin in 1971 [8].

Theorem (Sjölin). *If $f \in L_p(\mathbb{T}^N)$, $p > 1$, then one has $\lim_{k \rightarrow \infty} M_k f(x) = f(x)$ a.e. on \mathbb{T}^N .*

Note, in fact Sjölin proved this theorem for even wider class of functions f .

It is not hard to verify, that if f_1 is Kolmogorov's function, then square partial sums of the function $f(x_1, \dots, x_N) = f_1(x_1) \in L_1(\mathbb{T}^N)$ diverge a.e. on \mathbb{T}^N .

Thus the problem of convergence a.e. of square partial sums of multiple Fourier series is completely solved in classes $L_p(\mathbb{T}^N)$, $p \geq 1$: if $p > 1$ then we have the convergence and if $p = 1$, then the convergence a.e. fails. Hence, convergence a.e. of square partial sums of L_p -functions does not depend on the number of dimensions and result has exactly the same formulation for $N > 1$ as in one dimensional case. However, for rectangular partial sums due to C. Fefferman [9] we have a completely new phenomena.

Theorem (Charles Fefferman (1971)). *There exists a function $f \in C(\mathbb{T}^2)$, such that*

$$\overline{\lim}_{\min\{k, l\} \rightarrow \infty} |R_{k, l} f(x)| = +\infty$$

everywhere on \mathbb{T}^2 .

The divergence in Fefferman's theorem is achieved precisely on partial sums of the form (2.4). Since $C(\mathbb{T}^2) \subset L_2(\mathbb{T}^2)$, at first glance this contradicts equality (2.4). But of course there is no contradiction here. In Fefferman's example at each point $(x_1, x_2) \in Q \subset \subset \mathbb{T}^2$ partial sums $\widetilde{M}_k f(x_1, x_2)$ with $m_1(k) = [kx_1]$ and $m_2(k) = [kx_2]$ diverge at (x_1, x_2) , but according to (2.4), converge a.e. on \mathbb{T}^2 .

Obviously, if f_1 is the function from this theorem, then function $f(x_1, \dots, x_N) = f_1(x_1, x_2)$ gives a counterexample for any dimensional case. Since $C(\mathbb{T}^N) \subset L_p(\mathbb{T}^N)$ for any $p \geq 1$, then for rectangular partial sums the problem of a.e. convergence has the following solution: for any $p \geq 1$ the convergence a.e. fails in $L_p(\mathbb{T}^N)$ classes.

It may be worth mentioning that for rectangular partial sums there are examples of functions with high smoothness for which the classical principle of localization is not valid (see [1]). Therefore, for these partial sums too (as in case of spherical partial sums) the principle of generalized localization may become a replacement for the classical principle of localization.

I. Bloshanskiy intensively and in detail studied the generalized localization for rectangular partial sums (see [1]). In particular he proved [10], that generalized localization of rectangular partial sums belongs to those properties which discriminate between two- and three- dimensional Fourier series. Namely, he showed that generalized localization holds in the classes $L_p(\mathbb{T}^N)$, $p > 1$, for $N = 2$ but not for $N \geq 3$.

If $p = 1$, generalized localization for rectangular partial sums does not hold for any $N \geq 2$. To prove this statement it is sufficient to consider function $f(x) = f_1(x_1) \cdot f_2(x_2, \dots, x_N)$, where f_1 is Kolmogorov's example of function in $L_1(\mathbb{T})$ with an unboundedly divergent Fourier series and f_2 is a L_1 -function which is equal to zero in a neighborhood of the origin.

3. GENERALIZED LOCALIZATION PRINCIPLE FOR THE MULTIPLE FOURIER INTEGRALS

1. The Carbery-Soria theorem. For the spherical partial sums (1.1) the Luzin's conjecture is still open. But, as we will see the next section, the problem of generalized localization principle has a positive solution.

By the definition of Fourier coefficients, one can write

$$S_\lambda f(x) = \int_{\mathbb{T}^N} \theta(x - y, \lambda) f(y) dy,$$

where the kernel has the form

$$\theta(x, \lambda) = (2\pi)^{-N} \sum_{|n|^2 < \lambda} e^{inx}.$$

To investigate convergence of $S_\lambda f(x)$ when $\lambda \rightarrow \infty$ one has to investigate the asymptotic behavior of $\theta(x, \lambda)$ for large λ . Of course, this is not an easy problem. But if we change the sum into an integral, then we have the explicit form of the kernel:

$$e(x, \sqrt{\lambda}) = (2\pi)^{-N} \int_{|\xi|^2 < \lambda} e^{ix\xi} d\xi = C \frac{J_{\frac{N}{2}}(|x|\sqrt{\lambda})}{(|x|\sqrt{\lambda})^{N/2}},$$

where $J_\nu(t)$ is the Bessel function. On the other hand, $e(x, \lambda)$ is the kernel of the integral operator

$$(3.1) \quad E_\lambda f(x) = \int_{\mathbb{R}^N} e(x - y, \lambda) f(y) dy = (2\pi)^{-\frac{N}{2}} \int_{|\xi| < \lambda} \hat{f}(\xi) e^{ix\xi} d\xi,$$

which is the spherical partial integrals of the multiple Fourier integral of a function $f \in L_2(\mathbb{R}^N)$. Here a Fourier transform of function f is defined as

$$\hat{f}(\xi) = (2\pi)^{-\frac{N}{2}} \int_{\mathbb{R}^N} f(x) e^{-ix\xi} dx.$$

For the spherical partial integrals of multiple Fourier integrals the generalized localization principle in $L_p(\mathbb{R}^N)$ has been investigated by many authors from 1983 till 2016 (Sjölin, P., Carbery, A., Soria F., Rubio de Francia, J. L., Vega, L., Bastys A., and others (see [11] - [18]).

In particular, in the fundamental paper [13] the following theorem has been proved

Theorem (Carbery, Soria (1988)). *Let $f \in L_p(\mathbb{R}^N)$, $2 \leq p < 2N/(N - 1)$ and $f = 0$ on an open set $\Omega \subset \mathbb{R}^N$. Then*

$$(3.2) \quad \lim_{\lambda \rightarrow \infty} E_\lambda f(x) = 0$$

holds a.e. on Ω .

An earlier variant of this theorem where $f \in L_2(\mathbb{R}^N)$ was assumed to have a compact support was obtained by Sjölin, [11].

We prove Sjölin's theorem here by the method of Carbery and Soria; the proof of the above theorem (Carbery and Soria) is significantly more complicated. It may be worth mentioning that the original proof of Sjölin is completely different.

Let $f \in L_2(\mathbb{R}^N)$ and f have a compact support. In these conditions we must prove that the equality (3.2) holds a.e. on $\Omega = \mathbb{R}^N \setminus \text{supp} f$. If $x \in \Omega$ an arbitrary point, then to do this it suffices to show validity of (3.2) a.e. on a ball with center at x and sufficiently small radius R , so that this ball belongs to Ω . Therefore without loss of generality we may suppose, that f is supported outside of this ball or assuming $\{|x| < R\} \subset \Omega$, f is supported in $\{R \leq |x| \leq A\}$, and prove convergence to zero of $E_\lambda f(x)$ a.e. on the ball $\{|x| < r\}$ for any $r < R$. But this statement can be proved by a standard technique based on the following theorem (see [21]).

Theorem 3.1. *Let $E_* f(x) = \sup_{\lambda > 0} |E_\lambda f(x)|$ be the maximal operator. Then for any $r < R$ there exists a constant $C = C(R, r)$, such that*

$$(3.3) \quad \int_{|x| \leq r} |E_* f(x)|^2 dx \leq C \int_{\mathbb{R}^N} |f(x)|^2 dx.$$

First we prove some auxiliary assertions. Let $\chi(x) \in C_0^\infty(\mathbb{R}^N)$ be a radial function, such that

$$\chi(x) = \begin{cases} 0, & |x| < \frac{1}{3}(R-r), \\ 1, & \frac{2}{3}(R-r) < |x| \leq 2A. \end{cases}$$

If we denote $e_\lambda(x) = e(x, \lambda)\chi(x)$, then we have

$$E_\lambda f(x) = \int_{\mathbb{R}^N} e_\lambda(x-y)f(y)dy, \text{ for all } x, \text{ with } |x| \leq r,$$

since f is supported in $\{R \leq |x| \leq A\}$. Therefore to prove estimate (3.3) it suffices to obtain the inequality

$$(3.4) \quad \int_{\mathbb{R}^N} \sup_{\lambda > 0} |e_\lambda * f|^2 dx \leq C \int_{\mathbb{R}^N} |f(x)|^2 dx.$$

Now we need some estimates for the Fourier transform of the function $e_\lambda(x)$, which we denote by $\widehat{e}_\lambda(\eta)$. These estimates were obtained in [13] and for the convenience of readers we will give the proof from [13] here.

Lemma 3.2. *For $j = 0, 1$ and for an arbitrary $l \in \mathbb{N}$ there exists a constant C_l , depending on l, r and R , such that for all $\lambda \in \mathbb{R}$ and $\eta \in \mathbb{R}^N$ one has*

$$\left| \frac{d^j}{d\lambda^j} \widehat{e}_\lambda(\eta) \right| \leq \frac{C_l}{(1 + \|\eta\| - \lambda)^l}.$$

Proof. We shall consider the case $j = 0$, the case $j = 1$ being handled similarly (see [13]). By the definition of the Fourier transform we have

$$\begin{aligned} \widehat{e}_\lambda(\eta) &= (2\pi)^{-\frac{3}{2}N} \int_{\mathbb{R}^N} \left(\int_{|\xi| < \lambda} e^{ix\xi} d\xi \chi(x) \right) e^{-ix\eta} dx = (2\pi)^{-N} \int_{|\xi| < \lambda} (2\pi)^{-\frac{N}{2}} \int_{\mathbb{R}^N} \chi(x) e^{-ix(\eta-\xi)} dx d\xi = \\ &= (2\pi)^{-N} \int_{|\xi| < \lambda} \widehat{\chi}(\eta - \xi) d\xi = (-2\pi)^{-N} \int_{|\eta-\xi| < \lambda} \widehat{\chi}(\xi) d\xi. \end{aligned}$$

First suppose $|\eta| < \lambda$. Since $\chi(0) = 0$ and for any integer j one has the estimate $|\widehat{\chi}(\xi)| \leq C_j(1 + |\xi|)^{-j}$, then

$$\begin{aligned} |\widehat{e}_\lambda(\eta)| &= \left| -(-2\pi)^{-N} \int_{|\eta-\xi| \geq \lambda} \widehat{\chi}(\xi) d\xi \right| \leq C_j \int_{|\eta-\xi| \geq \lambda} (1 + |\xi|)^{-j} d\xi \leq \\ &\leq C_j \int_{|\xi| > \lambda - |\eta|} (1 + |\xi|)^{-j} d\xi \leq \frac{C_l}{(1 + \|\eta\| - \lambda)^l}. \end{aligned}$$

Similarly, if $|\eta| > \lambda$,

$$|\widehat{e}_\lambda(\eta)| = \left| (-2\pi)^{-N} \int_{|\eta-\xi| < \lambda} \widehat{\chi}(\xi) d\xi \right| \leq C_j \int_{|\xi| > |\eta| - \lambda} (1 + |\xi|)^{-j} d\xi \leq \frac{C_l}{(1 + \|\eta\| - \lambda)^l}.$$

□

It is not hard to verify that from this lemma one has

Corollary 3.3. *Let $j = 0, 1$. Uniformly in $\eta \in \mathbb{R}^N$ we have*

$$\int_0^\infty \left| \frac{d^j}{d\lambda^j} \widehat{e}_\lambda(\eta) \right|^2 d\lambda < C.$$

Now we are ready to prove estimate (3.4). Since

$$[e_\lambda * f]^2 = 2 \int_0^\lambda (e_t * f) \frac{d}{dt} (e_t * f) dt$$

and $2ab \leq a^2 + b^2$

$$\sup_{\lambda > 0} |e_\lambda * f|^2 \leq \int_0^\infty |e_t * f|^2 dt + \int_0^\infty \left| \frac{d}{dt} e_t * f \right|^2 dt,$$

or

$$\int_{\mathbb{R}^N} \sup_{\lambda > 0} |e_\lambda * f|^2 \leq \int_{\mathbb{R}^N} |\widehat{f}(\eta)|^2 \int_0^\infty |\widehat{e}_t(\eta)|^2 dt d\eta + \int_{\mathbb{R}^N} |\widehat{f}(\eta)|^2 \int_0^\infty \left| \frac{d}{dt} \widehat{e}_t(\eta) \right|^2 dt d\eta.$$

Finally, making use of Corollary 3.3 and the fact that f is an L_2 -function we obtain estimate (3.4).

Thus theorem of Carbery and Soria is proved for functions $f \in L_2(\mathbb{R}^N)$ with compact support.

Let $A(\xi) = \sum_{|\alpha|=m} a_\alpha \xi^\alpha$ be an arbitrary elliptic homogeneous polynomial on $\xi \in \mathbb{R}^N$ with

constant coefficients a_α , i.e. $A(\xi) > 0$ for all $\xi \neq 0$.

With this polynomial $A(\xi)$ we can define a partial sums of the multiple Fourier integrals as

$$E(A, \lambda, f)(x) = (2\pi)^{-\frac{N}{2}} \int_{A(\xi) < \lambda} \widehat{f}(\xi) e^{ix\xi} d\xi.$$

Note, if $A(\xi) = |\xi|^2$, then the partial sums $E(A, \lambda, f)(x)$ coincides with the spherical partial sums $E_{\sqrt{\lambda}} f(x)$.

In the paper [22], using the method of Carbery-Soria [13], the following theorem is proved.

Theorem 3.4. *Let $A(\xi)$ be an arbitrary elliptic polynomial. Let $f \in L_p(\mathbb{R}^N)$, $2 \leq p < 2N/(N-1)$ and $f = 0$ on an open set $\Omega \subset \mathbb{R}^N$. Then $\lim_{\lambda \rightarrow \infty} E(A, \lambda, f)(x) = 0$ holds a.e. on Ω .*

So in case of an arbitrary elliptic polynomial $A(\xi)$ we have the same result as for the spherical partial sums $E_\lambda f(x)$. We note that, unlike to the generalized localization, the conditions for the classical Riemann localization of $E(A, \lambda, f)(x)$ strictly depends on the geometry of the set $\{\xi \in \mathbb{R}^N : A(\xi) < 1\}$, namely on the number of nonzero curvatures of the level surface $\{\xi \in \mathbb{R}^N : A(\xi) = 1\}$ (see [23] and [24]).

2. Generalized localization for distributions. We recall the Schwartz space S is the class $C^\infty(\mathbb{R}^N)$ being equipped with a family of seminorms

$$d_{\alpha, \beta}(\phi) = \sup_{x \in \mathbb{R}^N} |x^\alpha D^\beta \phi(x)|.$$

We also consider a class of tempered distributions S' defined as dual to S .

Along with S we also consider \mathcal{E} , i.e. $C^\infty(\mathbb{R}^N)$ being equipped with a family of seminorms

$$\rho_{\alpha, K}(\phi) = \sup_{x \in K} |D^\alpha \phi(x)|.$$

As usual we denote its conjugate space by \mathcal{E}' . Distributions from \mathcal{E}' have the following important properties:

- every $f \in \mathcal{E}'$ has a finite support,
- for any $f \in \mathcal{E}'$ one has $\widehat{f} \in C^\infty$ (the Paley-Wiener theorem),
- for every $f \in \mathcal{E}'$ there is $l \in \mathbb{R}$ such that $f \in H^l$.

Definition 3.5. *We say that tempered distribution f belongs to the Sobolev class H^l if \widehat{f} is a regular distribution such that*

$$\|f\|_{H^l}^2 = \int |\widehat{f}(\xi)|^2 (1 + |\xi|^2)^l d\xi < \infty.$$

Example. Consider $D^\alpha \delta(x) \in \mathcal{E}'$, where $\delta(x)$ is the Dirac delta-function. We have

$$(D^{\widehat{\alpha}} \delta)(\xi) = (-1)^{|\alpha|} \xi^\alpha.$$

Therefore $D^\alpha \delta(x) \in H^{-l}$ with $l > \frac{N}{2} + |\alpha|$.

Since for any distribution $f \in \mathcal{E}'$ its Fourier transform is infinitely differentiable function, then we can define the Riesz means of order $s \geq 0$ of the spherical partial integrals as follows

$$E_\lambda^s f(x) = (2\pi)^{-\frac{N}{2}} \int_{|\xi| < \lambda} \left(1 - \frac{|\xi|^2}{\lambda^2}\right)^s \widehat{f}(\xi) e^{ix\xi} d\xi.$$

The authors of the paper [18] obtained the necessary and sufficient conditions for the generalized localization of the Riesz means of distributions to be hold. These conditions are formulated in the following two theorems.

Theorem 3.6. *Let $f \in \mathcal{E}' \cap H^{-l}$, $l \geq 0$. Then for $s \geq l$, equality*

$$\lim_{\lambda \rightarrow \infty} E_\lambda^s f(x) = 0$$

holds true a.e. on $\mathbb{R}^N \setminus \text{supp } f$.

Theorem 3.7. *For any s and l such that $0 \leq s < l$ there are $f \in H^{-l} \cap \mathcal{E}'$ and a set K of positive measure such that $\text{supp } f \cap K = \emptyset$ and*

$$\limsup_{\lambda \rightarrow \infty} |E_\lambda^s f(x)| = \infty, \text{ a.e. } x \in K.$$

In Theorem 3.6 it is actually proved that the action on a distribution of the Riesz means of order s makes it roughly s -smoother.

Observe, as mentioned above for $E_\lambda^0 f(x) \equiv E_\lambda f(x)$ the generalized localization holds true for L_2 -functions. The following theorem (proved in [18]) states, that for a negative order Riesz means the generalized localization fails.

Theorem 3.8. *For any $\epsilon > 0$ there are compactly supported $g \in L_2(\mathbb{R}^N)$ and set $K : \text{mes}(K) > 0$ and $\text{supp } g \cap K = \emptyset$ such that*

$$\sup_{\lambda > 0} |E_\lambda^{-\epsilon} g(x)| = \infty, x \in K.$$

The arguments of the proof here essentially consist in reducing the problem to the generalized localization of spherical means $E_\lambda g(x)$ on L_p with $p < 2$, for which there is a counterexample due to Bastys [15].

We note, that the proof of Theorem 3.7 is essentially based on Theorem 3.8.

3. Generalized localization for continuous wavelet decompositions.

In the paper [19], the authors studied the generalized localization principle for multidimensional spherically symmetric continuous wavelet decompositions. It should be noted that the pointwise convergence of one-dimensional and multidimensional continuous wavelet decompositions has been investigated by many authors (a detailed survey of their work can be found in the paper [20] by Ashurov and Butaev). In this case, to ensure the almost-everywhere convergence of multidimensional (and even one-dimensional) spherically symmetric continuous wavelet decompositions, it is necessary to impose an additional restriction on the rate of decay at infinity of the corresponding wavelets. In the above paper [19], using the result of Carbery-Soria [13], it was shown that the generalized localization principle holds for multidimensional spherically symmetric continuous wavelet decompositions (with completely arbitrary wavelets) of functions $f \in L_p(\mathbb{R}^N)$, provided $2 \leq p < 2N/(N-1)$.

4. GENERALIZED LOCALIZATION PRINCIPLE FOR THE MULTIPLE FOURIER SERIES

1. The main result. If we turn back to the multiple Fourier series (1.1) and consider the classes $L_p(\mathbb{T}^N)$ when $1 \leq p < 2$, then as A. Bastys [15] has proved, following Fefferman in making use of the Kakeya's problem, that the generalized localization for S_λ is not valid, i.e. there exists a function $f \in L_p(\mathbb{T}^N)$, such that on some set of positive measure, contained in $\mathbb{T}^N \setminus \text{supp } f$, we have

$$\overline{\lim}_{\lambda \rightarrow \infty} |S_\lambda f(x)| = +\infty.$$

It should be noted that in [15] this result is also proved for the spherical partial integrals $E_\lambda f(x)$.

The following result gives a complete solution of the generalized localization problem for the spherical partial sums of the multiple Fourier series. It may be worth mentioning that this problem was first formulated in 1976 in a review paper [2].

Theorem 4.1. *Let $f \in L_2(\mathbb{T}^N)$ and $f = 0$ on an open set $\Omega \subset \mathbb{T}^N$. Then the equality $\lim_{\lambda \rightarrow \infty} S_\lambda f(x) = 0$ holds a.e. on Ω .*

Thus the problem of generalized localization for S_λ is completely solved in classes $L_p(\mathbb{T}^N)$, $p \geq 1$: if $p \geq 2$ then we have the generalized localization and if $p < 2$, then the generalized localization fails.

Recall that in order for classical Riemann localization to take place for $S_\lambda f$, the expandable function f must have $\frac{N-1}{2}$ "derivatives", i.e. f must belong to the Sobolev class $H^l(\mathbb{T}^N)$ with $l \geq \frac{N-1}{2}$ (see [2]).

Let us introduce the maximal operator

$$S_* f(x) = \sup_{\lambda > 0} |S_\lambda f(x)|.$$

The proof of Theorem 4.1 is based on the following estimate of this operator.

Theorem 4.2. *Let Ω be an open subset of \mathbb{T}^N . Then for any compact set $K \subset \Omega$ there exists a constant $C_K > 0$ such that for any function $f \in L_2(\mathbb{T}^N)$ with $\text{supp } f \subset \mathbb{T}^N \setminus \Omega$ one has*

$$(4.1) \quad \|S_* f(x)\|_{L_2(K)} \leq C_K \|f\|_{L_2(\mathbb{T}^N)}.$$

The formulated theorems are easily transferred to the case of non-spherical partial sums of multiple Fourier series (see [22], [25]).

We should also note, that in the remarkable paper of C. Kenig and P. Tomas [26] the authors proved, by making use of transference techniques, the equivalence of convergence a.e. of spherical partial sums of multiple Fourier series and integrals. It may be worth mentioning that Theorem 4.1 on the generalized localization does not follow from the Carbery - Soria theorem on Fourier integrals [13], by application of the C. Kenig and P. Tomas theorem [26]. The reason here is that, we integrate on the left hand side of (4.1) only over a compact K , where the function vanishes (not over whole domain \mathbb{T}^N), and therefore we can not apply the duality technique, which is the key step in the proof of C. Kenig and P. Tomas. Nevertheless, to prove Theorem 4.1 we have used many of original ideas from A. Carbery and F. Soria [13].

2. Auxiliary assertions. It is not hard to verify, that Theorem 4.2 is an easy corollary of the following theorem:

Theorem 4.3. *Let $f \in L_2(\mathbb{T}^N)$ and $f = 0$ on the ball $\{|x| < R\}$, $R < 1$. Then for any $r < R$ there exists a constant $C = C(R, r)$, such that*

$$(4.2) \quad \int_{|x| \leq r} |S_* f(x)|^2 dx \leq C \int_{\mathbb{T}^N} |f(x)|^2 dx.$$

The proof of Theorem 4.3 can be found in [5]. We present here the main ideas of the proof.

So we assume that $f = 0$ on the fixed ball $\{|x| < R\}$ and fix a number $r < R < 1$.

Let $\chi_b(t)$ be the characteristic function of the segment $[0, b]$. We denote by $\varphi_1(t)$ a smooth function with $\chi_{(R-r)/3}(t) \leq \varphi_1(t) \leq \chi_{2(R-r)/3}(t)$ and put $\varphi_2(t) = 1 - \varphi_1(t)$. Now we define a new function $\psi(x)$ as follows: $\psi(x) = \varphi_2(|x|)$, when $x \in \mathbb{T}^N$ and otherwise it is a 2π -periodical on each variable x_j function.

Let us denote $\theta_\lambda(x) = \theta(x, \lambda)\psi(x)$. Then we have

$$S_\lambda f(x) = \int_{\mathbb{T}^N} \theta_\lambda(x-y)f(y)dy, \text{ for all } x, \text{ with } |x| \leq r,$$

since f is supported in $\{|x| \geq R\}$. Therefore, if we denote the last integral by $\theta_\lambda * f$, then to prove the estimate (4.2) it suffices to obtain the inequality

$$(4.3) \quad \int_{\mathbb{T}^N} \sup_{j > 0} |\theta_j * f|^2 dx \leq C \int_{\mathbb{T}^N} |f(x)|^2 dx,$$

where sup is taken over all positive integers.

Remark 4.4. *Of course, by analogy with the multiple Fourier integrals (see (3.1)), in partial sums (1.1), one could sum over the set $\{|n| < \lambda\}$. But then we would have to take the sup in (4.3) over all positive real numbers, instead of all positive integers.*

Now we need some estimates for the Fourier coefficients of the function $\theta_j(x)$, which we denote by $(\theta_j)_n$, $j \in \mathbb{N}$ (the set of positive integers), $n \in \mathbb{Z}^N$.

Lemma 4.5. *For an arbitrary $l \in \mathbb{N}$ there exists a constant C_l , depending on l, r and R , such that for all $j \in \mathbb{N}$ and $n \in \mathbb{Z}^N$ one has*

$$(4.4) \quad |(\theta_j)_n| \leq \frac{C_l}{(1 + ||n| - \sqrt{j}|)^l}.$$

Let $\{\psi_m\}$ be the Fourier coefficients of function $\psi(x)$: $\psi_m = (2\pi)^{-N} \int_{\mathbb{T}^N} \psi(y) e^{-imy} dy$. Then for any integer $q \geq 0$ there exists a constant c_q , depending on $(R - r)$, such that

$$|\psi_m| \leq \frac{c_q}{(1 + |m|)^q}.$$

Lemma is an easy consequence of this estimate (see the proof of Lemma 3.2).

It should be noted the importance of considering the function $\theta_j(x)$ instead of $\theta(x, j)$. The Fourier coefficients $(\theta(x, j))_n$ of function $\theta(x, j)$ have the form

$$(\theta(x, j))_n = \begin{cases} 1, & j > |n|, \\ 0, & j \leq |n|. \end{cases}$$

These numbers do not decrease as $j \rightarrow \infty$, while the numbers $(\theta_j)_n$ have a sufficiently strong decrease (see estimate (4.4)).

Let $(\Theta_j)_n = (\theta_{j+1})_n - (\theta_j)_n$, that is,

$$(\Theta_j)_n = (2\pi)^{-N} \sum_{|m|^2=j} \psi_{m-n} = (2\pi)^{-N} \sum_{|n-m|^2=j} \psi_m$$

(if the Diophantine equation $|m - n|^2 = j$ does not have a solution, then $(\Theta_j)_n = 0$). Using the same arguments as above, one can prove the following estimate.

Lemma 4.6. *For any k and $l \in \mathbb{N}$, there exists a constant C_l such that*

$$(4.5) \quad \sum_{k \leq \sqrt{j} < k+1} |(\Theta_j)_n|^2 \leq \frac{C_l}{(1 + ||n| - k|)^l}.$$

If we sum this estimate with respect to k from 0 to ∞ , then we obtain the next corollary.

Corollary 4.7. *Uniformly in $n \in \mathbb{Z}^N$ one has*

$$\sum_{j=0}^{\infty} |(\Theta_j)_n|^2 = \sum_{k=0}^{\infty} \sum_{k \leq \sqrt{j} < k+1} |(\Theta_j)_n|^2 \leq C.$$

Now we try to estimate the same sum for $(\theta_j)_n$. Let $k \leq \sqrt{j} < k + 1$, i.e. $j = k^2 + p$, $p = 0, 1, \dots, 2k$. From the estimate (4.4) we have

$$\sum_{k \leq \sqrt{j} < k+1} |(\theta_j)_n|^2 = \sum_{p=0}^{2k} |(\theta_{k^2+p})_n|^2 \leq \frac{C_l \cdot k}{(1 + ||n| - k|)^l},$$

therefore

$$(4.6) \quad \sum_{j=0}^{\infty} |(\theta_j)_n|^2 = \sum_{k=0}^{\infty} \sum_{k \leq \sqrt{j} < k+1} |(\theta_j)_n|^2 \leq C \cdot |n|.$$

But this estimate is not sufficient for our purpose. Indeed, let $\Theta_j(x) = \theta_{j+1}(x) - \theta_j(x)$. Then $\theta_{j+1} * f + \theta_j * f = 2\theta_j * f + \Theta_j * f$. Note the Fourier coefficients of the function $\Theta_j(x)$ are the numbers $(\Theta_j)_n$, introduced above.

Now we use the idea of the proof of estimate (3.4). If for a sequence of numbers $\{F_q\}$ we have $F_0 = 0$, then

$$F_q^2 = \sum_{j=0}^{q-1} [F_{j+1} - F_j][F_{j+1} + F_j], \quad q \geq 1.$$

Hence

$$[\theta_q * f]^2 = \sum_{j=0}^{q-1} [\Theta_j * f]^2 + 2 \sum_{j=0}^{q-1} [\Theta_j * f][\theta_j * f],$$

or

$$(4.7) \quad \sup_{q>0} |\theta_q * f|^2 \leq \sum_{j=0}^{\infty} |\Theta_j * f|^2 + 2 \sum_{k=0}^{\infty} \sum_{p=0}^{2k} |\Theta_{k^2+p} * f| |\theta_{k^2+p} * f|.$$

Integrating over T^N and making use of the inequality $2ab \leq a^2 + b^2$ one has

$$(4.8) \quad \int_{T^N} \sup_{q>0} |\theta_q * f|^2 \leq 2 \sum_n |f_n|^2 \sum_{j=0}^{\infty} |(\Theta_j)_n|^2 + \sum_n |f_n|^2 \sum_{j=0}^{\infty} |(\theta_j)_n|^2.$$

By virtue of Corollary 4.7 and the fact that f is an L_2 -function, we can estimate first sum on the right hand side of (4.8) by $C \sum_n |f_n|^2 = C \int_{\mathbb{T}^N} |f(x)|^2 dx$. But the second sum we can estimate, using (4.6), only by $C \sum_n |f_n|^2 |n|$. Thus we must have a better estimate for the sum (4.6).

To prove Theorem 4.3 we need the same estimate as (4.5) for the sum $\sum_{p=0}^{2k} (p+1)^2 |(\Theta_{k^2+p})_n|^2$ (that is, the value under the sum (4.5) is multiplied by $(p+1)^2$). To obtain this estimate we must show, that $|(\Theta_{k^2+p})_n|^2$ decreases fast enough in p . Unfortunately this is not always the case. Nevertheless if we change the order of summation in the last sum (since the summands are positive, this is always possible), say first take the sum over some sets Q_q^k , $q = 0, 1, \dots$ of parameters p , and then take the sum over $q = 0, 1, \dots$, we can succeed to prove the necessary estimate.

Lemma 4.8. *There exist sets Q_q^k , $q = 0, 1, \dots, 2k-1$, of integers p , $0 \leq p \leq 2k$, such, that for any l , one has an estimate with a constant C_l :*

$$(4.9) \quad \sum_{q=0}^{2k-1} (q+1)^2 \sum_{p \in Q_q^k} |(\Theta_{k^2+p})_n|^2 \leq \frac{C_l}{(1 + \sqrt{|n| - k})^l}.$$

Our next aim is to define the sets Q_q^k , $q = 0, 1, \dots$ with above property.

Denote by y_0 the intersection point of the ball $\{x \in \mathbb{R}^N : |x - n| \leq k + 1\}$ with the straight line On that passes through the origin and point n (the nearest one to the origin). Let T_{y_0} be the tangential hyperplane to the ball $\{x \in \mathbb{R}^N : |x - n| \leq k + 1\}$ at the point y_0 . Let $B_0 := \{y \in T_{y_0} : |y - y_0| < 1\}$ and $B_j := \{y \in T_{y_0} : \sqrt{j} \leq |y - y_0| < \sqrt{j+1}\}$, where $j = 1, 2, \dots, 2k-1$. Let C_j^k , $j = 0, 1, \dots, 2k-1$, be the N -dimensional cylinders with the base B_j and with the axis parallel to On and the length $|n|$. Consider the ring $K = \{x \in \mathbb{R}^N : k \leq |x - n| < (k+1)\}$ and divide it in to the following sets: $P_j^k = K \cap C_j^k$, $j = 0, 1, \dots, 2k-1$.

Let us define the sets Q_q^k , $q = 0, 1, \dots, 2k-1$, as follows. Let Q_q^k be the set of those integers p , $0 \leq p \leq 2k$, for which the Diophantine equation $|m - n|^2 = k^2 + p$ has a solution in P_q^k . If P_q^k does not contain any of solutions of equation $|m - n|^2 = k^2 + p$, for any p , then we assign to the set Q_q^k one of those parameters p that are not included in the previous sets Q_j^k , $j = 0, 1, \dots, q-1$. If there are no such p 's left, then we define Q_j^k , $j = q, q+1, \dots, 2k-1$ as the empty set.

In the proof of Lemma 4.10 we need to know how many at most parameters p does the set Q_q^k contain. Since the sets P_q^k are sufficiently small, then there should not be many such parameters. Indeed, the length of the projection of P_q^k on the axis of Ox_1 is at most $2\sqrt{q+1}$. Consequently, if, for a fixed p , there is a solution of the Diophantine equation $|m - n|^2 = k^2 + p$, provided $m \in P_q^k$, then the first coordinates m_1 of the numbers m , take at most $[2\sqrt{q+1}]$ ($[a]$ is the integer part of the number a) different values. When p varies from 0 to $2k$, each of these numbers m_1 can repeat at most 2^{N-1} times (that is the number of all vectors $m = (m_1, m_2, \dots, m_N) \in \mathbb{Z}^N$ within unit cube with fixed m_1). Hence each set Q_q^k has at most $2^N [\sqrt{q+1}]$ parameters p with the above property (one can give a better estimate for this number, but for our purpose this estimate is sufficient).

If we denote $S_p = \{m \in \mathbb{Z}^N : |m - n|^2 = k^2 + p\}$ ($p = 0, 1, \dots, 2k$), then with this choice of Q_q^k we have the following estimates.

If $|n| \geq k + 1$, then

$$\min_{m \in S_p, p \in Q_q^k} |m| \geq \sqrt{(|n| - k - 1)^2 + q}.$$

If $k < |n| < k + 1$, then

$$\min_{m \in S_p, p \in Q_q^k} |m| \geq \sqrt{q}.$$

If $|n| \leq k$, then

$$\min_{m \in S_p, p \in Q_q^k} |m| \geq \frac{1}{2} \sqrt{(|n| - k)^2 + q}.$$

Lemma 4.8 is a consequence of these estimates.

Corollary 4.9. *Uniformly in $n \in \mathbb{Z}^N$, one has*

$$\sum_{k=0}^{\infty} \sum_{q=0}^{2k-1} (q+1)^2 \sum_{p \in Q_q^k} |(\Theta_{k^2+p})_n|^2 \leq C.$$

Now we turn back to the Fourier coefficients $(\theta_j)_n$. From Lemma 4.5 we have the following estimate.

Lemma 4.10. *Uniformly in $n \in \mathbb{Z}^N$, one has*

$$\sum_{k=0}^{\infty} \sum_{q=0}^{2k-1} (q+1)^{-2} \sum_{p \in Q_q^k} |(\theta_{k^2+p})_n|^2 \leq C.$$

Proof. As mentioned above, each Q_q^k has at most $2^N [\sqrt{q+1}]$ parameters p . Therefore, by virtue of Lemma 4.5, one has

$$\sum_{k=0}^{\infty} \sum_{q=0}^{2k-1} (q+1)^{-2} \sum_{p \in Q_q^k} |(\theta_{k^2+p})_n|^2 \leq \sum_{k=0}^{\infty} \frac{C_l}{(1+|n|-k)^l} \sum_{q=0}^{2k-1} \frac{2^N [\sqrt{q+1}]}{(q+1)^2} \leq C.$$

□

3. Proofs of Theorems. Now we rewrite estimate (4.7) as

$$\sup_{q \in \mathbb{N}} |\theta_q * f|^2 \leq \sum_{j=0}^{\infty} |\Theta_j * f|^2 + 2 \sum_{k=0}^{\infty} \sum_{q=0}^{2k-1} \sum_{p \in Q_q^k} |\Theta_{k^2+p} * f| (q+1) |\theta_{k^2+p} * f| (q+1)^{-1}.$$

Integrating over \mathbb{T}^N and making use of Corollaries 4.7, 4.9 and Lemma 4.10, we have

$$\begin{aligned} \int_{\mathbb{T}^N} \sup_{q \in \mathbb{N}} |\theta_q * f|^2 &\leq \sum_n |f_n|^2 \sum_{j=0}^{\infty} |(\Theta_j)_n|^2 + \\ &+ \sum_n |f_n|^2 \sum_{k=0}^{\infty} \sum_{q=0}^{2k-1} (q+1)^2 \sum_{p \in Q_q^k} |(\Theta_{k^2+p})_n|^2 + \\ &+ \sum_n |f_n|^2 \sum_{k=0}^{\infty} \sum_{q=0}^{2k-1} (q+1)^{-2} \sum_{p \in Q_q^k} |(\theta_{k^2+p})_n|^2 \leq C \sum_n |f_n|^2 = C \int_{\mathbb{T}^N} |f(x)|^2 dx. \end{aligned}$$

Thus, the estimate (4.3) and, consequently, Theorem 4.2 is proved.

Theorem 4.1 can be proved by a standard technique based on Theorem 4.2 (see [21]). Indeed, let $f \in L_2(\mathbb{T}^N)$ and $f = 0$ on $\Omega \subset \mathbb{T}^N$. We must prove

$$\lim_{\lambda \rightarrow \infty} S_\lambda f(x) = 0 \text{ a.e. on } \Omega.$$

Let $K \subset \Omega$ be an arbitrary compact. Then for an arbitrary $\delta > 0$ there exist two 2π -periodical on each argument functions f_1 and f_2 with the following properties: $f = f_1 + f_2$ and $f_1 = f_2 = 0$ on Ω ; $f_1 \in C^\infty(\mathbb{T}^N)$ and $\|f_2\|_{L_2(\mathbb{T}^N)} < \delta$. Moreover for an arbitrary $\varepsilon > 0$ and $\lambda > \lambda_0(\varepsilon)$ one has $|S_\lambda f_1(x)| < \frac{\varepsilon}{2}$ on Ω and $M_\varepsilon = \text{mes}\{x \in K \mid |S_\lambda f_2(x)| > \frac{\varepsilon}{2}\} < \varepsilon$ (this inequality follows from Theorem 4.2).

Therefore, if $\lambda > \lambda_0(\varepsilon)$ then on $K \setminus M_\varepsilon$ one has

$$|S_\lambda f| \leq |S_\lambda f_1| + |S_\lambda f_2| < \varepsilon.$$

Obviously, this means $S_\lambda f(x) \rightarrow 0$ a.e. on Ω .

5. ACKNOWLEDGEMENT

The author conveys thanks to Sh. A. Alimov for discussions of this result and gratefully acknowledges S. Umarov (University New Haven, USA) for support and hospitality.

The author was supported by Foundation for Support of Basic Research of the Republic of Uzbekistan (project number is OT-F4-88).

REFERENCES

1. Alimov, Sh.A., Ashurov, R.R., Pulatov, A.K.: Multiple Fourier Series and Fourier Integrals. Commutative Harmonic Analysis, vol. IV, pp. 197. Springer, Berlin (1992)
2. Alimov, Sh.A., Ilin, V.A., Nikishin, E.M.: Convergence problems of multiple trigonometric series and spectral decompositions. I. Russian Math. Surveys, **31**, 29-86 (1976)
3. Mityagin, B.S., Nikishin, E.M.: On almost-everywhere divergence of Fourier series. Sov. Math. Dokl. **14**, 677-680, (1973)
4. Il'in, V.A.: On a generalized interpretation of the principle of localization for Fourier series with respect to fundamental systems of functions, Sib. Mat. Zh., **9**, 1093-1106 (1968)
5. Ashurov, R., R.: Generalized localization for spherical partial sums of multiple Fourier series, J. Fourier Anal. Appl., 1-10, 2019, /doi.org/10.1007/s00041-019-09697-7
6. Raussen, M., Skau, Ch.: Interview with Abel Prize recipient Lennart Carleson. Notices of the American Mathematical Society, **54** (2): 223-229 (2007)
7. Tevzadze, N. R.: On the convergence of the double Fourier series of a square summable function. Soobshch. Akad. Nauk Gruz. SSR. **2**, 277-279 (1970)
8. Sjölin, P.: Convergence almost-everywhere of certain singular integrals, Ark. Mat. **1**, 65-90 (1971)
9. Fefferman, C.: On the divergence of multiple Fourier series, Bull. Am. Math. Soc. **77**, 191-195 (1971)
10. Bloshanskiy, I. L.: On the uniform convergence of multiple trigonometric series and Fourier integrals, Math. Notes **18**, 675-684 (1975)
11. Sjölin, P.: Regularity and integrability of spherical means, Monatsh. Math. **96**, 277-291 (1983)
12. Bastys, A.J.: The generalized localization principle for an N-fold Fourier integral, Sov. Math. Dokl., **278**, 777-778 (1984)
13. Carbery, A., Soria F.: Almost everywhere convergence of Fourier integrals for functions in Sobolev spaces, and an L_2 -localization principle, Revista Mat. Iberoamericana **4**, 319- 337 (1988)
14. Carbery, A., Rubio de Francia, J. L., Vega, L.: Almost everywhere summability of Fourier integrals, J. London Math. Soc. **38**, 513-524 (1988)
15. Bastys, A.J.: Generalized localization of Fourier series with respect to the eigenfunctions of the Laplace operator in the classes L_p , Litovskii Matematicheskii Sbornik, **31**, 387-405 (1991)
16. Carbery, A., Romera, E., Soria, F.: Radial weights and mixed norm inequalities for the disc multiplier, J. Funct. Anal. **109**, 52-75 (1992)
17. Carbery, A. Soria, F.: Pointwise Fourier inversion and localization in R^n , J. Fourier Anal. Appl., **3**, Special Issue, 847-858 (1997)
18. Ashurov, R.R., Butaev, A.: On Generalized Localization of Fourier Inversion for Distributions, Contemporary Mathematics, USA, **672**, 33-50 (2016)
19. Ashurov, R. R., Faiziev, Yu. E.: Generalized localization principle for continuous wavelet decompositions, Math. Notes **106**, 75-81 (2019)
20. Ashurov, R.R., Butaev, A.: On pointwise convergence of continuous wavelet transforms, Uzbek. Math. J., **1**, 426 (2018).
21. Stein, E.M., Weiss, G.: Introduction to Fourier analysis on Euclidean spaces, Princeton University Press, Princeton, New Jersey, (1971)
22. Ashurov, R. R., Ahmedov, A., Ahmad Rodzi b. Mahmud,: The generalized localization for multiple Fourier integrals, J. Math. Anal. Appl., **371** 832-841(2010)
23. Ashurov, R., R.: On conditions for localization of spectral expansions corresponding to elliptic operators with constant coefficients, Math. Notes, **33**, 434-439 (1983)
24. Ashurov, R., R.: On localization conditios for multiple trigonometric Fourier series, Sov. Math. Dokl. **31**, 496-499 (1985)
25. Ashurov, R.R., Butaev, A.: On the Pinsky phenomenon, J. Fourier Anal. Appl. **16**, 804-812 (2010)
26. Kenig, C. E., Tomas, P.A.: Maximal operators defined by Fourier multipliers// Studia Math. **68**, 79-83 (1980).

INSTITUTE OF MATHEMATICS, UZBEKISTAN ACADEMY OF SCIENCE

Current address: Institute of Mathematics, Uzbekistan Academy of Science, Tashkent, 81 Mirzo
Ulugbek str. 100170

E-mail address: ashurovr@gmail.com